TITLE
V3 Interneurons Regulate Locomotor Vigor by Recruitment of Spinal Motor
Neurons During Fictive Swimming in Larval Zebrafish

ABBREVIATED TITLE
V3 Interneurons Regulate Locomotor Vigor

AUTHOR NAMES AND AFFILIATIONS
Timothy D. Wiggin[†], Jacob E. Montgomery[†], Amanda J. Brunick, Jack H. Peck,
Mark A. Masino

University of Minnesota, Department of Neuroscience, Minneapolis, MN 55455

[†] Equal contributions.

Corresponding author email: masino@umn.edu

Number of Pages: 47
Number of Figures: 7
Number of Tables: 1
Number of Words for:
        Abstract   206
        Introduction   521
        Discussion   1,310

Conflict of Interest Statement:
The authors declare no competing financial interests.

Acknowledgements:
The authors appreciate the helpful suggestions and feedback from Drs. Ronald
Calabrese and Ronald Harris-Warrick. We also thank Kayce Vanpelt for technical
support and Marc Tye and the staff of the University of Minnesota Zebrafish Core
Facility for animal care. Finally, we thank Drs. Koichi Kawakami and Misha
Ahrens for kindly providing *Tg(UAS:GCaMP6s)^{nk13a}* fish.

Financial Support:
This work was supported by National Institutes of Health (http://www.nih.gov/)
R01 NS065054 (MAM), R01 NS094176 (MAM), F31 NS083110 (TDW), R25
NS083059 (to Dr. Robert Meisel), Regenerative Medicine Minnesota (JEM), and
the Grant-in-Aid of Research, Artistry, and Scholarship (MAM).

ABSTRACT

Survival for vertebrate animals is dependent on the ability to successfully find food, locate a mate, and avoid predation. Each of these behaviors requires fine motor control, which is set by a combination of kinematic properties. For example, the frequency and amplitude (vigor; strength) of motor output combine to determine features of locomotion such as distance traveled and speed. Although there is a good understanding of how different populations of excitatory spinal interneurons establish locomotor frequency, there is not a mechanistic understanding for how locomotor vigor is established. Recent evidence indicates that locomotor vigor is regulated in part by subsets of identified excitatory spinal interneurons (INs), such as the V2a neuronal population in adult zebrafish. Here we provide evidence that the majority of V3 interneurons (V3-INs), which are a developmentally and genetically defined population of ventromedial glutamatergic spinal neurons, are active during fictive swimming. Further, that targeted ablation of V3-INs reduces the proportion of active MNs during fictive swimming, but ablation does not affect the locomotor frequencies produced. These data are consistent with a role of V3-INs in providing excitatory drive to spinal motor neurons during swimming in larval zebrafish, which suggests that locomotor vigor (but not locomotor frequency) may be regulated, in part, by V3-INs.

SIGNIFICANCE STATEMENT

Currently, there is a fundamental lack of knowledge about the cellular and spinal network properties that produce locomotor vigor in vertebrates. Here we show, directly for the first time, that V3 interneurons in zebrafish larvae are active during *in vivo* fictive locomotion, and that targeted ablation of the spinal V3 interneuron population reduces the probability of motoneuron firing during fictive swimming. In contrast to V2a interneurons, ablation of V3 interneurons does not affect locomotor frequency, the fictive neural correlate of speed, which clarifies their role in motor control rather than rhythm generation. Thus, we propose that the V3 interneuron subpopulation is a source of excitation in the vertebrate locomotor neural circuitry that regulates locomotor vigor independently of speed.

INTRODUCTION

Efficient locomotion is set by a combination of kinematic properties, including the frequency and vigor of rhythmic motor output. These properties combine to determine various locomotor features, such as distance traveled and speed (Webb, 1975; Jayne and Lauder, 1995; Donley and Dickson, 2000; Tytell, 2004). Locomotor frequency (e.g. limb or tail-beat rhythmic activity) is a well-studied kinematic variable, which has been shown to increase linearly with speed (Bainbridge, 1958; Hunter and Zweifel, 1971; Webb et al., 1984; Müller and van Leeuwen, 2004). Locomotor vigor is defined as the strength of the motor output and measured as the amplitude of limb displacement in limbed animals during locomotor events (walking, trotting, galloping, hopping (Heglund et al., 1974)) or the amplitude of tail-bend in non-limbed animals (fishes) during undulatory swimming (Wardle et al., 1995; Videler et al., 1999). There is a fundamental understanding of how identified subsets of excitatory spinal interneurons help to establish the frequency of motor output (Crone et al., 2009; Zhong et al., 2011; Ausborn et al., 2012; Eklöf-Ljunggren et al., 2012; Kimura et al., 2013; Ampatzis et al., 2014; Ljunggren et al., 2014; Menelaou et al., 2014; Bellardita and Kiehn, 2015). A critical gap in our knowledge, however, is a mechanistic understanding of the cellular and network properties that regulate locomotor vigor in vertebrate animals.

Across vertebrate species from fish to mammals, the neural circuitry that controls locomotion is located in the ventromedial spinal cord (Grillner and Jessell, 2009; Kiehn, 2011). In vertebrates, developmental gene expression

specifies at least five ventral classes of neuronal progenitor cells. These

progenitors give rise to motor neurons (MNs) and four classes of ventral

interneurons, referred to as V0 to V3 neurons (Ericson et al., 1997; Pierani et al.,

1999). Over the past 10 years, a subset of identified excitatory spinal

interneurons (V2a-INs) has been shown to control the speed of locomotion in a

variety of vertebrates (Zhong et al., 2011; Ausborn et al., 2012; Ampatzis et al.,

2014; Gosgnach et al., 2017). A recent study indicated that V2a-INs in adult

zebrafish were involved in both regulating locomotor speed and vigor (Song et

al., 2018). To date, however, a subpopulation of spinal interneurons that

selectively regulates locomotor vigor has not been identified. In the mammalian

spinal cord, a genetically defined subset of pre-motor glutamatergic neurons,

called V3 interneurons (V3-INs), contribute to limb coordination, motor burst

duration, and muscle spasms, but do not have a clear role in locomotor pattern

generation (Chopek et al., 2018; Danner et al., 2019; Lin et al., 2019).

This work shows, directly for the first time, that V3-INs in zebrafish larvae

are active during *in vivo* fictive locomotion and, in part, contribute exclusively to

locomotor vigor and not to locomotor frequency, which clarifies their role in fine

motor control rather than rhythm generation. This is conceptually important

because our results support that the V3-IN subpopulation regulates locomotor

vigor independently of speed. This difference is a major functional distinction

between the V3-INs and V2a-INs since the latter regulate both speed and vigor

(Zhong et al., 2011; Ausborn et al., 2012; Ampatzis et al., 2014; Gosgnach et al.,

2017; Song et al., 2018).

MATERIALS AND METHODS

Fish Lines and Maintenance

All procedures were approved by the Institutional Animal Care and Use
Committee of the University of Minnesota Twin Cities. Wild type (Segrest Farms,
Gibsonton, FL) and transgenic adult zebrafish (*Danio rerio*) were maintained in
the University of Minnesota Zebrafish Core Facility. The transgenic lines used in
these experiments were: *Tg(nkx2.2a:mEGFP)[vu17]* (Kirby et al., 2006),
Tg(vglut2a:DsRed)[nns9] (Miyasaka et al., 2009), *Tg(vglut2a:Gal4ff)[nns20]* (Satou et
al., 2013), *Tg(UAS:GCaMP6s)[nk13a]* (Muto et al., 2017), *Tg(UAS:Kaede)[s1999t]*
(Scott et al., 2007), and *parga[mn2Et]* (Balciunas et al., 2004). In-crossed and out-
crossed transgenic larvae at 4 – 6 days post fertilization (dpf) were used in these
experiments. Larval zebrafish were maintained in petri dishes filled with embryo
water (60 μg/ml Instant Ocean® salt mix, Cincinnati, OH) and 0.00005%
Methylene Blue in a 28.5°C incubator with a 14:10 light:dark cycle.

Kaede Photoconversion

In vivo photoconversion of genetically encoded Kaede was performed in
Tg(vglut2a:Gal4ff)[nns20];Tg(UAS:Kaede)[s1999t] larvae using a fixed-objective, upright
microscope (Olympus FV-1000, Center Valley, PA). Larvae were anesthetized in
zebrafish ringers solution (Wiggin et al., 2012, 2014) with 0.02% Tricaine-S (MS-
222; Western Chemical, Ferndale, WA), embedded laterally in a fluorodish
(World Precision Instruments, Sarasota, FL) filled with 1.5% low-melting point

agarose in zebrafish ringers solution and covered with additional saline

containing anesthetic. Photoconversion of Kaede was performed using the

bleach function of the FV-1000 to direct a 405 µm laser spot onto a Kaede-

expressing cell targeted for conversion (5% laser intensity, 7 s, single pulse). The

embedding procedure resulted in small variations in the orientation of the spinal

cord such that the larvae were not precisely level and square with respect to the

imaging plane. Following confocal stack acquisition, Fiji (Schindelin et al., 2012)

was used to rotate the image in order to correct these errors.

V3-IN Soma Distribution Analysis

Quantification of the distribution of V3-IN cell bodies was performed using

Fiji and custom Matlab code (Mathworks, Natick, MA). Confocal microscopy was

used to collect overlapping stacks of images documenting the expression of

Tg(vglut2a:DsRed)^{nns9} and transmitted light morphology. Images were acquired

at high magnification (40X / NA 0.8 objective) and a z-axis resolution of 1.3 µm.

The fields of view in each stack overlapped with their neighbors by at least 10%

of the width to permit accurate stitching, and included the full width of the spinal

cord and dorsal root ganglia (DRG) in the field of view. The collection of stacks

was stitched into a single 3-D image of the caudal hindbrain and spinal cord

(Preibisch et al., 2009). The Fiji Cell Counter tool was used to identify the

coordinates of all V3 neurons in the rostral 2 mm of the larval spinal cord (out of

a total length of ~3 mm), the coordinates of the DRG neuron clusters, and the

position of the dorsal and ventral edges of the spinal cord. The markers tracing

the dorsal and ventral edges of the spinal cord were used to calculate local

tangent lines to the dorsal and ventral edges of the spinal cord at the

rostrocaudal position of each V3-IN. The equation of a line perpendicular to the

ventral tangent and intersecting the V3-IN was calculated, referred to as the "V3

plumb line." The rostrocaudal distance of each V3 neuron was calculated as the

distance along the ventral surface of the spinal cord between the spinal

cord/hindbrain junction (Segment 2/3 boundary) and the V3 plumb line

intersection with the ventral surface. The dorsoventral height of the spinal cord at

the location of each V3-IN was calculated as the distance between the

intersections of the V3 plumb line with the dorsal surface and the ventral surface,

respectively. The distance of the V3-IN from the ventral surface of the spinal cord

was also measured along the plumb line. The z-axis position of DRG neurons on

each side of the spinal cord were used as boundary points of the lateral edges of

the spinal cord, and a linear fit to the DRG neurons on each side of the animal

was used to calculate the edges of the spinal cord at each rostrocaudal position

in the imaging area. The midline slice and spinal cord width were calculated at

the rostrocaudal position of each V3-IN, along with the distance from midline of

the V3-IN.

Whole-mount Immunohistochemistry

Immunohistochemical labeling was conducted using established

procedures (Montgomery et al., 2016). Five dpf *parga^{mn2Et}* larvae were sacrificed

with 0.2% Tricaine-S and fixed overnight in 4% paraformaldehyde at 4°C. Whole-

mount larvae were permeabilized with Proteinase K, post-fixed, and blocked with

0.2% BSA, 10% normal donkey serum, 0.5% Triton X-100 in PBS. Goat

polyclonal anti-choline acetyltransferase (ChAT; 1:50; AB144P) was added to the

blocking solution and incubated overnight at 4°C. Fixed larvae were incubated

with Alexa Fluor 633 donkey anti-goat secondary antibody (1:500; A-21082,

Thermo Fisher Scientific Inc., Waltham, MA) in secondary blocking solution

(0.2% BSA, 1% normal donkey serum, 0.5% Triton X-100 in PBS) for three

nights at 4°C.

Anti-ChAT labeled *parga^{mn2Et}* larvae were embedded laterally in low

melting-point agarose. An Olympus Fluoview FV1000 confocal microscope was

used to collect images through a 60x objective centered on body segment 15.

Confocal stacks contained the entire dorsoventral extent of the spinal cord and

spanned approximately three body segments (285 μm). Images were processed

in Fiji (open source image processing package) and labeled cells from a single

side of the spinal cord (approximately three hemisegments) were quantified using

the Cell Counter plugin. Photoshop CS5 (Adobe Systems, San Jose, CA) was

used to adjust brightness and contrast across entire z-projected images.

In Vivo Rhod-2 AM Dye Loading of Motor Neurons

For *in vivo* characterization of MNs in *parga^{mn2Et}* larvae, a 0.25% Rhod-2

AM (Life Technologies, Grand Island, NY) stock was made by dissolving 50 μg of

dye in 20 μL Pluronic F-127:DMSO (1:4; Life Technologies, Grand Island, NY).

Next, the skin was removed from the midbody region and a 1:50 dilution of the

Rhod-2 AM stock was bath applied to the exposed muscles to permit

sequestration of indicator dye by MNs via the peripheral motor axons. Sensory

neurons were not strongly loaded by this procedure. Following one-hour

incubation, the Rhod-2 AM solution was thoroughly washed-out using zebrafish

ringers solution. Confocal images were collected and cells were quantified as

described for whole-mount immunohistochemistry (above).

Peripheral Nerve Recordings

218   Peripheral nerve (PN) recordings were performed and analyzed as

previously described (Masino and Fetcho, 2005). In brief, the larvae were

anesthetized, the skin was carefully removed from the recording region, and

larvae were paralyzed with of 0.1 mM α-bungarotoxin (Tocris, Ellisville, MO) to

prevent muscle contractions. The anesthetic was removed to allow the larvae to

perform fictive behavior, and a suction electrode (tip diameter: 9-15 μm) was

placed over the intermyotomal cleft to record MN axons. Fictive motor bursts

were identified using custom Matlab code (Wiggin et al., 2012).

Calcium Imaging and Analysis

228   For calcium imaging of V3-INs,

Tg(vglut2a:Gal4ff)^{nns20};Tg(UAS:GCaMP6s)^{nk13a} zebrafish larvae were used. For

calcium imaging of MNs, Calcium Green-1 AM (Life Technologies, Grand Island,

NY) dye was loaded into MNs using the protocol for Rhod-2 AM dye loading

described above. Larvae were transferred to a fixed-stage Olympus BX51 WI

upright microscope with either a 20x / NA 0.5 or a 40x / NA 0.8 water dipping

objective lens. A field of view containing *GCaMP6s*-expressing V3-INs in

transgenic larvae or Calcium Green-1 AM-loaded MNs in wild type larvae was

selected for calcium imaging and a simultaneous peripheral nerve (PN) recording

was acquired. Calcium indicator fluorescence was excited with an E-Cite Series

120 Q lamp (Excelitas Technologies, Waltham, MA) and acquired at 20 Hz with a

Retiga EXi camera and QCapture Pro 6.0 software (QImaging, Surry, BC). Data

were acquired during spontaneous fictive swimming in 10-60 s durations for both

V3-IN and MN trials. Multiple trials (typically 4) were acquired for each field of

view, with at least five min between trials. Movement artifacts during data

acquisition were corrected in Fiji by registering frames from all trials to a common

reference image (Preibisch et al., 2009). Optical and electrophysiological data

acquisition were synchronized by using the sync output of the stimulator to

trigger acquisition of the PN recording.

For all calcium imaging experiments, raw fluorescence values were

determined by drawing a region of interest around each identified V3-IN or MN

soma and using the Plot Z-axis Profile command in Fiji. Fluorescence from a

region of interest containing background signal was subtracted from raw soma

fluorescence measurements to reduce the effect of photobleaching. Baseline

fluorescence (F_0) was defined as the mean fluorescence in the first 10 frames of

the recording. To calculate $\Delta F/F$, F_0 was subtracted from fluorescence for each

frame ($F_{(t)}$), then divided by F_0; $\Delta F/F = (F_{(t)} - F_{(0)}) / F_{(0)}$. A neuron was considered

active if $\Delta F/F$ was 10% or greater.

Laser Ablation of V3-INs

Acute ablation of the V3-INs was performed using a Micropoint pulsed

nitrogen pumped dye laser (Andor Technology, Belfast, Northern Ireland). Light

(λ = 435 bm) was delivered via an Olympus BX51 WI microscope using a 60x /

NA 1.0 water dipping objective lens. The laser was focused to apply maximal

light intensity to the imaging plane of the microscope, the intensity of the laser

was calibrated to produce a 4.5 – 5.0 µm hole in a mirror slide (this optimal

intensity was determined empirically), and the x-y coordinates of the laser point

in the field of view were indicated on a digital display. Three to six dpf

Tg(vglut2a:DsRed)[nns9] zebrafish larvae were anesthetized in 0.02% Tricaine in

zebrafish ringers solution and embedded laterally in a gel composed of 3%

methylcellulose and 0.02% Tricaine in zebrafish ringers saline. A coverslip was

placed over the gel and secured at the edges with agarose to stabilize the

position of the larvae.

Targeted, bilateral laser ablation of DsRed[+] V3-INs in the ventral spinal

cord was performed by positioning the microscope objective such that a neuron

was in the x-y position at which the laser was focused. Laser pulses were applied

to each V3-IN soma at 33 Hz for 10 s. Laser intensity was empirically determined

a priori such that >95% of the targeted V3-INs were not present 24 h post-

ablation. Following the train of laser pulses, the microscope was repositioned to

focus on another identified V3-IN, and the protocol was repeated. Larvae were

embedded for approximately 50 min in order to perform bilateral ablation of ~140

identified V3-INs between bod segment (S) 5 and S20. To confirm that laser

ablation of targeted V3-INs was effective and persistent, the lack of DsRed

fluorescence in the lesioned spinal cord of *Tg(vglut2a:DsRed)^{nns9}* larvae was

confirmed ~24 hours post-ablation. Sham larvae were treated identically to V3

ablation larvae, but the x-y position of the laser was offset by 5 – 20 µm from

each V3 cell, producing non-specific damage to neurons other than the V3

population.

Statistical Analysis

Statistical analysis of data was performed using two-sample *t*-test,

ANOVA test, Holm-Sidak post-hoc test, Kolmogorov-Smirnov test, and least-

squares linear regression. Statistical tests were carried out using SigmaPlot 12

and 14 software (SyStat Software, San Jose, CA), Matlab (Mathworks, Natick,

MA), or Microsoft Excel (Microsoft, Seattle, WA). An α level of 0.05 was used to

determine statistical significance. Data are reported as means and standard

deviation (Mean, SD).

RESULTS

V3-INs are Individually Identifiable in the Larval Zebrafish Spinal Cord

To characterize the functional roles of V3-INs *in vivo*, it was necessary to

reliably target the neurons for recording and perturbation. While there are no

published transgenic zebrafish lines that specifically label neurons expressing

sim1a (the zebrafish V3-IN marker), it is possible to infer V3-IN identity from their

other characteristics. V3-INs proliferate ventral to the central canal and are

glutamatergic (Zhang et al., 2008; Yang et al., 2010). In zebrafish, there is >90%

overlap of *vesicular glutamate transporter 2* (*vglut2a*) and *sim1a* expression in

the ventral spinal cord at the embryonic stage (1.5 dpf) (Yang et al., 2010).

However, it is possible that the embryonic ventral glutamatergic neurons migrate

or are replaced by a non-V3 population between 1.5 dpf and the free-swimming

larval stage (>3 dpf).

To test the hypothesis that the ventral DsRed+ neurons observed in 4 – 6

dpf *Tg(vglut2a:DsRed)^{nns9}* larvae are V3-INs, we crossed this line to

Tg(nkx2.2a:mEGFP)^{vu17}, which marks p3 progenitor neurons (Briscoe et al.,

2000; Schäfer et al., 2007) (Fig. 1A). At 4 – 6 dpf, *nkx2.2a:mEGFP* expression

was restricted to the region ventral to the central canal (Fig. 1C). The p3 domain

of the zebrafish spinal cord also produces Kolmer-Agduhr (KA) neurons (Yang et

al., 2010), and likely other neuronal types, so it was expected that there would be

more *nkx2.2a:mEGFP*-expressing neurons than *vglut2a:DsRed*-expressing

neurons (Fig. 1B). Approximately 94% of *vglut2a:DsRed*-expressing neurons

ventral to the central canal also expressed *nkx2.2a:mEGFP* (*n* = 6 larvae, 162

neurons), and were therefore confirmed to be V3-INs (Fig. 1D). There was not a

significant difference between the dorsoventral position of the *vglut2a:DsRed-*

expressing neurons that co-expressed *nkx2.2a:mEGFP* and those that did not

(two-sample t-test; *t* = 1.6; *p* = 0.10), so there was not a way to reliably segregate

the identified V3-INs, from those that did not express *nkx2.2a:mEGFP*. However,

the overwhelming majority of the DsRed^{+} neurons ventral to the central canal

were confirmed to be V3-INs. The lack of confirmed transgene overlap in the

remaining cells may be due to variegated transgene expression rather than the

presence of a non-V3-IN ventral glutamatergic neuronal population. Based on

this experiment, we treat the ventral *vglut2a:DsRed*-expressing neurons as the

V3-IN population, with the caveat that a small fraction (~6%) may be

misidentified.

V3-INs are Located in the Ventromedial Spinal Cord and are Distributed Along

the Rostrocaudal Axis of the Spinal Cord

In mice, the V3-INs migrate following differentiation to form spatially

segregated dorsal and ventral populations that have different cellular properties

and may be necessary for different behaviors (Borowska et al., 2013). To

determine if there are multiple populations of spatially segregated V3-INs in

zebrafish larvae, we used confocal microscopy to create 3-D reconstructions of

the spinal cords of 4 – 6 dpf *Tg(vglut2a:DsRed)^{nns9}* zebrafish larvae (Fig. 2A-B).

The center of each V3-IN soma in the most rostral 2 mm of the spinal cord was

tagged, and the distribution of the neurons within the spinal cord was calculated

based on the imputed borders of the spinal cord (see *Methods*). Neurons in the

far caudal spinal cord were not counted because the dimensions of the spinal

cord taper to the extent that it was difficult to determine the location of the central

canal and Mauthner axons. All of the neurons (*n* = 4 larvae; 797 neurons) were

located medial of the Mauthner axon (Fig. 2C). V3-IN density was highest in the

rostral spinal cord but remained close to 1 cell per 10 μm throughout the cord

(Fig. 2D). The V3-IN somas were approximately 5 µm in diameter in the

rostrocaudal axis, so a density of ~1 cell per 10 µm thoroughly tiled the ventral

spinal cord. The majority of V3-INs were located in the most medial and most

ventral 25% of the spinal cord (Figs. 2E & F). There was no parasagittal plane

through the medial spinal cord that was unoccupied by V3-INs (Fig. 2E), which

suggested that the V3 population may be considered a midline population.

However, the density of neurons was not highest at the sagittal midline of the

spinal cord, and when cell density was plotted along the population axis of the

V3-INs, the point of highest density was adjacent to the spinal cord midline (Fig.

2G). Therefore, we propose that distinct left and right populations of V3-INs are

present, but that there are not multiple ipsilateral populations of spatially

segregated V3-INs in larval zebrafish.

V3-INs Project Bilaterally and Have Descending Processes

To characterize the morphologies and putative synaptic connectivity of the

V3-INs, we examined the 3-D morphology of individual V3-INs using confocal

microscopy (Fig. 3). The density of DsRed fluorescence is too high in the

Tg(vglut2a:DsRed)^{nns9} transgenic line to accurately trace the processes of a

single neuron, so we expressed a photoconvertible protein (Kaede) in

Tg(vglut2a:Gal4)^{nns20};Tg(UAS:Kaede)^{s1999t} larvae to identify and photoconvert

single V3-INs (Fig. 3A-B). Following photoconversion, it was possible to obtain

morphological images of the V3-INs (*n* = 5 neurons, 4 larvae; Fig. 3C). Based on

this sample of V3 morphologies, it was evident that variation in the projection

patterns of the V3-INs was present. For example, the V3-IN imaged in S10 had

minimal projection density dorsal or rostral to the cell body, while the V3-IN in

S18 predominantly projected rostrally and dorsally (Fig. 3C, Lateral z-Projection).

Despite these differences, there were some common features of this sample of

V3-IN morphologies. First, the imaged V3-INs possessed bilateral projections,

but had a higher density of processes ipsilateral to the cell body (Fig. 3C,

Horizontal z-Projection). Second, V3-INs had the greatest density of processes

close to the cell body and longer caudal projections than rostral projections (Fig.

3C). Third, while the full extent of the descending processes of the S9 and S14

neurons is not presented in Fig. 3C, V3-INs had a maximum detectable

projection distance shorter than 300 µm. The limited projection length we

observed may be due to the Kaede photoconversion method used and should be

considered a lower bound on the full extent of the projections of the V3-INs. The

projection pattern of the V3-INs appeared to innervate the ventral spinal cord,

which is populated by MNs, V2a-INs (Eklöf-Ljunggren et al., 2012; Menelaou and

McLean, 2013), other V3-INs, KA neurons, and serotonergic neurons (McLean

and Fetcho, 2004; Yang et al., 2010).

The Majority of V3-INs are Active During Spontaneous Fictive Swimming

In mice, there is only indirect evidence supporting temporal correlation

between V3-IN activity and locomotion (Borowska et al., 2013). To test the

hypothesis that activity in the V3-INs is appropriately timed to provide excitatory

drive to MNs during locomotion, we monitored the activity of the V3-INs during

spontaneous fictive swimming using a genetically-encoded calcium indicator

(GCaMP6s; Fig. 4). The activity of V3-INs and fictive motor output during

spontaneous swimming was measured via simultaneous calcium imaging and

peripheral nerve recording using 4 - 6 dpf

Tg(vglut2a:Gal4ff)^{nns20};Tg(UAS:GCaMP6s)^{nk13a} larvae (Fig. 4A-B). We found that

the majority, 86 of 93 (92%), of V3-INs were active ($\Delta F/F > 10\%$) during

spontaneous fictive swimming episodes (n = 6 larvae, 15 trials). Therefore, this is

the first direct evidence that V3-INs are active during fictive locomotion, which

supports the participation of V3-INs in producing locomotor activity.

Ablation of V3-INs Does Not Affect Episodic or Burst Properties of Spontaneous

Fictive Swimming

The previous experiments demonstrated that the V3-INs have

morphology, neurotransmitter phenotype, and activity consistent with providing

excitatory drive to motor neurons but did not demonstrate the function of the V3

neurons or their necessity for behavior. To determine whether canonical features

of fictive swimming (episode duration, burst (tail-beat) frequency, and burst

duration) are modulated by V3-IN activity, we performed laser ablation of a large

percentage of the V3-IN population (>95%; ~140 V3-identified somas across

body segments 5 to 20; Fig. 5). Three experimental groups were used: 1) The V3

ablation group, in which all V3-INs on both sides of the spinal cord between S5

and S20 were laser ablated (n = 6 larvae; 145 (SD 9.6) neurons per larva).

Representative examples of targeted laser ablation efficacy within a restricted

region (Fig. 5A-B) and spatial localization (Fig. 5C-D) are shown. 2) The sham

ablation group, in which laser pulses were delivered 5 – 20 µm rostral to each

identified V3-IN cell body (n = 6 larvae; 130 (SD 10.8) sham targets per larva). 3)

The unmanipulated control group was not embedded or exposed to laser pulses

(n = 6 larvae).

The day following laser ablation or sham ablation, fictive swimming was

measured using PN recordings. Episode duration, burst frequency, and burst

duration were not significantly different between unmanipulated, sham and

ablation experimental groups during spontaneous swimming (one-way ANOVAs;

all F < 1.3; all p > 0.30; Table 1). These results indicate that the population of

spinal V3-INs do not play a significant role in establishing episode duration, burst

frequency (a proxy for tail-beat frequency), or burst duration during fictive

locomotion in larval zebrafish.

Ablation of V3-INs Reduces the Proportion of Active Motor Neurons but Does Not

Affect Peak Locomotor Frequency During Spontaneous Fictive Swimming

Since the ablation of spinal V3-INs in zebrafish larvae did not affect

episode duration, burst frequency, or burst duration (Table 1), we tested the

hypothesis that ablation of V3-INs reduces locomotor vigor by decreasing the

number of active MNs during spontaneous fictive swimming. In order to ablate

V3-INs without also photo-damaging or bleaching motor neurons, we used a two-

step neuronal targeting strategy. First, V3-INs were identified and ablated using

the *Tg(vglut2a:DsRed)* [nns9] transgenic line. Following recovery from laser

ablation, motor neurons were loaded with a cell-permanent calcium indicator,

Calcium Green-1, AM. To verify that spinal neurons loaded with AM dye (Calcium

Green-1 and/or Rhod-2) were MNs, we first confirmed motor neuronal specificity

of GFP expression in the *parga*[*mn2Et*] enhancer trap line (Balciunas et al., 2004).

Parga[*mn2Et*] larvae immunolabeled for the cholinergic marker choline

acetyltransferase (ChAT) revealed that the majority of GFP-expressing neurons

co-labeled with ChAT antibodies (*n* = 5; 91.3% (SD2.1); Fig. 6A-A''). A small

percentage of dorsally-positioned ChAT+ cells did not colocalize with GFP (4.1%

(SD2.8); Fig. 6A-A'', arrowheads); these cells were considered members of a

subset of cholinergic spinal interneurons that modulate MN excitability recently

identified in zebrafish (Bertuzzi and Ampatzis, 2018). Thus, GFP expression in

the *parga*[*mn2Et*] enhancer trap line was specific to MNs.

Next, to confirm that loading of AM dye was restricted to MNs, the skin

was removed from both sides of the body from *parga*[*mn2Et*] larvae, in which GFP

expression was shown to be limited to MNs (Fig. 6A-A''), and larvae were

incubated in zebrafish ringers solution containing Rhod-2 AM dye (*n* = 4; Fig. 6B-

B''). Rhod2-AM, which was not used for calcium imaging experiments because it

did not produce a strong calcium signal, was used as a proxy for Calcium Green-

1 AM since the excitation/emission wavelengths of Calcium Green-1 AM

overlapped with, and could not be disambiguated from, GFP expressed in

parga[*mn2Et*] MNs. The majority of GFP-expressing MNs co-labeled with Rhod-2

AM (84.3% (SD4.3); Fig. 6B-B''). A small percentage of GFP-negative cells

contained Rhod-2 AM (5.4% (SD0.5); Fig. 6B-B'', arrowheads). Taken together,

these results confirmed that MNs were selectively labeled with AM dye and,

therefore, the cell permeable calcium indicator (Calcium Green-1 AM) could be

used to monitor *in vivo* calcium activity in spinal MNs.

To examine the effects of V3-IN ablation on the production of fictive

locomotor vigor, we determined the proportion of spinal MNs that were active

during spontaneous fictive swimming episodes. MN activity in

Tg(vglut2a:DsRed)nns9 larvae was measured using *in vivo* calcium imaging by

loading MNs with Calcium Green-1 AM, while peak burst frequency during

spontaneous fictive locomotor activity was measured using extracellular PN

recordings (Fig. 7A-B). We quantified the proportion of active MNs and peak

burst frequencies during spontaneous fictive swimming for Control (50 swim

episodes, 4 larvae), Sham Ablated (14 swim episodes, 2 larvae), and V3-IN

Ablated (59 swim episodes, 3 larvae) groups. Importantly, there were no

differences in the properties of the fictive swimming produced by these larvae,

consistent with our prior experiments (Table 1). Neither the distribution of peak

burst frequencies (Kolmogorov-Smirnov test < 0.33, $p > 0.14$ for all group

comparisons; Fig. 7C), nor the mean peak burst frequencies, were different

between Control, Sham, and Ablation Groups (Control = 39.0 Hz (SD8.5); Sham

= 36.4 Hz (SD10.7); Ablated = 39.3 Hz (SD8.9); one-way ANOVA, F = 5.0, p =

0.08; Fig. 7D).

Assessment of the probability distributions showed that the proportion of

active MNs in the Ablated group was significantly reduced compared to both the

Control (Kolmogorov-Smirnov test = 0.68, $p < 0.001$) and Sham (Kolmogorov-

Smirnov test = 0.54, p = 0.001) groups (Fig. 7E). The proportion of active MNs

was not significantly different between the Control and Sham groups

(Kolmogorov-Smirnov test = 0.24, p = 0.50; Fig. 7E). Further, V3-IN ablation had

a main effect of significantly reducing the proportion of active MNs during

spontaneous fictive swimming (Fig. 7F; Control mean = 65.2% (SD16.7); Sham

mean = 56.6% (SD24.2); Ablated mean = 41.2% (SD15.6); one-way ANOVA, F =

25.3; p < 0.001). Pairwise multiple comparisons (Holm-Sidak; Fig. 7D) revealed

significant differences for the proportion of active MNs between both the Control

and Ablated groups (t = 7.02, p < 0.001) and the Sham and Ablated groups (t =

2.98, p = 0.007), but significant differences were not found between Control and

Sham groups (t = 1.54, p = 0.10). These results indicate that V3-INs participate in

regulating locomotor vigor by recruitment of spinal MNs but do not regulate burst

frequency during spontaneous fictive locomotion.

DISCUSSION

The goal of this study was to characterize the potential role of V3-INs in

producing locomotor vigor in genetically tractable and optically accessible larval

zebrafish. Locomotor vigor, the strength of a locomotor output, is regulated by

the recruitment of motor units appropriate to the desired movement: small units

for fine behaviors and large units for large movements (Zajac and Faden, 1985;

Mendell, 2005). This recruitment pattern of small to large motor units, the so-

called size principle, is well established in both mammalian and swimming

locomotor systems (Henneman et al., 1965; Gustafsson and Pinter, 1984a,

1984b; Cope and Pinter, 1995; McLean et al., 2007; Gabriel et al., 2011).

However, investigations into the pre-motor neural mechanisms that contribute to

the regulation of locomotor vigor have been difficult in mammals for two primary

reasons. First, mammalian spinal locomotor circuits are complicated by the

requirements to control and coordinate limbs and joints, rather than simply

produce a smooth gradient of motion intensity (Grillner, 2003; Grillner and El

Manira, 2020). Second, despite recent progress in tool development, *in vivo*

functional manipulations of mammalian spinal locomotor circuits using mini-

scopes and fiber-optics are both challenging and somewhat limited in use

(Montgomery et al., 2015; Samineni et al., 2017; Canales et al., 2018). Zebrafish

larvae provide several advantages over mammalian models for *in vivo* studies:

the spinal cords of fish are less complicated than mammals, zebrafish are

recognized as a powerful neurogenetic model system, and the translucent nature

of zebrafish larvae makes them particularly appropriate for *in vivo* optical

methods of investigation. In this study, we found that, unlike the mammalian

spinal circuit, V3-INs in the larval zebrafish spinal cord are contained within a

single, ventral anatomical domain (Fig. 1-2) and are not necessary to fictive

locomotor rhythm (Table 1) (Zhang et al., 2008; Danner et al., 2019). Instead,

zebrafish V3-INs are swim-active neurons that are necessary for recruitment of

motor neurons during fictive locomotion (Fig. 4, 7), and therefore are most likely

necessary for the recruitment of motor units during free-moving locomotion.

V3-INs are a population of previously uncharacterized ventral VGlut2a+ neurons

in larval zebrafish

Here, we report that almost all ventrally-located neurons labeled by the

Tg(vglut2a:DsRed)[nns9] and *Tg(vglut2a:Gal4ff)[nns20]* transgenic lines are V3-IN cells

(Figs. 1-5). The V3-INs, a numerous ventromedial population, are distributed

across the rostrocaudal axis of the spinal cord (Fig. 2) and project to both sides

of the spinal cord, with a richer innervation of the ipsilateral side of the spinal

cord (Fig. 3). In this dataset, there are several different V3-IN projection patterns

(Fig. 3), but none of these anatomical projection patterns resemble previously

characterized ventral neuron cell types, such Kolmer-Agduhr cells (Dale et al.,

1987; Bernhardt et al., 1992; Montgomery et al., 2016), ventral serotonergic

neurons (Van Raamsdonk et al., 1996; Brustein et al., 2003; McLean and Fetcho,

2004; Montgomery et al., 2018), or VeMes (Hale et al., 2001). V3-INs are a

morphologically distinct cell type or collection of cell types that have not been

characterized through previous experimental approaches.

The observed diversity of cell morphology in the V3-IN cell population is

similar to the diversity of V3-INs in the mammalian spinal cord (Borowska et al.,

2013; Deska-Gauthier et al., 2020). In mammals, the morphology of V3-IN

subtypes is correlated with their circuit function (Borowska et al., 2013).

However, we have not found evidence for differential recruitment of V3-INs

during fictive swimming or functional heterogeneity. It may be that larval

zebrafish V3-INs are a primitive class of excitatory spinal INs, as has been

previously proposed for *Engrailed1*-expressing inhibitory neurons in zebrafish

(Higashijima et al., 2004). However, analysis of cell morphology in zebrafish

larvae is complicated by ongoing development in the nervous system and the

rostrocaudal developmental gradient (Cole and Ross, 2001). Further experiments

are necessary to clarify if these morphologies represent distinct sub-types of V3-

INs, developmental stages of a single morphological family, or regional

specializations of the V3-INs.

Zebrafish V3-INs are active during fictive locomotion

Results indicating that V3-IN activity is directly correlated to fictive

locomotor activity in vertebrate preparations has not been convincingly

presented. In mice, there is indirect evidence indicating temporal correlation

between V3-IN activity and locomotion (Borowska et al., 2013). Through

simultaneous optical and electrophysiological recordings, we showed that activity

in the majority of V3-INs is correlated to fictive swimming (Fig. 4) and is therefore

appropriately timed to provide excitatory drive to MNs during locomotion. Thus,

this is the first direct evidence in any vertebrate preparation that V3-INs are

active during fictive locomotion, which provides supporting evidence that they

may participate in locomotor activity.

V3-IN activity is necessary for motor neuron recruitment, but not necessary for

rhythm generation

Given that V3-INs have been shown to affect both the regularity and

coordination of locomotor activity in mice (Zhang et al., 2008), it was surprising

that laser ablation of V3-INs in larval zebrafish had no effect on episode duration,

mean burst frequency, or burst duration during fictive swimming in larval

zebrafish (Table 1). One interpretation of this finding is that despite similarities in

gene expression and neurotransmitter profile, pre-motor interneuron function is

divergent between axial and tetrapod locomotion. The latter argument has been

advanced regarding the V2a interneurons (Dougherty and Kiehn, 2010;

Dougherty et al., 2013); the different effects (or lack thereof) of V3-IN ablation in

zebrafish and mice may be a second example of the trend. An important

difference between the current study and mouse knock-out experiments is the

mechanism of ablation. In the current study, V3-INs were ablated acutely, and

little time (~24 hours) was given for circuit compensation. In the mouse

experiments, on the other hand, V3-INs were ablated or silenced chronically

(Zhang et al., 2008; Rabe Bernhardt et al., 2012), leaving opportunity for

physiological compensation. Acute silencing experiments using optogenetics

could be performed in both species (Arrenberg et al., 2009; Hägglund et al.,

2013), and may demonstrate that this difference between species is due to the

experimental manipulation rather than the underlying neural circuit.

Over the last several years, a subset of identified excitatory spinal

interneurons, called V2a interneurons, has been shown to control the speed of

locomotion in a variety of vertebrates (Zhong et al., 2011; Ausborn et al., 2012;

Ampatzis et al., 2014; Gosgnach et al., 2017). Of particular interest, a recent

study also ascribed a role for regulating locomotor vigor to V2a-INs in adult

zebrafish (Song et al., 2018). Taken together, these results indicate that V2a-INs

are involved in regulating both locomotor speed and vigor. To date, however, a

subpopulation of spinal interneurons that selectively regulate locomotor vigor has

not been identified. Our work provides evidence that, in zebrafish larvae, V3-INs

contribute exclusively to locomotor vigor. Specifically, we show that targeted

ablation of V3-INs (Fig. 5) reduces the probability of MN firing during fictive

swimming (Fig. 7E-F), but does not affect the range of locomotor frequencies

produced (Fig. 7C-D). This suggests that vigor, but not frequency, may be

determined, in part, by the proportion of active V3-INs. Thus, we propose that

V3-INs are a source of excitation in the vertebrate locomotor neural circuitry that

regulate the vigor of locomotor output.

Conclusion

The work presented here addressed an important concept regarding

motor control; specifically, a fundamental lack of understanding of the cellular

properties that produce locomotor vigor in vertebrates. We showed, directly for

the first time, that V3-INs in zebrafish larvae are active during *in vivo* fictive

locomotion and contribute, at least in part, exclusively to locomotor vigor. Further,

targeted ablation of the V3-IN spinal cord population does not affect locomotor

frequency (speed), which clarifies their role in fine motor control rather than

rhythm generation. Thus, we propose that the V3-IN subpopulation regulates

locomotor vigor independently of speed, which is a functional difference between

the V3-INs and V2a-INs since the latter regulate both speed and vigor (Zhong et

al., 2011; Ausborn et al., 2012; Ampatzis et al., 2014; Gosgnach et al., 2017;

Song et al., 2018).

REFERENCES

Ampatzis K, Song J, Ausborn J, El Manira A (2014) Separate microcircuit
modules of distinct v2a interneurons and motoneurons control the speed of
locomotion. Neuron 83:934–943 Available at:
http://linkinghub.elsevier.com/retrieve/pii/S0896627314006291 [Accessed
October 19, 2017].

Arrenberg AB, Del Bene F, Baier H (2009) Optical control of zebrafish behavior
with halorhodopsin. Proc Natl Acad Sci U S A 106:17968–17973 Available
at: http://www.pnas.org/cgi/doi/10.1073/pnas.0906252106 [Accessed
October 23, 2017].

Ausborn J, Mahmood R, El Manira A (2012) Decoding the rules of recruitment of
excitatory interneurons in the adult zebrafish locomotor network. Proc Natl
Acad Sci U S A 109:E3631-9 Available at:
http://www.pnas.org/cgi/doi/10.1073/pnas.1216256110 [Accessed October
19, 2017].

Bainbridge R (1958) The Speed of Swimming of Fish as Related to Size and to
the Frequency and Amplitude of the Tail Beat. J Exp Biol 35.

Balciunas D, Davidson AE, Sivasubbu S, Hermanson SB, Welle Z, Ekker SC
(2004) Enhancer trapping in zebrafish using the Sleeping Beauty
transposon. BMC Genomics 5.

Bellardita C, Kiehn O (2015) Phenotypic Characterization of Speed-Associated
Gait Changes in Mice Reveals Modular Organization of Locomotor
Networks. Curr Biol 25:1426–1436 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/25959968 [Accessed October 23,
2017].

Bernhardt RR, Patel CK, Wilson SW, Kuwada JY (1992) Axonal trajectories and
distribution of GABAergic spinal neurons in wildtype and mutant zebrafish
lacking floor plate cells. J Comp Neurol 326:263–272.

Bertuzzi M, Ampatzis K (2018) Spinal cholinergic interneurons differentially
control motoneuron excitability and alter the locomotor network operational
range. Sci Rep 8.

Borowska J, Jones CT, Zhang H, Blacklaws J, Goulding M, Zhang Y (2013)
Functional subpopulations of V3 interneurons in the mature mouse spinal
cord. J Neurosci 33:18553–18565 Available at:
http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.2005-13.2013
[Accessed October 23, 2017].

Briscoe J, Pierani A, Jessell TM, Ericson J (2000) A homeodomain protein code
specifies progenitor cell identity and neuronal fate in the ventral neural tube.
Cell 101:435–445 Available at:

http://www.ncbi.nlm.nih.gov/pubmed/10830170 [Accessed October 22, 2017].

Brustein E, Chong M, Holmqvist B, Drapeau P (2003) Serotonin Patterns Locomotor Network Activity in the Developing Zebrafish by Modulating Quiescent Periods. J Neurobiol.

Canales A, Park S, Kilias A, Anikeeva P (2018) Multifunctional Fibers as Tools for Neuroscience and Neuroengineering. Acc Chem Res 51.

Chopek JW, Nascimento F, Beato M, Brownstone RM, Zhang Y (2018) Sub-populations of Spinal V3 Interneurons Form Focal Modules of Layered Pre-motor Microcircuits. Cell Rep 25.

Cole LK, Ross LS (2001) Apoptosis in the developing zebrafish embryo. Dev Biol 240:123–142.

Cope T, Pinter M (1995) The Size Principle: Still Working After All These Years. Physiology 10:280–286.

Crone SA, Zhong G, Harris-Warrick R, Sharma K (2009) In mice lacking V2a interneurons, gait depends on speed of locomotion. J Neurosci 29:7098–7109 Available at: http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.1206-09.2009 [Accessed October 23, 2017].

Dale N, Roberts A, Ottersen OP, Storm-Mathisen J (1987) The morphology and distribution of "Kolmer-Agduhr cells", a class of cerebrospinal-fluid-contacting neurons revealed in the frog embryo spinal cord by GABA immunocytochemistry. Proc R Soc Lond B Biol Sci.

Danner SM, Zhang H, Shevtsova NA, Borowska-Fielding J, Deska-Gauthier D, Rybak IA, Zhang Y (2019) Spinal V3 Interneurons and Left–Right Coordination in Mammalian Locomotion. Front Cell Neurosci.

Deska-Gauthier D, Borowska-Fielding J, Jones CT, Zhang Y (2020) The temporal neurogenesis patterning of spinal p3–V3 interneurons into divergent subpopulation assemblies. J Neurosci 40.

Donley JM, Dickson KA (2000) Swimming kinematics of juvenile kawakawa tuna (Euthynnus affinis) and chub mackerel (Scomber japonicus). J Exp Biol 203:3103–3116 Available at: http://www.ncbi.nlm.nih.gov/pubmed/11003821 [Accessed October 23, 2017].

Dougherty KJ, Kiehn O (2010) Firing and Cellular Properties of V2a Interneurons in the Rodent Spinal Cord. J Neurosci 30:24–37 Available at: http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.4821-09.2010.

Dougherty KJ, Zagoraiou L, Satoh D, Rozani I, Doobar S, Arber S, Jessell TM, Kiehn O (2013) Locomotor Rhythm Generation Linked to the Output of Spinal Shox2 Excitatory Interneurons. Neuron 80:920–933.

Eklöf-Ljunggren E, Haupt S, Ausborn J, Dehnisch I, Uhlén P, Higashijima S, El
Manira A (2012) Origin of excitation underlying locomotion in the spinal
circuit of zebrafish. Proc Natl Acad Sci U S A 109:5511–5516 Available at:
http://www.pnas.org/cgi/doi/10.1073/pnas.1115377109 [Accessed October
19, 2017].

Ericson J, Rashbass P, Schedl A, Brenner-Morton S, Kawakami A, van
Heyningen V, Jessell TM, Briscoe J (1997) Pax6 controls progenitor cell
identity and neuronal fate in response to graded Shh signaling. Cell 90:169–
180 Available at: http://www.ncbi.nlm.nih.gov/pubmed/9230312 [Accessed
October 22, 2017].

Gabriel JP, Ausborn J, Ampatzis K, Mahmood R, Eklöf-Ljunggren E, El Manira A
(2011) Principles governing recruitment of motoneurons during swimming in
zebrafish. Nat Neurosci 14:93–99 Available at:
http://www.nature.com/doifinder/10.1038/nn.2704 [Accessed October 19,
2017].

Gosgnach S, Bikoff JB, Dougherty KJ, El Manira A, Lanuza GM, Zhang Y (2017)
Delineating the diversity of spinal interneurons in locomotor circuits. J
Neurosci 37.

Grillner S (2003) The motor infrastructure: From ion channels to neuronal
networks. Nat Rev Neurosci 4.

Grillner S, El Manira A (2020) Current principles of motor control, with special
reference to vertebrate locomotion. Physiol Rev 100.

Grillner S, Jessell TM (2009) Measured motion: searching for simplicity in spinal
locomotor networks. Curr Opin Neurobiol 19:572–586 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/19896834 [Accessed October 20,
2017].

Gustafsson B, Pinter MJ (1984a) An investigation of threshold properties among
cat spinal alpha-motoneurones. J Physiol 357:453–483.

Gustafsson B, Pinter MJ (1984b) Relations among passive electrical properties
of lumbar alpha-motoneurones of the cat. J Physiol 356:401–431.

Hägglund M, Dougherty KJ, Borgius L, Itohara S, Iwasato T, Kiehn O (2013)
Optogenetic dissection reveals multiple rhythmogenic modules underlying
locomotion. Proc Natl Acad Sci U S A 110:11589–11594.

Hale ME, Ritter DA, Fetcho JR (2001) A confocal study of spinal interneurons in
living larval zebrafish. J Comp Neurol 437:1–16 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/11477593 [Accessed October 19,
2017].

Heglund NC, Taylor CR, Mcmahon TA (1974) Scaling stride frequency and gait
to animal size: Mice to horses. Science (80-) 186.

HENNEMAN E, SOMJEN G, CARPENTER DO (1965) Functional Significance of
Cell Size in Spinal Motoneurons. J Neurophysiol 28:560–580.

Higashijima S, Masino MA, Mandel G, Fetcho JR (2004) Engrailed-1 expression
marks a primitive class of inhibitory spinal interneuron. J Neurosci 24:5827–
5839 Available at: http://www.ncbi.nlm.nih.gov/pubmed/15215305 [Accessed
October 19, 2017].

Hunter JR, Zweifel JR (1971) Swimming speed, tail beat frequency, tail beat
amplitude, and size in jack makerel, Trachurus symmetricus, and other
fishes. Fish Bull 69:253–267 Available at:
http://swfsc.noaa.gov/publications/CR/1971/7114.PDF [Accessed October
23, 2017].

Jayne BC, Lauder GV (1995) Speed effects on midline kinematics during steady
undulatory swimming of largemouth bass, Micropterus salmoides. J Exp Biol
198:585–602 Available at: http://www.ncbi.nlm.nih.gov/pubmed/9318295
[Accessed October 23, 2017].

Kiehn O (2011) Development and functional organization of spinal locomotor
circuits. Curr Opin Neurobiol 21:100–109 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/20889331 [Accessed October 20,
2017].

Kimura Y, Satou C, Fujioka S, Shoji W, Umeda K, Ishizuka T, Yawo H,
Higashijima S (2013) Hindbrain V2a Neurons in the Excitation of Spinal
Locomotor Circuits during Zebrafish Swimming. Curr Biol 23:843–849
Available at: http://www.ncbi.nlm.nih.gov/pubmed/23623549 [Accessed
October 20, 2017].

Kirby BB, Takada N, Latimer AJ, Shin J, Carney TJ, Kelsh RN, Appel B (2006) In
vivo time-lapse imaging shows dynamic oligodendrocyte progenitor behavior
during zebrafish development. Nat Neurosci.

Lin S, Li Y, Lucas-Osma AM, Hari K, Stephens MJ, Singla R, Heckman CJ,
Zhang Y, Fouad K, Fenrich KK, Bennett DJ (2019) Locomotor-related V3
interneurons initiate and coordinate muscles spasms after spinal cord injury.
J Neurophysiol.

Ljunggren EE, Haupt S, Ausborn J, Ampatzis K, El Manira A (2014) Optogenetic
activation of excitatory premotor interneurons is sufficient to generate
coordinated locomotor activity in larval zebrafish. J Neurosci 34:134–139
Available at: http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.4087-
13.2014 [Accessed October 19, 2017].

Masino MA, Fetcho JR (2005) Fictive swimming motor patterns in wild type and
mutant larval zebrafish. J Neurophysiol 93:3177–3188 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/15673549 [Accessed October 19,
2017].

McLean DL, Fan J, Higashijima S, Hale ME, Fetcho JR (2007) A topographic
map of recruitment in spinal cord. Nature 446:71–75 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/17330042 [Accessed October 19,
2017].

McLean DL, Fetcho JR (2004) Ontogeny and innervation patterns of
dopaminergic, noradrenergic, and serotonergic neurons in larval zebrafish. J
Comp Neurol 480:38–56 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/15515022 [Accessed October 19,
2017].

Mendell LM (2005) The size principle: A rule describing the recruitment of
motoneurons. J Neurophysiol 93:3024–3026.

Menelaou E, McLean DL (2013) Speed control: spinal interneurons with crossed
purposes. Curr Biol 23:R716-8 Available at:
http://linkinghub.elsevier.com/retrieve/pii/S0960982213009251 [Accessed
October 19, 2017].

Menelaou E, VanDunk C, McLean DL (2014) Differences in the morphology of
spinal V2a neurons reflect their recruitment order during swimming in larval
zebrafish. J Comp Neurol 522:1232–1248 Available at:
http://doi.wiley.com/10.1002/cne.23465 [Accessed October 19, 2017].

Miyasaka N, Morimoto K, Tsubokawa T, Higashijima S, Okamoto H, Yoshihara Y
(2009) From the olfactory bulb to higher brain centers: genetic visualization
of secondary olfactory pathways in zebrafish. J Neurosci 29:4756–4767
Available at: http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.0118-
09.2009 [Accessed October 20, 2017].

Montgomery JE, Wahlstrom-Helgren S, Wiggin TD, Corwin BM, Lillesaar C,
Masino MA (2018) Intraspinal serotonergic signaling suppresses locomotor
activity in larval zebrafish. Dev Neurobiol.

Montgomery JE, Wiggin TD, Rivera-Perez LM, Lillesaar C, Masino MA (2016)
Intraspinal serotonergic neurons consist of two, temporally distinct
populations in developing zebrafish. Dev Neurobiol 76:673–687 Available at:
http://doi.wiley.com/10.1002/dneu.22352 [Accessed October 19, 2017].

Montgomery KL, Yeh AJ, Ho JS, Tsao V, Iyer SM, Grosenick L, Ferenczi EA,
Tanabe Y, Deisseroth K, Delp SL, Poon ASY (2015) Wirelessly powered,
fully internal optogenetics for brain, spinal and peripheral circuits in mice.
Nat Methods 12.

Müller UK, van Leeuwen JL (2004) Swimming of larval zebrafish: ontogeny of
body waves and implications for locomotory development. J Exp Biol
207:853–868 Available at: http://www.ncbi.nlm.nih.gov/pubmed/14747416
[Accessed October 22, 2017].

Muto A, Lal P, Ailani D, Abe G, Itoh M, Kawakami K (2017) Activation of the

hypothalamic feeding centre upon visual prey detection. Nat Commun.

Pierani A, Brenner-Morton S, Chiang C, Jessell TM (1999) A sonic hedgehog-
independent, retinoid-activated pathway of neurogenesis in the ventral spinal
cord. Cell 97:903–915 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/10399918 [Accessed October 22,
2017].

Preibisch S, Saalfeld S, Tomancak P (2009) Globally optimal stitching of tiled 3D
microscopic image acquisitions. Bioinformatics.

Rabe Bernhardt N, Memic F, Gezelius H, Thiebes A-L, Vallstedt A, Kullander K
(2012) DCC mediated axon guidance of spinal interneurons is essential for
normal locomotor central pattern generator function. Dev Biol 366:279–289
Available at: http://linkinghub.elsevier.com/retrieve/pii/S0012160612001807
[Accessed October 20, 2017].

Samineni VK et al. (2017) Fully implantable, battery-free wireless optoelectronic
devices for spinal optogenetics. Pain 158.

Satou C, Kimura Y, Hirata H, Suster ML, Kawakami K, Higashijima S -i. (2013)
Transgenic tools to characterize neuronal properties of discrete populations
of zebrafish neurons. Development 140:3927–3931 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/23946442 [Accessed October 20,
2017].

Schäfer M, Kinzel D, Winkler C (2007) Discontinuous organization and
specification of the lateral floor plate in zebrafish. Dev Biol 301:117–129
Available at: http://linkinghub.elsevier.com/retrieve/pii/S001216060601219X
[Accessed October 25, 2017].

Schindelin J, Arg I, Arganda-Carreras I, a-Carreras, Frise E, Kaynig V, Longair
M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y,
White D, Hartenstein V, Eliceiri K, Tomancak P, Cardona A (2012) Fiji: an
open-source platform for biological-image analysis. Nat Methods 9.

Scott EK, Mason L, Arrenberg AB, Ziv L, Gosse NJ, Xiao T, Chi NC, Asakawa K,
Kawakami K, Baier H (2007) Targeting neural circuitry in zebrafish using
GAL4 enhancer trapping. Nat Methods.

Song J, Dahlberg E, El Manira A (2018) V2a interneuron diversity tailors spinal
circuit organization to control the vigor of locomotor movements. Nat
Commun.

Tytell ED (2004) The hydrodynamics of eel swimming II. Effect of swimming
speed. J Exp Biol 207:3265–3279 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/15326203 [Accessed October 23,
2017].

Van Raamsdonk W, Bosch TJ, Smit-Onel MJ, Maslam S (1996) Organisation of

860 the zebrafish spinal cord: distribution of motoneuron dendrites and 5-HT
861 containing cells. Eur J Morphol.

862 Videler JJ, Müller UK, Stamhuis EJ (1999) Aquatic vertebrate locomotion: Wakes
863 from body waves. J Exp Biol 202.

864 Wardle, Videler, Altringham (1995) Tuning in to fish swimming waves: body form,
865 swimming mode and muscle function. J Exp Biol 198.

866 Webb PW (1975) Hydrodynamics and energetics of fish propulsion. Bull Fish Res
867 Board Canada 190:1–159 Available at: http://www.dfo-
868 mpo.gc.ca/library/1486.pdf [Accessed October 23, 2017].

869 Webb PW, KOSTECKI PT, STEVENS ED (1984) The Effect of Size and
870 Swimming Speed on Locomotor Kinematics of Rainbow Trout. J Exp Biol
871 109.

872 Wiggin TD, Anderson TM, Eian J, Peck JH, Masino MA (2012) Episodic
873 swimming in the larval zebrafish is generated by a spatially distributed spinal
874 network with modular functional organization. J Neurophysiol 108:925–934
875 Available at: http://jn.physiology.org/cgi/doi/10.1152/jn.00233.2012
876 [Accessed October 19, 2017].

877 Wiggin TD, Peck JH, Masino MA (2014) Coordination of fictive motor activity in
878 the larval zebrafish is generated by non-segmental mechanisms. Bene F
879 Del, ed. PLoS One 9:e109117 Available at:
880 http://dx.plos.org/10.1371/journal.pone.0109117 [Accessed October 19,
881 2017].

882 Yang L, Rastegar S, Strähle U (2010) Regulatory interactions specifying Kolmer-
883 Agduhr interneurons. Development 137:2713–2722 Available at:
884 http://dev.biologists.org/cgi/doi/10.1242/dev.048470 [Accessed October 20,
885 2017].

886 Zajac FE, Faden JS (1985) Relationship among recruitment order, axonal
887 conduction velocity, and muscle-unit properties of type-identified motor units
888 in cat plantaris muscle. J Neurophysiol 53:1303–1322.

889 Zhang Y, Narayan S, Geiman E, Lanuza GM, Velasquez T, Shanks B, Akay T,
890 Dyck J, Pearson K, Gosgnach S, Fan CM, Goulding M (2008) V3 Spinal
891 Neurons Establish a Robust and Balanced Locomotor Rhythm during
892 Walking. Neuron 60:84–96 Available at:
893 http://linkinghub.elsevier.com/retrieve/pii/S0896627308008064 [Accessed
894 October 20, 2017].

895 Zhong G, Sharma K, Harris-Warrick RM (2011) Frequency-dependent
896 recruitment of V2a interneurons during fictive locomotion in the mouse spinal
897 cord. Nat Commun 2:274 Available at:
898 http://www.nature.com/doifinder/10.1038/ncomms1276 [Accessed October
899 19, 2017].

902
Figure 1. V3-INs are identified by co-localization of *vglut2a:DsRed* and

***nkx2.2a:mEGFP* transgene expression in neurons located in the**

ventromedial spinal cord. (A) Whole-mount

Tg(nkx2.2a:mEGFP)^{vu17};Tg(vglut2a:DsRed)^{nns9} double transgenic zebrafish larva.

(B-D) A representative single confocal optical section of the spinal cord

exhibiting: (B) *nkx2.2a:mEGFP* expression, (C) *vglut2a:DsRed* expression, and

(D) overlayed *nkx2.2a:mEGFP* and *vglut2a:DsRed* expression. The boundaries

of the spinal cord and central canal are illustrated by solid and dashed white

lines, respectively, in panel B. White triangles indicate neurons that co-express

both transgenes and the black triangle indicates the only *vglut2a:DsRed* neuron

ventral to the central canal that did not co-express *nkx2.2a:mEGFP* in the field of

view. Scale bars: 1 mm in A and 10 μm in B-D.

916
917
Figure 2. Distribution of V3-IN Cell Bodies in the Spinal Cord. (A) Maximum

intensity projection of a confocal stack of the spinal cord of a 5 dpf

Tg(*vglut2a:DsRed*)[nns9] zebrafish larva. The dashed white outline indicates the

region of the cord in which cells were quantified. The solid white outline is the

region of the image shown at higher magnification in (B). (B) A higher

magnification field of view. White circles indicate V3 neurons, black triangles

indicate DRG neurons located lateral of the spinal cord. (C) Distribution of

location of all quantified V3 neurons (red dots) projected in the coronal plane of

the spinal cord. The schematic outline of the spinal cord (solid grey line), as well

as the location of the central canal (cc) and Mauthner axon (m), are approximate.

The heavy black line indicates the line of best fit that best describes the axis of

the V3 neuronal population. (D-G) The windowed moving average of V3 density

along the rostrocaudal axis (D; window width: 250 µm), mediolateral axis (E;

window width: 15% SC radius), dorsoventral axis (F; window width: 15% SC

radius), and along the population axis illustrated in panel C (G; window width:

15% SC radius). Black lines are the mean, grey area indicates the standard

deviation. SC: spinal cord. Scale bar: 100 μm in A and 19 μm in B.

Figure 3. V3-INs are descending, bilaterally projecting cells. Kaede was

expressed in *vglut2a*-expressing neurons in

Tg(vglut2a:Gal4ff)^{nns20};*Tg(UAS:Kaede)^{s1999t}* double transgenic larvae (A). A

single V3-IN was exposed to UV laser light, converting the Kaede protein from

green fluorescence to red (B). (C) Inverted contrast projections of Kaede-red

fluorescence in 5 neurons. Each row of images is a neuron shown in a

parasagittal projection (left) and horizontal projection (right). On the horizontal

projections, the center of the cell body is indicated with white triangles to account

for optical bloom and contralateral projections are indicated with black triangles.

946   Lateral projections: Rostral left, dorsal up. Horizontal projections: Rostral left,

947   lateral up. Scale bar: 20 μm.

948
949
Figure 4. V3-IN activity is temporally correlated to fictive locomotion. (A)

Each V3 neuron in the field of view in

Tg(*vglut2a:Gal4ff*)^{nns20};Tg(*UAS:GCaMP6s*)^{nk13a} double transgenic larvae was

assigned a ROI during spontaneous fictive swimming. Panels in (A) show raw

GCaMP6s fluorescence (F) and correspond to numbered time-points in (B).

Bottom panel in A shows calculated ΔF/F between 3 and 7 s time-points. Dashed

lines represent ventral boundary of the spinal cord. (B) Synchronized recordings

of GCaMP6s fluorescence (top) and fictive swimming (bottom) allowed

identification of swimming-related neuronal activity.

959
960
961

Figure 5. Targeted ablation of V3-INs is both efficient and spatially

restricted. (A-B) Confocal stacks of the same Tg(*vglut2a:DsRed*)^{nns9} zebrafish

larva before (A) and 12 hours after (B) laser ablation of the V3-INs in the region

indicated (red dashed lines). (C-D) A magnified view of two V3-INs before (left)

and after (right) a single cell laser ablation. Scale bars: 20 µm in A and B and 5

µm in C and D.

Figure 6. Spinal neurons loaded with rhodamine-2 AM dye are definitively

identified as MNs. (A) The EGFP-expressing *parga^{mn2Et}* enhancer trap line (A,

green) was labeled with antibodies to choline acetyl transferase (ChAT; A′, red).

Colabeling (A″) of EGFP with ChAT antibodies indicated that EGFP expression in

the *parga^{mn2Et}* line is a marker of spinal motor neurons. White arrowheads

indicate ChAT-positive cells that do not contain EGFP (non-MNs). (B) Live

parga^mn2Et larvae were anesthetized, skinned, and incubated in rhodamine-2 AM

dye (Rhod-2, AM). EGFP expression (B, green) and Rhod-2, AM labeling (B′,

red), showed that Rhod-2, AM specifically labeled all motor neurons (B″). White

arrowheads indicate two Rhod-2, AM-labeled cells that did not contain EGFP.

Scale bar: 20 μm.

982
983
984
Figure 7. V3-IN laser ablation reduces the proportion of motor neurons that

are active during fictive swimming. MNs in *Tg(vglut2a:DsRed)^{nns9}* larvae were

loaded with Calcium Green-1 AM dye and assigned ROIs during recording.

Synchronous MN fluorescence and fictive swimming recordings were compared

between Control, V3-IN Ablated, and Sham ablated preparations. (A and B) Raw

Calcium Green-1 AM fluorescence panels correspond to numbered time-points in

aligned ΔF/F (each gray line represents a MN) and peripheral nerve traces in a

Control preparation (A) and a V3-IN Ablated preparation (B). (C-F) Cumulative

probability distributions and mean Peak Burst Frequencies (C and D) and

percent of MNs active (10% ΔF/F threshold) during fictive swimming episodes (E

and F). Asterisks represent significant differences. Error bars represent SD.

		Episode Duration (msec)	Burst Frequency (Hz)	Burst Duration (msec)
Control (*n* = 6 larvae)		720, SD713	25.7, SD3.4	10.4, SD2.8
Ablated (*n* = 6 larvae)		524, SD223	28.3, SD1.9	11.8, SD2.2
Sham (*n* = 6 larvae)		878, SD642	28.2, SD3.8	11.4, SD2.7
ANOVA	**F**	0.58	1.30	0.42
	p	0.57	0.30	0.67

997 Table 1. Measures of the episodic and burst properties from peripheral nerve

998 recordings during fictive locomotion in zebrafish larvae. All values presented as

999 mean, standard deviation.