Large scale genomic rearrangements in selected

Arabidopsis thaliana T-DNA lines are caused by T-DNA

insertion mutagenesis

Boas Pucker[1,2+], Nils Kleinbölting[3+], and Bernd Weisshaar[1*]

1 Genetics and Genomics of Plants, Center for Biotechnology (CeBiTec), Bielefeld University, Sequenz 1, 33615 Bielefeld, Germany

2 Evolution and Diversity, Department of Plant Sciences, University of Cambridge, Cambridge, United Kingdom

3 Bioinformatics Resource Facility, Center for Biotechnology (CeBiTec, Bielefeld University, Sequenz 1, 33615 Bielefeld, Germany

+ authors contributed equally

* corresponding author: Bernd Weisshaar

BP: bpucker@cebitec.uni-bielefeld.de, ORCID: 0000-0002-3321-7471
NK: nkleinbo@cebitec.uni-bielefeld.de, ORCID: 0000-0001-9124-5203
BW: bernd.weisshaar@uni-bielefeld.de, ORCID: 0000-0002-7635-3473

Abstract

Background

Experimental proof of gene function assignments in plants is based on mutant analyses. T-DNA

insertion lines provided an invaluable resource of mutants and enabled systematic reverse

genetics-based investigation of the functions of *Arabidopsis thaliana* genes during the last decades.

Results

We sequenced the genomes of 14 *A. thaliana* GABI-Kat T-DNA insertion lines, which eluded

flanking sequence tag-based attempts to characterize their insertion loci, with Oxford Nanopore

Technologies (ONT) long reads. Complex T-DNA insertions were resolved and 11 previously

unknown T-DNA loci identified, resulting in about 2 T-DNA insertions per line and suggesting that

this number was previously underestimated. T-DNA mutagenesis caused fusions of chromosomes

along with compensating translocations to keep the gene set complete throughout meiosis. Also, an

inverted duplication of 800 kbp was detected. About 10% of GABI-Kat lines might be affected by

chromosomal rearrangements, some of which do not involve T-DNA. Local assembly of selected

reads was shown to be a computationally effective method to resolve the structure of T-DNA

insertion loci. We developed an automated workflow to support investigation of long read data from

T-DNA insertion lines. All steps from DNA extraction to assembly of T-DNA loci can be completed

within days.

Conclusion

Long read sequencing was demonstrated to be an effective way to resolve complex T-DNA

insertions and chromosome fusions. Many T-DNA insertions comprise not just a single T-DNA, but

complex arrays of multiple T-DNAs. It is becoming obvious that T-DNA insertion alleles must be

characterized by exact identification of both T-DNA::genome junctions to generate clear genotype-

to-phenotype relations.

Keywords: long read sequencing, genome assembly, structural variants, translocations,

chromosome fusions, reverse genetics, chromosomal rearrangements, GABI-Kat

Background

T-DNA insertion lines contributed substantially to the high-value knowledge about the functions of

plant genes. This knowledge has been produced by the plant research community on the basis of

gene structures predicted from genome sequences. T-DNA insertional mutagenesis emerged as an

effective mechanism for the generation of knock-out alleles for use in reverse genetics and targeted

gene function search [1, 2]. Since targeted integration of DNA into plant genomes via homologous

recombination was at least technically challenging [3], large collections of sequence-indexed T-

DNA integration lines with random insertion sites were used to provide knock-out alleles for the

majority of genes [4]. Knowledge about the inserted sequences is an advantage over other

mutagenesis methods, because localization of the insertion within the mutagenized genome based

on the generation of flanking sequence tags (FSTs) is possible [5, 6]. CRISPR/Cas technology now

offers technically feasible alternatives for access to mutant alleles for reverse genetics [7].

However, thousands of T-DNA insertion mutants have been characterized and represent today the

main or reference mutant allele for (lack of) a given gene function.

Agrobacterium tumefaciens is a Gram-negative soil bacterium with the ability to transfer DNA into

plant cells and to integrate this T-DNA stably and at random positions into the nuclear genome [8,

9]. The tumor inducing (Ti) plasmid, which is naturally occurring in *Agrobacteria* and enables them

to induce the formation of crown galls in plants, contains the T-DNA that is transferred into plant

cells [10]. The T-DNA is enclosed by 25 bp long imperfect repeats that were designated left (LB)

and right border (RB) [9]. The T-DNA sequence between LB and RB can be modified to contain

resistance genes for selection of successfully transformed plants [11]. T-DNAs from optimized

binary plasmids are transformed into *A. thaliana* plants via floral dip to generate stable lines [12]. T-

DNA transfer into the nucleus of a plant cell is supported by several VIR proteins which are, in the

biotechnologically optimized system, encoded on a separate helper plasmid. It is assumed that host

proteins are responsible for integration of the T-DNA into the genome, most likely as a DNA double

strand into a double strand break (DSB) using host DNA repair pathways and DNA polymerase

theta [9, 13, 14]. T-DNA integration resembles DNA break repair through non-homologous end-

joining (NHEJ) or microhomology-mediated end-joining (MMEJ) and is often accompanied by the

presence of filler DNA or microhomology at both T-DNA::genome junctions [9, 13]. Chromosomal

inversions and translocations are commonly associated with T-DNA insertions in numerous plant

species, not only in *A. thaliana* [15-21], suggesting that often more than just one DSB is associated

with T-DNA integration [9].

The most important collections of T-DNA lines for the model plant *Arabidopsis thaliana* are SALK

(150,000 lines) [6], GABI-Kat (92,000 lines) [22, 23], SAIL (54,000 lines) [24], and WISC (60,000

lines) [25]. In total, over 700,000 insertion lines have been constructed [4]. GABI-Kat lines were

generated through the integration of a T-DNA harboring a sulfadiazine resistance gene for selection

of transformed lines [22]. Additionally, the T-DNA contains a 35S promoter at RB causing

transcriptional up-regulation of plant genes next to the integration site if the right part of the T-DNA

next to RB stays intact during integration [1]. Integration sites were predicted based on FSTs and

allowed access to knock-out alleles of numerous genes. At GABI-Kat, T-DNA insertion alleles were

confirmed by an additional "confirmation PCR" using DNA from the T2 generation [26] prior to the

release of a mutant line and donation of the line to the Nottingham Arabidopsis Stock Centre

(NASC). Researchers could identify available T-DNA insertion lines via SimpleSearch on the GABI-

Kat website [27]. Since 2017, SimpleSearch uses Araport11 annotation data [28]. Araport11 is

based on the *A. thaliana* Col-0 reference genome sequence from TAIR9 which includes about 96

annotated gaps filled with Ns as a sign for unknown sequences [29], among them the centromers

and several gaps in the pericentromeric regions.

The prediction of integration sites based on bioinformatic evaluations using FST data does often

not reveal the complete picture. Insertions might be masked from FST predictions due to truncated

borders [13], because of repetitive sequences or paralogous regions in the genome [30], or even

lack of the true insertion site in the reference sequence used for FST mapping [31, 32]. Also,

confirmation by sequencing an amplicon that spans the predicted insertion site at one of the two

expected T-DNA::genome junctions is not fully informative. Deletions and target site duplications at

the integration site can occur and can only be detected by examining both borders of the inserted

T-DNA [13]. In addition, several studies have reported more complex T-DNA insertions in

association with large deletions, insertions, inversions or even chromosomal translocations [13, 18,

19, 21, 33, 34]. A relation of T-DNA insertion mutagenesis to chromosomal translocations has not

only been reported for *A. thaliana*. Also for transgenic rice (*Oryza sativa*) [35] and transgenic birch

(*Betula platyphylla* x *B. pendula*) [36] changes at the genome level have been reported. Binary

vector backbone (BVB) sequences have been detected at insertion sites [19, 37] as well as

fragments of *A. tumefaciens* chromosomal DNA [38]. Recombination between two T-DNA loci was

described as a mechanism for deletion of an enclosed genomic fragment [39].

Chromosomal rearrangements can occur during the repair of DSBs, e.g. via microhomology-

mediated end joining or non-allelic homologous recombination [reviewed by 40, 41]. Evidently,

regions with high sequence similarity like duplications are especially prone to chromosomal

rearrangements [40].

While usually one T-DNA locus per line was identified by FSTs, the number of T-DNA insertion loci

per line is usually higher. For GABI-Kat, it was estimated that about 50% of all lines (12,018 of

21,049 tested, according to numbers from the end of 2019) display a single insertion locus. This

estimation is based on segregation analyses using sulfadiazine resistance as a selection marker

[22]. Other insertion mutant collections report similar numbers [4]. The average number of T-DNA

insertions per line was reported to be about 1.5, but this is probably an underestimation since the

kanamycin and BASTA selection marker genes applied to determine the numbers are known to be

silenced quite often [4]. For these reasons, it is required that insertion mutants (similar to mutants

created by e.g. chemical mutagenesis) are backcrossed to wild type prior to phenotyping a

homozygous line.

The FSTs produced for the different mutant populations by individual PCR and Sanger-sequencing

allowed usually access to a single T-DNA insertion locus per line, although for GABI-Kat there are

several examples with up to three confirmed insertion loci based on FST data (e.g. line GK-011F01,

see [27]). This leaves a significant potential of undiscovered T-DNA insertions in lines already

available at the stock centers, which has been exploited by the group of Joe Ecker by applying

TDNA-Seq (Illumina technology) to the SALK and a part of the GABI-Kat mutant populations

(unpublished, all data available from GenBank/ENA). Essentially the same technology has

subsequently been used to set up a sequence indexed insertion mutant library of *Chlamydomonas*

reinhardtii [42]. With the fast development of new DNA sequencing technologies, the

comprehensive characterization of T-DNA insertion lines comes into reach.

Several studies already harnessed high throughput sequencing technologies to investigate T-DNA

insertion and other mutant lines [43, 44]. Oxford Nanopore Technologies (ONT) provides a cost-

effective and fast approach to study *A. thaliana* genomes, since a single MinION/GridION Flow Cell

delivers sufficient data to assemble one genotype [45]. Here, we present a method to fully

characterize T-DNA insertion loci and additional genomic changes of T-DNA insertion lines through

ONT long read sequencing. We selected 14 lines that contain confirmed T-DNA insertion alleles.

The first border or first T-DNA::genome junction has been confirmed by sequencing an amplicon

across one junction, but the insertion eluded characterization of the second T-DNA::genome

junction. We refer to the additional T-DNA::genome junction that is expected to exist after

confirmation of one T-DNA::genome junction as "2^{nd} border". Within this set of lines that is enriched

for unusual relations between the two expected T-DNA::genome junctions, we detected several

chromosome fragment or chromosome arm translocations, a duplication of 800 kbp and also an

insertion of DNA from the chloroplast (plastome), all related to T-DNA insertion events. The results

clearly demonstrate the importance of characterizing both T-DNA::genome junctions for reliable

selection of suitable alleles for setting up genotype/phenotype relations for gene function search. In

parallel to data evaluation, we created an automated workflow to support long-read-based analyses

of T-DNA insertion lines and alleles.

154   Results

In total, 14 GABI-Kat T-DNA insertion lines (Table 1, Additional file 1) were selected for genomic

analysis via ONT long read sequencing. This set of lines was selected based on prior knowledge

which indicated that the insertion locus addressed in the respective line was potentially unusual.

The specific feature used for selection was the (negative) observation that creation of confirmation

amplicons which span the T-DNA::genome junction failed for one of the two junctions, operationally

that means that the 2nd border could not be confirmed. T-DNA insertion loci in the selected lines

were assessed by *de novo* assembly of the 14 individual genome sequences, and by a

computationally more effective local assembly of selected reads.

Table 1: Key findings summary of ONT-sequenced GABI-Kat T-DNA insertion lines.

Line ID[a]	number of insertions				Summary of observation
	FST pred.	PCR- conf.	total found	ONT found	
GK-038B07	2	0 (1)[b]	4[c]	3	FST-predicted insertion in Chr5 is part of a fusion of Chr3 and Chr5, 2 T-DNA arrays detected at translocation fusion points of which one contains in addition an inversion of ~2 Mbp fused with another T-DNA array, additional insertion of a complex T-DNA array in Chr1.
GK-089D12	1	1	2	1	Fusion of Chr3 and Chr5, FSTs are derived from the single DNA::genome junction that contains LB, both translocation fusions contain mostly canonical T-DNAs, failure to confirm 2nd T-DNA::genome junction explained by shortened T-DNA.
GK-430F05	1	1	3	2	Three T-DNA insertion sites, two complex T-DNA arrays with one containing more than 5 kbp BVB[d], the confirmed T-DNA insertion in Chr3 is complex, no theoretical explanation for failure of confirmation PCR at 2nd DNA::genome junction, one insertion in the pericentromeric region of (probably) Chr4 containing a rearranged RB region.
GK-654A12[d]	1	1	2[c]	1	Fusion of Chr1 and Chr4 with a complex T-DNA array that also contains BVB, Chr1-part of the fusion predicted by FST, the compensating fusion of Chr4 and Chr1 does not contain T-DNA at the translocation fusion point, translocation explains failure to confirm 2nd T-DNA::genome junction, additional insertion of a T-DNA array in Chr2 with a 162 bp duplicated inversion at the integration site.

GK-767D12	1	1	2	1	Large segmental duplication and inversion at the predicted insertion site, a long T-DNA array also containing BVB present at the southern end of the segmental duplication, inversion explains failure to confirm 2nd T-DNA::genome junction, the northern fusion point of the inverted segmental duplication does not contain T-DNA, another canonical T-DNA insertion in Chr2 but with 4 Mbp distance.
GK-909H04	1	1	3	2	The predicted T-DNA insertion contains an inverted duplication of 20 kbp at the integration site explains why the 2nd T-DNA::genome junction could not be confirmed, integration of 652 bp derived from the plastome at the northern fusion point of the duplication but no T-DNA, two additional canonical T-DNA insertions in Chr1 and Chr2.
GK-947B06	1	1	2	1	Complex T-DNA array insertion of 3 T-DNAs at the predicted insertion site in Chr1, no theoretical explanation for failure of the confirmation PCR at the 2nd DNA::genome junction, additional complex insertion containing 2 T-DNAs and BVB on Chr2.

[a] Lines with newly detected insertions, all lines are listed in Additional file 2.

[b] The T-DNA insertion used for line selection was a false positive case not detected by ONT seq.

[c] One locus in a line with a chromosome translocation causes the FSTs from this single locus to

map to two places in the reference sequence. These two places might also be the source of

potentially existing FST from the compensating chromosome fusion. If the compensating chro-

mosome fusion does not contain a T-DNA, the number of insertions is lower than expected.

[d] BVB, binary vector backbone

A tool designated "loreta" (long read-based t-DNA analysis) has been developed during the

analyses and might be helpful for similar studies (see methods for details). The results of both

approaches demonstrate that a full *de novo* assembly is not always required if only certain regions

in the genome are of interest. The 14 GABI-Kat lines harbor a total of 26 T-DNA insertions resulting

in an average of 1.86 insertions per line. A total of 11 insertion loci detected in seven of 14 lines

were not revealed by previous FST based attempts to detect T-DNA insertions (Table 1, Additional

file 2 and 3). In case of GK-038B07, the lack of re-detection of the expected insertion allele of

At4g19510 was explained by a PCR template contamination during the initial confirmation, the line

that contains the real insertion (source of the contamination) is most probably GK-159D11. A DNA

template contamination is also likely in case of GK-040A12 where the insertion allele of At1g52720

expected from the confirmation was also not found in the ONT data. At least, the error detected fits

to the selection criteria, because the 2nd T-DNA::genome junction can obviously not be detected if

the T-DNA insertion allele as such is not present in the line.

Six of the 14 lines studied by ONT whole genome sequencing displayed chromosomal

rearrangements. To visualize these results, we created ideograms of the five *A. thaliana*

chromosomes with a color code for each chromosome. The colors allow to visually perceive

information on chromosome arm translocations, and the changeover points indicate presence or

absence of T-DNA sequences.

Chromosome fusions

In four lines, fusions of different chromosomes were detected. These fusions result from

chromosome arm translocations which were, in all four cases, compensated within the line by

reciprocal translocations.

The T-DNA insertion on chromosome 5 (Chr5) of GK-038B07 is part of a complex chromosome

arm translocation (Fig. 1). A part of Chr5 is fused to Chr3, the replaced part of Chr3 is fused to an

inversion of 2 Mbp on Chr5. This inversion contains T-DNAs at both ends, one of which is the

insertion predicted by FSTs.

Fig. 1: Organization of the nuclear genome of GK-038B07 with a focus on translocations, inversions

and T-DNA structures.

A: Ideograms of the chromosomes display the reciprocal fusion of Chr3 and Chr5 as well as a 2

Mbp inversion between two T-DNA arrays at the fusion sites; numbers indicate end points of

pseudochromosome fragments according to TAIR9. Chromosomes depicted in grey match the

Col-0 sequence. The insert displays the color codes used for all five chromosomes; N, northern end

of chromosome; S, southern end of chromosome. B-E: Visualization of the four T-DNA insertion loci

of GK-038B07 resolved by local assembly. LB and RB, T-DNA left and right border; dark red, *bona*

fide T-DNA sequences located between the borders; light red, sequence parts from the binary

vector backbone (BVB); numbers above the red bar indicate nucleotide positions with position 1

placed at the left end of LB in the binary vector which makes position 4 the start of the transferred

DNA [13]; numbers below the colored bars indicate pseudochromosome positions according to

TAIR9.

For line GK-082G09, two FST predictions had been generated and one FST lead to the prediction

of an insertion at Chr3:6,597,745 which was confirmed by PCR. Confirmation of the expected

corresponding 2nd border failed. Another FST-based prediction at Chr5:23,076,864 was not

addressed by PCR. ONT sequencing confirmed both predictions (Fig. 2). There is only one

complex insertion consisting of multiple T-DNA copies and BVB in GK-082G09 that fuses the south

of Chr5 to an about 6.6 Mbp long fragment from the north of Chr3 (Fig. 2C). This translocation is

compensated by a fusion of the corresponding parts of both chromosomes without a T-DNA at the

fusion point. The second fusion point of Chr3 and Chr5, that was detected in the *de novo* assembly

of the genome sequence of GK-082G09, was validated by generating and sequencing a PCR

     amplicon spanning the translocation fusion point (Fig. 2B, see Additional file 1 for

sequences/accession numbers and Additional file 4 for the primer sequences).

Fig. 2: Organization of the nuclear genome of GK-082G09 with a focus on translocations,

inversions and T-DNA structures. A-C: For a description of the figure elements see legend to Fig. 1.

D: Read coverage depth analyses of the region of Chr3 that is involved in the fusions which

confirms a deletion of about 4 kbp from Chr3. The reads that cover the deleted part were derived

from the wild type allele present in the segregating population (see methods).

Line GK-089D12 harbors two T-DNA insertions (Fig. 3) and both were predicted by FSTs, one in

Chr3 and one in Chr5. Since fragments of Chr3 and Chr5 are exchanged in a reciprocal way with

no change in sequence direction (southern telomeres stay at the southern ends of the

chromosomes), PCR confirmation would have usually resulted in "fully confirmed" insertion alleles.

Only long read sequencing allowed to determine the involvement of translocations. The line was

studied because the shortened T-DNA at 089D12-At5g51660-At3g63490 (Fig. 3B, see Additional

file 3 for designations of insertions) caused failure of formation of the confirmation amplicon.

Fig. 3: Organization of the nuclear genome of GK-089D12 with a focus on translocations,

inversions and T-DNA structures. A-C: For a description of the figure elements see legend to Fig. 1.

FSTs from line GK-654A12 indicated a T-DNA insertion on Chr1. ONT sequencing revealed a

translocation between Chr1 and Chr4 that explained failure to generate the confirmation amplicon

at the 2nd border (Fig. 4). The southern arms of Chr1 and Chr4 are exchanged, with a T-DNA array

inserted at the fusion point of the new chromosome that contains CEN1 (centromere of Chr1). The

fusion point of the new chromosome that contains CEN4 does not contain T-DNA sequences. Also

this T-DNA-free fusion point (designated 654A12-FCAALL-0-At1g45688) was validated by

generating and sequencing a PCR amplicon which spanned the fusion site (Fig. 4C, Additional files

1 and 4). The T-DNA array at 654A12-At1g45688-FCAALL contains BVB sequences, interestingly

as an independent fragment and not in an arrangement that is similar to the binary plasmid

construction which provided the T-DNA (Fig. 4B).

Fig. 4: Organization of the nuclear genome of GK-654A12 with a focus on translocations, inversions and T-DNA structures. A-C: For a description of the figure elements see legend to Fig. 1. See Additional file 2 for an explanation of pGABI1. D: The T-DNA insertion in Chr2 is associated with a small duplicated inversion of about 160 bp as already described for a fraction of all T-DNA::genome junctions [13].

Intrachromosomal rearrangements and a large duplication

For line GK-433E06 the FST data indicated four insertions, one T-DNA insertion (433E06-At1g73770-F9L1) at Chr1:27,742,275 has been confirmed by amplicon sequencing. ONT sequencing revealed an intrachromosomal translocation that exchanged the two telomeres of Chr1

together with about 5 Mbp DNA. The FSTs that indicated two T-DNA insertions in chromosome 1

were derived from one T-DNA array (Fig. 5). Once more, the compensating fusion point,

designated 433E06-F9L1-0-At1g73770, does not contain T-DNA sequences which was validated

by amplicon sequencing (Fig. 5C, Additional files 1 and 4).

Fig. 5: Organization of the nuclear genome of GK-433E06 with a focus on translocations, inversions

and T-DNA structures. A-C: For a description of the figure elements see legend to Fig. 1.

In line GK-767D12, a large duplication of a part of Chr2 that covers about 800 kbp was detected

(Fig. 6). The duplication is apparent from read coverage analyses (Fig. 6B) based on read mapping

against the TAIR9 reference genome sequence (Col-0) which was performed for all lines studied

(Additional file 5). The duplicated region is inserted in reverted orientation (inversion) next to the T-

DNA insertion 767D12-At2g19210-At2g21385. This insertion was predicted by an FST at

Chr2:8,338,072 and has been confirmed by PCR, the 2[nd] border confirmation for the T-DNA

insertion failed because of reversed orientation. The other end of the duplicated inversion of Chr2 is

fused to Chr2:8,338,361 (designated 767D12-At2g21385-0-At2g19210) without T-DNA sequences

(Fig. 6D). Also this T-DNA-free fusion point was validated by sequencing a PCR amplicon which

spanned the fusion site (Additional files 1 and 4). The T-DNA array at 767D12-At2g19210-

At2g21385 is the largest we detected in this study. It consists of 8 almost complete T-DNA copies

arranged in diversified configurations and includes a large BVB fragment (Fig. 6C).

FST analyses detected only one T-DNA insertion in line GK-909H04. This insertion, designated

909H04-At1g54390, had been confirmed by PCR but failed for the 2nd border. ONT sequencing

revealed an inverted duplication of about 20 kbp next to the T-DNA insertion site (Fig. 7). The

fusion between this inverted duplication and the remaining part of Chr1 does not contain T-DNA

sequences, but a 652 bp fragment derived from the plastome (Fig. 7C). The cpDNA insertion was

validated by generating and sequencing a PCR amplicon spanning the insertion and both junctions

to the genome (Additional files 1 and 4). ONT sequencing also revealed an additional insertion of a

truncated T-DNA (909H04-At1g38212 at about 14.3 Mbp of Chr1) which is in the pericentromeric

region not far from CEN1 (CEN1 is located at 15,086,046 to 15,087,045 and marked in the TAIR9

reference sequence by a gap of 1,000 Ns). Initial analyses indicated that this insertion might be

associated with a deletion of about 45 kbp. However, the predicted deletion was not indicated by

the read coverage depth analyses, and the region is rich in TEs (also At1g38212 is annotated as

"transposable element gene"). We assembled a new genome sequence of the Col-0 wild type used

at GABI-Kat (assembly designated Col-0_GKat-wt, see below) and studied the structure of 909H04-

At1g38212 on the basis of this assembly. The results indicated that the deletion predicted on the

basis of the TAIR9 assembly is a tandemly repeated sequence in TAIR9 which is differently

represented in Col-0_GKat-wt (Additional file 6). The 3'-end of an example read from line GK-

909H04 maps continuously to Col-0_GKat-wt and also to a sequence further downstream in TAIR9.

The evidence collected clearly shows that there are only 13 bp deleted at the T-DNA insertion at

14.3 Mbp of Chr1 (Fig. 7D), and that the initially predicted deletion is caused by errors in the TAIR9

assembly in this pericentromeric region.

Fig. 7: Organization of the nuclear genome of GK-909H04 with a focus on insertions and T-DNA

structures. A-D: For a description of the figure elements see legend to Fig. 6. A: The read coverage

depth plot includes two zoom-in enlargements of the regions at 14.3 and 20.3 Mbp of Chr1. These

two display variable coverage in the region of the truncated T-DNA insertion 909H04-At1g38212

(see text), and increased coverage next to the T-DNA insertion 909H04-At1g54390. C: Duplicated

inversion detected in the local assembly of GK-909H04 which is also supported by read coverage

depicted in Panel A. Green block, sequence part from the plastome (cpDNA).

The six junctions (five sequences) that contained no T-DNA, three from compensating chromosome

fusions, one from the 800 kbp inversion and two at both ends of the cpDNA insertion (see

Additional file 3), were analyzed for specific features at the junctions. The observations made were

fully in line with what has already been described for T-DNA insertion junctions: some short filler

DNA and microhomology was found (Additional file 7). A visual overview over the T-DNA insertion

structures of all 14 lines, including those not displaying chromosomal rearrangements, is presented

in Additional file 8.

Detection of novel T-DNA insertions and T-DNA array structures

As mentioned above, 11 T-DNA insertion loci were newly detected in 7 of 14 lines studied,

indicating that these were missed by FST-based studies (Table 1, Additional file 2 and 3). The

primer annealing sites for FST generation at LB seem to be present in all 11 T-DNA insertions only

found by ONT sequencing. Analysis of the data on T-DNA::genome junctions summarized in

Additional file 3 revealed that a majority of the T-DNA structures have LB sequences at both T-

DNA::genome junctions (14 of 26). The bias for the T-DNA::genome junctions involving LB is

increased by the fact that several of the RB junctions were truncated, and also by some other

junctions which involve BVB sequences. True T-DNA::genome junctions involving intact RB were

not in the dataset, and in 14 out of 26 cases an internal RB::RB fusion was detected.

While about 40% (8 to 10 of 26, depending on judgement of small discontinuous parts) of the

insertions contain a single T-DNA copy (referred to here as "canonical" insertions), some of which

even further truncated and shortened, there are often cases of complex arrays of T-DNA copies

inserted as T-DNA arrays. We observed a wide variety of configurations of the individual T-DNA

copies within the complex arrays. In six out the 26 cases BVB sequences were detected, in the

case of 038B07-At1g14080, 082G09-At5g57020-At3g19080, 430F05-At4g23850 and 947B06-

T7M7 even almost complete vector sequences.

Sequence read quality decreased in T-DNA arrays

During the analyses of T-DNA insertion sequences, we frequently faced regions without sequence

similarity to any sequence in the *A. thaliana* genome sequence, the sequence of the Ti-plasmid (T-

DNA and BVB), or the *A. tumefaciens* genome sequence. Analysis of the read quality (Phred score)

in these regions compared to other regions on the same read revealed a quality drop (Fig. 8).

Consequently, the number of miscalled bases in these regions is especially high. These miscalls

prevent matches in BLAST searches where a perfect match of several consecutive bases is

required as seed for a larger alignment. In some cases, entire reads displayed an extremely low

quality. Reads that display such locally increased error rates were found in the context of T-DNA

array structures which involve head-to-head or tail-to-tail configurations that have the ability to form

foldback structures. It is important to note that the second T-DNA unit was sequenced with lower

quality independent of the read direction (Fig. 8).

Fig. 8: Decrease of Phred score in ONT reads when running from genomic sequence into a T-DNA

array. The insertion used as example is from line GK-038B07 and is shown in Fig. 1C. A/B: Read

quality data from two ONT reads with opposite direction. The reads are displayed in 5' to 3'

orientation. Green and blue bars indicate the read parts matching Chr3 and Chr5, respectively. The

red bar indicates the position of the T-DNA array which is arranged as inverted repeat. Read IDs of

the two examples are shown above each panel. The read quality drop to a Phred score of about 7

occurs when the second part of the inverted repeat is sequenced.

Independent Col-0 assembly resolves misassemblies

As mentioned above for the insertion allele 909H04-At1g38212, the detection of rearrangements in

the insertion lines is not only dependent on the quality of the reads from or assemblies of the

genomes of the lines studied. The results are also dependent on the correctness of the reference

sequence. While the quality of the Col-0 reference sequence (the sequence from TAIR9 is still the

most recent, see Introduction) is generally of high quality, there are some sequence regions that

are not fully resolved. We used a subset of our ONT data, namely ultra-long (> 100 kbp, see

Methods) T-DNA free reads, to *de novo* assemble the Col-0 genome sequence. The assembly,

designated Col-0_GKat-wt, comprises 35 contigs after polishing and displays an N50 of 14.3 Mbp

(GCA_905067165, see Additional file 9). The Col-0_GKat-wt assembly is about 4 Mbp longer than

TAIR9 but still does not reach through any of the centromers. Comparison to the TAIR9 sequence

indicated that the main gain in assembly length was reached in the pericentromeric regions.

Our collection of ONT sequencing datasets from the GABI-Kat lines provides a combined coverage

of over 500x for the TAIR9 reference genome sequence of Col-0. In addition to using ultra-long

reads for generating an assembly, the reads were also used for identification of potentially

problematic regions in the reference sequence. We identified conflicting regions by evaluating read

alignments to assemblies and obtained a list of 383 candidate regions (Additional file 10). We

compared selected regions against our *de novo* genome assembly and focused first on the locus

At1g38212 (at about 14.3 Mbp of Chr1, see Fig. 7). The differences in this region of the TAIR9

assembly, which were detected when analyzing the T-DNA insertion allele 909H04-At1g38212

(Additional file 6), did show up again. Together with nine other examples selected across all

chromosomes, Additional file 11 displays regional comparisons of TAIR9 to Col-0_GKat-wt. The 96

gaps containing various numbers of Ns which are reported for TAIR9 are frequently detected

(Additional files 10 and 11).

Discussion

By sequencing GABI-Kat T-DNA insertion lines with ONT technology, we demonstrate the power of

long read sequencing for the characterization of complex T-DNA insertion lines. The complexity of

these lines has, at least, four aspects: (i) the number of different insertion loci present in a given

line in different regions of the nuclear genome, (ii) the variance of the structures of one or several

T-DNA copies appearing at a given insertion locus, (iii) the changes in the genome sequence in the

direct vicinity of the T-DNA, and (iv) the changes of chromosome topology related to T-DNA

integration.

Number of T-DNA insertion loci per *A. thaliana* insertion line

The average number of T-DNA insertions per *A. thaliana* T-DNA insertion line is assumed to be

about 1.5 [4]. However, the insertion lines available at the stock centers like NASC or ABRC list in

almost all cases only one insertion per line. In our limited dataset of 14 lines, 11 new insertions

were detected among a total of 26, indicating that one should expect an average of about 2

insertions per line. The 11 new insertions all contain sufficiently intact LB sequences that should

have allowed generation of FSTs. The reason for the lack of detection is probably that the FST data

generated at GABI-Kat in total have not reached the saturation level, although several insertions

are predicted per line at GABI-Kat [23]. The potential of existing T-DNA insertion lines for finding

additional knock-out alleles in existing T-DNA insertion lines is also indicated by the fact that TDNA-

Seq revealed additional insertion loci in established and FST-indexed lines (see Introduction).

Clearly, analysis by ONT sequencing can effectively reveal additional insertions and can

successfully be used to fully characterize the genomes of T-DNA insertion lines. This approach is

faster, less laborious, more comprehensive and compared to the level of reliability also significantly

cheaper than PCR- or short-read based methods.

Structure of the inserted T-DNA or T-DNA array

The variance of the T-DNA structures that were resolved by ONT sequencing spans a really wide

range of configurations and lengths. Tandem repeats as well as inverted repeats [46] are occurring.

Insertion length starts with 2.7 kbp for 909H04-At1g38212 (Fig. 7B) and reaches up to about 50 kbp

for 767D12-At2g19210-At2g21385 (Fig. 6C). The lines were selected to contain a T-DNA by

checking for resistance to sulfadiazine [22] which is provided by the T-DNA used at GABI-Kat.

However, since there are regularly several insertions per line, also T-DNA fragments with a

truncated selection marker gene are to be expected - given that resistance is provided in trans. For

SALK, SAIL and WISC insertion lines, T-DNA arrays sizes of up to 236 kbp have been reported

[19]. We hypothesize that the complexity of T-DNA arrays might correlate with the tendency of

selection marker silencing, which could mechanistically be implemented via siRNA [19]. The

comparably reduced complexity of T-DNA arrays derived from pAC161 (the binary vectors mostly

used at GABI-Kat) could thus explain why the sulfadiazine selection marker stays active for many

generations.

Inclusion of BVB sequences in T-DNA array structures has been reported repeatedly for various

species [37, 47, 48]. For the studied GABI-Kat lines, BVB sequences were structurally resolved as

internal components of T-DNA arrays as well as at the junction to genomic sequences. A total of six

T-DNA arrays with BVB sequences were detected among 26 cases, indicating that about 20% of all

insertions, and an even higher percentage of lines, contain inserted BVB sequences.

We detected only few intact right border sequences in contrast to left border sequences, which fits

the empirical observation that FST-generation for characterization of insertion populations is much

more productive at LB than at RB [4-6]. In turn, the lines studied here are selected from insertions

detected by using LB for FST generation, which introduces a bias. When insertions accessed via

FSTs from RB were studied, RB is found to be more precisely cut than LB [13, 49], which is

explained by protection of RB by VirD2 [9]. Nevertheless, within the longer T-DNA arrays and also

in the insertions newly detected by ONT sequencing, most of the RBs are lost. This does not fit well

to current models for the integration mechanism and explanations for the observed internal "right

end to right end" (without RB) fusions in T-DNA arrays and requires further investigation.

Changes in the genome sequence at the insertion site

Changes in the genome sequence in the direct vicinity of the T-DNA insertion site have already

been described in detail [13]. However, that study relied on data from PCR amplicon sequences

and could, therefore, not detect events that affect distances longer than the length of an average

amplimer of about 2 kbp. In addition, amplicons from both T-DNA::genome junctions were required.

Here, we addressed insertions that failed to fulfill the "amplicon sequences from both junctions

available" criterion. This allowed to focus on a set of GABI-Kat lines that has a higher chance of

showing genomic events (Table 1). The T-DNA::genome junctions studied here fall, with one

exception, into the range already described for DSB-based integration and repair by NHEJ, with

filler sequences and microhomology at the insertion site [9]. The exception is 909H04-At1g54385-

cp-At1g54440 (Fig. 7C), an insertion allele that displays an about 20 kbp long duplicated inversion

and in addition 652 bp derived from the plastome at the additional breakpoint that links the

inversion back to the chromosome. During repair of the initial DSBs and in parallel to T-DNA

integration, also cpDNA is used to join broken ends of DNA at insertion loci. Inversions require

more than one repaired DSB in the DNA at the insertion site to be explained, and that cpDNA is

available in the nucleus has been demonstrated experimentally [50] and also in the context of

horizontal gene transfer [51].

Genome level changes and translocations related to T-DNA integration

Our analyses revealed five lines with chromosome arm translocations, either exchanged within one

chromosome (GK-433E06, Fig. 5) or exchanged between two different chromosomes (Figures 1 to

4). In addition, line GK-767D12 displayed a chromosomal rearrangement that resulted in an

inverted duplication of 0.8 Mbp. This aligns well with previous reports of interchromosomal

structural variations, translocations, and chromosome fusions in T-DNA insertion lines [18, 19, 33,

34]. Because of the bias for complex cases in the criteria we used for selection of the lines

investigated, we cannot deduce a reliable value for the frequency of chromosomal rearrangements

in the GABI-Kat population. However, an approximation taking into account that the 6 cases are

from 14 lines sequenced, and the 14 lines sequenced are a subset of 342 out of 1,818 lines with

attempted confirmation of both borders but failure at the 2nd T-DNA::genome junction, ends up with

about one of 10 GABI-Kat lines that may display chromosome-level rearrangements (~10%). It

remains to be determined if this estimation holds true, but the approximation fits somehow to the

percentage of T-DNA insertion lines that show Mendelian inheritance of mutant phenotypes (88%)

while 12% do not [52]. For the SALK T-DNA population, 19% lines with chromosomal translocations

have been reported [18], based on genetic markers and lack of linkage between markers from

upstream and downstream of an insertion locus.

Although the number of investigated lines with chromosome arm translocations is small, the high

proportion of fusions between Chr3 and Chr5 in our dataset is conspicuous (3 out of 5, see

Additional file 3). Also for the line SAIL_232 a fusion of Chr3 and Chr5 was reported [19]. This work

addressed 4 T-DNA insertion lines (two SALK, one WISC and SAIL_232) by ONT sequencing and

Bionano Genomics (BNG) optical genome maps. Translocations involving chromosomes other than

Chr3 and Chr5 were observed in our study and have also been reported before [16-18, 33], but it is

possible that translocations between Chr3 and Chr5 occur with a higher rate than others. Full

sequence characterization of the genomes of (many) more T-DNA insertion lines by long read

sequencing have the potential to reveal hot spots of translocations and chromosome fusions, if

these exist.

ONT reads are best suited for the identification of large scale structural variants, for this reason we

did not address single nucleotide variants (SNV) or small insertions/deletions (InDels). T-DNA-

independent structural variants that were identified, except chromosome fusions, are mostly

explained by potential errors in the TAIR9 reference sequence as discussed below. One may

conclude that the remaining 8 out of 14 lines for which we have not detected large structural

variation are identical in genome organization to the Col-0 wild type. However, this conclusion is

based on negative results and should be taken with care.

Compensating translocations

The chromosome arm translocations detected are all "reciprocal" translocations, which involve two

breakpoints and exchange parts of chromosomes. Both rearranged chromosomes are equally

detected in the sequenced DNA of the line. The combination of both rearranged chromosomes in

the offspring is maintained because homozygosity of only one of the two rearranged chromosomes

is expected to be lethal due to imbalance of gene dose for large chromosomal regions. However, if

both rearranged chromosomes can be transmitted together in one gametophyte, both rearranged

chromosomes might exist in offspring in homozygous state [53]. The induction of chromosome

mutations, chromosomal rearrangements and chromosome arm translocations by T-DNA insertion

mutagenesis has since a long time received attention in *A. thaliana* genetics. One reason is that

these types of mutations cause distorted segregation among offspring since similar segregation

results are also indicative of genes essential for gametophyte development [53-55]. With regard to

the chromosome arm translocations we detected, it is important to note that some of the arms are

fused without integrated T-DNA. The case of line GK-082G09 (one T-DNA insertion locus, still

reciprocal chromosome arm translocation) is relevant in this context, because the presence of a

single T-DNA insertion per line was used as a criterion to select valid candidates for gametophyte

development mutants.

Our analyses of the sequences of T-DNA free chromosomal junctions (e.g. 082G09-At5g57020-0-

At3g19080) did not result in the detection of specialties that make these junctions different from T-

DNA::genome junctions. We cannot fully exclude that the T-DNA free junctions are the result of

recombination of two loci that initially both contained T-DNA, and that one T-DNA got lost during

recombination at one of two loci. However, it is also possible that the translocations are the direct

result of DSB repair, similar to what has been found after targeted introduction of DSBs [56]. We

speculate that both, T-DNA containing and T-DNA free junction cases, result from

DSB/integration/repair events that involve genome regions which happen to be in close contact,

even if different chromosomes are involved. It is evident that several DSB breaks are required, and

repair of these DSB can happen with the DNA that is locally available, might it be cpDNA (see

above), T-DNA that must have been delivered to DSB repair sites, or different chromosomes that

serve as template for fillers [13] or as target for fusion after a DSB happened to occur.

ONT sequencing of mutants offers easy access to data on presence or absence of translocations

and other structural variants. We demonstrated that long read sequencing is a suitable method for

characterizing lines that failed in previous characterization attempts based on FSTs. For example,

the investigation of T-DNA insertion alleles/lines that display deformed pollen phenotypes, which

was impacted by chromosome fusions and uncharacterized T-DNA insertions [55], can now be

combined with long read sequencing to reveal all insertion and structural variation events with high

resolution. Clearly, comprehensive characterization of T-DNA insertion lines, independent from the

population from which the mutant originates, as well as other lines used for forward and reverse

genetic experiments, can prevent unnecessary work and questionable results. The entire workflow

from DNA extraction to the final genome sequence can be completed in less than a week.

Estimated costs per line for ONT sequencing are $200-300 if multiple samples are consecutively

analyzed on one GridION flow cell or if barcoding and multiplexing are applied. The application of

"loreta" supports the inspection of T-DNA insertions as soon as the read data are generated.

Therefore, we recommend to characterize T-DNA insertion alleles by long read sequencing.

Analyses of inverted duplicated DNA sequences by ONT sequencing

Decreased quality (Phred scores) was previously described for ONT sequence reads as

consequence of inverted repeats which might form secondary structures and thus interfere with the

DNA translocation through the nanopore [57]. Complex T-DNA arrays are a challenge to ONT

sequencing and probably all other current sequencing technologies because of the frequently

included inverted repeat structures. We observed that the second part of an inverted repeat has low

sequence quality, while the first part (probably not engaging in secondary structure) is of good

sequence quality. Evidence has been provided that changed translocation speed is responsible for

the low Phred scores in inverted repeats [57]. The quality decrease needs to be considered

especially when performing analyses at the single read level. These stretches of sequence with bad

quality also pose a challenge for the assembly, especially since the orientation of the read

determines which part of the inverted repeat is of good or poor quality. However, we were able to

solve the problem to a satisfying level by manual consideration of reads from the opposite direction

which contain the other part of the inverted repeat in good sequence quality.

Conclusions

This study presents a comprehensive characterization of multiple GABI-Kat lines by long read

sequencing. The results argue strongly for full characterization of mutant alleles to avoid

misinterpretation and errors in gene function assignments. If an insertion mutant and the T-DNA

insertion allele in question are not characterized well at the level of the genotype, the phenotype

observed for the mutant might be due to a complex integration locus, and not causally related to the

gene that is expected to be knocked-out by the insertion. Structural changes at the genome level,

including chromosome translocations and other large rearrangements with junctions without T-

DNA, may have confounding effects when studying the genotype to phenotype relations with T-

DNA lines. This conclusion must also consider that during the last 20 to 30 years, many T-DNA

alleles have been used in reverse genetic experiments. Finally, and similar to the ONT sequence

data that resulted from the analyses of four SALK and SAIL/WISC T-DNA insertion lines [19], the

ONT sequence data from this study allowed to detect and correct many non-centromeric

misassemblies in the current reference sequence.

Methods

Plant material

The lines subjected to ONT sequencing were chosen from a collection of GABI-Kat lines which

were studied initially to collect statistically meaningful data about the structure of T-DNA insertion

sites at both ends of the T-DNA insertions [13]. In this context and also after 2015, confirmation

amplicon sequence data from both T-DNA::genome junctions of individual T-DNA insertions were

created at GABI-Kat, which was successful for 1,481 cases from 1,476 lines by the end of 2019

(1,319 cases were successfully completed for both junctions in the beginning of 2015). To generate

this dataset, 1,835 individual T-DNA insertions from 1,818 lines with one T-DNA::genome junction

already confirmed were addressed, meaning that there were 354 cases from 342 lines which failed

at the 2nd T-DNA::genome junction. From these 354 cases, we randomly selected the 14 insertions

(in 14 different lines) that were studied here, with good germination as additional criterion for

effective handling (Additional file 1). Since the focus of interest in insertion alleles was always to

identify null alleles of genes, all 14 insertions addressed are CDSi insertions (insertions in the

coding sequence or enclosed introns, that means between ATG and Stop codons on genomic

DNA). A total of 100 T2 seeds of each line were plated with sulfadiazine selection as described

[26]. Surviving T2 plantlets contain at least one integrated T-DNA, either in hemizygous or in

homozygous state. Sulfadiazine-resistant plantlets were transferred to soil, grown to about 8-leaf

stage and pooled for DNA extraction. For a single locus with normal heritability, statistically 66% of

the chromosomes in the pool contain the T-DNA. The T-DNA in GK-654A12 is from pGABI1 [38],

the other lines contain T-DNA from pAC161 [22].

DNA extraction, size enrichment, and quality assessment

Genomic DNA was extracted from young plantlets or young leaves through a CTAB-based protocol

(Additional file 12) modified from [22, 58]. We observed, like others before [4], that the quality of

extracted DNA decreased with the age of the leaf material processed. Young leaves lead to the

best results in our hands. The cause might be increasing cell and vacuole size containing more

harmful metabolites which may cause reduced quality and yield of DNA. As DNA quality for ONT

sequencing also decreases with storage time, we processed the DNA as soon as possible after

extraction. DNA quantity and quality was initially assessed based on NanoDrop (Thermo Scientific)

measurement, and on an agarose gel for DNA fragment size distribution. Precise DNA

quantification was performed via Qubit (Thermo Fisher) measurement using the broad range buffer

following the supplier's instructions. Up to 9 µg of genomic DNA were subjected to an enrichment of

long fragments via Short Read Eliminator kit (Circulomics) according to the suppliers' instructions.

Library preparation and ONT sequencing

DNA solutions enriched for long fragments were quantified via Qubit again. One µg DNA (R9.4.1

flow cells) or two µg (R10 flow cells) were subjected to library preparation following the LSK109

protocol provided by ONT. Sequencing was performed on R9.4.1 and R10 flow cells on a GridION.

Real time base calling was performed using Guppy v3.0 on the GridION (R9.4.1 flow cells) and on

graphic cards in the de.NBI cloud [59] (R10 flow cells), respectively.

De novo genome sequence assemblies

Reads of each GK line were assembled separately to allow validation of other analysis methods

(see below). Canu v1.8 [60] was deployed with previously optimized parameters [61]. Assembly

quality was assessed based on a previously developed Python script (Table 2). No polishing was

performed for assemblies of individual GABI-Kat lines as these assemblies were only used to

analyze large structural variants and specifically T-DNA insertions.

Through removal of all T-DNA reads from the combined ONT read dataset from all insertion lines

(Col-0 background [22]) and size filtering, a comprehensive data set of ultra-long reads (longer than

100,000 nt) was generated. This dataset is available from ENA/GenBank with the ID ERS5246674

(SAMEA7490021). The assembly of these ultra-long reads was computed as described above.

Polishing was performed with Racon v.1.4.7 [62] and medaka v.0.10.0 as previously described [58].

Potential contamination sequences were removed based on sequence similarity to the genome

sequences of other species, and contigs smaller than 100 kbp were discarded as previously

described [58, 63]. To ensure accurate representation of the Col-0 wild type genome structure, the

assembly was checked for the chromosome fusion events reported for GK-082G09, GK-433E06,

and GK-654A12 as well as for the chloroplast DNA integration of GK-909H04. The Sanger reads of

the validation amplicons generated for these loci were subjected to a search via BLASTn [64] using

default settings. BLASTn was also used to validate the absence of any T-DNA or plasmid

sequences in this assembly using pSKI015 (AF187951), pAC161 (AJ537514), and pROK2 [6] as

query.

Analyses of the Col-0_GKat-wt assembly and comparison to TAIR9

To identify differences to the TAIR9 reference genome sequence, the Col-0_GK-wt contigs were

sorted and orientated using pseudogenetic markers derived from TAIR9. The TAIR9 sequence was

split into 500 bp long sequence chunks which were searched against the Col-0_GK-wt contigs via

BLAST. Unique hits with at least 80% of the maximal possible BLAST score were considered as

genetic markers. The following analysis with ALLMAPS [65] revealed additional and thus

unmatched sequences of Col-0_GK-wt around the centromeres.

ONT reads were aligned to the TAIR9 reference genome sequence via Minimap2 v2.1-r761 [66].

Mappings were converted into BED files with bedtools v2.26 [67]. The alignments were evaluated

for the ends of mapping reads, and these ends were quantified in genomic bins of 100 bp using a

dedicated tool designated Assembly Error Finder (AEF) v0.12 (Table 2) with default parameters.

Neighboring regions with high numbers of alignment ends were grouped if their distance was

smaller than 30 kbp. Regions with outstanding high numbers of alignment ends indicate potential

errors in the targeted assembly. A selection of these regions from TAIR9 was compared against

Col-0_GKat-wt through dot plots [32].

Analysis of T-DNA insertions

The T-DNA insertions of each line were analyzed in a semi-automatic way. A tool was developed,

written in Python and designated "loreta" (Table 2), that needs as input: reads in FASTQ format, T-

DNA sequences in FASTA format, a reference file containing sequences for assembly annotation in

FASTA format (for this study: sequences of T-DNA and vector backbone, the *A. thaliana* nuclear

genome, plastome, chondrome, and the *A. tumefaciens genome*), and – if available – precomputed

de novo assemblies (as described above). The output are HTML pages with annotated images

displaying models of T-DNA insertions and their neighborhood. Partial assemblies of reads

containing T-DNA sequences are computed, and the parts of the *de novo* assemblies containing T-

DNA sequences are extracted. All resulting sequences as well as all individual reads containing T-

DNA sequences are annotated using the reference file. If run on a local machine, a list of tools that

need to be installed is given in the github repository (Table 2). For easier access to the tool on a

local machine, there is also a Docker file available that can be used to build a Docker image.

Reads containing T-DNA sequences were identified by BLASTn [64, 68] using an identity cutoff of

80% and an e-value cutoff of 1e-50. All identified reads were then assembled using Canu v1.8 with

the same parameters as for the *de novo* assemblies (see above) and in addition some parameters

to facilitate assemblies with low coverage: correctedErrorRate=0.17, corOutCoverage=200,

stopOnLowCoverage=5 and an expected genome size of 10 kbp. From the precomputed *de novo*

assemblies, fragments were extracted that contain the T-DNA insertion and 50 kbp up- and

downstream sequence. The resulting fragments, contigs from the Canu assembly, the contigs

marked as "unassembled" by Canu as well as all individual reads (converted to FASTA using

Seqtk-1.3-r106 [69]) were annotated using the reference sequences. For this purpose, a BLASTn

search was performed (contig versus reference sequences) with the same parameters as for the

identification of T-DNA reads. These BLAST results were mapped to the sequence (contig/read) as

follows: BLAST hits were annotated one after another, sorted by decreasing score. If the overlap of

a BLAST hit with a previously annotated one exceeds 10 bp, the second BLAST hit was discarded.

For further analysis, reads were mapped back to the assembly. All reads were mapped back to the

fragments of the *de novo* assembly, reads containing T-DNA were mapped back to (1) the Canu

contigs, (2) "unassembled" contigs and (3) individual reads containing T-DNA sequences. Mapping

was performed using Minimap2 [66] with the default options for mapping of ONT sequencing data.

To further inspect the chromosome(s) sequences prior to T-DNA insertion, the same analysis was

performed using the *A. thaliana* sequences neighboring the T-DNA insertion. A FASTA file was

generated that contains these flanking sequences using bedtools [67], reads containing this part of

A. thaliana sequence (and no T-DNA) are again identified using BLAST, assembled and annotated

as described above. Infoseq from the EMBOSS package [70] was used to calculate the length of

different sequences in the pipeline.

All information is summarized in HTML files containing images; these images display all annotated

sequences along with mapped reads and details on the BLAST results. These pages were used for

manual inspection and final determination of the insertion structures. For lines with a single

canonical insertion, such as 050B11-At5g64610 in GK-050B11 (see Additional file 3), the

assembled and annotated contig of the partial assembly was sufficient. In more complex cases like

GK-038B07, the exact insertion structure was not clear based on the assembled contigs or based

on sections of the *de novo* assembly. If, based on the read mappings shown in the visualization,

the partial assembly looked erroneous (many partial mappings), individual reads were used for the

determination of the insertion structure. These individual reads were also considered for

determination of exact T-DNA positions, if these positions were not clear from contigs / sections.

This was often the case for head-to-head or tail-to-tail configurations of T-DNA arrays, where one of

the T-DNAs was represented by sequence of low quality and could not be annotated (and led to

misassemblies in assembled contigs). These cases were resolved by identification of reads

orientated in the other direction, because then the sequence derived from the other T-DNA was of

low quality and by combining the annotated results, a clear picture could be derived. If different

reads contradicted each other in exact positions of the T-DNA, we chose the "largest possible T-

DNA" that could be explained by individual reads.

Mapping of ONT reads for detection of copy number variation

ONT reads from each line were aligned against the TAIR9 Col-0 reference genome sequence using

Minimap v2.10-r761 [71] with the options '-ax map-ont --secondary=no'. The resulting mappings

were converted into BAM files via samtools v1.8 [72] and used for the construction of coverage files

with a previously developed Python script [73]. Coverage plots (see Additional file 5) were

constructed as previously described [61] and manually inspected for the identification of copy

number variations.

Sequence read quality assessment

Reads containing T-DNA sequence were annotated based on sequence similarity to other known

sequences based on BLASTn [64, 68] usually matching parts of the Ti-plasmid or *A. thaliana*

genome sequence. Reads associated with complex T-DNA insertions were considered for

downstream analysis if substantial parts (>1 kbp) of the read sequence were not matched to any

database sequences via BLASTn. Per base quality (Phred score) of such reads was assessed

based on a sliding window of 200 nt with a step size of 100 nt.

Chromosome fusion and cpDNA insertion validation via PCR and Sanger sequencing

Chromosomal fusions without a connecting T-DNA were analyzed via PCR using manually

designed flanking primers (Additional file 4). Amplicons were generated using genomic DNA

extracted from plants of the respective line as template with Q5 High-Fidelity DNA polymerase

(NEB) following supplier's instructions. PCR products were separated on a 1% agarose gel and

visualized using ethidiumbromide and UV light. Amplicons were purified with Exo-CIP Rapid PCR

Cleanup Kit (NEB) following supplier's instructions. Sanger sequencing was performed at the

Sequencing Core Facility of the Center for Biotechnology (Bielefeld University, Bielefeld, Germany)

using BigDye terminator v3.1 chemistry (Thermo Fisher) on a 3730XL sequencer. The resulting

Sanger sequences were merged using tools from the EMBOSS package [70]. After transferring

reverse reads to their reverse complement using revseq, a multiple alignment was generated using

MAFFT [74]. One consensus sequence for each amplicon was extracted from the alignments using

em_cons with option -plurality 1, and the resulting sequences were submitted to ENA (see

Additional file 1 for accession numbers).

Analyses of T-DNA free chromosome fusion junctions

The five junction sequences were analyzed by BLAST essentially as described [13]. Briefly,

searches were performed against all possible target sequences (*A. tumefaciens*; *A. thaliana*

nucleome, plastome and chondrome; T-DNA and vector backbone) using BLASTn default

parameters. If the complete query was not covered, the unmatched part of the query sequence was

classified as filler and extracted. Subsequently, this sequence was used in a BLAST search with an

e-value cutoff of 10, a word-size of 5 and the '-task "blastn-short"' option activated to detect smaller

and lower quality hits. If this was not successful (as in GK-909H04), the filler sequence was

extended by 10 bases up- and downstream and the procedure described above was repeated.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

Sequence read datasets generated and analyzed during this study were made available at ENA

under the accession PRJEB35658. Individual run IDs are included in Additional file 1. The Col-0

genome sequence assembly of the GABI-Kat Col-0 genetic background (Col-0_GKat-wt) is

available at ENA under the accession GCA_905067165.

Table 2: Availability of scripts.

Description	URLs
Previously developed scripts for general tasks	https://github.com/bpucker/script_collection
Tool for the analysis of ONT datasets ("loreta")	https://github.com/nkleinbo/loreta
Scripts for the analyses presented in this study	https://github.com/bpucker/GKseq

779   **Competing interest**

780   The authors declare that they have no competing interests.

Funding

783   We acknowledge support for the Article Processing Charge by the Open Access Publication Fund

784   of Bielefeld University.

Authors' contribution

787   BP performed DNA extraction and sequencing. BP and NK performed bioinformatic analyses. BP,

788   NK, and BW interpreted the results and wrote the manuscript.

Acknowledgements

We are very grateful to the Bioinformatics Resource Facility support team of the CeBiTec and to

de.NBI for providing computing infrastructure and excellent technical support. We also thank the

Sequencing Core Facility of the CeBiTec for granting access to the ONT infrastructure. Many

thanks to Tobias Busche, Christian Rückert, and Jörn Kalinowski for general support related to the

ONT sequencing, and to Prisva Viehöver for high quality Sanger sequencing. We thank Andrea

Voigt for excellent technical support. The bioinformatic work was supported in part by grants from

the German Federal Ministry of Education and Research (BMBF) for the project "Bielefeld-Gießen

Center for Microbial Bioinformatics–BiGi" (grant no. 031A533) within the German Network for

Bioinformatics Infrastructure (de.NBI).

Supplementary information

Additional file 1: Summary of GABI-Kat insertion line data, segregation data for the F2 families after

selection for sulfadiazine resistance, sequencing data including run IDs from submission to

ENA/SRA, and accession numbers of T-DNA free junction sequences.

Additional file 2: Extended version of Table 1 covering all 14 lines.

Additional file 3: Overview of the T-DNA insertions and associated structural variants in the

investigated GABI-Kat lines.

Additional file 4: Sequences of oligonucleotides used for the validation of fusion points of

chromosomal translocations and other large structural variants.

Additional file 5: Read coverage of all analyzed lines in relation to the TAIR9 reference genome

sequence.

Additional file 6: Structure of genomic locus around one insertion in GK-909H04.

Additional file 7: Analysis results of genomic fusion junction sequences without T-DNA insertion.

Additional file 8: Visual overview over all insertions detected.

Additional file 9: Assembly statistics of Col-0_GK-wt.

Additional file 10: Potential errors in the TAIR9 reference genome sequence of Col-0.

Additional file 11: Dot plots between TAIR9 and Col-0_GK-wt for potential errors in the reference

sequence.

Additional file 12: Protocol for the extraction of genomic DNA from *A. thaliana* for ONT sequencing.

References

1.   Ulker B, Peiter E, Dixon DP, Moffat C, Capper R, Bouche N, Edwards R, Sanders D, Knight

H, Knight MR: **Getting the most out of publicly available T-DNA insertion lines**. *The Plant*

Journal 2008, **56**(4):665-677.

2.   OMalley RC, Ecker JR: **Linking genotype to phenotype using the Arabidopsis unimutant**

collection. *The Plant Journal* 2010, **61**(6):928-940.

3.   Fauser F, Roth N, Pacher M, Ilg G, Sanchez-Fernandez R, Biesgen C, Puchta H: **In planta**

gene targeting. *Proceedings of the National Academy of Sciences of the United States of*

America 2012, **109**(19):7535-7540.

4.   OMalley RC, Barragan CC, Ecker JR: **A user's guide to the Arabidopsis T-DNA insertion mutant collections**. In: *Methods in Molecular Biology.* Edited by Alonso J, Stepanova A, vol. 1284, 2015/03/12 edn. New York, NY: Humana Press; 2015: 323-342.

5.   Strizhov N, Li Y, Rosso MG, Viehoever P, Dekker KA, Weisshaar B: **High-throughput generation of sequence indexes from T-DNA mutagenized Arabidopsis thaliana lines**. *BioTechniques* 2003, **35**(6):1164-1168.

6.   Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, Stevenson DK, Zimmerman J, Barajas P, Cheuk R *et al*: **Genome-wide insertional mutagenesis of Arabidopsis thaliana**. *Science* 2003, **301**(5633):653-657.

7.   Puchta H: **Applying CRISPR/Cas for genome engineering in plants: the best is yet to come**. *Current Opinion in Plant Biology* 2017, **36**:1-8.

8.   Smith EF, Townsend CO: **A Plant-Tumor of Bacterial Origin**. *Science* 1907, **25**(643):671-673.

9.   Gelvin SB: **Integration of Agrobacterium T-DNA into the Plant Genome.** *Annual Review of Genetics* 2017, **51**:195-217.

10.   Zambryski P, Holsters M, Kruger K, Depicker A, Schell J, Van Montagu M, Goodman H: **Tumor DNA structure in plant cells transformed by A. tumefaciens**. *Science* 1980, **209**(4463):1385-1391.

11.   Hernalsteens J, Van Vliet F, De Beuckeleer M, Depicker A, Engler G, Lemmers M, Holsters M, Van Montagu M, Schell J: **The Agrobacterium tumefaciens Ti plasmid as a host vector system for introducing foreign DNA in plant cells**. *Nature* 1980, **287**(5783):654-656.

12.   Clough SJ, Bent AF: **Floral dip: a simplified method for Agrobacterium-mediated transformation of** *Arabidopsis thaliana*. *The Plant Journal* 1998, **16**(6):735-743.

13.   Kleinboelting N, Huep G, Appelhagen I, Viehoever P, Li Y, Weisshaar B: **The Structural**

858   **Features of Thousands of T-DNA Insertion Sites Are Consistent with a Double-Strand Break**

859   **Repair-Based Insertion Mechanism.** *Molecular Plant* 2015, **8**(11):1651-1664.

860   14.   van Kregten M, de Pater S, Romeijn R, van Schendel R, Hooykaas PJ, Tijsterman M: **T-**

861   **DNA integration in plants results from polymerase-θ-mediated DNA repair.** *Nature Plants*

862   2016, **2**(11):16164.

863   15.   Castle LA, Errampalli D, Atherton TL, Franzmann LH, Yoon ES, Meinke DW: **Genetic and**

864   **molecular characterization of embryonic mutants identified following seed transformation in**

865   **Arabidopsis.** *Molecular Genetics and Genomics* 1993, **241**(5-6):504-514.

866   16.   Forsbach A, Schubert D, Lechtenberg B, Gils M, Schmidt R: **A comprehensive**

867   **characterization of single-copy T-DNA insertions in the Arabidopsis thaliana genome.** *Plant*

868   *Molecular Biology* 2003, **52**(1):161-176.

869   17.   Lafleuriel J, Degroote F, Depeiges A, Picard G: **A reciprocal translocation, induced by a**

870   **canonical integration of a single T-DNA, interrupts the HMG-I/Y Arabidopsis thaliana gene.**

871   *Plant Physiology and Biochemistry* 2004, **42**(3):171-179.

872   18.   Clark KA, Krysan PJ: **Chromosomal translocations are a common phenomenon in**

873   **Arabidopsis thaliana T-DNA insertion lines.** *The Plant Journal* 2010, **64**(6):990-1001.

874   19.   Jupe F, Rivkin AC, Michael TP, Zander M, Motley ST, Sandoval JP, Slotkin RK, Chen H,

875   Castanon R, Nery JR *et al*: **The complex architecture and epigenomic impact of plant T-**

876   **DNA insertions.** *PLoS Genetics* 2019, **15**(1):e1007819.

877   20.   Min Y, Frost JM, Choi Y: **Gametophytic Abortion in Heterozygotes but Not in Homozygotes:**

878   **Implied Chromosome Rearrangement during T-DNA Insertion at the ASF1 Locus in**

879   **Arabidopsis.** *Molecules and Cells* 2020, **43**(5):448-458.

21.   Krispil R, Tannenbaum M, Sarusi-Portuguez A, Loza O, Raskina O, Hakim O: **The Position**

and Complex Genomic Architecture of Plant T-DNA Insertions Revealed by 4SEE.

International Journal of Molecular Sciences 2020, **21**(7):ijms21072373.

22.   Rosso MG, Li Y, Strizhov N, Reiss B, Dekker K, Weisshaar B: **An *Arabidopsis thaliana* T-**

DNA mutagenised population (GABI-Kat) for flanking sequence tag based reverse genetics.

Plant Molecular Biology 2003, **53**(1-2):247-259.

23.   Kleinboelting N, Huep G, Kloetgen A, Viehoever P, Weisshaar B: **GABI-Kat SimpleSearch:**

new features of the Arabidopsis thaliana T-DNA mutant database. *Nucleic Acids Research*

2012, **40**(Database issue):D1211-D1215.

24.   Sessions A, Burke E, Presting G, Aux G, McElver J, Patton D, Dietrich B, Ho P, Bacwaden

J, Ko C *et al*: **A High-Throughput Arabidopsis Reverse Genetics System**. *The Plant Cell*

2002, **14**(12):2985-2994.

25.   Sussman MR, Amasino RM, Young JC, Krysan PJ, Austin-Phillips S: **The Arabidopsis**

knockout facility at the University of Wisconsin-Madison. *Plant Physiology* 2000,

124(4):1465-1467.

26.   Li Y, Rosso MG, Viehoever P, Weisshaar B: **GABI-Kat SimpleSearch: an Arabidopsis**

thaliana T-DNA mutant database with detailed information for confirmed insertions. *Nucleic*

Acids Research 2007, **35**(Database issue):D874-D878.

27.   Kleinboelting N, Huep G, Weisshaar B: **Enhancing the GABI-Kat Arabidopsis thaliana T-**

DNA Insertion Mutant Database by Incorporating Araport11 Annotation. *Plant and Cell*

Physiology 2017, **58**(1):e7.

28. Cheng CY, Krishnakumar V, Chan A, Thibaud-Nissen F, Schobel S, Town CD: **Araport11: a**

complete reannotation of the Arabidopsis thaliana reference genome. *The Plant Journal*

2017, **89**(4):789-804.

29. Lamesch P, Berardini TZ, Li D, Swarbreck D, Wilks C, Sasidharan R, Muller R, Dreher K,

Alexander DL, Garcia-Hernandez M *et al*: **The Arabidopsis Information Resource (TAIR):**

improved gene annotation and new tools. *Nucleic Acids Research* 2012, **40**(Database

issue):D1202-D1210.

30. Huep G, Kleinboelting N, Weisshaar B: **An easy-to-use primer design tool to address**

paralogous loci and T-DNA insertion sites in the genome of Arabidopsis thaliana. *Plant*

Methods 2014, **10**:28.

31. Vukasinovic N, Cvrckova F, Elias M, Cole R, Fowler JE, Zarsky V, Synek L: **Dissecting a**

hidden gene duplication: the Arabidopsis thaliana SEC10 locus. *PLoS ONE* 2014,

9(4):e94077.

32. Pucker B, Holtgräwe D, Stadermann KB, Frey K, Huettel B, Reinhardt R, Weisshaar B: **A**

chromosome-level sequence assembly reveals the structure of the Arabidopsis thaliana Nd-

1 genome and its gene set. *PLoS One* 2019, **14**(5):e0216233.

33. Nacry P, Camilleri C, Courtial B, Caboche M, Bouchez D: **Major chromosomal**

rearrangements induced by T-DNA transformation in Arabidopsis. *Genetics* 1998,

149(2):641-650.

34. Tax FE, Vernon DM: **T-DNA-associated duplication/translocations in Arabidopsis.**

Implications for mutant analysis and functional genomics. *Plant Physiology* 2001,

126(4):1527-1538.

35.   Wei FJ, Kuang LY, Oung HM, Cheng SY, Wu HP, Huang LT, Tseng YT, Chiou WY, Hsieh-

Feng V, Chung CH *et al*: **Somaclonal variation does not preclude the use of rice**

transformants for genetic screening. *The Plant Journal* 2016, **85**(5):648-659.

36.   Gang H, Liu G, Zhang M, Zhao Y, Jiang J, Chen S: **Comprehensive characterization of T-**

DNA integration induced chromosomal rearrangement in a birch T-DNA mutant. *BMC*

Genomics 2019, **20**(1):311.

37.   Krizkova L, Hrouda M: **Direct repeats of T-DNA integrated in tobacco chromosome:**

characterization of junction regions. *The Plant Journal* 1998, **16**(6):673-680.

38.   Ulker B, Li Y, Rosso MG, Logemann E, Somssich IE, Weisshaar B: **T-DNA-mediated**

transfer of Agrobacterium tumefaciens chromosomal DNA into plants. *Nature Biotechnology*

2008, **26**(9):1015-1017.

39.   Seagrist JF, Su SH, Krysan PJ: **Recombination between T-DNA insertions to cause**

chromosomal deletions in Arabidopsis is a rare phenomenon. *PeerJ* 2018, **6**:e5076.

40.   Schmidt C, Schindele P, Puchta H: **From gene editing to genome engineering: restructuring**

plant chromosomes via CRISPR/Cas. *aBIOTECH* 2020, **1**(1):21-31.

41.   Pellestor F, Gatinois V: **Chromoanagenesis: a piece of the macroevolution scenario**.

Molecular Cytogenetics 2020, **13**:3.

42.   Li X, Zhang R, Patena W, Gang SS, Blum SR, Ivanova N, Yue R, Robertson JM, Lefebvre

PA, Fitz-Gibbon ST *et al*: **An Indexed, Mapped Mutant Library Enables Reverse Genetics**

Studies of Biological Processes in Chlamydomonas reinhardtii. *The Plant Cell* 2016,

28(2):367-387.

43.   Inagaki S, Henry IM, Lieberman MC, Comai L: **High-Throughput Analysis of T-DNA Location**

and Structure Using Sequence Capture. *PLoS One* 2015, **10**(10):e0139672.

44. Jiang N, Lee YS, Mukundi E, Gomez-Cano F, Rivero L, Grotewold E: **Diversity of genetic**

lesions characterizes new Arabidopsis flavonoid pigment mutant alleles from T-DNA

collections. *Plant Science* 2020, **291**:110335.

45. Michael TP, Jupe F, Bemm F, Motley ST, Sandoval JP, Lanz C, Loudet O, Weigel D, Ecker

JR: **High contiguity Arabidopsis thaliana genome assembly with a single nanopore flow cell.**

Nature Communications 2018, **9**(1):541.

46. Jorgensen R, Snyder C, Jones JDG: **T-DNA is organized predominantly in inverted repeat**

structures in plants transformed with Agrobacterium tumefaciens C58 derivatives. *Molecular*

Genetics and Genomics 1987, **207**(2-3):471-477.

47. Wu H, Sparks CA, Jones HD: **Characterisation of T-DNA loci and vector backbone**

sequences in transgenic wheat produced by Agrobacterium-mediated transformation.

Molecular Breeding 2006, **18**(3):195-208.

48. Rajapriya V, Kannan P, Sridevi G, Veluthambi K: **A rare transgenic event of rice with**

Agrobacterium binary vector backbone integration at the right T-DNA border junction. *J*

Plant Biochem Biot 2021.

49. Zambryski P, Depicker A, Kruger K, Goodman HM: **Tumor induction by Agrobacterium**

tumefaciens: analysis of the boundaries of T-DNA. *Journal of Molecular and Applied*

Genetics 1982, **1**(4):361-370.

50. Huang CY, Ayliffe MA, Timmis JN: **Direct measurement of the transfer rate of chloroplast**

DNA into the nucleus. *Nature* 2003, **422**(6927):72-76.

51. Bock R: **The give-and-take of DNA: horizontal gene transfer in plants.** *Trends in Plant*

Science 2009, **15**(1):11-22.

52.   Feldmann KA: **T-DNA insertion mutagenesis in** *Arabidopsis*: **mutational spektrum**. *The Plant*

969   *Journal* 1991, **1**(1):71-82.

970   53.   Curtis MJ, Belcram K, Bollmann SR, Tominey CM, Hoffman PD, Mercier R, Hays JB:

971   **Reciprocal chromosome translocation associated with TDNA-insertion mutation in**

972   **Arabidopsis: genetic and cytological analyses of consequences for gametophyte**

973   **development and for construction of doubly mutant lines**. *Planta* 2009, **229**(4):731-745.

974   54.   Bonhomme S, Horlow C, Vezon D, de Laissardiere S, Guyon A, Ferault M, Marchand M,

975   Bechtold N, Pelletier G: **T-DNA mediated disruption of essential gametophytic genes in**

976   **Arabidopsis is unexpectedly rare and cannot be inferred from segregation distortion alone**.

977   *Molecular Genetics and Genomics* 1998, **260**(5):444-452.

978   55.   Ruprecht C, Carroll A, Persson S: **T-DNA-induced chromosomal translocations in feronia**

979   **and anxur2 mutants reveal implications for the mechanism of collapsed pollen due to**

980   **chromosomal rearrangements.** *Molecular Plant* 2014, **7**(10):1591-1594.

981   56.   Schmidt C, Fransz P, Ronspies M, Dreissig S, Fuchs J, Heckmann S, Houben A, Puchta H:

982   **Changing local recombination patterns in Arabidopsis by CRISPR/Cas mediated**

983   **chromosome engineering**. *Nature Communications* 2020, **11**(1):4418.

984   57.   Spealman P, Burrell J, Gresham D: **Inverted duplicate DNA sequences increase**

985   **translocation rates through sequencing nanopores resulting in reduced base calling**

986   **accuracy**. *Nucleic Acids Research* 2020, **48**(9):4940-4945.

987   58.   Siadjeu C, Pucker B, Viehöver P, Albach DC, Weisshaar B: **High Contiguity De Novo**

988   **Genome Sequence Assembly of Trifoliate Yam (Dioscorea dumetorum) Using Long Read**

989   **Sequencing.** *Genes* 2020, **11**(3):E274.

990   59.   Belmann P, Fischer B, Kruger J, Prochazka M, Rasche H, Prinz M, Hanussek M, Lang M,

991   Bartusch F, Glassle B *et al*: **de.NBI Cloud federation through ELIXIR AAI**. *F1000Research*

992   2019, **8**:842.

993   60.   Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM: **Canu: scalable and**

994   **accurate long-read assembly via adaptive k-mer weighting and repeat separation**. *Genome*

995   *Research* 2017, **27**(5):722-736.

996   61.   Pucker B, Ruckert C, Stracke R, Viehover P, Kalinowski J, Weisshaar B: **Twenty-Five Years**

997   **of Propagation in Suspension Cell Culture Results in Substantial Alterations of the**

998   **Arabidopsis Thaliana Genome**. *Genes* 2019, **10**(9):671.

62.   Vaser R, Sović I, Nagarajan N, Šikić M: **Fast and accurate de novo genome assembly from**

long uncorrected reads. *Genome Research* 2017, **27**(5):737-746.

63.   Pucker B, Holtgräwe D, Rosleff Sörensen T, Stracke R, Viehöver P, Weisshaar B: **A De**

Novo Genome Sequence Assembly of the Arabidopsis thaliana Accession Niederzenz-1

Displays Presence/Absence Variation and Strong Synteny. *PLoS ONE* 2016,

11(10):e0164321.

64.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ: **Basic local alignment search tool**.

Journal of Molecular Biology 1990, **215**(3):403-410.

65.   Tang H, Zhang X, Miao C, Zhang J, Ming R, Schnable JC, Schnable PS, Lyons E, Lu J:

ALLMAPS: robust scaffold ordering based on multiple maps. *Genome Biology* 2015, **16**:3.

66.   Li H: **Minimap2: pairwise alignment for nucleotide sequences**. *Bioinformatics* 2018,

34(18):3094-3100.

67.   Quinlan AR, Hall IM: **BEDTools: a flexible suite of utilities for comparing genomic features**.

Bioinformatics 2010, **26**(6):841-842.

68.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL:

1014   BLAST+: architecture and applications. *BMC Bioinformatics* 2009, **10**:421.

1015   69.   Seqtk: a fast and lightweight tool for processing FASTA or FASTQ sequences

1016   [https://github.com/lh3/seqtk]

1017   70.   Rice P, Longden I, Bleasby A: EMBOSS: The European Molecular Biology Open Software

1018   Suite. *Trends in Genetics* 2000, **16**(6):276-277.

1019   71.   Li H: Minimap and miniasm: fast mapping and de novo assembly for noisy long sequences.

1020   *Bioinformatics* 2016, **32**(14):2103-2110.

1021   72.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin

1022   R: The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 2009, **25**(16):2078-

1023   2079.

1024   73.   Pucker B, Brockington SF: Genome-wide analyses supported by RNA-Seq reveal non-

1025   canonical splice sites in plant genomes. *BMC Genomics* 2018, **19**(1):980.

1026   74.   Katoh K, Standley DM: MAFFT multiple sequence alignment software version 7:

1027   improvements in performance and usability. *Molecular Biology and Evolution* 2013,

1028   **30**(4):772-780.

1029