Complementary Phenotyping of Maize Root Architecture by Root Pulling Force and X-Ray Computed Tomography

Mon-Ray Shao[1], Ni Jiang[1], Mao Li[1], Anne Howard[2], Kevin Lehner[2], Jack L. Mullen[2], Shayla L. Gunn[1], John K. McKay[2], Christopher N Topp[1]

[1] Donald Danforth Plant Science Center, Saint Louis, MO, USA
[2] Department of Agricultural Biology, Colorado State University, Fort Collins, CO 80523, USA

ABSTRACT

The root system is critical for the survival of nearly all land plants and a key target for improving abiotic stress tolerance, nutrient accumulation, and yield in crop species. Although many methods of root phenotyping exist, within field studies one of the most popular methods is the extraction and measurement of the upper portion of the root system, known as the root crown, followed by trait quantification based on manual measurements or 2D imaging. However, 2D techniques are inherently limited by the information available from single points of view. Here, we used X-ray computed tomography to generate highly accurate 3D models of maize root crowns and created computational pipelines capable of measuring 71 features from each sample. This approach improves estimates of the genetic contribution to root system architecture, and is refined enough to detect various changes in global root system architecture over developmental time as well as more subtle changes in root distributions as a result of environmental differences. We demonstrate that root pulling force, a high-throughput method of root extraction that provides an estimate of root biomass, is associated with multiple 3D traits from our pipeline. Our combined methodology can therefore be used to calibrate and interpret root pulling force measurements across a range of experimental contexts, or scaled up as a stand-alone approach in large genetic studies of root system architecture.

INTRODUCTION

In maize, the entirety of primary, seminal, lateral, crown, and brace roots together form a complex architecture which controls the plant's ability to effectively acquire water, scavenge nutrients, and resist lodging (Hochholdinger, 2009). As a result, root growth and development are fundamental to overall plant development and competitiveness (Lynch, 1995), and several prominent large-effect, loss-of-function mutants in cereal seedling root development have been identified and reviewed previously (Hochholdinger et al., 2017; Bray and Topp, 2018). However, root system architecture of mature, field-grown plants at the quantitative level has been understudied and underutilized due to the relative difficulty in obtaining measurements, with significant tradeoffs intrinsic to any particular phenotyping method (Pauli et al., 2016; Topp et al., 2016). Nevertheless, because root growth is highly plastic and affected by environmental conditions such as substrate moisture and texture (Sharp, 2004; Rich and Watt, 2013), field-based studies are valuable despite their challenges.

In its simplest form, root phenotyping of crop species such as maize or rice can be performed by manual measurement of a limited set of amenable traits, such as root biomass, length, width, or the growing angle, either in soil or soil-free conditions. Currently known genes controlling quantitative root system architecture traits in rice were identified using such measurements, including *PSTOL1* (Gamuyao et al., 2012), *DRO1* (Uga et al., 2013), and a recent *DRO1 homolog* (Kitomi et al., 2020). In field conditions, additional techniques for quantifying roots exist, such as the use of minirhizotrons, soil core sampling, and measuring of root pulling force (Holbert and Koehler, 1924; Bohm and Böhm, 1979; Samson and Sinclair, 1994; Wasson et al., 2014). Historically, root pulling force (RPF) has been useful as a field assay because of its simplicity and potential for scalability, and has been applied to both monocots and dicots (Ortman et al., 1968; O'Toole and Soemartono, 1981; Donovan et al., 1982; Bailey et al., 2002; Landi et al., 2002; Fletcher et al., 2015). RPF is generally correlated with greater root biomass and branching, but more nuanced interpretations and its relationship with recently tractable architectural measurements have yet to be established.

More intricate phenotyping of root system architecture can be performed upon two-dimensional images of either field-excavated root crowns, or young gel-media grown root systems, followed by analysis with specialized software (Le Bot et al., 2010; Lobet et al., 2011; Galkovskyi et al., 2012; Colombi et al., 2015; Das et al., 2015; Symonova et al., 2015; Delory et al., 2018; Seethepalli et al., 2020). Such methods have been used to quantify root system architecture in diverse crops such as maize, wheat, rice, and cowpea (Bucksch et al., 2014; Canè et al., 2014; Burridge et al., 2017; Wedger et al., 2019). However, 2D-based measurements have a limitation in that images are typically taken from only one or two camera perspectives, with information lost from roots occluding each other in the image.

As a result, interest and capacity towards three-dimensional root phenotyping has been increasing, driven in part by technical advances and interdisciplinary approaches (Morris et al., 2017). For example, young cereal plants grown in a gel-based media can be imaged over a 360° rotation, allowing digital reconstruction in 3D and high-throughput feature extraction (Iyer-

Pascuzzi et al., 2010; Clark et al., 2011). By scaling this technique to mapping populations,
studies have identified new univariate or multivariate root QTLs, demonstrating the value of
high-throughput and high-information-content trait capture for dissection of plant architecture
(Topp et al., 2013; Zurek et al., 2015). Other 3D-based solutions include the use of X-ray
computed tomography (XRT), which is capable of imaging any plant structure, including roots
within soil based upon physical density properties (Mairhofer et al., 2012; Mooney et al., 2012;
Bao et al., 2014; Rogers et al., 2016; Duncan et al., 2019; Li et al., 2019; Li et al., 2020;
Helliwell et al.). While XRT has been applied to plant physiology in some form for nearly two
decades, instrument accessibility and technical limitations typically restrict its use to small plant
structures, low throughput, and/or limited fields of view.
98
Here we integrate two protocols, first sampling via RPF and washing mature, lignified, field-
grown maize root crowns, followed by imaging via XRT and trait quantification for over 290 roots
across multiple field seasons. By imaging the roots absent of soil or other media, scanning and
segmentation times were significantly reduced such that replication across two environments
and/or two time points was possible. We extracted up to 71 3D features for each root crown
sample, including up to 65 traits with significant variation between genotypes, as well as root
shape or distributional traits, which showed differences between experimental contexts. The
median broad-sense heritability across all traits ranged from 0.23 to 0.56, depending on the
germplasm and conditions. Finally, we examined covariance between 3D traits and RPF values
to identify correlations between high-resolution phenomics and high-throughput field data. This
study illustrates the contributions of both phenotyping approaches, and provides insights into
the root architectural attributes that influence RPF, which can be used for the mapping of root
traits, multi-environment studies, and crop breeding.
112
113
114

MATERIALS AND METHODS

Plant Germplasm and Experimental Design

All plants were grown at the Colorado State University Agricultural Research Development &
Education Center in Fort Collins, CO, USA (40.649 N, -105.000 W) in 2017 and 2018. The
Genomes 2 Fields (G2F) germplasm (https://www.genomes2fields.org/) was planted in May
2017 in a split-plot design with full irrigation or limited irrigation (drought) treatments, with two
field replicates per treatment for a total of 1060 plots total. Prior to planting, the field was
fertilized with nitrogen at 65 lbs per acre. From 260 genotypes planted as part of a large field
trial, 30 genotypes were selected for root imaging in this study across both irrigation treatments
(Supplemental Table 1), for a total sample size of N = 107 roots.
127
The Shoot Apical Meristem (SAM) diversity panel (Leiboff et al., 2015) along with 11 hybrid and
4 inbred check lines were planted in May 2018 using a split-plot design with full irrigation or
limited irrigation (drought) treatments, with three field replicates per treatment. Prior to planting,
the field was fertilized with nitrogen at 190 lb per acre. From 390 genotypes planted as part of a

large field trial, 20 genotypes were subsampled for root imaging in this study across both
irrigation treatments (Supplemental Table 1). Root systems were harvested at 9 weeks after
planting (time point 1) and again at 16 weeks after planting (time point 2), for a total sample size
of $N = 187$ roots.
136
In both the G2F and SAM experiments, each plot consisted of two 12-foot rows with 30-inch
spacing between rows and 9-inch spacing between plants within rows. The irrigated treatments
received approximately 1 inch of water per week, while the drought treatments were irrigated
until well established (approx. 5 weeks after planting) and then received only natural
precipitation (103.8 mm and 69.9 mm in the 2017 and 2018 growing seasons, respectively),
except at the root harvesting when it also received irrigation to homogenize the root harvesting
process.
144

145 **Field Phenotyping and 2D Root Imaging**

146
The protocol used for root pulling and harvesting was similar to that in (Fletcher et al., 2015).
Briefly, all plants were irrigated 24 hours prior to sampling to homogenize soil conditions at root
harvest. Maize plants were tied at the base of the stem, just above the root crown, with a rope
attached to a dynamometer. The root system was extracted from the soil by vertical manual
pulling, with the required force (Kg) needed measured using a hand-held Imada DS2 digital
force gauge (Imada Inc., Northbrook, IL, USA). Within each field treatment (full vs limited
irrigation), two roots per genotype were harvested from the G2F population and an average of 4
roots per genotype (across two time points) were harvested for the SAM population. After
pulling, root samples were washed to remove all remaining soil and allowed to dry before
imaging.
157
158   Roots from the G2F 2017 experiment were also imaged in 2D using a photography station
159   equipped with a Sony a7 II mirrorless camera. Roots were placed horizontally on a flat surface
160   with a black background, and the resulting images were then cropped and analyzed using DIRT
161   (Das et al., 2015).
162

163 **3D Root Imaging and Feature Extraction**

164
For 3D phenotyping in the G2F and SAM populations, root samples were clamped at the stem
with a small vise and imaged using a North Star X5000 X-ray system (North Star Imaging, MN,
USA). The sample was continuously imaged while rotated using the North Star efX-CT software
system, generating 1800 radiographs per sample (approx. 3 minutes). To provide an internal
calibration of the image geometry, a fixed standard (15 mm large tool, North Star Imaging) was
imaged with each sample batch. The radiographs were then reconstructed using efX-CT and
exported as an unadjusted RAW volume, resulting in a voxel size of 109-113 µm depending on
the sample batch. For each sample, the RAW volume was converted to 2D slices using the
custom Python script *raw2img*. The slices were then thresholded and binarized using the
custom script *batch-segmentation*, which subsequently converted each sample to a 3D point
cloud and quantified 19 root traits adapted from (Galkovskyi et al., 2012). Finally, each sample

was converted to a 3D skeleton using the custom script *batch-skeleton*, which then quantified
an additional root 52 traits. The raw phenotype data is available in Supplemental File 1.
178
A list and description of root features measured using *batch-segmentation* and *batch-skeleton*
are available in Supplemental Table 2. Mean, standard deviation, skewness, kurtosis, energy,
entropy, and smoothness from the distributions of root biomass (volume), convex hull, and
solidity were calculated using the method described in Malik and Baharudin, 2012. Fractal
dimension, which measures the degree to which root subsections approximate a smaller copy of
the whole root crown, was estimated by taking the 2D projection of the 3D volume, then
calculated using a similar approach to that described in Grift et al., 2011. DensityS features are
computationally similar to plant compactness traits described in Yang et al., 2014. Scripts used
for image processing and feature extraction are available at https://github.com/Topp-Roots-Lab/
188
189 **Statistical Analysis**
190
All downstream (i.e. post feature extraction) analysis was performed in the R statistical
computing environment. Initially, principal component analysis using all 71 3D roots traits was
used to identify large outliers, leading to the removal of 2 samples in the G2F 2017 data and 3
samples in the SAM 2018 data. Additionally, for all univariate analysis, outliers within each trait
were identified and omitted if they were beyond the 1st quartile minus 1.5 * interquartile-range
or the 3rd quartile plus 1.5 * interquartile-range.
197
After univariate outlier removal, analysis of variance (ANOVA) was performed for each trait
using the *car* package (Fox and Weisberg, 2018). Subsequently, the ANOVA p-values were
adjusted using the Benjamini-Hochberg method. Individual two-sample comparisons as seen in
boxplots were performed using Mann-Whitney U tests. Correlations between root traits were
calculated using Spearman's correlation coefficient. Linear regressions were performed using
the *lm* function in R.
204
Variance components were estimated by using the *lme4* package (Bates et al., 2015) to fit the
linear model $Y_{ijk} \sim G_i + E_j + (G*E)_{ij} + e_{ijk}$, where Y is the phenotypic value, G_i is the i^{th} genotype,
E_j is the j^{th} environment, $(G*E)_{ij}$ is the interaction between the i^{th} genotype and the j^{th}
environment, and e_{ijk} is the residual error of the k^{th} sample from the i^{th} genotype and j^{th}
environment. Broad-sense heritability was calculated using the equation
$H^2 = \frac{\sigma_G}{\sigma_G + \sigma_{GxE}/e + \sigma_{residual}/re}$ where σ_G is the estimated phenotypic variance due to genotype,
σ_{GxE} is the estimated phenotypic variance due to genotype x environment, $\sigma_{residual}$ is the residual
variance, e is the number of environments, and re is the average number of biological replicates
per genotype across both environments (Nyquist and Baker, 1991). This heritability estimator is
optimized as a predictor of the response to selection. For the SAM 2018 data, variance
components and broad-sense heritability were calculated separately for the two time points.
216
Principal component analysis (PCA) of the 3D root data within the G2M and SAM experiments
was performed using the base R *prcomp* function. For PCA-LDA, PCA was performed upon
each genotype subset, and the number of principal components required to explain 90% of the

trait variance was used as inputs into the LDA function from the *MASS* package (Venables and
Ripley, 2002).

The *randomForest* (Liaw et al., 2002) and *caret* (Kuhn and Others, 2008) R packages were
used for random forest classification, with *mtry* and *ntree* parameters found using a grid search
approach between every combination of *mtry* between 1 to 20 and *ntree* values of 500, 1000,
2500, 5000. Parameters giving the best accuracy were kept, as calculated by 10-fold cross
validation repeated 3 times. From the final random forest models, the proximity matrix was
calculated and non-metric multidimensional scaling was used to visualize the distances between
samples.

RESULTS

Field and 3D Phenotyping Capture Variation in Maize Root System Architecture

In each of two field seasons, we sampled 50 maize genotypes (30 from the G2F panel and 20
from the SAM panel) that were grown under two different irrigation treatments, providing two
environments in terms of soil moisture. At the designated time point(s) for sampling (see
Methods), root crowns were excavated by tying the base of the stem with rope to a digital force
gauge, and manually placing a vertical force on the plant until the root crown was ruptured and
lifted free from the soil. The force gauge attached to the root system measured the kilogram of
force required for this process, also known as the root pulling force (RPF).

Field-pulled root samples, which maintain their 3D structure due to lignification, were washed
clean and subsequently imaged using a Northstar X5000 X-Ray computed tomography system
(Figure 1, Supplemental Figure 1A). Scans were then exported as vertical (y-axis) image slices,
thresholded using an automated algorithm, and converted to a skeletonized image and a point
cloud image for trait analysis (Supplemental Figure 1B-D). These were analyzed for 19 3D traits
using an established 3D-analysis pipeline (Bray and Topp, 2018) as well as 52 additional traits
predominantly focused on 3D root distribution traits using a newly developed series of feature
extraction tools. In total, 71 traits from the 3D volumes were extracted and used for analysis
(Supplemental Table 2).

In the G2F data set, root samples were also photographed for 2D image analysis via DIRT
using recommended protocols (Supplemental Figure 1E) (Das et al., 2015). Correlations
between the most directly comparable 2D and 3D traits were as we expected - for example, 2D
area and 3D surface area had a Pearson correlation coefficient of 0.754.

To assess the degree to which traits derived from field-pulled root crown samples would
respond to selection, we estimated broad-sense heritability (H^2) in the G2F experiment for each
3D and 2D trait, as well as for RPF (Supplemental Figure 2). Traits related to overall root crown
size showed similar H^2 values between 3D measurements (e.g. 3D surface area $H^2 = 0.47$) and

2D measurements (e.g. 2D area H^2 = 0.44). The traits with highest heritability, however, were
3D-derived maximum root count (H^2 = 0.76) and average root radius (H^2 = 0.74), illustrating
where 3D root phenotyping is particularly adept. Among 2D traits, high heritability did not
necessarily result in high association with RPF, although some traits such as 2D area had a
strong positive association (Supplemental Figure 2E-F). Nevertheless, depending on the
experimental conditions and amount of replication, it is probable that many root traits - though
computationally extractable - have questionable value due to high background noise and
sensitivity to sampling variation. In total, for example, 19 3D traits and 33 2D traits had
calculated H^2 values of less than 0.05; therefore, both 3D and 2D root traits must be screened
and evaluated for a given data set before drawing conclusions. Reassuringly, however, RPF
itself had a H^2 value of 0.67 in the G2F experiment, which is remarkably high for a physical field-
based root measurement, even when compared to the architectural traits measured from 3D
images.
277
In the SAM experiment, broad-sense heritability for 3D traits were calculated separately within
each time point (Supplemental Figure 3A-C). In general, H^2 values here were higher than in the
G2F experiment, which reflects a combination of the field conditions, genetic variation, and
sample sizes. As an average across both time points, root crown width ("HorEqDiameter", mean
H^2 = 0.81), fractal dimension top view (mean H^2 = 0.81), maximum root count (mean H^2 = 0.80),
convex hull volume (mean H^2 = 0.79), and surface area (mean H^2 = 0.79) were among the
most heritable traits, although the individual performance of these traits fluctuated depending on
the time point. However, the average heritability across all 3D traits was only slightly higher at
first time point (0.510) than at second time point (0.505), suggesting that heritability of most
roots traits is relatively static over this time span. One interesting exception to this is RPF itself,
which had a H^2 value of only 0.59 at the first time point, but significantly increased to a H^2 value
of 0.85 by the second time point. This indicates that RPF measurements taken later in the plant
life cycle may be more informative and reliable for the purposes of distinguishing genotypic
differences in maize root system architecture, as well as for breeding. Overall, however, traits
with higher average heritability across time points tended to also have a greater correlation in
measurements between time points (Supplemental Figure 3D).
294
Focusing on 3D root traits and RPF, we subsequently wanted to examine whether genotype and
environment effects were significant factors on a trait-by-trait basis (Figure 2). Using analysis of
variance (ANOVA), in the G2F experiment we detected 21 root traits where genotype had a
significant effect, and 35 traits where the environment (irrigation regime) had a significant effect
(Figure 2B, Supplemental Figure 4A, Supplemental Figure 5). Root traits affected by both
genotype and environment include RPF, average root radius, median/maximum number of
roots, convex hull skewness, and solidity in several regions along the middle of the root crown.
Non-parametric tests for differences between environments confirmed that RPF, average root
radius, and convex hull volume, for example, were higher in the high irrigation environment,
whereas solidity was higher in the low irrigation environment (Figure 2C). These meet
expectations of soil moisture effects on root system architecture (e.g. more expansive growth
under higher moisture availability), providing confidence to our 3D phenotyping.
307

In the SAM experiment, the situation was somewhat reversed: in support of the overall higher
trait heritability, a remarkable 65 root traits had a significant effect from genotype, but only 17
traits had a significant effect from environment, while 37 traits had a significant effect from time
point (Figure 2B, Supplemental Figure 4A, Supplemental Figure 6). Traits such as RPF, surface
area, volume, root crown depth, fractal dimension side view, and biomass distribution skewness
were affected by all three variables. Again, non-parametric tests for differences in RPF, volume,
and fractal dimension side, for example, confirmed the impacts of environment and time point as
detected by ANOVA (Figure 2D). The somewhat divergent trends in 3D root phenotypes
between the G2F and SAM experiments, however, indicate that additional generalizations about
root system architecture and how growth plasticity relates to it may be difficult to come by, as
root variation is highly dependent on the experimental conditions and/or genotypes. Indeed,
although the sample sizes here precluded strong statistical power to test genotype-environment
interactions using ANOVA, variance component analysis suggests that such interactions may
have a significant influence on a number of root architecture traits (Supplemental Figure 2B;
Supplemental Figure 3B-C).
323
Next, we applied supervised multivariate classification methods to determine which traits were
most closely associated with differences in genotype, environment, or time (Supplemental Table
3). Because of the high number of genotypes (18 in the G2F set and 16 in the SAM set, after
filtering for genotypes with the least missing data), in both experiments the data was split into
every possible combination of three genotypes, generating 816 different genotype combinations
in the G2F set and 560 different genotype combinations in the SAM set. We performed PCA-
LDA for genotype classification upon each three-genotype data subset, in each case using the
minimum number of principal components to explain 90% of the variance (5-7 PC's with a
median of 6 in G2F data; 9-13 PC's with a median of 11 in SAM data) as the inputs for LDA
(Figure 3A-B). Across all genotype combination subsets, the average classification accuracy
using leave-one-out cross-validation was 54.6% in the G2F and 67.2% in the SAM, both
significantly higher than the 1/3 expected by random chance, particularly considering the
realistic possibility that numerous genotypes may in actuality be phenotypically similar.
337
We examined what traits were most important towards PCA-LDA classification across all
genotype combinations (Supplemental Figure 7-8). In both the G2F and SAM populations,
maximum root count, average root radius, and specific root length tended to be very important
for genotype discrimination. Additionally, the median root count, number of root tips, elongation,
and average edge length were important in genotypic classification among the G2F population,
while several root solidity and density traits were important in genotypic classification among the
SAM population. Although RPF was not among the top traits for genotypic classification, it was
still well above average, ranking 12th overall in the G2F set and 19th overall in the SAM set.
Interestingly, traits related to overall root size such as volume or surface area did not seem to
be important factors overall in discriminating between genotypes, suggesting that these metrics,
although intuitive and undoubtedly important in other contexts, are by themselves insufficient to
distinguish between multiple and often subtly distinct genotypes, highlighting the need for the
more comprehensive phenotyping described here.
351

To evaluate the overall effect of the environment (influential in the G2F experiment) and time
(highly influential in the SAM experiment), we performed random forest classification to
distinguish between the two possible levels of each variable upon root system architecture. For
these classifications we included all genotypes, which increased the sample size for each
model. Using 10-fold cross validation, the best model parameters resulted in a classification
accuracy of 81.0%, indicating that while the environment had an effect which was detectable
using classification techniques, the contrasting irrigation regimes were not so dramatic as to
result in a shift in root system architecture obvious across every sample (Figure 3C).
Nevertheless, changes in density and solidity distributions, as well as root crown depth, were
the most distinguishing features, with RPF being less important (Figure 3D). Here, the
importance of solidity distributions in the upper half of the root crown (including SolidityVHist 02-
11) is consistent with ANOVA analysis (Supplemental Figure 4A); in particular, the low broad-
sense heritability of SolidityVHist 05-10, coupled with disproportionately high variance from
environment and genotype-environment effects, indicates that these are more determined by
environmental factors than by genetics in this experiment (Supplemental Figure 2A-B). On the
other hand, DensityS5 (a measure of relative compactness), which had a moderately high
heritability in this experiment, is still important for distinguishing the effect of environment,
suggesting that this trait is strongly affected by both genotype and environment.
370
For classifying roots based on time point in the SAM data, using 10-fold cross validation the
best model parameters resulted in a random forest classification accuracy of 78.6% (Figure 3E),
which was reasonable when considering that samples across both environmental conditions
were included. Here, differences in convex hull volume, volume, depth, root crown width, and
solidity distribution were the most distinguishing features, with RPF closely behind these and
other important traits (Figure 3F). Furthermore, solidity distribution features at the very top and
bottom of the root crown (SolidityVHist 01 and 17-19) appear to be relevant. Again, these
results are unsurprising given the expectation of increasing root crown size over time, and are
largely consistent with ANOVA results on time point effects upon these traits (Figure 2B).
380
381 ***Root Architecture Relationships and Correspondence to Root Pulling Force***
382
Root pulling force has been used historically and recently as a proxy for root biomass and root
volume. Nevertheless, to have additional and more detailed information on the architectural
changes that RPF measures would increase its utility as a field assay. We first calculated
correlations between RPF and 3D root phenotype across all measured samples, irrespective of
genotype and environment, or time point in the case of the SAM data. In both experiments, RPF
was most correlated with root volume, fractal dimension, surface area, total root length, root
crown width, number of bifurcating clusters, and number of root tips. (Figure 4A-B,
Supplemental Table 4). Traits negatively correlated with RPF were generally weaker and less
consistent between the two experiments, but did include convex hull energy (a measure of root
system uniformity) and Density S5 in both cases.
393
Associations between root pulling force and root system architecture traits are most useful if
they are not only significantly correlated, but also exhibit a close linear relationship. Regression

analysis between RPF and positively correlated 3D architecture traits (as observed in the G2F
experiment), such as fractal dimension and surface area, showed a reasonably good fit (Figure
4C-E). In contrast, there was a relatively poor fit with convex hull kurtosis, the most negatively
correlated trait (Figure 4F). These may in part be due to sampling error or noise, but also
because multiple root characteristics that may not be strongly correlated to each other
nevertheless each contribute to RPF in various ways. Nevertheless, the regression fit between
physical root biomass (i.e. root crown weight) in the SAM experiment and RPF or other
positively correlated 3D architecture traits was extremely high, while again relatively poor with
negatively correlated traits such as solidity in the upper root crown (Figure 4G-J). This high
goodness-of-fit was not a by-product of regression between two time points; rather, regression
between physical root biomass and these 3D traits remained high even when observing trends
and regressions within each time point (Supplemental Figure 4B-J). Furthermore, regression fit
was typically higher in time point 1 than in time point 2, which might be due to root crown traits
beginning to diverge in ways more independent of biomass, such as in architectural and spatial
orientation, which could nonetheless contribute to RPF.
411
To explore the degree to which trends across multiple traits may be associated with RPF, we
performed principal component analysis (PCA) from the G2F and SAM data using the 3D-based
root phenotypes alone (Supplemental Figure 9A, E). In the G2F data, RPF was more tightly
associated with principal component 2 (Supplemental Figure 9B), which was primarily
composed of traits related to overall size, i.e. surface area, volume, total root length, and
number of root tips, but also significantly composed of 3D biomass distribution traits
(Supplemental Figure 9I). On the other hand, in the SAM data, RPF was more tightly associated
with principal component 1 (Supplemental Figure 9F), which as with the G2F data was primarily
composed of traits related to overall size, including surface area, volume, and total root length,
and additionally fractional dimension side/top, but notably not of 3D biomass distribution traits
(Supplemental Figure 9J). Both PC1 and PC2 were statistically different between the two
environmental conditions in the G2F data and between the two time points in the SAM data
(Supplemental Figure 9C-H), but the differences between the G2F and SAM results here likely
derive from the fact that much of the phenotypic variation in the SAM data is greatly affected by
sampling time point, which an unsupervised method such as PCA does not distinguish.
427
428 As a whole, multiple analytical approaches corroborate the conclusion that distinct sets of root
429 traits are relevant depending on the germplasm, environment, and developmental time stage,
430 reinforcing the relevance of high-dimensional root phenotyping. This also demonstrates not only
431 the complexity of root architectures, but its propensity to change under different contexts and
432 genetic influences, and its ability to adapt to various conditions.
433
434
435

DISCUSSION

437
We have shown that using X-ray computed tomography, changes in 3D root architecture
between different treatments and conditions in the field can be measured in a biologically

interpretable way, while also with high precision and detail. For example, soil moisture
conditions affect the solidity of the root system, particularly in the mid-portion of the crown. In
contrast, changes over time influence not only the overall size of the root crown, but also solidity
in the upper and lower portions of the crown. Interesting, in both contexts, the depth (i.e., the
length of the vertical axis) of the root crown also was a distinguishing feature. This is likely
specific to root crowns excavated using the root pulling method, as under standard shovel
excavation, the root depth would be arbitrary. By using root pulling, however, the depth is a
function of the root system and the soil conditions, as these determine where the root crown
breaks and therefore encapsulates some information. Finally, in many instances the 3D
architectural measurements were easily sufficient to distinguish different maize genotypes,
primarily using an entirely different set of traits including specific root length, median/maximum
root count, average root radius, and so on.
452
To date, 2D imaging has been by far the most popular form of quantifying root system
architecture, whether in *Arabidopsis* grown on media plates, or root crowns excavated from the
field. While relatively straightforward and convenient, 2D imaging does not represent true root
system architecture in its natural form, and therefore may omit important information. More
recently, optical imaging platforms have been developed to perform 3D imaging of plants
growing in gel media (Clark et al., 2011; Topp et al., 2013; Jiang et al., 2019). Here, we present
a new approach to quantifying hundreds of field-excavated root crowns using X-ray CT, which is
typically restricted to very small sample sizes or reconstituted soils from pot experiments (Bao et
al., 2014). Our results suggest that 3D imaging and the root architectural traits derived from it
have higher heritability, and therefore may be more informative, than methods using 2D
imaging. Therefore, we anticipate that future studies and breeding efforts in quantitative root
system architecture will increasingly utilize 3D phenotyping.
465
Nonetheless, the significant overhead associated with 3D imaging and analysis of roots will be a
limitation to many researchers for the foreseeable future. We addressed this by making explicit
comparisons between high information content 3D phenotypes and root pulling force, which is
accessible and can be scaled to high throughput levels. RPF measurements had several
significant positive correlations with 3D architecture traits including volume, surface area, total
root length, number of roots, and fraction dimension. Indeed, fractal dimension was a
surprisingly powerful trait, not only being highly correlated with RPF, but also having high
heritability and contributing significantly to differentiating root system architecture over time. This
is additional evidence that fractal dimension of root systems (Tatsumi et al., 1989; Eghball et al.,
1993; Nielsen et al., 1997; Eshel, 1998; Grift et al., 2011), is indeed a useful feature for
quantifying maize root crowns under a variety of scenarios.
477
However, there remains sufficient sources of variance among features such that future
improvements could strengthen the relationship between the above traits and RPF values. For
example, we performed root pulling manually by hand, but a field robot or other form of
mechanical assistance may result in more consistent RPF measurements (Mayer, 2019).
Furthermore, while the fields were flooded just prior to root pulling to standardize the soil
moisture conditions at the time of RPF sampling, local heterogeneity in soil texture or

compactness may influence the measurements. This can be addressed in part by integrating
larger studies whereby spatial effects can be modeled, which likewise would be facilitated with
mechanization of the root pulling process. Finally, it should be noted that several traits not
measurable by our XRT system could contribute to RPF, including the abundance of root hairs
and variation in rhizosheath formation. Indeed, the fact that the heritability of RPF here
increased later in development suggests that it may be controlled by different genetic factors
over time, and therefore it may also be possible to select and breed for early and later root
phenotypes at least partially independently.
492
A better understanding of the exact nature and the potential interactions between the many root
traits investigated here, such as exactly how fractal dimension and root volume together affect
RPF, as well as the addition of more traits that could theoretically be calculated from 3D imaging
(such as those relating to topology), may lead to additional insight into the relationship between
root system architecture and RPF. Fully resolving this relationship would be particularly
beneficial for multi-environment phenotyping, which requires high sample sizes to which the
RPF method is well-suited for. Indeed, our study suggests that many more field-scale studies,
utilizing wide-ranging conditions and germplasms, will be needed to fully characterize and
understand quantitative root system architecture and genotype-environment interactions in
diverse plant species such as maize.
503
504
505

REFERENCES

Bailey PHJ, Currey JD, Fitter AH (2002) The role of root system architecture and root hairs in
promoting anchorage against uprooting forces in Allium cepa and root mutants of
Arabidopsis thaliana. Journal of Experimental Botany **53**: 333–340

**Bao Y, Aggarwal P, Robbins NE 2nd, Sturrock CJ, Thompson MC, Tan HQ, Tham C, Duan
L, Rodriguez PL, Vernoux T, et al** (2014) Plant roots use a patterning mechanism to
position lateral root branches toward available water. Proc Natl Acad Sci U S A **111**: 9319–
9324

Bates D, Maechler M, Bolker B (2015) Walker., S. Fitting linear mixed-effects models using
lme4. J Stat Softw **67**: 1–48

Bohm W, Böhm W (1979) Methods of studying root systems. Springer verlag

Bray AL, Topp CN (2018) The Quantitative Genetic Control of Root Architecture in Maize. Plant
Cell Physiol **59**: 1919–1930

Bucksch A, Burridge J, York LM, Das A, Nord E, Weitz JS, Lynch JP (2014) Image-based
high-throughput field phenotyping of crop roots. Plant Physiol **166**: 470–486

Burridge JD, Schneider HM, Huynh B-L, Roberts PA, Bucksch A, Lynch JP (2017)
Genome-wide association mapping and agronomic impact of cowpea root architecture.
Theor Appl Genet **130**: 419–431

Canè MA, Maccaferri M, Nazemi G, Salvi S, Francia R, Colalongo C, Tuberosa R (2014)
Association mapping for root architectural traits in durum wheat seedlings as related to
agronomic performance. Mol Breed **34**: 1629–1645

Clark RT, MacCurdy RB, Jung JK, Shaff JE, McCouch SR, Aneshansley DJ, Kochian LV
(2011) Three-dimensional root phenotyping with a novel imaging and software platform.
Plant Physiol **156**: 455–465

Colombi T, Kirchgessner N, Le Marié CA, York LM, Lynch JP, Hund A (2015) Next
generation shovelomics: set up a tent and REST. Plant Soil **388**: 1–20

Das A, Schneider H, Burridge J, Ascanio AKM, Wojciechowski T, Topp CN, Lynch JP,
Weitz JS, Bucksch A (2015) Digital imaging of root traits (DIRT): a high-throughput
computing and collaboration platform for field-based root phenomics. Plant Methods **11**: 51

Delory BM, Li M, Topp CN, Lobet G (2018) archiDART v3.0: A new data analysis pipeline
allowing the topological analysis of plant root systems. F1000Res **7**: 22

Donovan LS, Jui P, Kloek M, Nicholls CF (1982) AN IMPROVED METHOD OF MEASURING
ROOT STRENGTH IN CORN (Zea mays L.). Canadian Journal of Plant Science **62**: 223–
227

Duncan KE, Bray AL, Dowd TG, Topp CN (2019) Using 3D X-ray Microscopy to Study Crown
Root Development and Primary Root Tip Growth in Diverse Maize (Zea mays L.) Lines.
Microsc Microanal **25**: 1032–1033

Eghball B, Settimi JR, Maranville JW, Parkhurst AM (1993) Fractal analysis for
morphological description of corn roots under nitrogen stress. Agron J **85**: 287–289

Eshel A (1998) On the fractal dimensions of a root system. Plant Cell Environ **21**: 247–251

Fletcher RS, Mullen JL, Heiliger A, McKay JK (2015) QTL analysis of root morphology,
flowering time, and yield reveals trade-offs in response to drought in Brassica napus. J Exp
Bot **66**: 245–256

Fox J, Weisberg S (2018) An R Companion to Applied Regression. SAGE Publications

Galkovskyi T, Mileyko Y, Bucksch A, Moore B, Symonova O, Price CA, Topp CN, Iyer-
Pascuzzi AS, Zurek PR, Fang S, et al (2012) GiA Roots: software for the high throughput
analysis of plant root system architecture. BMC Plant Biol **12**: 116

Gamuyao R, Chin JH, Pariasca-Tanaka J, Pesaresi P, Catausan S, Dalid C, Slamet-Loedin
I, Tecson-Mendoza EM, Wissuwa M, Heuer S (2012) The protein kinase Pstol1 from
traditional rice confers tolerance of phosphorus deficiency. Nature **488**: 535–539

Grift TE, Novais J, Bohn M (2011) High-throughput phenotyping technology for maize roots.
Biosystems Eng **110**: 40–48

Helliwell JR, Sturrock CJ, Mairhofer S, Craigon J, Ashton RW, Miller AJ, Others The
emergent rhizosphere: imaging the development of the porous architecture at the root-soil
interface. Sci Rep 2017; 7 (1): 14875.

Hochholdinger F (2009) The Maize Root System: Morphology, Anatomy, and Genetics.

Handbook of Maize: Its Biology 145–160

Hochholdinger F, Yu P, Marcon C (2017) Genetic Control of Root System Development in
Maize. Trends Plant Sci. doi: 10.1016/j.tplants.2017.10.004

Holbert JR, Koehler B (1924) Anchorage and extent of corn root systems. J Agric Res **27**: 71–
78

Iyer-Pascuzzi AS, Symonova O, Mileyko Y, Hao Y, Belcher H, Harer J, Weitz JS, Benfey
PN (2010) Imaging and analysis platform for automatic phenotyping and trait ranking of
plant root systems. Plant Physiol **152**: 1148–1157

Jiang N, Floro E, Bray AL, Laws B, Duncan KE, Topp CN (2019) Three-Dimensional Time-
Lapse Analysis Reveals Multiscale Relationships in Maize Root Systems with Contrasting
Architectures. The Plant Cell **31**: 1708–1722

Kitomi Y, Hanzawa E, Kuya N, Inoue H, Hara N, Kawai S, Kanno N, Endo M, Sugimoto K,
Yamazaki T, et al (2020) Root angle modifications by the DRO1 homolog improve rice
yields in saline paddy fields. Proc Natl Acad Sci U S A **117**: 21242–21250

Kuhn M, Others (2008) Building predictive models in R using the caret package. J Stat Softw
28: 1–26

Landi P, Sanguineti MC, Darrah LL, Giuliani MM, Salvi S, Conti S, Tuberosa R (2002)
Detection of QTLs for vertical root pulling resistance in maize and overlap with QTLs for
root traits in hydroponics and for grain yield under different water regimes. Maydica **47**:
233–243

Le Bot J, Serra V, Fabre J, Draye X, Adamowicz S, Pagès L (2010) DART: a software to
analyse root system architecture and development from captured images. Plant Soil **326**:
261–273

Leiboff S, Li X, Hu H-C, Todt N, Yang J, Li X, Yu X, Muehlbauer GJ, Timmermans MCP, Yu
J, et al (2015) Genetic control of morphometric diversity in the maize shoot apical
meristem. Nat Commun **6**: 8974

Liaw A, Wiener M, Others (2002) Classification and regression by randomForest. R news **2**:
18–22

Li M, Klein LL, Duncan KE, Jiang N, Chitwood DH, Londo JP, Miller AJ, Topp CN (2019)
Characterizing 3D inflorescence architecture in grapevine using X-ray imaging and
advanced morphometrics: implications for understanding cluster density. J Exp Bot **70**:
6261–6276

Li M, Shao M-R, Zeng D, Ju T, Kellogg EA, Topp CN (2020) Comprehensive 3D phenotyping
reveals continuous morphological variation across genetically diverse sorghum
inflorescences. New Phytol **226**: 1873–1885

Lobet G, Pagès L, Draye X (2011) A novel image-analysis toolbox enabling quantitative
analysis of root system architecture. Plant Physiol **157**: 29–39

Lynch J (1995) Root Architecture and Plant Productivity. Plant Physiol **109**: 7–13

Mairhofer S, Zappala S, Tracy SR, Sturrock C, Bennett M, Mooney SJ, Pridmore T (2012) RooTrak: automated recovery of three-dimensional plant root architecture in soil from x-ray microcomputed tomography images using visual tracking. Plant Physiol **158**: 561–569

Malik F, Baharudin B (2012) Quantized histogram color features analysis for image retrieval based on median and Laplacian filters in DCT domain. 2012 International Conference on Innovation Management and Technology Research. doi: 10.1109/icimtr.2012.6236471

Mayer, A (2019) Making Agriculture Part of the Climate Change Solution: Researchers seek new ways to sustainably increase food production. BioScience **69**: 771-777

Mooney SJ, Pridmore TP, Helliwell J, Bennett MJ (2012) Developing X-ray Computed Tomography to non-invasively image 3-D root systems architecture in soil. Plant Soil **352**: 1–22

Morris EC, Griffiths M, Golebiowska A, Mairhofer S, Burr-Hersey J, Goh T, von Wangenheim D, Atkinson B, Sturrock CJ, Lynch JP, et al (2017) Shaping 3D Root System Architecture. Curr Biol **27**: R919–R930

Nielsen KL, Lynch JP, Weiss HN (1997) Fractal geometry of bean root systems: correlations between spatial and fractal dimension. Am J Bot **84**: 26–33

Nyquist WE, Baker RJ (1991) Estimation of heritability and prediction of selection response in plant populations. Crit Rev Plant Sci **10**: 235–322

Ortman EE, Peters DC (1968) Vertical-pull technique for evaluating tolerance of corn root systems to northern and western corn rootworms. J Econ Entomol **61**: 373-375

O'Toole JC, Soemartono (1981) Evaluation of a simple technique for characterizing rice root systems in relation to drought resistance. Euphytica **30**: 283–290

Pauli D, Chapman SC, Bart R, Topp CN, Lawrence-Dill CJ, Poland J, Gore MA (2016) The Quest for Understanding Phenotypic Variation via Integrated Approaches in the Field Environment. Plant Physiol **172**: 622–634

Rich SM, Watt M (2013) Soil conditions and cereal root system architecture: review and considerations for linking Darwin and Weaver. J Exp Bot **64**: 1193–1208

Rogers ED, Monaenkova D, Mijar M, Nori A, Goldman DI, Benfey PN (2016) X-Ray Computed Tomography Reveals the Response of Root System Architecture to Soil Texture. Plant Physiol **171**: 2028–2040

Samson BK, Sinclair TR (1994) Soil core and minirhizotron comparison for the determination of root length density. Plant Soil **161**: 225–232

Seethepalli A, Guo H, Liu X, Griffiths M, Almtarfi H, Li Z, Liu S, Zare A, Fritschi FB, Blancaflor EB, et al (2020) RhizoVision Crown: An Integrated Hardware and Software Platform for Root Crown Phenotyping. Plant Phenomics **2020**: 3074916

Sharp RE (2004) Root growth maintenance during water deficits: physiology to functional genomics. Journal of Experimental Botany **55**: 2343–2351

Symonova O, Topp CN, Edelsbrunner H (2015) DynamicRoots: A Software Platform for the
Reconstruction and Analysis of Growing Plant Roots. PLoS One **10**: e0127657

Tatsumi J, Yamauchi A, Kono Y (1989) Fractal Analysis of Plant Root Systems. Ann Bot **64**:
499–503

Topp CN, Bray AL, Ellis NA, Liu Z (2016) How can we harness quantitative genetic variation
in crop root systems for agricultural improvement? Journal of Integrative Plant Biology **58**:
213–225

Topp CN, Iyer-Pascuzzi AS, Anderson JT, Lee C-R, Zurek PR, Symonova O, Zheng Y,
Bucksch A, Mileyko Y, Galkovskyi T, et al (2013) 3D phenotyping and quantitative trait
locus mapping identify core regions of the rice genome controlling root architecture. Proc
Natl Acad Sci U S A **110**: E1695–704

Uga Y, Sugimoto K, Ogawa S, Rane J, Ishitani M, Hara N, Kitomi Y, Inukai Y, Ono K,
Kanno N, et al (2013) Control of root system architecture by DEEPER ROOTING 1
increases rice yield under drought conditions. Nat Genet **45**: 1097–1102

Venables WN, Ripley BD (2002) Modern Applied Statistics with S. Statistics and Computing.
doi: 10.1007/978-0-387-21706-2

Wasson AP, Rebetzke GJ, Kirkegaard JA, Christopher J, Richards RA, Watt M (2014) Soil
coring at multiple field environments can directly quantify variation in deep root traits to
select wheat genotypes for breeding. J Exp Bot **65**: 6231–6249

Wedger MJ, Topp CN, Olsen KM (2019) Convergent evolution of root system architecture in
two independently evolved lineages of weedy rice. New Phytol **223**: 1031–1042

Yang W, Guo Z, Huang C, Duan L, Chen G, Jiang N, Fang W, Feng H, Xie W, Lian X, et al
(2014) Combining high-throughput phenotyping and genome-wide association studies to
reveal natural genetic variation in rice. Nat Commun **5**: 5087

Zurek PR, Topp CN, Benfey PN (2015) Quantitative trait locus mapping reveals regions of the
maize genome controlling root system architecture. Plant Physiol **167**: 1487–1496

Figure 1: Pipeline for 3D root imaging using X-ray computed tomography. Samples are excised from the soil in the field using the root pulling force method, and the measurement recorded. The root crown is washed, dried, and then imaged using the NorthStar Imaging X5000 (see Methods) to generate radiographs as the sample is rotated 360º across the vertical axis. From the radiographs, a 3D reconstruction is generated using the FDK algorithm. Slices along the vertical axis are exported for automated thresholding, from which a skeleton and point cloud model of the root crown are generated. 3D root traits are then measured from the skeleton and point cloud, and analyzed. See Supplementary Figure 1 for large example images of 3D models.

Root Pulling → X-Ray CT → Reconstruction & Thresholding → 3D Skeleton, Point Cloud → Feature Extraction

Figure 2: RPF and 3D global root system architecture traits are affected by genotype, environment, and developmental time point. (A) 3D reconstructions from X-ray imaging of genotype Tx601 root crowns in the SAM 2018 experiment, at the two time points (9 vs 16 weeks) and from each environment (limited vs full irrigation). (B) ANOVA for genotype, environment, and time point (in SAM 2018) effects upon RPF and 3D root traits (adjusted p < 0.05); for legibility, G2F 2017 and SAM 2018 experiments were separately scaled, and non-significant features were set at a -log10(p) value of 0. (C) Boxplot of selected traits significantly different between the two environmental conditions in the G2F 2017 experiment (Mann-Whitney U Test p < 0.05). (D) Boxplot of selected traits significantly different between the two development time points and/or the two environmental conditions in the SAM 2018 experiment (Mann-Whitney U Test p < 0.05).

Figure 3: Classification based on genotype, environment, and time point using 3D root system architecture traits and RPF. Examples of highly distinguishable genotypes by PCA-LDA in the G2F 2017 data (A) and SAM 2018 data (B). Random forest classification of all samples based on environment in the G2F 2017 data (C) and importance of the 12 most influential traits plus RPF (D). Random forest classification of all samples based on time point in the SAM 2018 data (E) and importance of the 12 most influential traits plus RPF (F).

Figure 4: RPF and 3D root system architecture traits are strongly associated with each other and root biomass. Heatmap of RPF and its 20 most correlated traits (10 most positive and 10 most negative) in the G2F 2017 (A) and SAM 2018 (B) datasets. Regression of example traits positively correlated to RPF such as fractal dimension side/top (C, D) and surface area (E), and example traits negatively correlated to RPF such as convex hull volume (F), within the G2F 2017 dataset. Regression of example traits positively correlated to root biomass such as RPF (G), volume (H), and number of tips (I), and example traits negatively correlated to root biomass such as solidity vertical histogram-01 (J), within the SAM 2018 dataset. Adjusted r-squared values for C-J shown in respective plot insets.

Parsed Citations

Bailey PHJ, Currey JD, Fitter AH (2002) The role of root system architecture and root hairs in promoting anchorage against uprooting forces in Allium cepa and root mutants of Arabidopsis thaliana. Journal of Experimental Botany 53: 333–340
  Google Scholar: Author Only Title Only Author and Title

Bao Y, Aggarwal P, Robbins NE 2nd, Sturrock CJ, Thompson MC, Tan HQ, Tham C, Duan L, Rodriguez PL, Vernoux T, et al (2014) Plant roots use a patterning mechanism to position lateral root branches toward available water. Proc Natl Acad Sci U S A 111: 9319–9324
  Google Scholar: Author Only Title Only Author and Title

Bates D, Maechler M, Bolker B (2015) Walker., S. Fitting linear mixed-effects models using lme4. J Stat Softw 67: 1–48
  Google Scholar: Author Only Title Only Author and Title

Bohm W, Böhm W (1979) Methods of studying root systems. Springer verlag
  Google Scholar: Author Only Title Only Author and Title

Bray AL, Topp CN (2018) The Quantitative Genetic Control of Root Architecture in Maize. Plant Cell Physiol 59: 1919–1930
  Google Scholar: Author Only Title Only Author and Title

Bucksch A, Burridge J, York LM, Das A, Nord E, Weitz JS, Lynch JP (2014) Image-based high-throughput field phenotyping of crop roots. Plant Physiol 166: 470–486
  Google Scholar: Author Only Title Only Author and Title

Burridge JD, Schneider HM, Huynh B-L, Roberts PA, Bucksch A, Lynch JP (2017) Genome-wide association mapping and agronomic impact of cowpea root architecture. Theor Appl Genet 130: 419–431
  Google Scholar: Author Only Title Only Author and Title

Canè MA, Maccaferri M, Nazemi G, Salvi S, Francia R, Colalongo C, Tuberosa R (2014) Association mapping for root architectural traits in durum wheat seedlings as related to agronomic performance. Mol Breed 34: 1629–1645
  Google Scholar: Author Only Title Only Author and Title

Clark RT, MacCurdy RB, Jung JK, Shaff JE, McCouch SR, Aneshansley DJ, Kochian LV (2011) Three-dimensional root phenotyping with a novel imaging and software platform. Plant Physiol 156: 455–465
  Google Scholar: Author Only Title Only Author and Title

Colombi T, Kirchgessner N, Le Marié CA, York LM, Lynch JP, Hund A (2015) Next generation shovelomics: set up a tent and REST. Plant Soil 388: 1–20
  Google Scholar: Author Only Title Only Author and Title

Das A, Schneider H, Burridge J, Ascanio AKM, Wojciechowski T, Topp CN, Lynch JP, Weitz JS, Bucksch A (2015) Digital imaging of root traits (DIRT): a high-throughput computing and collaboration platform for field-based root phenomics. Plant Methods 11: 51
  Google Scholar: Author Only Title Only Author and Title

Delory BM, Li M, Topp CN, Lobet G (2018) archiDART v3.0: A new data analysis pipeline allowing the topological analysis of plant root systems. F1000Res 7: 22
  Google Scholar: Author Only Title Only Author and Title

Donovan LS, Jui P, Kloek M, Nicholls CF (1982) AN IMPROVED METHOD OF MEASURING ROOT STRENGTH IN CORN (Zea mays L.). Canadian Journal of Plant Science 62: 223–227
  Google Scholar: Author Only Title Only Author and Title

Duncan KE, Bray AL, Dowd TG, Topp CN (2019) Using 3D X-ray Microscopy to Study Crown Root Development and Primary Root Tip Growth in Diverse Maize (Zea mays L.) Lines. Microsc Microanal 25: 1032–1033
  Google Scholar: Author Only Title Only Author and Title

Eghball B, Settimi JR, Maranville JW, Parkhurst AM (1993) Fractal analysis for morphological description of corn roots under nitrogen stress. Agron J 85: 287–289
  Google Scholar: Author Only Title Only Author and Title

Eshel A (1998) On the fractal dimensions of a root system. Plant Cell Environ 21: 247–251
  Google Scholar: Author Only Title Only Author and Title

Fletcher RS, Mullen JL, Heiliger A, McKay JK (2015) QTL analysis of root morphology, flowering time, and yield reveals trade-offs in response to drought in Brassica napus. J Exp Bot 66: 245–256
  Google Scholar: Author Only Title Only Author and Title

Fox J, Weisberg S (2018) An R Companion to Applied Regression. SAGE Publications
  Google Scholar: Author Only Title Only Author and Title

Galkovskyi T, Mileyko Y, Bucksch A, Moore B, Symonova O, Price CA, Topp CN, Iyer-Pascuzzi AS, Zurek PR, Fang S, et al (2012) GiA Roots: software for the high throughput analysis of plant root system architecture. BMC Plant Biol 12: 116
  Google Scholar: Author Only Title Only Author and Title

Gamuyao R, Chin JH, Pariasca-Tanaka J, Pesaresi P, Catausan S, Dalid C, Slamet-Loedin I, Tecson-Mendoza EM, Wissuwa M, Heuer S (2012) The protein kinase Pstol1 from traditional rice confers tolerance of phosphorus deficiency. Nature 488: 535–539
  Google Scholar: Author Only Title Only Author and Title

Grift TE, Novais J, Bohn M (2011) High-throughput phenotyping technology for maize roots. Biosystems Eng 110: 40–48
  Google Scholar: Author Only Title Only Author and Title

Helliwell JR, Sturrock CJ, Mairhofer S, Craigon J, Ashton RW, Miller AJ, Others The emergent rhizosphere: imaging the development of the porous architecture at the root-soil interface. Sci Rep 2017; 7 (1): 14875.
  Google Scholar: Author Only Title Only Author and Title

Hochholdinger F (2009) The Maize Root System: Morphology, Anatomy, and Genetics. Handbook of Maize: Its Biology 145–160
  Google Scholar: Author Only Title Only Author and Title

Hochholdinger F, Yu P, Marcon C (2017) Genetic Control of Root System Development in Maize. Trends Plant Sci. doi: 10.1016/j.tplants.2017.10.004
  Google Scholar: Author Only Title Only Author and Title

Holbert JR, Koehler B (1924) Anchorage and extent of corn root systems. J Agric Res 27: 71–78
  Google Scholar: Author Only Title Only Author and Title

Iyer-Pascuzzi AS, Symonova O, Mileyko Y, Hao Y, Belcher H, Harer J, Weitz JS, Benfey PN (2010) Imaging and analysis platform for automatic phenotyping and trait ranking of plant root systems. Plant Physiol 152: 1148–1157
  Google Scholar: Author Only Title Only Author and Title

Jiang N, Floro E, Bray AL, Laws B, Duncan KE, Topp CN (2019) Three-Dimensional Time-Lapse Analysis Reveals Multiscale Relationships in Maize Root Systems with Contrasting Architectures. The Plant Cell 31: 1708–1722
  Google Scholar: Author Only Title Only Author and Title

Kitomi Y, Hanzawa E, Kuya N, Inoue H, Hara N, Kawai S, Kanno N, Endo M, Sugimoto K, Yamazaki T, et al (2020) Root angle modifications by the DRO1 homolog improve rice yields in saline paddy fields. Proc Natl Acad Sci U S A 117: 21242–21250
  Google Scholar: Author Only Title Only Author and Title

Kuhn M, Others (2008) Building predictive models in R using the caret package. J Stat Softw 28: 1–26
  Google Scholar: Author Only Title Only Author and Title

Landi P, Sanguineti MC, Darrah LL, Giuliani MM, Salvi S, Conti S, Tuberosa R (2002) Detection of QTLs for vertical root pulling resistance in maize and overlap with QTLs for root traits in hydroponics and for grain yield under different water regimes. Maydica 47: 233–243
  Google Scholar: Author Only Title Only Author and Title

Le Bot J, Serra V, Fabre J, Draye X, Adamowicz S, Pagès L (2010) DART: a software to analyse root system architecture and development from captured images. Plant Soil 326: 261–273
  Google Scholar: Author Only Title Only Author and Title

Leiboff S, Li X, Hu H-C, Todt N, Yang J, Li X, Yu X, Muehlbauer GJ, Timmermans MCP, Yu J, et al (2015) Genetic control of morphometric diversity in the maize shoot apical meristem. Nat Commun 6: 8974
  Google Scholar: Author Only Title Only Author and Title

Liaw A, Wiener M, Others (2002) Classification and regression by randomForest. R news 2: 18–22
  Google Scholar: Author Only Title Only Author and Title

Li M, Klein LL, Duncan KE, Jiang N, Chitwood DH, Londo JP, Miller AJ, Topp CN (2019) Characterizing 3D inflorescence architecture in grapevine using X-ray imaging and advanced morphometrics: implications for understanding cluster density. J Exp Bot 70: 6261–6276
  Google Scholar: Author Only Title Only Author and Title

Li M, Shao M-R, Zeng D, Ju T, Kellogg EA, Topp CN (2020) Comprehensive 3D phenotyping reveals continuous morphological variation across genetically diverse sorghum inflorescences. New Phytol 226: 1873–1885
  Google Scholar: Author Only Title Only Author and Title

Lobet G, Pagès L, Draye X (2011) A novel image-analysis toolbox enabling quantitative analysis of root system architecture. Plant Physiol 157: 29–39
  Google Scholar: Author Only Title Only Author and Title

Lynch J (1995) Root Architecture and Plant Productivity. Plant Physiol 109: 7–13
  Google Scholar: Author Only Title Only Author and Title

Mairhofer S, Zappala S, Tracy SR, Sturrock C, Bennett M, Mooney SJ, Pridmore T (2012) RooTrak: automated recovery of three-dimensional plant root architecture in soil from x-ray microcomputed tomography images using visual tracking. Plant Physiol 158: 561–569
  Google Scholar: Author Only Title Only Author and Title

Malik F, Baharudin B (2012) Quantized histogram color features analysis for image retrieval based on median and Laplacian filters in DCT domain. 2012 International Conference on Innovation Management and Technology Research. doi: 10.1109/icimtr.2012.6236471

Google Scholar: Author Only Title Only Author and Title

Mayer, A (2019) Making Agriculture Part of the Climate Change Solution: Researchers seek new ways to sustainably increase food production. BioScience 69: 771-777
Google Scholar: Author Only Title Only Author and Title

Mooney SJ, Pridmore TP, Helliwell J, Bennett MJ (2012) Developing X-ray Computed Tomography to non-invasively image 3-D root systems architecture in soil. Plant Soil 352: 1–22
Google Scholar: Author Only Title Only Author and Title

Morris EC, Griffiths M, Golebiowska A, Mairhofer S, Burr-Hersey J, Goh T, von Wangenheim D, Atkinson B, Sturrock CJ, Lynch JP, et al (2017) Shaping 3D Root System Architecture. Curr Biol 27: R919–R930
Google Scholar: Author Only Title Only Author and Title

Nielsen KL, Lynch JP, Weiss HN (1997) Fractal geometry of bean root systems: correlations between spatial and fractal dimension. Am J Bot 84: 26–33
Google Scholar: Author Only Title Only Author and Title

Nyquist WE, Baker RJ (1991) Estimation of heritability and prediction of selection response in plant populations. Crit Rev Plant Sci 10: 235–322
Google Scholar: Author Only Title Only Author and Title

Ortman EE, Peters DC (1968) Vertical-pull technique for evaluating tolerance of corn root systems to northern and western corn rootworms. J Econ Entomol 61: 373-375
Google Scholar: Author Only Title Only Author and Title

O'Toole JC, Soemartono (1981) Evaluation of a simple technique for characterizing rice root systems in relation to drought resistance. Euphytica 30: 283–290
Google Scholar: Author Only Title Only Author and Title

Pauli D, Chapman SC, Bart R, Topp CN, Lawrence-Dill CJ, Poland J, Gore MA (2016) The Quest for Understanding Phenotypic Variation via Integrated Approaches in the Field Environment. Plant Physiol 172: 622–634
Google Scholar: Author Only Title Only Author and Title

Rich SM, Watt M (2013) Soil conditions and cereal root system architecture: review and considerations for linking Darwin and Weaver. J Exp Bot 64: 1193–1208
Google Scholar: Author Only Title Only Author and Title

Rogers ED, Monaenkova D, Mijar M, Nori A, Goldman DI, Benfey PN (2016) X-Ray Computed Tomography Reveals the Response of Root System Architecture to Soil Texture. Plant Physiol 171: 2028–2040
Google Scholar: Author Only Title Only Author and Title

Samson BK, Sinclair TR (1994) Soil core and minirhizotron comparison for the determination of root length density. Plant Soil 161: 225–232
Google Scholar: Author Only Title Only Author and Title

Seethepalli A, Guo H, Liu X, Griffiths M, Almtarfi H, Li Z, Liu S, Zare A, Fritschi FB, Blancaflor EB, et al (2020) RhizoVision Crown: An Integrated Hardware and Software Platform for Root Crown Phenotyping. Plant Phenomics 2020: 3074916
Google Scholar: Author Only Title Only Author and Title

Sharp RE (2004) Root growth maintenance during water deficits: physiology to functional genomics. Journal of Experimental Botany 55: 2343–2351
Google Scholar: Author Only Title Only Author and Title

Symonova O, Topp CN, Edelsbrunner H (2015) DynamicRoots: A Software Platform for the Reconstruction and Analysis of Growing Plant Roots. PLoS One 10: e0127657
Google Scholar: Author Only Title Only Author and Title

Tatsumi J, Yamauchi A, Kono Y (1989) Fractal Analysis of Plant Root Systems. Ann Bot 64: 499–503
Google Scholar: Author Only Title Only Author and Title

Topp CN, Bray AL, Ellis NA, Liu Z (2016) How can we harness quantitative genetic variation in crop root systems for agricultural improvement? Journal of Integrative Plant Biology 58: 213–225

Topp CN, Iyer-Pascuzzi AS, Anderson JT, Lee C-R, Zurek PR, Symonova O, Zheng Y, Bucksch A, Mileyko Y, Galkovskyi T, et al (2013) 3D phenotyping and quantitative trait locus mapping identify core regions of the rice genome controlling root architecture. Proc Natl Acad Sci U S A 110: E1695–704
Google Scholar: Author Only Title Only Author and Title

Uga Y, Sugimoto K, Ogawa S, Rane J, Ishitani M, Hara N, Kitomi Y, Inukai Y, Ono K, Kanno N, et al (2013) Control of root system architecture by DEEPER ROOTING 1 increases rice yield under drought conditions. Nat Genet 45: 1097–1102
Google Scholar: Author Only Title Only Author and Title

Venables WN, Ripley BD (2002) Modern Applied Statistics with S. Statistics and Computing. doi: 10.1007/978-0-387-21706-2

Google Scholar: Author Only Title Only Author and Title

Wasson AP, Rebetzke GJ, Kirkegaard JA, Christopher J, Richards RA, Watt M (2014) Soil coring at multiple field environments can directly quantify variation in deep root traits to select wheat genotypes for breeding. J Exp Bot 65: 6231–6249
Google Scholar: Author Only Title Only Author and Title

Wedger MJ, Topp CN, Olsen KM (2019) Convergent evolution of root system architecture in two independently evolved lineages of weedy rice. New Phytol 223: 1031–1042
Google Scholar: Author Only Title Only Author and Title

Yang W, Guo Z, Huang C, Duan L, Chen G, Jiang N, Fang W, Feng H, Xie W, Lian X, et al (2014) Combining high-throughput phenotyping and genome-wide association studies to reveal natural genetic variation in rice. Nat Commun 5: 5087
Google Scholar: Author Only Title Only Author and Title

Zurek PR, Topp CN, Benfey PN (2015) Quantitative trait locus mapping reveals regions of the maize genome controlling root system architecture. Plant Physiol 167: 1487–1496
Google Scholar: Author Only Title Only Author and Title