LCK Regulates Homologous Recombination DNA Repair Identifying a New Target

for Sensitizing PARP Inhibitors in HR Proficient Ovarian Cancer

Goutam Dey[1+], Rashmi Bharti[1+], Chad Braley[1], Ravi Alluri[1], Emily Esakov[1], Katie Crean-

Tate[2], Keith McCrae[1,5], Amy Joehlin-Price[3], Peter G. Rose[2], Justin Lathia[1,5], Zihua

Gong[4,5], Ofer Reizes[1,5*]

1 Department of Cardiovascular and Metabolic Sciences, Lerner Research Institute,

Cleveland Clinic Foundation, Cleveland, Ohio, USA

2 Division of Gynecologic Cancer, Women's Health Institute, Cleveland Clinic

3 Pathology and Lab Medicine, Cleveland Clinic

4 Cancer Biology, Lerner Research Institute, Cleveland Clinic

5 Case Comprehensive Cancer Center, Cleveland, OH

[+] co-first authors
* Address correspondence to:
Ofer Reizes, Ph.D.
Department of Cardiovascular and Metabolic Sciences
Lerner Research Institute
9500 Euclid Avenue
Cleveland Clinic
Cleveland, OH 44195
Email: reizeso@ccf.org
Direct: 216-445-0880
24

25 **Short title**: LCK regulates homologous recombination repair

26 **Conflict of interest:** Authors declare no conflict of interest.

29 **Graphical Abstract**

Abstract

Poly-ADP Ribose Polymerase (PARP) targeted therapy is clinically approved for the treatment of homologous recombination (HR) repair deficient tumors. The remarkable success in treatment of HR repair deficient cancers has not translated to HR-proficient cancer. Our studies identify the mechanism of non-receptor lymphocyte-specific protein tyrosine kinase (LCK) in HR repair in endometrioid epithelial ovarian cancer (eEOC). LCK expression is induced and activation in the nucleus in response to DNA damage insult. LCK inhibition attenuates expression of RAD51, BRCA1, and BRCA2 proteins necessary for HR-mediated DNA repair, sufficient to suppress RAD51 foci formation, and augments γH2AX foci formation. Mechanistically, DNA damage leads to direct interaction of LCK with RAD51 and BRCA1 in a kinase dependent manner. Attenuation of LCK sensitized HR-proficient eEOC cells to PARP inhibitor in cell culture and pre-clinical mouse studies. These findings identify a new mechanism for expanding utility of PARP inhibitors in HR proficient ovarian cancer.

Abbreviations: eEOC: endometrioid epithelial ovarian cancer, HR: homologous recombination, LCK: lymphocyte-specific protein tyrosine kinase, PARP: poly(ADP-ribose) polymerase, DDR: DNA damage response, NHEJ: non–homologous end joining.

Introduction

Epithelial ovarian cancer (EOC) is the second most common gynecologic malignancy in the United States, but the leading cause of gynecologic cancer death. It is estimated that in 2022, ~19,880 women in the U.S. will be diagnosed with epithelial ovarian cancer (EOC) and 12,810 will die from their disease[1]. Poly-(ADP-ribose) polymerase inhibitors (PARPi) have emerged as new therapeutic options in the treatment of ovarian cancer[2,3,4]. Recent studies show that this treatment has prolonged median recurrence-free survival after primary therapy by more than 24 months[5]. While the benefit of PARPi is greatest in BRCA1/2-mutant or deficient tumors, those with HR deficiencies also experience a benefit from this therapy[5]. Conversely, PARPi and chemotherapy have so far shown limited efficacy in HR-proficient ovarian cancers. Further, platinum resistance is associated with HR proficiency in EOC[6,7]. This limited efficacy of both platinum and PARPi therapy highlights an unmet clinical need in ovarian cancer patients.

Several strategies have been assessed to expand the utility of PARPi in HR-proficient cancers[2,8,9,10]. RAD51, BRCA1, and BRCA2 are critical components of the HR repair complex. Studies have focused on disrupting this complex. Cyclin-dependent kinase (CDK) proteins demonstrably regulate the HR repair pathway in a lung cancer model[11]. Indeed, the CDK inhibitor dinaciclib is able to attenuate the expression of RAD51 and BRCA proteins resulting in the inhibition of HR repair capacity and potentiation of the pharmacological effect of PARPi[9]. However, there is no clinically approved drug for combination with PARPi for HR-proficient cancers.

Approximately 80% of endometrial cancers and 10% of ovarian cancers demonstrate endometrioid tumor histology (eEOC)[12]. A small but clinically significant proportion of

eEOC display high-grade histology, advanced stage (FIGO stage III-IV), and a poor 5-

year survival of 6-24%. These traits are similar to those of the more aggressive high-

grade serous type of ovarian cancer [13]. Moreover, somatic and/or germline mutations in

HR genes occur in only a third of ovarian tumors, indicating the majority of eEOC are

HR-proficient. Of note, eEOC show a considerably higher rate of resistance to platinum-

based chemotherapy[14] compared to serous carcinomas and do not commonly respond

to targeted therapies such as PARP inhibitors.

We previously determined that intracellular, non-receptor tyrosine kinase, LCK regulates

genes implicated in DNA repair machinery in eEOC[15]. We also demonstrated the

pharmacologic inhibition of LCK attenuated expression of homologous recombination

DNA damage repair genes leading to sensitization of eEOC cells to cisplatin[16]. In

contrast, increased expression of LCK led to upregulation of DNA damage-repair genes

and increased resistance to cisplatin. As LCK modulates RAD51, BRCA1, and BRCA2

expression, we hypothesized that blocking LCK expression or inhibiting kinase activity

would sensitizes eEOC to PARPi. Here, we elucidate the mechanism of LCK in

regulating HR DNA damage repair and a therapeutic approach to sensitize HR-

proficient eEOC to PARP inhibitors.

Results

Regulation of Homologous Recombination (HR) DNA repair protein expression by

LCK in eEOC.

We tested whether LCK inhibition is sufficient to inhibit HR DNA repair genes RAD51,

BRCA1, and BRCA2 at the protein level. We inhibited LCK expression using shRNA

and CRISPR in the CP70 and SKOV3 cell lines, both of which are cisplatin resistant and

known HR-proficient eEOC cells. Cells were transduced with lentivirus containing

shRNA control (shCon) or LCK-targeted shRNA (KD1, KD2). Additionally, we generated

LCK knock-out (KO) CP70 cells via CRISPR/Cas9. LCK inhibition was confirmed by

immunoblotting followed by analysis of expression of BRCA1, BRCA2, and RAD51 via

western blot analysis. In CP70 cells, we observed that KD1, KD2, and KO displayed

attenuated protein expression of BRCA1, BRCA2 and RAD51 when compared to shCon

(Fig. 1A, and Supplementary Fig. S1A and C). The protein expression levels of BRCA1,

BRCA2, and RAD51 were similarly attenuated in LCK knock-down SKOV3 cells (Fig.

1A, and Supplementary Fig. S1B).

In complementary studies, we tested whether LCK overexpression would increase

RAD51, BRCA1, and BRCA2 protein expression in eEOC. LCK overexpression led to

induction of RAD51, BRCA1, and BRCA2 protein expression in CP70 and SKOV3 cells

(Fig. 1A, and Supplementary Fig. S1D and E). To test the hypothesis that LCK

increases the stability of BRCA1 and RAD51 proteins, empty vector (EV) and OE cells

were treated with cycloheximide and harvested at various time-points to assess protein

expression levels (Fig.1B, C, Supplementary Fig. S1F). The half-lives of BRCA1 and

RAD51 were significantly increased in OE when compared to control. Half-lives of

RAD51 in CP70 EV and OE cells treated with cycloheximide were 3.2 and 5.6 h,

respectively. Half-lives of BRCA1 in CP70 EV and OE cells treated with cycloheximide

were 54 min and 3.4 h, respectively. These studies indicate that LCK is sufficient to

regulate BRCA1 and RAD51 protein expression via protein stabilization.

To test whether pharmacologic inhibition of LCK can attenuate expression of DNA

damage repair proteins, we used PP2, a cell-permeable, small-molecule inhibitor of

LCK kinase[17,18]. We tested the efficacy of PP2 in CP70 and SKOV3 OE cells. PP2

attenuated pLCK at Y394, the autophosphorylation site of LCK in these cells (Fig. 1D,

and Supplementary Fig. S1G and H). γH2AX, a marker of DNA damage and replication

stress, was elevated by PP2 treatment. This is indicative of either increased damage or

reduced repair of DNA damage due to attenuation of BRCA1 and RAD51 expression.

LCK inhibition also attenuates expression of RAD51, BRCA1, and BRCA2 in parental

CP70 and SKOV3 as well as in the CRL1978 clear-cell EOC cell line (Supplementary

Fig. S2A).

LCK inhibition attenuates HR DNA damage repair in eEOC.

The inhibition of DNA damage repair genes led us to test whether LCK inhibition impairs

HR-dependent DNA repair. We utilized the DR-GFP reporter assay established in

U2OS cells to measure repair efficiency[19] (Fig. 2A). U2OS cells with/without DR-GFP

reporter system express endogenous LCK protein expression as shown by western blot

analysis (Supplementary Fig. S2B). U2OS cells treated with PP2 leads to a dose-

dependent reduction of the GFP-positive cell population when compared to DMSO-

treated cell population, indicating reduced DNA repair as a consequence of LCK

inhibition (Fig. 2B). Likewise, shRNA silencing of LCK led to a reduction in the GFP-

positive population compared to shCon transduced cells (Fig. 2B). This indicates that

LCK inhibition attenuates HR repair efficiency in cancer cells.

DNA damage leads to activation of several repair pathways including PARP, HR, and

NHEJ [20]. As our studies indicated LCK inhibition attenuates HR repair proteins, we

assessed LCK's impact on the expression of alternative DNA repair pathways, including

PARP and NHEJ, in CP70 and SKOV3 cells. The LCK inhibitor PP2 did not inhibit

PARylation in CP70 and SKOV3 cells (Fig. 2C). The Ku80 and Ku70 proteins are a

critical component of the NHEJ pathway[21]. After PP2 treatment, Ku80 protein

expression was elevated in CP70, but not in SKOV3 (Fig. 2D). Furthermore, Ku70

protein expression was not changed in either CP70 or SKOV3 after PP2 exposure (Fig.

2D). In parallel, Ku70 and Ku80 expression levels in CRL1978 cells did not change

following PP2 treatment (Supplementary Fig. S2C). These findings indicate that LCK

inhibition targets HR repair proteins independent of induction of NHEJ repair

mechanisms (Fig. 2E).

DNA damage induces LCK dependent BRCA1 expression.

We next assessed the effects of DNA damage on LCK expression and activation. DNA

damage in ovarian cancer cells was induced using either etoposide, ultraviolet radiation,

or methyl methanesulfonate (MMS). Dose-dependent treatment of CP70 cells with

etoposide or MMS led to increased LCK protein expression (Fig. 3A and Supplementary

Fig. S3A and B). Etoposide or MMS treatment in SKOV3 cells led to increased

phosphorylation of LCK at pY394, while total levels of LCK protein remained unchanged

(Supplementary Fig. S4A). Likewise, ultraviolet radiation of CP70 cells was sufficient to

increase LCK phosphorylation (Supplementary Fig. S4B). BRCA1 and γH2AX

expression was induced by etoposide or MMS, indicating increased DNA damage (Fig.

3A). Because DNA damage, particularly double strand breaks (DSB), and its repair

machinery are concentrated in the nucleus[22], we investigated the effects that DNA

damage could induce on the accumulation of LCK in the nucleus. We found increased

total LCK and pLCK in the nucleus of etoposide-treated cells (Fig. 3B and

Supplementary Fig. S3C). Sub-cellular fractionation and immunofluorescence analysis

both showed that pLCK was predominately localized to the nucleus of etoposide-treated

cells (Fig. 3C). These findings were replicated in SKOV3 cells (Supplementary Fig.

S4C). No pLCK was detectable in the nucleus of DMSO-treated cells (Fig. 3C). We next

tested whether inhibition of LCK would be sufficient to block BRCA1 expression in

etoposide-treated cells. Etoposide treatment in shCon cells showed increased BRCA1

protein expression, whereas this treatment attenuated BRCA1 expression in KD cells

(Fig. 3D and Supplementary Fig. S3D). We repeated these studies in CRISPR/CAS9

KO cells and observed similar attenuation of BRCA1 in KO CP70 cells and no

attenuation in parental cells (Fig. 3E and Supplementary Fig. S3E). These findings

indicate that the induction of BRCA1 expression in response to DNA damage is

disrupted by LCK inhibition.

LCK regulation of DNA double strand break repair.

As γH2AX and RAD51 are markers of DNA damage and repair of DSB, we tested for

foci formation in control and etoposide-treated cells. KO and OE CP70 (LCK

overexpression in CRISPR/Cas9-background) cells were treated in presence or

absence etoposide then subjected to immunofluorescence analysis to detect and

quantify γH2AX foci (Fig. 4A, B and Supplementary Fig. S5A). In the absence of

etoposide, no γH2AX foci formation was detected. Parental CP70 (WT) cells treated

with etoposide led to increased γH2AX foci compared to DMSO treatment. KO cells

treated with etoposide exhibited 4-5 fold increased foci formation compared to WT. Foci

formation was nearly completely blocked in etoposide treated OE cells (Fig. 4A and B).

In parallel, we assessed RAD51 foci formation in KO and OE CP70 cells treated in the

absence or presence of etoposide (Fig. 4C, D and Supplementary Fig. S5B). As with

γH2AX, no RAD51 foci were observed in WT, KO, or OE cells treated with DMSO. In

contrast, etoposide treatment led to a significant increase in RAD51 foci in WT CP70

cells that was significantly suppressed in KO cells (Fig. 4C, D, and Supplementary Fig.

S5B). RAD51 foci formation was significantly increased in OE cells treated with

etoposide. This data supports the conclusion that LCK can regulate HR repair during

DNA damage response (DDR).

LCK complexes with RAD51 and BRCA1 in response to DNA damage.

As pLCK is localized in the nucleus in response to DNA damage and can induce

BRCA1 and RAD51, we tested whether LCK directly interacts with RAD51 and BRCA1

in nuclear extracts. CP70 and SKOV3 cells were transduced with a myc-tagged LCK

(Fig. 5A and Supplementary Fig. S6). We treated cells in the absence or presence of

etoposide, isolated nuclei, and performed an IP (immunoprecipitation) assay with myc

antibodies. In untreated cells, neither BRCA1 nor RAD51 co-precipitated with mycLCK.

In contrast, etoposide treatment resulted in co-precipitation of RAD51 and BRCA1 with

mycLCK (Fig. 5B and C). In parallel, RAD51 could co-immunoprecipitate with LCK from

etoposide-treated OE SKOV3 cells. LCK and BRCA1 were detected in RAD51

immunoprecipitates (Fig. 5D). These findings indicate that in response to DNA damage,

LCK interacts with a complex containing RAD51 and BRCA1 (Fig. 5E).

LCK kinase activity is essential for HR repair.

LCK interacts with BRCA1 and RAD51 in response to DNA damage, so we tested

whether kinase activity and autophosphorylation of LCK is necessary for activity and

DNA repair. We generated LCK mutants at lysine 273, which is necessary for kinase

activity (K273R); tyrosine 394, an autophosphorylation and activation site (Y394F)[23];

and tyrosine 192, a SH2 adaptor protein binding site (Y192F)[24] and transduced them

into LCK KO CP70 cells (Fig. 6A). OE and Y192F mutants retained kinase activity, while

K273R and Y394F mutants lacked kinase activity. We performed IP studies in

etoposide-treated cells and determined that OE and Y192F cells were able to co-

immunoprecipitate BRCA1 and RAD51, whereas K273R and Y394F failed to co-

immunoprecipitated BRCA1 and RAD51 (Fig 6B). We next assessed γH2AX and

RAD51 foci formation in the OE, K273R, Y192F, and Y394F transduced cells. OE and

Y192F exhibited similar level of foci formation in response to etoposide (Fig. 6C, D, E,

and F), whereas K273R and Y394F showed increased γH2AX foci and reduced RAD51

foci in etoposide treated cells (Fig. 6C, D, E, and F and Supplementary Fig. S7A, B).

We further assayed the cells for etoposide sensitivity. We determined that naïve CP70

cells exhibited an IC50 of 5μM to etoposide. OE (IC50=10.03μM, 95% CI is 7.894 to

12.67) and Y192F (IC50=8.70μM, 95% CI is 7.044 to 10.72) exhibited increased

resistance to etoposide, whereas K273R (IC50=2.28μM, 95% CI is 1.938 to 2.692) and

Y394F (IC50=2.01μM, 95% CI is 1.655 to 2.455) showed increased sensitivity to

etoposide (Fig. 6G). We performed a timed experiment with etoposide treated CP70

cells and determined that the extent of γH2AX foci was lower in OE and LCK Y192F

mutant cells than in WT cells at 0 hours. Moreover, KO, Y394F, and K273R cells

exhibited the highest number of γH2AX foci at all time points (Fig. 6H, and IF in

Supplementary Fig. S8). These findings lead to the hypothesis that LCK kinase activity

is essential for interaction with RAD51 and BRCA1 during DNA damage response and

also facilitated HR repair.

LCK inhibition augments PARPi induced DNA damage and genomic instability.

We performed single cell gel electrophoresis (alkaline COMET) assay to quantify the

extent of double and single strand DNA breaks by visualizing tail area[25]. CP70 and

SKOV3 cells were incubated with either PP2, olaparib, or both. Treated cells were

processed and stained with SYBR Gold to detect and measure the tail moment (Fig.

7A). PP2 and olaparib alone displayed a comparable increase in comet tails compared

to DMSO (Fig. 7B). The combination of PP2 and olaparib induced a fourfold increase in

comet tail area compared to monotherapy (Fig. 7B).

PARP inhibitors have been reported to induce genomic instability, leading to

chromosomal aberration and DNA damage in cancer cells[26,27]. Chromosomal damage

can be detected by chromosomal breaks, gaps, and radial formations. We identified

multiple breaks, gaps, and radial formation in PP2 and olaparib-treated cells (Fig. 7C).

PP2 and olaparib displayed a comparable increase in chromosomal damage when

compared to DMSO (Fig. 7D). The combination of PP2 and olaparib displayed

increased chromosomal damage in both CP70 and SKOV3 cells (Fig. 7D).

Based on this analysis, we assessed whether the LCK inhibitor PP2 could synergize

with a PARPi, olaparib, to augment the DNA damage response (Fig. 7E). Olaparib

treatment led to an increase in BRCA1 expression and a detectable increase in γH2AX

expression in SKOV3 cells (Fig. 7E). Co-treatment with PP2 was sufficient to suppress

BRCA1 expression and significantly augment γH2AX expression in a dose-dependent

manner (Fig. 7E). Our findings indicate LCK inhibition leads to HR deficiency. As proof

of concept, we tested the impact of LCK silencing on the efficacy of olaparib in SKOV3

and CP70 cells via colony formation assay. Olaparib sensitivity was analyzed in

parental (WT), KO, OE (Fig. 7F). We quantified colony formation and determined that

sensitivity of CP70 to olaparib is greater in LCK KO than in parental CP70 cells (Fig. 7F,

Supplementary Fig. S9). We determined that, in CP70 cells, olaparib resistance

increased 3-fold in LCK OE compared to LCK KO. We replicated these findings by

silencing with shRNA in CP70 and SKOV3 cells (Supplementary Fig. S10A and B).

shCon, KD1, and KD2 cells were treated with various concentrations of olaparib and

plated for colony formation. In CP70 cells, silencing LCK inhibited colony formation with

greater efficiency in olaparib-treated cancer cells than in shCon treated cells

(Supplementary Fig. S10A, B). Similarly, in SKOV3 cells, the number of colonies were

significantly decreased after olaparib treatment in KD1 and KD2 cells as compared to

shCon cells (Supplementary Fig. S10C, D). These findings support the hypothesis that

olaparib has a higher efficacy in LCK-deficient cancer cells and indicate that LCK

inhibition is sufficient to sensitize eEOC to PARPi.

LCK inhibition potentiates therapeutic efficacy of PARPi in *in vivo*

To test whether LCK impacts olaparib efficacy in pre-clinical models of eEOC, we

injected KO and OE CP70 cells into mice and once tumors were detected, we treated

with 3 course of 5 day treatment of Olaparib (Fig. 8A). KO and OE CP70 exhibited

nearly identical tumor growth in vehicle-treated mice (Fig. 8B and C). Olaparib treatment

led to suppression of tumor growth in OE mice and to complete suppression of tumor

growth in KO mice (Fig. 8B and C). These findings indicate that LCK inhibition

potentiates olaparib synthetic lethality.

We performed a molecular analysis on tumor sections from OE and KO cells treated

with and without olaparib. TUNEL assay to detect apoptotic DNA fragmentation

indicated no positive cells (green fluorescence) in OE and KO tumors, indicating no

apoptotic cells (Fig.8D, Supplementary Fig. S11). Tumors from olaparib-treated mice

exhibited a few TUNEL-positive cells present in OE tumors, whereas the majority of

cells were TUNEL-positive in KO tumors. Tumors were assessed for presence of γH2AX

in tissue sections by immunohistochemistry (Fig. 8E). Vehicle treated mice exhibited low

levels of γH2AX in both KO and OE cohort (Fig. 8E). Tumors from olaparib-treated mice

exhibited low levels of γH2AX expression in OE, whereas the majority of cells were

positive in KO group (Fig. 8E). These findings indicate a large number of DNA double-

strand breaks were generated due to suppression of LCK and inhibition of PARP. We

next assayed for expression of CD31, an indicator of microvessel (angiogenesis)

density and of tumor mass and growth. Vehicle treated mice exhibited high CD31

positive staining in both KO and OE cohort (Fig. 8F). Tumors from olaparib-treated mice

exhibited high levels of CD31 expression in OE, whereas there was no detectable CD31

in the KO group (Fig. 8F). These findings suggested that olaparib was also sufficient to

inhibit tumor angiogenesis, corroborating previous findings which showed that PARP

facilitates tumor vascularization by augmenting CD31 and VEGF[28].

Discussion

Endometrioid ovarian cancer, while rare, lacks effective therapeutic strategies and are HR proficient. Our studies identified a strategy to induce HR deficiency in endometrioid ovarian cancer. We discovered that inhibition or disruption of the non-receptor tyrosine kinase LCK attenuates the expression of HR proteins RAD51, BRCA1, and BRCA2 in eEOC. This complements our previous study showing that LCK overexpression or LCK inhibition modulated the mRNA levels of HR DNA repair genes including RAD51, BRCA1, and BRCA2 [15]. Here, we showed that LCK modulates HR genes at the protein level. This leads to functional consequences as the inhibition of LCK, via shRNA or pharmacologic inhibitor, leads to inhibition of DNA damage repair as assessed using the established DR-GFP assay in U2OS osteosarcoma cells. LCK does not impact PARP and NHEJ repairs, the alternate mechanisms for repair of double strand breaks by direct ligation independent of an homologous template[21]. The NHEJ repair proteins, Ku70 and Ku80, did not have their expression levels impacted by PP2 treatment. LCK kinase activity is necessary for maintaining HR proficiency. Finally, we demonstrated that LCK disruption is sufficient to sensitize endometrioid ovarian cancer cells to olaparib. These findings are consistent with our data indicating that LCK inhibition leads to chemosensitization to cisplatin treatment in endometrioid ovarian cancer.

We determined for the first time that the LCK protein is upregulated in response to DNA damage in eEOC. DNA damage by etoposide, methyl methanesulfonate (MMS), and by UV radiation induces LCK protein expression. This is also corroborated by previous findings that indicate fractionated radiation induces stem cell populations in human gliomas to display LCK activation [29]. Phosphorylation of LCK (pY394) was elevated in

etoposide, MMS, and UV-treated SKOV3 cells and in UV-treated CP70 cells as

compared to untreated cells. We found that DNA damage led to nuclear localization of

both total and pY394 LCK protein, a finding supported by immunofluorescence analysis

of pY394 LCK following DNA damage. This finding is unprecedented as LCK is

localized to the inner leaflet of the cell membrane on microdomains[30]. Previous studies

have found constitutively active LCK in the nucleus where it binds at the promoter

region of LIM domain only 2 (LMO2) leading to gene expression[31]. Our findings are

significant as we show that LCK is activated by DNA damage, leading to nuclear

translocation and subsequent activation of HR repair pathways.

Here we confirm nuclear translocation of LCK in response to DNA damage as shown by

co-immunoprecipitation with BRCA1 and RAD51. Interaction of LCK with BRCA1 and

RAD51 requires kinase activity or phosphorylation on Y394, indicating active LCK is

necessary for complex formation[23]. In contrast, phosphorylation on Y192 is not required

for complex formation. Moreover, kinase activity and autophosphorylation are necessary

for functional DNA repair, as shown by γH2AX and RAD51 foci formation assays. These

findings indicate that LCK kinase activity and autophosphorylation is essential to allow

for interaction with HR repair proteins BRCA1 and RAD51 during DNA damage

response.

LCK regulates HR repair in response to DNA damage and its inhibition potentiates the

activity of PARPi to induce synthetic lethality. The simultaneous inhibition of LCK and

PARP with pharmacological agents PP2 and olaparib showed significantly more DNA

damage and chromosomal aberration compared to only either PP2 or olaparib

treatment in eEOC cells. PP2 treatment was sufficient to attenuate DNA repair,

augmenting the effect of olaparib in ovarian cancer cells. Finally, in vivo studies showed

that olaparib efficacy was enhanced in CP70 LCK KO compared to OE tumor bearing

mice. This provides evidence for proof of concept for utility of LCK inhibitors to disrupt

HR DNA damage repair. Indeed, several strategies are currently being explored in the

clinic to increase use of PARP targeted therapy in HR proficient cancers. CDK1 and

CDK12 inhibition led to HR deficiency by decreasing HR repair proteins RAD51,

BRCA1, and BRCA2 in lung cancer[8]. Further, inhibition of BET proteins also led to

attenuation of RAD51 and BRCA1 proteins in breast, ovarian, and prostate cancer

models [2]. PI3K inhibition is also sufficient to reduce BRCA1 and BRCA2 expression,

hampering HR repair in triple-negative breast cancer[32]. Other reported targets are

HSP90 [33] and VEGFR3 [34] to attenuate RAD51, BRCA1, and BRCA2 expressions in

ovarian cancer. Concurrent inhibition of LCK enhanced the efficacy of PARPi in above

HR proficient cancer models in preclinical settings. Clinical trials are now going on to

assess the efficacy of PARPi in combination with CDK1/12 inhibitors, PI3K inhibitor, and

VEGFR3 inhibitors[3]. Our findings complement these studies and identify a new

signaling pathway for enhancing PARP targeted therapy in eEOC.

These findings provide an innovative new strategy for inducing an HR-deficient status in

an otherwise HR-proficient tumor. We identify targeted therapies that compromise HR

repair genes and augment sensitivity to PARPi. Our study defines the mechanistic

impact of LCK and potentially other non-receptor tyrosine kinases in regulation of HR

repair that is apparently crucial to ovarian cancer's response to chemotherapy and

PARP inhibitors. This study highlights new clinical applications that target LCK,

expanding PARPi utility.

Methods

Cell lines and culture conditions

Cisplatin resistant eEOC cancer cells CP70 were a gift from Analisa Difeo (University of

Michigan) and SKOV3 were purchased from American Type Culture Collection (ATCC).

Others cell lines used in this study are mentioned in resource table. Cells were grown in

DMEM and McCoy's 5A media respectively, supplemented with 10% fetal bovine serum

at 37°C in humidified incubator in 5% CO_2. Cells were tested and confirmed as

mycoplasma contamination negative on a quarterly basis. Cells were passaged by

treatment with trypsin/EDTA solution when they reached 80-90% confluence and further

passaged or seeded for experiments.

Chemicals and reagents

We used a number of pharmacological agents in our study. The PARP inhibitor

(Olaparib)[35], the LCK inhibitor (PP2)[17,18] and the radiomimetic drug, etoposide[36] were

purchased as shown in the resource table. Inhibitors were dissolved in 100% DMSO to

make stock concentrations and kept at -20°C until use. The details of chemical,

reagents, primary antibodies, and secondary antibody details are outlined in the

resource table.

Plasmid construct mutants

Myc-tagged LCK containing plasmid was generated using pENTR/D-TOPO cloning kit

(Thermo Scientific) according to manufacturer instructions. Briefly, Myc-LCK gene block

was purchased from Integrated DNA Technologies (IDT, USA). Myc-LCK was cloned

into pENTR/D-TOPO vector. The entry clone was further transferred into a destination

vector, pLenti CMV Puro DEST (Addgene). The plasmid was validated by DNA

sequencing (Eurofins). LCK mutants 192F, Y394F, 273R were generated by site

directed mutagenesis and sequenced. Each mutant was cloned into a lenti viral vector,

pLenti CMV Puro DEST (Addgene) for subsequent use.

Lentivirus production

Lentiviral particles for LCK silencing were generated using established lab protocols[15].

Briefly HEK293T cells were seeded into 6 well plates. The next day cells were

transfected with pRSV-Rev, pMDLg/pRRE, pMD2.G and lentivral vector expressing

shRNA for targeting LCK (KD1, TRCN0000426292, KD2, TRCN0000001599). Following

24h incubation, transfection media was replaced with fresh DMEM medium. 48h post

transfection, lentiviral particle containing media was filtered to remove cell debris and

added to CP70 and SKOV3 cells. Fresh media was subsequently added to the HEK-

293T transfection plates and incubated for an additional 24 hours followed by filtration

and addition to further CP70 and SKOV3 cells. Transduced CP70 and SKOV3 cells

were identified using 1.5ug/ml and 2ug/ml puromycin (Thermo Scientific) selection

respectively.

Generation of CRISPR/Cas9 KO cells

CP70 cells were used to generate LCK CRISPR/Cas9 knockout cells according to the

manufacturer protocol (Santa Cruz Biotechnology). Briefly, cells were transfected with

GFP labelled LCK CRISPR/Cas9 plasmid using lipofectamine 3000 (Thermo Scientific)

in the presence of antibiotic-free, FBS-enriched, Optimem media. Following

transfection, cells were kept in transfection medium for 24h, then replaced with fresh

culture media. After an additional 24h, transfected cells were screened for GFP

expression using a flow cytometer, and the GFP$^{+/high}$ population was isolated and plated

as single cells into a 96 well plate. Cells were grown and expanded in accordance with

standard culture techniques as stated above, followed by western blotting for LCK

protein expression with anti-LCK antibody (0.5 µg/mL, R & D Systems). Clones with the

lowest LCK expression compared to parental cells were considered LCK KO cells.

Western blot analysis

Western blot analysis was performed as reported with modifications as follows[37,38].

Briefly, cancer cells were washed with chilled Dulbecco's phosphate buffered saline

(PBS) two times at the end of treatment. NP-40 lysis buffer (Invitrogen) was added

dropwise to the plates and placed on ice for 10 minutes. The NP-40 lysis buffer contains

50 mM Tris, pH 7.4, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1 mM Na3VO4, 1%

Nonidet™ P40 (NP40), 0.02% NaN3 and was supplemented with 1mM PMSF and 2

µg/ml protease cocktail inhibitor (PCI) (Sigma Aldrich). Cells were then collected in a 1.5

mL centrifuge tube by scraping, and incubated on ice for one hour with occasional

vortexing. Lysates were centrifuged at 10,000rpm for 10 min at 4°C. Protein

concentration was measured of each lysate supernatant by BCA kit analysis (Thermo

Scientific). Protein samples were then prepared using 6x Laemmli dye containing BME

(β-mercapto ethanol) and boiled for 5-10min. Protein samples were subjected to SDS-

PAGE gel electrophoresis using pre-made gradient gels (4-20%, Biorad). Proteins were

transferred by wet transfer to a PVDF membrane (Millipore) at 4°C overnight.

Membranes were then blocked in 5% BSA in TBST for one hour at room temperature,

and subsequently, incubated overnight at 4°C in the following primary antibodies: T-LCK

(1:1000 R&D Systems), T-LCK (1:1000 Proteintech), P-LCK 394 (1:1000 R&D

Systems), RAD51 (1:1000, Proteintech), BRCA1 (1:500, EMD Millipore), BRCA2 (1:500,

EMD Millipore), γH2AX (1:1000 Cell Signaling Technology), GAPDH (1:5000

Proteintech), and β-actin (1:4000 Proteintech). After primary antibody incubation

membranes were washed three times with TBST (Tris-buffered saline containing 0.1%

tween 20) washing buffer on a platform shaker. Membranes were incubated with HRP-

conjugated rabbit (1:25000) or mouse (1:25000) secondary antibodies for one hour at

room temperature, followed by three washes with TBST buffer. Chemiluminescence

reagent (PerkinElmer) was added to detect immobilized proteins in PVDF membranes

utilizing the ChemiDoc imaging system. Densitometry was performed using Image J

software.

Nuclear protein isolation and co-immunoprecipitation analysis

CP70 and SKOV3 cells transduced with Myc-tagged LCK were treated with etoposide

(10μM) or DMSO for 24h followed by replacement with fresh serum-enriched media for

an additional 24h. Cells were collected and washed with cold PBS two times, scraped

and centrifuged. Cell pellets were then lysed with cytoplasmic and nuclear extraction

buffers according to manufacturer protocols (NE-PER Nuclear and Cytoplasmic

Extraction Kit, Thermo Scientific). Protein concentrations of nuclear lysates were

estimated using the BCA method outlined above. For co-immunoprecipitation, nuclear

protein lysates were incubated with 3μg anti-Myc antibody (Proteintech) or 3μg control

antibody (Cell Signaling Technology) overnight at 4°C with gentle rocking. Pre-cleaned

protein A/G agarose beads (Thermo Scientific) were added to the lysates and incubated

for 4h at 4°C on a rotating mixer. Beads were then collected by centrifugation and

washed three times with chilled NP-40 lysis buffer. 6x Laemmli buffer (Alfa Aesar)

containing BME was added and beads were boiled for 5 minutes. Samples were

separated on SDS-PAGE and processed for western blot analysis as outlined above.

Further, LCK overexpression (OE) SKOV3 (without Myc tagged) cells were treated with

etoposide (10µM) for 24h. Then, serum-enriched media was added to replace drug-

containing media and kept for another 24h. Then, cells were collected, and nuclear

lysates were prepared. Further immunoprecipitation/co-immunoprecipitation was

performed after RAD51 pulled down as described above.

Gene conversion assay

Gene conversion assay or DR-GFP assay was performed according to reported

methods[39]. Human osteosarcoma U2OS cells stably transfected with DR-GFP plasmid

and I-SceI endonuclease expression vector pCBASce were kindly provided by Maria

Jasin at Memorial Sloan-Kettering Cancer Center. Cells were treated with, PP2 (5, 7,

and 10µM) or DMSO vehicle for 48h. Cells were then transfected with I-SceI

endonuclease expression vector pCBASce using Lipofectamine 3000. In a separate set

of experiments, U2OS cells with DR-GFP integration were transfected with shCon, LCK

KD1 or KD2 for 24h and incubated in serum enriched medium for another 24h. Cells

were further transfected with I-SceI plasmid for 24h followed by incubation with serum

enriched medium for 24h. Live cells (Live/Dead dye kit, Thermo Scientific) were

analyzed with a flow cytometer to estimate the percentage of GFP-positive cells.

RAD51 and γH2AX nuclei staining and analysis

Laser scanning confocal microscopy was performed to detect RAD51 and γH2AX foci in

cancer cells. Briefly, ovarian cancer cell, CP70 were grown on coverslips and treated

with 10µM etoposide or vehicle for 24h followed by an additional 24h in drug-free media.

Coverslips were washed with PBS and fixed with 4% paraformaldehyde (Electron

Microscopy Sciences) in PBS. Cells were permeabilized with 0.01% triton-X 100 (Fisher

Scientific) for 5 minutes followed by a wash with chilled PBS and blocked with 3% goat

serum (Thermo Scientific) for 1h at room temperature. Cells were incubated with anti-

RAD51 (1:250, Abcam) or anti-γH2AX (1:300, Cell Signaling Technology) antibodies

overnight at 4°C in a humidified chamber. Next, cover slips were washed 3X with PBS.

Alexa fluorescent conjugated secondary antibodies were added to the coverslips and

incubated for 1h. Coverslips were washed 3x and mounted with DAPI containing

Vectashield (Vector Lab). Images were captured by confocal microscope at 63x

magnification in oil emersion (Leica SP8 confocal microscope). RAD51 and γH2AX foci

were counted on 20 representative cells by image J software.

Metaphase spread analysis

Metaphase spread analysis was performed on LCKi and PARPi treated cells using

established methods [27]. Briefly, cells were treated with LCK inhibitor, PP2 (5µM)

(Selleck Chemicals) and PARP inhibitor, olaparib (3µM) (Selleck Chemicals) for 48h.

After treatment, cells were harvested. Cells were treated with colcemid (50ng/ml)

(Sigma) for 1.5h then washed with PBS and placed in 0.075 mol/L KCl (Sigma) solution

for 20min. Subsequently, cells were washed with PBS and fixed in carnoy fixative

solution (Methanol: acetic acid 3:1) added dropwise followed by one hour incubation.

Cell pellets were collected, and fixative solution was added and incubated at 4°C for

24h. Cell pellets were collected and small amount fixative solution was added and cell

suspension was slowly dropped on glass slides and allowed to dry at 37°C. Slides were

then stained with Giemsa solution (Sigma Aldrich). Images were captured at 100X

magnification with a bright field microscope. Abnormalities in chromosomes were

quantified (Chromosomal break, gap, radial formation) by visually counting five nuclei per treatment group.

Single Cell Electrophoresis Assay

Single cell electrophoresis assay or comet assay was performed according to a previously reported method[40]. This experiment was performed following the manufacturer's instructions (Trevigen). Briefly, CP70 and SKOV3 eEOC cells were treated with 5µM PP2 and/or 3µM Olaparib for 48h. Comet LMAgarose was melted at 90°C for 10min in a water bath then cooled for 20min to 37°C. Treated cells were detached from plates using trypsin. Serum enriched media was added to neutralize the trypsin. Cell suspension was washed twice with chilled 1X Ca^{++} and Mg^{++} free PBS, and subsequently suspended at 1 X 10^5 cells/ml in chilled 1X PBS buffer (Free of Ca++ and Mg++). For the alkaline comet assay, the cell suspension was mixed with molten LMAgarose at 37°C. Immediately, 50µL LMAgarose mix was spread on glass microscope slides and incubated at 4°C for 30 min in the dark. Slides were incubated in lysis solution (provided in kit) overnight at 4°C. The next day comet slides were incubated in alkaline unwinding solution at room temperature for 20 min. Agarose gel electrophoresis (21volts for 30min) was performed using alkaline electrophoresis protocol. After electrophoresis, slides were briefly immersed in distilled water twice and then immersed in 70% ethyl alcohol for 5 min. Slides were dried and 100µl of SYBR gold (excitation/emission is 496 nm/522 nm) was added on agarose and kept for 30 min in the dark. Slides were washed with water, dried, and visualized by fluorescence microscopy.

Colony formation assay

The pharmacological effect of olaparib in cancer cells was investigated by colony

formation assay according to the earlier reported method[27]. Briefly, ovarian cancer cells

CP70 and SKOV3 cells (WT, sh, LCK OE, LCK KO, LCK KD and mutants) were placed

on 12 well plates. The next day, cells were treated with olaparib in a dose dependent

manner for 12 days. During this time the media was changed every day using fresh

drug. At the end of experiment, PBS was added to the well to wash the colony. Then,

cells were fixed with 4% paraformaldehyde solution for 10min. Cells were then washed

two times with PBS and incubated in 0.2% crystal violet solution for one hour at room

temperature. After incubation cells were washed three times with PBS. Then, the

images of six well plates were captured and number of colony forming area was

analyzed by Image J software.

Cell titer glo viability assay

CP70 (WT, OE, KD, KO and LCK mutants, Y394F, K273R and Y192F) cells were

collected after trypsinization. Cells were counted and 4000 cells were plated in each

well of a 96 well plate. Cells were then treated with etoposide for 48h in a dose

dependent manner. Control cells were treated with vehicle (DMSO). After the drug

treatment, Cell TiterGlo® Luminescent Cell viability assay reagent (Promega) mixture

was prepared and added to the cells to lyse them for 10 minutes shaking and

luminescence was measured via luminometer. Cell viability percentage was calculated

as luminescence of treated group/luminescence of vehicle treated group×100.

NHEJ gene expression

To check for NHEJ expression, CP70 and SKOV3 cells were grown on 60mm petri

dishes until 70% confluent. Cells were then treated with PP2 in a dose dependent

manner (5, 7 and 10μM) for 48h and collected by scraping and Western blot was

performed to assess protein expression of NHEJ markers Ku70 and Ku80.

***In vivo* animal study in NSG mice**

In vivo antitumor efficacy of Olaparib was tested in NSG (NOD severe combined

immunodeficient (SCID) IL2R gamma) mice. This study was approved (IACUC

Protocol# 2707) by Institutional Animal Care and Use Committee (IACUC), Cleveland

Clinic Lerner Research Institute. Animals were procured from BRU (Biological

Response Unit) facility of the Cleveland Clinic. All cells used in this study were

transfected (Lentiviral transfection) with pCDH-EF1a-eFFly-mCherry plasmid. Mice were

injected with CP70 LCK knockout cells or CP70 LCK overexpression cells

intraperitoneally (Cells: 0.5×10^6). After that mice were divided into four groups-**1**. Mice:

CP70 LCK KO: Vehicle (n=8), **2**. Mice: CP70 LCK KO: Olaparib (50mg/kg) (n=8), **3**.

Mice: CP70 LCK OE: Vehicle (n=8), **4.** Mice: CP70 LCK KO: Olaparib (50mg/kg) (n=8).

Olaparib was dissolved in ddH$_2$O containing 4% DMSO and 30% polyethelene glycol

(PEG300) and injected intraperitoneally. Animals were treated with olaparib for

5days/week. Bioluminescence of the tumor were measured by *in vivo* imaging system

(IVIS Spectrum CT, PerkinElmer). At the end of the experiment, mice were sacrificed

according to the protocol of Institutional Animal Care and Use Committee (IACUC),

Cleveland Clinic Lerner Research Institute. Tumor tissues were collected and preserved

in 10% formalin solution.

Immunohistochemistry and TUNEL assay

Immunohistochemical (IHC) analysis and TUNEL assay of tumor sections were

performed according to the earlier reported method[41]. Formalin fixed tissues were sent

to histology core to make thin slice (around 5micron) of tissue embedded on glass

slides. Next, slides were dipped in Histo-Clear to deparaffinize. Then sections were

rehydrated in gradient ethanol (100%, 95%, 80% and 60% ethanol 5 min for each bath).

For antigen retrieval, sections were then put in Tris-EDTA buffer (pH9) and boiled in a

pressure cooker for 10min. Then slides were put for an hour to cool. Slides were then

put into 0.1% triton X-100 solution for 10 min to permeabilize. After washing with PBS

three times, tissue sections were blocked with blocking buffer (5% goat serum and 0.1%

triton X-100 in 1X TBST) at room temperature for one hour. Then antibodies were added

to the tissues and incubated for overnight at 4°C. Next day, sections were washed three

times with PBS. Then diluted 100 µl Peroxidase Labeled Polymer was added to the

tissues and incubated for 30 min at room temperature. Now sections were washed three

times with PBS. Then diluted DAB chromogen was added to the tissue and incubated

for 2-5min until desired color was generated. Sections were washed three times and

then hematoxylin staining was performed. After washing with PBS, tissue sections were

dried and mounted with Cytoseal. Images were captured using upright microscope at

20x magnification.

For TUNEL assay, tissue sections were processed for antigen retrieval and

permeabilization as discussed earlier. Sections were washed three times with PBS.

Then enzyme solution was prepared according to the manufacturer instructions. Slides

were then incubated in the enzyme for 60min at 37°C. Sections were washed three

times with PBS and mounted with Vectashield containing DAPI.

Software and Statistics

Graph pad prism software was utilized for graph preparation and to determine statistical

significance (detailed in each figure legend). Image J was used for quantification of

data. Each experiment was performed at least three times. p-value less than 0.05 was

considered significant.

Acknowledgments

Authors thank the members of Reizes and Lathia Laboratories for helping to improve the scientific quality of the manuscript. We would like to thank Alex Myers for extensive editing the manuscript. We thank Dr. Gauravi Deshpande for helping in image acquisition by confocal microscopy, Amy Graham and Eric Schultz for flow cytometry, and Drs. Debjit Khan and Krishnendu Khan from Fox Laboratory for insights on complex formation assays. We would also like to thank Dr. Suparna Mazumdar (Cleveland Clinic) for giving constructive comments of this study. Research in the Reizes laboratory is funded by Center of Research Excellence in Gynecologic Cancer from the Cleveland Clinic Foundation, VeloSano Bike to Cure, and the Laura J. Fogarty Endowed Chair for Uterine Cancer Research. Dr. Gong is supported by and NIH/NCI grant (R01 CA222195).

References

633   1.   Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2022. *CA Cancer J*
634   *Clin* 2022, **72**(1)**:** 7-33.
635

636   2.   Yang L, Zhang Y, Shan W, Hu Z, Yuan J, Pi J*, et al.* Repression of BET activity
637   sensitizes homologous recombination-proficient cancers to PARP inhibition. *Sci*
638   *Transl Med* 2017, **9**(400).
639

640   3.   Konstantinopoulos PA, Ceccaldi R, Shapiro GI, D'Andrea AD. Homologous
641   Recombination Deficiency: Exploiting the Fundamental Vulnerability of Ovarian
642   Cancer. *Cancer Discov* 2015, **5**(11)**:** 1137-1154.
643

644   4.   O'Connor MJ. Targeting the DNA Damage Response in Cancer. *Mol Cell* 2015,
645   **60**(4)**:** 547-560.
646

647   5.   Moore K, Colombo N, Scambia G, Kim BG, Oaknin A, Friedlander M*, et al.*
648   Maintenance Olaparib in Patients with Newly Diagnosed Advanced Ovarian
649   Cancer. *N Engl J Med* 2018, **379**(26)**:** 2495-2505.
650

651   6.   Galluzzi L, Senovilla L, Vitale I, Michels J, Martins I, Kepp O*, et al.* Molecular
652   mechanisms of cisplatin resistance. *Oncogene* 2012, **31**(15)**:** 1869-1883.
653

654   7.   Tumiati M, Hietanen S, Hynninen J, Pietila E, Farkkila A, Kaipio K*, et al.* A
655   Functional Homologous Recombination Assay Predicts Primary Chemotherapy
656   Response and Long-Term Survival in Ovarian Cancer Patients. *Clin Cancer Res*
657   2018, **24**(18)**:** 4482-4493.
658

659   8.   Johnson N, Li YC, Walton ZE, Cheng KA, Li D, Rodig SJ*, et al.* Compromised
660   CDK1 activity sensitizes BRCA-proficient cancers to PARP inhibition. *Nat Med*
661   2011, **17**(7)**:** 875-882.
662

663   9.   Alagpulinsa DA, Ayyadevara S, Yaccoby S, Shmookler Reis RJ. A Cyclin-
664   Dependent Kinase Inhibitor, Dinaciclib, Impairs Homologous Recombination and
665   Sensitizes Multiple Myeloma Cells to PARP Inhibition. *Mol Cancer Ther* 2016,
666   **15**(2)**:** 241-250.
667

668   10.   Abbotts R, Topper MJ, Biondi C, Fontaine D, Goswami R, Stojanovic L*, et al.*
669   DNA methyltransferase inhibitors induce a BRCAness phenotype that sensitizes
670   NSCLC to PARP inhibitor and ionizing radiation. *Proc Natl Acad Sci U S A* 2019,
671   **116**(45)**:** 22609-22618.
672

673   11.   Johnson SF, Cruz C, Greifenberg AK, Dust S, Stover DG, Chi D*, et al.* CDK12
674   Inhibition Reverses De Novo and Acquired PARP Inhibitor Resistance in BRCA
675   Wild-Type and Mutated Models of Triple-Negative Breast Cancer. *Cell Rep* 2016,
676   **17**(9)**:** 2367-2381.

678   12.   Meinhold-Heerlein I, Fotopoulou C, Harter P, Kurzeder C, Mustea A, Wimberger
679   P*, et al.* The new WHO classification of ovarian, fallopian tube, and primary
680   peritoneal cancer and its clinical implications. *Arch Gynecol Obstet* 2016, **293**(4)**:**
681   695-700.
682

683   13.   Winterhoff B, Hamidi H, Wang C, Kalli KR, Fridley BL, Dering J*, et al.* Molecular
684   classification of high grade endometrioid and clear cell ovarian cancer using
685   TCGA gene expression signatures. *Gynecol Oncol* 2016, **141**(1)**:** 95-100.
686

687   14.   Ren T, Sun TT, Wang S, Sun J, Xiang Y, Shen K*, et al.* Clinical analysis of
688   chemo-resistance risk factors in endometriosis associated ovarian cancer. *J
689   Ovarian Res* 2018, **11**(1)**:** 40.
690

691   15.   Saygin C, Wiechert A, Rao VS, Alluri R, Connor E, Thiagarajan PS*, et al.* CD55
692   regulates self-renewal and cisplatin resistance in endometrioid tumors. *J Exp
693   Med* 2017, **214**(9)**:** 2715-2732.
694

695   16.   Crean-Tate KK, Braley C, Dey G, Esakov E, Saygin C, Trestan A*, et al.*
696   Pretreatment with LCK inhibitors chemosensitizes cisplatin-resistant
697   endometrioid ovarian tumors. *J Ovarian Res* 2021, **14**(1)**:** 55.
698

699   17.   Zhu X, Kim JL, Newcomb JR, Rose PE, Stover DR, Toledo LM*, et al.* Structural
700   analysis of the lymphocyte-specific kinase Lck in complex with non-selective and
701   Src family selective kinase inhibitors. *Structure* 1999, **7**(6)**:** 651-661.
702

703   18.   Nyakeriga AM, Garg H, Joshi A. TCR-induced T cell activation leads to
704   simultaneous phosphorylation at Y505 and Y394 of p56(lck) residues. *Cytometry
705   A* 2012, **81**(9)**:** 797-805.
706

707   19.   Nakanishi K, Cavallo F, Brunet E, Jasin M. Homologous recombination assay for
708   interstrand cross-link repair. *Methods Mol Biol* 2011, **745:** 283-291.
709

710   20.   Scully R, Panday A, Elango R, Willis NA. DNA double-strand break repair-
711   pathway choice in somatic mammalian cells. *Nat Rev Mol Cell Biol* 2019, **20**(11)**:**
712   698-714.
713

714   21.   Chang HHY, Pannunzio NR, Adachi N, Lieber MR. Non-homologous DNA end
715   joining and alternative pathways to double-strand break repair. *Nat Rev Mol Cell
716   Biol* 2017, **18**(8)**:** 495-506.
717

718   22.   Zhu S, Paydar M, Wang F, Li Y, Wang L, Barrette B*, et al.* Kinesin Kif2C in
719   regulation of DNA double strand break dynamics and repair. *Elife* 2020, **9**.
720

721 23. Liaunardy-Jopeace A, Murton BL, Mahesh M, Chin JW, James JR. Encoding
722   optical control in LCK kinase to quantitatively investigate its activity in live cells.
723   *Nat Struct Mol Biol* 2017, **24**(12)**:** 1155-1163.
724
725 24. Kastle M, Merten C, Hartig R, Kaehne T, Liaunardy-Jopeace A, Woessner NM*, et*
726   *al.* Tyrosine 192 within the SH2 domain of the Src-protein tyrosine kinase
727   p56(Lck) regulates T-cell activation independently of Lck/CD45 interactions. *Cell*
728   *Commun Signal* 2020, **18**(1)**:** 183.
729
730 25. Xiao M, Guo J, Xie L, Yang C, Gong L, Wang Z*, et al.* Let-7e Suppresses DNA
731   Damage Repair and Sensitizes Ovarian Cancer to Cisplatin through Targeting
732   PARP1. *Mol Cancer Res* 2020, **18**(3)**:** 436-447.
733
734 26. Lloyd RL, Wijnhoven PWG, Ramos-Montoya A, Wilson Z, Illuzzi G, Falenta K*, et*
735   *al.* Combined PARP and ATR inhibition potentiates genome instability and cell
736   death in ATM-deficient cancer cells. *Oncogene* 2020, **39**(25)**:** 4869-4883.
737
738 27. Kim H, George E, Ragland R, Rafail S, Zhang R, Krepler C*, et al.* Targeting the
739   ATR/CHK1 Axis with PARP Inhibition Results in Tumor Regression in BRCA-
740   Mutant Ovarian Cancer Models. *Clin Cancer Res* 2017, **23**(12)**:** 3097-3108.
741
742 28. Yasukawa M, Fujihara H, Fujimori H, Kawaguchi K, Yamada H, Nakayama R*, et*
743   *al.* Synergetic Effects of PARP Inhibitor AZD2281 and Cisplatin in Oral
744   Squamous Cell Carcinoma in Vitro and in Vivo. *Int J Mol Sci* 2016, **17**(3)**:** 272.
745
746 29. Kim RK, Yoon CH, Hyun KH, Lee H, An S, Park MJ*, et al.* Role of lymphocyte-
747   specific protein tyrosine kinase (LCK) in the expansion of glioma-initiating cells
748   by fractionated radiation. *Biochem Biophys Res Commun* 2010, **402**(4)**:** 631-636.
749
750 30. Bijlmakers MJ, Marsh M. Trafficking of an acylated cytosolic protein: newly
751   synthesized p56(lck) travels to the plasma membrane via the exocytic pathway. *J*
752   *Cell Biol* 1999, **145**(3)**:** 457-468.
753
754 31. Venkitachalam S, Chueh FY, Yu CL. Nuclear localization of lymphocyte-specific
755   protein tyrosine kinase (Lck) and its role in regulating LIM domain only 2 (Lmo2)
756   gene. *Biochem Biophys Res Commun* 2012, **417**(3)**:** 1058-1062.
757
758 32. Ibrahim YH, Garcia-Garcia C, Serra V, He L, Torres-Lockhart K, Prat A*, et al.*
759   PI3K inhibition impairs BRCA1/2 expression and sensitizes BRCA-proficient
760   triple-negative breast cancer to PARP inhibition. *Cancer Discov* 2012, **2**(11)**:**
761   1036-1047.
762
763 33. Choi YE, Battelli C, Watson J, Liu J, Curtis J, Morse AN*, et al.* Sublethal
764   concentrations of 17-AAG suppress homologous recombination DNA repair and
765   enhance sensitivity to carboplatin and olaparib in HR proficient ovarian cancer
766   cells. *Oncotarget* 2014, **5**(9)**:** 2678-2687.

34. Lim JJ, Yang K, Taylor-Harding B, Wiedemeyer WR, Buckanovich RJ. VEGFR3 inhibition chemosensitizes ovarian cancer stemlike cells through down-regulation of BRCA1 and BRCA2. *Neoplasia* 2014, **16**(4)**:** 343-353 e341-342.

35. Yap TA, Kristeleit R, Michalarea V, Pettitt SJ, Lim JSJ, Carreira S*, et al.* Phase I Trial of the PARP Inhibitor Olaparib and AKT Inhibitor Capivasertib in Patients with BRCA1/2- and Non-BRCA1/2-Mutant Cancers. *Cancer Discov* 2020, **10**(10)**:** 1528-1543.

36. Best OG, Gardiner AC, Majid A, Walewska R, Austen B, Skowronska A*, et al.* A novel functional assay using etoposide plus nutlin-3a detects and distinguishes between ATM and TP53 mutations in CLL. *Leukemia* 2008, **22**(7)**:** 1456-1459.

37. Jimenez-Pascual A, Hale JS, Kordowski A, Pugh J, Silver DJ, Bayik D*, et al.* ADAMDEC1 Maintains a Growth Factor Signaling Loop in Cancer Stem Cells. *Cancer Discov* 2019, **9**(11)**:** 1574-1589.

38. Bharti R, Dey G, Das AK, Mandal M. Differential expression of IL-6/IL-6R and MAO-A regulates invasion/angiogenesis in breast cancer. *Br J Cancer* 2018, **118**(11)**:** 1442-1452.

39. Weinstock DM, Nakanishi K, Helgadottir HR, Jasin M. Assaying double-strand break repair pathway choice in mammalian cells using a targeted endonuclease or the RAG recombinase. *Methods Enzymol* 2006, **409:** 524-540.

40. Pillay N, Tighe A, Nelson L, Littler S, Coulson-Gilmer C, Bah N*, et al.* DNA Replication Vulnerabilities Render Ovarian Cancer Cells Sensitive to Poly(ADP-Ribose) Glycohydrolase Inhibitors. *Cancer Cell* 2019, **35**(3)**:** 519-533 e518.

41. Bharti R, Dey G, Ojha PK, Rajput S, Jaganathan SK, Sen R*, et al.* Diacerein-mediated inhibition of IL-6/IL-6R signaling induces apoptotic effects on breast cancer. *Oncogene* 2016, **35**(30)**:** 3965-3975.

809 **KEY RESOURCES TABLE**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-LCK antibody	Proteintech	12477-1-AP
Anti-LCK antibody	R&D System	MAB3704
Anti-P-LCK (Y394) antibody	R & D system	MAB7500
Anti-RAD51 antibody	Proteintech	14961-1-AP
Anti-RAD51 antibody	Santa Cruz	sc-398587
Anti-P-H2A.X (Ser139) Antibody	Cell signaling	2577S
Anti-BRCA1 antibody	EMD Millipore	OP-92-100UG
Anti-BRCA2 antibody	EMD Millipore	OP-95-100UG
Anti-Myc antibody	Proteintech	60003-2-Ig
Anti-GAPDH antibody	Proteintech	HRP-60004
Anti-Lamin A/C antibody	Proteintech	10298-1-AP
Alpha Tubulin Monoclonal antibody	Proteintech	HRP-66031
Anti-Ku70 antibody	Proteintech	10723-1-AP
Anti-Ku80 antibody	Proteintech	16389-1-AP
Rabbit IgG XP® Isotype Control	Cell signaling	3900S
Mouse IgG XP® Isotype Control	Cell signaling	5415S
Anti-Rabbit IgG (H+L), HRP antibody	Promega	W4018
Anti-Mouse IgG (H+L), HRP antibody	Promega	W4028
Goat anti-Rabbit IgG Alexa Fluor 488	Thermo	A32731
Goat anti-mouse IgG Alexa Fluor 568	Thermo	A11031
Chemicals and reagents		
Precision Plus Protein™ Kaleidoscope	Biorad	1610375
Olaparib	Selleck chemicals	S1060
PP2	Selleck chemicals	S7008
Etoposide	Selleck chemicals	S1225
Cisplatin	Fesenius Kabi	401572I
Colcemid	Sigma	10295892001
Laemmli SDS sample buffer, reducing (6X)	Alfa Aesar	J61337
Pierce™ Protein A/G Plus Agarose	Thermo	20423
Precision Plus Protein™ Kaleidoscope™	1610375	Biorad
SYBR™ Gold Nucleic Acid Gel Stain	Thermo	S11494
Trypan Blue	Fisher scientific	25900CI
Immobilon-P PVDF Membrane	Merck Millipore	IPVH00010
Critical Commercial Assays		
VECTASHIELD® Mounting Medium	Vector lab	H-1200
NP40 Lysis buffer	Thermo	FNN0021
Pierce™ IP Lysis Buffer	Thermo	87788
Comet Assay Kit	Trevigen	4250-050-K
NuPAGE™ Protein Gel	Thermo	NP0329BOX
4–20% Mini-PROTEAN Protein Gels	Biorad	4568096
Pierce BCA Protein Assay Kit	Thermo	23225
Lipofectamin 3000	Thermo	L3000001
CellTiter-Glo® 2.0 Cell Viability Assay	Promega	G9241

Live/Dead assay kit	Thermo	L23105
KaryoMAX™ Giemsa Stain Solution	Thermo	10092013
Giemsa Stain, Modified Solution	32884-1L	Sigma
NP40 lysis buffer	Invitrogen	FNN0021
Protease Inhibitor Cocktail	Sigma	04693159001
Cytoplasmic and nuclear protein isolation kit	Thermo	78835
Experimental Models: Cell Lines		
CP70	Dr. Analisa Difeo	NA
SKOV3	ATCC	NA
U2OS	Dr. Zihua Gong	NA
U2OS DRGFP reporter cells	Dr. Zihua Gong	NA
HEK293T	ATCC	NA
CRL1978	ATCC	NA
Recombinant DNA		
pLenti CMV Puro DEST	Addgene	NA
Myc-LCK pLenti CMV Puro DEST	In house	
Myc-LCK Y394F pLenti CMV Puro DEST	In house	NA
Myc-LCK K273R pLenti CMV Puro DEST	In house	NA
Myc-LCK Y192F pLenti CMV Puro DEST	In house	NA
I-SceI plasmid	Dr. Zihua Gong	NA
ShRNA targeting LCK	Sigma Aldrich	TRCN0000426292
ShRNA targeting LCK	Sigma Aldrich	TRCN0000001600
ShRNA targeting LCK	Sigma Aldrich	TRCN0000001598
ShRNA targeting LCK	Sigma Aldrich	TRCN0000001599
LCK CRISPR/Cas9 KO Plasmid	Santa Cruz	SC-400434-KO-2
Software		
FlowJo	BD Bioscience	NA
Graph Pad prism	www.graphpad.com	NA
ImageJ	imagej.nih.gov	NA

814 **Figures**

Fig. 1

Fig 1: LCK modulates expression of HR repair proteins. (A) Western blot of CP70 and SKOV3 cells

containing various lentiviral LCK KD, KO, and OE to determine effects on LCK, BRCA1, BRCA2 and

RAD51 expression. **(B and C)** CP70 EV and CP70 LCK OE cells were treated with cycloheximide in a

time dependent manner. Then, immunoblot analysis was performed to evaluate the expression of RAD51

and BRCA1 proteins. Half-lives were determined from digitized images. **(D)** Western blot analysis of

CP70 and SKOV3 LCK OE cells treated with PP2 in a dose dependent manner for 48hrs, demonstrating

the effects of a pharmacological inhibitor of LCK on P-LCK, T-LCK, BRCA1, BRCA2, RAD51 and γH2AX

protein expression.

Fig 2: LCK inhibition attenuates the HR repair pathway in eEOC cells. (A) Schematic of DNA repair

assay in U2OS osteosarcoma cells stably transduced with the DR-GFP reporter system. This reporter

system contains upstream gene-encoding mutated GFP, and downstream truncated GFP. Transfection of

I-SceI endonuclease induces double strand breaks in the upstream gene. Following efficient HR repair,

GFP expression is restored and can be utilized to indicate HR-efficient cells for quantification. **(B)** U2OS

cells were treated with varying concentrations of the LCKi PP2 for 48h or transfected with Sh Con, LCK

KD1 or KD2 for 24h and incubated in serum enriched medium for another 24h. Cells were then subjected

to DR-GFP assay. **(C)** CP70 and SKOV3 cells were treated with PP2 or Olaparib for 48h and immunoblot

experiment was performed to check PARylation. (**D**) CP70, and SKOV3 cells were treated with increasing

concentrations of PP2 for 48h and cells were harvested, lysed, and immunoblotted for Ku70 and Ku80

protein expression. GAPDH was used as loading control. **(E)** Schematic model summarizing LCK

inhibition specificity for HR DNA repair. (Unpaired t test, p*<0.05, p**<0.01, p***<0.001).

Fig. 3

Fig 3: LCK is activated in the nucleus in response to DNA damage inhibition suppresses BRCA1

expression during DNA damage response. **(A)** Western blot analysis of T-LCK, BRCA1 and γH2AX

expression following 24hrs etoposide or MMS treatment, followed by 24hrs recovery in CP70 cells. **(B)**

Etoposide treated CP70 LCK OE cells were used for cytoplasmic and nuclear proteins extraction followed

by western blot analysis. **(C)** Immunofluorescence study of etoposide treated CP70 cells. **(D)** Western blot

analysis of etoposide/DMSO treated CP70 cells (Sh Con or LCK KD1 and KD2). **(E)** Western blot analysis

of etoposide/DMSO treated CP70 cells (WT and LCK KO).

Fig. 4

Fig 4: LCK regulates DNA damage and repair. (A and B) CP70 WT, LCK KO via CRISPR/Cas9 (KO),

and LCK OE on CRISPR background (OE) cells were treated with DMSO/etoposide 10μM for 24h. Then

cells were kept in drug-free media for another 24h. Immunofluorescence was performed to visualize

γH2AX foci formation in different groups. Scale bar represents 10μm. γH2AX foci was counted by image J

software and 20 cells were counted and plotted. **(C and D)** CP70 WT, LCK KO via CRISPR/Cas9 (KO),

and LCK OE on CRISPR background (OE) cells were treated with DMSO/etoposide 10μM for 24h. Cells

were incubated in drug free media for another 24h. Immunofluorescence was performed to visualize

RAD51 foci formation in different groups. RAD51 foci in 20 cells were counted by image J software. Level

of significance indicated on graph as determined by Graph pad Prism software ($p^* < 0.05$, $p^{**} < 0.01$,

$p^{***} < 0.001$).

Fig. 5

Fig 5: LCK interacts with RAD51 and BRCA1 during DNA damage response. (A) Schematic of Myc

tagged LCK construct. **(B)** Myc tagged LCK expressing CP70 cells were treated with vehicle (DMSO) or

etoposide. Cells were harvested, lysed, and nuclei were purified. Immunoprecipitation with Myc followed

by immunoblotting for Myc, BRCA1, and RAD51. **(C)** Myc tagged LCK expressing SKOV3 cells were

treated with vehicle or etoposide. Myc protein was pull down to determine the interaction of LCK with

RAD51 and BRCA1 by co-immunoprecipitation study. **(D)** LCK overexpressing SKOV3 cells were treated

with etoposide. RAD51 was pulled down from protein lysate. Then the expression of LCK and RAD51

was checked in pulled down protein sample by co-immunoprecipitation study. **(E)**   Schematic of LCK

binding partners during DNA damage response.

Fig. 6

Fig 6: LCK kinase activity is essential for HR repair. (A) Structure of MYC labelled LCK construct.

LCK Y394F, LCK K273R, LCK Y192F mutants were generated by site directed mutagenesis. These

mutants were further introduced in CP70 cells having LCK KO background. **(B)** CP70 cells (MYC tagged

LCK, LCK Y394F, LCK K273R, and LCK Y192F) were treated with etoposide/DMSO for 24h. Then cells

were put in drug free media for 24h. Cells were collected and nuclear proteins were isolated. MYC

antibodies were used for co-immunoprecipitation study to visualize the complex formation with RAD51

and BRCA1 proteins. **(C, E)** CP70 cells (LCK OE, LCK Y394F, LCK K273R, and LCK Y192F) were

treated with etoposide for 24h**.** Cells were then kept in drug-free media for 24h. Immunofluorescence was

performed to visualize γH2AX and RAD51 foci formation. **(D, F)** Quantification of γH2AX and RAD51 foci

formation in CP70 cells by Image J software. **(G)** CP70 cells (WT, LCK OE, LCK KO, LCK Y394F, LCK

K273R, and LCK Y192F) were treated with etoposide in a dose dependent manner for 48h. Then cell titer

glow viability assay was performed to check cell viability. **(H)** CP70 cells (LCK, LCK Y394F, LCK K273R,

and LCK Y192F in LCK knock out background) were grown on cover slips and treated with etoposide for

24h followed by incubation for 0, 2, 4, 8 and 24h. Cells were then subjected to immunofluorescence

analysis to visualize γH2AX foci formation. Then γH2AX foci were counted and plotted to visualize the

H2AX decay kinetics. $p* < 0.05$, $p** < 0.01$, $p*** < 0.001$ based on Graphpad Prism analysis.

Fig. 7

Fig 7: LCKi promotes genomic instability and augments PARPi induced genomic instability in ovarian cancer cells. **(A)** Single cell electrophoresis or COMET assay was used to validate and independently quantify the observed DNA damages. CP70 and SKOV3 cells were treated with PP2 5µM and/or Olaparib 3µM for 48h. Cells (1×10^5 Cells/ml) were then collected and mixed with LMAgarose (1:10 V/V). Then, 50L of LMAgarose solution was put on COMET slide and subjected to single-cell gel electrophoresis. **(B)** Extent of DNA damage was estimated based on measurement of COMET tail area using Image J. **(C)** CP70 and SKOV3 cells treated with PP2 5µM and/or Olaparib 3µM for 48h then had the chromosomal aberration assay performed. The arrow indicates the presence of abnormalities in chromosomes including breaks, gaps, and radials. **(D)** Abnormities in chromosomes were quantified

(Chromosomal break, gap, radial formation) by counting by visual observation. **(E)** CP70 and SKOV3

cells were treated with Olaparib and PP2 for 48h. Cells were harvested, lysed, and blotted for BRCA1 and

γH2AX. **(F)** CP70 Parental (WT), KO, and OE (in KO background) cells were treated with Olaparib in

dose dependent manner for 12 days. To identify colonies, plates were stained with crystal violet and

images were captured. Colonies formed were counted and plotted as percentage of control formation in

the gra. p*<0.05, p**<0.01, p***<0.001 based on Graph pad Prism analysis.

Fig 8: Disruption of LCK leads to inhibition ovarian tumor treated with Olaparib (A) Schematic

model of animal study. CP70 LCK KO and LCK OE cells were injected intraperitoneally in NSG mice.

After 12 days, PARP inhibitor Olaparib i.p. (50mg/kg) was administered 5days/week. **(B)** After three

weeks of Olaparib treatment, tumor volume was detected by IVIS imaging. **(C)** Tumor growth kinetics. **(D)**

TUNEL assay to detect DNA fragmentation in tumor tissue sections. TUNEL positive cells were counted

from five images and plotted in graph (Supplemental file S11 A). **(E)** IHC staining of γH2AX of tumor

sections from different groups. γH2AX positive cells were counted from five images and plotted in graph

(Supplemental file S11 B). **(F)** CD31 expression (Indicator of microvessel density and growth) of tumor

sections from different group of mice. Microvessel density was counted from five images and plotted in

graph (Supplemental file S11 C). Images are representative of two tumors from each cohort. We

quantified the staining from 5 fields from each mouse. Images were captured at 20X magnification.

Supplementary File

Supplementary Fig. S1

Supplementary Fig. S1: (A) Protein quantification T-LCK, BRCA1, BRCA2 and RAD51 in shCon, LCK

KD1, and LCK KD2 CP70 cells as observed in western blot analysis (Main Fig.1A, Left panel). **(B)**

Quantification of T-LCK, BRCA1, BRCA2 and RAD51proteins in shCon, LCK KD1, and LCK KD2 SKOV3

cells (Main Fig.1A, second panel from left). **(C)** Quantification of T-LCK, BRCA1, BRCA2, RAD51 protein

expression in CP70 WT and LCK KO cells (Main Fig.1A, third panel from left). **(D and E)** Quantification of

T-LCK, BRCA1, BRCA2 and RAD51 proteins in CP70 EV and LCK OE (Main Fig.1A, fourth panel from

left), and SKOV3 EV and LCK OE cells (Main Fig.1A, fifth panel from left). **(F)** Stability study of BRCA1

and RAD51 proteins in CP70 EV and LCK OE cells. **(G and H)** Protein quantification of P-LCK, BRCA1,

BRCA2, RAD51 and γH2AX in CP70 LCK OE and SKOV3 LCK OE cells treated with PP2 for 48h in a

dose dependent manner (Main Fig. 1D).

Supplementary Fig. S2

Supplementary Fig. S2: (A) Pharmacological inhibition of LCK attenuates HR repair proteins in ovarian cancer cells. CP70, SKOV3 and CRL1978 cells were treated with the LCKi, PP2 for 48h. Cells were harvested, lysed, and analyzed by immunoblot to assess protein expression of BRCA1, BRCA2, RAD51, and γH2AX. **(B)** Western blot analysis in different cells to check the expression of LCK and BRCA1. U2OS is osteosarcoma cell line which was used in DR GFP assay. U2OS and U2OS: DR-GFP cells were examined for checking LCK and BRCA1 expression. These cells were also found to express LCK and BRCA1 like CP70 and SKOV3 cells. **(C)** CRL1978 cells were treated with increasing concentrations of PP2 for 48h and cells were harvested, lysed, and immunoblotted for Ku70, and Ku80 protein expression. GAPDH was used as loading control.

Supplementary Fig. S3

Supplementary Fig. S3 (A, B) Protein quantification of T-LCK, BRCA1, and γH2AX in CP70 cells treated

with etoposide and MMS (Main Fig. 2A). **(C)** Quantification of P-LCK and T-LCK in CP70 LCK OE cells

treated with etoposide (Main Fig. 2B). **(D)** Quantification of BRCA1 protein in CP70 shCon, LCK KD1 and

LCK KD2 cells treated with/without etoposide (Main Fig. 3D). **(E)** Quantification of BRCA1 protein

expression in CP70 WT and LCK KO cells treated with/without etoposide (Main Fig. 3E).

Supplementary Fig. S4

Supplementary Fig. S4: **(A)** SKOV3 cells were treated with etoposide, and MMS for 24. After that cells

were put in 24h in drug free media. Cells were then subjected to western blot analysis for checking

protein expression. **(B)** DNA damage by UV radiation upregulates LCK phosphorylation. CP70 cells were

treated with UV radiation for 1min, 2min and 4min. Cells were kept in serum enriched media for 24h.

Then cells were subjected to western blot analysis to check the expression of P-LCK, T-LCK and BRCA1

expression. **(C)** SKOV3 LCK OE cells were treated with etoposide for 24h. Cells were then put in drug

free media for another 24h. Cells were collected, and cytoplasmic and nuclear proteins were extracted for

western blot analysis.

Supplementary Fig. S5

1001

1002

1003

1004

Supplementary Fig. S5 (A) CP70 WT, LCK KO (CRISPR/Cas9) and LCK OE (In CRISPR background)

cells were treated with DMSO/etoposide 10µM for 24h. Then cells were kept in drug free media for

another 24h. Then immunofluorescence study was performed to visualize γH2AX foci formation in

different groups. **(B)** CP70 WT, LCK KO (CRISPR/Cas9) and LCK OE (In CRISPR background) cells

were treated with DMSO/etoposide 10µM for 24h. Cells were put in drug free media for another 24h.

Then immunofluorescence study was performed to visualize RAD51 foci formation in different groups.

1011

1012

1013

1014 **Supplementary Fig. S6**

1015

1016

1017 **Supplementary Fig. S6:** Transfection efficiency of Myc tagged LCK in SKOV3 and CP70. SKOV3 and

1018 CP70 cells were transduced with EV or Myc tagged LCK plasmid by using lentiviral particle. Then, cells

1019 were checked for Myc and LCK expression.

1020

1021

1022

1023

1024

1025

1026

1027 **Supplementary Fig. S7**

1028

1029

1030

Supplementary Fig. S7: (A) CP70 cells (LCK OE, LCK Y394F, LCK K273R, and LCK Y192F, all

constructs were introduced into the CP70 LCK KO cells) were treated with etoposide for 24h**.** Cells were

then kept in drug free media for 24h. Immunofluorescence study was performed to visualize γH2AX foci

formation. **(B)** CP70 cells (LCK OE, LCK Y394F, LCK K273R, and LCK Y192F) were treated with

etoposide for 24h**.** Cells were then kept in drug free media for 24h. Immunofluorescence study was

performed to visualize RAD51 foci formation.

1037

1038

1039

1040

1041 **Supplementary Fig. S8**

1042

1043

1044 **Supplementary Fig. S8:** CP70 cells (LCK, LCK Y394F, LCK K273R, and LCK Y192F

1045 in LCK knock out background) were grown on cover slips and treated with etoposide for

1046 24h followed by incubation for 0, 2, 4, 8 and 24h. Cells were then subjected to

1047 immunofluorescence analysis to visualize H2AX foci formation.

1048

1049

1050

1051 **Supplementary Fig. S9**

A Ola(μM)   0   0.25   0.5   1   1.5   2

WT

KO

OE

B

CP70 cells	IC$_{50}$: Ola (μM)
WT	0.97 (95% CI is 0.6371 to 1.294)
OE	1.34 (95% CI is 0.8383 to 1.903)
KO	0.13 95% CI is -0.4857 to 0.5532

1052

1053

1054 **Supplementary Fig. S9: (A)** CP70 Parental cells and CP70 LCK KO and CP70 CD55

1055 OE (In KO background) cells were treated with Olaparib in dose dependent manner for

1056 12 days. After that colonies were stained with crystal violet and images were captured.

1057 **(B)** Number of Colony formation was counted and plotted as percentage of colony

1058 formation in the graph (Main fig 6G). IC50 values were shown in the table.

1059

1060

1061   **Supplementary Fig. S10**

1062

1063   **Supplementary Fig. S10: (A)** CP70 Sh Con or LCK knock down cells were treated with Olaparib in a
1064   dose dependent manner for 12 days. Number of colonies was counted and plotted using graph pad
1065   prism. **(B)** SKOV3 Sh Con or LCK knock down cells were treated with Olaparib in dose dependent
1066   manner for 12 days. Number of Colony formation was counted and plotted in the graph.

1067

1068

1069

1070

1071

1072

1073

1074 **Supplementary Fig. S11**

1075

Supplementary Fig. S11: (A) TUNEL assay to detect DNA fragmentation in tumor tissue sections.

TUNEL positive cells were counted from five images and plotted in graph (Main fig. 8D). **(B)** IHC staining

of ☐H2AX of tumor sections from different groups. γH2AX positive cells were counted from five images

and plotted in graph (Main fig. 8E).). **(C)** CD31 expression (Indicator of microvessel density and growth)

of tumor sections from different group of mice. Microvessel density was counted from five images and

plotted in graph (Main fig. 8F). Images are representative of two tumors from each cohort. We quantified

the staining from 5 fields from each mouse. Images were captured at 20X magnification.

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105