Suppression of PIK3CA-driven epileptiform activity by acute pathway control

Achira Roy[1,*], Victor Z. Han[1,2], Angela M. Bard[1], Devin T. Wehle[1,3], Stephen E. P. Smith[1,4], Jan-Marino Ramirez[1,4,5,6], Franck Kalume[1,5,7], Kathleen J. Millen[1,4,*]

[1]Center for Integrative Brain Research, Seattle Children's Research Institute, Seattle, Washington, USA; [3]Graduate Program in Neuroscience and Departments of [2]Biology, [4]Pediatrics, [5]Neurological Surgery, [6]Physiology and Biophysics, and [7]Pharmacology, University of Washington, Seattle, Washington, USA.

*Correspondence: Kathleen J. Millen, PhD

Seattle Children's Research Institute,

Center for Integrative Brain Research

1900 9th Avenue, Mailstop: JMB-10,

Seattle, WA, USA. ZIP: 98101

Email: kathleen.millen@seattlechildrens.org; Tel: +1-206-884-3225

Achira Roy, PhD

Seattle Children's Research Institute

Center for Integrative Brain Research

1900 9th Avenue, Mailstop: JMB-10,

Seattle, WA, USA. ZIP: 98101

24   Email: achira.roy@gmail.com; Tel: +1-206-884-5490

Competing Interests Statement: All authors read and approved the final manuscript.
None of the authors have competing interests.

Short title: Acute suppression of PIK3CA-driven epilepsy

Keywords: PI3K; epilepsy; mouse model; preclinical drug screens; BKM120; RAD001;
AZD5363; intractable; pyramidal neuron; hippocampus; electrophysiology

Number of Figures: 5

Number of Tables: 1

Number of Figure supplements: 7

ABSTRACT

Patients harboring mutations in the PI3K-AKT-MTOR signaling pathway often develop a spectrum of neurodevelopmental disorders including epilepsy. A significant proportion of them remain unresponsive to conventional anti-seizure medications. Understanding mutation-specific pathophysiology is thus critical for molecularly targeted therapies. We previously determined that mouse models expressing patient-related activating mutation in *PIK3CA* are epileptic and acutely treatable with PI3K inhibition, irrespective of dysmorphology (Roy et al. 2015). Using the same mutant model, we have now identified physiological mechanisms underlying the dysregulated neuronal excitability and its acute attenuation. We show that Pik3ca-driven hyperexcitability in hippocampal pyramidal neurons is mediated by changes in multiple non-synaptic, cell-intrinsic properties. These are distinct from mechanisms driving epilepsy in TSC/RHEB models. Further, we report that acute inhibition of PI3K or AKT, but not MTOR, suppresses the intrinsic epileptiform nature of the mutant neurons. These data represent an important step towards precision therapeutics against intractable epilepsy, using pathway drugs originally developed as anti-cancer agents.

INTRODUCTION

Mutations in the PI3K-AKT-MTOR signaling pathway, long studied for roles in cancer (Madsen 2020; Yang et al. 2019), also cause clinically important developmental brain overgrowth syndromes. Affected individuals display phenotypes ranging from dysplastic megalencephaly, hemimegalencephaly and focal cortical dysplasia, as well as comorbidities including hydrocephalus, autism and intellectual disability (Bast et al. 2006; Blumcke et al. 2011; Mirzaa 2018; Stafstrom and Carmant 2015; Dobyns and Mirzaa 2019; Crino 2016). These mutations also cause focal epilepsy, representing 25-50% of all cases of intractable (treatment-resistant) epilepsy in children (Bast et al. 2006; Blumcke et al. 2011; Kim and Lee 2019; Mirzaa 2018; Stafstrom and Carmant 2015). Traditionally, anti-seizure drugs targeting generic ion channels are identified by evoking acute seizures in wild-type rodents (Wilcox, West, and Metcalf 2020; Kehne et al. 2017; Barker-Haliski and White 2019), which do not mimic clinical epilepsies associated with specific genetic mutations and hence are largely ineffective. Since PI3K pathway mutations are now known to cause intractable epilepsy, repurposing pathway-targeted anti-cancer drugs offers a tantalizing opportunity to fundamentally shift the therapeutic approach towards intractable epilepsy. Presently, the MTOR inhibitor rapamycin and its analogs are the sole pathway-related drugs used clinically to treat epilepsy, primarily in patients with rare *TSC1/2* deletion mutations which enhance downstream MTOR signaling(Stafstrom 2019; Crino 2016; Kim and Lee 2019). Yet, these treatments remain effective only to a modest degree, suggesting that MTOR activation may not be the sole arm of this complex signaling pathway accounting for all PI3K pathway-driven epilepsies (Cho 2011; Meng et al. 2013; Roy et al. 2015).

Here we assess acute mechanisms driving epileptiform activity in *Nestin-*

cre;Pik3ca^{E545K} mouse model harboring a patient-related activating mutation in *Pik3ca*,

encoding the p110a catalytic subunit of PI3K (Roy et al. 2015). We report that Pik3ca

overactivation causes intrinsic neuronal changes leading to hyperexcitability. We further

establish that this epileptiform activity is acutely dependent on AKT, but not MTOR,

regulation. Our study defines key aspects of acute neuronal dysregulation in an

important genetic epilepsy model, which we leverage to repurpose the available PI3K

pathway inhibitors against intractable epilepsy.

RESULTS

In vivo recordings demonstrate network hyperexcitability in mutant CA1

Pik3ca mutant mouse models recapitulate patient phenotypes including brain overgrowth, cortical dysplasia, hydrocephalus and epilepsy, with phenotypic severity dependent on the mutant allele and its time of activation (Roy et al. 2019; Roy et al. 2015). Moreover, this developmental epilepsy is dissociable from dysmorphology and acutely suppressible by 1-hour *in vivo* administration of the pan-PI3K inhibitor BKM120 (Maira et al. 2012; Roy et al. 2015). To identify the underlying dynamic mechanisms, we performed *in vivo* and *in vitro* electrophysiological studies primarily focused on hippocampal pyramidal neurons, since the connection between epilepsy and hippocampal pathophysiology is well documented in patients and model systems, including ours (Roy, Millen, and Kapur 2020; Chatzikonstantinou 2014; Mazumder, Patial, and Singh 2019; Berdichevsky et al. 2013; Roy et al. 2015).

We investigated *in vivo* changes in the hippocampal CA1 network activity of ~P70 *Nestin-cre;Pik3ca^{E545K}* mutant mice relative to control littermates using local field potential (LFP) recordings, while simultaneously monitoring cortical surface activity by electrocorticography (ECoG). These recordings showed spontaneous interictal spike activity either restricted to the cortex or the hippocampus exclusively, or concurrent to the two regions in the mutant mice (Figure 1A-C; Figure 1 – figure supplement 1A-D). A broad spectrum of spike patterns was observed, consisting of single or groups of interictal spikes, the simplest identifiable unit of epileptiform activity (McCormick and Contreras 2001), as well as trains of sharp spikes, low-frequency slow waves or high-frequency low-amplitude "brushing events". These epileptiform events were observed

exclusively in the mutant mice. Within the mutant group, the event frequency was

significantly higher in the hippocampus than in the neocortex (Figure 1D). Power

spectrum analyses revealed that the mutant hippocampi exhibit significantly higher

power in the gamma frequency bands (Figure 1E-H), which is often indicative of seizure

onset (Hughes 2008; Lee, Spencer, and Spencer 2000). These observations indicate

that Pik3ca overactivation caused significant neural hyperexcitability in our mice,

predominantly in the hippocampus.

Mutant CA1 has elevated, acutely regulatable network excitability *in vitro*

To assess the physiological aspects of Pik3ca-driven epilepsy at the tissue level,

we performed hippocampal extracellular field recordings in acute forebrain slices. First,

we conducted these recordings in P16-20 CA1 pyramidal layer, across a potassium

concentration gradient in the recording buffer ($[K^+]$: 3 mM to 12 mM; Figure 2A-C). This

standard model shifts wild-type neuronal network from a silent state to tonic spiking and

then to bursting, followed by a depolarization block at very high levels of $[K^+]$, thus

mimicking a seizure-like response (Filatov et al. 2011). Compared to controls, mutant

slices exhibited a narrow range of hypersynchronous bursting and an elevated mean

baseline measured from the integrated trace, especially from 8mM to 12mM $[K^+]$ (Figure

2B,C; Figure 2 – figure supplement 2A). Mutant slices also showed significantly

enhanced relative peak amplitudes (at 8-9 mM $[K^+]$; Figure 2 – figure supplement 2B).

Thus, the mutation lowers the excitability threshold and a depolarization block is

reached faster than in controls. Further, acute extracellular application of BKM120

suppressed the hypersynchronous ictal bursts in the mutant CA1 field in 20-30 min, with

minimal physiological effect on control brains (Figure 2D,E). These data provide

evidence of a Pik3ca-driven active mechanism.

Mutant CA1 and CA3 pyramidal neurons are hyperexcitable

Next, we investigated mechanisms driving the mutant epileptiform activity at the

cellular level, using whole-cell patch-clamp recordings from CA1 and CA3 pyramidal

neurons in P16-20 control and mutant brains (Figure 3A). Silent and spontaneously

firing, tonic and burst-generating neurons were detected in both control and mutant

slices (Figure 3B,C). However, mutant slices had a significantly higher proportion of

burst-generating cells in both regions and lower proportion of tonic-firing cells in mutant

CA1 compared to controls (Figure 3D,E). Additionally, we observed significantly higher

tonic spike frequencies in mutant CA1 and CA3 and higher burst frequency in mutant

CA1, compared to controls (Figure 3F,G). Burst-generating cells demonstrated multiple

burst types. Specifically, we identified burst clusters and two types of plateau-bursts:

paroxysmal depolarization shift (PDS) and non-PDS waveforms (Figure 3 – figure

supplement 1A). We defined burst cluster as a multi-spike burst activity of random

nature without a plateau potential; these were only observed in mutant slices (Figure 3 –

figure supplement 1B). Plateau-bursts with depolarization shift resulting in sodium-spike

inactivation were termed as PDS "bursting cells". These depolarization shifts have

previously been implicated as the intracellular correlate of *in vivo* interictal spikes

(Marcuccilli et al. 2010; McCormick and Contreras 2001; Kubista, Boehm, and Hotka

2019; Tryba et al. 2019). We defined non-PDS bursts as those where plateau potential

developed in absence of prominent sodium-spike inactivation. No significant differences

between control and mutant hippocampal pyramidal cells were observed with respect to

average burst duration or inter-burst interval (Figure 3 – figure supplement 1C-E).

However, the plateau potential shift (PPS), defined here as the difference of the steady

state plateau potential and the resting membrane potential (RMP), was significantly

larger in mutant CA1 but not in CA3, relative to respective controls (Figure 3H,H'). This

was despite similar RMP across control and mutant hippocampal neurons (Figure 3 –

figure supplement 2A-C). The evoked current-clamp recordings further validated the

spontaneous activity results, especially by displaying significantly higher percentage of

burst-generating pyramidal neurons in the mutant hippocampus (Figure 3I-M).

Incidentally, no significant difference in membrane intrinsic properties like input

resistance, rheobase current and burst-threshold current, as well as evoked tonic spike

frequencies (for the tested 0-90pA range), was observed between control and mutant

cells (Table 1; Figure 3 – figure supplement 2D-F). But the decay time constant for

mutant CA3 neurons was significantly longer than that in mutant CA1 and control

groups (Table 1). With similar resistance, this implied that mutant CA3 neurons have

higher membrane capacitance than other cell groups. Together, these data demonstrate

that Pik3ca overactivation results in intracellular hyperexcitability, with some distinct cell

type-specific effects.

To determine whether the Pik3ca-related hyperexcitability is driven by altered

synaptic interactions or other intrinsic properties, we assessed the effects of channel

and receptor blockers on mutant hippocampal physiology. Blocking glutamatergic inputs

by extracellular administration of NMDA and non-NMDA receptor-antagonists, 3-((±)2-

carboxypiperazin-4yl) propyl-1-phosphate (CPP) and 6-cyano-7-nitroquinoxaline-2,3-

dione (CNQX) respectively, had no overt physiological effect on the majority of mutant

hippocampal neurons (Figure 4 – figure supplement 1A-C,E,F,H-K). Similarly, blocking

inhibitory synaptic inputs with gabazine did not significantly alter the mutant firing

patterns, spike frequencies or PPS (Figure 4 – figure supplement 1A,D,G,L). The

proportion of mutant CA1 neurons affected by these channel blockers was

comparatively less than that in CA3. Our data lead to the important conclusion that

Pik3ca-driven epileptiform activity is primarily not dependent on synaptic transmission.

In contrast, inhibition of calcium (Ca^{2+})-dependent inward current by extracellular

cadmium (Cd^{2+}) attenuated the paroxysmal bursts and reduced burst frequency and

PPS (Figure 4A-D), indicating a calcium channel-dependent mechanism underlying the

Pik3ca-dependent epileptiform activity. Intracellular cesium blocked potassium channels

and related currents, altering the intrinsic firing pattern in both control and mutant

hippocampal slices. Specifically, compared to regular baseline recordings, intracellular

cesium considerably reduced the proportion of tonic-firing cells in both CA1 and CA3

(Figure 4E-H, compare to Figure 3D,E). Unlike the spontaneous recordings, intracellular

cesium prompted the burst frequency in mutant CA1 to normalize and in mutant CA3 to

significantly rise, relative to respective controls (Figure 4I, compare to Figure 3G).

Intracellular cesium also normalized the mutant PPS to control levels (Figure 4J,

compare to Figure 3H). No overt effect on RMP or input resistance was seen in cesium-

treated control and mutant cells (Figure 4K,L). We conclude that Pik3ca-related

epileptiform activity is primarily caused by a complex set of altered non-synaptic cell-

intrinsic properties.

Acute treatment of epileptiform activity with PI3K pathway drugs

The acute suppression of epileptiform activity in *Nestin-cre;Pik3ca*^{E545K} mice by

BKM120, both *in vivo* (Roy et al. 2015) and at the tissue level, prompted us to dissect

the downstream pathway dynamics using inhibitors at the cellular level, in order to

coarsely determine their mechanistic roles and identify new therapeutic targets to treat

intractable epilepsy (Figure 5A). Acute extracellular administration of BKM120 or the

pan-AKT inhibitor AZD5363 (Davies et al. 2012) resulted in a large-scale and significant

alteration of firing pattern, frequencies and a relative reduction of PPS, sometimes

leading to gradual silencing of the recorded neurons (Figure 5B-I,I'). Neither drug had

any significant effect on neuronal RMP or tonic rheobase current (Figure 5 – figure

supplement 1A,B,E,F). In contrast to BKM120, acute AKT downregulation by AZD5363

significantly reduced the mutant burst duration and inter-burst interval (Figure 5 – figure

supplement 1C,D,G,H). Further, AZD5363 altered the physiology of all mutant CA3

neurons much faster (~8 min) than BKM120 (~32 min); however, both drugs changed

the intracellular activity of ~50% cells within 4 min (Figure 5 – figure supplement 1I,J).

AZD5363 also blocked spontaneous bursts in mutant neurons faster than BKM120

(Figure 5 – figure supplement 1K). Thus, acute regulation of both PI3K and AKT activity

directly suppressed *Pik3ca*^{E545K}-driven epileptiform neuronal activity. In contrast, acute

administration of the MTOR inhibitor RAD001 (everolimus) (Cho 2011; Krueger et al.

2010) showed no overt effect on the mutant RMP, tonic or burst firing patterns,

frequencies, average PPS, burst duration, inter-burst interval and rheobase current

(Figure 5J-M, Figure 5 – figure supplement 2A-D). This was despite western blot

confirmation that acute treatment of forebrain slices with RAD001 modulated direct

downstream targets of MTOR (Figure 5 – figure supplement 2E-G). In summary, our

study demonstrates that epileptiform activity caused by Pik3ca overactivation is based

on multiple intrinsic electrophysiological characteristics that are acutely modifiable by

downregulating PI3K and AKT activity, but not MTOR.

DISCUSSION

Activating mutations in the PI3K-AKT-MTOR pathway commonly cause pediatric

intractable epilepsy (Kim and Lee 2019; Dobyns and Mirzaa 2019); yet the underlying

mechanisms remain largely undefined. Using our *Nestin-cre;Pik3ca^{E545K}* model, we now

establish that Pik3ca-dependent neuronal hyperexcitability in hippocampal pyramidal

neurons is driven primarily by intrinsic neuronal mechanisms. Acute attenuation of

epileptiform activity by inhibiting AKT but not MTOR also reveals a previously unknown

role of PI3K signaling in neuronal homeostasis. Our mouse model of human intractable

epilepsy is now validated across whole animal, tissue, and cellular levels.

Epileptiform activity is typically marked by increased tonic depolarizations and/or

quasiperiodic bursts, mediated by intrinsic membrane properties, ephaptic or

synaptic/circuit-level interactions (McCormick and Contreras 2001; Dudek, Yasumura,

and Rash 1998). We confirmed enhanced interictal spike frequency and diverse,

synchronized epileptiform spike patterns in the mutant brains *in vivo*, predominantly in

the hippocampus. This is consistent with studies of *PIK3CA* variants in brain tumors and

mouse models with patient-related *Pik3r2* mutations, which are less common causes of

intractable epilepsy (Yu et al. 2020; Shi et al. 2020). Our *in vitro* analyses revealed

intracellular correlates of epileptiform activity and enhanced firing frequencies in the

mutant hippocampal pyramidal neurons. Relative to controls, *Pik3ca^{E545K}* neurons

exhibited significantly higher proportions of bursting cells and higher heterogeneity in

burst patterns, including burst clusters and PDSs. A prior study showed that human

neurons from surgically resected neocortical samples of pediatric patients with focal

cortical dysplasia (FCD) exhibited remarkably similar burst characteristics (Marcuccilli et

al. 2010). These samples were not genotyped; however, since most cases of FCD and

epilepsy result from variants in PI3K pathway genes (Mirzaa 2018), such cross-species

phenotypic correlation is remarkable and emphasizes the clinical relevance of our

mouse model.

The PI3K-AKT-MTOR pathway is also known to influence synaptogenesis and

neuronal plasticity (Sanchez-Alegria et al. 2018). For example, neocortical samples

from patient and mouse models with *TSC* mutations exhibit synaptic input-mediated

epileptogenic hyperexcitability (Wang et al. 2007). However, blocking glutamatergic

(NMDA and non-NMDA) or GABAergic synaptic inputs showed no significant effect on

the neuronal activity in our mutant brain slices. We acknowledge that altered neuronal

circuitry may be relevant to Pik3ca-mediated epilepsy, but our data demonstrate that

altered synaptic interactions are at least not primary mediators of dynamic hippocampal

neuronal hyperexcitability.

Significant reduction of burst-generating cells, burst frequency and PPS by

blocking voltage-dependent calcium or potassium channels indicate that $Pik3ca^{E545K}$-

driven epileptiform activity is predominantly mediated by multiple altered cell-intrinsic

properties. Yet, some cell-intrinsic properties, such as burst frequencies, PPS and time

constants, are distinctly different between mutant CA1 and CA3 neurons. Since all

neurons do not react uniformly to either pathway overactivation or Cd^{2+}/Cs^+-dependent

channel inhibition, current anti-seizure drugs that typically target single ion channels

cannot address the heterogeneous dysregulation, thus potentially explaining epileptic

intractability in this patient population. It is intriguing to note that intercellular signaling

heterogeneity involving PI3K-AKT-MTOR pathway was also reported by a recent

quantitative study that highlighted its potential relevance to human disorders, such as

insulin resistance (Norris et al. 2021).

PI3K-AKT-MTOR inhibitors developed to treat cancer potentially represent novel

anti-seizure therapeutics (Cardamone et al. 2014; Lindhurst et al. 2015; Maira et al.

2012; Zou et al. 2020; Forde et al. 2021; Venot et al. 2018). Rapamycin and its analogs,

that are currently in clinical use to treat epilepsy in TSC patients, curbed epileptiform

activity in mouse models harboring *TSC1/2*, *RHEB, MTOR* or *PTEN* mutations when

administered long-term (>7-day) (Curatolo and Moavero 2013; Crino 2016; Lasarge and

Danzer 2014; Stafstrom 2019; Kim and Lee 2019). In contrast, we show acute inhibition

of epileptiform activity by BKM120 *in vitro*, confirming our previous *in vivo* finding (Roy

et al. 2015). We also demonstrate that acute AKT inhibition with AZD5363 has similar

effects. This first preclinical study using AZD5363 to treat epilepsy shows that acute

Pik3ca-driven epileptiform mechanism is predominantly AKT-dependent, limiting roles

for other PI3K targets. Intriguingly, rapamycin analog RAD001 had minimal acute effect

in our model. While elevated MTOR activity is known to be epileptogenic (Crino 2016;

Kim and Lee 2019), our data suggest that alternate PI3K-AKT targets acutely regulate

neuronal hyperexcitability. This study establishes an important foundation to determine

active pathway-driven cellular mechanisms and to assess the plethora of available PI3K

pathway inhibitors, facilitating new molecularly rational therapeutic interventions for

intractable epilepsy.

MATERIALS AND METHODS

Mice

The following mouse lines were used: *Nestin-cre* (Jackson Labs, Bar Harbor, Maine, USA; Stock #003771, RRID: IMSR_JAX:003771), conditional *Pik3ca^{E545K}* knock-in (Robinson et al. 2012). *Nestin-cre* and *Pik3ca^{E545K}* mouse lines were maintained in C57BL/6 and FVB backgrounds respectively. We have designated *Nestin-cre;Pik3ca^{E545K}* conditional mutant mice as "*Pik3ca* mutants" or "mutants" throughout the manuscript.

 All mice were housed in Optimice cages with aspen bedding at the Seattle Children's Research Institute's specific pathogen-free (SPF) vivarium facility (light "ON": 6am – 8pm). Noon of the day of vaginal plug was designated as embryonic day 0.5 (E0.5). The day of birth was designated as postnatal day 0 (P0). Genotyping by PCR was done using separate sets of primers for the *Cre* coding region and the *Pik3ca^{E545K}* allele, as previously described(Roy et al. 2015). All mouse procedures were approved and conducted in accordance with the guidelines laid down by the Institutional Animal Care and Use Committees (IACUC) of Seattle Children's Research Institute, Seattle, WA, USA. ARRIVE guidelines have been followed for reporting work involving animal research.

In vivo electrophysiology

5 control and 6 *Nestin-cre;Pik3ca^{E545K}* mutant mice (age: ~P70) were used for *in vivo* regular and depth-electrode electrophysiology experiments. We saw no sex-dependent data correlation (Roy et al. 2015).

Electrode implantation surgery: Mice underwent survival surgery to implant ECoG,

EMG, and hippocampal depth electrodes under isoflurane anesthesia using similar

procedures as reported in our prior work (Bolea et al. 2019; Roy et al. 2015). The ECoG

electrodes consisted of silver wire (diameter: 130 μm bare; 180 μm coated) attached to

micro screws. The ECoG electrodes were implanted bilaterally through small cranial

burr holes above the somatosensory cortices. A similar reference electrode was placed

above the cerebellum following the same procedure. The depth electrodes were made

of 2 fine twisted tungsten wires (30um nylon coated, California fine wire) and implanted

in the hippocampal CA1 region (coordinates: -2.0 mm anteroposterior, 1.5 mm

mediolateral, 1.9 mm dorsoventral in reference to bregma) to record local field potential

(LFP). All electrodes were connected to interface connector and fixed to the skull with

dental cement (Lang Dental Manufacturing Co., Inc., Wheeling, IL). Mice were allowed

to recover from surgery for 2-3 days.

Video, electroencephalagraphy and local field potential (Video-EEG-LFP) recording:
Simultaneous video-EEG-LFP recordings were collected from conscious mice on

PowerLab 8/35 and 16/30 data acquisition units, using LabChart 7.3.3 software (AD

Instruments, Colorado Spring, USA). ~6 hours of baseline ECoG tracings were visually

reviewed for the presence of spontaneous epileptiform events, as previously studied

(Kalume 2013; Liautard et al. 2013; Roy et al. 2015). All bioelectrical signals were

acquired at a 20 KHz sampling rate. The ECoG signals were processed with a 1-70 Hz

bandpass filter and the LFP signal with a 5 Hz high-pass filter. Interictal spikes were

identified as transient, clearly distinguished from background activity, with pointed peak

and slow wave. Myoclonic seizures were identified as shock-like muscular jerks on

video, associated with a spike or polyspike-wave complex on EEG. Power spectral

analysis and visual inspection of the data were conducted to characterize the EEG

activity in different frequency bands and identify epileptiform event on the ECoG and

local CA1 field recordings using the Labchart 8.2 software.

In vitro electrophysiology

Slice preparation: Postnatal (P16-20) pups were anesthetized briefly in a closed

chamber by administering isoflurane (5% flow rate) or CO_2 (constant flow rate: 10-30%

of chamber vol/min); then perfused transcardially with ice-cold low Na^+-buffer ("slicing

solution," which included the following: 252 mM sucrose, 2 mM KCl, 2 mM $MgCl_2$, 2.6

mM $CaCl_2$, 1.2 mM NaH_2PO_4, 26 mM $NaHCO_3$, and 15 mM glucose, with the pH

adjusted to 7.4 and the osmolarity to 310 ± 5 mOsm). Brain was dissected out by

separating the head and cutting along the skull sutures using fine scissors and forceps.

The forebrain was isolated in ice-cold, oxygenated (95% O_2, 5% CO_2) slicing solution. A

slanted (~15° from vertical) agar block was secured on a specimen tray as a support for

the brain during slicing. The isolated forebrain was glued with cyanoacrylate in the

orientation depending on the type of slices we wanted; and then placed in the vibratome

for slicing. Slicing was proceeded until hippocampal landmark was visible. For coronal

slices (extracellular recordings), the forebrain was placed such that its antero-posterior

(A-P) axis was perpendicular to the specimen tray/vibratome blade, and olfactory bulb

distal to the tray surface. For horizontal slices (whole-cell recordings), the forebrain was

oriented such that it's A-P axis was parallel to the specimen tray and ventral side

(hippocampal side) was on the top, distal to the tray surface. Once the hippocampus

became visible, thick, acute slices (thickness: 350μm for extracellular recordings, 200-

250μm for whole-cell patch clamp recordings) were taken and incubated in ice-cold

slicing solution continuously oxygenated with 95% O_2/5% CO_2. After cutting, slices were

immediately returned to the same solution and maintained in a warm bath (28 ± 0.5°C)

for recovery. After 30 min, they were transferred into regular artificial cerebrospinal fluid

(aCSF), composed of the same components as slicing solution except for the

replacement of sucrose with 126 mM NaCl. Slices were then kept at room temperature,

continuously superfused with oxygenated aCSF, until recording. Given that $Pik3ca^{E545K}$

mutant is megalencephalic, a greater number of hippocampal/forebrain sections were

obtained. However, care was taken so that the dorso-ventral plane(s) used for recording

was always comparable between control and mutant slices. No randomization was

used. Tissue collection was not performed blind since the mice were subjected to

genotyping and drug administration.

Recording: Slices were transferred into a recording chamber, continuously superfused

with oxygenated aCSF, for extracellular or intracellular whole-cell recordings. The

pClamp software suite (Molecular Devices; RRID:SCR_011323) was used for data

acquisition and analysis. Signals were amplified (MultiClamp 700A, Axon Instruments,

Molecular Devices, USA), digitized (D1322A, Axon Instruments, Molecular Devices,

USA), and stored in a computer for post-hoc analysis.

   a)   *Extracellular recording:* To determine if hippocampal neuronal populations in

$Pik3ca^{E545K}$ mutants have different electrochemical signatures than controls,

extracellular population recordings were performed on CA1 hippocampal pyramidal

neuronal layer in acute 350 μm-thick coronal brain slices of ~P16-20 $Pik3ca^{E545K}$ mutant

and control littermates (n=6). To induce epileptiform activity in the slices, we increased

potassium concentration ([K$^+$]) (Borck and Jefferys 1999; Leschinger et al. 1993) from 3

mM to 12 mM, in 1mM increments at 30^{o}C. This standard model induces wild-type

neurons to transition from a silent neuronal state to tonic spiking and then to bursting,

followed by depolarization block at very high [K$^+$] (Filatov et al. 2011).

b)   *Intracellular recording:* Horizontal hemi-forebrain slices with prominent

hippocampus were used for intracellular whole-cell visual patch-clamp experiments.

Slices transferred to the recording chamber were maintained at 30-34°C, constantly

superfused with oxygenated aCSF. Borosilicate glass capillaries/pipettes for patch-

clamp recording had electrode resistance (R_e) optimally kept around 5-8 MΩ, after being

filled with internal solution containing the following: potassium gluconate (~132 mM),

KCl (5 mM), HEPES (10 mM), EGTA (5 mM), $CaCl_2.H_2O$ (0.5 mM), $MgCl_2$ (2 mM),

disodium phosphocreatine (5 mM), disodium-ATP (4 mM), trisodium-GTP (0.5 mM),

EGTA (5 mM). Cells were visualized under brightfield optics using the 40X water-

immersion objective of an upright microscope (Olympus, BX51WI). The patch electrode

was advanced toward the target cells by a micromanipulator (MP-225, Sutter Instrument

Company, USA) and 1 GΩ seal was established, typically by a small negative pressure,

with the membrane ruptured by gentle suction and/or zap pulses. Whole-cell patch-

clamp recording was performed from cell bodies of pyramidal neurons of hippocampal

CA1 and CA3 respectively. Following intracellular recording protocols were also used:

i.   Spontaneous gap-free recordings in current (I)-clamp: cells were typically tested

at resting conditions (without current injection) unless noticed otherwise.

ii.   Evoked I-clamp steps protocol: current steps start at -50 pA, incremental 10 pA,

duration; 300 ms; 15 steps (-90 to +50 pA) were recorded across experiments.

Acute chemical assays

Channel blockers: Chemicals inhibiting specific ion channels were introduced in the

bath/recording chamber during *in vitro* whole-cell recordings to compare different

components contributing to the neural activity in control and *Pik3ca^{E545K}* mutant mice. To

block all fast-synaptic excitatory transmission, 3-((±)2-carboxypiperazin-4yl) propyl-1-

phosphate (CPP, NMDA receptor antagonist, 20 µM; Tocris Bioscience, UK) and 6-

cyano-7-nitroquinoxaline-2,3-dione (CNQX, AMPA/kainite (non-NMDA) receptor

antagonist, 20 µM, diluted in DMSO; Alomone Labs, Israel) were introduced in the bath

(Lehmann et al. 1987; Lester et al. 1989). SR95531 hydrobromide (Gabazine, GABA$_A$

receptor antagonist, 10 µM, Tocris Bioscience, UK) were used to block inhibitory

synaptic transmission. Divalent cation Cd^{2+} ($CdCl_2$, 100 µM; Sigma-Aldrich, USA)

(Boissard et al. 2002) was used to depress synaptic transmission and block inward

calcium-selective current, isolating the outward K^+ current. To block voltage-dependent

K^+ channels, intracellular administration of cesium (Covey, Kauer, and Casseday 1996)

was done by replacing potassium gluconate with cesium gluconate in the internal

solution while maintaining the same osmolarity.

PI3K pathway drugs: Drugs regulating PI3K pathway were added to the circulating

recording buffer, thus getting acutely administered in the brain slice placed in the

recording chamber. For protein analyses, these were added at identical concentration to

the brain slices incubated in oxygenated aCSF for ~1hr. The following PI3K pathway

inhibitors were dissolved in 100% DMSO and used *in vitro*: pan-PI3K inhibitor BKM120

(Buparlisib; Novartis, Switzerland; working concentration: ~3.5 µM); pan-AKT inhibitor

AZD5363 (Capivasertib; Selleckchem, USA; working concentration: 0.5 µM); MTOR

inhibitor RAD001 (Everolimus, Chem Express Cat# 159351-69-6; working

concentration: 0.52 µM).

Western blotting

Horizontal forebrain slices from 5 control and 5 mutant mice were obtained and

recovered in oxygenated aCSF as detailed above. Slices showing nice hippocampal

morphology were selected, treated for 1 hour with DMSO, BKM120, AZD5363 and

RAD001 respectively, in the same concentrations used for recording. The treated slices

were then flash-frozen in liquid nitrogen and stored at -80°C for post-processing. These

brain slices were lysed in NP-40 lysis buffer (150 mM NaCl, 50 mM Tris (pH 7.4), 1%

NP-40, 10 mM NaF, 2 mM sodium orthovanadate + protease/phosphatase inhibitor

cocktails [Sigma, USA]). Samples were normalized to equal protein concentrations

(0.333 mg/ml) using Pierce BCA protein assay (Thermo Fisher Scientific, USA).

Standards were made through serial dilutions of 10 mg/ml BSA. Samples were diluted

into a final concentration of 1x Laemmli sample buffer and boiled at 95°C for ten

minutes. 7.5% 10-well gels were prepared, and samples were run at 120V for 1.5 hours.

Gels were transferred onto polyvinylidene difluoride (Millipore) membranes and run

either overnight at 25V or for 2 hours at 60V. Membranes were blocked in 4% milk in

TBST (0.05 M Tris, 0.15 M NaCl, pH7.2, 0.1% (v/v) Tween20) for one hour at room

temperature, and primary antibodies were applied overnight at 4°C in blocking medium.

Primary antibodies for western blots were diluted as follows: rabbit anti-Phospho-AKT

Ser473 (D9E) (Cell Signaling Technology, USA; RRID: AB_2797780; 1:2000), mouse

anti-Pan-AKT (40D4) (Cell Signaling Technology, USA; Cat# 2920; 1:2000), rabbit anti-

Phospho-S6 Ribosomal Protein Ser235/236 (Cell Signaling Technology, USA; RRID:

AB_2721245, 1:2000), mouse anti-S6 Ribosomal Protein (54D2) (Cell Signaling

Technology, USA; RRID: AB_2238583, 1:1000), rabbit anti-beta(b)-Actin (GeneTex,

USA; RRID: AB_1949572, 1:10000). After washing and probing respectively with goat

anti-rabbit (RRID: AB_2313567) and anti-mouse (RRID: AB_10015289) horseradish

peroxidase (HRP)-conjugated secondary antibodies (1:10000; Jackson

ImmunoResearch Labs, USA), blots were imaged using Femto chemiluminescent

detection reagents (Thermo Fisher Scientific, USA; Cat# 34095) in a FluorChem R

western blot imaging system (ProteinSimple, Bio-Techne, USA). 8-bit images were used

as a representative western blot. 16-bit images were used to quantify the intensity of

each band using ImageJ v1.53. Regions of interest were drawn in each sample's lane.

After quantifying the lane in a histogram, the peak representing the band of interest was

isolated and the area of the region was measured as the band's quantification.

Quantitative and Statistical Analyses

Number of mice used was consistent with previous experiments completed and

published by us and other investigators. For extracellular field potential analyses,

baseline potential from integrated traces were measured for each [K^+] and normalized to

the value at 3 mM for each slice per genotype; relative peak amplitude was similarly

measured for control and mutant slices for 3,7,8,9 mM of [K^+]. In whole-cell patch-clamp

recording, plateau potential shift (PPS) was calculated by subtracting the baseline

potential from the highest plateau/burst potential in a paroxysmal depolarizing event.

Burst duration and inter-burst interval were measured as shown in Figure 3 – figure

supplement 1 and averaged across multiple bursts per cell. Input resistance was

measured from evoked current-clamp recordings by dividing the voltage difference

(measured at -10pA to -30pA I-steps) by the current interval (20pA). The decay

membrane time constant was obtained by recording the membrane response to 20pA

hyperpolarizing current pulses (300 ms duration, 1 Hz) and fitting the response to a

single exponential curve. We chose 20pA of hyperpolarizing current because such a

current intensity did not produce sag. Evoked spike frequencies and rheobase current

were calculated exclusively from tonic-firing cells. Burst threshold current was

calculated from burst-generating cells as the first current step which induced burst. *In*

vitro electrophysiological analyses were performed using Clampfit 10.7 and 11 (pClamp,

Molecular Devices, USA). For Western Blots, each band intensity was recalculated

relative to its respective b-actin band intensity, and then normalized across average

intensity per protein lane. Normalized ratios of phosphoproteins over total proteins were

thereafter obtained.

Statistical significance was assessed using 2-tailed unpaired t-tests with Welch's

correction (EEG power analyses, whole-cell RMP, burst duration, inter-burst interval,

Cd^{2+} data), 2-tailed paired t-tests (field potential baseline measurement, rheobase

current, drug-treated analyses) and ANOVA followed by Tukey post-tests (EEG

interictal spike frequency, field potential relative peak amplitude; whole-cell tonic spike

and burst frequencies, cell proportions, PPS, evoked spike frequency and time

constant; Western blots). Normal distribution was assumed for the data analyses, when

required. For EEG-ECoG-LFP experiments, quantitative and statistical data analysis

was performed in Labchart 8.2 software (AD Instruments, Colorado Spring, USA) and

Igor Pro 6.37 (WaveMetrics Inc., USA); final graphs were made in GraphPad Prism v7.0

(GraphPad Software Inc., San Diego, USA). For the remaining data, statistical analyses

and graph plotting were done using GraphPad Prism v7.0 and Microsoft Excel.

Differences were considered significant at $p < 0.05$.

Acknowledgments

We thank Suzanne J. Baker for gifts of mouse lines (*Nestin-cre* and *Pik3ca*^{E545K}); Leon

Murphy (Novartis) for BKM120; Rory M. Murphy, Amanda P. Tran Hartman, Nikhil Sahai

and Jiyun Ryu for technical assistance; Aguan D. Wei and Paul Wakenight for

discussions. This work is funded by the NIH grants 1R01NS099027 (K.J.M.),

R01MH113545 (S.E.P.S.), R01NS102796 (F.K.), and CURE Sleep and Epilepsy Grant

(F.K.).

Author Contributions: A.R., F.K., J.M.R. and K.J.M. contributed to the study

conception and design. A.R., V.H., A.M.B., D.T.W. and F.K. contributed to the data

collection. A.R. and F.K. contributed to in vitro and in vivo data analyses respectively.

D.T.W. and S.E.P.S. contributed to Western blot data analyses and interpretation. A.R.,

F.K., J.M.R., K.J.M. contributed to the overall data interpretation. S.E.P.S., F.K. and

K.J.M. provided the funding resources. A.R. wrote the first draft of the manuscript and

others commented on previous versions of the manuscript. All authors read and

approved the final manuscript.

Ethics

Animal experimentation: All animal experimentation was conducted in accordance with

the guidelines laid down by the Institutional Animal Care and Use Committees (IACUC)

of Seattle Children's Research Institute, Seattle, WA, USA (protocol IDs: IACUC00006

(K.J.M.); IACUC00108 (F.K.)).

REFERENCES

Barker-Haliski, M., and H.S. White. 2019. 'Validated animal models for antiseizure drug (ASD) discovery: Advantages and potential pitfalls in ASD screening', *Neuropharmacology*: 107750.

Bast, T., G. Ramantani, A. Seitz, and D. Rating. 2006. 'Focal cortical dysplasia: prevalence, clinical presentation and epilepsy in children and adults', *Acta Neurol Scand*, 113: 72-81.

Berdichevsky, Y., A. M. Dryer, Y. Saponjian, M. M. Mahoney, C. A. Pimentel, C. A. Lucini, M. Usenovic, and K. J. Staley. 2013. 'PI3K-Akt signaling activates mTOR-mediated epileptogenesis in organotypic hippocampal culture model of post-traumatic epilepsy', *J Neurosci*, 33: 9056-67.

Blumcke, I., M. Thom, E. Aronica, D. D. Armstrong, H. V. Vinters, A. Palmini, T. S. Jacques, G. Avanzini, A. J. Barkovich, G. Battaglia, A. Becker, C. Cepeda, F. Cendes, N. Colombo, P. Crino, J. H. Cross, O. Delalande, F. Dubeau, J. Duncan, R. Guerrini, P. Kahane, G. Mathern, I. Najm, C. Ozkara, C. Raybaud, A. Represa, S. N. Roper, N. Salamon, A. Schulze-Bonhage, L. Tassi, A. Vezzani, and R. Spreafico. 2011. 'The clinicopathologic spectrum of focal cortical dysplasias: a consensus classification proposed by an ad hoc Task Force of the ILAE Diagnostic Methods Commission', *Epilepsia*, 52: 158-74.

Boissard, R., D. Gervasoni, M. H. Schmidt, B. Barbagli, P. Fort, and P. H. Luppi. 2002. 'The rat ponto-medullary network responsible for paradoxical sleep onset and maintenance: a combined microinjection and functional neuroanatomical study', *Eur J Neurosci*, 16: 1959-73.

Bolea, I., A. Gella, E. Sanz, P. Prada-Dacasa, F. Menardy, A. M. Bard, P. Machuca-Marquez, A. Eraso-Pichot, G. Modol-Caballero, X. Navarro, F. Kalume, and A. Quintana. 2019. 'Defined neuronal populations drive fatal phenotype in a mouse model of Leigh syndrome', *Elife*, 8.

Borck, C., and J. G. Jefferys. 1999. 'Seizure-like events in disinhibited ventral slices of adult rat hippocampus', *J Neurophysiol*, 82: 2130-42.

Cardamone, M., D. Flanagan, D. Mowat, S. E. Kennedy, M. Chopra, and J. A. Lawson. 2014. 'Mammalian target of rapamycin inhibitors for intractable epilepsy and subependymal giant cell astrocytomas in tuberous sclerosis complex', *J Pediatr*, 164: 1195-200.

Chatzikonstantinou, A. 2014. 'Epilepsy and the hippocampus', *Front Neurol Neurosci*, 34: 121-42.

Cho, C. H. 2011. 'Frontier of epilepsy research - mTOR signaling pathway', *Exp Mol Med*, 43: 231-74.

Covey, E., J. A. Kauer, and J. H. Casseday. 1996. 'Whole-cell patch-clamp recording reveals subthreshold sound-evoked postsynaptic currents in the inferior colliculus of awake bats', *J Neurosci*, 16: 3009-18.

Crino, P. B. 2016. 'The mTOR signalling cascade: paving new roads to cure neurological disease', *Nat Rev Neurol*, 12: 379-92.

Curatolo, P., and R. Moavero. 2013. 'mTOR inhibitors as a new therapeutic option for epilepsy', *Expert Rev Neurother*, 13: 627-38.

Davies, B. R., H. Greenwood, P. Dudley, C. Crafter, D. H. Yu, J. Zhang, J. Li, B. Gao, Q. Ji, J. Maynard, S. A. Ricketts, D. Cross, S. Cosulich, C. C. Chresta, K. Page, J. Yates, C. Lane, R. Watson, R. Luke, D. Ogilvie, and M. Pass. 2012. 'Preclinical pharmacology of AZD5363, an inhibitor of AKT: pharmacodynamics, antitumor activity, and correlation of monotherapy activity with genetic background', *Mol Cancer Ther*, 11: 873-87.

Dobyns, W. B., and G. M. Mirzaa. 2019. 'Megalencephaly syndromes associated with mutations of core components of the PI3K-AKT-MTOR pathway: PIK3CA, PIK3R2, AKT3, and MTOR', *Am J Med Genet C Semin Med Genet*, 181: 582-90.

Dudek, F. E., T. Yasumura, and J. E. Rash. 1998. ''Non-synaptic' mechanisms in seizures and epileptogenesis', *Cell Biol Int*, 22: 793-805.

Filatov, G., G. P. Krishnan, N. F. Rulkov, and M. Bazhenov. 2011. 'Dynamics of epileptiform activity in mouse hippocampal slices', *J Biol Phys*, 37: 347-60.

Forde, K., N. Resta, C. Ranieri, D. Rea, O. Kubassova, M. Hinton, K. A. Andrews, R. Semple, A. D. Irvine, and V. Dvorakova. 2021. 'Clinical experience with the AKT1 inhibitor miransertib in two children with PIK3CA-related overgrowth syndrome', *Orphanet J Rare Dis*, 16: 109.

Hughes, J. R. 2008. 'Gamma, fast, and ultrafast waves of the brain: their relationships with epilepsy and behavior', *Epilepsy Behav*, 13: 25-31.

Kalume, F. 2013. 'Sudden unexpected death in Dravet syndrome: respiratory and other physiological dysfunctions', *Respir Physiol Neurobiol*, 189: 324-8.

Kehne, J. H., B. D. Klein, S. Raeissi, and S. Sharma. 2017. 'The National Institute of Neurological Disorders and Stroke (NINDS) Epilepsy Therapy Screening Program (ETSP)', *Neurochem Res*.

Kim, J. K., and J. H. Lee. 2019. 'Mechanistic Target of Rapamycin Pathway in Epileptic Disorders', *J Korean Neurosurg Soc*, 62: 272-87.

Krueger, D. A., M. M. Care, K. Holland, K. Agricola, C. Tudor, P. Mangeshkar, K. A. Wilson, A. Byars, T. Sahmoud, and D. N. Franz. 2010. 'Everolimus for subependymal giant-cell astrocytomas in tuberous sclerosis', *N Engl J Med*, 363: 1801-11.

Kubista, H., S. Boehm, and M. Hotka. 2019. 'The Paroxysmal Depolarization Shift: Reconsidering Its Role in Epilepsy, Epileptogenesis and Beyond', *Int J Mol Sci*, 20.

Lasarge, C. L., and S. C. Danzer. 2014. 'Mechanisms regulating neuronal excitability and seizure development following mTOR pathway hyperactivation', *Front Mol Neurosci*, 7: 18.

Lee, S. A., D. D. Spencer, and S. S. Spencer. 2000. 'Intracranial EEG seizure-onset patterns in neocortical epilepsy', *Epilepsia*, 41: 297-307.

Lehmann, J., J. Schneider, S. McPherson, D. E. Murphy, P. Bernard, C. Tsai, D. A. Bennett, G. Pastor, D. J. Steel, C. Boehm, and et al. 1987. 'CPP, a selective N-methyl-D-aspartate (NMDA)-type receptor antagonist: characterization in vitro and in vivo', *J Pharmacol Exp Ther*, 240: 737-46.

Leschinger, A., J. Stabel, P. Igelmund, and U. Heinemann. 1993. 'Pharmacological and electrographic properties of epileptiform activity induced by elevated K+ and lowered Ca2+ and Mg2+ concentration in rat hippocampal slices', *Exp Brain Res*, 96: 230-40.

Lester, R. A., M. L. Quarum, J. D. Parker, E. Weber, and C. E. Jahr. 1989. 'Interaction of 6-cyano-7-nitroquinoxaline-2,3-dione with the N-methyl-D-aspartate receptor-associated glycine binding site', *Mol Pharmacol*, 35: 565-70.

Liautard, C., P. Scalmani, G. Carriero, M. de Curtis, S. Franceschetti, and M. Mantegazza. 2013. 'Hippocampal hyperexcitability and specific epileptiform activity in a mouse model of Dravet syndrome', *Epilepsia*, 54: 1251-61.

Lindhurst, M. J., M. R. Yourick, Y. Yu, R. E. Savage, D. Ferrari, and L. G. Biesecker. 2015. 'Repression of AKT signaling by ARQ 092 in cells and tissues from patients with Proteus syndrome', *Sci Rep*, 5: 17162.

Madsen, R. R. 2020. 'PI3K in stemness regulation: from development to cancer', *Biochem Soc Trans*, 48: 301-15.

Maira, S. M., S. Pecchi, A. Huang, M. Burger, M. Knapp, D. Sterker, C. Schnell, D. Guthy, T. Nagel, M. Wiesmann, S. Brachmann, C. Fritsch, M. Dorsch, P. Chene, K. Shoemaker, A. De Pover, D. Menezes, G. Martiny-Baron, D. Fabbro, C. J. Wilson, R. Schlegel, F. Hofmann, C. Garcia-Echeverria, W. R. Sellers, and C. F. Voliva. 2012. 'Identification and characterization of NVP-BKM120, an orally available pan-class I PI3-kinase inhibitor', *Mol Cancer Ther*, 11: 317-28.

Marcuccilli, C. J., A. K. Tryba, W. van Drongelen, H. Koch, J. C. Viemari, F. Pena-Ortega, E. L. Doren, P. Pytel, M. Chevalier, A. Mrejeru, M. H. Kohrman, R. E. Lasky, S. M. Lew, D. M. Frim, and J. M. Ramirez. 2010. 'Neuronal bursting properties in focal and parafocal regions in pediatric neocortical epilepsy stratified by histology', *J Clin Neurophysiol*, 27: 387-97.

Mazumder, A. G., V. Patial, and D. Singh. 2019. 'Mycophenolate mofetil contributes to downregulation of the hippocampal interleukin type 2 and 1beta mediated PI3K/AKT/mTOR pathway

hyperactivation and attenuates neurobehavioral comorbidities in a rat model of temporal lobe
epilepsy', *Brain Behav Immun*, 75: 84-93.
McCormick, D. A., and D. Contreras. 2001. 'On the cellular and network bases of epileptic seizures', *Annu*
Rev Physiol, 63: 815-46.
Meng, X. F., J. T. Yu, J. H. Song, S. Chi, and L. Tan. 2013. 'Role of the mTOR signaling pathway in epilepsy',
J Neurol Sci, 332: 4-15.
Mirzaa, G., Roy, A., Dobyns, W.B., Millen, K.J., Hevner, R.F. 2018. 'Hemimegalencephaly and Dysplastic
Megalencephaly' ' in Golden JA Adle-Biassette H, Harding BN (ed.), *Developmental*
Neuropathology (Wiley Blackwell).
Norris, D., P. Yang, S. Y. Shin, A. L. Kearney, H. J. Kim, T. Geddes, A. M. Senior, D. J. Fazakerley, L. K.
Nguyen, D. E. James, and J. G. Burchfield. 2021. 'Signaling Heterogeneity is Defined by Pathway
Architecture and Intercellular Variability in Protein Expression', *iScience*, 24: 102118.
Robinson, G., M. Parker, T. A. Kranenburg, C. Lu, X. Chen, L. Ding, T. N. Phoenix, E. Hedlund, L. Wei, X.
Zhu, N. Chalhoub, S. J. Baker, R. Huether, R. Kriwacki, N. Curley, R. Thiruvenkatam, J. Wang, G.
Wu, M. Rusch, X. Hong, J. Becksfort, P. Gupta, J. Ma, J. Easton, B. Vadodaria, A. Onar-Thomas, T.
Lin, S. Li, S. Pounds, S. Paugh, D. Zhao, D. Kawauchi, M. F. Roussel, D. Finkelstein, D. W. Ellison, C.
C. Lau, E. Bouffet, T. Hassall, S. Gururangan, R. Cohn, R. S. Fulton, L. L. Fulton, D. J. Dooling, K.
Ochoa, A. Gajjar, E. R. Mardis, R. K. Wilson, J. R. Downing, J. Zhang, and R. J. Gilbertson. 2012.
'Novel mutations target distinct subgroups of medulloblastoma', *Nature*, 488: 43-8.
Roy, A., K. J. Millen, and R. P. Kapur. 2020. 'Hippocampal granule cell dispersion: a non-specific finding in
pediatric patients with no history of seizures', *Acta Neuropathol Commun*, 8: 54.
Roy, A., R. M. Murphy, M. Deng, J. W. MacDonald, T. K. Bammler, K. A. Aldinger, I. A. Glass, and K. J.
Millen. 2019. 'PI3K-Yap activity drives cortical gyrification and hydrocephalus in mice', *Elife*, 8.
Roy, A., J. Skibo, F. Kalume, J. Ni, S. Rankin, Y. Lu, W. B. Dobyns, G. B. Mills, J. J. Zhao, S. J. Baker, and K. J.
Millen. 2015. 'Mouse models of human -related brain overgrowth have acutely treatable
epilepsy', *Elife*, 4:e12703.
Sanchez-Alegria, K., M. Flores-Leon, E. Avila-Munoz, N. Rodriguez-Corona, and C. Arias. 2018. 'PI3K
Signaling in Neurons: A Central Node for the Control of Multiple Functions', *Int J Mol Sci*, 19.
Shi, X., Y. Lim, A. K. Myers, B. L. Stallings, A. McCoy, J. Zeiger, J. Scheck, G. Cho, E. D. Marsh, G. M.
Mirzaa, T. Tao, and J. A. Golden. 2020. 'PIK3R2/Pik3r2 Activating Mutations Result in Brain
Overgrowth and EEG Changes', *Ann Neurol*, 88: 1077-94.
Stafstrom, C. E. 2019. *Developmental Epilepsy - From Clinical Medicine to Neurobiological Mechanisms*
(World Scientific).
Stafstrom, C. E., and L. Carmant. 2015. 'Seizures and epilepsy: an overview for neuroscientists', *Cold*
Spring Harb Perspect Med, 5.
Tryba, A. K., E. M. Merricks, S. Lee, T. Pham, S. Cho, D. R. Nordli, Jr., T. L. Eissa, R. R. Goodman, G. M.
McKhann, Jr., R. G. Emerson, C. A. Schevon, and W. van Drongelen. 2019. 'Role of paroxysmal
depolarization in focal seizure activity', *J Neurophysiol*, 122: 1861-73.
Venot, Q., T. Blanc, S. H. Rabia, L. Berteloot, S. Ladraa, J. P. Duong, E. Blanc, S. C. Johnson, C. Hoguin, O.
Boccara, S. Sarnacki, N. Boddaert, S. Pannier, F. Martinez, S. Magassa, J. Yamaguchi, B.
Knebelmann, P. Merville, N. Grenier, D. Joly, V. Cormier-Daire, C. Michot, C. Bole-Feysot, A.
Picard, V. Soupre, S. Lyonnet, J. Sadoine, L. Slimani, C. Chaussain, C. Laroche-Raynaud, L.
Guibaud, C. Broissand, J. Amiel, C. Legendre, F. Terzi, and G. Canaud. 2018. 'Targeted therapy in
patients with PIK3CA-related overgrowth syndrome', *Nature*, 558: 540-46.
Wang, Y., J. S. Greenwood, M. E. Calcagnotto, H. E. Kirsch, N. M. Barbaro, and S. C. Baraban. 2007.
'Neocortical hyperexcitability in a human case of tuberous sclerosis complex and mice lacking
neuronal expression of TSC1', *Ann Neurol*, 61: 139-52.

Wilcox, K. S., P. J. West, and C. S. Metcalf. 2020. 'The current approach of the Epilepsy Therapy
Screening Program contract site for identifying improved therapies for the treatment of
pharmacoresistant seizures in epilepsy', *Neuropharmacology*, 166: 107811.
Yang, J., J. Nie, X. Ma, Y. Wei, Y. Peng, and X. Wei. 2019. 'Targeting PI3K in cancer: mechanisms and
advances in clinical trials', *Mol Cancer*, 18: 26.
Yu, K., C. J. Lin, A. Hatcher, B. Lozzi, K. Kong, E. Huang-Hobbs, Y. T. Cheng, V. B. Beechar, W. Zhu, Y.
Zhang, F. Chen, G. B. Mills, C. A. Mohila, C. J. Creighton, J. L. Noebels, K. L. Scott, and B. Deneen.
2020. 'PIK3CA variants selectively initiate brain hyperactivity during gliomagenesis', *Nature*, 578:
166-71.
Zou, Z., T. Tao, H. Li, and X. Zhu. 2020. 'mTOR signaling pathway and mTOR inhibitors in cancer: progress
and challenges', *Cell Biosci*, 10: 31.

TABLE

Table 1: Summary table of intrinsic membrane properties

Summary of basic intrinsic membrane properties quantitated and compared between

control and mutant CA1 and CA3 neurons. No significant differences in RMP, input

resistance and rheobase current (minimum injected current required to elicit the first

action potential) were observed between control and mutant neurons, in both CA1 and

CA3. Decay time constant of mutant CA3 neurons was significantly longer than that of

control CA1 ($p < 0.0001$) and CA3 ($p = 0.0071$) cells, as well as of mutant CA1 neurons

($p < 0.0001$) ($F=31.21$, $df=33$). Data is represented as mean ± SEM; differences were

considered significant at $p<0.05$.

#	Intrinsic properties	CA1		CA3	
		Control	**Mutant**	**Control**	**Mutant**
1	RMP (mV)	-55.36 ± 0.58	-54.64 ± 0.64	-56.07 ± 0.82	-56.49 ± 0.92
2	Input resistance (MΩ)	284.27 ± 21.6	262.82 ± 22.72	331.73 ± 23.54	272.99 ± 39.14
3	Time constant (ms)	21.79 ± 1.76	23.58 ± 1.85	32.23 ± 7.33	44.20 ± 4.56
4	Rheobase (pA)	11.54 ± 3.17	18.95 ± 3.66	18.95 ± 3.32	19.33 ± 4.19

FIGURE LEGENDS

Figure 1: *Nestin-cre;Pik3ca^E545K* mutant brains show higher neural excitability

(**A**) Schematic shows electrode placement for EEG-ECoG recordings in ~P70 *Nestin-cre;Pik3ca^E545K* and control littermates. LFP, local field potential. (**B,C**) Compared to controls, the mutants showed significantly higher frequency of regional (r) spikes and train of spikes/polyspikes in neocortex (black traces) and hippocampus (blue traces). (**D**) In the mutant, interictal spike frequency was significantly higher in hippocampus (blue box), compared to those generated in neocortex (orange box) or generalized in both regions (green box) (F=46.65, degrees of freedom (df)=27). (**E,F**) Power spectrum analysis displayed increased activity in mutant, as emphasized in the representative magnified segments. (**G,H**) Mutant hippocampus demonstrated significant higher activity in the gamma frequency range (F=3.932, df=27; neocortex: F=26.93, df=27). Data is represented as mean ± SEM scatter plots; differences were considered significant at p<0.05; ns, not significant. Scalebars: 0.2s,20µV (B,C); 1s,50µV (E,F). See also Figure 1 – figure supplement 1.

Figure 2: Extracellular mutant CA1 recordings demonstrate hyper-synchronization

(**A**) Schematized flowchart summarizes the acute coronal brain slicing for extracellular field potential recording. (**B,C**) Representative control (black) and mutant (red) voltage and integrated voltage traces across a [K$^+$] gradient; comparison indicated significantly higher excitability in the mutant brain. (**D,E**) Acute extracellular BKM120 administration at constant 7mM [K$^+$] aCSF suppressed synchronized ictal bursts in mutant CA1 field

within ~20-30min, with minimal effect on the control hippocampal slices. Scalebars:

2min,0.5V (B-E). See also Figure 2 – figure supplement 1.

Figure 3: Mutant hippocampal neurons produce increased epileptiform burst

activity

(**A**) Flowchart shows acute horizontal brain slicing for whole-cell recording. (**B-E**) Traces

represent silent, tonic, and burst categories of CA1 and CA3 neurons based on

spontaneous cellular activity; respective pie charts marked proportion of recorded cells.

Mutant CA1 and CA3 exhibited significantly higher proportions of burst-firing cells

compared to controls. Significantly fewer tonic-firing cells were observed in mutant CA1.

(**F,G**) relative to controls, spontaneous tonic spike frequencies were significantly higher

in mutant CA1 (t=2.175, df=43.71) and CA3 (t=4.063, df=19.78) cells; burst frequency

was significantly higher in mutant CA1 (CA1: t=2.244, df=29.38; CA3: t=0.1561,

df=6.754). (**H**) Plateau potential shift (PPS) in mutant bursting cells, as depicted in (**H'**),

was significantly higher in CA1 (t=2.586, df=8.118) but similar in CA3 (t=0.2436,

df=5.110), compared to respective controls. (**I-L**) Representative evoked voltage traces

of control and mutant showed tonic and burst firing in response to current steps. (**M**)

Evoked recording also marked significantly higher bursting cell proportions in the

mutant. Data is represented as pie charts, % bar graphs and mean ± SEM scatter plots;

differences were considered significant at p<0.05; ns, not significant. Scalebars:

1s,20mV (B,D); 0.1s,20mV (H', I-L). See also Figure 3 – figure supplements 1-2.

Figure 4: Pik3ca-related burst characteristics are dependent on cell-intrinsic

calcium and potassium channel regulation

(**A-D**) Blocking calcium channels by extracellular cadmium (Cd^{2+}) significantly altered

intrinsic bursting pattern of mutant neurons including burst frequency (t=3.267, df=3)

and PPS (t=4.172, df=3). (**E-H**) Representative traces and respective pie charts show

types of cellular activity in control and mutant CA1 and CA3 neurons after administering

cesium intracellularly. In response to cesium, tonic firing cells were significantly reduced

and original differences in CA1 and CA3 cell proportions between control and mutant

were normalized. (**I**) Cs^+-induced burst frequency was significantly higher in mutant CA3

than those in respective controls (F= 26.31, df=21); unlike normal spontaneous state, no

such difference was observed in CA1. (**J-L**) Intracellular cesium lowered the

spontaneously enhanced mutant PPS in CA1 to control levels (F=4.003, df=20), while

having no effect on RMP (CA1: t=1.308, df=27.32; CA3: t=0.2616, df=6.816) or input

resistance (F= 2.936, df=42) in both control and mutant neurons. Data is represented as

pie charts, % bar graphs and mean ± SEM scatter plots; differences were considered

significant at p<0.05; ns, not significant. Scalebars: 1s,20mV (A,E,F). See also Figure 4

– figure supplement 1.

Figure 5: Acute inhibition of Pik3ca or Akt suppresses hyperactivity in mutant

cells

(**A**) Schematic of simplified PI3K-AKT-MTOR pathway, marking targets of action for

BKM120, AZD5363 and RAD001. Representative traces demonstrate differential action

of BKM120 (**B**), AZD5363 (**F**) and RAD001 (**J**) on *Nestin-cre;Pik3ca^{E545K}* hippocampal

neurons, ranging from being unaffected to change in frequency/pattern to silencing, as

quantitated respectively in (**C,G,K**). (**D,E,H,I**) BKM120 and AZD5363 significantly

reduced mutant burst frequency (BKM120: F=8.937, df=38; AZD5363: F=43.50, df=20)

and PPS (BKM120: F=24.39, df=36; AZD5363: t=6.386, df=6) in both CA1 and CA3.

(**E',I'**) Insets show activity traces of mutant bursting cell before (red) and after BKM120

(blue) or AZD5363 (pink) administration. (**L,M**) Acute RAD001 treatment had no overt

effect on the mutant burst frequency (t=0.4414, df=4) or PPS (t=0.7692, df=4). Data is

represented as % bar graphs and mean ± SEM scatter plots; differences were

considered significant at p<0.05; ns, not significant. Scalebars: 1s,20mV (B,F,J);

0.05s,20mV (E',I'). See also Figure 5 – figure supplements 1-2.

Figure Supplements

Figure 1 – figure supplement 1: ECoG-LFP traces show different types of epileptiform activity in *Nestin-cre;Pik3ca^E545K* brains

(**A,B**) Examples of different types of isolated spiking events in the neocortex and hippocampus taken from 2 different mutant mice showed a varied range of spike nature – in frequency, source of origin and amplitude. (**C**) Example of train of spiking events in the hippocampus and left cortex. (**D**) Example of low amplitude, high frequency brushing events. g, generalized; r, regional. Scalebars: 0.5s,50mV (A,B); 2s,50mV (C,D).

Figure 2 – figure supplement 1: Enhanced neuronal network excitability in mutant CA1 across an extracellular potassium gradient

(**A**) Bar graphs depicted normalized baseline shift of integrated field potential traces in the mutant with increasing [K$^+$] (refer to Figure 2). The mutant baseline was significantly higher than that of control especially in the 9-11mM range (t=3.343, degrees of freedom (df)=9). (**B**) Relative peak amplitude of mutant spiking was significantly higher than controls at higher potassium concentrations (F=266.1; df=564). Data is represented as mean ± SEM bar graphs; differences were considered significant at p<0.05; ns, not significant.

Figure 3 – figure supplement 1: Spontaneous burst-firing features of control and mutant pyramidal neurons

(**A**) Representative traces for subtypes of burst firing, namely burst cluster, paroxysmal depolarization shift (PDS) and non-PDS plateau bursts. (**B**) Proportion of burst

subcategories were not overtly different in control and mutant CA1 and CA3 regions (F=

$7.355e^{-12}$; df=6); burst clusters were only seen in mutant cells. (**C**) Representative trace

demonstrates how burst duration and inter-burst interval were calculated. (**D,E**) In both

CA1 and CA3, average (avg.) burst duration (CA1: t=0.9644, df=6.793; CA3: t=0.3508,

df=6.229) and inter-burst interval (CA1: t=0.1263, df=22.31; CA3: t=1.485, df=10.73)

were not significantly different between control and mutant neurons. Data is represented

as mean ± SEM scatter plots; differences were considered significant at p<0.05; ns, not

significant; PPS, plateau potential shift. Scalebars: 0.2s,20mV (a), 1s,20mV (c).

Figure 3 – figure supplement 2: Resting membrane potentials and threshold

currents for hippocampal neuronal firing remain unchanged by Pik3ca

overactivation

(**A-C**) Resting membrane potential (RMP) was not significantly different between control

and mutant neurons, both in CA1 (t=0.8331, df=94.99) and CA3 (t=0.3438, df=78.53),

as well as among different firing types (CA1: F=0.9136, df=90; CA3: F=2.829, df=75).

(**D,E**) No significant differences in evoked tonic spike frequencies were observed in the

tested range of input depolarizing current (0-90pA), between control and mutant CA1

and CA3 neurons (CA1: F=13.07, df=320; CA3: F=4.844, df=270). (**F**) No overt

differences in the current inducing the first burst were observed between control and

mutant (CA1: t=1.744, df=20.29; CA3: t=0.8493, df=2.439). Data is represented as

mean ± SEM scatter and line plots; differences were considered significant at p<0.05;

ns, not significant.

Figure 4 – figure supplement 1: Pik3ca-related epileptiform activity is not

dependent on glutamatergic or GABAergic inputs

(**A**) Representative spontaneous mutant trace and traces of the same cell post-

treatment with glutamatergic receptor antagonists (CPP, CNQX) and with GABA

antagonist gabazine showed no significant change in firing pattern. (**B-D**) Large

proportion of mutant cells remained unaffected after being blocked from external

glutamatergic or GABAergic inputs. (**E-L**) No significant differences in normalized

plateau potential shift (PPS; CPP: F=3.391, df=12; CNQX: F=1.432, df=12; Gabazine:

t=0.3904, df=4) or in tonic (CPP: t=1.876, df=7; CNQX: F=0.2766, df=20; Gabazine:

F=0.0001130, df=16) and burst frequencies (CPP: F=0.5140, df=16; CNQX: F=5.218,

df=16) were observed before and after administration of channel blockers. Data is

represented as % bar graphs and mean ± SEM scatter plots; differences were

considered significant at p<0.05; ns, not significant. Scalebar: 1s,20mV (A).

Figure 5 – figure supplement 1: Acute treatment with BKM120 and AZD5363 has

differential effect on *Nestin-cre;Pik3ca^{E545K}* neuronal physiological properties

(**A-D**) Acute BKM120 treatment did not significantly affect RMP (F=0.9268, df=42), tonic

rheobase current (t=0.000, df=8), average burst duration (CA1: t=1.264, df=8; CA3:

t=0.6150, df=11) or inter-burst interval (CA1: t=1.889, df=8; CA3: t=1.160, df=10) in

mutant hippocampal neurons. (**E-H**) Acute AKT inhibition by AZD5363 had no effect on

the mutant RMP (F=0.3316, df=20) or rheobase current (t=0.3859, df=7), but

significantly attenuated the average burst duration (t=4.052, df=5) and inter-burst

interval (t=4.597, df=6), largely by suppressing bursts or silencing the cells. (**I,J**) Plots

marked percentage of mutant CA3 "bursting" cells undergoing changes as a function of

time, in response to acute administration of BKM120 (I) and AZD5363 (J). (**K**) Mutant

plateau frequency significantly dropped as a function of time, where 0min marked the

time of administration of BKM120/AZD5363 onto the brain slice. Data is represented as

mean ± SEM scatter plots; differences were considered significant at $p<0.05$; ns, not

significant.

Figure 5 – figure supplement 2: Acute treatment with RAD001 has no overt effect

on mutant neuronal physiological properties

(**A-D**) Acute BKM120 treatment did not significantly affect the RMP ($t=1.327$, $df=4$),

average burst duration ($t=1.073$, $df=4$), inter-burst interval ($t=0.4023$, $df=4$) or rheobase

current ($t=1.000$, $df=4$) in mutant CA1 neurons. (**E**) Representative Western blots bands

of pAKT_S473, pan-AKT, pS6-S235/236 and RPS6 (total S6) are demonstrated across

different treatment (DMSO, BKM120, AZD5363, RAD001). (**F,G**) Quantifications of

normalized ratios of phosphoproteins over total proteins showed significant reduction of

pAKT_S473 in response to acute BKM120 and RAD001 treatment but not post-

AZD5363 ($F=22.98$, $df=32$); pS6_S235/236 in mutant had a decreasing trend in

response to all pathway drugs ($F=4.485$, $df=32$). Data is represented as mean ± SEM

scatter plots and bar graphs; differences were considered significant at $p<0.05$; ns, not

significant.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Nestin-cre;Pik3ca^E545K

Figure 1 - figure supplement 1

Figure 2 - figure supplement 1

Figure 3 – figure supplement 1

Figure 3 - figure supplement 2

Figure 4 - figure supplement 1

Figure 5 - figure supplement 1

Figure 5 - figure supplement 2