A modeling framework for determining modulation of neural-level tuning from non-invasive human fMRI data

Patrick Sadil*, Rosemary A. Cowell, David E. Huber

University of Massachusetts, Amherst

Corresponding Author: Patrick Sadil

Email: psadil@gmail.com

1   **Abstract**

Many neuroscience theories assume that tuning modulation of individual neurons underlies
changes in human cognition. However, non-invasive fMRI lacks sufficient resolution to visualize
this modulation. To address this limitation, we developed an analysis framework called Inferring
Neural Tuning Modulation (INTM) for "peering inside" voxels. Precise specification of neural
tuning from the BOLD signal is not possible. Instead, INTM compares theoretical alternatives
for the form of neural tuning modulation that might underlie changes in BOLD across
experimental conditions. The most likely form is identified via formal model comparison, with
assumed parametric Normal tuning functions, followed by a non-parametric check of
conclusions. We validated the framework by successfully identifying a well-established form of
modulation: visual contrast-induced multiplicative gain for orientation tuned neurons. INTM can
be applied to any experimental paradigm testing several points along a continuous feature
dimension (e.g., direction of motion, isoluminant hue) across two conditions (e.g., with/without
attention, before/after learning).

15   **Introduction**

16   Neurons in mammalian sensory cortex often exhibit tuning functions, responding
selectively to a narrow range of values of a stimulus feature such as line orientation[1]. Moreover,
experimental manipulations of perceptual or cognitive state can modulate these tuning functions,
providing crucial insights into the neural mechanisms of perception and cognition. For example,
spatial attention achieves performance gains by multiplicative rescaling of orientation tuning
functions in macaque V4[2], rather than through a bias shift (additive shift) or increased selectivity
(sharpening). As another example, Figure 1A shows a neuron in mouse primary visual cortex [3],
tuned to motion direction, that appears to undergo multiplicative rescaling as spatial frequency is
manipulated (Figure 1A). However, single-cell electrophysiology is rarely feasible in humans,
and non-invasive techniques such as functional magnetic resonance imaging (fMRI) reflect the
activity of many cells: a patch of striate cortex the size of a typical fMRI voxel (2 mm^3) reflects
the activity of approximately 300,000 – 500,000 neurons[4]. Nonetheless, fMRI studies reveal
feature-selective tuning in voxels (e.g., in which a voxel's BOLD response varies systematically
as a function of a well-defined feature such as stimulus orientation)[5,6], which is assumed to arise
from a non-uniform distribution of tuning preferences across the neurons contributing to a
voxel[7].

Figure 1 A) Modulation of neural tuning. In this neuron (mouse primary visual cortex)[3], direction tuning interacts multiplicatively with spatial frequency (cpd: cycles per degree; $\Delta F/_F$: average change in fluorescence). **B)** Convolution ($*$) of neural tuning functions with a weight distribution leads to voxel tuning functions. The modeling procedure of INTM assumes that the observed voxel tuning functions reflect the shape of the neural tuning functions as applied across the distribution of preferred orientations (the weight distribution) within each voxel. Compare with Equation 3. **C)** An additive shift in the neural tuning function causes an additive shift in the voxel tuning function. **D)** Multiplicative gain in the neural tuning function causes multiplicative gain in the voxel tuning function. Compare C and D with Equation 4.

Like neural tuning functions, voxel tuning functions are modulated by manipulations of
perceptual state[8–11], and so it is tempting to infer the form of *neural-level* tuning modulation
directly from observed changes in voxel tuning. But, as many have acknowledged, directly
inferring neural behavior from voxel behavior might be misleading[10,12–14]. The relationship
between voxel and neural tuning functions presents a many-to-one inverse problem: the same
voxel tuning function can arise from many different combinations of the shape of the underlying
neural tuning functions and the distribution of neurons across different preferred stimulus values
(e.g., the number of neurons preferring vertical, horizontal, or oblique orientations). Thus, fMRI
tuning does not uniquely specify tuning at the neural level[14].
Rather than *quantitatively* solving this inverse problem, which is probably not possible,
we developed a framework that allows drawing *qualitative* conclusions about changes in neural
level tuning from the Blood Oxygen Level Dependent (BOLD) signal. We term the framework
Inferring Neural Tuning Modulation (INTM). Building on encoding models[15–17], INTM assumes
that voxel tuning emerges from the combined activity of feature-tuned neurons. Unlike many
existing encoding models, which assume that voxel activity is driven by a discrete number of
neural "subpopulations", INTM conceptualizes the underlying neural preferences as existing
along a continuum (Figure 1B). This continuous distribution resembles population receptive field
mapping, but whereas population receptive field mapping is generally concerned with

uncovering the quantitative parameters of tuning (e.g., the size of receptive neural fields, or the
population's preferred location in the visual field)[18], INTM is designed to uncover the
modulation of tuning. To do this, INTM leverages the observation that some forms of
modulation produce patterns of BOLD data that are distinct from other forms (Figure 1C-D), and
so the BOLD data provide information about which form of tuning modulation is most likely.
This framework encompasses many kinds of analyses, but, in short, INTM prescribes building
multiple encoding models of voxel data that are parameterized by continuous distributions at the
neural level, followed by comparison between the models in light of the observed BOLD data to
uncover the most likely form of tuning modulation.
We validated INTM by applying it to a new dataset for which the "ground truth"
modulation of tuning was known from single-cell recordings: changes in visual contrast induce
multiplicative scaling in orientation-tuned neurons[19–21]. This simple test-case demonstrates the
advantages of INTM, considering that existing methods, if interpreted as indicating the
underlying neural tuning functions, can easily reach the wrong conclusion about the form of
neural tuning modulation when applied to changes in visual contrast [22,c.f., 14]. We used INTM to
design two complementary analyses that differ in the assumptions required for the statistical
models underlying the analyses. The first analysis uses Bayesian estimation and quantitative
model comparison by making parametric assumptions for the shape of neural tuning (a circular
Normal distribution). The second makes no assumptions about the shape of neural tuning (i.e., a
non-parametric analysis), providing a qualitative check of whether similar conclusions are
reached. Critically, both techniques are needed, with the parametric model not only providing
statistical model comparison, but also indicating the expected qualitative data patterns for the
non-parametric check. As applied to BOLD data collected from the primary visual cortex of
human participants viewing oriented gratings at high versus low contrast, the techniques
correctly implicated multiplicative scaling, despite working with only voxel-level data.

Parametric Analysis: Linking Neural Tuning to Voxel Tuning
We first use INTM to build a parametric model of voxel tuning for inferring how changes
in stimulus contrast modulate neural tuning. The resulting model provides a concrete
demonstration of the ways in which different kinds of neural tuning modulation should produce
different behaviors at the level of the voxel.
The parametric modeling approach involves assuming specific forms of the neural tuning
functions, and further specifying the contribution of these tuning functions to a voxel's activity
with a continuous distribution. For application to orientation preferences, which lie on a circular
measurement scale, the contributions are assumed to follow a circular Normal distribution
(Figure 2). The circular Normal distribution resembles a typical Normal distribution but is
periodic, thereby accommodating the circular nature of orientations. The distribution's density
function has a period of 2π, but orientations have a period of π, so the values of the orientations
used in an experimental design are doubled when applying the model. For an orientation, r, the
density function is
89

90    $$f(r|\phi, \kappa_1) = \frac{\exp(\kappa_1 \cos(r - \phi))}{2\pi I_0(\kappa_1)} \qquad (1)$$

92   where κ_1 is a concentration parameter, ϕ is the orientation of the peak value of the
93   function, and $I_0(\cdot)$ is the modified Bessel function of the first kind, of order 0. The numerator of

Preferred Orientation

0   90   180

Figure 2 Orientation preferences of simulated voxels can be approximated with unimodal distributions. **Left:** A Kohonen network was used to simulate orientation preference maps. Each panel represents a separate voxel. **Right:** Weight distribution for each voxel. The histograms represent the tallies of each preferred orientation in the left panels. The black line is the best fitting (maximum likelihood estimation) circular Normal density function (Equation 1). The horizontal gray (shaded) ribbons give the 99% confidence interval for a flat distribution (i.e., the range of heights that we would expect when a simulated voxel contains an equal number of neurons tuned to all orientations). Given the amount of cortex sampled by each stimulated voxel, most weight distributions tend to be flat, but the fitted von Mises density function can capture individual modes.

Equation 1 produces a periodic bell shape peaking at ϕ, with the concentration parameter
determining its "sharpness".
In the parametric model, there are two uses of the circular Normal distribution. First,
Equation 1 describes the relative contribution to a voxel's activity of different subpopulations of
neurons that prefer different orientations, and so we refer to it as a "weight distribution". With a
flat weight distribution (low κ_1), the neurons within a voxel are in equal proportion (resulting in
a complete absence of voxel tuning even if the contributing neurons are well-tuned). With a
peaked weight distribution (high κ_1), most of a voxel's activity is driven by neurons tuned to
orientations around ϕ (resulting in sharp voxel tuning if the contributing neurons are also well-
tuned).
A circular Normal density function fits single cell orientation tuning functions
adequately[23] and so, in INTM, a second circular Normal distribution function is used to specify
the individual neural tuning functions (Equation 2). Although this equation relies on the
distribution function, the interpretation is not a probability density but instead a description of
neurons' responses to orientations. In this equation, the concentration parameter indicates the
sharpness of a single cell's tuning. To account for a neuron's baseline activity and
responsiveness, we included additional parameters. The baseline activity parameter reflects the
activity of a neuron in the absence of stimulation and is implemented as an additive offset, α.
The parameter for responsiveness models how well a neuron responds to stimulation and is
implemented as a gain parameter, γ. With the addition of these baseline activity and

responsiveness parameters, Equation 2 specifies the neural tuning function, $NTF(\cdot)$ for
orientation of a presented stimulus relative to that neuron's preferred orientation.
116
117 $$NTF(r|\kappa_2, \gamma, \alpha) = \alpha + \gamma f(r|0, \kappa_2) \qquad (2)$$
118
Within each voxel (but not across voxels), all neural tuning functions are assumed to take
on the same circular Normal parameter values. The voxel tuning function is obtained by
convolving Equations 1 and 2 (Figure 1B). This results I   n Equation 3, the voxel tuning
function, $VTF(\cdot)$, in a baseline condition (e.g., low-contrast stimuli). It is critical to note that the
choice of baseline condition is arbitrary (e.g., the high-contrast condition could be labeled as the
baseline) considering that the model is applied to both conditions conjointly. The only difference
in terms of which condition is labeled as baseline is whether the modulation parameters are
increases or decreases.
127
128 $$VTF_{baseline}(r|\kappa_v, \gamma_v, \alpha_v, \phi_v) = \alpha_v + \gamma_v f(r|\phi_v, \kappa_v) \qquad (3)$$
129
The subscript v indicates parameters that are estimated separately for each voxel. The
concentration parameter, κ_v, is equal to neither κ_1 nor κ_2, but is a function of both[24]. Thus, the
model does not distinguish between voxels containing a concentrated distribution of orientation-
tuned neurons each of which has diffuse tuning (high κ_1 and low κ_2), and voxels containing a
diffuse distribution of orientation-tuned neurons each with concentrated tuning (low κ_1 and high
κ_2). The key insight behind INTM is the realization that there is no need to differentiate between
such possibilities, if the measurement goal is identification of the qualitative form of neural
tuning modulation rather than precise specification of neural tuning functions.
Within the INTM framework, each form of tuning modulation is instantiated by
incorporating additional parameters that modulate one or more parameters in the baseline voxel
tuning function (Equation 3, Figure 1C-D). In this validation test case as applied to visual
contrast manipulations, we considered two forms of tuning modulation that are capable of
producing an average increase in neural activity: multiplicative gain and additive shift. Other
forms of neural tuning modulation, such as bandwidth changes and tuning preference shifts,
could be implemented within the INTM framework, but these forms were not included in the
candidate set of models for this test case because they do not change average neural activity (i.e.,
they are not viable models for the neural modulation that underlies changes in visual contrast).
In this test case, multiplicative gain is known to be "ground truth"—that is, changes in
visual contrast induce multiplicative gain in single-neuron tuning functions[19–21]. However, in
seeming contradiction to multiplicative modulation, an examination of raw voxel tuning
functions suggests an additive increase with increases in visual contrast (Figure 4A,B).
Nevertheless, averages can be misleading, and a formal model comparison between additive shift
and multiplicative gain is required that considers each voxel separately, particularly when
considering that many of the voxels contributing to the average exhibit poor tuning. For
completely untuned voxels, additive shift and multiplicative gain are indistinguishable and so the
average will necessarily look additive. If INTM works, it should leverage differences between
voxels (i.e., capitalize on the small proportion of well-tuned voxels), to reach the conclusion that
multiplicative gain is the more likely form of tuning modulation despite what the average BOLD
signal appears to show.

To model a change in multiplicative gain, we included a voxel-specific gain multiplier
parameter, g_v (Equation 4a). To model an additive shift change, we included another, voxel-
specific additive shift parameter, a_v (Equation 4b). In summary, Equation 3 captures the
situation for low contrast in both models, whereas either Equation 4a or 4b captures the situation
for high contrast.
164
165   *Multiplicative*: $VTF_{modulated}(r|\kappa_v, \gamma_v, \alpha_v, \phi_v, g_v) = \alpha_v + g_v\gamma_v f(r|\phi_v, \kappa_v)$ *(4a)*
166   *Additive*: $VTF_{modulated}(r|\kappa_v, \gamma_v, \alpha_v, \phi_v, a_v) = a_v + \alpha_v + \gamma_v f(r|\phi_v, \kappa_v)$ *(4b)*
167
For other applications, shifts in orientation preferences could be modeled by allowing ϕ_v
to change across conditions, and a change in concentration could be modeled by allowing κ_v to
change across conditions. In the current dataset, these forms were not pursued because neither
allows contrast to alter the average activity of a voxel across orientations – an effect that was
clearly present in the empirical data (Figure 4A) – and so these forms could be rejected without
explicit modeling.
Equations 3, 4a, and 4b model voxel activity at each tested orientation, at each level of
contrast. However, there is variability in voxel activation across runs, and this variability varies
across voxels (some voxels are noisier than others, Supplementary Figure 2). These sources of
variability were included in the model, and all voxel-specific parameters were estimated
hierarchically (Supplementary Methods). Hierarchical estimation of voxel tuning functions
combines information across voxels, allowing model comparison to automatically down-weight
the noisier voxels and magnify the well-tuned voxels[25].
The parametric model assumes that all neurons within a voxel share a common shape for
their tuning function (homogeneity of tuning shape), which is unlikely to hold strictly true.
However, some degree of tuning shape heterogeneity should not invalidate inferences drawn
about the form of tuning modulation, provided that tuning function shape does not differ
systematically as a function of the orientation preference of the neurons. That is, there could
exist a range of tuning function shapes, but if that heterogeneity occurs to the same extent for
neurons centered on all preferred orientations, then it should not lead to scenarios where one
form of modulation mimics another at the voxel level. Nevertheless, to assess whether such
heterogeneity posed a problem when adjudicating between the forms of tuning modulation, we
also developed a non-parametric check that does not make this assumption. If the non-parametric
check supports the same qualitative conclusion as the parametric model, this suggests that the
homogeneity of tuning shape assumption was adequate.

Model Recovery
INTM judges the relative plausibility of each form of neural tuning modulation (e.g.,
Equation 4a versus 4b) with model comparison based on empirical data. But under what
circumstances should model comparison be trusted? To assess the extent to which inherent
properties of the models (e.g., their flexibility) allow reliable model comparison, we assessed
model recovery via two kinds of simulation: data-informed and data-uninformed[26]. Model
recovery entails producing simulated data from each model, asking whether the model that
produced the simulated data is better able to capture its own data than competitor models. If a
model is overly flexible, then it might provide a better explanation of data generated by other
models than do the generating models themselves.
Data-uninformed model recovery asks whether the model comparison technique is well-
calibrated to the relative flexibility of each model in general. First, synthetic data were generated

with each model using "weakly informative" priors (Supplementary Methods). These priors were
not designed to reflect precise experimental knowledge but instead simply provide a rough scale
for each parameter individually. Next, the models were applied to these synthetic data, keeping
in mind the "ground truth", data-generating model. The model comparison technique has built-in
adjustments for flexibility by approximating the ability of each model to predict held-out data
(i.e., cross-validation). However, this approximation might be inadequate, providing too much or
too little penalty for model flexibility. Data-uninformed recovery asks whether this model
comparison is operating as it should in general.
213   There is no guarantee that a specific real dataset will be diagnostic regarding a particular
model comparison. For instance, if none of the voxels is sufficiently well-tuned, then the two
forms of tuning modulation will make the same prediction. Similarly, if the manipulation of
visual contrast is too weak, it would not be possible to reach a reliable conclusion. Whether the
real dataset is sufficient for differentiating between the candidate models is assessed using data-
informed model recovery by repeating the recovery process using what has been learned about
the parameters of the models from the empirical data.
220   Whereas data-uninformed model recovery assesses the fairness of model comparison
across a wide range of individual parameters that are plausible given the prior distribution, data-
informed recovery generates synthetic data by sampling from the posterior distribution obtained
by estimating the models with empirical data. Generating synthetic data that are consistent with
real data constrains the parameters (e.g., tuning width and modulation) to a much narrower and
more realistic range of values. Indeed, the behavior from the posterior parameter values closely
matches the weak tuning effects seen in the real data at the level of separate voxels (e.g.,
Supplementary Figure 5). Because the data-uninformed model recovery has priors that are only
weakly informative, it can allow extreme parameter values for estimated beta-weights, and hence
implausibly sharp tuning functions; this increases the chance that each generating model will
produce a unique signature pattern in the data, making model recovery likely to succeed. By
contrast, the more constrained data-informed model recovery presents more of a challenge to
model recovery, providing greater reassurance in the case that it succeeds.

Non-parametric Check: An Orthogonal Regression Slope Analysis

234   The parametric model assumed normal functions for the NTFs and weight distributions
and additionally assumed homogeneity of NTF shape within a voxel. These assumptions will
often be violated, but such violations might not pose a problem for model comparison,
particularly if the violations of these assumptions are uniformly applied across the stimulus
dimension of interest. Nevertheless, as a complement to parametric model comparison, we
developed a more qualitative non-parametric check of the empirical data. If this check supports
the same conclusion as the parametric model, this suggests that any violations of parametric
assumptions were insufficient to alter the theoretical conclusions.
242   The parametric model is crucial to the development of the non-parametric check and the
specific parametric models under consideration should first be used to generate predictions for
the check. The consequences of parametric assumptions are then considered in light of these
predictions. In the case of a comparison between multiplicative gain versus additive shift, these
two forms of tuning modulation can be distinguished using voxel-wise regression; if the
modulation is multiplicative, a within-voxel plot comparing high- versus low-contrast should
have a slope greater than 1, whereas if the tuning modulation is additive, this plot should reveal a
slope equal to 1. Furthermore, these predictions do not rely on parametric assumptions. For
instance, if a voxel is multimodal, preferring more than one orientation, this would serve only to

Figure 3 Plotting a voxel's response to high versus low contrast orientation uncovers the form of tuning modulation. **Left:** Simulated voxel tuning functions in which higher levels of contrast induce either an additive (top) or multiplicative (bottom) tuning modulation. The eight vertical lines are eight hypothetical orientations at which these voxel tuning functions might be probed, which would produce eight responses per level of contrast, High vs. Low. Note that, whereas the parametric model assumes that the weight distribution and neural tuning function follow specific shapes (e.g., unimodal distributions), the slope test does not (e.g., the voxel tuning function can be bimodal, as displayed). **Right:** The two kinds of neural tuning modulation reveal different signatures when the responses to high contrast stimuli are plotted against the responses to low-contrast stimuli. The diagonal line corresponds to no effect of contrast. A line drawn through the points produced by the additive model necessarily has a slope equal to 1 (top); under this form of modulation, the effect of contrast does not depend on the orientation. A line drawn through the points produced by the multiplicative model necessarily has a slope greater than 1 (bottom); under this form of modulation, the effect of contrast is largest at those orientations that are closest to the voxel's preferred orientation.

rearrange the order of the orientations in the regression plot, but the qualitative distinction
between a slope of 1 versus a slope greater than 1 would still map onto additive shift versus
multiplicative gain. The generality of the slope test across all shapes of tuning function and all
shapes of weight distribution is mathematically proved in Supplementary Methods Equations 1-
4.
The key difference between these two forms of tuning modulations it that the
multiplicative model allows the effect of contrast to vary by orientation whereas the additive
model does not. Additive modulation corresponds to an increase in neural activity at all
orientations. Thus, regardless of the weight distribution and regardless of the shape of the neural
tuning functions, additive tuning modulation causes the low-contrast tuning function to shift
upwards uniformly across orientations (Figure 3, top left). Hence, a scatterplot of the voxel's
response to high-contrast stimuli against its response to low-contrast stimuli has a slope of 1
(Figure 3, top right). In contrast, multiplicative tuning modulation corresponds to a greater
increase in neural activity at the most preferred orientations (Figure 3, bottom left), and a
scatterplot of a voxel's response to high versus low-contrast stimuli has a slope larger than 1
(Figure 3, bottom right). Therefore, the models can be differentiated by plotting high-contrast
activity against low-contrast activity and calculating the slope of the best fitting line (Figure 3,
right). A slope of one implies additive shift, but a slope greater than one implies multiplicative

gain (see also Supplementary Methods). The only assumption made in this analysis is that the
magnitude of the additive shift or multiplicative gain is the same for all neurons contributing to a
particular voxel or, more specifically, that if the magnitude of modulation does vary across
neurons, it does not vary systematically with neurons' preferred orientation (see Supplementary
Information).
274   Analyzing the slope is not necessarily straightforward. First, the slope between high- and
low-contrast cannot be determined using standard linear regression (i.e., estimating the line that
minimizes the ordinary least squares error—the squared vertical distance between the line and
the data). Standard linear regression estimates a relation between two sets of observations, such
as the activity at low and high contrast. But when *both* of those observations are corrupted by
noise, the estimated relationship is "diluted" and biased to zero[27]. To mitigate this dilution, the
slope was estimated using *orthogonal* regression[27], by finding the line that minimizes the squared
Euclidean distances from the data (Methods, Supplementary Methods). Second, the slope
analysis cannot be done across voxels because there may be covariance between a voxel's low
contrast (baseline) activation and the effect of contrast on the voxel's activity (e.g., voxels with a
larger response in the low-contrast condition may receive a larger contrast effect, and this
tendency would bias an across-voxel analysis toward a slope greater than 1). To mitigate this
confound, slopes were determined separately for every voxel, followed by an analysis of the
distribution of slopes. Finally, even a within-voxel slope analysis produces results that are
difficult to interpret because most voxels are only weakly tuned, responding to all orientations
almost equivalently. For those voxels, the data for the slope analysis will be an uncorrelated
cloud of points, and an orthogonal regression for a cloud of points can take on nearly any slope
value, including positive or negative infinity for the case of vertical slopes. Therefore, we report,
not the slope of the line, but instead the angle of the slope.
293   In using INTM, we recommend first developing and comparing parametric models of
294   how stimuli give rise to voxel activity during manipulations of cognitive or perceptual state, and
295   then, in addition, to consider ways of relaxing those parametric assumptions to provide a non-
296   parametric check of the model comparison results, as with the slope test for comparisons of
297   multiplicative versus additive changes.

298   **Results**

299   **Binning Voxels to Plot Average Voxel Tuning**
300   Voxel tuning functions were first analyzed with a standard approach of "averaging"
across circularly aligned voxels (Figure 4)[10], which is an appropriate technique if all voxels have
similarly shaped tuning functions. This "binning" alignment technique appears to reveal an
additive shift with contrast, in contrast to what is known from electrophysiology. This apparent
additive shift is caused by the inclusion of many voxels with poor orientation tuning. When the
BOLD response is aggregated across a many such "undiagnostic" voxels, the true form of tuning
modulation is obscured (i.e., it appears additive rather than multiplicative). One approach to
mitigating this variability would be to adopt experiment-specific thresholds, excluding some
subset of "relatively" untuned or unresponsive voxels. However, it would be difficult to justify
such thresholding for each experiment, and more importantly excluding voxels throws away
information (e.g., about the average noise in voxels' responses). The two techniques of INTM

Figure 4 Average voxel tuning functions visually suggest an additive shift even though the multiplicative model is better in its posterior predictions of individual voxel data. **A)** Average voxel tuning at low and high contrast using binning to average (see Methods). Error bars give 95% confidence interval for the mean of the 1,010 V1 voxels across seven participants. **B)** Examples of individual voxel BOLD data in comparison to posterior predictions of the multiplicative gain model. The panels show the 16 voxels with the largest average difference in activity across levels of contrast. Vertical error bars span 95% confidence intervals (within-subjects), and the width of the lines indicate the 95% highest density interval of the posterior predictive distribution for the multiplicative model (additive model looks similar). **C)** Overall comparison between each model ($\hat{\beta}$) and observed data (β). Each point corresponds to a single voxel's average activity at a given orientation at a given level of contrast. Panels show the average of the posterior for predictions from either the additive or multiplicative model.

avoid thresholding, even while using the well-tuned "diagnostic" voxels to identify the most
likely form of tuning modulation.

Data-Uninformed Model Recovery
To evaluate the ability of the INTM parametric model to recover both kinds of
modulation, we conducted model recovery by simulating datasets from the prior distribution for
each candidate model, applying both models to both sets of generated data, and comparing the

results using an approximation to cross-validation[26,28] (see Methods). The models did not exhibit
mimicry: model comparison picked the true data-generating model in every simulated dataset,
providing strong evidence that the technique recovers the appropriate class of modulation.

Application of the Parametric Model to the Observed Data

We next asked whether the parametric INTM analysis recovers the correct form of tuning
modulation when applied to the observed fMRI data. Specifically, it is known from
electrophysiology that the tuning functions of orientation-sensitive neurons undergo
multiplicative tuning modulation with an increase in stimulus contrast[19–21]. Given this
knowledge, model comparison on the collected BOLD data ought to favor the multiplicative
model. Visual comparison of the posterior predictive distribution from both models revealed that
both models largely captured individual voxel tuning functions adequately (Figure 4B,C). The
models were then compared to each other with cross-validation[28]. The multiplicative model
provided the best account of the data: a difference of 14.5 standard errors, in units of expected
log pointwise-predictive density[28]. a significance threshold (e.g., $\alpha = 0.05$ or 1.65 standard
errors) would suggest that this difference in predictive ability is poorly accounted for by a null
model in which the two forms of tuning modulation are equally predictive.

Data-Informed Model Recovery

Exploiting what was learned about the plausible parameter values by observing the data,
we further probed the validity of model comparison through data-informed model recovery[26,28].
Synthetic datasets were generated from the posterior distributions of the multiplicative and
additive models. Next, the multiplicative and additive models were applied to each of these
simulated datasets. All datasets were best accounted for by the model whose posterior was used
to generate the data even though, as seen in Figure 4, the differences between the models were
very subtle.

Orthogonal Regression

The parametric model correctly indicated that an increase in visual contrast caused an
increase in multiplicative gain, but this result relied on assumptions of circular Normal
distributions for the neural tuning function and distribution of weights for each voxel, as well as
homogeneity of tuning shape for each voxel. However, we determined that the two candidate
forms of tuning modulation should be distinguishable based on what they predict for the slope of
a plot of high-contrast activity against low-contrast activity (Figure 3, Introduction,
Supplementary Methods), without relying on these parametric assumptions and the assumed
homogeneity of tuning shape. As predicted by multiplicative tuning modulation, this non-
parametric orthogonal regression slope analysis revealed slopes larger than 1, corresponding to
slope angles greater than 45 degrees (Figure 5). Figure 5B shows the distribution of within-voxel
slopes, as estimated with orthogonal regression of the high- on the low-contrast activity
(Introduction, Methods). The median slope had an angle greater than 45° (i.e., slope of 1). This
tendency for slopes to be larger than 1, as predicted by the multiplicative model, was further
supported by a hierarchical Bayesian estimation of the slopes (Supplementary Methods).

Figure 5 Non-parametric check of model comparison using orthogonal regression analysis. **A)** Orthogonal regression on data and model predictions. Each line corresponds to a single voxel. Most voxels are weakly responsive at each level of contrast and are only weakly influenced by the experimental manipulation (i.e., lines cluster in the lower left corner, close to the diagonal). Across voxels (i.e., comparing separate lines), both models capture larger visual contrast effects for more responsive voxels: predictions lying further rightward along the x-axis are located further away from the diagonal. However, only the multiplicative model captures the observed interaction between orientation and contrast within voxels. Compare to Figure 3. **B)** Distribution of within-voxel slopes. As allowed by multiplicative but not additive modulation, the distribution of slopes is shifted above 45° (i.e., slope of 1).

Discussion

Many theories of cognition predict specific kinds of neural tuning modulation with
different manipulations [e.g., 29,30]. For instance, attention researchers do not only consider neural
responses when people are in a state of high attention; this state is compared and contrasted to
one of low attention, and theories of attention are informed by this comparison [e.g., 31,32]. However,
these predictions for tuning modulation – the manner in which neural tuning functions change –
are rarely tested in humans because there are no non-invasive techniques for recording from
individual neurons. Tuning functions, and modulation of tuning functions, can be measured non-
invasively using fMRI, but the response of a voxel reflects hundreds of thousands of neurons.
Thus, voxel tuning functions have been interpreted as population-level properties of stimulus
representations in the brain, population-level responses that cannot be used to infer properties of
the underlying, individual neurons[10,12–14].
Rather than attempting a quantitative measurement of neural-level tuning from fMRI
data, we developed a framework, INTM, for identifying the functional form of neural-level
tuning modulation. A key component underlying the INTM framework is a modeled link
between the neural and voxel tuning functions (Figure 1B-D). This link justifies inferences of
neural tuning modulation from voxel tuning modulation; even though the BOLD signal reflects
the aggregated activity of many neurons, different forms of modulation can produce identifiable
signatures in activity of the voxels (i.e., Equation 4a vs 4b, Figure 3), thereby allowing
identification of the most likely form of modulation by applying each model to the BOLD data.
We tested that link and validated INTM with a test case in which the neural-level "ground truth"
modulation was known to be multiplicative gain[19–21]. This test case was particularly challenging

378 because the average voxel results appeared to indicate a clear additive shift rather than
379 multiplicative gain. We used INTM to design two techniques: 1) parametric assumptions of
380 tuning function shapes and hierarchical Bayesian estimation of the models followed by formal
381 model comparison; and 2) a qualitative, non-parametric check using orthogonal regression,
382 without relying on parametric shape assumptions. Both techniques of the INTM framework
383 correctly implicated multiplicative gain of neural tuning by leveraging subtle effects that were
384 more apparent in some voxels than others.
385   Several techniques have previously been developed to peer inside the voxel, but none of
386 them statistically compares alternative theories of tuning modulation using a model that
387 explicitly links neural tuning with voxel tuning. For example, biophysical models specify how
388 the BOLD signal arises from the often nonlinear coupling between neural activity and
389 vasculature[33,34]. But, despite this biological realism, these models have not yet been used to
390 examine changes in neural tuning functions. In contrast to biophysical models, encoding models
391 of fMRI data delve inside the voxel by modeling how components comprising a voxel (e.g., sub-
392 voxel "channels") transform stimuli into voxels' activity[5,15–17], and, promisingly, enable
393 researchers to use the BOLD signal to uncover features of neural tuning[18]. Encoding models can
394 be applied to multiple conditions – for example through simulation[35], or through fitting to data
395 and inverting[36] – thereby estimating how the channels are modulated by an experimental
396 manipulation. Inverting an encoding model uses it as a decoder, estimating channel responses for
397 other stimuli[15,16,37] or estimating how channels are modulated in other conditions[38]. But this use
398 of inverted encoding models requires hard-wired (and arbitrary) assumptions about the shape of
399 the channel responses in baseline conditions[39,40]. These assumptions are even more restrictive
400 than those made by the parametric model in INTM considering that they not only assume a
401 specific class of shape (e.g., a circular Normal), but they require adopting a specific, user-
402 selected parameter value for that shape (e.g., a particular value for the width of the circular
403 Normal). These assumptions mean that the inferred channel modulation cannot be taken to
404 reflect the modulations in underlying neural functions that would be measured with
405 electrophysiology, as users of such models readily acknowledge[14,39]. As a result, inverted
406 encoding models have been used to pursue very different goals than the aim of the present
407 study[14] (i.e., understanding population- rather than neural-level representations). Beyond
408 inverted encoding models, other researchers have compared the qualitative predictions of
409 different forms of tuning modulation[35,41,42], analogous to the logic underlying INTM's non-
410 parametric check. However, these techniques neither fit the alternative encoding models to
411 empirical data, nor account for model flexibility.
412   These previous attempts to peer inside voxels are combined in INTM. Like biophysical
413 models, INTM connects the latent neural activity to the observed BOLD. It does so by assuming
414 a linear coupling that is justified given our experimental design and the goals of our modeling
415 procedure (we discuss potential violations of this assumption in more detail below)[43–46]. But
416 unlike these biophysical models, INTM uses an encoding model (Figure 1B-D, Equations 3, 4a,
417 4b). Like inverted encoding models, the parametric analysis in INTM assumes a specific class of
418 neural tuning shape (circular Normal). But unlike inverted encoding models, it avoids the need to
419 assume specific parameter values for the shape (e.g., tuning width). The non-parametric slope
420 analysis in INTM takes things a step further by avoiding the need to assume a specific class of
421 neural tuning shape. In applying the framework to data, we used a quantitative measure of the
422 predictive abilities of each model (a form of cross-validation), a measure that is sensitive to both
423 the flexibility of each model and the signal-to-noise ratio of the data.

INTM is a general framework for inferring neural tuning modulation, but the two
techniques developed in this report have a crucial feature that ought to be retained in further
applications of the framework: analyses were within-voxel. As previously reported for voxel-
wise orientation tuning[6], most of the voxels in our study were poorly tuned, and they were
additionally only weakly affected by stimulus contrast. Although we restricted the analyses to
only those voxels whose population receptive fields overlapped with the stimulus (see Methods),
most voxels were only weakly responsive (e.g., falling near the origin in Figure 5A) and were
minimally affected by visual contrast (e.g., lying near the diagonal in Figure 5A). These voxel
differences and other sources of voxel variation – including partial voluming[47], proximity of each
voxel to blood vessels[48], the responsiveness of neurons within each voxel, etc. – are present in
any fMRI experiment. Analyses that are insensitive to weak signals risk conflating different
forms of tuning modulation, particularly when the noise differs across conditions
(Supplementary Methods – Non-Parametric Check)[22]. In the current case, because most voxels
exhibited nearly flat tuning, this altered the appearance of the average voxel tuning function
(Figure 4A), incorrectly suggesting that changing visual contrast causes an additive shift. This
was because an additive shift is equivalent to multiplicative gain for a voxel with poor tuning.
Both INTM techniques avoided this problem by assessing the voxels individually, identifying
(and exploiting) the voxels that exhibited the unique combination of being well-tuned and
possessing a strong visual contrast effect. Voxels with this combination of behaviors were more
diagnostic for the comparison between different forms of tuning modulation.
The techniques developed here were tailored to uncover changes in orientation tuning
caused by stimulus contrast, relying on assumptions that will not be appropriate in all studies.
We highlight these assumptions here and use them to clarify the distinction between the
framework more generally versus specific applications of the framework. First, the models
assumed that voxel tuning functions are a linear combination of the activity of the underlying
neural tuning functions. This assumption is reasonable in some experiments[43,44], but the
relationship between neural firing rate and BOLD signal is not perfectly linear[46,49]. Second, the
techniques implicitly assumed, through the general linear model used to estimate voxel activity,
a hemodynamic response function that is shared across all voxels in all participants. Although
canonical, that assumption is erroneous [e.g., 50]. Next, the parametric analysis approximated the
distribution of neurons tuned to orientations within each voxel with a circular Normal
distribution, which is unimodal and periodic. This approximation provided a convenient formula
for deriving the voxel tuning function, and the resulting function matched the data (e.g.,
individual voxel tuning functions in Figure 4B appear unimodal). However, this assumption may
not hold for other types of stimuli, other brain regions, or for other scan parameters (e.g., an
aperiodic stimulus like pitch would require an aperiodic tuning function).
Violations of these assumptions may bias model comparison in some cases but,
fortunately, none of the assumptions is intrinsic to the framework, and there are several ways of
relaxing each of them. For example, more complex weight distributions such as multimodal
functions could be used in the parametric model. Alternatively, we have shown that assumptions
about the weight distributions can be bypassed by a technique like the non-parametric slope
analysis. As an another example, although nonlinear relationships between the neural firing rate
and the BOLD signal have been documented, multiple mathematical accounts have been
proposed to explain these nonlinearities[49,51,52], and these relationships could be incorporated into
INTM.

Despite its generality, INTM has two limitations that must be plainly acknowledged. The
first is that INTM requires that the models under consideration make distinct predictions at the
level of the voxel. For instance, the parametric and non-parametric INTM techniques both
assume that the magnitude of tuning modulation is the same for all neurons that contribute to the
voxel response. These techniques may be robust to relaxing this assumption if modulation
magnitude differences apply uniformly across the weight distribution that maps neural
preferences onto voxel preferences. But if the magnitude of modulation varies systematically, the
models might no longer make distinct predictions at the level of the voxel. For example, if the
magnitude of the additive shift is larger for neurons that prefer the same orientation as the voxel,
this unlikely circumstance could produce a change in the voxel tuning function that is
indistinguishable from a constant magnitude of multiplicative gain. The inability to discriminate
these two scenarios exemplifies the fundamental inverse problem that INTM cannot solve:
INTM cannot determine the parameters of tuning functions for individual neurons (indeed, it is
not designed to). Instead, INTM operates upon distributions of neurons (grouped by, e.g., region-
of-interest, voxel, feature preferences), and so it can compare only models whose distributions of
tuning functions change in ways that are distinct. With the example of an additive shift whose
magnitude happens to covary with the weight distribution, this might occur for some voxels, but
is unlikely to occur with most voxels, demonstrating the need to use all of the voxels in the
model comparison.
The second key limitation is that INTM can differentiate only between models that are
formally included in the model comparison process. If the "true" form of tuning modulation is
not included in the set of possibilities, the results may mislead researchers into conflating the
winning form of modulation with the true form. For instance, in the present study, the technique
contrasted tuning modulation models in which just one form of modulation occurred but did not
consider more complex situations in which multiple forms of tuning modulation could have
occurred. For example, it is known from electrophysiology that with extended adaptation,
orientation tuning functions can both widen and shift their preferred orientation[53]. Since that
more complex model, in which tuning widened and shifted within every trial, was not included
in model comparison, INTM is silent about whether that effect occurred. Note that such
combinations of tuning modulation could be included in the model comparison process, but
when attempting to adjudicate between complex models it would be critical to use model
recovery simulations, specifically data-informed recovery, to determine if the collected data were
sufficiently constraining.
In summary, using a simple test case – modulation of orientation tuning by stimulus
contrast – we presented and validated the INTM method for identifying the form of neural tuning
modulation from a BOLD dataset. The method is applicable to a broad range of domains and
manipulations. The main insight of the framework is that there are reasonable assumptions,
which can be made in many neuroimaging studies, that enable inference about neural tuning
modulation. The main requirement for using the technique is an experimental paradigm that: (1)
produces measurable voxel tuning by testing the BOLD response at different levels of a single
stimulus dimension (e.g., direction of motion, color, pitch), and (2) includes some cognitive
manipulation of interest (e.g., with and without attention, before and after perceptual learning,
complex versus simple stimuli) that modulates the voxel tuning from one condition to another.
Now that INTM has been validated, it can be used to study modulations of neural tuning where
electrophysiology has not yet provided an answer (e.g., studies of tuning modulation underlying
tasks that cannot be taught to animals).

<div align="center">

Methods

</div>

Participants

Seven participants (22 – 31 years old; 3 female, 2 did not report) completed three sessions for monetary compensation ($50 per 2-hour session). All participants had normal or corrected-to-normal vision. One additional participant completed a single session but exhibited substantial motion; their data were excluded. The procedure was approved by the University of Massachusetts Institutional Review Board.

Behavioral Stimulation and Recording

Stimuli were presented to participants with a gamma corrected 32" LCD monitor at 120 Hz refresh rate (Cambridge Research Systems). The experiment was designed using the Psychophysics toolbox (Version 3.0.14)[54] and custom MATLAB code (2018b, MathWorks). Behavioral responses were collected with a button box (Current Design). Eye-tracking data were recorded at a rate of 1000 Hz with the Eyelink 1000 Plus system on a long range mount (SR Research), controlled using the Eyelink Toolbox extension to Psychtoolbox[55]. Due to a technical error, the button presses of one participant were not recorded.

     Orientation scans. For the main experiment, participants completed 18 functional runs across three sessions. During each run, oriented grayscale gratings were presented twice at each of two levels of contrast (eight orientations at 50% or 100% Michelson contrast in all runs for six out of seven participants, eight orientations at 20% or 80% Michelson contrast in 12 runs for one participant, and seven orientations at 20% or 80% Michelson contrast in that participant's remaining 6 runs). Grating parameters replicated those of Rademaker et al.[56]. Gratings (spatial frequency of 2 cycles per degree) were masked with annuli (1.2° inner and 7° outer radii). The annulus edges were smoothed with an isotropic 2D Gaussian kernel (1° kernel, 0.5° standard deviation). Throughout each run, a magenta fixation dot was presented in the center of the screen (0.2°, RGB: 0.7843, 0, 0.8886).

     In each trial, a counterphasing (5 Hz) grating was presented for five seconds. In the middle three seconds of each trial, the spatial frequency of the grating either increased or decreased (1 cycle per degree) for 200 ms. Participants were instructed to indicate via button press the direction of change as soon as they noticed it. Per run, gratings were presented at multiple orientations, twice at each combination of orientation and contrast. In most runs, there were eight orientations, but in one session of one participant (totaling six runs), only seven orientations were presented. Inter-stimulus intervals ranged from 8000 – 12000 ms in steps of 200 ms. A five-second fixation period preceded the first trial, and a fifteen-second fixation period succeeded the final trial. Each run lasted 490 seconds.

     Population Receptive Field (pRF) Mapping Scans. We mapped a circular area of the visual field, of radius 8° centered on a central fixation point. pRF mapping scans followed the protocol of Benson et al.[57]. Briefly, natural images[58] were overlaid on pink noise and viewed through a series of circular apertures (8° radius). Within one run per session, the apertures enabled view of either moving bars (2°) or rotating wedges (1/4 aperture) and rings that expanded and contracted (see stimulus software for details). In bar runs, a bar traversed the central region in cycles. During each cycle, the bar was visible for 28 seconds, followed by a 4-second blank period. The bar moved in one of eight directions (east, north, west, south, northeast, northwest, southwest, or southeast, in that order). A 16-second blank period preceded the first cycle, a 12-second blank period followed the fourth cycle, and there was a 16-second blank period at the end of all cycles (300 seconds in total).

In the second pRF scan, the apertures were either wedges that rotated clockwise or
counterclockwise, or they were rings that expanded or contracted. These runs started with a 16-
second blank period, followed by two, 32-second cycles of a counterclockwise rotating wedge,
two 28-seconds of expanding rings (each followed by a 4-second blank period), two 32-second
clockwise wedge rotations, and two 28-second cycles of contracting rings (followed by 4 and 26
seconds of blank, respectively). The total run time was 300 seconds.
Throughout the pRF scans, the color of a central fixation dot (0.3°) changed between
black, white, and red. Participants were instructed to monitor the color of the fixation dot and
press a button when the dot turned red. To help participants maintain fixation, a circular fixation
grid was presented throughout.

fMRI Data Acquisition

MRI data were collected on a 3T Siemens Skyra scanner with a 64-channel head coil. In
each of the three sessions we collected field-mapping scans, functional scans, and a T1-weighted
anatomical scan (MPRAGE, FOV 256 × 256, 1 mm isotropic, TE 2.13 ms, Flip Angle 9°). The
anatomical scan was used to align field-mapping and functional images parallel to the calcarine
sulcus. Gradient recall echo scans estimated the magnetic field. The pRF and primary functional
data were collected with the same scan parameters (TR 1000 ms, TE 31 ms, flip angle 64°, FOV
94 × 94, 2.2 mm isotropic, interleaved acquisition, no slice gap, Multiband Acceleration Factor
4). To aid alignment of functional and anatomical images, single-band reference images were
collected before each functional run for all but three participants (TR 8000 ms, TE 65.4 ms, flip
angle 90°, FOV 94 × 94, 2.2 mm isotropic, interleaved acquisition, no slice gap).

MRI Preprocessing

582   Preprocessing of images was performed with *fMRIPrep* 1.4.0, which relies on *Nipype*
583   1.2.0 and *Nilearn* 0.5.2 (RRID:SCR 001362).

Anatomical Data

The T1-weighted (T1w) images were corrected for intensity non-uniformity (INU) with
N4BiasFieldCorrection, distributed with ANTs 2.2.0. The T1w images were then skull-stripped
with a *Nipype* implementation of the *antsBrainExtraction.sh* workflow (from ANTs), using
OASIS30ANTs as target template. A T1w-reference map was computed after registration of the
individual T1w images (after INU-correction) using *mri_robust_template* (FreeSurfer 6.0.1;
RRID: SCR 001847). Brain surfaces were reconstructed using *recon-all*, and the brain mask
estimated previously was refined with a custom variation of the method to reconcile ANTs-
derived and FreeSurfer-derived segmentations of the cortical gray-matter of Mindboggle (RRID:
SCR 002438). Brain tissue segmentation of cerebrospinal fluid (CSF), white-matter (WM) and
gray-matter (GM) was performed on the brain-extracted T1w using *fast* (FSL 5.0.9, RRID:SCR
002823).

Functional Data.

597   For each of the functional runs, the following preprocessing was performed. First, a
reference volume and its skull-stripped version were generated using a custom methodology of
fMRIPrep. A deformation field to correct for susceptibility distortions was estimated based on a
field map that was co-registered to the BOLD reference, using a custom workflow of fMRIPrep
derived from D. Greve's *epidewarp.fsl* script
(www.nmr.mgh.harvard.edu/~greve/fbirn/b0/epidewarp.fsl) and further improvements of
Human Connectome Project Pipelines. Based on the estimated susceptibility distortion, an
unwarped BOLD reference was calculated for a more accurate co-registration with the

anatomical reference. The BOLD reference was then co-registered to the T1w reference using
bbregister (FreeSurfer) which implements boundary-based registration. Co-registration was
configured with nine degrees of freedom to account for distortions remaining in the BOLD
reference. Head-motion parameters with respect to the BOLD reference (transformation
matrices, and six corresponding rotation and translation parameters) were estimated before any
spatiotemporal filtering using *mcflirt* (FSL 5.0.9). The BOLD time-series were resampled to
surfaces on the following spaces: fsaverage and fsnative (FreeSufer). The BOLD time-series
were resampled onto their original, native space by applying a single, composite transform to
correct for head-motion and susceptibility distortions. These resampled BOLD time-series will
be referred to as "preprocessed BOLD". A reference volume and its skull-stripped version were
generated using a custom methodology of fMRIPrep. A set of physiological regressors were
extracted to allow for component-based noise correction[59]. Principal components are estimated
after high-pass filtering the pre-processed BOLD time-series (using a discrete cosine filter with
128 s cut-off) for the anatomical CompCor (aCompCor). The time-series entering the CompCor
analyses are derived from a mask at the intersection of subcortical regions with the union of CSF
and WM masks calculated in T1w space, after their projection to the native space of each
functional run (using the inverse BOLD-to-T1w transformation). Gridded (volumetric)
resamplings were performed using *antsApplyTransforms* (ANTs), configured with Lanczos
interpolation to minimize the smoothing effects of other kernels. Non-gridded (surface)
resamplings were performed using *mri_vol2surf* (FreeSurfer).
To estimate voxel-wise responses to each orientation, a general linear model (GLM) was
fit using SPM12 (version 7487, RRID:SCR 007037) to the time-series of each voxel during each
orientation run. Fitting the GLM can be viewed as a preprocessing step to reduce the
dimensionality of the data; the method presented here could be configured to run on the raw
timeseries, but working with beta weights of a GLM rather than the raw timeseries drastically
reduced the computational requirements of the Bayesian estimation. Prior to fitting the GLM,
each voxel's timeseries was converted into a percent signal change, relative to the average signal
within a run (across voxels). Design matrices were convolved with the canonical hemodynamic
response function, parameterized with the SPM12 defaults, and additionally contained both six
motion (three translation and three rotation) and multiple aCompCor regressors. For each run,
the number of components was determined by the broken-stick method.

Statistics and Reproducibility

Population Receptive Field Mapping

To mitigate the effects of stimulus vignetting – whereby orientation information differs
between the edge and central portions of a grating[60,61] – we restricted the analyses to only striate
voxels whose population receptive fields (pRFs) did not overlap with the stimulus edges. We
estimated the pRFs of each voxel with standard methods. First, the preprocessed functional data
for the pRF scans were converted into percent mean signal change within a run (across voxels).
The compressive spatial summation model was fit to each voxel using analyzePRF HCP7TRET
[52,57]. Following Benson et al., the compressive exponent of this model was set to 0.05. The
resulting pRF parameters were combined with an anatomical prior for a Bayesian estimation of
the parameters (neuropythy 0.94)[62]. Only the parameters of voxels for which the pRF explained
more than 10% of the variance of the run were used as empirical parameters for Bayesian
estimation; the remaining voxels' posterior pRF parameters were determined entirely by the
prior.

The resulting pRF parameters determined whether a voxel would be retained for
analyses. The pRF resembles an isotropic, bivariate Gaussian. The three pRF sessions were
analyzed separately, resulting in three sets of three pRF parameters per voxel. Within a set of
parameters, two indicate the center location of the pRF, and the third determines its size – the
standard deviation of the Gaussian. A voxel was retained only if a circle centered on its pRF with
radius equal to two standard deviations was entirely contained by the grating stimulus in each of
the three sessions.

Estimated Average Voxel Tuning Function
The voxel tuning function (Figure 4) was estimated according to a "binning" method[10].
First, each voxel's activity was z-scored, separately for each voxel. Then, one run was held out
and the voxel's preferred orientation was calculated with the remaining runs. The preferred
orientation was estimated as the orientation that produced the largest activity, averaged across
runs and levels of contrast. Next, the activity in the held-out run was plotted against stimulus
orientation translated into a function of distance from the voxel's preferred orientation. This
procedure was repeated once for each available run, holding out a different run on each
repetition. Averages and repeated-measures confidence intervals, across voxels and ignoring
participants, were then calculated[63], treating contrast and distance from preferred orientation as
within-voxel factors.
In Figure 4, one session of one participant was excluded; the figure shows only sessions
in which the participant was shown eight orientations.

Bayesian Estimation of Parametric Circular Normal Models
Each of the voxel-specific parameters (baseline tuning function, tuning modulation, and
variability in the output of the tuning function) were estimated with a hierarchical Bayesian
model. All models were estimated with R (R Core Team) and its interface (RStan, 2.18.2) to the
Stan language[64]. The Stan language provides a robust variation of Hamiltonian Monte Carlo[65,66],
which efficiently and accurately approximates Bayesian posterior distributions. Two diagnostics
were used to assess the accuracy of the estimation. First, we calculated the split-$\hat{R}$ for each
parameter in the model[67]. This value is analogous to an F-score in an analysis of variance, and
values close to 1 give no evidence that the different Markov chains are sampling from different
distributions. The split-$\hat{R}$ was below 1.1 for all parameters[25]. Second, chains were monitored for
divergences, whose presence would indicate that the samples, even if they are from a common
distribution, likely misrepresent the true posterior distribution. There were no divergences when
applying the multiplicative and additive models to real data.
In all applications of models to data, four chains were initialized with random values for the
parameters (RStan defaults), and the sampling algorithm was given 1000 draws to adapt. After
adaptation, each chain was used to draw 2000 samples from the posterior distribution. All model
comparison was done with Pareto-smooth importance sampling, leave-one-out cross-validation
plus, a technique for approximating cross-validation[28]. We refer readers to the original report for
details of this technique.

Model Recovery
For data-uninformed model recovery, 100 datasets were simulated from the priors of both
models. The datasets contained one participant contributing 100 voxels measured in 18 runs at 8
levels of orientation. Both the additive and multiplicative models were applied to each simulated
dataset, resulting in 200 simulated datasets and 400 model applications.

Data informed model recovery was implemented analogously to the data-uninformed
model recovery. One hundred datasets were generated from the posterior of the multiplicative
and additive models. The datasets contained six participants, each contributing 200 voxels
measured in 18 runs at 8 levels of orientation. The multiplicative and additive models were
applied to these simulated datasets and model comparison was conducted as above.

Orthogonal Regression Estimate of Slope
A derivation of the slope for orthogonal regression can be found in Casella and Berger,
chapter 12[27]. Here, we report the result. Let x_i be the activity of a voxel at low contrast on trial i,
out of n trials; let y_i be the activity of a voxel at high contrast on an analogous trial (i.e., the beta
weight for the same orientation in the same run); let $\bar{x}$ be the voxel's average activity at low-
contrast; and let $\bar{y}$ be the voxel's average activity at high-contrast. The following are the sums of
squares and the cross-product.
706

$$
\begin{aligned}
S_{xx} &= \sum_{i}^{n}(x_i - \bar{x})^2 \\
S_{yy} &= \sum_{i}^{n}(y_i - \bar{y})^2 \\
S_{xy} &= \sum_{i}^{n}(x_i - \bar{x})(y_i - \bar{y})
\end{aligned}
$$

708
709   The slope, g, that solves the total least squares problem is given by
710

$$
g = \frac{-(S_{xx} - S_{yy}) + \sqrt{(S_{xx} - S_{yy})^2 + 4S_{xy}^2}}{2S_{xy}} \tag{5}
$$

712
As stated in the main text, a separate slope was estimated for each voxel. In one session,
one participant was shown seven rather than eight orientations. When calculating the slopes for
this participant, the activity from those sessions were excluded.

Orientation Preference Simulation
To show that there are circumstances in which a von Mises density function is a
reasonable weight distribution, we simulated cortical orientation preference maps (Figure 2).
Each voxel was constructed with a Kohonen network. We tailored the network parameters so
that, following self-organization, each voxel would contain a roughly accurate number of
pinwheels for a voxel of size 2.2mm^3 – the voxel sized used in this study – approximately two
pinwheels per millimeter[68]. However, this simulation is meant as a "proof of concept", rather
than a demonstration that the von Mises distribution is an ideal weight distribution. That is, it
does not consider many factors that likely influence how neuronal activity will be reflected in
voxel activity[48]. For example, it does not consider the variability in pinwheel density across V1,
vasculature, nor how cortical folding would allow disparate parts of cortex to be sampled by the
same voxel. Nevertheless, it shows why we might expect weight distributions in most voxels to

be relatively flat, with the deviations from uniformity being captured adequately by the single
mode of the von Mises distribution.
Each network was simulated as follows. First, a grid of 67 x 67 "neurons" was initialized
to have a random orientation preference. Then, for 20000 iterations, the grid was presented with
a random orientation stimulus. The neuron that preferred an orientation closest to the stimulus
was selected, and the weights were updated. Weights were updated based on the proximity of a
neuron in the grid to the winning neuron. Specifically, on iteration $i + 1$, the weight, w, of
neuron j was given by the following.
736

737    $$w_{j,i+1} = w_{j,i} + \frac{\psi(d_j, \sigma_i)}{\psi(0, \sigma_i)} \delta_i \qquad (6)$$

738
In Equation 6, $\psi(x, \sigma)$ is the density of x as assigned by the density function of a normal
distribution with mean 0 and standard deviation σ. The variable d_j is the distance on the grid
between neuron j and the winning neuron (Euclidean distance, as calculated after setting a city-
block distance of 0.03 between adjacent neurons). The learning rate δ_i is the difference between
the neuron's preferred orientation and the presented stimulus. Finally, σ_i changes the effective
distance between neurons over the course of learning, altering the size of the update to preferred
orientation for neurons that neighbor the winning neuron. On each iteration, σ_i is equal to $0.06 *$
$((20000 - i + 1)/20000)$.

Acknowledgements

748   The work was supported by NIH award 1RF1MH114277-01 to R.A.C. and D.E.H.

Data Availability

750   The imaging and behavioral data that support the findings of this study are available as a
751   repository on the Open Science Framework (10.17605/OSF.IO/93YNM).

Code Availability

753   Software to perform the analyses is available on the Open Science Framework
754   (10.17605/OSF.IO/93YNM).

References

1.   Hubel, D. H. & Wiesel, T. N. Receptive fields of single neurones in the cat's striate cortex.
J. Physiol. **148**, 574–91 (1959).
2.   McAdams, C. J. & Maunsell, J. H. Effects of attention on orientation-tuning functions of
single neurons in macaque cortical area v4. *J. Neurosci.* **19**, 431–41 (1999).
3.   Allen Institute for Brain Science. Allen Brain Observatory. observatory.brain-
map.org/visualcoding/ (2016).
4.   Leuba, G. & Garey, L. J. Comparison of neuronal and glial numerical density in primary
and secondary visual cortex of man. *Exp. Brain Res.* **77**, (1989).
5.   Dumoulin, S. O. & Wandell, B. A. Population receptive field estimates in human visual
cortex. *NeuroImage* **39**, 647–660 (2008).
6.   Kamitani, Y. & Tong, F. Decoding the visual and subjective contents of the human brain.
Nat. Neurosci. **8**, 679–685 (2005).

7.   Boynton, G. M. Imaging orientation selectivity: decoding conscious perception in v1. *Nat.*
Neurosci. **8**, 541–542 (2005).
8.   Saproo, S. & Serences, J. T. Spatial attention improves the quality of population codes in
human visual cortex. *J. Neurophysiol.* **104**, 885–895 (2010).
9.   Scolari, M. & Serences, J. T. Basing perceptual decisions on the most informative sensory
neurons. *J. Neurophysiol.* **104**, 2266–2273 (2010).
10.   Serences, J. T., Saproo, S., Scolari, M., Ho, T. & Muftuler, L. T. Estimating the influence
of attention on population codes in human visual cortex using voxel-based tuning functions.
NeuroImage **44**, 223–231 (2009).
11.   Zhang, J., Meeson, A., Welchman, A. E. & Kourtzi, Z. Learning alters the tuning of
functional magnetic resonance imaging patterns for visual forms. *J. Neurosci.* **30**, 14127–
14133 (2010).
12.   Çukur, T., Nishimoto, S., Huth, A. G. & Gallant, J. L. Attention during natural vision warps
semantic representation across the human brain. *Nat. Neurosci.* **16**, 763–70 (2013).
13.   Nevado, A., Young, M. P. & Panzeri, S. Functional imaging and neural information coding.
NeuroImage **21**, 1083–1095 (2004).
14.   Sprague, T. C. *et al.* Inverted Encoding Models Assay Population-Level Stimulus
Representations, Not Single-Unit Neural Tuning. *Eneuro* **5**, ENEURO.0098-18.2018
(2018).
15.   Brouwer, G. J. & Heeger, D. J. Decoding and Reconstructing Color from Responses in
Human Visual Cortex. *J. Neurosci.* **29**, 13992–14003 (2009).
16.   Kay, K. N., Naselaris, T., Prenger, R. J. & Gallant, J. L. Identifying natural images from
human brain activity. *Nature* **452**, 352–355 (2008).
17.   Mitchell, T., Shinkareva, S. & Carlson, A. Predicting human brain activity associated with
the meanings of nouns. *Science* **320**, 1191–1195 (2008).
18.   Keliris, G. A., Li, Q., Papanikolaou, A., Logothetis, N. K. & Smirnakis, S. M. Estimating
average single-neuron visual receptive field sizes by fMRI. *Proc. Natl. Acad. Sci.* **116**,
201809612 (2019).
19.   Alitto, H. J. & Usrey, W. M. Influence of Contrast on Orientation and Temporal Frequency
Tuning in Ferret Primary Visual Cortex. *J. Neurophysiol.* **91**, 2797–2808 (2004).
20.   Sclar, G. & Freeman, R. D. Orientation selectivity in the cat's striate cortex is invariant
with stimulus contrast. *Exp. Brain Res.* **46**, 457–461 (1982).
21.   Skottun, B. C., Bradley, A., Sclar, G., Ohzawa, I. & Freeman, R. D. The effects of contrast
on visual orientation and spatial frequency discrimination: a comparison of single cells and
behavior. *J. Neurophysiol.* **57**, 773–786 (1987).
22.   Liu, T., Cable, D. & Gardner, J. L. Inverted encoding models of human population response
conflate noise and neural tuning width. *J. Neurosci.* **38**, 398–408 (2018).
23.   Swindale, N. V. Orientation tuning curves: empirical description and estimation of
parameters. *Biol. Cybern.* **78**, 45–56 (1998).
24.   Mardia, K. V. & Jupp, P. E. *Directional statistics*. (John Wiley & Sons, Inc., 1999).
25.   Gelman, A. *et al. Bayesian Data Analysis*. (Chapman and Hall/CRC, 2013).
26.   Wagenmakers, E.-J., Ratcliff, R., Gomez, P. & Iverson, G. J. Assessing model mimicry
using the parametric bootstrap. *J. Math. Psychol.* **48**, 28–50 (2004).
27.   Casella, G. & Berger, R. L. *Statistical inference*. (Duxbury Press, 2001).
28.   Vehtari, A., Gelman, A. & Gabry, J. Practical Bayesian model evaluation using leave-one-
out cross-validation and WAIC. *Stat. Comput.* **27**, 1413–1432 (2017).

29.   Dosher, B. A. & Lu, Z.-L. Perceptual learning reflects external noise filtering and internal
noise reduction through channel reweighting. *Proc. Natl. Acad. Sci.* **95**, 13988–13993
(1998).
30.   Ling, S., Liu, T. & Carrasco, M. How spatial and feature-based attention affect the gain and
tuning of population responses. *Vision Res.* **49**, 1194–1204 (2009).
31.   Treisman, A. M. & Gelade, G. A feature-integration theory of attention. *Cognit. Psychol.*
12, 97–136 (1980).
32.   Schneider, W. & Shiffrin, R. M. Controlled and automatic human information processing: I.
Detection, search, and attention. *Psychol. Rev.* **84**, 1–66 (1977).
33.   Buxton, R. B., Wong, E. C. & Frank, L. R. Dynamics of blood flow and oxygenation
changes during brain activation: The balloon model. *Magn. Reson. Med.* **39**, 855–864
(1998).
34.   Stephan, K. E., Harrison, L. M., Penny, W. D. & Friston, K. J. Biophysical models of fMRI
responses. *Curr. Opin. Neurobiol.* **14**, 629–635 (2004).
35.   Alink, A., Abdulrahman, H. & Henson, R. N. Forward models demonstrate that repetition
suppression is best modelled by local neural scaling. *Nat. Commun.* **9**, (2018).
36.   Brouwer, G. J. & Heeger, D. J. Cross-orientation suppression in human visual cortex. *J.*
Neurophysiol. **106**, 2108–2119 (2011).
37.   Naselaris, T., Kay, K. N., Nishimoto, S. & Gallant, J. L. Encoding and decoding in fMRI.
NeuroImage **56**, 400–410 (2011).
38.   Sprague, T. C., Saproo, S. & Serences, J. T. Visual attention mitigates information loss in
small- and large-scale neural codes. *Trends Cogn. Sci.* **19**, 215–226 (2015).
39.   Gardner, J. L. & Liu, T. Inverted encoding models reconstruct an arbitrary model response,
not the stimulus. *eNeuro* **6**, ENEURO.0363-18.2019 (2019).
40.   Sprague, T. C., Boynton, G. M. & Serences, J. T. The importance of considering model
choices when interpreting results in computational neuroimaging. *eNeuro* **6**,
ENEURO.0196-19.2019 (2019).
41.   Ramirez, F. M., Cichy, R. M., Allefeld, C. & Haynes, J.-D. The Neural Code for Face
Orientation in the Human Fusiform Face Area. *J. Neurosci.* **34**, 12155–12167 (2014).
42.   Weiner, K. S., Sayres, R., Vinberg, J. & Grill-Spector, K. fMRI-adaptation and category
selectivity in human ventral temporal cortex: regional differences across time scales. *J.*
Neurophysiol. **103**, 3349–3365 (2010).
43.   Heeger, D. J., Huk, A. C., Geisler, W. S. & Albrecht, D. G. Spikes versus BOLD: what
does neuroimaging tell us about neuronal activity? *Nat. Neurosci.* **3**, 631–633 (2000).
44.   Rees, G., Friston, K. & Koch, C. A direct quantitative relationship between the functional
properties of human and macaque v5. *Nat. Neurosci.* **3**, 716–723 (2000).
45.   Boynton, G. M., Engel, S. A., Glover, G. H. & Heeger, D. J. Linear systems analysis of
functional magnetic resonance imaging in human v1. *J. Neurosci.* **16**, 4207–21 (1996).
46.   Vazquez, A. L. & Noll, D. C. Nonlinear Aspects of the BOLD Response in Functional
MRI. *NeuroImage* **7**, 108–118 (1998).
47.   González Ballester, M. Á., Zisserman, A. & Brady, M. Segmentation and measurement of
brain structures in MRI including confidence bounds. *Med. Image Anal.* **4**, 189–200 (2000).
48.   Kriegeskorte, N., Cusack, R. & Bandettini, P. How does an fMRI voxel sample the
neuronal activity pattern: compact-kernel or complex spatiotemporal filter? *NeuroImage* **49**,
1965–1976 (2010).

49.   Buxton, R. B. & Frank, L. R. A Model for the Coupling between Cerebral Blood Flow and
860   Oxygen Metabolism during Neural Stimulation. *J. Cereb. Blood Flow Metab.* **17**, 64–72
861   (1997).
862   50.   Handwerker, D. A., Ollinger, J. M. & D'Esposito, M. Variation of BOLD hemodynamic
863   responses across subjects and brain regions and their effects on statistical analyses.
864   *NeuroImage* **21**, 1639–1651 (2004).
865   51.   Bao, P., Purington, C. J. & Tjan, B. S. Using an achiasmic human visual system to quantify
866   the relationship between the fMRI BOLD signal and neural response. *eLife* **4**, 1–21 (2015).
867   52.   Kay, K. N., Winawer, J., Mezer, A. & Wandell, B. A. Compressive spatial summation in
868   human visual cortex. *J. Neurophysiol.* **110**, 481–494 (2013).
869   53.   Dragoi, V., Sharma, J. & Sur, M. Adaptation-induced plasticity of orientation tuning in
870   adult visual cortex. *Neuron* **28**, 287–298 (2000).
871   54.   Brainard, D. H. The Psychophysics Toolbox. *Spat. Vis.* **10**, 433–436 (1997).
872   55.   Cornelissen, F. W., Peters, E. M. & Palmer, J. The Eyelink Toolbox: eye tracking with
873   MATLAB and the Psychophysics Toolbox. *Behav. Res. Methods Instrum. Comput.* **34**,
874   613–617 (2002).
875   56.   Rademaker, R. L., Chunharas, C. & Serences, J. T. Coexisting representations of sensory
876   and mnemonic information in human visual cortex. *Nat. Neurosci.* **22**, 1336–1344 (2019).
877   57.   Benson, N. C. *et al.* The Human Connectome Project 7 Tesla retinotopy dataset:
878   Description and population receptive field analysis. *J. Vis.* **18**, 23 (2018).
879   58.   Kriegeskorte, N. *et al.* Matching Categorical Object Representations in Inferior Temporal
880   Cortex of Man and Monkey. *Neuron* **60**, 1126–1141 (2008).
881   59.   Behzadi, Y., Restom, K., Liau, J. & Liu, T. T. A component based noise correction method
882   (CompCor) for BOLD and perfusion based fMRI. *NeuroImage* **37**, 90–101 (2007).
883   60.   Roth, Z. N., Heeger, D. J. & Merriam, E. P. Stimulus vignetting and orientation selectivity
884   in human visual cortex. *eLife* **7**, 1–25 (2018).
885   61.   Carlson, T. A. Orientation Decoding in Human Visual Cortex: New Insights from an
886   Unbiased Perspective. *J. Neurosci.* **34**, 8373–8383 (2014).
887   62.   Benson, N. C. & Winawer, J. Bayesian analysis of retinotopic maps. *eLife* **7**, 0–45 (2018).
888   63.   Morey, R. D. Confidence intervals from normalized data: a correction to cousineau (2005
889   ). *Tutor. Quant. Methods Psychol.* **4**, 61–64 (2008).
890   64.   Carpenter, B. *et al.* Stan : A Probabilistic Programming Language. *J. Stat. Softw.* **VV**, 1–43
891   (2016).
892   65.   Duane, S., Kennedy, A. D., Pendleton, B. J. & Roweth, D. Hybrid Monte Carlo. *Phys. Lett.
893   B* **195**, 216–222 (1987).
894   66.   Hoffman, M. & Gelman, A. The No-U-Turn sampler: adaptively setting path lengths in
895   Hamiltonian Monte Carlo. *J. Mach. Learn. Res.* **15**, 1593–1623 (2014).
896   67.   Gelman, A. & Rubin, D. B. Inference from Iterative Simulation Using Multiple Sequences.
897   *Stat. Sci.* **7**, 457–472 (1992).
898   68.   Yacoub, E., Harel, N. & Ugurbil, K. High-field fMRI unveils orientation columns in
899   humans. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 10607–12 (2008).
900