Homothorax Controls a Binary Rhodopsin Switch in *Drosophila* Ocelli

Abhishek Kumar Mishra[1]*, Cornelia Fritsch[1], Ruben Voutev[2], Richard S. Mann[2] and Simon G. Sprecher[1,3]*

[1]Institute of Cell and Developmental Biology, Department of Biology, University of Fribourg, Chemin du Musée 10, 1700, Fribourg, Switzerland, [2]Department of Biochemistry and Molecular Biophysics and Neuroscience, Mortimer B. Zukerman Mind Brain Behavior Institute, Columbia University, New York, NY 10027, USA

[3]Lead Contact

*Correspondence: simon.sprecher@gmail.com; abhishekkumar.mishra@unifr.ch

Keywords: *Drosophila*, Photoreceptor, Homothorax, Ocelli, Compound eye, Rhodopsin, Gene duplication, Evolution, cell fate

Running title: Hth regulates binary rhodopsin switch in *Drosophila* ocelli

Abstract

Visual perception of the environment is mediated by specialized photoreceptor (PR) neurons of the eye. Each PR expresses photosensitive opsins, which are activated by a particular wavelength of light. In most insects, the visual system comprises a pair of compound eyes that are mainly associated with motion detection, color or polarized light perception and a triplet of ocelli that are thought to be critical during flight to detect horizon and movements. It is widely believed that evolutionary diversification of compound eye and ocelli in insects occurred from an ancestral visual organ around 500 million years ago. Concurrently, opsin genes were also duplicated to provide distinct spectral sensitivities to different PRs of compound eye and ocelli. In the fruit fly *Drosophila melanogaster*, Rhodopsin1 (Rh1) and Rh2 are closely related opsins that are originated from the duplication of a single ancestral gene. However, in the visual organs, Rh2 is uniquely expressed in ocelli whereas Rh1 is uniquely expressed in outer PRs of the compound eye. It is currently unknown how this differential expression of Rh1 and Rh2 in the two visual organs is controlled to provide unique spectral sensitivities to ocelli and compound eyes. Here, we show that Homothorax (Hth) is expressed in ocelli and confers proper *rhodopsin* expression. We find that Hth controls a binary rhodopsin switch in ocelli to promote Rh2 expression and repress Rh1 expression. Genetic and molecular analysis of *rh1* and *rh2* supports that Hth acts through their promoters to regulate rhodopsin expression in the ocelli. Finally, we also show that when ectopically expressed in the retina, *hth* is sufficient to induce Rh2 expression only at the outer PRs in a cell autonomous manner. We therefore propose that the diversification of *rhodpsins* in the ocelli and retinal outer PRs occurred by duplication of an ancestral gene, which is under the control of Homothorax.

Introduction

The ability to perceive and discriminate a broad range of environmental stimuli

in nature is essential for many aspects of life. Animals rely heavily on visual cues to

perform complex tasks such as navigation to find food, mates and shelter as well as

social interactions. Visual cues are perceived by visual organs that contain

photoreceptors (PRs) as light-sensing structures. PRs are specialized cells that gather

information from the surrounding world which is subsequently processed by the brain.

Each PR expresses a unique photosensitive *opsin/rhodopsin* and they define the

wavelength of light by which a PR will be activated.

It is believed that eyes have evolved separately due to fundamental differences

between visual organs of different animals [1]. However, it is also known that eye

development in different animal phyla share a common genetic network initiated by

Pax6 gene orthologs [2]. Finding similarities in the gene regulatory network that

controls eye development further strengthened the idea that phylogenetically diverse

eye types may share a conserved eye developmental program [3-5]. Insects are

among the largest and most diverse animal groups. Decoding eye development in

insects offers great opportunity to unravel developmental insights that lead to the

emergence of evolutionary complexity.

In most insects, compound eyes represent the prominent visual organ that is

responsible for providing major share of the visual information. In the fruit fly

Drosophila melanogaster, each compound eye consists of approximately 850

ommatidia and each ommatidium houses eight PRs: six outer PRs and two inner PRs

[6]. Additionally, winged insects (such as *Drosophila*) possess ocelli that are

comparatively simple photosensory organs embedded in the dorsal head cuticle [7].

In *Drosophila*, a triplet of ocelli (one medial and two lateral) are arranged in a triangular

shape between the two compound eyes and the dorsal vertex of the head [8]. It is

believed that insect compound eye- and ocellus-like precursor structures have segregated from an ancestral eye over 500 million years ago [9, 10]. In order for these visual organs to perform different functions that require distinct spectral sensitivities, new opsin genes evolved by gene duplication during evolution [11]. In *Drosophila,* phylogenetic analysis supports that Rhodopsin1 (Rh1), Rh2 and Rh6 have originated from a common ancestral gene [12]. A first gene duplication may have separated Rh6 from Rh1/Rh2 and a second gene duplication may have separated the closely related Rh1 and Rh2 [12]. The green sensitive Rh6 is expressed in the inner PRs of the compound eye and is critically involved in color perception [13-15]. The blue-green sensitive Rh1 is expressed in the outer PRs of the compound eye and is mainly associated with motion detection [16-18]. Conversely, the UV-blue sensitive Rh2 is expressed in PRs of the ocelli [19-21] and is proposed to be involved in horizon sensing and flight stabilization [22, 23]. While it is known that *rh1, rh2* and *rh6* are genetically linked [24], it is still unknown how they are differentially expressed in different PRs in *Drosophila*.

Here we show that the homeodomain transcription factor Homothorax (Hth) regulates Rh2 expression in the ocelli. We demonstrate that Hth is expressed in ocellar PRs and controls a binary *rhodopsin* switch by promoting Rh2 expression and repressing Rh1 expression in ocelli. We also demonstrate that misexpression of Hth forces outer PR of the retina to induce Rh2 expression and clonal expression in the retina suggest that this process is cell autonomous. Furthermore, genetic and molecular analysis of *rh1* and *rh2* shows that the *rhodopsin* switch in ocelli is transcriptionally controlled by Hth and that it may act directly through *rh1* and *rh2* promoter sequences. Finally, we argue that while Hth maintains Rh3 fate in the DRA [25], it initiates Rh2 fate in the ocelli. The results presented here greatly adds to our

97   understanding of how genetically linked opsins are spatiotemporally controlled to

98   provide distinct spectral sensitivity to different visual organs.

Results

Differential expression of *rhodopsins* in the compound eye and ocelli

It is believed that the hexapod ancestor of extant insects only had a single visual organ (also referred as ancestral eye) and that other visual organs (such as compound eye and ocelli) evolved as a result of a morphological bifurcation event over 500 million years ago [10, 11] (Figure 1A). It is hypothesised that the different functions of compound eye and ocelli in conjunction of gene duplication events of opsins led to the emergence of an ocelli specific *rhodopsin* gene, with a distinct spectral sensitivity [11]. In most insects, three different opsin subfamilies are present that show distinct spectral sensitivity to different wavelengths of light and are associated with different functions. Ultraviolet-sensing (UV) opsins are sensitive to 300-400 nm range of wavelength whereas blue-sensing (B) opsins are sensitive to a wavelength of 400-500 nm. Long wavelength-sensing (LW) opsins are sensitive to a range between 500-600 nm [26-28]. In the *Drosophila* compound eye, brightness detection is achieved by inputs from six outer-PRs (R1-R6) that express the LW-opsin Rhodopsin 1 (Rh1). *Drosophila* uses outer PRs mainly for motion detection and dim light vision [16-18]. Color vision requires the two inner PRs (R7 and R8) that encode either one of the UV- (Rh3 or Rh4) or the B- (Rh5) or LW-opsins (Rh6) [29, 30]. In most insects, ocelli express a LW-opsin, which is different from the one present in the compound eye [11, 21, 31]. In *Drosophila*, PRs in the ocelli express the LW-opsin Rh2 that is sensitive to a wavelength range of 418-506 nm [19-21]. The different spectral sensitivity of ocelli is also reflected by their involvement to perform different functions than the compound eye and they are believed to detect horizon, control head orientation and stabilize flight posture while flying [22]. To get an insight of *rhodopsin* gene duplications in *Drosophila*, we compared the nucleotide sequences of *rh1* to *rh6* by generating a

phylogenetic tree (by using MUSCLE online tool; phylogeny.fr) [32]. The phylogenetic

tree made by maximum likelihood method showed two clades with branching support

value of 1 each (Figure 1B). Clade-I showed tandem gene duplication that separated

Rh5 from the closely related Rh3/Rh4. Clade-II represents LW-opsin gene

duplications in *Drosophila* where first a tandem gene duplication separated Rh6 from

Rh1/Rh2 (Figure 1B). Rh6 subsequently got expressed in the inner PRs of the

compound eye (Figure 1H). A further gene duplication led to the separation of the blue

shifted rhodopsins Rh1 and Rh2 (Figure 1B). While Rh1 subsequently got expressed

in the outer PRs of the compound eye (Figure 1F), Rh2 got exclusively expressed in

ocelli (Figure 1D).

Homothorax regulates a binary Rhodopsin switch in ocelli

In the compound eye, the homeodomain transcription factor Homothorax (Hth)

is expressed in the dorsal rim area (DRA) of inner PRs, where it has been shown to

be both necessary and sufficient to regulate Rh3 expression and thereby critically

contributes to the polarized-light sensing system [25, 33]. We found that Hth is also

expressed in all PRs of the ocelli (Figure 2A). Since ocellar PRs expresse Rh2, we

first analysed if Hth is involved in regulating Rh2 expression. We performed

knockdown of *hth* using the pan-photoreceptor driver *lGMR*-Gal4 and observed a

complete loss of Rh2 expression in PRs of the ocelli (Figure 2B). Additionally, we also

observed that in *hth* knockdown, the loss of Rh2 expression in the ocelli is

compensated by gain of Rh1 (Figure 2B), a Rhodopsin that is normally exclusively

expressed in the outer PRs of the compound eye. Thus, Hth regulates a binary switch

of Rhodopsin in the ocelli where it promotes Rh2 expression and represses Rh1

expression.

In the retina of the compound eye, a feedback mechanism has been described

that allows Rh6 to transcriptionally repress Rh5 in yellow R8 PRs [34]. Since in *hth*

knockdown, ectopic Rh1 expression in the ocelli replaces Rh2, we speculated that Hth

may repress Rh1 in order to allow Rh2 expression. Knockdown of *hth* in turn removed

this repression and as a result Rh1 gets activated and represses Rh2 in ocelli. To

check this hypothesis, we misexpressed Rh1 in ocelli by using *lGMR-Gal4* and found

that ectopic expression of Rh1 was not sufficient to repress Rh2 and in fact in this case

ocelli co-expressed both Rh1 and Rh2 (Supplemental figure 1A). Thus, Rh2 loss in

hth knockdown seems to be independent of Rh1 repression.

To further test whether switching Rhodopsin expression might also alter PR

identity based on other molecular markers of outer PRs in the retina, we monitored

Seven-up (svp) and BarH1 expression in the ocelli in *hth* knockdown. Svp is a steroid

hormone receptor and is expressed in R3/R4 and R1/R6 pairs whereas BarH1 is a

homeobox transcription factor expressed in the R1/R6 pair of the developing

compound eye [35, 36]. We found that neither Svp nor BarH1 are expressed in

wildtype nor in *hth* knockdown ocelli (Supplemental figure 2A, 2B, 2C and 2D).

Therefore, although ocellar PRs gained Rh1 and lost Rh2 expression when *hth* was

knocked down, they do not appear to undergo an identity change towards retinal outer

PRs.

We next investigated whether regulation of this Rhodopsin switch by Hth occurs

at the transcriptional level by using ocelli specific *rh2-lacZ* [20] and outer PR specific

rh1-lacZ [36] reporter lines. In wildtype control animals, β-gal expression was

specifically observed in ocelli in case of *rh2-lacZ* (Figure 2C) whereas no expression

was observed in ocelli in the case of *rh1-lacZ* (Figure 2E). In *hth* knockdown

background, we found that *β*-gal expression from *rh2-lacZ* is completely abolished

from ocelli (Figure 2D), whereas ectopic *β*-gal expression from *rh1-lacZ* is now seen

in ocellar PRs (Figure 2F). Therefore, Hth is indeed involved in regulating a binary

rhodopsin switch in the ocelli by promoting transcription of *rh2* and repressing

transcription of *rh1*.

We have previously shown that the homeodomain transcription factor Hazy

(Flybase: Pph13 for PvuII-PstI homology 13) controls expression of Rh2 in the ocelli

[37, 38]. We therefore next asked if Hazy and Hth act jointly to regulate the rhodopsin

switch in ocelli. However, we find that in *hazy$^{-/-}$* null mutant flies, the absence of Rh2

expression is not accompanied with the ectopic expression of Rh1 in ocellar PRs

(Supplemental figure 1B). Moreover, we found that Hth is still expressed in the ocelli

of *hazy$^{-/-}$* mutant flies (Supplemental figure 1C). Also, the expression of Hazy remains

unchanged in ocellar PRs when knocking down *hth* (Supplemental figure 1D). Thus,

the Hth-dependent Rhodopsin switch in ocelli does not depend on Hazy.

Hth controls a binary rhodopsin switch in ocelli by acting through the promoters

of *rh1* and *rh2*

Since Hth encodes a homeodomain transcription factor we next investigated if

Hth may act by directly regulating the *rhodopsin* promotors. We identified three

potential Hth binding sites in the minimal promoter region of *rh2* (-293/+55) [20],

whereas the *rh1* minimal promoter region (-247/+73) [36] contains two potential Hth

binding sites. To test if the Hth binding sites in the *rh1* and *rh2* promoter regions may

be involved in the rhodopsin switch in ocelli, we created transgenic flies containing

transgenes with the minimal promoter regions of *rh1* or *rh2* driving GFP (*rh1-GFP* and

rh2-GFP [37]). Next, in order to abolish Hth binding, we introduced point mutations in

the Hth binding regions of the *rh1* and *rh2* promoters and created *rh1(hth mut)-GFP*

and *rh2(hth mut)-GFP* transgenic flies (Figure 3A and 3D) (See material and methods

for details). In support with the previous observations, we do observe *rh2-GFP*

expression in the ocelli (Figure 3B) whereas *rh1-GFP* is not expressed in ocellar PRs

(Figure 3E). However, by mutatingthe Hth binding sites in the promoter region of *rh2*

(*Rh2 (hth mut)-GFP*), we observed a loss of GFP expression in the ocelli (Figure 3C).

Conversely, we found that deleting the Hth binding sites in the promoter region of *rh1*

(*rh1 (hth mut)-GFP)* leads to ectopic GFP expression in ocelli (Figure 3F).

Hth is sufficient to induce Rh2 expression in the outer PRs of retina

In the dorsal rim area of the retina, Hth is required in inner PRs to promote Rh3

expression and ectopic expression of Hth under the control of *lGMR*-Gal4 was

sufficient to block the expression of inner PR rhodopsins (Rh4, Rh5 and Rh6) and to

induce Rh3 expression in all inner PRs. However, in outer PRs, expression of Rh1

was not affected suggesting that only inner PRs were responsive to Hth misexpression

[25]. We next investigated if misexpression of Hth in the retina was also sufficient to

induce Rh2 expression. In the wildtype retina, Rh1 was uniquely expressed in outer

PRs whereas Rh2 was absent (Figure 4A, 4A' and 4A''). When Hth was misexpressed

in the retina under the control of *lGMR*-Gal4, we observed that Rh2 was now

ectopically co-expressed in all outer PRs. However, we found that Rh1 expression

was unaffected and it was still expressed in all outer PRs (Figure 4B, 4B' and 4B'').

We also found that ectopic Rh2 expression was only limited to the outer PRs (and not

inner PRs) suggesting that only Rh1 expressing PRs were competent to induce Rh2

expression in the retina, while inner PRs were not.

To investigate if this process is cell autonomous, we generated outer PR versus

inner PR Hth gain-of-function clones in the retina using the "flip-out" technique [39]

(material and methods for details). Flip-out clones were marked by presence of GFP

expression. We found that when clones were generated in the inner PRs, Rh2

expression was not induced (Figure 5A, 5A' and 5A''). However, when clones were

generated in the outer PRs, Rh2 expression was ectopically induced in all GFP

expressing clones (Figure 5B, 5B' and 5B''). Thus, ectopic Rh2 expression in Rh1

expressing PRs of the retina is a cell-autonomous process. Expression of Hth in inner

PR clones did not have any effect on Rh1 expression in the neighbouring outer PRs

(Figure 5C, 5C' and 5C'') and Rh1 expression was maintained when clones were

generated in the outer PRs (Figure 5D, 5D' and 5D'').

Taken together, we showed that Hth regulates a binary rhodopsin switch in

ocelli by promoting Rh2 expression at the cost of Rh1. We found that regulation of

Rh1 and Rh2 is transcriptionally controlled by Hth. We also found that Hth regulates

the rhodopsin switch by acting through the promoters of *rh1* and *rh2*. We further

demonstrated that misexpression of Hth in the retina modifies outer PRs to acquire

Rh2 expression resulting in outer PRs co-expressing Rh1 and Rh2 and we show that

this process is cell-autonomous (Figure 6).

Discussion

Homothorax expression provides new insights to understand Rhodopsin fate in

Drosophila

In insects, ocelli represent a fundamentally simpler visual organ whose spectral

sensitivity is different from the compound eye. The unique spectral sensitivity of ocellar

PRs in *Drosophila* is provided by the UV-blue sensing Rh2 [21]. The presence of this

particular Rhodopsin exclusively in ocelli and not in the retina may explain why ocelli

perform different functions than the compound eye. We have characterized the roles

of Hth during terminal differentiation of ocellar PRs and showed that Hth regulates a

binary rhodopsin switch in ocelli that promotes Rh2 expression and represses Rh1

expression. Hth is known to be expressed in the ocellar primordium of the early third

instar laval (L3) eye antennal imaginal disc but gets downregulated later at mid- to

late-L3 [40]. In our study, we show that Hth is expressed in all mature and terminally

differentiated PRs of ocelli.

Hth performs similar function in PRs of the ocelli and the DRA of the retina: in

ocelli, it induces Rh2 expression by repressing Rh1 whereas in the DRA of the retina

it induces Rh3 by repressing inner-PR Rhodopsins [25]. Loss of *hth* transforms the

DRA into odd-coupled ommatidia where Rh3 is expressed in R7 and Rh6 in R8

suggesting that the Rh3/Rh6 pair represents the default state of rhodopsins in inner

PRs of the retina [25]. In ocelli, we find ectopic expression of Rh1 by loss of *hth*

suggesting that Rh1 may be the default state in ocellar PRs. Loss of Hth leads to

ectopic expression of the R8 specific transcription factor Senseless in the DRA, which

is otherwise not expressed in the wildtype DRA suggesting a fate change of the DRA

to become odd-coupled ommatidia [33]. Interestingly, Hth loss in ocelli does not induce

expression of outer PR-specific transcription factors (such as Svp and BarH1)

suggesting that the ocellar PR fate may not have been changed.

Genetic and molecular dissection of Hth in the ocelli during PR development

The DNA binding property of Hth is required for proper DRA fate in the retina

[33]. Similarly, we observed that Hth regulates binary rhodopsin switch in ocelli by

transcriptionally controlling *rh1* and *rh2* expression through binding to their upstream

DNA sequences. The homeodomain transcription factor Hazy controls rhodopsin

expression in the retina and in ocelli [37, 38]. However, epistatic analysis showed that

both Hth and Hazy acted independently to control Rh2 expression in ocelli.

Misexpression of *hth* in all developing PRs of the retina is sufficient to induce a fate

switch where inner PRs are transformed into PRs of DRA [33]. Rh3 is expanded to all

inner PRs whereas specific loss of Rh4, Rh5 and Rh6 was seen. However, outer PR

fate was not changed suggesting that only those PRs, which were previously

committed to become inner PRs during development, were responsive to Hth [33]. We

additionally found that outer PRs, too were responsive to *hth* misexpression since they

induce Rh2 expression while maintaining its default fate by expressing Rh1. However,

the genetic program activated by Hth misexpression to induce Rh2 expression in outer

PRs is still unknown.

Rhodopsin duplication and role of Hth during ocelli diversification.

Photoreception is achieved by expression of different opsins in conjunction with an

array of proteins of the phototransduction cascade. Each PR contains different opsins,

which are light sensing structures and they define the particular wavelength of light to

which a given PR will respond. In most insect species, opsins were categorized into

three major clades: 1) short-wavelength (SW) opsins, which are mostly UV-sensitive,

2) middle-wavelength (MW) opsins, which are mainly blue-sensitive and 3) long-

wavelength (LW) opsins, which are ranging from blue-violet such as Rh2 in *Drosophila*

[41] to green-red in some butterflies [42]. Presence of all three opsins in most insect

species may imply that the ancient retina was trichromatic [43]. Further phylogenetic analyses support ancestral trichromacy in insects and show that SW, MW and LW sensitive opsins must have derived from a single ancestral opsin as a result of gene duplications [12]. In many insect species, spectral sensitivity of opsin gene clades is represented by more than one paralog such as the LW-opsin clade consisting of three paralogs in *Drosophila* (Rh1, Rh2 and Rh6), four in the mosquito *Anopheles gambiae* and two in the honeybee *Apis mellifera* [12]. It is believed that a first tandem gene duplication in *Drosophila* separated Rh6 from its sister paralog Rh1/Rh2 and while Rh6 is retained in the inner PRs of compound eye, Rh1/Rh2 was associated with the outer PRs of retina and PRs of ocelli. A second gene duplication event led to the diversification of Rh1 and Rh2 by accumulation of further amino acid changes in their sequence. While Rh1 expression was retained in the outer PRs of the compound eye, Rh2 became associated with the PRs of the ocelli. Our results also suggest that the ability of Hth to bind to both promoters may have been a prerequisite during evolution to differentially regulate the expression of the two paralogs and thus, provide spectral sensitivity of ocelli by promoting Rh2 expression and repressing Rh1 expression. However, it would be interesting to know if a change in the spectral sensitivity of ocelli upon Hth loss would in turn change their specific functions. Also, if gene duplications of opsins occurred during evolution to allow the diversification of different spectral sensitivities in different visual organs, it would be interesting to know what kind of spectral sensitivity was present in the ancestral visual organ. We conclude that Hth may act as an evolutionary factor required by ocelli to provide their unique spectral identity to ocellar PRs that are marked by expression of Rh2. To maintain this unique spectral identity, Hth controls a binary rhodopsin switch to repress outer PR fate in ocelli by repressing Rh1 expression.

The opposing regulatory function of Hth on its direct targets.

We show here that Hth functions both as a transcriptional activator of *rh2* and as a transcriptional repressor of *rh1*. Usually, Hth in conjunction with its binding partner Extradenticle acts as a transcriptional activator. However, it has also been shown to function in gene repression, for example by recruiting the transcriptional repressor Engrailed to the regulatory elements of target genes [44]. In this case the formation of the repressor complex occurs directly at the regulatory regions of the repressed gene and depends on the proximity of DNA-binding sites for different components of the complex in the regulatory region. The same mechanism could also explain the opposing regulatory effect that Hth has on *rh1* in comparison to *rh2* in the ocelli. Analysis of transcription profiles of different PR types would provide a list of Hth interactors expressed in the ocelli versus the outer PRs of the retina. Binding analysis of these interactors to the *rh1* and rh2 minimal promoter sequences would help to identify the potential repressor complex forming on the *rh1* promoter, helping to better understand the regulatory functions of Hth.

Author contributions

AKM and SGS designed all experiments. AKM, CF and RV performed experiments and generated reagents/materials. AKM, CF, RV, RSM and SGS wrote the manuscript.

Acknowledgements

We thank C. Desplan, J. Rister, A. Salzberg, T. Cook, D. Vasiliauskas, Developmental

Studies Hybridoma Bank (DSHB) and Bloomington Stock Center for providing flies

and antibodies. We thank UniFr bioimage team for maintaining imaging facility and all

members of Egger and Sprecher lab for fruitful discussions. This work is supported by

the Swiss National Science foundation grant number 310030_188471 to SGS, the

Novartis foundation for biomedical research grant number 18A017 to SGS and the

National Institute of Health grant number 5R35GM118336-05 to RSM.

Materials and methods

Fly stocks

Wildtype Canton S flies have been used in this study. Other fly strains used are: UAS-

hth^RNAi (BL27655 and BL34637; we do see rhodopsin switch phenotype in both RNAi

lines. However, all the experiments were done in BL27655), *lGMR*-Gal4

(Bloomington's stock center). UAS-*rh1* [34], UAS-*hth* [45], *rh1*(-252/+57)-*lacZ* [36]*, rh2*

(-309/+32)-*lacZ* [20], *rh2* (-293/+55)-*GFP* [37], *hazy^-/-* [46], *hsFLP*; UAS-*mCD8::GFP*;

lGMR<wt>Gal4 (a gift from Claude Desplan lab). The following transgenic lines were

made in this study: *rh2 (hth mut)-GFP*, *rh1-GFP* and *rh1 (hth mut)-GFP*. Flies were

reared on standard food medium and at 25°C. Knockdown experiments were

performed at 29°C temperature. For *hth* overexpression clones by "flip-out" technique,

flies were grown at 25°C and heat shock at 37°C was performed for half an hour at

the late pupal stage. Vials were placed back at 25°C and freshly hatched flies were

dissected.

Generation of *rh1* and *rh2* reporter transgenes

The *rh1* minimal promoter (-247 to +73) was PCR amplified from genomic DNA with

primers "rh1 enh Kpn fw" (gcggtacCTGGAGACTCAAGAATAATACTCGGCCAG) and

"rh1 enh Xba re" (gatctagAGGGTTCCTGGATTCTGAATATTTCACTG) and cloned

into pBluescript vector using the KpnI and XbaI sites added to the primers. For cloning

of the *rh2* minimal promoter see [37]. The same *rh1* and *rh2* promoter fragments were

in vitro synthesized (BioCat) altering the sequences of the potential Hth binding sites.

The first Hth site in the *rh1* promoter (TGACAT) was changed to TaAgcT creating a

HindIII restriction site. The second Hth site in the *rh1* promoter (CTGTCG) was

changed to CTaaaG. The first Hth site in the *rh2* promoter (GGACAG) was changed

to GtttAG, the second Hth site in the *rh2* promoter (GTGTCA) was changed into

agcTgA and the third Hth site (CTGTCC) was changed to CTaaaC. Both versions of

the two enhancers were cloned into a GFP reporter plasmid containing *eGFP,* a

miniwhite marker *and an attB* site kindly provided by Jens Rister. The plasmids were

injected into nos-φC31; attP40 flies for integration on the second choromosome using

the φC31 site-specific integration system [47].

Immunohistochemistry

Adult ocelli were dissected and stained by a protocol published in [37] whereas

dissection and immunohistochemistry of adult retinas were done according to [48].

After the immunostainings, tissue samples were mounted by using Vectashield H-

1000 (Vector laboratories). Primary antibodies and their dilutions were as follows:

Rabbit anti-Rh2 1:100 [37], Rabbit anti-Rh6 1:10,000 [49], Rabbit anti-Hth 1:500 [50],

Chicken anti-βGal 1:1000 (Abcam), Chicken anti-GFP 1:2000 (Life technologies),

Rabbit anti-Hazy 1:500 [46], Mouse anti-Svp 1:100 [51], Rat anti-BarH1 1:200 [52],

Mouse anti-Rh1 1:20, Rat anti-Elav 1:20 and Mouse anti-Chp 1:20 (Developmental

Studies Hybridoma bank). Phalloidin conjugated with Alexa-568 and Alexa-647

(Sigma-Aldrich, Life Technologies) marks F-actin and were used (1:1000) during

incubation with secondary antibodies. The following secondary antibodies were used:

Goat anti-rabbit, Goat anti-mouse, Goat anti-rat and Goat anti-chicken that are

conjugated with Alexa-488, Alexa-555 and Alexa-647 (Jackson Immunoresearch). All

secondary antibodies are used at 1:200 dilution.

Confocal microscopy and Image analysis

Tissue samples were imaged with a Leica TCS SP5 confocal microscope at a

resolution of 1024x1024 pixels and optical sections were taken in the range of 1-2 µm

depending on the sample size. Images were further processed and analysed in

Fiji/ImageJ and Adobe photoshop 2020 software.

Figure legends

Figure 1: Model representing the evolutionary diversification of the insect

ancestral eye with rhodopsin duplication and its expression in the *Drosophila*

adult visual system. (A) A morphological bifurcation from an ancestral eye of a

hexapod ancestor to extent insects over 500 million years ago (mya) is believed to

form ocelli and compound eye. (B) Gene duplications of rhodopsin in *Drosophila* where

the blue dotted square box marks a clade consisting of Rh1, Rh2 and Rh6. A first

tandem gene duplication separated Rh6 from Rh1/Rh2 and a second gene duplication

separated the closely related Rh1 and Rh2. (C, F) Expression of Rh1 (green) in the

wildtype ocelli (C) and retina (F). Rh1 is exclusively expressed in the outer PRs of the

retina but not in ocelli. (D, G) Expression of Rh2 (green) in the wildtype ocelli (D) and

retina (G). Rh2 is uniquely expressed in all PRs of the ocelli but not in the retina. (E,

H) Expression of Rh6 (green) in ocelli (E) and retina (H). Rh6 is normally expressed

in the inner PRs in retina but not in ocelli. PRs of ocelli are marked by Elav (Red)

whereas rhabdomeres in the retina are marked by Phalloidin (Red).

Figure 2: Hth expression and phenotype in ocelli. (A) Antibody staining of anti-Hth

showed that it is expressed in all PRs of wildtype ocelli. Anti-Chp (green) and anti-Elav

(blue) mark the PRs of ocelli. (B) Antibody staining of anti-Rh2 (green) and anti-Rh1

(Red) in *hth* knockdown (*lGMR>*UAS-*hth^RNAi*) ocelli shows loss of Rh2 expression and

gain of Rh1 expression. (C, D) Antibody staining of anti-β-Galactosidase (β-Gal) in the

control (*rh2-lacZ; lGMR/+*) and in *hth* knockdown (*rh2-lacZ; lGMR>hth^RNAi*) ocelli. Anti-

βGal (green) expression is seen in control ocelli, also marked by anti-Rh2 (blue) (C).

However, anti-βGal (green) expression is absent in *hth* knockdown ocelli, marked by

anti-Rh1 (red) (D). (E, F) Antibody staining of βGal in the control (*rh1-lacZ; lGMR/+*)

and in *hth* knockdown (*rh1-lacZ; lGMR>hth^RNAi*) ocelli. In the control ocelli marked by

anti-Rh2 (blue), anti-βGal (green) expression is absent (E) whereas it is ectopically

expressed in case of *hth* knockdown ocelli marked by anti-Rh1 (red) (F).

Figure 3: Expression of *rh1* and *rh2* reporter constructs in ocelli. (A) Schematic

representation of the *rh2* minimal promoter used for making GFP reporter constructs

ranging from positions -293 to +55 of the transcription start (+1, arrow). The wildtype

promoter (*rh2-GFP*; in the top) contains three potential Hth binding sites (green) with

their corresponding sequences shown. In the *hth* mutant promoter (*rh2(hth mut)-GFP*;

in the bottom), the mutated sequences are shown with altered residues depicted in

red. (B, C) Expression of the wildtype *rh2-GFP* and mutant *rh2(hth mut)-GFP* reporter

lines (green) in ocelli stained with antibodies against GFP (green), Rh2 (red) and Rh1

(blue). Wildtype *rh2-GFP* (marked by anti-GFP) is expressed in PRs of ocelli (marked

by anti-Rh2) staining the entire rhabdomeres (B) whereas the mutant *rh2(hth-mut)-*

GFP (marked by anti-GFP) is not expressed in ocelli (C). (D) Schematic representation

of the *rh1* minimal promoter used for making the GFP reporter constructs ranging from

positions -247 to +73 of the transcription start (+1, arrow). The wildtype promoter (top)

contains two potential Hth binding sites (green; with its corresponding sequence

shown). In the *hth* mutant promoter (bottom), the mutated sequences are shown with

the altered residues depicted in red. (E, F) Expression of the wildtype *rh1-GFP* and

mutant *rh1(hth mut)-GFP* reporter lines in ocelli stained with antibodies against GFP

(green) Rh2 (red) and Rh1 (blue). (E) Like Rh1, the *rh1-GFP* reporter is not expressed

in the PRs of ocelli expressing only Rh2. (F) The mutant *rh1(hth-mut)-GFP* reporter

line is expressed in PRs of ocelli.

Figure 4: Hth misexpression phenotype in the retina. (A) Antibody staining to show

expression of Rh2 and Rh1 in the wildtype retina. Rh2 (green) is normally not

expressed in the retina (A') whereas Rh1 (blue) is expressed in the outer PRs of retina

(A''). PRs of the retina were stained with phalloidin (red). (B) Expression of Rh1 and

Rh2 in the *hth* overexpressing retina. Rh2 expression (green) can be seen now in the

outer PRs of retina (B') whereas Rh1 expression (blue) remained unaffected (B'').

Figure 5: Hth overexpression clones in the retina. (A, B, C, D) Antibody staining to

mark *hth* overexpression in clones (green) in the retina generated by the flip-out

technique. (A, A') Antibody staining against GFP to identify inner PR clones of the

retina, also co-stained by Rh2 (blue) and Phalloidin (red). Rh2 expression was not

induced (A") when clones were generated in the inner PRs. (B, B') Antibody staining

against GFP to identify outer PR clones of the retina, also co-stained by Rh2 (blue)

and Phalloidin (red). When clones were generated in the outer PRs, Rh2 expression

was induced (B"). White arrows marked inner and outer PR clones (A, A', B, B') and

their corresponding Rh2 expression. (C, C', D, D') Antibody staining against GFP to

identify inner and outer PR clones of the retina, also co-stained by Rh1 (blue) and

Phalloidin (red). When clones were either induced in the inner PRs (C, C') or outer

PRs (D, D') of the retina, Rh1 expression is unaffected (C", D").

Figure 6: Model of Rh2 cell fate in the ocelli. Schematic representation showing

origin of ocelli and compound eye from an ancestral eye in insects around 500 million

years ago. We showed that Rh2 is normally expressed in ocelli and gets independently

regulated by Hazy and Hth. Hth regulates a binary rhodopsin switch to promote Rh2

expression in ocelli while repressing Rh1. Overexpression of *hth* is sufficient to

promote Rh2 expression but only at the outer-PRs of retina.

Supplemental figure 1: Independent roles of Hazy and Hth to regulate Rh2

expression in ocelli. (A) Overexpression of *rh1* in all PRs of ocelli by using pan-

photoreceptor *lGMR*-Gal4. Antibody staining against Rh2 (green) and Rh1 (red) were

performed to show that overexpression of *rh1* does not inhibit Rh2 expression in ocelli.

(B) Antibody staining to show expression of Rh2 (green) and Rh1 (red) in *hazy^{-/-}* null

mutant ocelli. Rh2 expression is lost is absent in *hazy^{-/-}* mutants but they don't show

ectopic Rh1 expression in ocelli. (C) Antibody staining to show Hth expression (blue)

in *hazy⁻ᐟ⁻* mutant ocelli, also marked by anti-Chp (green). Hth expression is not affected

in ocellar PRs in *hazy⁻ᐟ⁻* mutants. (D) Antibody staining to show Hazy expression (blue)

in *hth* knockdown ocelli. Ocelli in *hth* knockdown is marked by staining against Rh1

(red). Hazy is expressed during rhodopsin switch in *hth* knockdown ocelli.

Supplemental figure 2: Knockdown of Hth does not alter ocellar PR cell fate. (A,

B) Antibody staining against Svp (green) in the wildtype and in *hth* knockdown ocelli.

Svp is normally expressed in R3/R4 and R1/R6 pairs of the developing retinal PRs

[36] and not in wildtype ocelli (A). During rhodopsin switch by *hth* knockdown in ocelli,

Svp is still not expressed in ocellar PRs (B). (C, D) Antibody staining against BarH1

(green) in the wildtype and in *hth* knockdown ocelli. BarH1 is normally expressed in

R1/R6 pair of the developing retina [35] and not in wildtype ocelli (C). Upon rhodopsin

switch by *hth* knockdown, BarH1 is still not expressed in the ocellar PRs (D). Ocellar

PRs are marked by antibody staining against Elav (blue, in A and B) and Chp (red, in

C and D).

References:

1.   von Salvini-Plawen L, Mayr E. On the Evolution of Photorecptors and Eyes:
Plenum Press; 1977.
2.   Quiring R, Walldorf U, Kloter U, Gehring WJ. Homology of the eyeless gene of
Drosophila to the Small eye gene in mice and Aniridia in humans. Science.
1994;265(5173):785-9. Epub 1994/08/05. doi: 10.1126/science.7914031. PubMed
PMID: 7914031.
3.   Gehring WJ. The genetic control of eye development and its implications for
the evolution of the various eye-types. Int J Dev Biol. 2002;46(1):65-73. Epub
2002/03/21. PubMed PMID: 11902689.
4.   Kumar JP. Signalling pathways in Drosophila and vertebrate retinal
development. Nat Rev Genet. 2001;2(11):846-57. Epub 2001/11/21. doi:
10.1038/35098564. PubMed PMID: 11715040.
5.   Zuker CS. On the evolution of eyes: would you like it simple or compound?
Science. 1994;265(5173):742-3. Epub 1994/08/05. doi: 10.1126/science.8047881.
PubMed PMID: 8047881.

6.   Wolff TaR, D.F. Pattern formation in the Drosophila retina. In: The Development
of Drosophila melanogaster Cold Spring Harbor Laboratory Press. 1993;2:1277-325.
7.   Yoon CS, Hirosawa K, Suzuki E. Studies on the structure of ocellar
photoreceptor cells of Drosophila melanogaster with special reference to
subrhabdomeric cisternae. Cell Tissue Res. 1996;284(1):77-85. Epub 1996/04/01.
doi: 10.1007/s004410050568. PubMed PMID: 8601298.
8.   Stark WS, Sapp R, Carlson SD. Ultrastructure of the ocellar visual system in
normal and mutant Drosophila melanogaster. J Neurogenet. 1989;5(2):127-53. Epub
1989/05/01. doi: 10.3109/01677068909066203. PubMed PMID: 2500507.
9.   Friedrich M. Continuity versus split and reconstitution: exploring the molecular
developmental corollaries of insect eye primordium evolution. Dev Biol.
2006;299(2):310-29. Epub 2006/09/16. doi: 10.1016/j.ydbio.2006.08.027. PubMed
PMID: 16973149.
10.   Waloszek D. The 'Orsten' window — a three-dimensionally preserved Upper
Cambrian meiofauna and its contribution to our understanding of the evolution of
Arthropoda. Paleontological Research. 2003;7(1):71-88, 18.
11.   Henze MJ, Dannenhauer K, Kohler M, Labhart T, Gesemann M. Opsin
evolution and expression in arthropod compound eyes and ocelli: insights from the
cricket Gryllus bimaculatus. BMC Evol Biol. 2012;12:163. Epub 2012/09/01. doi:
10.1186/1471-2148-12-163. PubMed PMID: 22935102; PubMed Central PMCID:
PMCPMC3502269.
12.   Friedrich M. Opsins and cell fate in the Drosophila Bolwig organ: tricky lessons
in homology inference. Bioessays. 2008;30(10):980-93. Epub 2008/09/19. doi:
10.1002/bies.20803. PubMed PMID: 18800378.
13.   Chou W-H, Hall KJ, Wilson DB, Wideman CL, Townson SM, Chadwell LV, et
al. Identification of a novel Drosophila opsin reveals specific patterning of the R7 and
R8 photoreceptor cells. Neuron. 1996;17(6):1101-15.
14.   Huber A, Schulz S, Bentrop J, Groell C, Wolfrum U, Paulsen R. Molecular
cloning of Drosophila Rh6 rhodopsin: the visual pigment of a subset of R8
photoreceptor cells. FEBS Lett. 1997;406(1-2):6-10. Epub 1997/04/07. doi:
10.1016/s0014-5793(97)00210-x. PubMed PMID: 9109375.
15.   Papatsenko D, Sheng G, Desplan C. A new rhodopsin in R8 photoreceptors of
Drosophila: evidence for coordinate expression with Rh3 in R7 cells. Development.
1997;124(9):1665-73. Epub 1997/05/01. PubMed PMID: 9165115.
16.   Hardie RC. Functional organization of the fly retina. Progress in sensory
physiology: Springer; 1985. p. 1-79.
17.   O'Tousa JE, Baehr W, Martin RL, Hirsh J, Pak WL, Applebury ML. The
Drosophila ninaE gene encodes an opsin. Cell. 1985;40(4):839-50. Epub 1985/04/01.
doi: 10.1016/0092-8674(85)90343-5. PubMed PMID: 2985266.
18.   Zuker CS, Cowman AF, Rubin GM. Isolation and structure of a rhodopsin gene
from D. melanogaster. Cell. 1985;40(4):851-8. Epub 1985/04/01. doi: 10.1016/0092-
8674(85)90344-7. PubMed PMID: 2580638.
19.   Feiler R, Harris WA, Kirschfeld K, Wehrhahn C, Zuker CS. Targeted
misexpression of a Drosophila opsin gene leads to altered visual function. Nature.
1988;333(6175):737-41. Epub 1988/06/23. doi: 10.1038/333737a0. PubMed PMID:
2455230.
20.   Mismer D, Michael WM, Laverty TR, Rubin GM. Analysis of the promoter of the
Rh2 opsin gene in Drosophila melanogaster. Genetics. 1988;120(1):173-80. Epub
1988/09/01. PubMed PMID: 2975615; PubMed Central PMCID: PMCPMC1203488.

21.   Pollock JA, Benzer S. Transcript localization of four opsin genes in the three
visual organs of Drosophila; RH2 is ocellus specific. Nature. 1988;333(6175):779-82.
Epub 1988/06/23. doi: 10.1038/333779a0. PubMed PMID: 2968518.
22.   Krapp HG. Ocelli. Curr Biol. 2009;19(11):R435-7. Epub 2009/06/12. doi:
10.1016/j.cub.2009.03.034. PubMed PMID: 19515345.
23.   Stange G, Stowe S, Chahl J, Massaro A. Anisotropic imaging in the dragonfly
median ocellus: a matched filter for horizon detection. Journal of Comparative
Physiology A. 2002;188(6):455-67.
24.   Bao R, Friedrich M. Molecular evolution of the Drosophila retinome: exceptional
gene gain in the higher Diptera. Mol Biol Evol. 2009;26(6):1273-87. Epub 2009/03/03.
doi: 10.1093/molbev/msp039. PubMed PMID: 19252076.
25.   Wernet MF, Labhart T, Baumann F, Mazzoni EO, Pichaud F, Desplan C.
Homothorax switches function of Drosophila photoreceptors from color to polarized
light sensors. Cell. 2003;115(3):267-79. Epub 2003/11/26. doi: 10.1016/s0092-
8674(03)00848-1. PubMed PMID: 14636555.
26.   Briscoe AD. Reconstructing the ancestral butterfly eye: focus on the opsins.
Journal of Experimental Biology. 2008;211(11):1805-13.
27.   Henze MJ, Oakley TH. The dynamic evolutionary history of pancrustacean
eyes and opsins. Integrative and comparative biology. 2015;55(5):830-42.
28.   Yuan F, Bernard GD, Le J, Briscoe AD. Contrasting modes of evolution of the
visual pigments in Heliconius butterflies. Molecular biology and evolution.
2010;27(10):2392-405.
29.   Gao S, Takemura SY, Ting CY, Huang S, Lu Z, Luan H, et al. The neural
substrate of spectral preference in Drosophila. Neuron. 2008;60(2):328-42. Epub
2008/10/30. doi: 10.1016/j.neuron.2008.08.010. PubMed PMID: 18957224; PubMed
Central PMCID: PMCPMC2665173.
30.   Yamaguchi S, Desplan C, Heisenberg M. Contribution of photoreceptor
subtypes to spectral wavelength preference in Drosophila. Proc Natl Acad Sci U S A.
2010;107(12):5634-9. Epub 2010/03/10. doi: 10.1073/pnas.0809398107. PubMed
PMID: 20212139; PubMed Central PMCID: PMCPMC2851746.
31.   Velarde RA, Sauer CD, Walden KK, Fahrbach SE, Robertson HM. Pteropsin:
a vertebrate-like non-visual opsin expressed in the honey bee brain. Insect Biochem
Mol Biol. 2005;35(12):1367-77. Epub 2005/11/18. doi: 10.1016/j.ibmb.2005.09.001.
PubMed PMID: 16291092.
32.   Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, Chevenet F, et al.
Phylogeny.fr: robust phylogenetic analysis for the non-specialist. Nucleic Acids Res.
2008;36(Web Server issue):W465-9. Epub 2008/04/22. doi: 10.1093/nar/gkn180.
PubMed PMID: 18424797; PubMed Central PMCID: PMCPMC2447785.
33.   Wernet MF, Desplan C. Homothorax and Extradenticle alter the transcription
factor network in Drosophila ommatidia at the dorsal rim of the retina. Development.
2014;141(4):918-28. Epub 2014/02/06. doi: 10.1242/dev.103127. PubMed PMID:
24496628; PubMed Central PMCID: PMCPMC3912834.
34.   Vasiliauskas D, Mazzoni EO, Sprecher SG, Brodetskiy K, Johnston RJ, Jr.,
Lidder P, et al. Feedback from rhodopsin controls rhodopsin exclusion in Drosophila
photoreceptors.   Nature.   2011;479(7371):108-12.   Epub   2011/10/11.   doi:
10.1038/nature10451.   PubMed   PMID: 21983964;   PubMed   Central   PMCID:
PMCPMC3208777.
35.   Higashijima S, Kojima T, Michiue T, Ishimaru S, Emori Y, Saigo K. Dual Bar
homeo box genes of Drosophila required in two photoreceptor cells, R1 and R6, and

primary pigment cells for normal eye development. Genes Dev. 1992;6(1):50-60. Epub 1992/01/01. doi: 10.1101/gad.6.1.50. PubMed PMID: 1346120.

36.   Mlodzik M, Hiromi Y, Weber U, Goodman CS, Rubin GM. The Drosophila seven-up gene, a member of the steroid receptor gene superfamily, controls photoreceptor cell fates. Cell. 1990;60(2):211-24. Epub 1990/01/26. doi: 10.1016/0092-8674(90)90737-y. PubMed PMID: 2105166.

37.   Mishra AK, Bargmann BOR, Tsachaki M, Fritsch C, Sprecher SG. Functional genomics identifies regulators of the phototransduction machinery in the Drosophila larval eye and adult ocelli. Dev Biol. 2016;410(2):164-77. Epub 2016/01/16. doi: 10.1016/j.ydbio.2015.12.026. PubMed PMID: 26769100.

38.   Mishra M, Oke A, Lebel C, McDonald EC, Plummer Z, Cook TA, et al. Pph13 and orthodenticle define a dual regulatory pathway for photoreceptor cell morphogenesis and function. Development. 2010;137(17):2895-904. Epub 2010/07/30. doi: 10.1242/dev.051722. PubMed PMID: 20667913; PubMed Central PMCID: PMCPMC2938920.

39.   Struhl G, Basler K. Organizing activity of wingless protein in Drosophila. Cell. 1993;72(4):527-40.

40.   Brockmann A, Domínguez-Cejudo MA, Amore G, Casares F. Regulation of ocellar specification and size by twin of eyeless and homothorax. Dev Dyn. 2011;240(1):75-85. Epub 2010/11/26. doi: 10.1002/dvdy.22494. PubMed PMID: 21104743.

41.   Hu KG, Reichert H, Stark WS. Electrophysiological characterization ofDrosophila ocelli. Journal of comparative physiology. 1978;126(1):15-24.

42.   Briscoe AD. Six opsins from the butterfly Papilio glaucus: molecular phylogenetic evidence for paralogous origins of red-sensitive visual pigments in insects. J Mol Evol. 2000;51(2):110-21. Epub 2000/08/19. doi: 10.1007/s002390010071. PubMed PMID: 10948267.

43.   Briscoe AD, Chittka L. The evolution of color vision in insects. Annu Rev Entomol. 2001;46:471-510. Epub 2000/12/09. doi: 10.1146/annurev.ento.46.1.471. PubMed PMID: 11112177.

44.   Kobayashi M, Fujioka M, Tolkunova EN, Deka D, Abu-Shaar M, Mann RS, et al. Engrailed cooperates with extradenticle and homothorax to repress target genes in Drosophila. Development. 2003;130(4):741-51. Epub 2002/12/31. doi: 10.1242/dev.00289. PubMed PMID: 12506004; PubMed Central PMCID: PMCPMC2692026.

45.   Inbal A, Halachmi N, Dibner C, Frank D, Salzberg A. Genetic evidence for the transcriptional-activating function of Homothorax during adult fly development. Development. 2001;128(18):3405-13. Epub 2001/09/22. PubMed PMID: 11566847.

46.   Zelhof AC, Koundakjian E, Scully AL, Hardy RW, Pounds L. Mutation of the photoreceptor specific homeodomain gene Pph13 results in defects in phototransduction and rhabdomere morphogenesis. Development. 2003;130(18):4383-92. Epub 2003/08/06. doi: 10.1242/dev.00651. PubMed PMID: 12900454.

47.   Bischof J, Maeda RK, Hediger M, Karch F, Basler K. An optimized transgenesis system for Drosophila using germ-line-specific phiC31 integrases. Proc Natl Acad Sci U S A. 2007;104(9):3312-7. Epub 2007/03/16. doi: 10.1073/pnas.0611511104. PubMed PMID: 17360644; PubMed Central PMCID: PMCPMC1805588.

48.   Hsiao HY, Johnston RJ, Jr., Jukam D, Vasiliauskas D, Desplan C, Rister J. Dissection and immunohistochemistry of larval, pupal and adult Drosophila retinas. J

Vis Exp. 2012;(69):4347. Epub 2012/11/28. doi: 10.3791/4347. PubMed PMID:
23183823; PubMed Central PMCID: PMCPMC3523422.
49. Tahayato A, Sonneville R, Pichaud F, Wernet MF, Papatsenko D, Beaufils P,
et al. Otd/Crx, a dual regulator for the specification of ommatidia subtypes in the
Drosophila retina. Dev Cell. 2003;5(3):391-402. Epub 2003/09/12. doi:
10.1016/s1534-5807(03)00239-9. PubMed PMID: 12967559.
50. Noro B, Culi J, McKay DJ, Zhang W, Mann RS. Distinct functions of
homeodomain-containing and homeodomain-less isoforms encoded by homothorax.
Genes Dev. 2006;20(12):1636-50. Epub 2006/06/17. doi: 10.1101/gad.1412606.
PubMed PMID: 16778079; PubMed Central PMCID: PMCPMC1482483.
51. Kanai MI, Okabe M, Hiromi Y. seven-up Controls switching of transcription
factors that specify temporal identities of Drosophila neuroblasts. Dev Cell.
2005;8(2):203-13. Epub 2005/02/05. doi: 10.1016/j.devcel.2004.12.014. PubMed
PMID: 15691762.
52. Charlton-Perkins M, Whitaker SL, Fei Y, Xie B, Li-Kroeger D, Gebelein B, et al.
Prospero and Pax2 combinatorially control neural cell fate decisions by modulating
Ras- and Notch-dependent signaling. Neural Dev. 2011;6:20-. doi: 10.1186/1749-
8104-6-20. PubMed PMID: 21539742.
690

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Supplemental figure 1

Supplemental figure 2