Intestine-enriched *apolipoprotein b* orthologs are required for stem cell differentiation and regeneration in planarians

Lily L. Wong[1],*, Christina G. Bruxvoort[1,2],*, Nicholas I. Cejda[1], Jannette Rodriguez Otero[3,4], and David J. Forsthoefel[1,5,#]

[1] Genes and Human Disease Research Program, Oklahoma Medical Research Foundation, Oklahoma City, Oklahoma

[2] Current address: Arthritis and Clinical Immunology Research Program, Oklahoma Medical Research Foundation; Department of Pathology, University of Oklahoma Health Sciences Center; and Department of Veteran Affairs Medical Center - Research Services, Oklahoma City, Oklahoma

[3] Howard Hughes Medical Institute, Department of Cell and Developmental Biology, University of Illinois at Urbana-Champaign, Urbana, Illinois

[4] Current address: Department of Education, Universidad Interamericana de Puerto Rico, San Juan, Puerto Rico

[5] Department of Cell Biology, University of Oklahoma Health Sciences Center, Oklahoma City, Oklahoma

* Authors contributed equally

[#] Correspondence:

David J. Forsthoefel, Ph.D.: david-forsthoefel@omrf.org

Abstract

Lipid metabolism plays an instructive role in regulating stem cell state and differentiation.

However, the roles of lipid mobilization and utilization in stem cell-driven regeneration are

unclear. Planarian flatworms readily restore missing tissue due to injury-induced activation of

pluripotent somatic stem cells called neoblasts. Here, we identify two intestine-enriched

orthologs of *apolipoprotein b, apob-1* and *apob-2,* which mediate transport of neutral lipid stores

from the intestine to target tissues including neoblasts, and are required for tissue homeostasis

and regeneration. Inhibition of *apob* function by RNAi causes head regression and lysis in

uninjured animals, and delays body axis re-establishment and regeneration of multiple organs in

amputated fragments. Furthermore, *apob* RNAi causes expansion of the population of

differentiating neoblast progeny and dysregulates expression of genes enriched in differentiating

and mature cells in eight major cell type lineages. We conclude that intestine-derived lipids

serve as a source of metabolites required for neoblast differentiation.

Introduction

Regeneration requires metabolites and energy for cell proliferation, differentiation, migration, and growth. Currently, specific mechanisms by which various metabolites are produced, transported, and utilized to promote regeneration are not well understood. In animals, neutral lipids (NLs) (e.g., triglycerides and cholesteryl esters) are stored in intracellular lipid droplets, which serve as central organelles for energy and lipid homeostasis[1,2]. Lipid stores can be mobilized either by NL packaging and export in lipoprotein particles (LPs), or by lipolysis and secretion as fatty acids (FAs)[3-5]. Mobilization enables transport to other tissues, for example between intestine, liver, and peripheral tissues such as brain and muscle in vertebrates, or between the fat body, intestine, nervous system, and imaginal discs in *Drosophila*[6-8]. Upon lipase-mediated hydrolysis of NLs, the FAs and cholesterol produced serve as building blocks for membrane biosynthesis, substrates for energy via fatty acid beta oxidation, sources of acetyl-CoA and other precursors for chromatin remodeling, and precursors of signaling molecules like eicosanoids and sphingolipids[9-11].

Considerable evidence indicates that disruption of lipid catabolism, biosynthesis, and transport can dysregulate stem cell pluripotency, proliferation, and differentiation[12-14]. For example, inhibiting fatty acid oxidation (FAO, the process of converting fatty acids to acetyl-CoA) causes symmetric production of differentiating progeny and stem cell depletion in the mammalian hematopoietic lineage[15]. By contrast, in the adult mouse hippocampus, inhibiting FAO promotes exit from quiescence and proliferation of neural stem/progenitor cells in the adult mouse hippocampus[16]. In *Drosophila*, compromising lipolysis and/or FAO causes intestinal stem cell necrosis[17], and loss of germline stem cells[18]. On the other hand, FAO-mediated production of acetyl-CoA (a precursor for histone acetylation) is required for differentiation, but not maintenance, of quiescent hematopoietic stem cells in *Drosophila* larvae, illustrating the growing recognition of the intersection of lipid metabolism and epigenetic regulation[19]. Perturbing lipid

synthesis and delivery also dysregulates stem cell state and differentiation dynamics. For

example, inhibiting fatty acid synthase (required for *de novo* lipogenesis) reduces proliferation

by adult mouse neural stem and progenitor cells[20]. Similarly, mutations in *hydroxysteroid (17-*

beta) dehydrogenase 7, a regulator of cholesterol biosynthesis, cause premature differentiation

of neural progenitors during development[21]. In culture, depriving human pluripotent stem cells of

extrinsic lipids promotes a "naive-to-primed" intermediate state, demonstrating the importance

of exogenous lipid availability for regulation of stem cell state[22].

Because stem and progenitor cell activation, proliferation, and differentiation are central

to regeneration, these diverse observations underline the potential importance of lipid

metabolism during regenerative growth. Currently, however, evidence that lipid transport and

utilization influence regeneration is limited. In zebrafish, *leptin b* (a hormonal regulator of

systemic lipid metabolism) is one of the most upregulated genes during fin and heart

regeneration[23], and *apolipoprotein E* (a regulator of NL transport in LPs) is upregulated during

fin regeneration[24]. In the axolotl *Ambystoma mexicanum*, genes involved in steroid, cholesterol,

and fatty acid metabolism are among the most upregulated at later stages of limb regeneration,

when differentiation of cartilage and muscle progenitors occurs[25]. In a primary cell culture model

of nerve injury, rat retinal ganglion cells regrow projections after axotomy more efficiently in the

presence of glial-derived LPs, reinforcing the role of cholesterol in axon regeneration[26]. In mice,

low density lipoprotein receptor deficiency delays liver regeneration and reduces hepatocyte

proliferation[27]. Similarly, elevated FA levels (induced by lipoprotein lipase overexpression)

causes lipotoxicity and compromises skeletal muscle regeneration[28], while inhibition of

peroxisomal FAO induces differentiation of myogenic satellite cells and muscle hypertrophy

during regeneration[29]. During skin wound repair, inhibiting triglyceride lipase-mediated lipolysis

by dermal adipocytes compromises recruitment of inflammatory macrophages and adipocyte

fate-switching to extracellular-matrix-secreting myofibroblasts[30].

Although these intriguing observations point to the importance of lipid metabolism, there

is scant functional understanding of how lipid transport and utilization influence stem cell

regulation during regeneration, particularly in emerging animal models with extensive

regenerative capacity. Planarians are freshwater flatworms capable of whole-body regeneration,

an ability conferred by pluripotent somatic stem cells called neoblasts that divide and

differentiate to replace damaged and lost tissues after amputation[31,32]. Diet-derived NLs are

stored in the planarian intestine in lipid droplets, suggesting the intestine is a major lipid storage

organ, as in *Drosophila* and *C. elegans*[6,33-35]. Planarian intestinal lipids have been proposed to

be a source of metabolites during extended fasting, as well as regeneration[34,36,37]. However,

mechanisms by which lipid secretion is controlled, and whether delivery to neoblasts or their

progeny is functionally required for regeneration, have not been investigated.

In this study, we show that two intestine-enriched *apolipoprotein b* (*apob*) orthologs are

required for NL transport from the intestine to neoblasts and their progeny, and that ApoB

function is required for stem cell differentiation and regeneration in the planarian *Schmidtea*

mediterranea. In mammals and insects, ApoB and ApoB-like proteins mediate trafficking of NLs

in LPs from digestive and storage organs to peripheral target tissues[5,38]. Here, we identify adult

stem cells and their differentiating progeny as additional target tissues for ApoB-mediated NL

transport in a regeneration competent animal, and propose utilizing planarians as models to

understand the influence of lipid metabolism on stem cells during regeneration.

Results

Planarian *apolipoprotein b* orthologs are expressed by intestinal cells

Previously, we demonstrated that knockdown of an intestine-enriched transcription factor, *nkx2.2*, inhibited neoblast proliferation and formation of the regeneration blastema, the unpigmented mass of tissue produced after amputation[39]. These observations suggested that the intestine could play a non-autonomous role in regulating stem cell dynamics. In an effort to identify intestine-enriched transcripts encoding neoblast regulators, we generated transcriptomes from control and *nkx2.2(RNAi)* planarians, in which neoblast proliferation and regenerative tissue production are severely inhibited (Supplementary Fig. 1a-c)[39]. An ortholog of human *apolipoprotein b* ("*apob-1*") encoding conserved Vitellogenin, DUF1943, and von Willebrand Factor D domains, was the second-most-downregulated transcript by significance and fold-change, while a paralog, *apob-2,* was also significantly downregulated (Fig. 1a; Supplementary Fig. 1d, e; Supplementary Data 1a-c; and Supplementary Data 2a-b). Consistent with single cell expression profiling[40] (Supplementary Fig. 1f), both transcripts were highly enriched in intestinal phagocytes, absorptive cells responsible for digestion, nutrient storage, and metabolite secretion (Fig. 1b and Supplementary Fig. 1f, g). In addition, both transcripts were weakly expressed in a small number of cells outside the intestine likely to be differentiating neoblast progeny (Fig. 1b and Supplementary Fig. 1f).

ApoB orthologs are required for viability and regulate neutral lipid transport

To determine whether ApoB orthologs were required for homeostatic tissue renewal and/or regeneration, we knocked down *apob-1* and *apob-2* using RNA interference[41]. In uninjured planarians, knockdown of either *apob-1* or *apob-2* individually had no phenotype (not shown), suggesting functional redundancy. However, after 3-5 double-stranded RNA (dsRNA) feedings, double knockdown *apob(RNAi)* animals displayed phenotypes that progressed from

mild (modest/regional pigmentation loss), to severe (animal-wide pigmentation loss and reduced

motility), to very severe (head regression, ventral curling, and eventual lysis) (Fig. 2a). Head

regression and ventral curling phenotypes are common in planarians lacking functional

neoblasts[42], and suggested that ApoB orthologs could have functional roles in the regulation of

planarian stem cells.

ApoB orthologs in vertebrates and insects regulate LP biogenesis and secretion[5,43]. To

test whether ApoB protein was secreted, we made a custom antibody recognizing the N-

terminus of Smed-ApoB-1 (Fig. 2b, c). ApoB-1 protein was enriched in the intestine, but also

throughout the parenchyma (tissues surrounding other organs, sandwiched between the

epidermis and intestine, where neoblasts also reside). This indicated that although *apob*

mRNAs were intestine-enriched (Fig. 1a, b), ApoB protein was robustly secreted and

transported to peripheral tissues (Fig. 2c). Expression was dramatically reduced in both regions

in *apob-1;apob-2(RNAi)* double knockdown planarians (*"apob(RNAi)"* hereafter for brevity),

demonstrating that knockdown effectively reduced ApoB protein levels (Fig. 2c).

ApoB orthologs facilitate NL secretion and transport via LPs[5,38], while ApoB binding by

receptors mediate NL uptake and metabolism by target cells[44]. To test whether planarian ApoB

functioned similarly, we first evaluated NL distribution in *apob(RNAi)* animals. In histological

sections, NLs (labeled with Oil Red O) were elevated in both the intestine as well as tissues

surrounding the intestine, suggesting that both LP secretion by the intestine and

uptake/metabolism by peripheral tissues were compromised (Fig. 2d). Using thin layer

chromatography, we also found significant elevation of cholesteryl esters and triglycerides in

lipid extracts from *apob(RNAi)* animals (Fig. 2e). These phenotypes predicted that delivery of

NLs to neoblasts and/or their progeny would be compromised by ApoB inhibition. To test this,

we quantified NL content in neoblasts and their progeny using a fluorescent NL probe, BODIPY-

493/503, by flow cytometry. Three major subpopulations of planarian cells can be distinguished

by their DNA content and sensitivity to X-irradiation[45,46] (Fig. 2f-h). The "X1" fraction/gate

includes >2C DNA content neoblasts in S/G2/M phase of the cell cycle, while "X2" includes 2C

neoblasts in G0/G1 and G0 post-mitotic progeny. "Xins," named for the fact that cells in this

gate are insensitive to X-irradiation, consists of later stage progeny and mature differentiated

cells. We found a dramatic reduction of fluorescence in both the X1 and X2 fractions in

apob(RNAi) animals vs. controls (Fig. 2i, j), indicating that a reduction of NL content in neoblasts

and their progeny was caused by ApoB inhibition. Together, these data demonstrated that

planarian ApoB proteins were produced by the intestine and were likely secreted as LPs to

transport NLs from the intestine to stem cells and their differentiating progeny.

apob and *lipoprotein receptor* genes are upregulated in regenerating fragments

Amputation induces changes in gene expression during planarian regeneration[47-49]. We

predicted that if *apob* and other genes involved in neutral lipid metabolism played functional

roles during regeneration, they would be up- or down-regulated in amputated tissue fragments.

Using quantitative PCR, we found that both *apob-1* and *apob-2* transcripts were upregulated in

tissue fragments at two distinct time points during earlier stages of regeneration commonly

associated with neoblast proliferation (1-3 days) and differentiation (4-5 days) (Supplementary

Fig. 2a, b). Upregulation of *apob-1* and *apob-2* was also observed in previously published RNA-

Seq data from a 14-day time course of whole-body planarian regeneration (Supplementary Fig.

2c)[49]. Using quantitative capillary-based Western blotting, we also found that ApoB-1 protein

levels increased by 3-5 days after amputation (Supplementary Fig. 2d). Consistently,

accumulation of neutral lipids was sustained in 3- and 6-day *apob(RNAi)* regenerates

(Supplementary Fig. 2e, f), suggesting that LP trafficking from the intestine was disrupted, as in

uninjured animals (Fig. 2d).

In addition, we found that three planarian orthologs of human lipoprotein receptors

(which bind apolipoproteins, enabling LP uptake/metabolism) were upregulated in the blastema,

and regenerating brain and pharynx (Supplementary Fig. 3a, b, d). These transcripts were

expressed in both *piwi-1*-mRNA-positive neoblasts as well as *piwi-1*-negative cells that were

likely to be differentiating neoblast progeny in the regenerating pharynx and blastema

(Supplementary Fig. 3d). Two lipoprotein receptor genes (*ldlr-1* and *ldlr-2*) were also

differentially expressed in the whole-body regeneration RNA-Seq dataset (Supplementary Fig.

3c)[49]. Using the Gene Ontology to mine the same dataset[49], we identified additional planarian

genes predicted to regulate lipid transport, triglyceride, and cholesterol metabolism, and found

that many of these transcripts were also up- and down-regulated during regeneration

(Supplementary Fig. 3e and Supplementary Data 3). Together, these observations indicate that

apob-1, apob-2, and other genes involved in neutral lipid metabolism were differentially

expressed in response to amputation, consistent with roles during regeneration.

apob paralogs are required for polarity re-establishment and organogenesis during

regeneration

To assess functional roles of *apob* paralogs during regeneration (Fig. 3a), we chose

uninjured animals after three dsRNA feedings that exhibited "mild" and "severe" phenotypes

(Fig. 2a), amputated their heads, and assessed blastema morphogenesis, which absolutely

requires neoblast proliferation and differentiation[50-52]. Controls, *apob-1(RNAi)*-only, and *apob-*

2(RNAi)-only fragments regenerated normally (Fig. 3b), again suggesting *apob* paralogs likely

functioned redundantly. However, *apob(RNAi)* double knockdown animals had reduced

posterior blastemas (head fragments) and anterior blastemas (trunk fragments) at 8 days post-

amputation (Fig. 3b). Blastema size was noticeably smaller in fragments from "severe" ("*apob-*

S") animals compared to "mild" ("*apob-M"*) animals (Fig. 3b), suggesting that neoblasts and/or

their progeny were more strongly affected over time.

Expression of so-called "position control genes" to re-establish axial polarity is an

essential early event required for whole-body regeneration[53]. We asked whether expression of

anterior (*notum*)[54] or posterior (*wnt11-2*)[55] was affected in *apob(RNAi)* regenerates. Indeed, at

three days post-amputation (3 dpa), the number of both posterior *wnt11-2*-expressing cells (Fig.

3c) and anterior *notum*-expressing cells (Fig. 3d) was reduced in *apob-M* and *apob-S*

fragments. By 6 dpa, although most *apob(RNAi)* fragments had regenerated more of these

cells, many fragments (especially *apob-S*) still had fewer than five cells compared to control

fragments (Fig. 3c, d). These data suggested that *apob* inhibition delayed, but did not

completely abolish, re-establishment of these cells.

apob RNAi also affected regeneration of planarian organs, including the brain, pharynx,

and intestine. Although both brain and pharynx regenerated, these organs were smaller than in

controls (Supplementary Fig. 2g, h). This suggesting regeneration was delayed, but not blocked,

similar to the phenotype for cells expressing polarity cues. Quantitatively, at 6 dpa, both brain

and pharynx were smaller in size, especially in *apob-S* fragments (Fig. 3e, f). To assess

intestine, we analyzed both head and tail regenerates, in which a combination of neoblast-

driven new cell production and remodeling of pre-existing, differentiated intestinal branches is

required for successful regeneration[56]. Newly regenerated posterior branches were significantly

shorter in *apob(RNAi)* head fragments (Fig. 3g). By contrast, in tail fragments, length of the new

anterior intestine was not significantly affected (Fig. 3h). Rather, these branches failed to fuse at

the anterior midline, leading to a "split" anterior branch phenotype in ~50% of *apob(RNAi)* tail

fragments (Fig. 3h). Together, these results suggested that ApoB reduction delayed

regeneration of multiple cell types and organs, and, in the case of the intestine, affected

differentiation of new cells as well as collective cell migration or other processes required for

remodeling.

apob knockdown causes accumulation of early neoblast progeny

Because *apob* knockdown disrupted multiple neoblast-driven processes during

regeneration, we asked whether defects in neoblast maintenance or proliferation were

responsible for the effects of *apob* RNAi on regeneration. First, we assessed neoblast numbers

by FISH using *piwi-1,* a pan-neoblast marker, and *tgs-1,* a marker for a more pluripotent

subpopulation (which also includes neural specialized neoblasts)[49,50,57]. In both uninjured

planarians and 7.5-day regenerates, neoblast numbers and distribution were grossly normal

(Supplementary Fig. 4a, b). We also assessed proliferation using anti-phospho-Histone H3

(Ser10) ("PS10") immunolabeling, which marks cells in late G2 and M phase of the cell

cycle[58,59]. In uninjured animals and 2-day head regenerates, the number of mitotic neoblasts

increased modestly in *apob-M,* but not *apob-S* samples (Fig. 4a, b). In 2-day trunk regenerates,

there was a more significant (~50%) increase in *apob-M* fragments, and a modest increase in

apob-S fragments (Fig. 4b). Together, these results suggested that ApoB reduction might cause

moderate hyperproliferation, or, alternatively, a modest mitotic delay, without dramatically

affecting *piwi-1+* or *tgs-1+* neoblast numbers. These mild phenotypes also raised the possibility

that ApoB reduction might preferentially dysregulate differentiation, rather than proliferation or

maintenance of actively cycling neoblasts.

We tested these possibilities quantitatively using flow cytometry. We dissociated

uninjured and regenerating planarians, and evaluated the proportions of cells in the X1 and X2

fractions (Fig. 4c). In uninjured planarians, as well as 2-day and 7-day regeneration fragments,

X1 neoblast numbers were unaffected by *apob* knockdown (Fig. 4d, e and Supplementary Fig.

4c), consistent with the moderate effect of *apob* RNAi on neoblast proliferation (Fig. 4a, b and

Supplementary Fig. 4a, b). By contrast, in uninjured planarians as well as 2- and 7-day trunk

fragments, the proportion of cells in the X2 gate increased significantly by ~20-40% (Fig. 4f and

Supplementary Fig. 4c) in *apob(RNAi)* animals vs. control. In 2 dpa head fragments, there was

also a modest, but statistically insignificant, increase (~15%) in the X2 fraction (Fig. 4f).

Because the X2 fraction includes both cycling G1-phase neoblasts and differentiating post-

mitotic progeny[60,61], these results suggested that *apob* inhibition might cause lengthening of G1

phase of the cell cycle, and/or a delay in differentiation of neoblast progeny, either of which

could increase the proportion of cells in X2.

In order to distinguish between these possibilities, we examined the X2 fraction in

uninjured planarians 24 hr after X-irradiation, which preferentially ablates over 95% radiation-

sensitive cycling neoblasts, without affecting many early neoblast progeny[49,52,60]. As expected,

the X1 fraction was almost completely eliminated in both control and *apob(RNAi)* samples (Fig.

4g, h). However, although irradiation reduced the proportion of X2 cells by ~48-56% in both

control and *apob(RNAi)* animals (Fig. 4i), the expansion of the X2 fraction persisted in *apob-M*

(~62.5% increase) and *apob-S* animals (~42.5% increase) relative to controls (Fig. 4i). This

increase in radiation-insensitive cells in X2 strongly suggested that the primary defect in *apob*

knockdown animals was a delay in differentiation of neoblast progeny.

Intriguingly, the increase in radiation-resistant X2 cells was transient: by 48 hr post-

irradiation, the proportion of X2 cells in *apob(RNAi)* samples was only modestly (but not

significantly) elevated (Fig. 4i). This suggested that differentiation was delayed, but not arrested,

in *apob(RNAi)* animals. One possible explanation for this observation is that, at later time points,

progeny had differentiated further, and resided in the Xins gate (rather than the X2 gate) as they

achieved a more mature cell state. Alternatively, some early progeny might have undergone

apoptosis or necrosis at later time points, reducing the apparent accumulation of neoblast

progeny in X2.

ApoB reduction preferentially dysregulates expression of transcripts enriched in

differentiating neoblast progeny and mature cell types

Our flow cytometry data suggested that the primary phenotype in uninjured and

regenerating *apob(RNAi)* animals was a delay in the differentiation of neoblast progeny. To test

this interpretation further, we performed whole animal RNA sequencing on control, *apob-M,* and

apob-S animals, and identified differentially expressed (DE) genes in the two *apob(RNAi)*

groups (Supplementary Fig. 5a-c and Supplementary Data 4). Then, to identify specific

biological processes affected by *apob* inhibition, we used the Gene Ontology (GO) to identify

patterns of dysregulation in specific functional categories (Supplementary Fig. 5d, e and

Supplementary Data 5). In addition, to determine whether *apob* knockdown disproportionately

dysregulated genes enriched in differentiating progeny states, we mapped the global *apob*

"dysregulation signature" to published bulk- and single-cell transcriptomes to determine which

cell types and states were most affected by *apob* inhibition.

apob knockdown dysregulated thousands of genes, causing upregulation of 842 (*apob*-

M) and 1960 (*apob-S*) transcripts, and downregulation of 1139 (*apob*-M) and 2547 *(apob*-S)

transcripts, relative to control (Supplementary Fig. 5a-c and Supplementary Data 4a-d). For

upregulated transcripts, "lipid metabolism" was the fifth-most over-represented Biological

Process (BP) term in *apob-M* animals, and was the most over-represented category in *apob-S*

animals (Supplementary Fig. 5d and Supplementary Data 5a, b). Enrichment of the

subcategories of acylglycerol, fatty acid, steroid, and glycerolipid metabolism was consistent

with known roles of ApoB orthologs, and suggested a possible compensatory gene expression

response to *apob(RNAi)*-induced disruption of NL transport. *apob(RNAi)* also downregulated

transcripts in additional metabolism categories, including gluconeogenesis, glycolysis, pyruvate,

and nucleotide metabolism (e.g., NADH and ADP), as well as ion transport, indicating wide-

ranging dysregulation of metabolite processing and trafficking, especially in *apob-S* animals

(Supplementary Fig. 5e and Supplementary Data 5c, d). Interestingly, *apob* inhibition

dysregulated many additional, non-metabolism-related transcripts. Upregulated functional

categories included differentiation and/or specification of multiple tissues including epidermis,

nervous system, and the eye, while downregulated categories included cilium morphogenesis

and function (possibly reflecting disruption of ciliated epidermal and/or protonephridial cell

numbers or physiology), muscle morphogenesis and function, and extracellular matrix

organization (Supplementary Fig. 5d, e and Supplementary Data 5a-d). Importantly, categories

related to cell cycle, mitosis, etc. were not enriched among up- or downregulated transcripts.

Together, these data suggested that ApoB reduction affected metabolism, cell/tissue

differentiation, and functions of mature cell types, but not processes required specifically for

cellular proliferation.

Next, to test whether *apob* disproportionately dysregulated genes expressed by

differentiating progeny and mature cells, we cross-referenced our DE transcript list with recently

published bulk transcriptomes from flow-sorted X1, X2, and Xins cell fractions (Fig. 4c)[49], as well

as cells sorted based on their expression of PIWI-1 protein, a widely used marker whose

expression marks states within planarian cell lineages (e.g., high PIWI-1 levels in neoblasts, low

PIWI-1 levels in some neoblasts and differentiating progeny, and negligible PIWI-1 levels in

mature cells)[49]. We first identified transcripts that were enriched in each cell fraction to generate

a "signature" transcript list for each fraction (Supplementary Fig. 6a, c). We then asked what

percent of these signature transcripts were dysregulated in *apob-M* and *apob-S* animals (Fig.

5a, b and Supplementary Fig. 6b, d). *apob* RNAi dysregulated signature transcripts in all six

fractions. However, compared to X1 or "PIWI-HI" signature transcripts, 2-4 times as many

transcripts were up- or down-regulated in X2 and "PIWI-LO" cell classes, which included both

neoblasts and early progeny (Fig. 5a, b). Relative to Xins and PIWI-NEG fractions, there were

also more dysregulated transcripts represented in the X2 and PIWI-LO fractions in *apob* RNAi

animals (Fig. 5a, b). To demonstrate the validity of this approach, we cross-referenced

transcriptomes from planarians 24 hours after irradiation[49], and found that dysregulated

transcripts overlapped mainly with X1 and PIWI-HI signature transcripts, and were primarily

downregulated, consistent with neoblast loss (Supplementary Fig. 6e, f). Together, this analysis

demonstrated that *apob* knockdown preferentially affected transcripts enriched in differentiating

progeny, an observation that is consistent with accumulation of cells in this state in flow

cytometry experiments (Fig. 4d-i).

To determine whether *apob* RNAi preferentially affected specific lineages or states within

individual lineages, we also compared the gene expression signature of *apob(RNAi)* animals

with recently published single cell transcriptomes[40]. Specifically, we cross-referenced transcripts

dysregulated by *apob* RNAi with transcripts enriched in cell state subclusters in eight planarian

lineages (Fig. 5c-i, Supplementary Fig. 8a, and Supplementary Fig. 8c-l). The resulting

"dysregulation signature" for each lineage showed how *apob* RNAi affected gene expression in

specific cell states during the progression from pluripotent cycling neoblasts ("N"), to early

transition states ("TS"), to differentiating progeny ("P") to mature cell states ("M") (see example

schematic for epidermal lineage in Fig. 5c). We observed a striking and consistent dysregulation

pattern in all eight lineages. In three well characterized lineages (epidermis, intestine, and

protonephridia) (Fig. 5c-i), the percentage of dysregulated "state-enriched" transcripts was

lowest for neoblast and early transition state subclusters, but higher for progeny and mature cell

states. Similar trends were observed for less characterized lineages such as muscle, pharynx,

and *cathepsin*-positive cells (Supplementary Fig. 8c-l). Validating this approach, irradiation[49]

primarily dysregulated neoblast and transition state transcripts (Supplementary Fig. 7a-g), and,

to a lesser degree for some lineages, progeny-enriched transcripts (Supplementary Fig. 8a and

Supplementary Fig. 8m-q). Similarly, and as expected, X1- and PIWI-HI-enriched transcripts

overlapped with neoblast and transition state transcripts, while X2-/PIWI-LO and Xins/PIWI-

NEG transcripts were enriched in progressively later cell state subclusters in each lineage

(Supplementary Fig. 7h-k, Supplementary Fig. 8b, and Supplementary Fig. 8r-v). Together,

these results provided further evidence that *apob* inhibition specifically dysregulated gene

expression in progeny and mature cell types/states in all eight lineages, and lent additional

support to the conclusion that ApoB was required for differentiation of planarian neoblasts.

Discussion

In this study, we discover a non-cell autonomous role for ApoB proteins in regulation of planarian neoblast differentiation. Our data suggest that ApoB transports NLs from the intestine to both neoblasts and their differentiating progeny via LPs (Fig. 6). In the absence of ApoB, neoblast proliferation and maintenance are largely unaffected, but differentiation is delayed, causing accumulation of differentiating progeny (Fig. 6) and slowing regeneration.

apob knockdown causes a ~40% increase in a flow cytometry fraction of 2C DNA content cells (X2) that includes both cycling neoblasts in G1, and post-mitotic differentiating progeny in G0[57,60]. This increase occurs without simultaneously affecting the >2C S/G2/M cell fraction (X1), and is sustained even in recently irradiated animals, indicating that the expansion is comprised primarily of G0 progeny, and not G1 neoblasts. We suggest that this phenotype indicates a disruption in the progression of differentiation after cell fate specification, and not specification itself. In planarians, specification likely begins during S-phase, because expression of fate-specific transcription factors is significantly higher in S/G2/M neoblasts than in G1 neoblasts[57,62]. Intriguingly, inhibition of other planarian genes required for differentiation (e.g., the transcription factor *mex3-1,* the extracellular matrix component *collagen4-1,* and the transcriptional co-activating protein *cbp-3*) also cause increases in neoblast numbers *in vivo*[52,63-65]. Furthermore, knockdown of *exocyst component 3 (exoc3),* a negative regulator of pluripotency whose mammalian homolog *Tnfaip3* promotes embryonic stem cell differentiation, causes expansion of the S/G2/M (X1) neoblast fraction[66]. We speculate that these genes may be required for cell fate specification, and that their inhibition shifts neoblast dynamics in favor of renewal divisions that expand the stem cell compartment. By contrast, *apob* RNAi does not cause accumulation of S/G2/M neoblasts, but rather increases post-mitotic progeny number, and causes greater dysregulation of transcripts associated with differentiation and mature cell states. Therefore, ApoB is likely required to drive post-specification stages of differentiation, and

inhibition delays late commitment steps (i.e., after cell cycle exit), and/or transitions to final

mature states (Fig. 6).

Differentiation requires extensive changes in gene expression that are often preceded by

genome-wide chromatin remodeling[67,68]. Intriguingly, FA oxidation is a significant source of

carbon for acetyl-CoA production and histone acetylation[69]. Consistent with a role for NLs in

planarian differentiation, knockdown of *exoc3* reduces triglyceride levels, causes expansion of

the neoblast population, inhibits organogenesis, and reduces expression of differentiation

markers; palmitic acid supplementation rescues these differentiation-associated phenotypes[66].

Furthermore, in the sexually reproducing *S. mediterranea* planarian biotype, inhibition of *nuclear*

hormone receptor 1 causes NL accumulation and blocks differentiation of gonads and

accessory reproductive tissues, a phenotype that is rescued by supplementation with either

acetyl-CoA or Acyl-CoA synthetase[70]. Because acetyl-CoA can also enter the citric acid cycle to

produce alpha-ketoglutarate, a substrate for histone demethylation[71], ApoB inhibition could

dysregulate epigenetic changes through multiple pathways. Histone acetylases, deacetylases,

methyltransferases, and demethylases are conserved in planarians, and their inhibition disrupts

stem cell maintenance, differentiation, and regeneration[64,65,72-75]. Because *apob* RNAi results in

widespread dysregulation of thousands of transcripts associated with differentiating progeny, it

is reasonable to suggest that in planarians, intestinal lipid stores serve as a ready carbon

source that is trafficked by ApoB-containing LPs to neoblasts and progeny to support epigenetic

modifications required for differentiation.

ApoB depletion may also delay differentiation by other mechanisms. For example, NL-

derived fatty acids may be utilized to produce ATP via beta-oxidation and the mitochondrial

electron transport chain (ETC) to support energy-dependent processes during differentiation.

Consistent with this idea, planarian mitochondrial mass is higher in differentiating progeny, and

pharmacological inhibition of the ETC promotes pluripotency and neoblast colony expansion,

which may also limit differentiation[76]. In mammals, disrupting the ETC blocks differentiation of

cardiomyocytes and mesenchymal stem cells[77,78], but the ETC is dispensable for differentiation

of mammalian epidermal progenitor cells and *Drosophila* ovarian stem cells[79,80]. Further study

will be needed to determine whether LP-transported NLs serve as a significant energy source

during planarian differentiation. Additionally, LP-mediated transport of morphogens like

Hedgehog or Wnt proteins[81], whose planarian orthologs play important roles in regulating axial

polarity and tissue differentiation[53], may be affected by *apob* knockdown. However, we find that

ApoB inhibition delays, but does not block or alter regeneration of axial polarity, and we find little

evidence of dysregulation of polarity-related transcripts (Supplementary Data 4 and 5),

suggesting that planarian LPs may not play a major role in planarian morphogen trafficking.

Similarly, fat-soluble vitamins known to influence stem cell dynamics are also transported in

LPs[82-87]. Although characterization of LP cargo may yield additional insights, the dramatic

dysregulation of lipid metabolism at the gene expression level, and the minimal disruption of

vitamin-related gene expression in *apob(RNAi)* animals (Supplementary Data 4 and 5), suggest

that LP-mediated vitamin transport may not play a significant role in planarian differentiation.

Lastly, we find that *apob* inhibition dysregulated genes associated with muscle differentiation

and function (Supplementary Data 5). In planarians, muscle cells not only secrete axial polarity

cues, but also serve a fibroblast-like role by secreting most components of the extracellular

matrix, whose functions are required to both spatially restrict the stem cell compartment, and

modulate proliferation and differentiation[63,88,89]. *apob* RNAi causes moderate downregulation of

most fibrillar collagens, as well as the basement membrane *collagen4-1,* which promotes

differentiation[63] (Supplementary Data 4 and 5). Thus, ApoB depletion may also delay

differentiation indirectly, by compromising the generation and/or function of muscle cells.

The effect of *apoB* knockdown on regeneration suggests several future directions. First,

although the existence of prominent LDs in planarian neoblasts has been known for decades[90],

the roles of this intriguing organelle have not been investigated. We did not assess neoblast LD

numbers or size, but *apoB* knockdown dramatically reduces NL content in both X1 (S/G2/M)

and X2 (G0/G1) neoblast fractions, suggesting that ApoB and LPs may influence neoblast LD

content and/or function. In addition, NL content is lower in G0/G1 cells, suggesting that a

primary role of LDs may be to support differentiation. Once thought of as static storage particles,

recent work has demonstrated that LDs in animal cells are dynamic, multi-functional organelles

that regulate nutrient sensing, cell stress responses, and even intracellular localization of

histones, transcription factors, and other proteins[2,10]. Additionally, although emerging data

suggest LDs as a potential therapeutic target in cancer stem cells [91,92], our knowledge of the

regulation and functions of LDs in stem cells, especially during regeneration, is limited. Studies

in planarians and other regeneration models could further illuminate the roles of this organelle.

Second, coordinated metabolic shifts between glycolysis and oxidative phosphorylation may be

a widespread aspect of stem cell transitions between quiescence, proliferation, and

differentiation, and extrinsic lipids can influence these states[11-14]. Our RNA-Seq data suggest

that transcription of regulators of amino acid metabolism, glycolysis, the tricarboxylic acid cycle,

and other metabolic pathways respond to *apob(RNAi)*, suggesting that planarian stem cell

metabolism is just as dynamic as in other animals. The fact that *apob(RNAi)* seems to primarily

affect post-mitotic states also suggests that planarian neoblasts might rely primarily on

glycolysis for energy and metabolite supply, and shift to lipid metabolism and oxidative

phosphorylation during differentiation, as in other systems[93-95]. Again, studies in animals with

high regenerative capacity could generate greater insights into whether and how injury can

induce metabolic switching. Third, we find that expression of *apob-1* and *apob-2,* as well as

numerous other regulators of NL metabolism are dynamically up- and down-regulated at the

transcript level during regeneration (Supplementary Fig. 3E). This suggests that coordination of

lipid metabolism is part of a genome-encoded program of regenerative gene expression.

Unraveling which transcription factors, chromatin modifiers, and other regulators are responsible

for such regulation is thus a third priority for future study.

Finally, because *apob-1* and *apob-2* are downregulated by inhibition of *nkx2.2,* an

intestine-enriched transcription factor also required for regeneration[39], our results provide a

specific example of how the planarian intestine can non-autonomously influence neoblast

dynamics. Intriguingly, unlike *nkx2.2* RNAi, *apob* knockdown does not reduce the abundance of

phosphoHistone-H3-S10-positive neoblasts, indicating that the proliferative defect in

nkx2.2(RNAi) animals is not caused by *apob* reduction, and additional downstream regulators of

proliferation remain to be discovered. Intriguingly, dozens of additional regulators of lipid

metabolism and transport of other metabolites are downregulated in *nkx2.2(RNAi)* animals

(Supplementary Data 1b), suggesting additional ways in which the intestine could influence

neoblasts and their niche.

In summary, we have identified *apolipoprotein B* orthologs and neutral lipid metabolism

as important regulators of stem cell differentiation and regenerative tissue growth. Since the

discovery of lipoproteins a century ago, their roles in lipid transport and disease have been

extensively investigated[96], but functions in stem cell regulation are not nearly as well

characterized[97-100]. Efforts to define functions of LPs and their metabolic derivatives more

precisely in planarians and other models will therefore improve our understanding of metabolic

requirements of stem cell-driven regeneration. In addition, because lipid metabolism is

amenable to pharmacological manipulation[101], further study may provide new insights relevant

to the dual goals of promoting repair of damaged human tissues, and inhibiting growth in

pathological contexts like cancer.

Methods

Ethics Statement

Anti-ApoB-1 antibodies were generated by GenScript USA (Piscataway, NJ), an OLAW,

AAALAC, and PHS-approved vendor. GenScript's animal welfare protocols were approved by

OMRF IACUC (17-58). No other vertebrate organisms were used in this study.

Planarian care

Asexual *Schmidtea mediterranea* (clonal line CIW4)[102] were maintained in 0.5 g/L Instant

Ocean salts with 0.0167 g/L sodium bicarbonate dissolved in Type I water[103], and fed with beef

liver paste. Planarians were starved 7-10 days prior to initiating RNAi. Animals were 2-5 mm in

length for most experiments except flow cytometry, for which 5-10 mm animals were used.

Uninjured, intact animals were randomly selected from large (300-500 animal) pools.

Cloning and expressed sequence tags

Transcripts were cloned as previously described[104]. Sequences were identified in the

dd_Smed_v6 transcriptome[105] and the Smed_ESTs3 library[106]. These included *nkx2.2*

(dd_2716_0_1/PL08007A2A07),

 apob-1 (dd_636_0_1/PL06004B2E09), *apob-2* (dd_194_0_1/PL08004B1B10), *ldlr-1*

(dd_9829_0_1), *ldlr-2* (dd_5596_0_1/PL04021A1C10), *vldlr-1* (dd_1510_0_1/PL05007B1H03),

notum (dd_24180_0_1), *wnt11-2* (dd_16209_0_1), *choline acetyltransferase/ChAT*

(dd_6208_0_1), *laminin* (dd_8356_0_1/PL030015A20A02), *cathepsin La/ctsla*

(dd_2567_0_1/PL06020B2D09), *piwi-1* (dd_659_0_1/PL06008A2C06), *tgs-1* (dd_10988_0_1),

solute carrier family 22 member 6/slc22a6 (dd_1159_0_1), and Niemann-Pick type C-2/*npc2*

(dd_73_0_1/PL030001B20C07). *S. mediterranea ldlr-1, ldlr-2,* and *vldlr-1* were identified by

BLAST homology and named after their top human refseq_protein BLASTX hits. Sequences of

primers and expressed sequence tags are available in Supplementary Data 6.

Domain organization and phylogenetic analysis

For ApoB-1, ApoB-2, Ldlr-1, Ldlr-2, and Vldrl-1, protein domains were identified using

HMMSCAN (https://www.ebi.ac.uk/Tools/hmmer/search/hmmscan) to search Pfam, TIGRFAM,

and Superfamily databases, with Phobius for transmembrane and signal peptide predictions

(conditional E-value cutoff of 1e-03)[107]. For the ApoB phylogenetic tree, N-terminal Vitellogenin

domains from ApoB and related proteins were aligned in Geneious using MAAFT (default

settings), and alignment was manually trimmed to the N- and C-terminal boundaries of human

Apo B-100. Phylogenetic tree was generated using PhyML 3.0[108] (http://www.atgc-

montpellier.fr/phyml/), using AIC for automatic selection of the LG substitution model, BioNJ

starting tree, NNI for tree topology improvement, and 100 bootstrap replicates. Accession

numbers for proteins used in domain diagrams and phylogenetic analysis are included in

Supplementary Data 6.

In situ hybridization

Riboprobe synthesis, WISH, and FISH were conducted as previously described[104].

Riboprobes were used at 0.05-1 ng/μl. Cryosections were generated after FISH as in[109].

RNAi

dsRNA synthesis and RNAi experiments were conducted as described[41,104] by mixing

one μg of in vitro-synthesized dsRNA with 8-9 μl of 1:10 food coloring:water mix, and 40 μl of

2:1 liver:water homogenate. For *nkx2.2* RNAi, animals were fed only once; RNA was extracted

for RNA-Seq after seven days. For *apob* RNAi, 2 μg control *egfp* dsRNA or 1 μg each *apob-1*

and *apob-2* (for simultaneous RNAi) were mixed with liver and food coloring, and animals were

fed 5 times for initial viability experiments, and 3-5 times for most other experiments, separating

animals with "mild" (*apob-M)* and "severe" *(apob-S)* phenotypes prior to fixation, amputation, or

flow cytometry. Non-eating planarians were always removed from the experiment if they refused

a second dsRNA feeding one day later.

pH3-PS10 immunolabeling

Mucus removal and fixation were conducted with 2% ice-cold HCl (3 min) and

methacarn at room temperature (RT, 20 min), followed by bleaching in 6% H_2O_2 in methanol as

in[109]. Fixed animals/regenerates were blocked (4 hr, RT) in IF block (1X PBS, 0.45% fish

gelatin, 0.6% IgG-free BSA, 0.3% Triton X-100), incubated in rabbit anti-phospho-Histone H3-

S10 at 1:2000 overnight (O/N, 4°C) (Cell Signaling 3377S), washed 8X in PBSTx (1X PBS plus

0.3% Triton X-100) (30 min each, RT), re-blocked for 1 hr, then incubated with goat anti-rabbit-

HRP (1:2000) (Jackson ImmunoResearch 111-035-003) (O/N, 4°C). Samples were again

washed 8X (20-30 min each, RT), then tyramide signal amplification (TSA) was conducted for

10 min as described[110] with TAMRA-tyramide. Samples were washed in PBSTx for two days,

then mounted in Vectashield (Vector Labs H-1000-10).

Immunolabeling (cryosections)

Cryosections (12 μm) of planarians relaxed in 0.66M $MgCl_2$ and fixed (O/N, 4°C) in 4%

formaldehyde (EM)/1X PBS were generated as described[109]. After rehydration, heat-mediated

antigen retrieval (10 min) in 10 mM sodium citrate, pH 6.0 was performed. Slides were

permeabilized for 30 min in 1X PBS/0.2% Tween-20, then blocked for 30 min at RT with 0.45%

fish gelatin and 0.6% BSA for 30 min in PBSTw (1X PBS, 0.05% Tween 20). Slides were

incubated with custom rabbit anti-ApoB-1 (1:1000, 0.59 μg/ml) and mouse 6G10 anti-muscle

(1:250 in block)[111] in blocking buffer (O/N, 4°C). Slides were washed three times (10 min, RT)

with PBSTw after antibody incubation. Slides were then incubated with goat anti-rabbit-HRP

(1:2000) (Jackson ImmunoResearch, 111-035-144) and goat anti-mouse-488 (1:250) (Jackson

ImmunoResearch, 115-545-146) at RT for 60 min. Slides were washed three times (in PBSTw),

with DAPI (2 µg/ml) counterstaining during the first wash. TSA was conducted for 10 min with

TAMRA-tyramide as described[110], followed by washes. Slides were mounted in Fluoromount G

(Southern Biotech).

Anti-ApoB-1 antibody generation

Sequence corresponding to the N-terminal Vitellogenin domain of ApoB-1 (dd_636,

nucleotides 45-1919, amino acids 1-625) was supplied to GenScript USA (Piscataway, NJ).

GenScript synthesized a plasmid encoding this region fused to KLH and a HIS tag, expressed

and purified the fusion protein, immunized New Zealand rabbits, affinity purified the antibody,

and assessed antibody titer and immunogenicity with both ELISA and Western blot.

Protein extraction and Simple Western analysis

5-10 planarians were rocked gently (40-50 rpm) for 7 min in 7.5% N-Acetyl-L-Cysteine

and rinsed 2X in 1X PBS. Samples were then homogenized using a motorized Kontes pestle

grinder in 250 µl RIPA (50 mM Tris pH 8.0, 150 mM NaCl, 1% NP-40, 0.5% sodium

deoxycholate, 0.5% SDS) with 40 mM DTT and 1X Halt Protease Inhibitor cocktail (Thermo

Scientific 78430). After 30 min on ice, samples were centrifuged (20,817 x *g,* 15 min, 4°C), and

supernatant was recovered and stored at -80°C. DTT concentration was reduced by buffer

exchange with RIPA (1 mM DTT) using Amicon Ultra 3 kDa columns (UFC500396), then protein

concentration was determined using a BCA kit (Pierce 23225) and DeNovix cuvet

spectrophotometer according to manufacturers' protocols. For Simple Western (ProteinSimple),

lysates were run according to the manufacturer's protocol on a Wes Instrument (running

Compass v4.0.0) at 0.1 mg/ml using the 66-440 kDa Wes Separation module (SM-W008) with

anti-ApoB-1 at 1:250 (~2.3 μg/ml). ApoB-1 peak areas were calculated with the following

settings: Range 10-600; Baseline Threshold 1.0, Window 15.0, and Stiffness 1.0; and Peak Find

Threshold 10.0 and Width 20.0 using the "Dropped Lines" method. Settings were identical for

total protein peak area, with these exceptions: Baseline Window 50.0 and Stiffness 0.5.

For the regeneration time course, ApoB-1 protein levels were run in biological triplicates,

normalized to total protein using the Wes Total Protein Detection Module (DM-TP01), and then

normalized to the "0 hour" time point in Excel.

Oil Red O staining

Planarians were relaxed in 0.66 M $MgCl_2$, fixed overnight (RT) in 4% formaldehyde (EM

grade) in 1X PBS, protected in sucrose, and cryosectioned (20 μm) onto SuperFrost Plus

slides[109]. Slides were rehydrated in deionized (DI) water (3 x 10 min, RT), then stained in Oil

Red O (Sigma O0625) solution (6 ml Whatman-filtered 0.5% Oil Red O in 100% isopropanol

plus 4 ml ultrapure water) for 15 min at RT. Slides were quickly dipped 3-5X in 60% isopropanol

to remove excess dye, then rinsed for 1 min in 60% isopropanol, then rinsed 1 min in DI water.

Slides were then rinsed in PBS-Tween-20 (0.01%, to prevent drying), then mounted in 90%

glycerol/1X PBS, and imaged within 2-3 days.

Thin Layer Chromatography

Ten planarians (5-8 mm) were placed in 1.7 ml microcentrifuge tubes with all planarian

salts removed, and the animals' mass was obtained. Lipids were then extracted using the Folch

method[112]. Briefly, 1 ml ice-cold 2:1 chloroform:methanol was added, then animals were

sonicated in ice water in a cup-horn sonicator (10 cycles of 5 second pulses at ~48-55W).

Samples were rocked at RT for 5 hr, then centrifuged (2 min at 16,000 x g, 4°C) to pellet

insoluble material. Supernatant was recovered to a new tube and stored at -80°C. 1 ml 2:1

chloroform:methanol was added to the pellet, re-sonicated, rocked overnight at RT, then

centrifuged as above. 0.2 volumes 0.9% NaCl (in water) was added to each extract, tubes were

inverted gently 10-15X to mix, vortexed for 10-15 sec, then centrifuged (2 min at 2,000 x g, RT).

Lower phases from each biological replicate (4 control and 4 *apob(RNAi)* replicates, 10 animals

each) were recovered, and speed-vacuumed (30°C x 60-90 min with spinning) to evaporate

solvent. Concentrated lipids were resuspended in 2 μl chloroform per mg animal mass (above)

and stored at -80°C (for less than 7 days). TLC was performed as previously described[113] with

slight modifications. 150Å silica gel HL 250 μl 20x20 cm plates (iChromatography 66011/Miles

Scientific P76011) were pre-equilibrated with 1:1 chloroform:methanol (~ 1 hr). After drying, 1 μl

lipids and 3 μl standards (30 μg mono-, di-, triglyceride mix, SUPELCO 1787-1AMP, plus 30 μg

cholesteryl palmitate, SIGMA C6072) were spotted onto the plates using TLC spotting

capillaries. Non-polar lipids were resolved with a 80:10:1 petroleum ether:ethyl ether:acetic acid

mix. After drying, TLC plates were sprayed with Primuline (SIGMA 206865) (1 mg/ml fresh stock

in ultrapure water, diluted 1:100 into 80 ml acetone plus 20 ml ultrapure water), dried, and

imaged on an Alpha Innotech chemiluminescent imager with Cy2 excitation/emission filters.

Peak areas in images were quantified in ImageJ/Fiji using the "Plot Lanes" function in the Gels

submenu (https://imagej.nih.gov/ij/docs/menus/analyze.html). Averages were calculated and

normalized to controls in Excel.

RNA extraction, library preparation, and RNA sequencing

Uninjured planarians and regenerating tissue fragments were homogenized in Trizol

using a motorized Kontes pestle grinder, and RNA was extracted using two chloroform

extractions and high-salt precipitation buffer according to the manufacturer's instructions. After

precipitation, solutions were transferred to Zymo RNA columns for DNAse treatment and

purification, according to manufacturer's instructions. RNA samples were analyzed using Agilent

RNA ScreenTape on an Agilent TapeStation 2200 according to manufacturer's protocol.

For analysis of gene expression in control vs. *nkx2.2(RNAi),* mRNA was enriched using

oligo-dT homopolymer beads, and libraries were generated using the Illumina Truseq Stranded

mRNA Library Prep Kit according to the manufacturer's protocol. Final libraries were assayed on

the Agilent TapeStation for appropriate size and quantity. Libraries were pooled in equimolar

amounts as ascertained by fluorometric analysis, then final pools were absolutely quantified

using qPCR on a Roche LightCycler 480 with Kapa Biosystems Illumina Library Quantification

Reagents. Paired-end (2x150 bp) sequence was generated on an Illumina NovaSeq 6000

instrument. 28M-43M reads were generated for each of three biological replicates per condition.

For analysis of gene expression in control, *apob-M,* and *apob-S* animals, total RNA for 4-6

biological replicates per condition was submitted to GENEWIZ (South Plainfield, NJ) for library

generation using NEB NEXT ULTRA library prep and RNA sequencing with standard Illumina

adapters. Paired-end (2x150 bp) sequence was generated on an Illumina HiSeq 4000

instrument; 19M - 26M reads were generated for each replicate.

Read mapping

For both *nkx2.2(RNAi)* and *apob(RNAi)* experiments, quality control and read mapping

to unique transcripts in dd_Smed_v6[104,105] were conducted with FastQC (v0.11.5)[114], BBDuk

(v35.66) (https://sourceforge.net/projects/bbmap/), and Bowtie2 (v2.3.1)[115]. BBDuk (v36.99)

settings for paired end reads: k=13 ktrim=r mink=11 qtrim=rl trimq=10 minlength=35 tbo tpe.

Bowtie2 (v2.3.1) for paired end reads was used for mapping, with "-a" multi-mapping and "--

local" soft-clipping allowed. For read summarization, the "featureCounts" utility in the Subread

package (v1.6.3)[116] was used with a custom ".SAF" file and options "-p -M -O -F SAF" to include

multi-mapping and multi-overlapping reads.

For mapping of X1/X2/Xins and PIWI-HI/-LO/-NEG bulk sequence, regeneration

fragment sequence, and whole animal 24 hr post-irradiation sequence[49], fastq files were

downloaded from NCBI GEO (GSE107874), mapped to dd_Smed_v6_unique using BBDuk and

Bowtie2, followed by count summarization using Samtools as previously described[104].

Differential expression analysis

Read counts matrices were imported into R, then analyzed in edgeR v3.8.6[117]. First, all

transcripts with counts per million (CPM) < 1 in three samples (*nxk2.2(RNAi),* Zeng X1/X2/Xins

data, and Zeng 24 hr irradiation data) or four samples (all others) (e.g., lowly expressed

transcripts) were excluded from further analysis. Next, after recalculation of library sizes,

samples were normalized using trimmed mean of M-values (TMM) method, followed by

calculation of common, trended, and tagwise dispersions. Finally, differentially expressed

transcripts were identified using the pairwise exact test (*nkx2.2(RNAi)* and *apob(RNAi)*

experiments, Zeng 24 hr irradiation data) or the generalized linear model (GLM) likelihood ratio

test (other Zeng datasets). Expression changes were considered to be significant if the false

discovery rate adjusted *p* value ("FDR") was <.05.

Gene Ontology analysis

GO analysis was conducted using BiNGO[118] using a custom *S. mediterranea* GO

annotation as previously described[104]. RefSeq protein collections used for BLASTX and

UniProtKB Biological Process GO terms used for annotation were downloaded for each

organism in April 2020.

Hierarchical clustering and heat maps

Hierarchical clustering of transcripts annotated with lipid-metabolism-related GO terms

was conducted using EdgeR-generated $\log_2$FC values in Cluster 3.0[119], with Euclidean distance

and complete linkage. Heat maps were generated with Java Treeview[120].

qRT-PCR

Total RNA was extracted from biological triplicates (5-10 animals/fragments per

replicate) using Trizol as for RNA-Seq samples. RNA was reverse transcribed using the iScript

cDNA Synthesis kit (BioRad 1708890). *apob-1* and *apob-2* levels were detected using the Fast

Start Essential Green DNA master mix (Roche 06924204001) on a Roche LightCycler 96

instrument. RNA levels were normalized to the geometric mean of endogenous controls *ef-2*

and *gapdh* using the Livak $\Delta\Delta$Ct method[121].

Flow cytometry

Planarians were dissociated and filtered in CMFB with collagenase as described[62]. Cells

were labeled at RT with Hoechst 33342 (50 μg/ml) for 45 min, followed by addition of propidium

iodide (1 μg/ml). For neutral lipid labeling, BODIPY 493/503 (Molecular Probes D3922) at 10

ng/ml was included with Hoechst. Cells were analyzed on a Becton Dickinson FACSCelesta

instrument with 405 nm, 488 nm, and 640 nm lasers. After gating for live cell singlets

(Supplementary Fig. 9a-c), X1, X2, and Xins gates were drawn using two criteria: cell

proportions were approximately 15% (X1) 25% (X2) and 60% (Xins), and reductions in X1 and

X2 fractions in 4-day post-irradiation animals were >95% and ~70%, respectively. Data were

analyzed and plots were generated in FloJo (v10.7.1). For irradiation, uninjured planarians were

dosed with 60 Grays (6,000 rads) using RS-2000 Biological Research X-Ray Irradiator (Rad

Source, Buford, GA).

Cross-referencing of *apob(RNAi)* RNA-Seq data with published transcriptomes

For comparison of dysregulated transcripts in *apob-M* and *apob-S* animals with bulk

neoblast transcriptome data[49], we first identified "signature" transcripts as follows. "X1

signature" transcripts were defined as those with a $\log_2 FC > 0$ (FDR<.05) compared to both "X2"

and "Xins." "X2 signature" transcripts had $\log_2 FC > 0$ (fdr<.05) vs. both "X1" and "Xins." "Xins

signature" had $\log_2 FC > 0$ (fdr<.05) vs. both "X1" and "X2." Similarly, "PIWI-HI signature"

transcripts were defined as those with a $\log_2 FC > 0$ (fdr<.05) compared to both "PIWI-LO" and

"PIWI-NEG." "PIWI-LO signature" transcripts had $\log_2 FC > 0$ (fdr<.05) vs. both "PIWI-HI" and

"PIWI-NEG." "PIWI-NEG signature" transcripts had $\log_2 FC > 0$ (fdr<.05) vs. both "PIWI-HI" and

"PIWI-LO." Next, we used the "merge" function and "VennDiagram" package in RStudio

(v1.2.1335) to identify signature transcripts in X1/X2/Xins or PIWI-HI/-LO/-NEG that were also

dsyregulated (up or down) in *apob-M* or *apob-S* animals, as shown in Fig. S6. "% of

dysregulated transcripts" was calculated as the number of overlapping transcripts divided by the

total number of X1/X2/Xins or PIWI-HI/-LO/-NEG transcripts.

For comparison of dysregulated transcripts in *apob-M* and *apob-S* animals with single

cell type/state data in[40], we again used the "merge" function in RStudio to identify the number of

transcripts enriched in individual lineage subclusters (Table S2)[40] that were also dysregulated

by *apob* RNAi. "% of dysregulated transcripts" was calculated as the number of overlapping

transcripts divided by the total number of transcripts in each individual subcluster. In Fig. 5, the

"N/TS" (Neoblast/Transition State) designation included subclusters with high *piwi-1* mRNA

expression thought to be neoblast/progenitor subpopulations in epidermal, intestine, and

protonephridia lineages based on the conclusions of Fincher et al. and other published

data[45,62,122-124]. Similarly, the "P" (Progeny) and "M" (Mature) designations were based on

conclusions from both single cell RNA-Seq data and previous work. For lineages that are less

well understood *in vivo* (Fig. S8), we designated subclusters/states using both transcript

dysregulation in 24 hr irradiated animals [49] and *piwi-1* mRNA levels in t-SNE plots[40]. "N/TS"

subclusters possessed the greatest number of irradiation-dysregulated transcripts and the

highest *piwi-1* expression; "P" subclusters possessed fewer (by proportion) irradation-

dysregulated transcripts and lower *piwi-1* expression; and "M" subclusters had the fewest

radiation-sensitive transcripts and negligible *piwi-1* expression.

Image Collection and Quantification

Epifluorescent images (FISH samples) were collected on a Zeiss AxioObserver.Z1 with

Excelitas X-Cite 120 LED Boost illumination and Zen 2.3 pro. For quantification of pharynx and

brain size, organ area and animal area were measured in ImageJ[125] and organ-to-body size

ratios were calculated. For intestine, length of anterior branch and posterior branches were

measured in ImageJ. Means of posterior branch length were calculated, and then posterior-to-

anterior (head fragments) or anterior-to-posterior (tail fragments) length ratio was calculated.

For tail fragments with split anterior branch, anterior branch length was measured from the

anterior of the pharynx to the tip of the anterior-most primary branch. For tail fragments, the split

anterior branch phenotype (failure to fuse at the midline) was scored if there was an obvious

gap between anterior branches for at least half the length of the anterior branch. For images of

apob-1 and *apob-2* FISH on cryosections, z-stacks were collected with an Apotome.2 for

generation of maximum orthogonal projections.

For anti-pH3-PS10-labeled samples, z-stacks were collected on a Zeiss

AxioObserver.Z1 at 10X magnification, followed by tile stitching, extended depth of focus

projection, and background subtraction (PS10 channel only) with a radius of 30. Control and

experimental samples to be compared were imaged at identical exposures. For quantification,

animal area and PS10+ nuclei were quantified using the Automated Segmentation tools in Zen

2.3 pro. Briefly, animal area was measured with Gaussian smoothing, no background

subtraction or sharpening, Morphology separation, and custom threshold settings that were the

same for all samples to be directly compared. PS10+ nuclei number was measured using

Lowpass smoothing, Rolling Ball background subtraction, Delineate sharpening, Watersheds

separation, with threshold settings and other parameters that were identical for all samples to be

directly compared. Number of mitoses per area were calculated for each animal/fragment.

Confocal images were collected on a Zeiss LSM 710 or LSM 880 laser scanning

microscope with 10X Plan NeoFluar, 20X Plan Apo, or 40X C-Apo objectives. For orthogonal

projections (anti-ApoB1 immunolabeling and *ldlr/vldlr* FISH), between three and ten z-planes

were collected at 1-2X "optimal" section thickness (based on objective NA). For *notum* and

wnt11-2 FISH, full fragment thickness stacks were projected to ensure that all mRNA-positive

cells were counted.

Images of Oil-Red-O-stained sections, live animals, live regenerates, and WISH samples

were collected on a Zeiss Stemi 508 with an Axiocam 105 color camera, or a Zeiss Axio

Zoom.V16 with an Axiocam 105 camera. In some cases brightness and/or contrast were

adjusted in Adobe Photoshop to improve signal contrast.

Statistics

Detailed data and information regarding statistical testing are included in Source Data 2.

For experiments with statistical analysis, n values are indicated exactly by the number of data

points in figure plots, along with definitions of error bars and *p* or *q* values; all tests were

conducted in Prism 9 (GraphPad Software, San Diego, CA). For one-way ANOVA, ordinary

ANOVA was performed unless Brown-Forsythe and Bartlett's tests indicated standard

deviations were significantly different, then Brown-Forsythe and Welch ANOVA tests were

performed. For two-way ANOVA (flow cytometry experiments with irradiation), *q* values (FDR-

adjusted *p* values) were reported when interaction between RNAi condition ("Genotype") and

irradiation was significant (X2 subpopulation). Otherwise, *p* values were derived using one-way

ANOVA. *p* values of $<.05$ (*), $<.01$ (**), $<.001$ (***), and <0.0001 (****) were annotated with

asterisks in figures.

Replicate information for other experiments: polarity marker analysis, 5-15 fragments per

condition; for WISH and FISH, images are representative of 3-6 individual animals or fragments;

for anti-ApoB1 and Oil Red O labeling, images are representative of sections from 2-3

fragments/animals, and of at least two repeated experiments.

For statistical testing of overlap between genes dysregulated in *apob(RNAi)* planarians

and other RNA-Seq data sets, the R Package GeneOverlap[126] was used to conduct Fisher's

exact test on each comparison. Total number of detected transcripts ("genome size" in

GeneOverlap) was determined conservatively by only including transcripts detected in both

apob RNA-Seq and bulk[49] or single cell[40] sequencing data. For sc-RNA-Seq, the digital

expression matrix in GEO GSE111764 was normalized in Seurat as in Fincher et al.[40]; only

transcripts with non-zero expression in 0.5% of cells (Fincher TableS1) in each subcluster were

considered to be detected. *p* values <0.05 are indicated with a caret (^) in figures.

Replicate and statistical information for RNA-Seq and other bioinformatics experiments

are detailed in appropriate Methods sections.

Data availability

Raw and processed RNA-Seq data associated with this study will be made available in

the NCBI Gene Expression Omnibus (GEO) upon publication. Other data supporting this study's

findings are available within the article and its Supplementary files, or are available from the

authors upon reasonable request.

Reporting summary

Further information on research design is available in the Nature Research Reporting

Summary linked to this article.

Competing interests

The authors declare no competing financial interests.

Acknowledgments

We are very grateful to Phil Newmark (HHMI and the Morgridge Institute for Research), in whose laboratory this project was initiated. We thank members of the Newmark and Forsthoefel labs, and OMRF colleagues Pat Gaffney, Linda Thompson, Dean Dawson, and Hui-Ying Lim for insightful discussions. We thank Jochen Rink, James Cleland, and Hanh Vu (Max Planck Institute for Biophysical Chemistry) for sharing planarian protein extraction protocols, Vasileios Morkotinis for *tgs-1* plasmid, and Rachel Roberts-Galbraith (Univ. of Georgia) for *notum* and *wnt11-2* plasmids. We thank Steve Farber (Carnegie Institution for Science) for advice on lipid extraction and TLC, and Jayhun Lee (Morgridge) for collaborative development of planarian lipid analysis protocols. We are grateful to Hadi Maktabi (ProteinSimple) and to OMRF colleagues Summer Wang and Lin Wang for help in developing Wes protocols; and to members of the OMRF Flow Cytometry Core (Jacob Bass and Diana Hamilton), the Quantitative Analysis Core (Lori Garman and Nathan Pezant), the Imaging and Histology Core, the Clinical Genomics Core, the Gnotobiotic Mouse Core (for use of X-irradiator), and IT/Research Computing Services for invaluable technical assistance. JRO was supported by the Summer Research Opportunities Program at the University of Illinois at Urbana-Champaign. This work was supported by NIH Centers of Biomedical Research Excellence (COBRE) GM103636 (Project 1 to DJF), and the Oklahoma Medical Research Foundation.

Author Contributions

Conception and design of the project: DJF and LLW; data collection: CGB, LLW, JRO, NIC, and DJF; data analysis and visualization: CGB, LLW, NIC and DJF; data interpretation: LLW, CGB, and DJF; RNA-seq analysis: DJF; manuscript preparation: DJF, LLW, and CGB.

References

1   Walther, T. C., Chung, J. & Farese, R. V., Jr. Lipid Droplet Biogenesis. *Annu. Rev. Cell.*
Dev. Biol. **33**, 491-510, doi:10.1146/annurev-cellbio-100616-060608 (2017).

2   Olzmann, J. A. & Carvalho, P. Dynamics and functions of lipid droplets. *Nat. Rev. Mol.*
Cell Biol. **20**, 137-155, doi:10.1038/s41580-018-0085-z (2019).

3   D'Andrea, S. Lipid droplet mobilization: The different ways to loosen the purse strings.
Biochimie **120**, 17-27, doi:10.1016/j.biochi.2015.07.010 (2016).

4   Guo, Y., Cordes, K. R., Farese, R. V., Jr. & Walther, T. C. Lipid droplets at a glance. *J.*
Cell Sci. **122**, 749-752, doi:10.1242/jcs.037630 (2009).

5   Olofsson, S. O., Asp, L. & Borén, J. The assembly and secretion of apolipoprotein B-
containing lipoproteins. *Curr Opin Lipidol* **10**, 341-346, doi:10.1097/00041433-
199908000-00008 (1999).

6   Heier, C. & Kühnlein, R. P. Triacylglycerol Metabolism in Drosophila melanogaster.
Genetics **210**, 1163-1184, doi:10.1534/genetics.118.301583 (2018).

7   Wong, L. H., Gatta, A. T. & Levine, T. P. Lipid transfer proteins: the lipid commute via
shuttles, bridges and tubes. *Nat. Rev. Mol. Cell Biol.* **20**, 85-101, doi:10.1038/s41580-
018-0071-5 (2019).

8   Lafontan, M. & Langin, D. Lipolysis and lipid mobilization in human adipose tissue. *Prog*
Lipid Res **48**, 275-297, doi:10.1016/j.plipres.2009.05.001 (2009).

9   Ruggles, K. V., Turkish, A. & Sturley, S. L. Making, baking, and breaking: the synthesis,
storage, and hydrolysis of neutral lipids. *Annu. Rev. Nutr.* **33**, 413-451,
doi:10.1146/annurev-nutr-071812-161254 (2013).

10   Welte, M. A. & Gould, A. P. Lipid droplet functions beyond energy storage. *Biochim*
Biophys Acta Mol Cell Biol Lipids **1862**, 1260-1272, doi:10.1016/j.bbalip.2017.07.006
(2017).

11   Ly, C. H., Lynch, G. S. & Ryall, J. G. A Metabolic Roadmap for Somatic Stem Cell Fate.
Cell Metab **31**, 1052-1067, doi:10.1016/j.cmet.2020.04.022 (2020).

12   Ito, K. & Suda, T. Metabolic requirements for the maintenance of self-renewing stem
cells. *Nat. Rev. Mol. Cell Biol.* **15**, 243-256, doi:10.1038/nrm3772 (2014).

13   Shyh-Chang, N. & Ng, H. H. The metabolic programming of stem cells. *Genes Dev.* **31**,
336-346, doi:10.1101/gad.293167.116 (2017).

14   Clémot, M., Sênos Demarco, R. & Jones, D. L. Lipid Mediated Regulation of Adult Stem
Cell Behavior. *Front Cell Dev Biol* **8**, 115, doi:10.3389/fcell.2020.00115 (2020).

15   Ito, K. *et al.* A PML-PPAR-delta pathway for fatty acid oxidation regulates hematopoietic
stem cell maintenance. *Nat. Med.* **18**, 1350-1358, doi:10.1038/nm.2882 (2012).

16   Knobloch, M. *et al.* A Fatty Acid Oxidation-Dependent Metabolic Shift Regulates Adult
Neural Stem Cell Activity. *Cell Rep* **20**, 2144-2155, doi:10.1016/j.celrep.2017.08.029
(2017).

17   Singh, S. R. *et al.* The lipolysis pathway sustains normal and transformed stem cells in
adult Drosophila. *Nature* **538**, 109-113, doi:10.1038/nature19788 (2016).

18   Sênos Demarco, R., Uyemura, B. S., D'Alterio, C. & Jones, D. L. Mitochondrial fusion
regulates lipid homeostasis and stem cell maintenance in the Drosophila testis. *Nat. Cell*
Biol. **21**, 710-720, doi:10.1038/s41556-019-0332-3 (2019).

19   Tiwari, S. K., Toshniwal, A. G., Mandal, S. & Mandal, L. Fatty acid beta-oxidation is
required for the differentiation of larval hematopoietic progenitors in Drosophila. *Elife* **9**,
doi:10.7554/eLife.53247 (2020).

20   Knobloch, M. *et al.* Metabolic control of adult neural stem cell activity by Fasn-dependent
lipogenesis. *Nature* **493**, 226-230, doi:10.1038/nature11689 (2013).

21   Driver, A. M., Kratz, L. E., Kelley, R. I. & Stottmann, R. W. Altered cholesterol
biosynthesis causes precocious neurogenesis in the developing mouse forebrain.
Neurobiol Dis **91**, 69-82, doi:10.1016/j.nbd.2016.02.017 (2016).

22   Cornacchia, D. *et al.* Lipid Deprivation Induces a Stable, Naive-to-Primed Intermediate
State of Pluripotency in Human PSCs. *Cell Stem Cell* **25**, 120-136 e110,
doi:10.1016/j.stem.2019.05.001 (2019).

23   Kang, J. *et al.* Modulation of tissue repair by regeneration enhancer elements. *Nature*
532, 201-206, doi:10.1038/nature17644 (2016).

24   Monnot, M. J. *et al.* Epidermal expression of apolipoprotein E gene during fin and scale
development and fin regeneration in zebrafish. *Dev. Dyn.* **214**, 207-215,
doi:10.1002/(SICI)1097-0177(199903)214:3<207::AID-AJA4>3.0.CO;2-5 (1999).

25   Voss, S. R. *et al.* Gene expression during the first 28 days of axolotl limb regeneration I:
Experimental design and global analysis of gene expression. *Regeneration (Oxf)* **2**, 120-
136, doi:10.1002/reg2.37 (2015).

26   Hayashi, H., Campenot, R. B., Vance, D. E. & Vance, J. E. Glial lipoproteins stimulate
axon growth of central nervous system neurons in compartmented cultures. *J. Biol.*
Chem. **279**, 14009-14015, doi:10.1074/jbc.M313828200 (2004).

27   Pauta, M. *et al.* Impaired liver regeneration in Ldlr-/- mice is associated with an altered
hepatic profile of cytokines, growth factors, and lipids. *J Hepatol* **59**, 731-737,
doi:10.1016/j.jhep.2013.05.026 (2013).

28   Tamilarasan, K. P. *et al.* Skeletal muscle damage and impaired regeneration due to LPL-
mediated lipotoxicity. *Cell Death Dis* **3**, e354, doi:10.1038/cddis.2012.91 (2012).

29   Pala, F. *et al.* Distinct metabolic states govern skeletal muscle stem cell fates during
prenatal and postnatal myogenesis. *J. Cell Sci.* **131**, doi:10.1242/jcs.212977 (2018).

30   Shook, B. A. *et al.* Dermal Adipocyte Lipolysis and Myofibroblast Conversion Are
Required for Efficient Skin Repair. *Cell Stem Cell* **26**, 880-895 e886,
doi:10.1016/j.stem.2020.03.013 (2020).

31   Newmark, P. A. & Sánchez Alvarado, A. Not your father's planarian: a classic model
enters the era of functional genomics. *Nat. Rev. Genet.* **3**, 210-219, doi:10.1038/nrg759
(2002).

32   Reddien, P. W. & Sánchez Alvarado, A. Fundamentals of planarian regeneration. *Annu.*
Rev. Cell Dev. Biol. **20**, 725-757 (2004).

33   Willier, B. H., Hyman, L. H. & Rifenburgh, S. A. A histochemical study of intracellular
digestion in triclad flatworms. *Journal of Morphology and Physiology* **40**, 299-340,
doi:DOI 10.1002/jmor.1050400205 (1925).

34   Jennings, J. B. Studies on feeding, digestion, and food storage in free-living flatworms
(Platyhelminthes: Turbellaria). *Biol. Bull.* **112**, 63-80 (1957).

35   Mak, H. Y. Lipid droplets as fat storage organelles in Caenorhabditis elegans: Thematic
Review Series: Lipid Droplet Synthesis and Metabolism: from Yeast to Man. *J. Lipid*
Res. **53**, 28-33, doi:10.1194/jlr.R021006 (2012).

36   Gamo, J. & Garcia-Corrales, P. Ultrastructural changes in the gastrodermal phagocytic
cells of the freshwater planarian *Dugesia gonocephala* s.l. during starvation. *J.*
Submicrosc. Cytol. **19**, 291-302 (1987).

37   Thommen, A. *et al.* Body size-dependent energy storage causes Kleiber's law scaling of
the metabolic rate in planarians. *Elife* **8**, doi:10.7554/eLife.38187 (2019).

38   Palm, W. & Rodenfels, J. Understanding the role of lipids and lipoproteins in
development. *Development* **147**, doi:10.1242/dev.186411 (2020).

39   Forsthoefel, D. J. *et al.* An RNAi screen reveals intestinal regulators of branching
morphogenesis, differentiation, and stem cell proliferation in planarians. *Dev. Cell* **23**,
691-704, doi:10.1016/j.devcel.2012.09.008 (2012).

40   Fincher, C. T., Wurtzel, O., de Hoog, T., Kravarik, K. M. & Reddien, P. W. Cell type
transcriptome atlas for the planarian *Schmidtea mediterranea*. *Science* **360**,
doi:10.1126/science.aaq1736 (2018).

41   Rouhana, L. *et al.* RNA interference by feeding in vitro-synthesized double-stranded
RNA to planarians: methodology and dynamics. *Dev. Dyn.* **242**, 718-730,
doi:10.1002/dvdy.23950 (2013).

42   Reddien, P. W., Bermange, A. L., Murfitt, K. J., Jennings, J. R. & Sánchez Alvarado, A.
Identification of genes needed for regeneration, stem cell function, and tissue
homeostasis by systematic gene perturbation in planaria. *Dev. Cell* **8**, 635-649,
doi:10.1016/j.devcel.2005.02.014 (2005).

43   Palm, W. *et al.* Lipoproteins in Drosophila melanogaster--assembly, function, and
influence on tissue lipid composition. *PLoS Genet.* **8**, e1002828,
doi:10.1371/journal.pgen.1002828 (2012).

44   Young, S. G. Recent progress in understanding apolipoprotein B. *Circulation* **82**, 1574-
1594, doi:10.1161/01.cir.82.5.1574 (1990).

45   Wagner, D. E., Wang, I. E. & Reddien, P. W. Clonogenic neoblasts are pluripotent adult
stem cells that underlie planarian regeneration. *Science* **332**, 811-816,
doi:10.1126/science.1203983 (2011).

46   Hayashi, T., Asami, M., Higuchi, S., Shibata, N. & Agata, K. Isolation of planarian X-ray-
sensitive stem cells by fluorescence-activated cell sorting. *Dev. Growth Differ.* **48**, 371-
380, doi:10.1111/j.1440-169X.2006.00876.x (2006).

47   Wenemoser, D. & Reddien, P. W. Planarian regeneration involves distinct stem cell
responses to wounds and tissue absence. *Dev. Biol.* **344**, 979-991,
doi:10.1016/j.ydbio.2010.06.017 (2010).

48   Wurtzel, O. *et al.* A generic and cell-type-specific wound response precedes
regeneration in planarians. *Dev. Cell* **35**, 632-645, doi:10.1016/j.devcel.2015.11.004
(2015).

49   Zeng, A. *et al.* Prospectively Isolated Tetraspanin(+) Neoblasts Are Adult Pluripotent
Stem Cells Underlying Planaria Regeneration. *Cell* **173**, 1593-1608 e1520,
doi:10.1016/j.cell.2018.05.006 (2018).

50   Reddien, P. W., Oviedo, N. J., Jennings, J. R., Jenkin, J. C. & Sánchez Alvarado, A.
SMEDWI-2 is a PIWI-like protein that regulates planarian stem cells. *Science* **310**, 1327-
1330, doi:10.1126/science.1116110 (2005).

51   Scimone, M. L., Meisel, J. & Reddien, P. W. The Mi-2-like *Smed-CHD4* gene is required
for stem cell differentiation in the planarian *Schmidtea mediterranea*. *Development* **137**,
1231-1241, doi:dev.042051 [pii]
10.1242/dev.042051 (2010).

52   Zhu, S. J., Hallows, S. E., Currie, K. W., Xu, C. & Pearson, B. J. A *mex3* homolog is
required for differentiation during planarian stem cell lineage development. *Elife* **4**,
e07025, doi:10.7554/eLife.07025 (2015).

53   Reddien, P. W. The Cellular and Molecular Basis for Planarian Regeneration. *Cell* **175**,
327-345, doi:10.1016/j.cell.2018.09.021 (2018).

54   Petersen, C. P. & Reddien, P. W. Polarized notum activation at wounds inhibits Wnt
function to promote planarian head regeneration. *Science* **332**, 852-855,
doi:10.1126/science.1202143 (2011).

55   Adell, T., Saló, E., Boutros, M. & Bartscherer, K. Smed-Evi/Wntless is required for β-
catenin-dependent and -independent processes during planarian regeneration.
Development **136**, 905-910, doi:10.1242/dev.033761 (2009).

56 Forsthoefel, D. J., Park, A. E. & Newmark, P. A. Stem cell-based growth, regeneration,
and remodeling of the planarian intestine. *Dev. Biol.* **356**, 445-459,
doi:10.1016/j.ydbio.2011.05.669 (2011).

57 Raz, A. A., Wurtzel, O. & Reddien, P. W. Planarian stem cells specify fate yet retain
potency during the cell cycle. *Cell Stem Cell*, doi:10.1016/j.stem.2021.03.021 (2021).

58 Hendzel, M. J. *et al.* Mitosis-specific phosphorylation of histone H3 initiates primarily
within pericentromeric heterochromatin during G2 and spreads in an ordered fashion
coincident with mitotic chromosome condensation. *Chromosoma* **106**, 348-360 (1997).

59 Newmark, P. A. & Sánchez Alvarado, A. Bromodeoxyuridine specifically labels the
regenerative stem cells of planarians. *Dev. Biol.* **220**, 142-153,
doi:10.1006/dbio.2000.9645 (2000).

60 Eisenhoffer, G. T., Kang, H. & Sánchez Alvarado, A. Molecular analysis of stem cells
and their descendants during cell turnover and regeneration in the planarian *Schmidtea*
mediterranea. *Cell Stem Cell* **3**, 327-339, doi:S1934-5909(08)00336-6 [pii]
10.1016/j.stem.2008.07.002 (2008).

61 Hayashi, T. *et al.* Single-cell gene profiling of planarian stem cells using fluorescent
activated cell sorting and its "index sorting" function for stem cell research. *Dev. Growth*
Differ. **52**, 131-144, doi:DGD1157 [pii]
10.1111/j.1440-169X.2009.01157.x (2010).

62 van Wolfswinkel, J. C., Wagner, D. E. & Reddien, P. W. Single-cell analysis reveals
functionally distinct classes within the planarian stem cell compartment. *Cell Stem Cell*
15, 326-339, doi:10.1016/j.stem.2014.06.007 (2014).

63 Chan, A., Ma, S., Pearson, B. J. & Chan, D. Collagen IV differentially regulates planarian
stem cell potency and lineage progression. *Proc Natl Acad Sci U S A* **118**,
doi:10.1073/pnas.2021251118 (2021).

64 Stelman, C. R., Smith, B. M., Chandra, B. & Roberts-Galbraith, R. H. CBP/p300
homologs CBP2 and CBP3 play distinct roles in planarian stem cell function. *Dev. Biol.*
473, 130-143, doi:10.1016/j.ydbio.2021.02.004 (2021).

65 Fraguas, S. *et al.* CREB-binding protein (CBP) gene family regulates planarian survival
and stem cell differentiation. *Dev. Biol.* **476**, 53-67, doi:10.1016/j.ydbio.2021.02.008
(2021).

66 Deb, S. *et al.* Tnfaip2/exoc3-driven lipid metabolism is essential for stem cell
differentiation and organ homeostasis. *EMBO Rep*, e49328,
doi:10.15252/embr.201949328 (2020).

67 Yadav, T., Quivy, J. P. & Almouzni, G. Chromatin plasticity: A versatile landscape that
underlies cell fate and identity. *Science* **361**, 1332-1336, doi:10.1126/science.aat8950
(2018).

68 Duncan, E. M. & Sánchez Alvarado, A. Regulation of Genomic Output and
(Pluri)potency in Regeneration. *Annu. Rev. Genet.* **53**, 327-346, doi:10.1146/annurev-
genet-112618-043733 (2019).

69 McDonnell, E. *et al.* Lipids Reprogram Metabolism to Become a Major Carbon Source
for Histone Acetylation. *Cell Rep* **17**, 1463-1472, doi:10.1016/j.celrep.2016.10.012
(2016).

70 Zhang, S. *et al.* A nuclear hormone receptor and lipid metabolism axis are required for
the maintenance and regeneration of reproductive organs. *bioRxiv*, 279364,
doi:10.1101/279364 (2018).

71 Tsukada, Y. *et al.* Histone demethylation by a family of JmjC domain-containing proteins.
Nature **439**, 811-816, doi:10.1038/nature04433 (2006).

72 Robb, S. M. & Sánchez Alvarado, A. Histone modifications and regeneration in the
planarian Schmidtea mediterranea. *Curr Top Dev Biol* **108**, 71-93, doi:10.1016/B978-0-
12-391498-9.00004-8 (2014).

73 Cao, P. L., Kumagai, N., Inoue, T., Agata, K. & Makino, T. JmjC Domain-Encoding
Genes Are Conserved in Highly Regenerative Metazoans and Are Associated with
Planarian Whole-Body Regeneration. *Genome Biol Evol* **11**, 552-564,
doi:10.1093/gbe/evz021 (2019).

74 Mihaylova, Y. *et al.* Conservation of epigenetic regulation by the MLL3/4 tumour
suppressor in planarian pluripotent stem cells. *Nat Commun* **9**, 3633,
doi:10.1038/s41467-018-06092-6 (2018).

75 Hubert, A. *et al.* Epigenetic regulation of planarian stem cells by the SET1/MLL family of
histone methyltransferases. *Epigenetics* **8**, 79-91, doi:10.4161/epi.23211 (2013).

76 Mohamed Haroon, M. *et al.* Mitochondrial state determines functionally divergent stem
cell population in planaria. *Stem Cell Reports*, doi:10.1016/j.stemcr.2021.03.022 (2021).

77 Chung, S. *et al.* Mitochondrial oxidative metabolism is required for the cardiac
differentiation of stem cells. *Nat. Clin. Pract. Cardiovasc. Med.* **4 Suppl 1**, S60-67,
doi:10.1038/ncpcardio0766 (2007).

78 Zhang, Y., Marsboom, G., Toth, P. T. & Rehman, J. Mitochondrial respiration regulates
adipogenic differentiation of human mesenchymal stem cells. *PLoS One* **8**, e77077,
doi:10.1371/journal.pone.0077077 (2013).

79 Teixeira, F. K. *et al.* ATP synthase promotes germ cell differentiation independent of
oxidative phosphorylation. *Nat. Cell Biol.* **17**, 689-696, doi:10.1038/ncb3165 (2015).

80 Baris, O. R. *et al.* The mitochondrial electron transport chain is dispensable for
proliferation and differentiation of epidermal progenitor cells. *Stem Cells* **29**, 1459-1468,
doi:10.1002/stem.695 (2011).

81   Panáková, D., Sprong, H., Marois, E., Thiele, C. & Eaton, S. Lipoprotein particles are
required for Hedgehog and Wingless signalling. *Nature* **435**, 58-65,
doi:10.1038/nature03504 (2005).

82   Blomhoff, R. & Blomhoff, H. K. Overview of retinoid metabolism and function. *J.*
Neurobiol. **66**, 606-630, doi:10.1002/neu.20242 (2006).

83   Rigotti, A. Absorption, transport, and tissue delivery of vitamin E. *Mol Aspects Med* **28**,
423-436, doi:10.1016/j.mam.2007.01.002 (2007).

84   Gil, A., Plaza-Diaz, J. & Mesa, M. D. Vitamin D: Classic and Novel Actions. *Ann Nutr*
Metab **72**, 87-95, doi:10.1159/000486536 (2018).

85   Fernández-Barral, A. *et al.* Vitamin D differentially regulates colon stem cells in patient-
derived normal and tumor organoids. *FEBS J.* **287**, 53-72, doi:10.1111/febs.14998
(2020).

86   Deng, S. *et al.* Vitamin E isomer delta-tocopherol enhances the efficiency of neural stem
cell differentiation via L-type calcium channel. *Neurosci. Lett.* **585**, 166-170,
doi:10.1016/j.neulet.2014.11.031 (2015).

87   Gudas, L. J. & Wagner, J. A. Retinoids regulate stem cell differentiation. *J Cell Physiol*
226, 322-330, doi:10.1002/jcp.22417 (2011).

88   Lindsay-Mosher, N., Chan, A. & Pearson, B. J. Planarian EGF repeat-containing genes
megf6 and hemicentin are required to restrict the stem cell compartment. *PLoS Genet.*
16, e1008613, doi:10.1371/journal.pgen.1008613 (2020).

89   Cote, L. E., Simental, E. & Reddien, P. W. Muscle functions as a connective tissue and
source of extracellular matrix in planarians. *Nat Commun* **10**, 1592, doi:10.1038/s41467-
019-09539-6 (2019).

90   Betchaku, T. Isolation of planarian neoblasts and their behavior in vitro with some
aspects of the mechanism of the formation of regeneration blastema. *J Exp Zool* **164**,
407-433, doi:10.1002/jez.1401640310 (1967).

91   Cruz, A. L. S., Barreto, E. A., Fazolini, N. P. B., Viola, J. P. B. & Bozza, P. T. Lipid
droplets: platforms with multiple functions in cancer hallmarks. *Cell Death Dis* **11**, 105,
doi:10.1038/s41419-020-2297-3 (2020).

92   Tirinato, L. *et al.* An Overview of Lipid Droplets in Cancer and Cancer Stem Cells. *Stem*
Cells Int **2017**, 1656053, doi:10.1155/2017/1656053 (2017).

93   Sinclair, J. W. *et al.* A metabolic shift to glycolysis promotes zebrafish tail regeneration
through TGF–β dependent dedifferentiation of notochord cells to form the blastema.
bioRxiv, 2020.2003.2003.975318, doi:10.1101/2020.03.03.975318 (2020).

94   Love, N. R., Ziegler, M., Chen, Y. & Amaya, E. Carbohydrate metabolism during
vertebrate appendage regeneration: what is its role? How is it regulated?: A postulation
that regenerating vertebrate appendages facilitate glycolytic and pentose phosphate

pathways to fuel macromolecule biosynthesis. *Bioessays* **36**, 27-33,
doi:10.1002/bies.201300110 (2014).

95   Shyh-Chang, N., Daley, G. Q. & Cantley, L. C. Stem cell metabolism in tissue
development and aging. *Development* **140**, 2535-2547, doi:10.1242/dev.091777 (2013).

96   Olson, R. E. Discovery of the lipoproteins, their role in fat transport and their significance
as risk factors. *J Nutr Sci* **128**, 439S-443S, doi:10.1093/jn/128.2.439S (1998).

97   Feng, Y. *et al.* Hematopoietic stem/progenitor cell proliferation and differentiation is
differentially regulated by high-density and low-density lipoproteins in mice. *PLoS One* **7**,
e47286, doi:10.1371/journal.pone.0047286 (2012).

98   Xu, J. *et al.* High density lipoprotein cholesterol promotes the proliferation of bone-
derived mesenchymal stem cells via binding scavenger receptor-B type I and activation
of PI3K/Akt, MAPK/ERK1/2 pathways. *Mol. Cell. Biochem.* **371**, 55-64,
doi:10.1007/s11010-012-1422-8 (2012).

99   Shen, H. *et al.* High density lipoprotein promotes proliferation of adipose-derived stem
cells via S1P1 receptor and Akt, ERK1/2 signal pathways. *Stem Cell Res Ther* **6**, 95,
doi:10.1186/s13287-015-0090-5 (2015).

100 Safina, D. *et al.* Low-density lipoprotein receptor-related protein 1 is a novel modulator of
radial glia stem cell proliferation, survival, and differentiation. *Glia* **64**, 1363-1380,
doi:10.1002/glia.23009 (2016).

101 Gaudet, D., Drouin-Chartier, J. P. & Couture, P. Lipid Metabolism and Emerging Targets
for Lipid-Lowering Therapy. *Can J Cardiol* **33**, 872-882, doi:10.1016/j.cjca.2016.12.019
(2017).

102 Sánchez Alvarado, A., Newmark, P. A., Robb, S. M. & Juste, R. The *Schmidtea
mediterranea* database as a molecular resource for studying platyhelminthes, stem cells
and regeneration. *Development* **129**, 5659-5665 (2002).

103 Roberts-Galbraith, R. H. & Newmark, P. A. Follistatin antagonizes Activin signaling and
acts with Notum to direct planarian head regeneration. *Proc Natl Acad Sci U S A* **110**,
1363-1368, doi:10.1073/pnas.1214053110 (2013).

104 Forsthoefel, D. J., Cejda, N. I., Khan, U. W. & Newmark, P. A. Cell-type diversity and
regionalized gene expression in the planarian intestine. *Elife* **9**, doi:10.7554/eLife.52613
(2020).

105 Brandl, H. *et al.* PlanMine--a mineable resource of planarian biology and biodiversity.
Nucleic Acids Res. **44**, D764-773, doi:10.1093/nar/gkv1148 (2016).

106 Zayas, R. M. *et al.* The planarian *Schmidtea mediterranea* as a model for epigenetic
germ cell specification: analysis of ESTs from the hermaphroditic strain. *Proc. Natl.
Acad. Sci. USA* **102**, 18491-18496 (2005).

107 Potter, S. C. *et al.* HMMER web server: 2018 update. *Nucleic Acids Res.* **46**, W200-
W204, doi:10.1093/nar/gky448 (2018).

108 Guindon, S. *et al.* New algorithms and methods to estimate maximum-likelihood
phylogenies: assessing the performance of PhyML 3.0. *Syst. Biol.* **59**, 307-321,
doi:10.1093/sysbio/syq010 (2010).

109 Forsthoefel, D. J., Waters, F. A. & Newmark, P. A. Generation of cell type-specific
monoclonal antibodies for the planarian and optimization of sample processing for
immunolabeling. *BMC Dev. Biol.* **14**, 45, doi:10.1186/s12861-014-0045-6 (2014).

110 King, R. S. & Newmark, P. A. *In situ* hybridization protocol for enhanced detection of
gene expression in the planarian *Schmidtea mediterranea*. *BMC Dev. Biol.* **13**, 8,
doi:10.1186/1471-213X-13-8 (2013).

111 Ross, K. G. *et al.* Novel monoclonal antibodies to study tissue regeneration in
planarians. *BMC Dev. Biol.* **15**, 2, doi:10.1186/s12861-014-0050-9 (2015).

112 Folch, J. & Lees, M. Proteolipides, a new type of tissue lipoproteins; their isolation from
brain. *J. Biol. Chem.* **191**, 807-817 (1951).

113 Carten, J. D., Bradford, M. K. & Farber, S. A. Visualizing digestive organ morphology
and function using differential fatty acid metabolism in live zebrafish. *Dev. Biol.* **360**, 276-
285, doi:10.1016/j.ydbio.2011.09.010 (2011).

114 Andrews, S. *FastQC: a quality control tool for high throughput sequence data.*,
<http://www.bioinformatics.babraham.ac.uk/projects/fastqc> (2010).

115 Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat.*
Methods **9**, 357-359, doi:10.1038/nmeth.1923 (2012).

116 Liao, Y., Smyth, G. K. & Shi, W. The R package Rsubread is easier, faster, cheaper and
better for alignment and quantification of RNA sequencing reads. *Nucleic Acids Res.* **47**,
e47, doi:10.1093/nar/gkz114 (2019).

117 Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-
140, doi:10.1093/bioinformatics/btp616 (2010).

118 Maere, S., Heymans, K. & Kuiper, M. BiNGO: a Cytoscape plugin to assess
overrepresentation of gene ontology categories in biological networks. *Bioinformatics* **21**,
3448-3449, doi:10.1093/bioinformatics/bti551 (2005).

119 de Hoon, M. J., Imoto, S., Nolan, J. & Miyano, S. Open source clustering software.
Bioinformatics **20**, 1453-1454, doi:10.1093/bioinformatics/bth078 (2004).

120 Saldanha, A. J. Java Treeview--extensible visualization of microarray data.
Bioinformatics **20**, 3246-3248, doi:10.1093/bioinformatics/bth349 (2004).

121 Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402-408,
doi:10.1006/meth.2001.1262 (2001).

122   Thi-Kim Vu, H. *et al.* Stem cells and fluid flow drive cyst formation in an invertebrate
excretory organ. *Elife* **4**, doi:10.7554/eLife.07405 (2015).

123   Scimone, M. L., Srivastava, M., Bell, G. W. & Reddien, P. W. A regulatory program for
excretory system regeneration in planarians. *Development* **138**, 4387-4398,
doi:10.1242/dev.068098 (2011).

124   Tu, K. C. *et al.* Egr-5 is a post-mitotic regulator of planarian epidermal differentiation.
Elife **4**, e10501, doi:10.7554/eLife.10501 (2015).

125   Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of
image analysis. *Nat. Methods* **9**, 671-675, doi:10.1038/nmeth.2089 (2012).

126   L, S. & ISoMaM, S. GeneOverlap: Test and visualize gene overlaps. *R package version*
1.28.0 (2021). <https://bioconductor.org/packages/release/bioc/html/GeneOverlap.html>.
1184

1185

Figures and Figure Legends

1187

Figure 1.

 Figure 1. Transcripts encoding planarian *apolipoprotein b* orthologs are intestine-

enriched and downregulated in *nkx2.2(RNAi)* animals. (a) *apob-1* and *apob-2* mRNA (ISH)

expression in control (top) and *nkx2.2(RNAi)* planarians. **(b)** *apob-1* and *apob-2* mRNA

expression (FISH) in sagittal sections. Arrows indicate *apob*-expressing cells (likely

differentiating phagocytes) outside the intestine (dotted yellow outline) in digitally brightened

images. Scale bars: 200 μm **(a)**; 50 μm **(b)**.

1196

1197

1198

Figure 2.

Figure 2. *apob* orthologs are required for viability and neutral lipid metabolism. (a)

Simultaneous RNAi-mediated knockdown of *apob-1* and *apob-2* caused mild (yellow) and

severe (orange) depigmentation, and very severe (pink) phenotypes including head regression,

ventral curling, and lysis. Knockdown was initiated with n = 15 animals per condition. **(b)** Simple

Western (capillary-based protein analysis) lane view of extracts from control and knockdown

planarians labeled with custom anti-ApoB-1. **(c)** ApoB-1 protein expression in sagittal sections.

Expression in intestine (dotted outline) and outside the intestine (arrows) was dramatically

reduced in "very severe" *apob-1(RNAi);apob-2(RNAi)* planarians (right panels). mAb 6G10

(green) labeled body wall, dorsoventral, and visceral muscle fibers (demarcating the intestine.)

(d) Neutral lipids accumulated in the intestine and parenchyma (arrows) of "mild" *apob(RNAi)*

planarians. Oil Red O labeling, sagittal sections. **(e)** Cholesteryl esters (CE) and triglycerides

(TG), but not diacylglycerides (DG) or monoglycerides (MG) were significantly elevated in "mild"

apob(RNAi) animals. Thin layer chromatography, intensities of lipid species measured in

ImageJ and normalized to controls. Student's t test (unpaired, two-tailed); p=.0017 (CE) and

p=.0303 (TG). Error bars: mean $\pm$ S.D., n=4. **(f)** Lineage schematic indicating cell types in X1,

X2, and Xins subpopulations. **(g)** Example flow plot for uninjured animals. **(h)** Example plot for

animals 4 days post irradiation ("4 dpi"), showing ablation of cells in X1 and depletion of cells in

X2. **(i-j)** *apob* knockdown caused reduction of neutral lipids in X1 and X2 cells. One-way

ANOVA with Tukey's multiple comparisons test. Error bars: mean $\pm$ S.D., n$\geq$6 biological

replicates per condition. Scale bars: 50 μm **(c)**; 200 μm **(d,** upper panels**)**; 100 μm **(d,** lower

panels**)**.

1220

1221

1222

1223   **Figure 3.**

Figure 3. *apob* inhibition delays regeneration. (a) Schematic of RNAi treatment and analysis

regimen. **(b)** *apob* double knockdown resulted in smaller blastemas (arrows) in head and trunk

regenerates (8 dpa). **(c-d)** Differentiation of cells expressing polarity genes *wnt11-2* (posterior)

and *notum* (anterior) was delayed in *apob(RNAi) animals.* FISH; max projections of confocal

images. n=5-15 fragments per condition were scored for number of positive cells. Control and

apob-M fragments shown as phenotype examples. **(e)** Pharynx regeneration was reduced in

apob(RNAi) head regenerates (6 dpa) (*laminin* mRNA FISH, epifluorescent images). p=.0021,

control vs. *apob-S.* **(f)** CNS regeneration was reduced in *apob(RNAi)* tail regenerates (6 dpa)

(*ChAT* mRNA FISH, epifluorescent images). p=.0101, control vs. *apob-S.* **(g-h)** *apob* RNAi

disrupted intestine regeneration (6 dpa) (*ctsla* mRNA FISH, epifluorescent images). New

branches were shorter in heads **(g)**. p=.0004 (*apob-M*); p=.0016 (*apob-S*). New branches often

failed to fuse in tails **(h)**. Significance testing in **(e-h)**: one-way ANOVA with Tukey's multiple

comparisons test. Error bars: mean +/- S.D., n≥7 per condition. DAPI (blue) in **(e-h)**. Scale bars:

500 μm **(b)**; 200 μm **(c-d);** 10 μm **(c-d** insets**)**; 200 μm **(e-h)**.

1238

1239

1240

1241   **Figure 4.**

Figure 4. *apob* inhibition causes accumulation of neoblast progeny. (a-b) PhosphoHistone-

H3-S10-positive (pH3-PS10) cells were modestly elevated in *apob-M,* but not *apob-S,* uninjured

animals **(a)** and 2-day regenerates **(b)**. One-way ANOVA with Dunnett's T3 multiple

comparisons test. Error bars: mean +/- S.D., n≥12 animals/fragments per condition **(a-b)**. **(c)**

Lineage schematic indicating cell types/states in X1, X2, and Xins subpopulations. **(d)** Examples

of flow dot plots from uninjured planarians indicating percentages of cells in each gate, with X2

increase in *apob(RNAi)* animals in red. **(e-f)** Percentage of cells in X1 **(e)** and X2 **(f)** in uninjured

(left), 2-day head (middle), and 2-day trunk (right) regenerates. Arrows indicate statistically

significant increases in X2. **(g)** Examples of flow plots from uninjured planarians, 24 hr post-

irradiation, indicating percentages of cells in each gate, with X2 increase in *apob(RNAi)* animals

in red. **(h-i)** Percentage of cells in X1 **(h)** and X2 **(i)** in uninjured planarians, 24 hr (left) and 48 hr

(right) post-irradiation. Arrows indicate significant increases in X2. One-way ANOVA and

Tukey's multiple comparisons test for non-irradiated samples and X1 irradiated samples (*p*

values in **e-f, h**, see Methods). Two-way ANOVA and two-stage linear step-up procedure of

Benjamini, Krieger and Yekutieli for multiple comparisons for X2 irradiated samples (*q* values in

i). Although differences between RNAi conditions in X1 at 48 hr post-irradiation were significant,

percent of cells in this gate was negligible (<0.7% in all samples). Error bars = mean ± S.D, n≥5

biological replicates per condition. Scale bars: 200 μm **(a-b)**.

1260

1261

1262

1263   **Figure 5.**

Figure 5. *apob* RNAi preferentially dysregulates transcripts in differentiating neoblast

progeny and mature cell states. (a-b) *apob* RNAi dysregulates greater proportions of X2 **(a)**

and PIWI-LO **(b)** signature transcripts (arrows in histograms). Venn diagrams at top show

analysis scheme: percent of X1/X2/Xins **(a)** and PIWI-HI/PIWI-LO/PIWI-NEG **(b)** signature

transcripts [49] that overlap with transcripts dysregulated in *apob-M/apob-S* animals (this study)

(see also Supp. Fig. 6a-d). Histograms show percentage of signature transcripts up- and down-

regulated in *apob-M/apob-S* animals. **(c)** Schematic example (for epidermal lineage) illustrating

how transcripts dysregulated in *apob(RNAi)* animals were cross-referenced with neoblast (N),

transition state (TS), progeny (P), and mature (M) cell state subclusters from[40]. See Methods for

details. **(d-f)** t-SNE plots (digiworm.wi.mit.edu) indicate subclusters and *piwi-1* mRNA

expression for each lineage. **(g-i)** *apob* knockdown dysregulated greater proportions of

transcripts enriched in progeny ("P") and mature ("M") cell subclusters in multiple cell type

lineages. Arrows indicate less-affected transcripts enriched in neoblasts/transition state ("N/TS")

subclusters. Carets (^) **(a, b, g-i)** indicate significant gene expression overlap (*p*<0.05, Fisher's

exact test, see Source Data 2 for individual *p* values).

1279

Figure 6.

Figure 6. A putative model for ApoB function in regulating differentiation of planarian

stem cell progeny. (a) Our data support a working model in which ApoB is expressed by

phagocytes in the intestine, a primary site of LP production and secretion. **(b)** ApoB mediates

secretion of neutral lipids in LPs from intestinal phagocytes to neoblasts and their progeny. **(c)**

In the absence of ApoB, lipids accumulate in the intestine, and LP delivery to neoblasts and

their progeny is disrupted, reducing their neutral lipid content. Neoblast proliferation and renewal

are largely unaffected by reduced ApoB function. Instead, differentiation and later maturation of

most, if not all, planarian cell lineages are slowed, causing an accumulation of differentiating

progeny and a delay in regeneration of multiple organs. Box in **(a)** represents region magnified

in **(b)** wild type and **(c)** *apob* knockdown conditions. Nuclei, blue. Lipid droplets, light green.

LPs, magenta. Apical/lumenal phagocyte surface is to the left and basal/mesenchymal

phagocyte surface is to the right in **(b, c)**.

1296

Supplementary Figures and Figure Legends

Supplementary Figure 1.

Supplementary Figure 1. *apob-1* **and** *apob-2* **encode intestine-enriched ApoB orthologs.**

(a) *nkx2.2* mRNA *in situ* expression patterns in uninjured control and *nkx2.2(RNAi)* planarians.

(b) Volcano plot showing downregulation of *apob-1, apob-2,* and *nkx2.2* in *nkx2.2(RNAi)*

animals*.* An offset of 1e-300 was added to all FDR-adjusted *p* values to enable plotting of

transcripts with *p*=0. **(c)** 470 downregulated (top) and 174 upregulated (bottom) transcripts in

nkx2.2(RNAi) animals exhibited intestine enrichment in a previous study[104]. Total numbers of

dysregulated transcripts in *nkx2.2(RNAi)* samples were slightly lower than in Supplementary

Data 1, because some were undetectable in the intestine data set. **(d)** Conserved domains in

human (*H. sap.),* chicken (*G. gal.*), fly (*D. mel.*), and planarian (*S. med.*) ApoB proteins. **(e)**

Phylogenetic relationship of planarian (*Smed*) ApoB-1 and ApoB-2 (based on similarity of N-

terminal Vitellogenin domains) with closely related protein families in human (*Hom_sap*), mouse

(*Mus_mus*), chicken (*Gal_gal*), fly (*Dros_mel*), honeybee (*Apis_mel*), frog (*X_trop*), and *C.*

elegans (*C_el*). Branch support is indicated. **(f)** t-SNE plots from single cell transcriptomes[40]

showing expression of *nkx2.2, apob-1,* and *apob-2* in the intestinal lineage. All transcripts were

enriched in differentiating progeny (subclusters 0/7, orange arrows) and mature phagocytes

(subcluster 4, black arrow); *nkx2.2* was also enriched in neoblasts/transition state cells

(subcluster 1). **(g)** Double FISH showing co-expression of *apob-1, apob-2,* and *nkx2.2* mRNA*.*

(h) Double FISH showing expression of *apob-1, apob-2* in phagocytes and basal cells (*apob-1*

only), but not goblet cells. Yellow boxes indicate regions magnified in insets. Scale bars: 500 µm

(a); 200 µm **(g-h)**; 10 µm **(g-h** insets, bottom left**).**

1321

1322

1323

Supplementary Figure 2.

Supplementary Figure 2. Further characterization of *apob* expression and RNAi phenotypes. (a) *apob-1* mRNA was upregulated in head and trunk regenerates (qRT-PCR). **(b)** *apob-2* mRNA was upregulated in head and trunk regenerates (qRT-PCR). **(c)** *apob-1* and *apob-2* mRNA levels were upregulated in whole fragment regeneration RNA-Seq data (Zeng et al., 2018). **(d)** ApoB-1 protein was upregulated during regeneration (3-5 dpa) in head and trunk fragments (Simple Western "Wes" capillary-based immunoassay). 7d-I, 7 day Intact (starved and uninjured) animals. **(e-f)** Neutral lipids accumulate in *apob*(RNAi) ("mild") regenerates. Dashed line indicates approximate plane of amputation. Anterior is left. **(g-h)** Brain (G) and pharynx (H) regeneration were delayed, but not blocked, by *apob* RNAi. Arrows indicate smaller organs in *apob-M* animals relative to controls (representative images). Significance testing (qPCR and Wes): one-way ANOVA (comparisons to 0 hr controls) with Dunnett's T3 multiple comparisons test. Error bars = mean +/- S.D. DAPI (blue) in **(d-g)**. Scale bars: 200 µm **(e-f, upper panels)**; 100 µm **(e-f, lower panels)**; 100 µm **(g-h)**.

1338

1339

1340

Supplementary Figure 3.

Supplementary Figure 3. Lipoprotein receptor expression is upregulated during planarian

regeneration. (a) Schematic of domains in LDLR and related homologs. Only domains common

to Ldlr-like proteins in multiple species are shown. Clr-like EGF-like domains are shown only if

they do not overlap with Ca-binding EGF-like domains. **(b)** WISH showing upregulation of

planarian *ldlr* homologs in blastemas (black arrows) or developing pharynges (yellow arrows) at

2 and 5 dpa. **(c)** *ldlr-1* and *ldlr-2* mRNA levels were upregulated in whole fragment regeneration

RNA-Seq data (Zeng et al., 2018). **(d)** *ldlr* homologs (magenta, mRNA FISH) were expressed in

piwi-1+ neoblasts (green, mRNA FISH) as well as differentiating (*piwi-1*-negative*)* cells in brain

and pharynx (arrows) in 5-day regenerates. **(e)** Heat maps showing numerous planarian

transcripts related to lipid metabolism that were up- and down-regulated during regeneration in

whole fragment regeneration RNA-Seq data (Zeng et al., 2018). See Supplementary Data 3 for

transcript IDs and expression data. Scale bars: 200 µm **(b)**; 200 µm **(d,** left panels**)**; 20 µm **(d,**

right panels**)**.

1354

Supplementary Figure 4.

Supplementary Figure 4. Further characterization of effects of *apob* inhibition on

neoblasts and neoblast progeny. (a) Distribution of *piwi-1*-expressing neoblasts in uninjured

(7 day starved, left panels) and 7.5 dpa head and tail regenerates (right panels). **(b)** Distribution

of *tgs-1*-expressing neoblasts in uninjured (7 day starved, left panels) and 7.5 dpa head and tail

regenerates (right panels). Whole-mount FISH **(a, b)**; epifluorescent images of representative

samples. Dotted lines, approximate amputation plane. **(c)** Percentage of cells in X1 and X2 in 7

day head (left) and trunk (right) regenerates. Arrows indicate significant increases in X2. One-

way ANOVA and Dunnett's (X1, 7 day heads) or Tukey's (all others) multiple comparisons test.

Error bars = mean +/- S.D. Scale bars: 200 μm **(a-b)**.

1368

1369

Supplementary Figure 5.

Supplementary Figure 5. *apob* RNAi dysregulates transcripts involved in metabolism-

and differentiation-related processes. (a-b) Volcano plots showing significantly

downregulated (blue) and upregulated (pink) transcripts in *apob-M* **(a)** and *apob-S* **(b)** uninjured

animals. **(c)** Multi-dimensional scaling plot showing similarity of control and RNAi sample

libraries, using the biological coefficient of variation method to calculate distances between each

library based on the 500 most variable transcripts across all samples. **(d-e)** Gene Ontology

Biological Process categories over-represented among transcripts upregulated **(d)** or

downregulated **(e)** by *apob* RNAi. Numbers of transcripts dysregulated indicated in parentheses

(*apob-M/apob-S*). NA, not applicable (GO category not enriched in *apob-M* or *apob-S*). -

$\log_{10}$(FDR) on X-axis.

1381

1382

1383

Supplementary Figure 6.

Supplementary Figure 6. *apob* **RNAi preferentially dysregulates transcripts in**

differentiating neoblast progeny and mature post-mitotic cells. (a) Venn diagram showing

no overlap between X1, X2, and Xins "signature" transcripts (defined as having logFC>0

(FDR<.05) vs. transcripts in both other fractions). **(b)** Venn diagrams showing overlap between

X1, X2, and Xins signature transcripts and transcripts up or down in *apob-M* and *apob-S*.

Percentages of X1/X2/Xins dysregulated are indicated. **(c)** Venn diagram showing no overlap

between PIWI-HI, PIWI-LO, and PIWI-NEG signature transcripts (defined as having logFC>0

(FDR<.05) vs. transcripts in both other fractions). **(d)** Venn diagrams showing overlap between

PIWI-HI, PIWI-LO, and PIWI-NEG signature transcripts and transcripts up or down in *apob-M*

and *apob-S*. Percentages of PIWI-HI/PIWI-LO/PIWI-NEG dysregulated are indicated. **(e-f)** X1-

enriched **(e)** and PIWI-HI-enriched **(f)** signature transcripts were preferentially downregulated in

planarians 24 hr post-irradiation. Venn diagrams at top show analysis schemes: percent of

X1/X2/Xins **(e)** and PIWI-HI/PIWI-LO/PIWI-NEG **(f)** signature transcripts that overlap with

transcripts dysregulated in planarians 24 hr post-irradiation. Histograms show percentage of

signature transcripts up- and down-regulated in planarians 24 hr post-irradiation. Analysis of

data from [49] (see Methods).

1401

1402

1403

1404

Supplementary Figure 7.

Supplementary Figure 7. Mapping of transcripts from 24 hr post-irradiation and cell-state-enriched subpopulations to single-cell subclusters in three lineages. (a) Schematic example (for epidermal lineage) illustrating how transcripts dysregulated in 24 hr post-irradiation animals[49] were cross-referenced with neoblast (N), transition state (TS), progeny (P), and mature (M) cell state subclusters[40]. See Methods for details. **(b-d)** t-SNE plots (digiworm.wi.mit.edu) indicate subclusters and *piwi-1* mRNA expression for each lineage. **(e-g)** Cross-referencing of scRNA-Seq data[40] with RNA-Seq data from whole planarians 24 hr post-irradiation[49]. Histograms showing that irradiation preferentially caused downregulation of transcripts enriched in neoblast ("N") and transition state ("TS") subclusters in multiple cell type lineages (arrows), by contrast to the effects of *apob* RNAi (see also Fig. 5). **(h)** Cell state schematic and Venn diagrams show analysis strategy to calculate proportion of subcluster-enriched transcripts[40] also enriched in sorted planarian cell subpopulations[49]. **(i-k)** Cross-referencing of scRNA-Seq data[40] with bulk sorted cell RNA-Seq data[49]. Histograms showing that X1/PIWI-HI ("P-HI") signature transcripts were primarily enriched in neoblast ("N") and transition state ("TS") subclusters, and that Xins/PIWI-NEG ("P-NEG") signature transcripts were enriched in progeny ("P") and mature ("M") cell state subclusters. In epidermal and intestinal lineages, X2/PIWI-LOW ("P-LO") signature transcripts were most highly enriched in neoblast/transition state and progeny subclusters. In the protonephridial lineage X2/PIWI-LO transcripts were more uniformly distributed, possibly due to fewer subclusters (and/or lower resolution of transition states) in this scRNA-Seq dataset.

Supplementary Figure 8.

Supplementary Figure 8. Additional examples of dysregulation of transcripts in

differentiating neoblast progeny and mature cells by *apob* RNAi. (a) Generic scheme used

to identify overlap of transcripts enriched in specific subclusters/cell states[40] that were

dysregulated in *apob(RNAi)* planarians (this study) or 24 hr post-irradiation[49]. **(b)** Generic

scheme to calculate proportion of cell-state-enriched transcripts[40] also enriched in sorted

planarian cell subpopulations[49]. **(c-g)** t-SNE plots (digiworm.wi.mit.edu) indicate subclusters and

piwi-1 mRNA expression for each lineage. **(h-l)** *apob* knockdown dysregulated greater

proportions of transcripts in progeny ("P") and mature ("M") subclusters in multiple cell type

lineages. Arrows indicate less-affected transcripts in neoblast/transition state ("N/TS")

subclusters. **(m-q)** Cross-referencing of scRNA-Seq data[40] with RNA-Seq data from whole

planarians 24 hr post-irradiation[49]. Transcripts enriched in neoblasts/transition state subclusters

were preferentially downregulated 24 hr post-irradiation (arrows), by contrast to the effects of

apob RNAi (H-L). **(r-v)** Cross-referencing of scRNA-Seq data[40] with bulk sorted cell RNA-Seq

data[49]. Histograms showing that X1/PIWI-HI ("P-HI") signature transcripts were primarily

enriched in neoblast ("N"), transition state ("TS"), and progeny ("P") subclusters; that X2/PIWI-

LOW ("P-LO") signature transcripts were most highly enriched in progeny subclusters; and that

Xins/PIWI-NEG ("P-NEG") signature transcripts were enriched in mature ("M") cell state

subclusters. Carets (^) indicate significant gene expression overlap (**h-l**, *p*<0.05, Fisher's exact

test, see Source Data 2 for individual *p* values).

1449

1450

1451   **Supplementary Figure 9.**

Supplementary Figure 9. Gating strategy for flow cytometry experiments. (a) Forward

scatter height (FSC-H) vs. forward scatter area (FSC-A) gate to limit to singlet events. **(b)**

Propidium iodide (PI-A) vs. forward scatter area (FSC-A) gate to limit to PI negative (e.g., non-

dead) events. **(c)** Hoechst 33342 blue (y-axis) vs. Hoechst 33342 red (x-axis) gates to limit to

Hoechst-positive (e.g., ≥2C DNA content) events. Percentages of events after each gating step

are indicated.

1458

Supplementary Data

Supplementary Data 1. Up- and downregulated transcripts in *nkx2.2(RNAi)* planarians. (a) RNA-Seq data for all transcripts detected in uninjured control vs. *nkx2.2(RNAi)* planarians, together with best BLAST hits for human/mouse/zebrafish/fly/*C. elegans*. **(b)** RNA-Seq data for 766 significantly downregulated transcripts in *nkx2.2(RNAi)* relative to control planarians. **(c)** RNA-Seq data for 719 significantly upregulated transcripts in *nkx2.2(RNAi)* relative to control planarians.

Supplementary Data 2. Gene Ontology Biological Process terms enriched for transcripts up- and downregulated by *nkx2.2* RNAi. (a) Biological Process term enrichment data for 766 transcripts downregulated in *nkx2.2(RNAi)* planarians. **(b)** Biological Process term enrichment data for 719 transcripts upregulated in *nkx2.2(RNAi)* planarians.

Supplementary Data 3. Transcripts annotated with lipid-related Gene Ontology terms in whole fragment planarian regeneration transcriptome. (a) Transcripts annotated with the term "lipid transport." **(b)** Transcripts annotated with the term "triglyceride." **(c)** Transcripts annotated with the term "cholesterol." Expression data are derived from NCBI GEO GSE107874 [49] mapped to the dd_Smed_v6 transcriptome [105] (see Methods). Some transcripts may be annotated with multiple GO terms sharing the key words indicated, and therefore have multiple entries.

Supplementary Data 4. Up- and downregulated transcripts in *apob(RNAi)* "mild" and "severe" planarians. (a) RNA-Seq data for all transcripts detected in uninjured control vs. *apob-1(RNAi);apob-2(RNAi)* planarians, together with best BLAST hits for

human/mouse/zebrafish/fly/*C. elegans*. **(b)** RNA-Seq data for 842 significantly upregulated

transcripts in *apob-M(RNAi)* relative to control planarians. **(c)** RNA-Seq data for 1960

significantly upregulated transcripts in *apob-S(RNAi)* relative to control planarians. **(d)** RNA-Seq

data for 1139 significantly downregulated transcripts in *apob-M(RNAi)* relative to control

planarians. **(e)** RNA-Seq data for 2547 significantly downregulated transcripts in *apob-S(RNAi)*

relative to control planarians.

1490

Supplementary Data 5. Gene Ontology Biological Process terms enriched for transcripts

up- and downregulated by *apob* RNAi. (a) Biological Process term enrichment data for 842

transcripts upregulated in *apob-M(RNAi)* planarians. **(b)** Biological Process term enrichment

data for 1960 transcripts upregulated in *apob-S(RNAi)* planarians. **(c)** Biological Process term

enrichment data for 1139 transcripts downregulated in *apob-M(RNAi)* planarians. **(d)** Biological

Process term enrichment data for 2547 transcripts downregulated in *apob-S(RNAi)* planarians.

1497

1498 **Supplementary Data 6. Gene and transcript identities used in phylogenetic analyses and**

1499 **gene expression studies. (a)** ApoB and related homologs used for domain diagram and

1500 phylogenetic analysis. **(b)** Alignment of N-terminal Vitellogenin domains used for phylogenetic

1501 tree. **(c)** Lipoprotein receptor homologs used for domain diagram. **(d)** Transcript identities,

1502 sequences, cloning primers and vectors used in this study.

1503

1504 **Source Data 1. Differential expression analysis of neoblast subpopulations from**

1505 **GSE107874.**

1506

1507 **Source Data 2. Numerical data and statistical analyses.**