IgM$^+$ and IgM$^-$ memory B cells represent heterogeneous populations capable of producing

class-switched antibodies and germinal center B cells upon re-challenge with *P. yoelii.*

Running title: Evaluation of MBCs after *P. yoelii* infection.

Susie L. Brown[1], Jonathan J. Bauer[1], Juhyung Lee[1], Enatha Ntirandekura[1], and Jason S.

Stumhofer[1].

[1]Department of Microbiology and Immunology, University of Arkansas for Medical Sciences,

Little Rock, AR, USA.

Correspondence should be addressed to J.S.S. (jstumhofer@uams.edu), University of Arkansas

for Medical Sciences, Department of Microbiology and Immunology, Room 521A, 4301 W.

Markham St., Little Rock, AR. Phone no. – 501-526-6180.

Summary: IgM$^+$ and IgM$^-$ MBCs that co-express CD73 and CD80 can differentiate into

plasmablasts and GC B cells after re-challenge with *P. yoelii*.

Keywords: B cells, memory B cells, antibody, germinal center

Abstract

   Memory B cells (MBCs) are essential for maintaining long-term humoral immunity to infectious organisms, including *Plasmodium*. MBCs are a heterogeneous population whose function can be dictated by isotype or expression of particular surface proteins. Here, aided by antigen-specific B-cell tetramers, MBC populations were evaluated to discern their phenotype and function in response to infection with a non-lethal strain of *P. yoelii*. Infection of mice with *P. yoelii* 17X resulted in the production of two predominant MBC populations: somatically hypermutated isotype-switched (IgM$^-$) and IgM$^+$ MBCs that co-expressed CD73 and CD80 that produced antigen-specific antibodies in response to secondary infection. Re-challenge experiments indicated that IgG-producing cells dominated the recall response over the induction of IgM-secreting cells, with both populations expanding with similar timing during the secondary response. Furthermore, using ZsGreen1 expression as a surrogate for activation-induced cytidine deaminase expression alongside CD73 and CD80 co-expression, ZsGreen1$^+$CD73$^+$CD80$^+$IgM$^+$ MBCs gave rise to class-switched IgG-producing plasmablasts that induced comparable titers of Ag-specific Abs as their IgM$^-$ counterparts after adoptive transfer and infection with *P. yoelii*. Moreover, ZsGreen1$^+$CD73$^+$CD80$^+$ IgM$^+$ and IgM$^-$ MBCs differentiated into B cells with a germinal center phenotype after adoptive transfer. A third population of B cells (ZsGreen1$^-$CD73$^-$CD80$^-$IgM$^-$) that emerges after infection responded poorly to reactivation in vitro and in vivo, indicating that these cells do not represent a population of MBCs. Together these data indicated that MBC function is not defined by immunoglobulin isotype, nor does co-expression of key surface markers limit the potential fate of MBCs after recall.

Introduction

Memory B cells (MBCs) represent a population of B cells that protect the host upon antigen (Ag) re-encounter. They can differentiate into antibody-secreting cells (ASCs) upon Ag recognition.[1] is additional evidence that indicates that MBCs can re-enter the germinal center (GC) for further rounds of somatic hypermutation.[2–4] While MBCs play an essential role in infection- and vaccine-induced protective immunity, many aspects of their generation, maintenance, and function in secondary responses remain vague. However, recent findings have shed new light on the biology of this cell type.[5] Once thought of as a homogeneous cell population that expresses class-switched, somatically hypermutated B cell receptors (BCRs) generated within a GC, MBCs are instead a diverse heterogeneous population of cells with a GC-dependent and -independent origin.[6,7] Nevertheless, interpreting these recent findings on the biology of MBCs has proved to be complex. For instance, the ability of MBCs to re-enter the GC remains a point of controversy,[8–10] as this event may be restricted to IgM$^+$ MBCs and depend on the presence of persistent GCs.[3,11] Also, recent evidence indicates that a bottleneck event occurs that restricts the ability of MBCs to re-enter the GC.[12] Conversely, others suggest that isotype-switched MBCs can re-enter GCs after challenge.[13] In contrast, other evidence indicates that functionality is defined based on the presence or absence of cell surface markers, such as CD80, PD-L2, and CD73, rather than the isotype of the BCR.[2,14,15] These findings suggest functional heterogeneity within the MBC pool. However, the contradicting results are likely the product of the models utilized to evaluate MBC function and highlight the need for additional studies to assess MBC heterogeneity on functional outcome, particularly in the context of infection-induced responses.

Studies in humans link *Plasmodium*-specific antibody (Ab) responses to protection from clinical disease.[16–20] Additionally, experimental mouse models of malaria indicate that B cells

and Abs are required for clearance of primary infections and protection against re-infection.[21–25]

Generation of long-lived plasma cells and MBCs during primary infection and their subsequent

maintenance are needed for protection from secondary infections. However, the acquisition of

Ab-mediated immunity against *Plasmodium* varies based on the number of exposures during

each transmission season.[26,27] Although there is evidence in humans that MBCs and parasite-

specific Abs can be maintained for long periods after infection,[28–33] there is also evidence that a

portion of MBCs generated against malarial antigens appear to be short-lived, and parasite-

specific Ab titers drop rapidly with low re-infection rates.[34–38] A similar drop in parasite-specific

MBCs also occurs in mice over time after *P. chabaudi* infection.[39,40] These findings highlight

many of the obstacles to developing an effective malaria vaccine.

MBCs are generated after *Plasmodium* infection in mice and humans; however, knowledge

of their specificity, phenotype, origin, and affinity for malarial Ags is limited, although recent

reports have provided novel insights into this subset of cells. For instance, MBCs produced after

primary infection give rise to a faster secondary Ab response upon re-infection with *P.*

chabaudi.[39–41] This secondary response is initially dominated by the activation of IgM$^+$ MBCs

and is partially dependent on CD4$^+$ T cells.[41] Here, utilizing a *P. yoelii* infection model and B

cell tetramers specific for two blood-stage Ags (MSP-1$_{42}$ and AMA-1), we sought to evaluate the

formation and diversity of the MBC compartment and determine which subpopulations of MBCs

respond upon re-exposure to Ag. We found that MBCs that co-express CD73 and CD80 within

the IgM$^+$ and the isotype-switched (IgM$^-$) MBC pools possess somatically hypermutated BCRs.

In contrast, a population of IgM$^-$CD73$^-$CD80$^-$ B cells that also emerge after infection display a

BCR that contains few to no mutations, indicating a more germline state.

Furthermore, using a fluorescent ZsGreen1 protein as a surrogate for activation-induced

cytidine deaminase (AID) expression and $CD73^+CD80^+$ co-expression, adoptive transfer of

$CD73^+CD80^+ZsGreen1^+$ IgM^+ or IgM^- MBCs into naïve mice followed by infection with *P.*

yoelii indicated that these MBC populations possess the ability to differentiate into plasmablasts

or GC B cells. In contrast, $CD73^-CD80^-ZsGreen1^-$ IgM^- B cells responded poorly to reactivation

in vitro compared to their $ZsGreen1^+$ counterparts, and they did not display the ability to

differentiate into plasmablasts or GC B cells after adoptive transfer, indicating that they are not a

functional MBC population. Collectively, these studies suggest that somatically hypermutated *P.*

yoelii-specific $CD73^+CD80^+$ IgM^+ and IgM^- MBCs respond to reactivation through cellular

expansion and differentiation into IgG-producing plasmablasts and GC B cells.

Materials and Methods

Ethics statement

All animal studies described were done in concordance with the principles set forth by the

Animal Welfare Act and the National Institutes of Health guidelines for the care and use of

animals in biomedical research. All animal studies were reviewed and approved by the

University of Arkansas for Medical Sciences Institutional Animal Care and Use Committee

(Protocol number 3970).

Mice

Male and female B6.Cg-Gt(ROSA)26Sor$^{tm6(CAG-ZsGreen1)Hze}$/J (Ai6), B6.129P2-

Aicda$^{tm(cre)Mnz}$/J (*aicda^{cre}*), B6.SJL-*Ptprc^{a}Pepc^b*/BoyJ (CD45.1) mice on a C57BL/6J

background and WT C57BL/6J mice were obtained from The Jackson Laboratory. *Aicda^{cre}*Ai6

reporter mice were generated by breeding homozygous *aicda^{cre}* and Ai6 mice. Male BALB/c mice

were obtained from Charles River Laboratories. All animals used in this study were between 7 and

12 weeks of age and were maintained and bred in specific pathogen-free facilities at the University

of Arkansas for Medical Sciences in accordance with institutional guidelines.

Parasites

Male BALB/c mice were infected with frozen parasite stock of murine *Plasmodium yoelii* 17X

(MRA-749, BEI Resources Repository, Manassas, VA). Subsequently, 10^5 parasitized

erythrocytes derived from the passage were intraperitoneally (i.p.) injected into experimental

groups of mice to establish infection. Mice were re-challenged at ≥ 90 days post-infection (p.i.)

with 10^6 parasitized erythrocytes, i.p. Male and female mice were used for the described

experiments (except for passaging parasite stock). Primary and secondary infections were carried

out in groups of mixed-sex mice. Adoptive transfer studies were performed separately for male

and female mice with the transfer of cells from donor mice only into recipient mice of the same

sex. During the primary infection, parasitemia was determined by flow cytometry as described,[42,43]

or after re-challenge by Giemsa-stained thin-blood smears.

Flow cytometry and antibodies

Single-cell suspensions were obtained by passing the spleen through a 40 μm cell strainer followed

by lysis of erythrocytes by re-suspension in a 0.86% NH_4Cl solution. Cells were maintained in

complete RPMI (RPMI 1640 supplemented with 10% FBS, 10% sodium pyruvate, 10% non-

essential amino acids, 10% L-glutamine, 10% Pen/Strep, and 1% β-mercaptoethanol). 3-5 × 10^6

splenocytes from individual mice were aliquoted for staining into wells of a 96-well round-bottom

plate. Cells were washed in FACS buffer (1 × PBS, 0.2% BSA and 0.2% 0.5M EDTA). Fc

receptors were blocked in FACs buffer with anti-mouse CD16/32 (clone 24G2; BioXCell,

Lebanon, NH) and normal mouse (ThermoFisher cat # 10410, Carlsbad, CA) and rat IgG

(ThermoFisher cat # 10710C) in the presence of eF780 fixable viability dye (ThermoFisher cat #

65-0865-14). Surface staining was conducted with the following antibodies: CD73-PE-Cy7, GL-

7-AF488, PD-L2-PE, IgM-BV710, B220-AF700, CD80-BV605, CCR6-APC, CD38-AF700,

CD19-BV650, CD19-BV510, B220-BV650, CD73-BV605, B220-BV785, CD138-BV711,

CD80-BV650, streptavidin-BV510 (BioLegend), LIFR-PE (R&D Systems, Minneapolis, MN),

CD38-eF450, CD39-PcP-Cy5.5, CD80-FITC, IgM-APC, GL-7-eF450, IgD-biotin, CD3e-APC-

eF780, CD11b-APC-eF780, CD11c-APC-eF780, Ter119-APC-eF780 (ThermoFisher). For

samples that did not require intracellular staining, cells were fixed with 4% paraformaldehyde.

Fluorescence minus one (FMO) controls were used to set the positive gates and indicate

background staining for histogram plots. Samples were acquired on an LSRIIFortessa (BD

Biosciences, San Jose, CA) and analyzed using FlowJo version X software (BD Biosciences,

Ashland, OR).

Tetramer preparation

Recombinant His-tagged merozoite surface protein-1 (MSP-1$_{42}$) (amino acids 1398-1754;

Accession P13828) and apical merozoite antigen-1 (AMA-1) (amino acids 26-478; Accession

AAC47193) proteins from *P. yoelii* 17X were produced in *E. coli* as previously described.[44,45]

After purification and refolding the MSP-1$_{42}$ and AMA-1 proteins, the His-tag was cleaved using

a Thrombin kit (Millipore Sigma, St. Louis, MO). Confirmation of the His-tag removal was

confirmed by W. blot using an anti-His Ab (ThermoFisher). MSP-1$_{42}$ and AMA-1 were

biotinylated and tetramerized with streptavidin-PE (SA-PE; Agilent, Santa Clara, CA) as

previously described.[46] A decoy reagent to gate out the non-MSP-1$_{42}$ or AMA-1–specific B cells

was constructed by conjugating SA-PE to DyLight 650 (ThermoFisher) followed by washing and

removal of any unbound DyLight 650 and incubating with an excess of biotin as described.[46]

Tetramer staining and enrichment

Single-cell suspensions of splenocytes were prepared as described above. Following red blood

cell lysis, cells were resuspended in MACs buffer (Miltenyi BioTec, Bergisch Gladbach,

Germany), and Fc receptors were blocked with anti-mouse CD16/32 (24G2; BioXCell) in a

buffer containing normal mouse and rat IgG (ThermoFisher) for 20 minutes at 4°C. Then 10 nM

of the prepared decoy reagent was incubated with the cells for 5 minutes at room temperature.

MSP-1 or AMA-1–PE tetramers were then added at a concentration of 10 nM of SA, and the

cells were incubated at 4°C for 30 minutes. Cells were subsequently washed in MACs buffer

before incubating with anti-PE microbeads (Miltenyi) for 5 minutes at 4°C. Finally, cells were

washed again in MACs buffer prior to positive selection on an AutoMACS Pro cell separator

(Miltenyi). Recovered cells were surface stained and ran on a BD LSRIIFortessa. Collected data

were analyzed using FlowJo version X software.

BCR sequencing

MSP-1$_{42}$ and AMA-1 tetramer-specific MBCs were sorted from day 90 *P. yoelii* or naïve

aicda^{cre}Ai6 reporter mice. Cells were pooled from the spleens of three individual mice then sub-

gated based on the following parameters: live, single cells, decoy$^-$tetramer$^+$CD19$^+$B220$^+$ and

sorted as follows: CD38$^+$GL-7$^-$IgM$^+$ZsGreen1$^-$ naïve B cells, CD38$^+$GL-7$^-$IgM$^+$ZsGreen1$^+$,

CD38$^+$GL-7$^-$IgM$^-$ZsGreen1$^+$, and CD38$^+$GL-7$^-$IgM$^-$ZsGreen1$^-$ B cells. cDNA from individual B

cells was used to generate a V(D)J library using a kit from 10x Genomics (Pleasanton, CA).

Libraries were sequenced on a NextSeq 150-cycle high-output flow cell (Illumina, San Diego,

CA). The Loupe V(D)J Browser (10x Genomics) was used to identify somatic mutations in the

heavy and light chain genes of individual B cells. Individual V_H and V_L chain sequences were

cross-checked and confirmed using the NCBI's IgBLAST tool.

Adoptive transfer experiments

Aicda^{cre}Ai6 mice were sacrificed at least 70 days after infection with *P. yoelii* 17X parasites.

Spleens were harvested and enriched for B cells by depleting other immune cell populations after

labeling with biotinylated Abs against CD11b, CD3ε, Ter119, and CD11c followed by incubation

with SA microbeads (Miltenyi Biotech) on an AutoMACS Pro cell separator (Miltenyi Biotech).

Negative fractions were combined and stained to identify populations of MBCs. Live, single-cell

$CD19^+B220^+CD38^+GL7^-$ MBCs were sorted based on the following parameters: IgM^+

($CD73^+CD80^+ZsGreen1^+IgG^-IgD^{+/-}$), or IgM^- ($CD73^+CD80^+ZsGreen1^+IgM^-IgD^-$ or $CD73^-CD80^-$

$ZsGreen1^-IgM^-IgD^{+/-}$) using a BD Biosciences FACSAria. FACS sorted cells were rested in

cRPMI at 37°C for at least 30 minutes before prepping the cells for transfer. Cells were washed

two times in sterile $1 \times$ PBS and then resuspended in sterile $1 \times$ PBS. Cells were loaded into a 1

ml syringe with a 30-gauge needle. Approximately $4\text{-}8 \times 10^4$ $CD73^+CD80^+ZsGreen1^+IgM^+$,

$CD73^+CD80^+ZsGreen1^+IgM^-$ or $CD73^-CD80^-ZsGreen1^-IgM^-$ B cells were transferred via the

retro-orbital sinus into congenic $CD45.1^+$ mice. Mice were challenged with 10^5 *P. yoelii* pRBCs

i.p. the following day. Transferred cells were recovered from infected mice on day 5, 8, or 28 p.i.

by first staining with anti-CD45.2-PE and then enriched with the use of anti-PE microbeads

(Miltenyi Biotech) followed by positive selection on an autoMACs Pro Separator.

ELISAs

High binding Immunlon HBX plates (ThermoFisher) were coated with 2.5μg/ml of recombinant

P. yoelii AMA-1 or $MSP-1_{42}$ proteins in sodium bicarbonate buffer overnight at 4°C. Plates were

blocked with 5% FBS (ThermoFisher) or FetalPlex (Gemini Bio, Calabasas, CA) in PBS for one

hour at 37°C. Serum isolated from mice was initially diluted 1:50 or 1:00 and then serially diluted

1:3 down the plate leaving the last row for blocking buffer to serve as the blank control. HRP-

conjugated IgG or IgM (Southern Biotech, Birmingham, AL) was incubated on the plate for 1 h at

37°C, and SureBlue substrate (Sera care, Milford, MA) was used for detection. The reactions were

neutralized with stop solution (Sera care) before reading the plates. The plates were read on a

FLUOStar Omega plate reader (BMG Labtech, Offenburg, Germany) at an absorbance of 450 nm.

ELISpot assays

Following activation with 35% EtOH, ELISpot plates (Millipore cat # MAHAS4510) were coated

overnight at 4°C with 2.5 µg/ml of *P. yoelii* recombinant MSP-1$_{42}$ or AMA-1 protein in PBS. The

following day, the plates were blocked for 30 minutes at room temperature with cRPMI.

Afterward, 10^5 splenocytes were plated and incubated overnight at 37°C in cRPMI. Next, plates

were washed with PBS and then blocked with 5% FBS in PBS for 1 h at room temperature.

Secondary IgM-AP (SouthernBiotech cat # 1020-04) or IgG-AP (SouthernBiotech cat # 1030-04)

conjugated Abs were then incubated for 1 hour at 37°C in 5% FBS in PBS. Finally, an ELISpot

developing reagent (consisting of BCIP and NBT; Sigma Millipore) was used to visualize spots;

subsequently, an AID ELISpot reader system (Autoimmun Diagnostika, Strassberg, Germany)

was utilized to calculate the number of ASCs.

For the MBC ELISpot assays, sorted populations of MBCs (CD73$^+$CD80$^+$ZsGreen1$^+$IgG$^-$

IgD$^{+/-}$, CD73$^-$CD80$^-$ZsGreen1$^-$IgG$^-$IgD$^{+/-}$, CD73$^+$CD80$^+$ZsGreen1$^+$IgM$^-$IgD$^-$, or CD73$^-$CD80$^-$

ZsGreen1$^-$IgM$^-$IgD$^{+/-}$) were plated at a concentration of 5,000/well in 96-well round-bottom plates.

MBCs were restimulated in the presence of the TLR7 agonist R848 (1 µg/ml; Invivogen, San

Diego, CA) and human IL-2 (10 ng/ml; Peprotech, Rocky Hill, NJ) with or without splenic feeder

cells depleted of B cells (50,000/well). Splenic feeder cells incubated with R848 and human IL-2

and MBCs incubated without restimulation or added splenic feeder cells served as controls. Cells

were incubated in vitro for three days at 37°C. After three days, the cells were transferred to

ELISpot plates pre-coated with 2.5 µg/ml of *P. yoelii* recombinant MSP-1$_{42}$ and AMA-1 or 10

233 µg/ml anti-mouse Kappa (Southern Biotech). Cells were incubated for 16-24 h at 37°C. Spots were

234 visualized as described above.

236 **Statistics**

237  GraphPad Prism 9 (San Diego, CA) was used for all statistical analyses. Specific tests of

238 statistical significance are detailed in figure legends.

Results

MBCs segregate into heterogeneous populations based on phenotype after *P. yoelii* infection

To phenotypically define MBC subsets after infection, C57BL/6 wild-type (WT) mice were infected with a non-lethal strain (17X) of *P. yoelii*. Periodically over the course of the infection and after clearance (Supplemental Fig. 1A), spleens were harvested and stained for several phenotypic markers associated with MBCs,[14,15,47,48] including proteins that regulate T cell activity (CD80, PD-L2), enzymes involved in the generation of adenosine from ATP (CD73, CD39), the chemokine receptor CCR6, and the receptor for leukemia inhibitory factor (LIFR). While the specific role of these proteins on MBC function is largely unknown, a loss of these proteins through germline deletion or inhibition of ligand-receptor interactions can negatively impact the immune system.[49–56]

As several effector B cell populations emerge after infection, we used a gating strategy to distinguish between $CD138^+CD38^{lo/-}$ plasmablasts (PBs), $CD38^-GL-7^+$ germinal center (GC) B cells, $CD38^+GL-7^+$ multipotent precursors,[6] and expanding $CD38^+GL-7^-$ MBCs in the spleen. The $CD38^+GL-7^+$ multipotent precursors represent a transitional stage in B cell differentiation that precedes the formation of GC B cells and MBCs and possesses the ability to differentiate into these populations.[6] Within the $CD138^-CD38^+GL-7^-$ population B cells were further subdivided into IgM^+ and IgM^- groups (Supplemental Figure 1B). It is important to recognize that a large proportion of the $CD38^+GL-7^-IgM^+$ B cells are naïve B cells, as they are indistinguishable from MBCs based on this phenotype. Subsequent analysis of the MBC associated markers CD73 and CD80 revealed the segregation of IgM^+ and IgM^- MBCs into two predominant populations: $CD73^+CD80^+$ and $CD73^-CD80^-$ (Fig. 1A). This CD73 and CD80

expression pattern resembled that seen for 4-hydroxy-3-nitrophenylacetyl (NP)-specific MBCs

after immunization with NP-chicken γ-globulin (NP-CGG)[15] or infection with *P. chabaudi*.[41]

Shortly after infection, expansion in the IgM$^+$CD73$^+$CD80$^+$ population occurred, followed by a

decline. Then a second expansion that peaked at day 70 post-infection (p.i.) was observed.

Similarly, IgM$^-$ CD73$^+$CD80$^+$ MBC numbers peaked at day 70. While IgM$^-$CD73$^+$CD80$^+$ MBC

numbers declined after day 70, the IgM$^+$CD73$^+$CD80$^+$ MBCs were steadily maintained through

day 350 p.i. (Fig. 1B). However, the decline in IgM$^-$CD73$^+$CD80$^+$ cell numbers was not as steep

as that seen for the IgM$^-$CD73$^-$CD80$^-$ cells, which saw a >30-fold reduction in numbers from

day 70 to day 350. Although, the overall number of IgM$^-$CD73$^-$CD80$^-$ MBCs was higher in the

spleen almost a year after infection than the IgM$^-$CD73$^+$CD80$^+$ MBCs (Fig. 1B). The IgM$^+$CD73$^-$

CD80$^-$ population encompasses MBCs and naïve B cells; it is not surprising that their numbers

remained steady past day 70 p.i.

Using the expression of CD80 and CD73 to define MBC populations, the expression of the

other aforementioned markers associated with MBCs was examined. The majority of the CD73$^-$

CD80$^-$ cells, regardless of their isotype, did not express PD-L2. Also, PD-L2 expression was not

uniform amongst the CD73$^+$CD80$^+$ cells; however, a higher frequency of IgM$^-$CD73$^+$CD80$^+$

MBCs expressed PD-L2 compared to IgM$^+$CD73$^+$CD80$^+$ MBCs (Fig. 1C). Regarding the

expansion of the CD73$^+$CD80$^+$PD-L2$^+$ MBCs, IgM$^+$, and IgM$^-$ populations displayed an early

peak in expansion with the IgM$^+$ cells peaking sooner (Fig. 1D). However, these peaks were

followed by a contraction event in both populations through day 42 p.i., before each population

expanded a second time. After this second peak plateaued at day 70 for the IgM$^-$ cells, they

contracted over time while the IgM$^+$ cell numbers remained steady.

Since CD39 assists CD73 in breaking down ATP to adenosine,[57] it was not surprising that

most CD73 expressing MBCs (IgM^+ and IgM^-) co-expressed CD39. Also, a high frequency

(>50%) of the $CD73^-$ MBCs (IgM^+ and IgM^-) expressed CD39 (Fig. 1E). The IgM^+

$CD73^+CD39^+$ cells peaked early after infection, then reduced between days 21 and 35 p.i. but

thereafter, the numbers remained steady over time (Fig. 1F). In contrast, the IgM^- $CD73^+CD39^+$

cells plateaued at day 28 p.i. before declining in numbers after day 70.

The chemokine receptor CCR6 is associated with MBC precursors found within the light

zone of the GC,[58] and $CD38^+GL\text{-}7^+$ multipotent precursors express CCR6.[6] Early on, after the

infection, the per-cell expression of CCR6 was the highest for the $CD38^+GL\text{-}7^+$ multipotent

precursors (Fig. 1G). As the infection progressed, the fluorescent intensity of CCR6 decreased on

this population, correlating with the decline in this population (data not shown). Within the MBC

pool, a small proportion of IgM^+ and IgM^- MBCs showed expression of CCR6, with a higher

expression on the IgM^+ cells, which includes naïve B cells. Still, most $CD73^+$ MBCs do not

express this chemokine receptor, possibly suggesting that MBCs lose expression of this receptor

with maturation or only a subset of them express it (Fig. 1G,H). Similarly, the expression of

CCR6 was low amongst GC B cells upon their initial appearance but increased slightly over time

(Fig. 1G). Signaling through the LIFR is linked to the self-renewal and survival of murine

hematopoietic stem cells,[54,59] and it is upregulated and responsive to LIF stimulation on MBCs.[47]

Here, a sizable proportion of IgM^+ and IgM^- $CD73^+CD80^+$ MBCs express the LIFR while very

few $CD73^-CD80^-$ MBCs express this receptor after *P. yoelii* infection (Fig. 1I).

Expression of CD73 by B cells is thought to be associated with the expression of activation-

induced cytidine deaminase (AID) through either a GC-dependent or -independent manner, as

the majority of $CD73^+$ MBCs have a high rate of somatic mutations in their BCR.[6,14,60] To

confirm that CD73 and CD80 co-expression coincides with AID expression, we crossed a mouse

in which cre recombinase (*aicda*[cre/+]) replaces the endogenous AID protein with a mouse

expressing the *R26*[ZsGreen1] allele (Ai6). Thus, cre-mediated recombination in the subsequent

offspring leads to permanent ZsGreen1 fluorescent protein marking upon expression of the *Aicda*

locus (*aicda*[cre]Ai6 mice).[61,62] Regardless of isotype, the CD73$^+$CD80$^+$ MBCs favored expression

of the ZsGreen1 protein, indicating they turned on AID transcription, whereas only a small

fraction of IgM$^+$ or IgM$^-$ CD73$^-$CD80$^-$ MBCs expressed ZsGreen1 (Fig. 2). Thus, while CD73

and CD80 co-expression did largely correlate with AID expression, it was not absolute.

Furthermore, while only a small proportion of IgM$^+$ and IgM$^-$ CD73$^-$CD80$^-$ B cells expressed

ZsGreen1, sizable numbers of these cells were present in the spleen, particularly for those that

expressed IgM, though the numbers for both populations were reduced on average compared to

the CD73$^+$CD80$^+$ cells (Fig. 2C).

Another layer of diversity in MBCs was recently described in the mouse model of *P.*

chabaudi infection. A proportion of the IgM$^+$ MBCs also expresses high amounts of IgD on their

surface.[41] Indeed, here, a sizable proportion of the IgM$^+$CD73$^+$CD80$^+$ B cells were IgD$^+$,

whereas only a small proportion of the IgM$^-$CD73$^+$CD80$^+$ were IgD$^+$ (Supplemental Fig. 1C).

Further phenotypic analysis indicated that most of the IgM$^+$CD73$^-$CD80$^-$ B cells were positive

for IgD expression, suggesting that these cells are predominantly naïve B cells. Similarly, the

majority of the IgM$^-$CD73$^-$CD80$^-$ B cells expressed IgD (Supplemental Fig. 1C). Since many

IgM$^-$CD73$^-$CD80$^-$ B cells do not express AID (based on the lack of ZsGreen1 fluorescence) and

are IgM$^{lo/-}$ and IgD$^+$, these cells largely represent a population of B cells that have not undergone

class switch recombination. Thus, it is unclear if this population represents an actual subset of

MBCs. Lastly, we noted that the expression of the B-cell co-receptor component CD19 was

significantly higher on IgM$^+$ and IgM$^-$ B cells that co-expressed CD73 and CD80 compared to

those that did not express these cell surface markers (Supplemental Fig. 1C).

Together these data indicate that based on differences in the expression of several cell surface

proteins, MBCs represent a population of cells, at least phenotypically, with multiple layers of

diversity. However, the layers of diversity seem less dependent on the isotype of the BCR but

rather dependent on the experiences of the B cell itself within the environment that forms after *P.*

yoelii infection, as similar phenotypic populations are present in IgM$^+$ and IgM$^-$ MBC pools.

Distinct MSP-1–specific plasmablasts, GC B cells, and MBCs are identifiable with a B-cell

tetramer after infection.

To substantiate our findings on MBCs from examining the total infection-induced B cell

response, we utilized a B cell tetramer to examine Ag-specific B cells. Here, a phycoerythrin

(PE)-conjugated B cell tetramer consisting of the 42 kD C-terminal end of the merozoite surface

protein-1 (MSP-1) from *P. yoelii* was generated. This reagent was subsequently used to label Ag-

specific B cells ex vivo, followed by magnetic bead-based enrichment of the labeled cells. As

components of the tetramer itself, such as PE, can be bound by B cells, splenocytes were first

stained with a decoy reagent to exclude them before incubation with the MSP-1 PE tetramer.[46]

Anti-PE-specific microbeads were then used to enrich the tetramer and decoy-specific B cells,

which were subsequently labeled with Abs for analysis and quantitation by flow cytometry. After

excluding dead cells, doublets, and non-B cells, B cell populations were identified and gated

based on B220 and CD138 expression. Decoy$^-$ MSP-1$^+$ B cells were identifiable within this

population before and after *P. yoelii* infection (Fig. 3A). Tracking MSP-1–specific B cells over

one year after infection indicated that MSP-1–specific B cells rapidly expanded and peaked in

cell numbers in the spleen shortly after infection and subsequently contracted by day 35.

Thereafter a second but smaller expansion in cell numbers occurred and peaked by day 70 p.i.

before contracting again by 100 days p.i. Hereafter the MSP-1–specific B cell numbers remained

consistent over one year p.i., but the numbers were higher than those seen in naïve mice (Fig.

3B).

To corroborate our results with the MSP-1 tetramer, a similar set of experiments was

performed using a B cell tetramer composed of another blood-stage Ag, apical merozoite

antigen-1 (AMA-1) (Supplemental Fig. 2A,B). Rather than peaking at day 7 p.i. as the MSP-1$^+$ B

cells, AMA-1–specific B cells continued to expand in number throughout the infection, reaching

their peak at day 28. Following a contraction phase, a second smaller wave of expansion

occurred at day 70 before slowly contracting over time, a pattern that resembled that for the

MSP-1$^+$ B cells.

MSP-1–specific B cells possessed a naïve phenotype (CD38$^+$GL-7$^-$IgM$^+$IgD$^+$) in uninfected

mice (Fig. 3C, *data not shown*), while distinct populations of effector B cells were noted in the

spleen after infection, including plasmablasts, MBCs, and GC B cells (Fig. 3C). Also, MSP-1–

specific CD38$^+$GL-7$^+$ precursor B cells[6] were observed at days 7 and 12 after infection but were

not maintained beyond this time (Fig.3C, *data not shown*). MSP-1–specific plasmablasts peaked

in cell numbers by day 7 p.i. and contracted quickly by day 12 p.i. (Fig. 3D). By day 7, a

transition from IgM$^+$ to class-switched MSP-1–specific plasmablasts was evident (Fig. 3E,F),

likewise for AMA-1–specific plasmablasts (Supplemental Fig. 2C,D). By day 12 p.i., the class-

switched plasmablasts dominated the response (Fig. 3E,F). Whereas a distinct second wave of

plasmablast formation started at 100 days after *P. chabaudi* infection and was visible out to 265

days p.i.,[41] this degree of expansion was not apparent after *P. yoelii* infection (Fig. 3C, D). MSP-

1–specific GC B cells peaked at day 28 p.i. and were still detectable at 100 days p.i., but by day

168, these cells were no longer prevalent. CD38$^+$GL-7$^-$ B cell numbers expanded at day 7 p.i. but

contracted at day 12. Then this population went through another small expansion and contraction

between day 28 and 35 p.i. before a large expansion took place after this time resulting in peak

numbers by day 70 p.i. Afterward, this population contracted before leveling out by 24 weeks p.i.

(Fig. 3C, D). Thus, the observed gains in cell numbers most likely represent MBC production

waves, GC-independent at day 7 and GC-dependent through day 70 p.i., rather than an influx of

naïve B cells.

Tetramer-specific MBCs display a similar heterogenic phenotype as polyclonal MBCs

To determine if the MSP-1–specific MBCs phenotypically resembled the overall infection-

induced MBC population, IgM$^+$ and IgM$^-$ tetramer$^+$ MBCs were examined for expression of

CD73 and CD80 throughout infection. Similar to findings in Figure 1, MSP-1–specific IgM$^+$ and

IgM$^-$ MBCs were segregated into two populations: CD73$^+$CD80$^+$ and CD73$^-$CD80$^-$ over time

(Fig. 4A). Few MSP-1–specific IgM$^+$ and IgM$^-$ cells upregulated CD73 by day 7 p.i., instead

most of these CD38$^+$GL-7$^-$ B cells were predominantly CD73$^-$CD80$^-$ or CD80$^+$ (Fig. 4B-C, *data

not shown*). Following an initial expansion and contraction in cell numbers, MSP-1–specific

IgM$^+$ and IgM$^-$ CD73$^+$CD80$^+$ and CD73$^-$CD80$^-$ MBCs peaked at day 70, after that contracting

with time (Fig. 4B). Except for the MSP-1–specific IgM$^-$ CD73$^+$CD80$^+$ MBCs, which were still

declining, all other populations of MSP-1–specific MBCs had stabilized by one year p.i.

(Fig.4B,C). This pattern of MSP-1–specific MBC expansion and contraction was in line with that

observed for the overall MBC population after infection (Fig. 1).

Examination of AMA-1–specific MBCs revealed some differences compared to the

MSP-1–specific MBCs. While the pattern of expansion and contraction of the CD73$^+$CD80$^+$

MBCs looked similar, the frequency and number of AMA-1–specific IgM$^+$ and IgM$^-$ CD73$^-$

CD80$^-$ MBCs were lower than those observed for the MSP-1–specific cells, and their cell

numbers were still declining a year after the infection had cleared (Supplemental Fig. 2E-G).

Overall, examining the total infection-induced and Ag-specific MBC response after *P. yoelii*

infection revealed that the spleen continues to generate IgM$^+$ and IgM$^-$ MBCs well after

clearance of the infection. While CD73$^+$CD80$^+$ IgM$^+$ MBC numbers reach a point of stability

after infection, there is slow attrition in CD73$^+$CD80$^+$ IgM$^-$ MBC numbers over time in the

spleen.

Further examination of MSP-1–specific MBCs after *P. yoelii* infection indicated that

most of the CD73$^-$CD80$^-$ MBCs expressed IgD, irrespective of IgM expression. Approximately

half of the IgM$^+$ CD73$^+$CD80$^+$ MBCs were IgD$^+$. In contrast, the IgM$^-$ CD73$^+$CD80$^+$ MBCs were

primarily IgD$^-$ (Supplemental Fig. 3). Examination of ZsGreen1 expression indicated that the

MSP-1–specific IgM$^-$ CD73$^+$CD80$^+$ MBCs were uniformly ZsGreen1$^+$ with very few ZsGreen1$^-$

cells present in this population, whereas there was more variability in ZsGreen1 expression in the

IgM$^+$ CD73$^+$CD80$^+$ population (Fig. 4D-F). Unlike the total infection-induced CD73$^-$CD80$^-$ B

cells (Fig. 2C), little to no MSP-1–specific ZsGreen1$^+$ cells accumulated in either the CD73$^-$

CD80$^-$ IgM$^+$ or IgM$^-$ pools. Lastly, the MBCs that co-expressed CD73 and CD80 also displayed a

higher MFI for CD19 expression than the CD73$^-$CD80$^-$ B cells (Supplemental Fig. 3), a pattern

that matched the overall infection-induced B cell response (Supplemental Fig. 1). Hence,

phenotypically distinct populations of Ag-specific MBCs exist in the spleen after infection with

P. yoelii. However, they largely resemble the general infection-induced MBC cell populations

and those generated in other mouse models of infection and immunization.

***Plasmodium*-specific MBC subsets are genetically distinct**

As B cells progress through the GC reaction, they accumulate mutations within their

BCR, some of which increase their overall affinity for binding Ag. These high-affinity B cells

can outcompete their brethren for T-cell help to maintain their survival; thus, allowing them to

undergo further rounds of replication and somatic mutation or differentiate into long-lived MBCs

and plasma cells. To explore the degree of somatic mutations in the BCR of various MBC

populations after clearance of *P. yoelii* infection, individual heavy and light-chains of MSP-1–

specific B cells (IgM$^+$ and IgM$^-$ ZsGreen1$^+$, IgM$^-$ ZsGreen1$^-$) were sequenced from *aicda*^{cre}Ai6

reporter mice. As shown in figure 4G-H, the IgM$^-$ ZsGreen1$^-$ B cells lacked mutations within

their heavy and light chain genes. Furthermore, although these cells did not express IgM on their

surface, they contained IgM transcripts. This finding supports the idea that these B cells are not

the product of the GC and instead may represent naïve, immature, anergic, or recently activated

B cells. They have downregulated IgM on their surface but have yet to undergo class switch

recombination or somatic hypermutation, hence the lack of ZsGreen1 expression. The ZsGreen1$^+$

IgM$^-$ MSP-1–specific MBCs accumulated a significant number of somatic mutations within their

V_H and V_L chains (Fig. 4G,H). Examination of their heavy chain constant region indicated that

these MBCs favored expression of IgG2c, but a high proportion also expressed IgG1 or IgG2b

(Fig. 4I). IgG3$^+$ clones composed the smallest fraction of the ZsGreen1$^+$ IgM$^-$ clones sampled.

While most of the ZsGreen1$^+$IgM$^+$ MBCs accumulated somatic mutations in their V_H or V_L

chain genes, the number of mutations seen was significantly lower than the ZsGreen1$^+$ IgM$^-$

MSP-1–specific MBCs (Fig. 4G,H).

MSP-1–specific MBCs rapidly expand after secondary infection

A hallmark of MBCs is their ability to differentiate into ASCs upon re-encounter with Ag. To

determine how MBCs function during secondary infection, mice infected 90 days prior were re-

challenged, i.p. with 10^6 *P. yoelii* 17X pRBCs. MSP-1 tetramer$^+$ B cells were examined on the

day of re-challenge and 3, 5, and 9 days after re-infection (Fig. 5A). Expansion in the number of

MSP-1$^+$ B cells in the spleen occurred five days after re-challenge, and the numbers continued to

rise over the next four days compared to the non-challenged mice (Fig. 5B). As expected,

plasmablasts contributed to a large part of the early expansion in Ag-specific B cells after re-

challenge. Their numbers expanded significantly over the first nine days after re-infection (Fig.

5A,C). An increase in the accumulation of MSP-1–specific IgM$^+$ and IgM$^-$ plasmablasts occurred

at day 5 p.c., though more IgM$^-$ plasmablasts were prevalent at this time (Fig. 5A,D). By day 9

p.c., the MSP-1–specific IgM$^-$ plasmablasts dominated the response, though expansion was still

occurring amongst the Ag-specific IgM$^+$ plasmablasts. The MSP-1$^+$ CD38$^+$GL-7$^-$ B cells,

encompassing naïve and MBCs, and the MSP-1$^+$ GC B cells also exhibited significant expansion

by days 5 and 9 p.c. respectfully (Fig. 5C).

There was little to no change in MSP-1–specific IgM titers in the serum after re-challenge

(Fig. 5E). In contrast, the MSP-1–specific IgG titers significantly increased by day 5 p.c., and

they remained significantly elevated at day 9 (Fig. 5E). By ELISpot assay, there was only a

minimal increase in MSP-1–specific IgM-secreting cells in some mice at day 5 p.c. (Fig. 5F),

mirroring the flow cytometry results (Fig. 5D), but no further increase in numbers occurred at

day 9 p.c. Alternatively, primary infected mice showed a significant expansion in MSP-1–

specific IgM-secreting cells at day 9. In contrast, the MSP-1–specific IgG-secreting cells showed

a significant expansion by day 5 p.c., and they remained elevated at day 9 (Fig. 5F), like the flow

cytometry results (Fig. 5D). Furthermore, MSP-1–specific ASCs were not detectable in the

spleen of mice with a primary *P. yoelii* infection until day 9 p.i. Their numbers were significantly

lower than those observed in the re-challenged mice. Overall, these data indicate that the MBC

response dominates the humoral response during secondary infection with *P. yoelii*. Together

these results suggest that Ag-specific MBCs generated during primary infection with *P. yoelii*

can differentiate into ASCs during secondary infection. Moreover, the IgG-producing ASCs

dominate the secondary response to *P. yoelii* with only a marginal input by IgM-producing

ASCs, which differs from the *P. chabaudi* re-challenge model findings.[41] In comparison, we

observed an expansion in B cells with a GC phenotype by day 9 p.c. However, it is unclear if

MBCs are responsible for this increase.

IgM$^+$ and IgM$^-$ *Plasmodium*-specific MBCs can differentiate into ASCs and GC B cells

upon reactivation.

The reduced expansion in Ag-specific IgM-secreting cells shortly after re-challenge indicates

that the IgM$^+$ MBCs may be non-responsive, possibly due to the high titers of MSP-1–specific

Abs still present in the serum before re-challenge, or they have the capacity to class switch and

become IgG-secreting cells after reactivation. A third possibility is that they preferentially

proceed to the GC. To explore these possibilities, we used an adoptive transfer approach. Using

P. yoelii infected *aicda*^{cre}Ai6 reporter mice, IgM$^+$ and IgM$^-$ MBC populations expressing CD73,

CD80, and the ZsGreen1 protein were sorted and adoptively transferred into naïve congenic mice

that were subsequently infected with *P. yoelii* (Fig. 6A, Supplemental Fig. 4A,B). The functional

capacity of the MBCs was evaluated 5, 8, and 28 days after infection.

490   The vast majority of the recovered donor cells at the early time points after infection

displayed a $CD38^{lo/-}CD138^+$ phenotype indicative of plasmablasts (Fig. 6B,C), signifying that

the transferred cells could differentiate into ASCs upon reactivation. Furthermore, while a

proportion of the donor IgM^+ MBCs differentiated into plasmablasts that expressed IgM, they

began downregulating IgM expression as early as day 5 p.i., and by day 8 p.i. they were

primarily IgM^- (Fig. 6B,D), suggesting their ability to undergo class switch recombination. This

was the case as congenic mice that received IgM^+ MBCs had detectable Ag-specific IgG in their

serum by day 8 compared to congenic mice that did not receive a donor cell transfer (Fig. 6E).

Thus, IgM^+ and IgM^- MBCs primarily differentiate into ASCs, with the IgM^+ MBCs undergoing

class switch recombination upon reactivation.

500   Transferred donor cells were still identifiable at day 28 p.i. (Fig. 6F). These donor cells

primarily displayed a GC B cell phenotype $(CD38^-GL\text{-}7^+)$ at this time (Fig. 6F,G), indicating that

IgM^+ and IgM^- MBCs possess the ability to differentiate into GC B cells. Furthermore, Ag-

specific $MSP\text{-}1^+$ B cells were found amongst donor-derived GC B cells from each transferred

MBC population (Supplemental Fig. 4C,D). Ag-specific IgG is readily detectable in the serum of

congenic mice that did not receive donor MBCs at day 28 (Supplemental Fig. 4E). Therefore, the

impact of the transferred IgM^+ and IgM^- MBCs on Ab production was not distinguishable from

the endogenous Ab response at this time. Overall, the transferred $CD73^+CD80^+ZsGreen1^+$ IgM^+

and IgM^- MBCs displayed a similar function upon recall, differentiating into ASCs early after

reactivation and eventually progressing to a phenotype associated with GC B cell differentiation.

IgM⁻ CD73⁻CD80⁻ MBCs are non-responsive to reactivation

While the above studies confirmed the function of the IgM⁺ and IgM⁻ CD73⁺CD80⁺ MBCs, they

did not address the function of the IgM⁻ CD73⁻CD80⁻ B cells, the other potential MBC

population identified in mice after *P. yoelii* infection (Fig. 1). To address the role of these B cells

in a recall response, we attempted a similar approach, as shown in Figure 6. Instead, we sorted

IgM⁻ B cells that were either CD73⁺CD80⁺ZsGreen1⁺ or CD73⁻CD80⁻ZsGreen1⁻ (Supplemental

Fig. 4B) and adoptively transferred them into congenic mice that were challenged with *P. yoelii*.

We hypothesized if these B cells are indeed MBCs, then like the IgM⁻ CD73⁺CD80⁺ZsGreen1⁺

MBCs, they should favor differentiation into plasmablasts upon reactivation. However, if they

behaved identical to the CD80⁻PD-L2⁻ MBCs described in other systems,[2] they should favor GC

B cell development. In either scenario, they would potentially upregulate AID expression and

hence express the ZsGreen1 fluorescent protein.

On day 8 p.i. cells recovered from congenic mice that received the IgM⁻

CD73⁺CD80⁺ZsGreen1⁺ donor B cells maintained ZsGreen1 expression. However, no

ZsGreen1-expressing cells were recovered from the mice that received the IgM⁻ CD73⁻CD80⁻

ZsGreen1⁻ donor B cells (Fig. 7A), indicating that AID expression was not induced in the donor

B cells. Furthermore, a negligible population of CD45.2⁺ cells that matched the background

numbers recovered from the no transfer control group was isolated from the recipients of the

IgM⁻ CD73⁻CD80⁻ZsGreen1⁻ donor B cells (Fig. 7B,C). Similar results were observed at day 28,

as the number of recovered IgM⁻ CD73⁻CD80⁻ZsGreen1⁻ donor B cells matched the background

amounts detected in the no transfer controls, and no ZsGreen1⁺ cells were recovered from these

mice (Fig. 7D-F).

The significantly reduced yields seen for the cells derived from the IgM$^-$CD73$^-$CD80$^-$

ZsGreen1$^-$ B cells may indicate that these cells have a reduced fitness upon transfer into a naive

host, possibly due to competition with naïve B cells for a survival signal. Therefore, as a separate

approach to evaluate their function, IgM$^-$CD73$^-$CD80$^-$ZsGreen1$^-$ B cells, naïve, and IgM$^+$ and

IgM$^-$CD73$^+$CD80$^+$ZsGreen1$^+$ B cells were sorted from *aicda*^{cre}Ai6 reporter mice that recovered

from a *P. yoelii* infection and restimulated in vitro. The IgM$^+$ CD73$^+$CD80$^+$ZsGreen1$^+$ B cells

and naïve B cells could differentiate into IgM-producing ASCs, and some of the secreted IgM

was Ag-specific (Fig. 8A,B). As seen in vivo, a proportion of the IgM$^+$ CD73$^+$CD80$^+$ZsGreen1$^+$

B cells could undergo class switch recombination after reactivation and differentiate into IgG-

producing ASCs with some producing Ag-specific IgG (Fig. 8C,D). In contrast, the ability of

naïve B cells to undergo class switch recombination and make IgG$^+$ ASCs during this stimulation

period was reduced compared to the IgM$^+$ CD73$^+$CD80$^+$ZsGreen1$^+$ MBCs. As expected, the

ability of the IgM$^-$ CD73$^+$CD80$^+$ZsGreen1$^+$ MBCs to differentiate into IgG-producing ASCs was

superior to the other B cell populations tested (Fig. 8C,D). The IgM$^-$ CD73$^-$CD80$^-$ZsGreen1$^-$ B

cells responded poorly under the chosen in vitro stimulation conditions producing few to no IgG$^+$

ASCs. Together this data indicates that IgM$^-$ CD73$^-$CD80$^-$ZsGreen1$^-$ B cells respond poorly to

reactivation in vivo and in vitro, suggesting that these B cells are not a true population of MBCs.

Discussion

Here, we indicate that MBC function upon recall with *P. yoelii* infection is not based on the

isotype of the BCR, as IgM$^+$ and IgM$^-$ MBCs possessed the ability to differentiate into ASCs and

GC B cells. These results contrast with the findings in other models that suggested that

reactivation of IgM$^-$ MBCs led predominantly to their differentiation into plasmablasts with very

few producing GC B cells.[3,4] In both cases, mice were immunized with a non-replicating Ag.

Hence, it is possible to speculate that the initial Ab burst due to the reactivation of transferred

IgM$^-$ MBCs resulted in the secretion of enough IgG to clear the Ag in these models, thus,

preventing MBCs from differentiating into GC B cells. Support for this idea comes from the

Pape et al. study, where the reactivation of IgM$^-$ MBCs prevented the differentiation of

endogenous B cells of the adoptive recipients from forming GCs.[4] Here, we utilized a replicating

parasite to investigate MBC function, which results in the maintenance of Ag well beyond

clearance of the infection, as seen by the presence of Ag-specific GC B cells at least 70 days

after resolution of the infection (Fig. 3D). Thus, our observation of IgM$^-$ MBCs giving rise to B

cells with a GC phenotype after transfer and challenge may partly be due to the inability of the

Abs produced from MBC-derived plasmablasts to completely clear the parasite on their own.

Thereby, a sufficient Ag load is available to allow for the accumulation of GC B cells.

Re-challenge of mice with *P. chabaudi* results in the rapid expansion of IgM-secreting

plasmablasts, a proportion of which are T-cell independent, that peaks before the expansion of

class-switched ASCs.[41] While secondary infection with *P. yoelii* did induce an increase in IgM-

secreting plasmablasts, it coincided with the class-switched ASC response, with the latter

dominating the secondary response. The reduced expansion of IgM$^+$ MBCs seen here with *P.*

yoelii mirrors the results of secondary immunization with PE, which the authors attributed to the

high serum quantities of anti-PE IgG circulating in these mice before re-exposure to PE.[4] Similar

to our findings here, the adoptive transfer of PE-specific IgM$^+$ MBCs into naïve mice that were

subsequently immunized with PE showed that these cells could recognize and respond to PE in

an environment lacking circulating anti-PE Abs giving rise to IgM$^-$ ASCs and GC B cells.[4]

Hence, we speculate that the ability of IgM$^+$ MBCs to expand in response to secondary infection

with *P. chabaudi* and not *P. yoelii* is due to differences in Ag-specific Ab quantity and/or quality

in these mice at the time of re-infection. An idea supported by the differences in *P. yoelii* and *P.*

chabaudi to establish secondary infections with blood parasitemia only detectable for the latter.

P. chabaudi establishes a persistent infection after re-challenge, just as it did after primary

infection but at a lower level of parasitemia.[63]

Thus, the Abs generated in response to primary infection with *P. yoelii* are more effective

at preventing re-establishment of infection than those produced in response to *P. chabaudi*

infection. Consequently, we speculate that critical epitopes recognized by the BCR of IgM$^+$

MBCs and naïve B cells are preferentially bound by affinity-matured circulating Ag-specific Abs

or high-affinity BCRs associated with IgM$^-$ MBCs after re-infection with *P. yoelii*. In contrast,

Ag is readily available for binding and reactivating IgM$^+$ MBCs during secondary infection with

P. chabaudi. The difference in the effectiveness of the Abs can be attributed to any number of

factors, including increased titers and affinity of the Abs and differences in the production of Abs

that target specific proteins essential for RBC binding and entry.

Importantly, our in vitro results indicate that the IgM$^+$ MBCs can give rise to Ag-specific

IgM-secreting plasmablasts, meaning that these MBCs can respond to restimulation. However, in

vivo, they favored differentiation into IgM$^-$ ASCs even without circulating Ag-specific Abs.

Thus, additional factors such as inflammatory cytokines and additional signals derived from

CD4$^+$ T cells may influence differentiation and class-switching of IgM$^+$ MBCs after reactivation

in the absence of Ag-specific Abs.

Instead of favoring a model where MBC isotype expression dictates effector function, our

results favor a model in which the expression of specific cell surface proteins defines their

functional activity. In this case, co-expression of CD73$^+$ and CD80$^+$ expression by IgM$^+$ and

IgM$^-$ MBCs represented a population of MBCs that had undergone extensive somatic

hypermutation of their BCR and could differentiate into ASCs and GC B cells. The ability of the

CD73$^+$CD80$^+$ MBCs to yield IgM$^-$ ASCs during secondary responses echoed findings seen with

CD80$^+$PD-L2$^+$ MBCs.[2] However, these CD80$^+$PD-L2$^+$ MBCs did not give rise to GC B cells

regardless of whether they expressed IgM or isotype-switched Ig, which differed from the results

seen here. Instead, populations of IgM-expressing CD80$^-$PD-L2$^-$ or CD80$^-$PD-L2$^+$ MBCs

preferentially gave rise to secondary GC B cells. While abundantly found after NP-CGG

immunization,[2,4] MBCs with a CD80$^-$PD-L2$^+$ phenotype were rarely present in the spleen after *P.*

yoelii infection regardless of isotype expression, as most PD-L2$^+$ MBCs also expressed CD80

(Figure 1). Although IgM$^+$ CD73$^-$CD80$^-$ MBCs were indistinguishable from naïve B cells in this

model, the IgM$^-$ CD73$^-$CD80$^-$ B cells found here did not respond to reactivation by

differentiating into ASCs or GC B cells, indicating that these cells are not MBCs. The lack of

somatic mutations in their BCR suggests that they are naive B cells that express IgD and low

amounts of IgM. Another possibility is that these cells are anergic or in a transitional state

following activation. Regardless of their origin or function, they are not a cell type contributing

to the host's immediate protection upon re-infection.

The cell markers used by immunologists to identify MBC populations suggest that

phenotypical and functional heterogeneity exists within the MBC pool. However, their biological

function on MBCs remains to be defined. For instance, the ectoenzymes CD39 and CD73 are

surface proteins that convert ATP to AMP and adenosine. Whereas CD39 is expressed on naïve

B cells,[49] CD73 is upregulated on Ag-experienced and GC B cells,[6,60] and its expression is

maintained on a subset of MBCs as seen here and by others.[2,6,14,15,41] Furthermore, Tfh cells

express high amounts of CD73,[50] suggesting that the microenvironment of the GC is rich in

ATP. However, its role in the GC is still unknown, as germline deletion of CD73 does not impair

the GC response and MBC formation in response to NP-CGG immunization.[50] Moreover, the

mutational content of the BCR and MBC function in a recall response was not impacted by the

loss of CD73. The only significant change observed in the absence of CD73 was reduced plasma

cell accumulation in the bone marrow.[50] Therefore, it is unclear why MBCs maintain expression

of this enzyme on their cell surface. Although work with human naïve B cells and IgM[+] MBCs

showed that the ability of CD39 and CD73 to hydrolyze ATP to adenosine was required for class

switch recombination.[64] Furthermore, patients with common variable immunodeficiency with

impaired class-switched Ab responses possess B cells that are selectively deficient in CD73

expression.[64]

Signaling through the LIFR receptor on hematopoietic stem cells is required for their

survival;[54] whether it promotes the survival of MBCs is unknown. However, the LIFR is

functional on MBCs. They can respond to stimulation by LIFR agonists in vitro by inducing

phosphorylation of Stat3.[47] MBCs are long-lived and can self-renew; therefore, they possess

attributes associated with hematopoietic stem cells. Another subset of B cells that can self-

renewal is B1 B cells. Interestingly, a subset of CD25[+] peritoneal B1 B cells expressed the LIFR

and were responsive to stimulation by LIF.[65] Thus, it is quite possible that LIF-LIFR signaling in

MBCs plays a role in the maintenance of these cells, but further work is needed to address this

possibility.

CCR6 expression is known to mark MBC precursors within the GC light zone; however,

CCR6 expression is not required for MBC formation.[58] Here, we found the highest expression of

CCR6 on CD38[+]GL-7[+] B cells early after infection with *P. yoelii*. Though a small proportion of

GC B cells expressed CCR6, only a tiny subset of CD38[+]GL-7[-] B cells also expressed this

chemokine receptor, but they primarily did not co-express CD73. This result differs from the

CD73[+] MBCs found in the Peyer's patch, where all IgA-bearing MBCs are positive for CCR6,

and most express CD73 and PD-L2.[53] CCR6 expression regulated the localization of MBCs

within the Peyer's Patch and promoted their survival. The lack of CCR6 expression on the

CD73[+] MBCs found here is intriguing given the additional finding that CCR6 expression is vital

for the localization of MBCs in the spleen but not their generation or maintenance.[56]

Furthermore, altered localization of MBCs within the spleen impacts the ability of MBCs to be

recalled to their cognate Ag in the absence of CCR6 expression, suggesting that the position of

MBCs within the spleen is important for subsequent interactions with other cells such as T cells,

which are implicated in the reactivation of some MBCs.[2,66] It is possible that infection with *P.*

yoelii, as opposed to immunization with protein in adjuvant, leads to differences in CCR6

expression on MBCs observed in these models.

Although we found no evidence of rapid induction of IgM[+] MBCs into IgM-secreting

cells, this does not diminish the potential importance of these cells to the recall response in

humans. In particular, Ag-specific IgM[+] MBCs are identifiable in the blood of individuals from

malaria-endemic regions,[38,41,67,68] and they can produce IgM Abs capable of neutralizing and

promoting opsonization.[67,68] These Ag-specific IgM[+] MBCs were more abundant than their IgM[-]

counterparts, even after years of repeated *Plasmodium* infections,[67] suggesting their capacity to

play a protective role from severe disease and illustrating the inefficient acquisition of protective

IgG responses against *Plasmodium*. Although, the reduced rate of somatically mutated BCRs in

IgM[+] MBCs may make them better suited to recognize a broader range of variant *P. falciparum*

Ags. Alternatively, the multimeric shape of IgM may improve its avidity for binding Ag and thus

enhance its ability to bind and restrict parasite invasion of host cells.[68]

Lastly, as markers used to identify MBC populations continue to be defined and their

function and biological role on MBCs is deduced, the challenge will be to determine if human

MBCs similarly express these markers and if they display a similar functional role. Evidence of

the expression of many of these markers on human MBCs is beginning to emerge. For instance, a

recent study examining MBCs from patients with SAR-CoV-2 or acute *P. falciparum* malaria

showed evidence of CD39 and CD73 expression by MBCs.[69] However, the frequency of cells

expressing CD39 and CD73 was reduced in all MBC populations in the malaria patients but not

the COVID-19 patients. Given a described role for CD73 in lymphocyte homing,[70] the homing

mechanism of MBCs may be disrupted, or alternative receptors are used to promote homing into

lymphoid tissues. It is also possible that given the link between CD73 and class-switching,[64,71]

reduction in adenosine production affects the ability of these MBCs to class switch upon

reactivation. Further understanding of the functional role of proteins expressed on MBC

populations may provide valuable insight into why B cells have difficulty building protective

memory against *P. falciparum*.

Acknowledgments

Plasmodium yoelii 17X was obtained through BEI Resources, NIAID, NIH: MRA-749, contributed by D. Walliker. This work was supported by the Arkansas Biosciences Institute and National Institutes of Health (NIH) Grant AI116653 (J.S.S.) and P20-GM103625-Project 3 (to J.S.S.). In addition, the flow cytometry core is supported by the Translational Research Institute (Grant UL1-TR000039; NIH National Center for Research Resources and National Center for Advancing Translation Sciences) and the UAMS Center for Microbial Pathogenesis and Host Inflammatory Responses (Grant P20-GM103625; NIH National Institute of General Medical Sciences Centers of Biomedical Research Excellence). Special thanks to Andrea Harris for her technical assistance as part of the flow cytometry core.

Author Contributions:

Conceived and designed the experiments: JSS.

Performed the experiments: SLB, JJB, JL, EN.

Analyzed the data: SLB, JJB, JSS.

Wrote the paper: JSS.

Conflict of Interest Disclosure: The authors report no conflict of interest.

References

1. Anderson SM, Tomayko MM, Shlomchik MJ. Intrinsic properties of human and murine
memory B cells. *Immunol Rev*. 2006;211:280–294.

2. Zuccarino-Catania GV, Sadanand S, Weisel FJ, et al. CD80 and PD-L2 define functionally
distinct memory B cell subsets that are independent of antibody isotype. *Nat Immunol*.
2014;15:631 637.

3. Dogan I, Bertocci B, Vilmont V, et al. Multiple layers of B cell memory with different effector
functions. *Nat Immunol*. 2009;10:1292–1299.

4. Pape KA, Taylor JJ, Maul RW, et al. Different B Cell Populations Mediate Early and Late
Memory During an Endogenous Immune Response. *Science*. 2011;331:1203–1207.

5. Cancro MP, Tomayko MM. Memory B cells and plasma cells: The differentiative continuum
of humoral immunity. *Immunol Rev*. 2021;303:72–82.

6. Taylor JJ, Pape KA, Jenkins MK. A germinal center–independent pathway generates
unswitched memory B cells early in the primary response. *J Exp Medicine*. 2012;209:597–606.

7. Obukhanych TV, Nussenzweig MC. T-independent type II immune responses generate
memory B cells. *J Exp Medicine*. 2006;203:305–310.

8. Shlomchik MJ. Do Memory B Cells Form Secondary Germinal Centers? *Csh Perspect Biol*.
2018;10:a029405.

9. Pape KA, Jenkins MK. Do Memory B Cells Form Secondary Germinal Centers? *Csh Perspect*
Biol. 2018;10:a029116.

10. McHeyzer-Williams LJ, Dufaud C, McHeyzer-Williams MG. Do Memory B Cells Form
Secondary Germinal Centers? *Csh Perspect Biol*. 2018;10:a028878.

11. Weill J-C, Gallou SL, Hao Y, et al. Multiple players in mouse B cell memory. *Curr Opin*
Immunol. 2013;25:334–338.

12. Mesin L, Schiepers A, Ersching J, et al. Restricted Clonality and Limited Germinal Center
Reentry Characterize Memory B Cell Reactivation by Boosting. *Cell*. 2019;180:92-106.e11.

13. McHeyzer-Williams LJ, Milpied PJ, Okitsu SL, et al. Class-switched memory B cells
remodel BCRs within secondary germinal centers. *Nat Immunol*. 2015;16:296 305.

14. Anderson SM, Tomayko MM, Ahuja A, et al. New markers for murine memory B cells that
define mutated and unmutated subsets. *J Exp Medicine*. 2007;204:2103–2114.

15. Tomayko MM, Steinel NC, Anderson SM, et al. Cutting Edge: Hierarchy of Maturity of
Murine Memory B Cell Subsets. *J Immunol*. 2010;185:7146–7150.

16. Metzger WG, Okenu DMN, Cavanagh DR, et al. Serum IgG3 to the Plasmodium falciparum
merozoite surface protein 2 is strongly associated with a reduced prospective risk of malaria.
Parasite Immunol. 2003;25:307 312.

17. McGregor IA. The Passive Transfer of Human Malarial Immunity. *Am J Trop Med Hyg*.
1964;13:SUPPL 237 9.

18. Cohen S, McGregor IA, Carrington S. Gamma-globulin and acquired immunity to human
malaria. *Nature*. 1961;192:733 737.

19. Riley EM, Allen SJ, Wheeler JG, et al. Naturally acquired cellular and humoral immune
responses to the major merozoite surface antigen (PfMSP1) of Plasmodium falciparum are
associated with reduced malaria morbidity. *Parasite Immunol*. 1992;14:321 337.

20. OSIER FHA, MURUNGI LM, FEGAN G, et al. Allele-specific antibodies to Plasmodium
falciparum merozoite surface protein-2 and protection against clinical malaria. *Parasite
Immunol*. 2010;32:193–201.

21. Weid T von der, Honarvar N, Langhorne J. Gene-targeted mice lacking B cells are unable to
eliminate a blood stage malaria infection. *Journal of immunology*. 1996;156:2510 2516.

22. Grun JL, Weidanz WP. Immunity to Plasmodium chabaudi adami in the B-cell-deficient
mouse. *Nature*. 1981;290:143 145.

23. Weinbaum FI, Evans CB, Tigelaar RE. Immunity to Plasmodium Berghei yoelii in mice. I.
The course of infection in T cell and B cell deficient mice. *J Immunol Baltim Md 1950*.
2012;117:1999–2005.

24. Langhorne J, Cross C, Seixas E, et al. A role for B cells in the development of T cell helper
function in a malaria infection in mice. *Proc Natl Acad Sci U S A*. 1998;95:1730 1734.

25. Meding SJ, Langhorne J. CD4+ T cells and B cells are necessary for the transfer of protective
immunity to Plasmodium chabaudi chabaudi. *Eur J Immunol*. 1991;21:1433–8.

26. Portugal S, Pierce SK, Crompton PD. Young Lives Lost as B Cells Falter: What We Are
Learning About Antibody Responses in Malaria. *J Immunol*. 2013;190:3039–3046.

27. Ly A, Hansen DS. Development of B Cell Memory in Malaria. *Front Immunol*. 2019;10:559.

28. Struik SS, Riley EM. Does malaria suffer from lack of memory? *Immunol Rev*. 2004;201:268
290.

29. Achtman AH, C. B P, Stephens R, et al. Longevity of the immune response and memory to
blood-stage malaria infection. *Current topics in microbiology and immunology*. 2005;297:71
102.

30. Ndungu FM, Olotu A, Mwacharo J, et al. Memory B cells are a more reliable archive for
historical antimalarial responses than plasma antibodies in no-longer exposed children. *Proc*
National Acad Sci. 2012;109:8247 8252.

31. Wipasa J, Suphavilai C, Okell LC, et al. Long-lived antibody and B Cell memory responses
to the human malaria parasites, Plasmodium falciparum and Plasmodium vivax. *Plos Pathog*.
2010;6:e1000770.

32. Nogaro SI, Hafalla JC, Walther B, et al. The Breadth, but Not the Magnitude, of Circulating
Memory B Cell Responses to P. falciparum Increases with Age/Exposure in an Area of Low
Transmission. *Plos One*. 2011;6:e25582.

33. Changrob S, McHenry AM, Nyunt MH, et al. Persistence of Long-lived Memory B Cells
specific to Duffy Binding Protein in individuals exposed to Plasmodium vivax. *Sci Rep-uk*.
2018;8:8347.

34. Wykes MN, Zhou YH, Liu XQ, et al. Plasmodium yoelii can ablate vaccine-induced long-
term protection in mice. *J Immunol*. 2005;175:2510 2516.

35. Xu H, Wipasa J, Yan H, et al. The mechanism and significance of deletion of parasite-
specific CD4(+) T cells in malaria infection. *The Journal of experimental medicine*.
2002;195:881 892.

36. Weiss GE, Traore B, Kayentao K, et al. The Plasmodium falciparum-Specific Human
Memory B Cell Compartment Expands Gradually with Repeated Malaria Infections. *Plos*
Pathog. 2010;6:e1000912 13.

37. Dorfman JR, Bejon P, Ndungu FM, et al. B cell memory to 3 Plasmodium falciparum blood-
stage antigens in a malaria-endemic area. *J Infect Dis*. 2005;191:1623 1630.

38. Triller G, Scally SW, Costa G, et al. Natural Parasite Exposure Induces Protective Human
Anti-Malarial Antibodies. *Immunity*. 2017;47:1197-1209.e10.

39. Nduati EW, Ng DHL, Ndungu FM, et al. Distinct kinetics of memory B-cell and plasma-cell
responses in peripheral blood following a blood-stage Plasmodium chabaudi infection in mice.
Plos One. 2010;5:e15007.

40. Ndungu FM, Cadman ET, Coulcher J, et al. Functional memory B cells and long-lived
plasma cells are generated after a single Plasmodium chabaudi infection in mice. *Plos Pathog*.
2009;5:e1000690.

41. Krishnamurty AT, Thouvenel CD, Portugal S, et al. Somatically Hypermutated Plasmodium-
Specific IgM+ Memory B Cells Are Rapid, Plastic, Early Responders upon Malaria Rechallenge.
Immunity. 2016;45:402–414.

42. Wikenheiser DJ, Brown SL, Lee J, et al. NK1.1 Expression Defines a Population of CD4+
Effector T Cells Displaying Th1 and Tfh Cell Properties That Support Early Antibody
Production During Plasmodium yoelii Infection. *Front Immunol*. 2018;9:2277.

43. Malleret B, Claser C, Ong ASM, et al. A rapid and robust tri-color flow cytometry assay for
monitoring malaria parasite development. *Sci Rep-uk*. 2011;1:118.

44. Shi Q, Lynch MM, Romero M, et al. Enhanced Protection against Malaria by a Chimeric
Merozoite Surface Protein Vaccine ▿ . *Infect Immun*. 2007;75:1349–1358.

45. Burns JM, Flaherty PR, Romero MM, et al. Immunization against Plasmodium chabaudi
malaria using combined formulations of apical membrane antigen-1 and merozoite surface
protein-1. *Vaccine*. 2003;21:1843–1852.

46. Taylor JJ, Martinez RJ, Titcombe PJ, et al. Deletion and anergy of polyclonal B cells specific
for ubiquitous membrane-bound self-antigen. *J Exp Medicine*. 2012;209:2065–2077.

47. Tomayko MM, Anderson SM, Brayton CE, et al. Systematic Comparison of Gene
Expression between Murine Memory and Naive B Cells Demonstrates That Memory B Cells
Have Unique Signaling Capabilities. *J Immunol*. 2008;181:27 38.

48. Bhattacharya D, Cheah MT, Franco CB, et al. Transcriptional Profiling of Antigen-
Dependent Murine B Cell Differentiation and Memory Formation. *J Immunol*. 2007;179:6808–
6819.

49. Kaku H, Cheng KF, Al-Abed Y, et al. A Novel Mechanism of B Cell–Mediated Immune
Suppression through CD73 Expression and Adenosine Production. *J Immunol*. 2014;193:5904–
5913.

50. Conter LJ, Song E, Shlomchik MJ, et al. CD73 Expression Is Dynamically Regulated in the
Germinal Center and Bone Marrow Plasma Cells Are Diminished in Its Absence. *Plos One*.
2014;9:e92009 10.

51. Good-Jacobson KL, Song E, Anderson S, et al. CD80 Expression on B Cells Regulates
Murine T Follicular Helper Development, Germinal Center B Cell Survival, and Plasma Cell
Generation. *J Immunol*. 2012;188:4217–4225.

52. Good-Jacobson KL, Szumilas CG, Chen L, et al. PD-1 regulates germinal center B cell
survival and the formation and affinity of long-lived plasma cells. *Nat Immunol*. 2010;11:535.

53. Lin Y-L, Ip P-P, Liao F. CCR6 Deficiency Impairs IgA Production and Dysregulates
Antimicrobial Peptide Production, Altering the Intestinal Flora. *Front Immunol*. 2017;8:805.

54. Escary JL, Perreau J, Duménil D, et al. Leukaemia inhibitory factor is necessary for
maintenance of haematopoietic stem cells and thymocyte stimulation. *Nature*. 1993;363:361 364.

55. Salazar-Gonzalez RM, Niess JH, Zammit DJ, et al. CCR6-Mediated Dendritic Cell
Activation of Pathogen-Specific T Cells in Peyer's Patches. *Immunity*. 2006;24:623–632.

56. Elgueta R, Marks E, Nowak E, et al. CCR6-Dependent Positioning of Memory B Cells Is
Essential for Their Ability To Mount a Recall Response to Antigen. *J Immunol*. 2015;194:505–
513.

57. Faas MM, Sáez T, Vos P de. Extracellular ATP and adenosine: The Yin and Yang in immune
responses? *Mol Aspects Med*. 2017;55:9–19.

58. Suan D, Kräutler NJ, Maag JLV, et al. CCR6 Defines Memory B Cell Precursors in Mouse
and Human Germinal Centers, Revealing Light-Zone Location and Predominant Low Antigen
Affinity. *Immunity*. 2017;47:1142 1153.e4.

59. Ying Q-L, Nichols J, Chambers I, et al. BMP Induction of Id Proteins Suppresses
Differentiation and Sustains Embryonic Stem Cell Self-Renewal in Collaboration with STAT3.
Cell. 2003;115:281–292.

60. Kaji T, Ishige A, Hikida M, et al. Distinct cellular pathways select germline-encoded and
somatically mutated antibodies into immunological memory. *J Exp Medicine*. 2012;209:2079–
2097.

61. Madisen L, Zwingman TA, Sunkin SM, et al. A robust and high-throughput Cre reporting
and characterization system for the whole mouse brain. *Nat Neurosci*. 2010;13:133 140.

62. Robbiani DF, Bothmer A, Callen E, et al. AID is required for the chromosomal breaks in c-
myc that lead to c-myc/IgH translocations. *Cell*. 2008;135:1028 1038.

63. Latham LE, Wikenheiser DJ, Stumhofer JS. ICOS signaling promotes a secondary humoral
response after re-challenge with Plasmodium chabaudi chabaudi AS. *Plos Pathog*.
2020;16:e1008527.

64. Schena F, Volpi S, Faliti CE, et al. Dependence of Immunoglobulin Class Switch
Recombination in B Cells on Vesicular Release of ATP and CD73 Ectonucleotidase Activity.
Cell Reports. 2013;3:1824–1831.

65. Tumang JR, Holodick NE, Vizconde TC, et al. A CD25− Positive Population of Activated
B1 Cells Expresses LIFR and Responds to LIF. *Front Immunol*. 2011;2:6.

66. Ise W, Inoue T, McLachlan JB, et al. Memory B cells contribute to rapid Bcl6 expression by
memory follicular helper T cells. *Proc National Acad Sci*. 2014;111:11792–11797.

872 67. Hopp CS, Sekar P, Diouf A, et al. Plasmodium falciparum–specific IgM B cells dominate in
873 children, expand with malaria, and produce functional IgM. *J Exp Med*. 2021;218:e20200901.

874 68. Thouvenel CD, Fontana MF, Netland J, et al. Multimeric antibodies from antigen-specific
875 human IgM+ memory B cells restrict Plasmodium parasites. *J Exp Med*. 2021;218:e20200942.

876 69. Wildner NH, Ahmadi P, Schulte S, et al. B cell analysis in SARS-CoV-2 versus malaria:
877 Increased frequencies of plasmablasts and atypical memory B cells in COVID-19. *J Leukocyte*
878 *Biol*. 2021;109:77–90.

879 70. Takedachi M, Qu D, Ebisuno Y, et al. CD73-Generated Adenosine Restricts Lymphocyte
880 Migration into Draining Lymph Nodes. *J Immunol*. 2008;180:6288–6296.

881 71. Kim E, Ackermann C, Tóth I, et al. Down-regulation of CD73 on B cells of patients with
882 viremic HIV correlates with B cell activation and disease progression. *J Leukocyte Biol*.
883 2017;101:1263–1271.

 Figure Legends

Figure 1 MBCs form a heterogeneous population after *P. yoelii* infection. (A-G) C57BL/6

mice were infected with *P. yoelii* 17X. To examine the expression of markers associated with

MBCs, B cells were gated on live, single-cell CD138$^-$CD19$^+$B220$^+$CD38$^+$GL-7$^-$ cells before

looking at IgM expression. (A) Representative flow plots of CD73 and CD80 expression on

IgM$^+$ or IgM$^-$ B cells at various times after *P. yoelii* infection. (B) Total number of IgM$^+$ and

IgM$^-$ CD73$^+$CD80$^+$ and CD73$^-$CD80$^-$ B cells over 350 days p.i. (C) Representative flow plots of

PD-L2 expression on IgM$^+$ and IgM$^-$ CD73$^+$CD80$^+$ and CD73$^-$CD80$^-$ B cells at day 167 p.i. (D)

Total number of IgM$^+$ and IgM$^-$ CD73$^+$CD80$^+$PD-L2$^+$ B cells over 350 days p.i. (E)

Representative flow plots of CD39 and CD73 expression on IgM$^+$ and IgM$^-$ B cells at day 28 p.i.

FMO = fluorescence minus one (F) Total number of IgM$^+$ and IgM$^-$ CD73$^+$CD39$^+$ B cells over

167 days p.i. (G) Representative histogram of CCR6 expression on CD38$^+$GL-7$^+$ B cells (solid

black line), IgM$^-$ (light gray shaded) and IgM$^+$ (dotted black line) CD38$^+$GL-7$^-$ B cells at day 7

p.i. CCR6 FMO (dark gray shaded). Graph denoting CCR6 median fluorescence intensity (MFI)

on splenic B cell populations at select times after *P. yoelii* infection. The CCR6 MFI for the

CD38$^+$GL-7$^-$IgM$^-$ B cells was assigned a value of 1.0 (dotted line), and the CCR6 MFI for all

other populations was normalized to the CCR6 MFI of the CD38$^+$GL-7$^-$IgM$^-$ B cells at each time

point. (H) Representative flow plots of CCR6 expression on IgM$^+$ and IgM$^-$ CD38$^+$GL-7$^-$ B cells

at day 28 p.i. (I) Representative histogram plots of LIFR expression on IgM$^+$ and IgM$^-$

CD73$^+$CD80$^+$ (solid black line) and CD73$^-$CD80$^-$ (dotted black line) B cells at day 28 p.i. LIFR

FMO (gray shaded). Graph showing the frequency of LIFR expression on MBC populations at

day 28 p.i. (B,D,F) Each data point shows the mean ± S.E.M. with 3-6 mice per time point.

Representative of three independent experiments. (H) Error bars denote the mean ± S.E.M. Data

are representative of three independent experiments with n = 3 mice per group.

(I) Each point represents an individual mouse, and the error bars denote the mean ± S.E.M. Data

are representative of three independent experiments with n = 3 mice per group.

Figure 2 CD73⁺CD80⁺ MBCs predominantly express the ZsGreen1 protein, a surrogate

marker for AID expression. *Aicda*^{cre}Ai6 reporter mice were infected with *P. yoelii*. To examine

ZsGreen1 expression, B cells were gated on live, single-cell CD138⁻CD19⁺B220⁺CD38⁺GL-7⁻

cells before looking at IgM expression. (A) Representative flow plots of ZsGreen1 expression in

IgM⁺ and IgM⁻ CD73⁻CD80⁻ and CD73⁺CD80⁺ CD38⁺GL-7⁻ B cells at day 167 p.i. (B)

Frequency and (C) total number of ZsGreen1⁺ IgM⁺ and IgM⁻ CD73⁻CD80⁻ and CD73⁺CD80⁺

CD38⁺GL-7⁻ B cells at day 167 p.i. Each point represents an individual mouse, and the error bars

denote the mean ± S.E.M. Data are representative of three independent experiments with n = 3-4

mice per group. Significance was calculated by two-way ANOVA with post hoc Holm-Sidak's

multiple comparisons test. **** $p < 0.0001$, n.s. not significant.

Figure 3 Identification and kinetics of MSP-1–specific B cells using a B cell tetramer. (A)

Representative flow plot showing gated splenic B cell populations identified after excluding dead

cells, doublets, non-B cells (CD3⁺CD11b⁺Ter119⁺), and enrichment with MSP-1 and decoy

tetramers. Representative flow plots showing gating of MSP-1⁺decoy⁻ B cells from naïve and

C57BL/6 mice infected for 7 days with *P. yoelii*. (B) Total number of MSP-1⁺ B cells over 365

days p.i. (C) Representative flow plots identifying CD138⁺MSP-1⁺ plasmablasts (left), CD138⁻

CD19⁺MSP-1⁺ B cells (middle), and CD38⁺GL-7⁻ (memory/naïve), CD38⁺GL-7⁺ (precursor),

and CD38$^-$GL-7$^+$ (GC) B cells sub-gated from the CD138$^-$CD19$^+$MSP-1$^+$ B cells (right). Splenic

B cell populations were enriched with MSP-1 and decoy tetramers and identified after excluding

dead cells, doublets, non-B cells (CD3$^+$CD11b$^+$Ter119$^+$), and decoy$^+$ cells. (D) Total number of

MSP-1$^+$ plasmablasts, memory/naïve, and GC B cells over 365 days p.i. (E) Representative flow

plots displaying IgM and IgD expression by MSP-1$^+$ plasmablasts at day 7 and 12 p.i. (F) The

frequency of IgM$^+$ and IgM$^-$ MSP-1$^+$ plasmablasts on days 7 and 12 p.i. in the spleen. (B,D,F)

Each data point shows the mean ± S.E.M. with 6-15 mice per time point. (B,D) Data are

combined from three independent experiments. (F) Data are representative of three independent

experiments. A non-parametric Mann-Whitney t test calculated significance. **$p < 0.01$, n.s. not

significant.

Figure 4 MSP-1–specific MBCs are phenotypically and genetically distinct populations. (A)

Splenic B cell populations were enriched with MSP-1 and decoy tetramers and identified after

excluding dead cells, doublets, non-B cells (CD3$^+$CD11b$^+$Ter119$^+$), and decoy$^+$ cells.

Representative gating strategy used to identify MSP-1$^+$ IgM$^-$ and IgM$^+$ MBCs based on CD73

and CD80 staining at day 70 p.i. Total number of MSP-1$^+$ IgM$^-$ or IgM$^+$ (B) CD73$^+$CD80$^+$ or (C)

CD73$^-$CD80$^-$ MBCs over 365 days p.i. Each data point shows the mean ± S.E.M. with 7-15 mice

per time point. Data are combined from three independent experiments. (D) Representative flow

plots showing the gates used to define ZsGreen1$^+$ B cells within the MSP-1–specific IgM$^+$ and

IgM$^-$ CD73$^+$CD80$^+$ and CD73$^-$CD80$^-$ MBC pools at day 100 p.i. (E) The frequency (F) and the

total number of ZsGreen1$^+$ B cells amongst the MSP-1–specific MBC populations at day 100 p.i.

Each data point shows the mean ± S.E.M. with n = 6 mice. Data are representative of two

independent experiments. Significance was calculated by two-way ANOVA with post hoc Holm-

Sidak's multiple comparisons test. **$p < 0.01$, ***$p < 0.001$, **** $p < 0.0001$, n.s. not

significant. Total number of mutations found in the CDR1 and CDR2 of the (G) heavy and (H)

light chain gene of individual MSP-1–specific B cell clones. Numbers in parentheses represent

the number of individual B cell clones assessed and graphed for each population. Dark dotted

line in the violin plot represents the median, while the light dashed lines represent the quartiles.

Significance was calculated by one-way ANOVA Kruskal-Wallis test with post hoc Dunn's

multiple comparisons test. **** $p < 0.0001$. (I) Frequency of IgG subclasses used by

individually sequenced ZsGreen1$^+$IgM$^-$ MBC clones.

Figure 5 IgG$^+$ ASCs dominate the secondary response to *P. yoelii*. Splenic B cell populations

were enriched with MSP-1 and decoy tetramers and identified after excluding dead cells,

doublets, non-B cells (CD3$^+$CD11b$^+$Ter119$^+$), and decoy$^+$ cells. (A) Representative flow plots

identifying MSP-1–specific CD138$^+$ plasmablasts, followed by sub-gating to look at IgM and

IgD expression, and MSP-1–specific CD19$^+$ B cells before and after re-challenge of mice at day

90 p.i. with 10^6 *P. yoelii* infected RBCs. Bold numbers represent the number of cells on the plot,

while non-bold numbers indicate frequency. (B) Total number of MSP-1$^+$ B cells before and

after re-challenge. C) Total number of MSP-1$^+$ naïve/MBCs, GC B cells, and plasmablasts

before and after re-challenge. (D) Total number of IgM$^+$ and IgM$^-$ MSP-1$^+$-specific plasmablasts

before and after re-challenge. (E) Serum IgM and IgG titers specific for MSP-1$_{42}$ were

determined by ELISA. (E) Splenocytes isolated before re-challenge (day 90) and post-challenge

(days 3, 5, and 9) were used to determine the number of MSP-1-specific IgM and IgG ASCs by

ELISpot. (B,C,D,F) Each point represents an individual mouse, and the error bars denote the

mean ± S.E.M. with 5 mice per time point. Data are representative of three independent

experiments. (B,C) Significance was calculated by one-way ANOVA Kruskal-Wallis test with

post hoc Dunn's multiple comparisons test. (D-F) Significance was calculated by two-way

ANOVA with post hoc Holm-Sidak's multiple comparisons test. *$p < 0.05$, **$p < 0.01$, ***$p <$

0.001, ****$p < 0.0001$, #$p < 0.05$, ##$p < 0.01$, n.s. not significant. In **E,** the absorbance values for

each dilution for the post-challenge samples were compared against the absorbance values for

the pre-challenge samples. # represents the comparison between pre-challenge and day 5 p.c.,

and * represents the comparison between pre-challenge and day 9 p.c.

Figure 6 ZsGreen1$^+$CD73$^+$CD80$^+$IgM$^+$ MBCs predominantly differentiate into class-

switched ASCs after reactivation. (A) Experimental model. CD73$^+$CD80$^+$ZsGreen1$^+$ IgM$^+$ or

IgM$^-$ MBCs were enriched and sorted from *aicda*^{cre}Ai6 reporter mice $\geq$ 70 days after infection.

4-8 $\times$ 10^4 MBCs were transferred into naïve congenic CD45.1$^+$ mice that were challenged 24 h

later with 10^5 *P. yoelii* infected RBCs. Donor cells were recovered 5, 8, and 28 days after

challenge. (B) Representative flow plots from day 8 p.i. showing gating strategy to identify

CD45.2$^+$ZsGreen1$^+$ donor cells (left). Recovered donor cells were analyzed for their ability to

differentiate into CD138$^+$CD38$^{lo/-}$ plasmablasts (middle) and the expression of IgM on the

plasmablasts (right). Gates in middle and right plots based on the endogenous B cell population

in the recipient mice. (C) Frequency of plasmablasts found within the recovered donor cell

populations. (D) Frequency of IgM$^+$ plasmablasts found within the recovered ZsGreen1$^+$

plasmablast population. A non-parametric Mann Whitney t test calculated significance, n.s. not

significant. (E) Serum IgG titers specific for MSP-1 and AMA-1 were determined by ELISA

from day 8 infected mice. Samples were combined from three independent experiments.

Significance was calculated by two-way ANOVA with post hoc Holm-Sidak's multiple

comparisons test. The absorbance values for each dilution for the transfer recipient samples were

compared against the absorbance values for the no transfer recipient samples. * represents the

comparison between IgM$^-$ donors and no transfer and # represents the comparison between IgM$^+$

donors and the no transfer. . *$p < 0.05$, **$p < 0.01$, #$p < 0.05$, ##$p < 0.01$, n.s. not significant (F)

Representative flow plots from day 28 p.i. showing gating strategy to identify

CD45.2$^+$ZsGreen1$^+$ donor cells (left). Recovered donor cells were analyzed for their ability to

differentiate into CD38$^-$GL-7$^+$ GC B cells (middle) and the expression of IgM and IgD on the

CD38$^-$GL-7$^+$ GC B cells (right). (G) Frequency of CD38$^-$GL-7$^+$ B cells found within the

recovered ZsGreen1$^+$ population derived from the donor IgM$^+$ or IgM$^-$ MBCs. (C,D) Each point

represents an individual mouse, and the error bars denote the mean $\pm$ S.E.M. with 5-10 mice per

time point. Data are pooled from two independent experiments. (G) Each point represents an

individual mouse, and the error bars denote the mean $\pm$ S.E.M. with 6-8 mice per time point.

Data are representative of three independent experiments.

1013

Figure 7 IgM$^-$ CD73$^-$CD80$^-$ZsGreen1$^-$ B cells fail to expand and upregulate ZsGreen1

expression after adoptive transfer and challenge with *P. yoelii*. (A) Representative flow plots

showing the gating strategy to identify ZsGreen1$^+$ cells derived from donor cells at day 8 p.i. (B)

Representative flow plots showing the gating strategy to identify CD45.2$^+$ cells derived from

donor cells at day 8 p.i. (C) Total number of recovered donor cells at day 8 p.i. (D)

Representative flow plots show the gating strategy to identify ZsGreen1$^+$ cells derived from

donors at day 28 p.i. (E) Representative flow plots showing the gating strategy to identify

CD19$^+$CD45.2$^+$ B cells derived from donor cells at day 28 p.i. (F) Total number of recovered

donor cells at day 28 p.i. (C, F) Each data point shows the mean $\pm$ S.E.M. with 2-6 mice per time

point from two independent experiments. Significance was calculated by one-way ANOVA

Kruskal-Wallis test with post hoc Dunn's multiple comparisons test. $*p < 0.05$, n.s. not

significant.

1026

Figure 8 Stimulation of IgM⁻ CD80⁻CD73⁻ B cells in vitro does not induce their

differentiation into ASCs. Splenocytes from *aicda*ᶜʳᵉAi6 reporter were isolated ≥ day 90 p.i to

enrich and sort populations of MBCs based on the presence or absence of CD73, CD80, and

ZsGreen1 expression and the presence or absence of IgM⁺ expression to determine their ability

to differentiate into ASCs after restimulation in vitro. Sorted populations of MBCs were left

unstimulated or restimulated with the TLR7 agonist R848 in the presence of recombinant IL-2

with or without B cell-depleted splenic feeder cells. (A) Total IgM⁺ or (B) IgM⁺ MSP-1/AMA-

1–specific ASCs and (C) total IgG⁺ or (D) IgG⁺ MSP-1/AMA-1–specific ASCs were determined

by ELISpot. Each point represents an individual mouse, and the error bars denote the mean ±

S.E.M. Data are representative of three combined experiments with n = 9 mice. Significance was

calculated by two-way ANOVA with post hoc Holm-Sidak's multiple comparisons test. $*p <$

0.05, $***p < 0.001$, $****p < 0.0001$, n.s. not significant.

1039

Supplemental Figure 1 Gating strategy used to denote different B cell populations in the

spleen. (A) Experimental model used in figures 1-4 and supplemental figures 1-3. C57BL/6 or

*aicda*ᶜʳᵉAi6 reporter mice were infected with 10^5 *P. yoelii* infected RBCs. Splenic B cell

populations were analyzed by flow cytometry at indicated times p.i. after excluding dead cells,

doublets, and non-B cells (CD3⁺CD11b⁺Ter119⁺). Experiments with tetramer enrichment also

gated out decoy- cells before gating on B cell populations. (B) Representative flow plots from

day 70 p.i. displaying the gating strategy used to identify IgM$^+$ and IgM$^-$ MBCs based on CD73

and CD80 expression patterns. Gates based on fluorescence minus one (FMO) controls. (C)

Representative flow plots (upper) showing the gates used to distinguish IgD$^+$ and IgD$^{lo/-}$ B cells

within the IgM$^+$ and IgM$^-$ CD73$^+$CD80$^+$ and CD73$^-$CD80$^-$ MBC pools at day 90 p.i. The graph

(upper right) displays the frequency of IgD$^+$ B cells amongst MBC populations at day 90 p.i.

Representative flow plots (lower) displaying the IgD$^+$ (orange) and IgD$^{lo/-}$ (blue) MBCs from the

gates in the upper plots overlayed over the total CD38$^+$GL-7$^-$ B cells (red) to illustrate the

increase in CD19 expression by the CD73$^+$CD80$^+$ MBCs regardless of whether they express IgD.

The graph (lower right) displays the MFI for CD19 expression for each population of MBCs.

Each data point on the graphs represents an individual mouse, and the error bars denote the mean

± S.E.M. Data are representative of three independent experiments with 3-6 mice per time point.

Significance calculated by two-way ANOVA with a post hoc Holm-Sidak's multiple

comparisons test. ****$p < 0.001$.

1059

Supplemental Figure 2 Detection and kinetics of AMA-1–specific B cells after *P. yoelii*

infection. Representative flow plot showing gated AMA-1–specific B cells from naïve and day

70 infected C57BL/6 mice identified after excluding dead, doublets, and non-B cells

(CD3$^+$CD11b$^+$Ter119$^+$) and enrichment with MSP-1 and decoy tetramers. (B) The total number

of AMA-1$^+$ B cells over 467 days p.i. (C) Representative flow plots displaying IgM and IgD

expression by AMA-1$^+$ plasmablasts at day 8 p.i. (D) The frequency of IgM$^+$ and IgM$^-$ AMA-1$^+$

plasmablasts at day 8 in the spleen. (E) Representative gating strategy used to identify AMA-1$^+$

isotype switched and IgM$^+$ MBCs based on CD73 and CD80 staining at day 56 p.i. Total number

of AMA-1$^+$ IgM$^+$ or IgM$^-$ (F) CD73$^+$CD80$^+$ or (G) CD73$^-$CD80$^-$ MBCs over 467 days p.i. Each

data point shows the mean ± S.E.M. with 3-5 mice per time point from two independent

experiments. Significance determined by a non-parametric Mann Whitney t test, n.s. not

significant.

1072

Supplemental Figure 3 Expression of IgD on different MBC populations after *P. yoelii*

infection. Representative flow plots showing the gates used to distinguish IgD$^+$ and IgD$^{lo/-}$ B

cells within the MSP-1–specific IgM$^+$ and IgM$^-$ CD73$^+$CD80$^+$ and CD73$^-$CD80$^-$ MBC pools at

day 90 p.i. The left graph displays the frequency of IgD$^+$ MBCs amongst the MSP-1–specific

MBC populations at day 90 p.i. The right graph displays the MFI for CD19 expression on the

different MBC populations. Each data point represents an individual mouse and shows the mean

± S.E.M. with n = 3 mice. Data are representative of two independent experiments. Significance

determined by two-way ANOVA with a post hoc Holm-Sidak's multiple comparisons test. **$p <$

0.01, ****$p < 0.0001$.

1082

Supplemental Figure 4 Gating strategy used to sort populations of MBCs from the spleen

of *P. yoelii* infected mice. Strategy used to sort (A) IgM$^-$IgD$^-$ CD73$^+$CD80$^+$ZsGreen1$^+$ or IgM$^-$

IgD$^{+/-}$ CD73$^-$CD80$^-$ZsGreen1$^-$ B cells, or (B) IgM$^+$IgD$^{+/-}$ CD73$^+$CD80$^+$ZsGreen1$^+$ B cells from

aicda^{cre}Ai6 reporter mice infected with 10^5 *P. yoelii* infected RBCs for ≥ 70 days. Gates based

on fluorescence minus one (FMO) controls. (C) Representative flow plots from day 28 p.i.

showing the gating strategy to identify CD45.2$^+$ZsGreen1$^+$ donor cells (left). Recovered donor

cells were analyzed for their ability to differentiate into CD38$^-$GL-7$^+$ GC B cells or retain their

CD38$^+$GL-7$^-$ MBC phenotype (middle). Donor cells that displayed a GC phenotype were

subsequently analyzed to determine their ability to bind MSP-1 (right). (D) Frequency of GC B

cells that specifically bound MSP-1 within the recovered donor cells. Each data point on the

graphs represents an individual mouse, and the error bars denote the mean ± S.E.M. Data are

representative of two independent experiments with 3-5 mice. Significance calculated by two-

way ANOVA with a post hoc Holm-Sidak's multiple comparisons test, n.s., not significant. (E)

ELISA from day 28 infected mice determined serum IgG titers specific for MSP-1 and AMA-1.

1097

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Supplemental Figure 1

Supplemental Figure 2

Supplemental Figure 3

Supplemental Figure 4