A chemical screen based on an interruption of zebrafish gastrulation identifies the

HTR2C inhibitor Pizotifen as a suppressor of EMT-mediated metastasis

Joji Nakayama[1,2,3,4,§], Lora Tan[1], Boon Cher Goh[2], Shu Wang[1, 5], Hideki Makinoshima[3,6]

and Zhiyuan Gong[1,§]

[1]Department of Biological Science, National University of Singapore, Singapore

[2]Cancer Science Institute of Singapore, National University of Singapore, Singapore

[3]Tsuruoka Metabolomics Laboratory, National Cancer Center, Tsuruoka, Japan

[4]Shonai Regional Industry Promotion Center, Tsuruoka, Japan.

[5]Institute of Bioengineering and Nanotechnology, Singapore

[6]Division of Translational Research, Exploratory Oncology Research and Clinical Trial

Center, National Cancer Center, Kashiwa, Japan.

[§]Corresponding authors:

Joji Nakayama

email: Zmetastasis@gmail.com

Zhiyuan Gong

email: dbsgzy@nus.edu.sg

Keywords: Metastasis, gastrulation, EMT, Phenotyping screening, zebrafish

Total words excluding supplemental figure legends and references section: 5468

Number of figures: 5

Number of tables: 1

Number of supplemental figures: 3

Number of supplemental tables: 6

Abstract

Metastasis is responsible for approximately 90% of cancer-associated mortality but few

models exist that allow for rapid and effective screening of anti-metastasis drugs. Current

mouse models of metastasis are too expensive and time consuming to use for rapid and

high-throughput screening. Therefore, we created a unique screening concept utilizing

conserved mechanisms between zebrafish gastrulation and cancer metastasis for

identification of potential anti-metastatic drugs. We hypothesized that small chemicals

that interrupt zebrafish gastrulation might also suppress metastatic progression of cancer

cells and developed a phenotype-based chemical screen to test the hypothesis. The screen

used epiboly, the first morphogenetic movement in gastrulation, as a marker and enabled

100 chemicals to be tested in five hours. The screen tested 1280 FDA-approved drugs and

identified Pizotifen, an antagonist for serotonin receptor 2C (HTR2C) as an epiboly-

interrupting drug. Pharmacologic and genetic inhibition of HTR2C suppressed metastatic

progression in a mouse model. Blocking HTR2C with Pizotifen restored epithelial

properties to metastatic cells through inhibition of Wnt-signaling. In contrast, HTR2C

induced epithelial to mesenchymal transition (EMT) through activation of Wnt-signaling

and promoted metastatic dissemination of human cancer cells in a zebrafish

xenotransplantation model. Taken together, our concept offers a novel platform for

discovery of anti-metastasis drugs.

Introduction

Metastasis, a leading contributor to the morbidity of cancer patients, occurs through multiple steps: invasion, intravasation, extravasation, colonization, and metastatic tumor formation (1-3). The physical translocation of cancer cells is an initial step of metastasis and molecular mechanisms of it involve cell motility, the breakdown of local basement membrane, loss of cell polarity, acquisition of stem cell-like properties, and EMT (4-6). These cell-biological phenomena are also observed during vertebrate gastrulation in that evolutionarily conserved morphogenetic movements of epiboly, internalization, convergence, and extension progress (7). In zebrafish, the first morphogenetic movement, epiboly, is initiated at approximately 4 hours post fertilization (hpf) to move cells from the animal pole to eventually engulf the entire yolk cell by 10 hpf (8, 9). The embryonic cell movements are governed by the molecular mechanisms that are partially shared in metastatic cell dissemination.

At least fifty common genes were shown to be involved in both metastasis and gastrulation progression: Knockdown of these genes in Xenopus or zebrafish induced gastrulation defects; conversely, overexpression of these genes conferred metastatic potential on cancer cells and knockdown of these genes suppressed metastasis (Table S1). This evidence led us to hypothesize that small molecules that interrupt zebrafish gastrulation may suppress metastatic progression of human cancer cells.

Here we report a unique screening concept based on the hypothesis. Pizotifen, an antagonist for HTR2C, was identified from the screen as a "hit" that interrupted zebrafish gastrulation. A mouse model of metastasis confirmed pharmacological and genetic inhibition of HTR2C suppressed metastatic progression. Moreover, HTR2C induced EMT and promoted metastatic dissemination of non-metastatic cancer cells in a zebrafish xenotransplantation model. These results demonstrated that this concept could offer a novel high-throughput platform for discovery of anti-metastasis drugs and can be converted to a chemical genetic screening platform.

Results

Small molecules interrupting epiboly of zebrafish have a potential to suppress metastatic progression of human cancer cells.

Before performing a screening assay, we conducted preliminary experiments to test the hypothesis. First, we examined whether hindering the molecular function of reported genes, whose knockdown induced gastrulation defects in zebrafish, might suppress cell motility and invasion of cancer cells. We chose protein arginine methyltransferase 1 (PRMT1) and cytochrome P450 family 11 (CYP11A1), both of whose knockdown induced gastrulation defects in zebrafish but whose involvement in metastatic progression is unclear (10, 11). Elevated expression of PRMT1 and CYP11A1 were observed in highly metastatic human breast cancer cell lines and knockdown of these genes through RNA interference suppressed the motility and invasion of MDA-MB-231 cells without affecting their viability (Figure S1A-C).

Next, we conducted an inverse examination of whether chemicals which were reported to suppress metastatic dissemination of cancer cells could interrupt epiboly progression of zebrafish embryos. Niclosamide and Vinpocetine are reported to suppress metastatic progression (12, 13) (14, 15). Either Niclosamide or Vinpocetine-treated zebrafish embryos showed complete arrest at very early stages or severe delay in epiboly progression, respectively (Figure S1D).

These results suggest that epiboly could serve as a marker for this screening assay and epiboly-interrupting drugs that are identified through this screening could have the potential to suppress metastatic progression of human cancer cells.

132 FDA-approved drugs induced delayed in epiboly of zebrafish embryos

We screened 1,280 FDA, EMA or other agencies-approved drugs (Prestwick, Inc) in our zebrafish assay. The screening showed that 0.9% (12/1280) of the drugs, including Actimycin A and Tolcapone, induced severe or complete arrest of embryonic cell

movement when embryos were treated with 10μM. 5.2% (66/1280) of the drugs, such as

Dicumarol, Racecadotril, Pizotifen and S(-) Eticlopride hydrochloride, induced either

delayed epiboly or interrupted epiboly of the embryos. 93.3% (1194/1280) of drugs has

no effect on epiboly progression of the embryos. 0.6% (8/1280) of drugs induced a toxic

lethality. Epiboly progression was affected more severely when embryos were treated

with 50μM; 1.7% (22/1280) of the drugs induced severe or complete arrest of it. 8.6%

(110/1280) of the drugs induced either delayed epiboly or interrupt epiboly of the

embryos. 4.3% (55/1280) of drugs induced a toxic lethality (Figure 1A and 1B, Table S2).

Among the epiboly-interrupting drugs, several drugs have already been reported to inhibit

metastasis-related molecular mechanisms: Adrenosterone or Zardaverine, which target

HSD11β1 or PDE3 and 4, respectively, are reported to inhibit EMT (16, 17);

Racecadotril, which targets enkephalinase, is reported to confer metastatic potential on

colon cancer cell (18); and Disulfiram, which targets ALDH, is reported to confer stem-

like properties on metastatic cancer cells (19). This evidence suggests that epiboly-

interrupting drugs have the potential for suppressing metastasis of human cancer cells.

Identified drugs suppressed cell motility and invasion of human cancer cells.

It has been reported that zebrafish have orthologues to 86% of 1318 human drug targets

(20). However, it was not known whether the epiboly-interrupting drugs could suppress

metastatic dissemination of human cancer cells. To test this, we subjected the 78 epiboly-

interrupting drugs that showed a suppressor effect on epiboly progression at a 10μM

concentration to *in vitro* experiments using a human cancer cell line. The experiments

examined whether the drugs could suppress cell motility and invasion of MDA-MB-231

cells through a Boyden chamber. Before conducting the experiment, we investigated

whether these drugs might effect viability of MDA-MB-231 cells using an MTT assay.

Sixteen of the 78 drugs strongly effected cell viability at concentrations less than 1μM

and were not used in the cell motility experiments. The remaining 62 drugs were assayed

in Boyden chamber motility experiments. Twenty of the 62 drugs inhibited cell motility

and invasion of MDA-MB-231 cells without effecting cell viability. Among the 20 drugs,

Hexachlorophene and Nitazoxanide were removed since the primary targets of the drugs,

D-lactate dehydrogenase and pyruvate ferredoxin oxidoreductase are not expressed in

mammalian cells. With the exception of Ipriflavone, whose target is still unclear, the

known primary targets of the remaining 17 drugs are reported to be expressed by

mammalian cells (Figure 2A and Table 1).

We confirmed if highly metastatic human cancer cell lines expressed the genes

that code for these targets using Western blotting analyses. Among the genes, serotonin

receptor 2C (HTR2C), which is a primary target of Pizotifen, was highly expressed in

only metastatic cell lines (Figure 2B). Pizotifen suppressed cell motility and invasion of

several highly metastatic human cancer cell lines in a dose-dependent manner (Figure

2C). Similarly, Dopamine receptor D2 (DRD2), which is a primary target of S(-)

Eticlopride hydrochloride, was highly expressed in only metastatic cell lines, and the drug

suppressed cell motility and invasion of these cells in a dose-dependent manner (Figure

S2).

These results indicate that a number of the epiboly-interrupting drugs also have

suppressor effects on cell motility and invasion of highly metastatic human cancer cells.

Pizotifen suppressed metastatic dissemination of human cancer cells in a zebrafish

xenotransplantation model.

While a number of the epiboly-interrupting drugs suppressed cell motility and invasion of

human cell lines *in vitro,* it was still unclear whether the drugs could suppress metastatic

dissemination of cancer cells in vivo. Therefore, we examined whether the identified

drugs could suppress metastatic dissemination of these human cancer cells in a zebrafish

xenotransplantation model. Pizotifen was selected to test since HTR2C was overexpressed

only in highly metastatic cell lines supporting the hypothesis that it could be a novel target

for blocking metastatic dissemination of cancer cells (Figure 2B). Red fluorescent

protein-labelled MDA-MB-231 (231R) cells were injected into the duct of Cuvier of *Tg*

(kdrl:eGFP) zebrafish at 2 dpf and then maintained in the presence of either vehicle or

Pizotifen. Twenty-four hours post-injection, the numbers of fish showing metastatic

dissemination of 231R cells were measured via fluorescence microscopy. In this model,

the dissemination patterns were generally divided into three categories: (i) head

dissemination, in which disseminated 231R cells exist in the vessel of the head part; (ii)

trunk dissemination, in which the cells were observed in the vessel dilating from the trunk

to the tail; (iii) end-tail dissemination, in which the cells were observed in the vessel of

the end-tail part (16).

Three independent experiments revealed that the frequencies of fish in the drug-

treated group showing head, trunk, or end-tail dissemination, significantly decreased to

55.3±7.5%, 28.5±5.0% or 43.5±19.1% when compared with those in the vehicle-treated

group; 95.8±5.8%, 47.1±7.7% or 82.6±12.7%. Conversely, the frequency of the fish in the

drug-treated group not showing any dissemination, significantly increased to 45.4±0.5%

when compared with those in the vehicle-treated group; 2.0±2.9% (Figure 2D and Table

S3).

Similar effects were observed in another xenograft experiments using an RFP-

labelled human pancreatic cancer cell line, MIA-PaCa-2 (MP2R). In the drug treated

group, the frequencies of the fish showing head, trunk or end-tail dissemination,

significantly decreased to 15.3±6.7%, 6.2±1.3%, or 41.1±1.5%; conversely, the frequency

of the fish not showing any dissemination significantly increased to 46.3±8.9% when

compared with those in the vehicle-treated group; 74.5±11.1%, 18.9±14.9%, 77.0±9.0%,

or 17.2±0.7% (Figure S3 and Table S4).

To eliminate the possibility that the metastasis suppressing effects of Pizotifen

might result from off-target effects of the drug, we conducted validation experiments to

determine whether knockdown of HTR2C would show the same effects. Sub-clones of

231R cells that expressed shRNA targeting either LacZ or HTR2C were injected into the

fish at 2 dpf and the fish were maintained in the absence of drug. In the fish that were

inoculated with shHTR2C 231R cells, the frequencies of the fish showing head, trunk,

and end-tail dissemination, significantly decreased to 6.7±4.9%, 6.7±0.7%, or

20.0±16.5%; conversely, the frequency of the fish not showing any dissemination,

significantly increased to 80.0±4.4% when compared with those that were inoculated with

shLacZ 231R cells; 80.0±27.1%, 20.0±4.5%, 90.0± 7.7%, or 0% (Figure 2E and Table

S5).

These results indicate that pharmacological and genetic inhibition of HTR2C

suppressed metastatic dissemination of human cancer cells in vivo.

Pizotifen suppressed metastasis progression of a mouse model of metastasis.

We examined the metastasis-suppressor effect of Pizotifen in a mouse model of

metastasis (21). Luciferase-expressing 4T1 murine mammary carcinoma cells were

inoculated into the mammary fat pads (MFP) of female BALB/c mice. On day two post

inoculation, the mice were randomly assigned to two groups and one group received once

daily intraperitoneal injections of 10mg/kg Pizotifen while the other group received a

vehicle injection. Bioluminescence imaging and tumor measurement revealed that the

sizes of the primary tumors in Pizotifen-treated mice were equal to those in the vehicle-

treated mice at the time of resection on day 10 post inoculation. Immunofluorescent

staining also demonstrated that the percentage of Ki67 positive cells in the resected

primary tumors of Pizotifen-treated mice were the same as those of vehicle-treated mice

(Figure 3A-C), additionally, both groups showed less than 1% cleaved caspase 3 positive

cells (data not shown). Therefore, no anti-tumor effect of Pizotifen was observed on the

primary tumor. After 70 days from inoculation, bioluminescence imaging detected light

emitted in the lungs, livers and lymph nodes of vehicle-treated mice but not those of

Pizotifen-treated mice (Figure 3C). Vehicle-treated mice formed 5 to 50 metastatic

nodules per lung in all 10 mice analyzed; conversely, Pizotifen-treated mice (n=10)

formed 0 to 5 nodules per lung in all 10 mice analyzed (Figure 3D). Histological analyses

confirmed that metastatic lesions in the lungs were detected in all vehicle-treated mice;

conversely, they were detected in only 2 of 10 Pizotifen-treated mice and the rest of the

mice showed metastatic colony formations around the bronchiole of the lung. In addition,

4 of 10 vehicle-treated mice exhibited metastasis in the liver and the rest showed

metastatic colony formation around the portal tract of the liver. In contrast, none of 10

Pizotifen-treated mice showed liver metastases and only half of the 10 mice showed

metastatic colony formation around the portal tract (Figure 3E). These results indicate that

Pizotifen can suppress metastasis progression without affecting primary tumor growth.

To eliminate the possibility that the metastasis suppressing effects of Pizotifen

might result from off-target effects, we conducted validation experiments to determine

whether knockdown of HTR2C would show the same effects. The basic experimental

process followed the experimental design described above except that sub-clones of 4T1

cells that expressed shRNA targeting either LacZ or HTR2C were injected into the MFP

of female BALB/c mice and the mice were maintained without drug. Histological

analyses revealed that all of the mice (n=5) that were inoculated with 4T1 cells expressing

shRNA targeting LacZ showed metastases in the lungs. The mean number of metastatic

lesions in a lung was 26.4±7.8. In contrast, only one of the mice (n=5) were inoculated

with 4T1 cells expressing shRNA targeting HTR2C showed metastases in the lungs and

the rest of the mice showed metastatic colony formation around the bronchiole of the

lung. The mean number of metastatic lesions in the lung significantly decreased to 10% of

those of mice that were inoculated with 4T1 cells expressing shRNA targeting LacZ

(Figure 3F-H).

Taken together, pharmacological and genetic inhibition of HTR2C showed an

anti-metastatic effect in the 4T1 model system.

HTR2C promoted EMT-mediated metastatic dissemination of human cancer cells

Although pharmacological and genetic inhibition of HTR2C inhibited metastasis

progression, a role for HTR2C on metastatic progression has not been reported.

Therefore, we examined whether HTR2C could confer metastatic properties on poorly

metastatic cells.

Firstly, we established a stable sub-clone of MCF7 human breast cancer cells

expressing either vector control or HTR2C. Vector control expressing MCF7 cells

maintained highly organized cell-cell adhesion and cell polarity; however, HTR2C-

expressing MCF7 cells led to loss of cell-cell contact and cell scattering. The cobblestone-

like appearance of these cells was replaced by a spindle-like, fibroblastic morphology.

Western blotting and immunofluorescence (IF) analyses revealed that HTR2C-expressing

MCF7 cells showed loss of E-cadherin and EpCAM, and elevated expressions of N-

cadherin, vimentin and an EMT-inducible transcriptional factor Zeb1. Similar effects

were validated through another experiment using an immortal keratinocyte cell line,

HaCaT cells, in that HTR2C-expressing HaCaT cells also showed loss of cell-cell contact

and cell scattering with loss of epithelial markers and gain of mesenchymal markers

(Figure 4A-C). Therefore, both the morphological and molecular changes in the HTR2C-

expressing MCF7 and HaCaT cells demonstrated that these cells had undergone an EMT.

Next, we examined whether HTR2C-driven EMT could promote metastatic

dissemination of human cancer cells. Boyden chamber assay revealed that HTR2C

expressing MCF7 cells showed an increased cell motility and invasion compared with

vector control-expressing MCF7 cells in vitro (Figure 4D). Moreover, we conducted in

vivo examination of whether HTR2C expression could promote metastatic dissemination

of human cancer cells in a zebrafish xenotransplantation model. Red fluorescence protein-

labelled MCF7 cells expressing either vector control or HTR2C were injected into the

duct of Cuvier of *Tg* (*kdrl:eGFP*) zebrafish at 2 dpf. Twenty-four hours post injection, the

frequencies of the fish showing metastatic dissemination of the inoculated cells were

measured using fluorescence microscopy. In the fish that were inoculated with HTR2C

expressing MCF7 cells, the frequencies of the fish showing head, trunk, and end-tail

dissemination significantly increased to 96.7±4.7%, 68.8±6.4%, or 89.5±3.4%;

conversely, the frequency of the fish not showing any dissemination decreased to 0%

when compared with those in the fish that were inoculated with vector control expressing

MCF7 cells; 33.1±18.5%, 0%, 56.9± 4.4%, or 43% (Figure 4E and Table S6).

These results indicated that HTR2C promoted metastatic dissemination of cancer

cells though induction of EMT, and suggest that the screen can easily be converted to a

chemical genetic screening platform.

Pizotifen induced mesenchymal to epithelial transition through inhibition of Wnt-

signaling.

Finally, we elucidated the mechanism of action of how Pizotifen suppressed metastasis,

especially metastatic dissemination of cancer cells. Our results showed that HTR2C

induced EMT and that pharmacological and genetic inhibition of HTR2C suppressed

metastatic dissemination of MDA-MB-231 cells that had already transitioned to

mesenchymal-like traits via EMT. Therefore, we speculated that blocking HTR2C with

Pizotifen might inhibit the molecular mechanisms which follow EMT induction. We

firstly investigated the expressions of epithelial and mesenchymal markers in Pizotifen-

treated MDA-MB-231 cells since the activation of an EMT program needs to be transient

and reversible, and transition from a fully mesenchymal phenotype to a epithelial-

mesenchymal hybrid state or a fully epithelial phenotype is associated with malignant

phenotypes (22). IF and FACS analyses revealed 20% of Pizotifen-treated MDA-MB-231

cells restored E-cadherin expression. Also, western blotting analysis demonstrated that

4T1 primary tumors from Pizotifen-treated mice has elevated E-cadherin expression

compared with tumors from vehicle-treated mice (Figure 5A-C). However, mesenchymal

markers did not change between vehicle and Pizotifen-treated MDA-MB-231 cells (data

not shown). We further analyzed E-cadherin positive (E-cad$^+$) cells in Pizotifen-treated

MDA-MB-231 cells. The E-cad$^+$ cells showed elevated expressions of epithelial markers

KRT14 and KRT19; and decreased expression of mesenchymal makers vimentin, MMP1,

MMP3, and S100A4. Recent research reports that an EMT program needs to be transient

and reversible and that a mesenchymal phenotype in cancer cells is achieved by

constitutive ectopic expression of Zeb1. In accordance with the research, the E-cad$^+$ cells

and 4T1 primary tumors from Pizotifen-treated mice had decreased Zeb1 expression

compared with vehicle-treated cells and tumors from vehicle-treated mice (Figure 5D). In

contrast, HTR2C-expressing MCF7 and HuMEC cells expressed Zeb1 but not vehicle

control MCF7 and HuMEC cells (Figure 4C). These results indicate that HTR2C-

mediated signaling induced EMT through up-regulation of Zeb1 and blocking HTR2C

with Pizotifen induced mesenchymal to epithelial transition through downregulation of

Zeb1.

We further investigated the mechanism of action of how blocking HTR2C with

Pizotifen induced down-regulation of Zeb1. In embryogenesis, serotonin-mediated

signaling is required for Wnt-dependent specification of the superficial mesoderm during

gastrulation (23). In cancer cells, overexpression of HTR1D is associated with Wnt-

signaling which enables induction of EMT (24, 25). This evidence led to a hypothesis that

HTR2C-mediated signaling might turn on transcriptional activity of β-catenin and that

might induce up-regulation of EMT-TFs. IF analyses revealed β-catenin was accumulated

in the nucleus of HTR2C-expressing MCF7 cells but it was located in the cytoplasm of

vector control-expressing cells (Figure 5E). Nuclear accumulation of β-catenin in

HTR2C-expressing MCF7 cells was confirmed by western blot (Figure 5F). In contrast,

Pizotifen-treated MDA-MB-231 cells showed β-catenin located in the cytoplasm of the

cells. Vehicle-treated cells showed β-catenin accumulated in the nucleus of the cells.

(Figure 5G), and western blotting analysis confirmed that it was located in the cytoplasm

of Pizotifen-treated MDA-MB-231 cells (Figure 5H). Furthermore,

immunohistochemistry and western blotting analyses showed that β-catenin accumulated

in the nucleus, and phospho-GSKβ and Zeb1 expression were decreased in 4T1 primary

tumors from Pizotifen-treated mice compared with vehicle-treated mice (Figure 5C).

These results indicated that HTR2C would regulate transcriptional activity of β-catenin

and Pizotifen could inhibit it.

Taken together, we conclude that blocking HTR2C with Pizotifen restored

epithelial properties to metastatic cells (MDA-MB-231 and 4T1 cells) through a decrease

of transcriptional activity of β-catenin and that suppressed metastatic progression of the

cells.

  Figures and figure legends

  Figure 1

  Figure 1. A chemical screen for identification of epiboly-interrupting drugs.

  (A) Cumulative results of the chemical screen in which each drug was used at either

  10μM (left) or 50μM (right) concentrations. 1,280 FDA, EMA or other agencies-

  approved drugs were subjected to this screening. Positive "hit" drugs were those that

  interrupted epiboly progression. (B) Representative samples of the embryos that were

  treated with indicated drugs.

339   **Figure 2**
340

Figure 2. Pizotifen, one of epiboly-interrupting drugs suppressed metastatic

dissemination of human cancer cells lines in vivo and vitro.

(A) Effect of the epiboly-interrupting drugs on cell motility and invasion of MBA-MB-

231 cells. MBA-MB-231 cells were treated with vehicle or each of the epiboly-

interrupting drugs and then subjected to Boyden chamber assays. Fetal bovine serum

(1%v/v) was used as the chemoattractant in both assays. Each experiment was performed

at least twice. (B) Western blot analysis of HTR2C levels (top) in a non-metastatic human

cancer cell line, MCF7 (breast) and highly metastatic human cancer cell lines, MDA-MB-

231 (breast), MDA-MB-435 (melanoma), PC9 (lung), MIA-PaCa2 (pancreas), PC3

(prostate) and SW620 (colon); GAPDH loading control is shown (bottom). (C) Effect of

pizotifen on cell motility and invasion of MBA-MB-231, MDA-MB-435 and PC9 cells.

Either vehicle or pizotifen treated the cells were subjected to Boyden chamber assays.

Fetal bovine serum (1%v/v) was used as the chemoattractant in both assays. Each

experiment was performed at least twice. (D) and (E) Representative images of

dissemination of 231R, shLacZ 231R or shHTR2C 231R cells in zebrafish

xenotransplantation model. The fish larvae that were inoculated with 231R cells, were

treated with either vehicle (top left) or the drug (lower left) (D). The fish larvae that were

inoculated with either shLacZ 231R or shHTR2C 231R cells (lower left) (E). White

arrows head indicate disseminated 231R cells. The images were shown in 4x

magnification. Scale bar, 100μm. The mean frequencies of the fish showing head, trunk,

or end-tail dissemination were counted (graph on right). Each value is indicated as the

mean ± SEM of two independent experiments. Statistical analysis was determined by

Student's t test. See also Figure S2 and S3, Table S3-S5.

Figure 3

Figure 3. Pizotifen suppressed metastatic progression in a mouse model of

metastasis.

(A) Mean volumes (n=10 per group) of 4T1 primary tumors formed in the mammary fat

pad of either vehicle or Pizotifen-treated mice at day 10 post injection. (B) Ki67

expression level in 4T1 primary tumors formed in the mammary fat pad of either vehicle

or Pizotifen-treated mice at day 10 post injection. The mean expression levels of Ki67

(n=10 mice per group) were determined and were calculated as the mean ration of Ki67

positive cells to DAPI area. (C) Representative images of primary tumors on day 10 post

injection (top panels) and metastatic burden on day 70 post injection (bottom panels)

taken using an IVIS Imaging System. (D) Representative images of the lungs from either

vehicle (top) or Pizotifen-treated mice (bottom) at 70 days past tumor inoculation.

Number of metastatic nodules in the lung of either vehicle or Pizotifen-treated mice

(right). (E) Representative H&E staining of the lung (top) and liver (bottom) from either

vehicle or Pizotifen-treated mice. (F) The mean number of metastatic lesions in step

sections of the lungs from the mice that were inoculated with 4T1-12B cells expressing

shRNA targeting for either LacZ or HTR2C. (G) Representative H&E staining of the lung

and liver from the mice that were inoculated with 4T1-12B cells expressing shRNA

targeting for either LacZ or HTR2C. Each value is indicated as the mean ± SEM.

Statistical analysis was determined by Student's *t* test.

Figure 4

Figure 4. HTR2C induced EMT-mediated metastatic dissemination of human cancer cells.

(A) The morphologies of the MCF7 and HaCaT cells expressing either the control vector or HTR2C were revealed by phase contrast microscopy. (B) Immunofluorescence staining of E-cadherin, EpCAM, Vimentin, and N-cadherin expressions in the MCF7 cells from Figure 4A. (C) Expression of E-cadherin, EpCAM, Vimentin, N-cadherin, and HTR2C was examined by western-blotting in the MCF7 and HaCaT cells; GAPDH loading control is shown (bottom). (D) Effect of HTR2C on cell motility and invasion of MCF7 cells. MCF7 cells were subjected to Boyden chamber assays. Fetal bovine serum (1%v/v) was used as the chemoattractant in both assays. Each experiment was performed at least twice. (E) Representative images of dissemination patterns of MCF7 cells expressing either the control vector (top left) or HTR2C (lower left) in a zebrafish xenotransplantation model. White arrows head indicate disseminated MCF7 cells. The mean frequencies of the fish showing head, trunk, or end-tail dissemination tabulated (right). Each value is indicated as the mean ± SEM of two independent experiments. Statistical analysis was determined by Student's *t* test.

See also Table S6.

Figure 5

Figure 5. Pizotifen restored mesenchymal-like traits of MDA-MB-231 cells into epithelial traits through blocking nuclear accumulation of β-catenin.

(A) IF staining of E-cadherin in either vehicle or Pizotifen-treated MDA-MB-231 cells.

(B) Surface expression of E-cadherin in either vehicle (black) or Pizotifen (red)-treated

MDA-MB-231 cells by FACS analysis. Non-stained controls are shown in gray. (C)

Protein expressions levels of E-cadherin, Zeb1, and β-catenin in the cytoplasm and

nucleus of 4T1 primary tumors from either vehicle or Pizotifen-treated mice are shown;

Luciferase, Histone H3, and β-tubulin are used as loading control for whole cell, nuclear

or cytoplasmic lysate, respectively. (D) Protein expression levels of epithelial and

mesenchymal markers and Zeb1 in either vehicle or Pizotifen-treated MDA-MB-231 cells

or E-cadherin positive (E-cad[+]) cells in Pizotifen-treated MDA-MB-231 cells are shown.

(E) Immunofluorescence staining of β-catenin in the MCF7 cells expressing either vector

control or HTR2C. (F) Expressions of β-catenin in the cytoplasm and nucleus of MCF7

424   cells. (G) IF staining of β-catenin in either vehicle or Pizotifen-treated MDA-MB-231

425   cells. (H) Expressions of β-catenin in the cytoplasm and nucleus of MDA-MB-231 cells

426   and the E-cad[+] cells.

428 **Table 1. Primary targets of the identified drugs**
429

The identified drugs	Primary targets of the identified drugs
Hexachlorophene	D-lactate dehydrogenase (D-LDH), not expressed in mammalian cells
Troglitazone	Agonist for Peroxisome proliferator-activated receptor α and γ (PPARα and γ)
Pizotifen malate	5-hydroxytryptamine receptor 2C (HTR2C)
Salmeterol	Adrenergic receptor beta 2 (ADRB2)
Nitazoxanide	Pyruvate ferredoxin oxidoreductase (PFOR), not expressed in mammalian cells
Valproic acid	Histone deacetylases (HDACs)
Dicumarol	NAD(P)H dehydrogenase quinone 1 (NQO1)
Loxapine succinate	Dopamine receptor D2 and D4 (DRD2 and DRD4)
Adrenosterone	Hydroxysteroid (11-beta) dehydrogenase 1 (HSD11β1)
Riluzole hydrochloride	Glutamate R and Voltage-dependent Na+ channel
Naftopidil dihydrochloride	5-hydroxytryptamine receptor 1A (HTR1A) and α1-adrenergic receptor (AR)
S(-)Eticlopride hydrochloride	Dopamine receptor D2 (DRD2)
Racecadotril	Membrane metallo-endopeptidase (MME)
Ipriflavone	Unknow
Flurbiprofen	Cyclooxygenase 1 and 2 (Cox1 and 2)
Zardaverine	Phosphodiesterase III/IV (PDE3/4)
Leflunomide	Dihydroorotate dehydrogenase (DHODH)
Olmesartan	Angiotensin II receptor alpha
Disulfiram	Aldehyde dehydrogenase (ALDH) Dopamine β-hydroxylase (DBH)
Zuclopenthixol dihydrochloride	Dopamine receptors D1 and D2 (DRD1 and 2)

Discussion

Reducing or eliminating mortality associated with metastatic disease is a key goal of

medical oncology, but few models exist that allow for rapid, effective screening of novel

compounds that target the metastatic dissemination of cancer cells. Based on accumulated

evidence that at least fifty genes play an essential role in governing both metastasis and

gastrulation progression (Table 1S), we hypothesized that small molecule inhibitors that

interrupt gastrulation of zebrafish embryos might suppress metastatic progression of

human cancer cells. We created a unique screening concept utilizing gastrulation of

zebrafish embryos to test the hypothesis. Our results clearly confirmed our hypothesis:

25.6% (20/76 drugs) of epiboly-interrupting drugs could also suppress cell motility and

invasion of highly metastatic human cell lines in vitro. In particular, Pizotifen which is an

antagonist for serotonin receptor 2C and one of the epiboly-interrupting drugs, could

suppress metastasis in a mouse model (Figure 3A-E). Thus, this screen could offer a novel

platform for discovery of anti-metastasis drugs.

There are at least two advantages to the screen described herein. One is that the

screen can easily be converted to a chemical genetic screening platform. Indeed, we have

provided the first evidence that HTR2C, which is a primary target of Pizotifen, induced

EMT and promoted metastatic dissemination of cancer cells (Figure 4A-E). In this

research, 1,280 FDA approval drugs were screened, this is less than a few percent of all of

druggable targets (approximately 100 targets) in the human proteome in the body. If

chemical genetic screening using specific inhibitor libraries were conducted, more genes

that contribute to metastasis and gastrulation could be identified. The second advantage is

that the screen enables one researcher to test 100 drugs in 5 hours with using optical

microscopy, drugs, and zebrafish embryos. That indicates this screen is not only highly

efficient, low-cost, and low-labor but also enables researchers who do not have high

throughput screening instruments to conduct drug screening for anti-metastasis drugs.

Acknowledgments

We sincerely appreciate Dr. Joshua Collins (NIH/NIDCR) and Dr. Diane Palmieri

(NIH/NCI) for helping this research. We thank Dr. Herrick (Albany medical collage) for

providing pCMV-h5TH2C-VSV with us. This study was funded by grants from National

465   Author Contributions

466   Design research; J.N. Conducting experiments; J.N. and L.T. Analyzing data: J.N.

467   Writing the paper; J.N. and Z.G. Funding Acquisition; Z.G, S.W. B.C.G., H.M. and

468   Supervision; Z.G.

470   Declaration of Interests

471   J.N., L.T., B.C.G., S.W., H.M. and G.Z. declare no conflict of interest.

473   Materials and Methods

474   Zebrafish embryo screening

Zebrafish embryos at two cell stage were collected at 20 mins after their fertilization.

Each drug was added to a well of a 24-well plate containing approximately 20 zebrafish

embryos per well in either 10 μM or 50 μM final concentration when the embryos reached

the sphere stage. Chemical treatment was initiated at 4 hours post-fertilization (hpf) and

approximately 20 embryos were treated with two different concentrations for each

compound tested. The treatment was ended at 9 hpf when vehicle (DMSO) treated

embryos as control reach 80-90% completion of the epiboly stage. The compounds which

induced delay (<50% epiboly) in epiboly were selected as hit compounds for in vitro

testing using highly metastatic human cancer cell lines. The study protocol was approved

by the Institutional Animal Care and Use Committee of the National University of

Singapore (protocol number: R16-1068).

487   Reagents

FDA, EMA and other agencies approved chemical libraries was purchased from

Prestwick Chemical (Illkirch, France). Pizotifen and S(-) Eticlopride hydrochloride were

purchased from Sigma-Aldrich (St. Louis, MO).

Cell culture and cell viability assay

MCF7, MDA-MB-231, MDA-MB-435, MIA-PaCa2, PC3, SW620, PC9 and HaCaT cells

were obtained from American Type Culture Collection (ATCC, Manassas, VA).

Luciferase-expressing 4T1 (4T1-12B) cells were provided from Dr. Gary Sahagian (Tufts

University, Boston, MA). All culture methods followed the supplier's instruction. Cell

viability assay was performed as previously described (16).

Plasmid

A DNA fragment coding for HTR2C was amplified by PCR with primers containing

restriction enzyme recognition sequences. The HTR2C coding fragment was amplified

from hsp70l:mCherry-T2A-CreERT2 plasmid (17).

Immunoblotting

Western blotting was performed as described previously (16). Anti-PRMT1, anti-

CYP11A1, anti-E-cadherin, anti-EpCAM, anti-Vimentin, anti-N-cadherin, anti-Zeb1,

anti-Histone H$_3$, anti-α-tubulin and anti-GAPDH antibodies were purchased from Cell

signaling Technology. Anti-HTR2C and anti-DRD2 antibodies were purchased form

Abcam. Anti-phospho-GSK3β (Ser9), anti-GSK3β, Anti-KRT18, anti-KRT19, anti-

MMP1, anti-MMP2, anti-S100A4, anti-Luciferase, and anti-β-catenin antibodies were

purchased from Santa Cruz.

Flow cytometry

Cells were stained with FITC-conjugated E-cadherin antibody (Biolegend, San Diego,

CA). Flow cytometry was performed as described (26) and analyzed with FlowJo

software (TreeStar, Ashland, OR).

shRNA mediated gene knockdown

The short hairpin RNA (shRNA)-expressing lentivirus vectors were

constructed using pLVX-shRNA1 vector (Clontech). PRMT1-shRNA_#3–targeting

sequence is GTGTTCCAGTATCTCTGATTA; PRMT1-shRNA_#4–targeting sequence

is TTGACTCCTACGCACACTTTG. CYP11A1-shRNA_#4–targeting sequence is

GCGATTCATTGATGCCATCTA; CYP11A1-shRNA_#4–targeting sequence is

GAAATCCAACACCTCAGCGAT. Human HTR2C-shRNA–targeting sequence is

TCATGCACCTCTGCGCTATAT. Mouse HTR2C-shRNA–targeting sequence is

CTTCATACCGCTGACGATTAT. LacZ-shRNA– targeting sequence is

CTACACAAATCAGCGATT.

Immunofluorescence

Immunofluorescence microscopy assay was performed by previously described (16). Goat

anti-mouse and goat anti-rabbit immunoglobulin G (IgG) antibodies conjugated to Alexa

Fluor 488 (Life Technologies) and diluted at 1:100 were used. Nuclei were visualized by

the addition of 2μg/ml of 4', 6-diamidino-2-phenylindole (DAPI) and photographed at

100x magnification by a fluorescent microscope BZ-X700 (KEYENCE, Japan).

Boyden chamber cell motility and invasion assay

These assays were performed by previously described (16). In Boyden chamber assay,

either $3x10^5$ MDA-MB-231, $1x10^6$ MDA-MB-435, or $5x10^5$ PC9 cells were applied to

each well in the upper chamber.

Zebrafish xenotransplantation model

Tg(kdrl:eGFP) zebrafish was provided by Dr. Stainier (Max Planck Institute for Heart and Lung Research). Embryos that were derived from the line were maintained in E3 medium containing 200μM 1-phenyl-2-thiourea (PTU). Approximately 100-400 Red fluorescence protein (RFP)-labelled MBA-MB-231 or MIA-Paca2 cells were injected into the duct of Cuvier of the zebrafish at 2dpf. The fish were randomly assigned to two groups. One group was maintained in the presence of pizotifen-containing E3 medium and the other group was maintained in vehicle-containing E3 medium.

Spontaneous metastasis mouse model

4T1-12B cells ($2x10^4$) were injected into the #4 mammary fat pad while the mice were anesthetized. To monitor tumor growth and metastases, mice were imaged biweekly by IVIS Imaging System (ParkinElmer). The primary tumor was resected 10 days after inoculation. The study protocol (protocol number: BRC IACUC #110612) was approved by A*STAR (Agency for Science, Technology and Research, Singapore).

Histological Analysis

All OCT embedded primary tumors, lungs, and livers of mice from the spontaneous metastasis 4T1 model were sectioned on a cryostat. Eight micron sections were taken at five hundred micron intervals through the entirety of the livers and lungs. Sections were subsequently stained with hematoxylin and eosin. Metastatic lesions were counted under a microscope in each section for both lungs and livers.

Statistics

Data were analyzed by Student's t test; $p < 0.05$ was considered significant.

References

1.   Chambers AF, Groom AC, & MacDonald IC (2002) Dissemination and growth
of cancer cells in metastatic sites. *Nature reviews. Cancer* 2(8):563-572.
2.   Steeg PS (2003) Metastasis suppressors alter the signal transduction of
cancer cells. *Nature reviews. Cancer* 3(1):55-63.
3.   Fidler IJ (2003) The pathogenesis of cancer metastasis: the 'seed and soil'
hypothesis revisited. *Nature reviews. Cancer* 3(6):453-458.
4.   Cavallaro U & Christofori G (2004) Cell adhesion and signalling by cadherins
and Ig-CAMs in cancer. *Nature reviews. Cancer* 4(2):118-132.
5.   Ellenbroek SI, Iden S, & Collard JG (2012) Cell polarity proteins and cancer.
Semin Cancer Biol 22(3):208-215.
6.   Tsai JH & Yang J (2013) Epithelial-mesenchymal plasticity in carcinoma
metastasis. *Genes & development* 27(20):2192-2206.
7.   Solnica-Krezel L (2005) Conserved patterns of cell movements during
vertebrate gastrulation. *Curr Biol* 15(6):R213-228.
8.   Latimer A & Jessen JR (2010) Extracellular matrix assembly and organization
during zebrafish gastrulation. *Matrix Biol* 29(2):89-96.
9.   Solnica-Krezel L (2006) Gastrulation in zebrafish -- all just about adhesion?
Curr Opin Genet Dev 16(4):433-441.
10. Tsai YJ*, et al.* (2011) The predominant protein arginine methyltransferase
PRMT1 is critical for zebrafish convergence and extension during
gastrulation. *FEBS J* 278(6):905-917.
11. Hsu HJ, Liang MR, Chen CT, & Chung BC (2006) Pregnenolone stabilizes
microtubules and promotes zebrafish embryonic cell movement. *Nature*
439(7075):480-483.
12. Weinbach EC & Garbus J (1969) Mechanism of Action of Reagents that
uncouple Oxidative Phosphorylation. *Nature* 221:1016.
13. Sack U*, et al.* (2011) Novel effect of antihelminthic Niclosamide on S100A4-
mediated metastatic progression in colon cancer. *Journal of the National*
Cancer Institute 103(13):1018-1036.
14. Huang EW*, et al.* (2012) Vinpocetine inhibits breast cancer cells growth in
vitro and in vivo. *Apoptosis : an international journal on programmed cell*
death 17(10):1120-1130.
15. Szilágyi G*, et al.* (2005) Effects of vinpocetine on the redistribution of cerebral
blood flow and glucose metabolism in chronic ischemic stroke patients: a PET
study. *Journal of the Neurological Sciences* 229-230:275-284.
16. Nakayama J, Lu J, Makinoshima H, & Gong Z (2020) A Novel Zebrafish Model
of Metastasis Identifies the HSD11β1 Inhibitor Adrenosterone as a
Suppressor of Epithelial-Mesenchymal Transition and Metastatic
Dissemination. *Mol Cancer Res* 18(3):477-487.
17. Kolosionek E*, et al.* (2009) Expression and activity of phosphodiesterase
isoforms during epithelial mesenchymal transition: the role of
phosphodiesterase 4. *Mol Biol Cell* 20(22):4751-4765.
18. Sasaki T*, et al.* (2014) Serum CD10 is associated with liver metastasis in
colorectal cancer. *J Surg Res.* 192(2):390-394.
19. Liu P*, et al.* (2013) Disulfiram targets cancer stem-like cells and reverses
resistance and cross-resistance in acquired paclitaxel-resistant triple-
negative breast cancer cells. *British journal of cancer* 109(7):1876-1885.
20. Gunnarsson L, Jauhiainen A, Kristiansson E, Nerman O, & Larsson DGJ (2008)
Evolutionary conservation of human drug targets in organisms used for
environmental risk assessments. *Environmental Science & Technology*
42(15):5807-5813.
21. Tao K, Fang M, Alroy J, & Sahagian GG (2008) Imagable 4T1 model for the
study of late stage breast cancer. *BMC cancer* 8:228.

22.   Kröger C, Afeyan, A.; Mraz, J.; Eaton, EN.; Reinhardt, F.; Khodor, YL.; Thiru, P.;
622   Bierie B.; Ye, X.; Burge, CB.; Weinberg RA (2019) Acquisition of a hybrid E/M
623   state is essential for tumorigenicity of basal breast cancer cells. *Proceedings*
624   *of the National Academy of Sciences of the United States of America*
625   116(15):7353-7362.
626   23.   Beyer TD, M.; Thumberger, T.; Vick, P.; Tisler, M.; Schneider, I.; Bogusch, S.;
627   Andre, P.; Ulmer, B.; Walentek, P.; Niesler, B.; Blum, M.; Schweickert, A.
628   (2012) Serotonin signaling is required for Wnt-dependent GRP specification
629   and leftward flow in Xenopus. *Curr Biol* 22(1):33-39.
630   24.   Sui HX, H.; Ji, Q.; Liu, X.; Zhou, L.; Song, H.; Zhou, X.; Xu, Y.; Chen, Z.; Cai, J.; Ji, G.;
631   Li, Q. (2015) 5-hydroxytryptamine receptor (5-HT1DR) promotes colorectal
632   cancer metastasis by regulating Axin1/β-catenin/MMP-7 signaling pathway.
633   *Oncotarget* 6(28):25975-25987.
634   25.   Zhan TR, N.; Boutros, M. (2017) Wnt signaling in cancer. *Oncogene*
635   36(11):1461–1473.
636   26.   Nakayama J*, et al.* (2009) BLNK suppresses pre-B-cell leukemogenesis
637   through inhibition of JAK3. *Blood* 113(7):1483-1492.
638

    Supplemental Information

    Figure S1.

644
645
Figure S1. Molecular mechanisms of epiboly in zebrafish overlap with those in

cancer metastasis.

(A) Western blot analysis of PRMT1 (upper) and CYP11A1 (middle) protein levels in

non-metastatic human cancer cell line (MCF7) and highly metastatic human cancer cell

lines (MDA-MB-231, MDA-MB-435, MIA-PaCa2, PC9, HCCLM3, PC3 and SW620); β-

actin loading control is shown (bottom). (B) Knockdown of PRMT1 or CYP11A1 in

MDA-MB-231 cells. MBA-MB-231 cells were transfected with a control shRNA

targeting LacZ, and one of four independent shRNAs targeting PRMT1 (clone #1 to

#4) or one of two independent shRNAs targeting CYP11A1 (clone #1 to #4). Reduced

PRMT1 and CYP11A1 expression, determined by western blot, in sub-cell lines of

MDA-MB-231 cells expressing PRMT1 shRNA (clone #3 and #4 or CYP11A1 (clone

#2 and #4), compared with controls (parental cell line MDA-MB-231 and control

shRNA cells); β-actin levels shown as a loading control. (C) Effect of shRNAs

targeting either PRMT1 or CYP11A1 on cell motility and invasion of MBA-MB-231

cells. Parental MDA-MB-231 cells and four sub-cell lines of MDA-MB-231 cells that

were transfected with either shRNA targeting either LacZ, two independent shRNAs

targeting PRMT1 (clone #3 and #4) or two independent shRNAs targeting CYP11A1

(clone #2 and #4), were subjected to Boyden chamber assays. (D) Zebrafish embryos

treated with either vehicle (DMSO), 10μM Niclosamide, or 50μM vinpocetine.

Approximately 20 embryos were treated with either DMSO as a vehicle control,

niclosamide, or vinpocetine. The treatment was started at 4 hpf when all of embryos

reached sphere stage and ended at 9 hpf when control embryos reached 80-90%

epiboly stage. Each experiment was performed at least twice. Statistical analysis was

determined by Student's *t* test.

Figure S2.

Figure S2. S (-) Eticlopride hydrochloride suppressed cell motility and invasion of

human cancer cells.

Related to Figure 2.

(A) Western blot analysis of DRD2 levels in non-metastatic human cancer cell line,

MCF7 (breast) and highly metastatic human cancer cell lines, MDA-MB-231 (breast),

MDA-MB-435 (melanoma), MIA-PaCa2 (pancreas), PC3 (prostate) and SW620 (colon);

GAPDH loading control is shown (bottom). (B) Effect of S (-) Eticlopride hydrochloride

on cell motility and invasion of MBA-MB-231, MDA-MB-435, and PC9 cells. Either

vehicle or pizotifen treated cells were subjected to Boyden chamber assays. Fetal bovine

serum (1%v/v) was used as the chemoattractant in both assays. Each experiment was

performed at least twice. Statistical analysis was determined by Student's *t* test.

Figure S3.

Figure S3. Pizotifen suppressed metastatic dissemination of human pancreatic

cancer cells in a zebrafish xenotransplantation model.

Related to Figure 2.

Representative images of dissemination of MIA-PaCa2 cells in zebrafish

xenotransplantation model. The fish were inoculated with MIA-PaCa2 cells, and treated

with either vehicle (top left) or drug (lower left). White arrows heads indicate

disseminated MIA-PaCa2 cells. The images were shown in 4x magnification. Scale bar,

100μm. The mean frequencies of the fish showing head, trunk, or end-tail dissemination

were tabulated (right). Each value is indicated as the mean ± SEM of two independent

experiments. Statistical analysis was determined by Student's *t* test.

702 **Table S1. A list of the genes that are involved between gastrulation and**
703 **metastasis progression**
704

Genes	Gastrulation Defects	Ref	Effects in Metastasis	Ref
BMP	Convergence and extension	(1)	EMT	(2)
WNT	Convergence and extension	(3)	Migration and Invasion	(4)
FGF	Convergence and extension	(5)	Invision	(6)
EGF	Epiboly	(7)	Migration	(8)
PDGF	Convergence and extension	(9)	EMT	(10)
CXCL12	Migration of endodermal cells	(11)	Migration and Invasion	(12)
CXCR4	Migration of endodermal cells	(11)	Migration and Invasion	(12)
PIK3CA	Convergence and extension	(13)	Migration and Invasion	(14)
YES	Epiboly	(15)	Migration	(16)
FYN	Epiboly	(17)	Migration and Invasion	(18)
MAPK1	Epiboly	(19)	Migration	(20)
SHP2	Convergence and extension	(21)	Migration	(22)
SNAI1	Convergence and extension	(23)	EMT	(24)
SNAI2	Mesoderm & Neural crest formation	(25)	EMT	(95)
TWIST1	Mesoderm formation	(26)	EMT	(27)
TBXT	Convergence and extension	(3)	EMT	(28)
ZEB1	Epiboly	(29)	EMT	(30)
GSC	Mesodermal patterning	(31)	EMT	(32)
FOXC2	Unclear, defects in gastrulation	(33)	EMT	(34)
STAT3	Convergence and extension	(35)	Migration	(36)
POU5F1	Epiboly	(37)	EMT	(38)
EZH2	Unclear, defects in gastrulation	(39)	Invasion	(40)
EHMT2	Defects in Neurogenesis	(41)	Migration and Invasion	(42)
BMI1	Defects in skelton formation	(43)	EMT	(44)
RHOA	Convergence and extension	(45)	Migration and Invasion	(46)
CDC42	Convergence and extension	(47)	Migration and Invasion	(48)
RAC1	Convergence and extension	(49)	Migration and Invasion	(50)
ROCK2	Convergence and extension	(51)	Migration and Invasion	(52)
PAR1	Convergence and extension	(53)	Migration	(54)
PRKCI	Convergence and extension	(53)	EMT	(55)
CAP1	Convergence and extension	(56)	Migration	(57)
EZR	Epiboly	(58)	Migration	(59)
EPCAM	Epiboly	(60)	Migration and Invasion	(61)
ITGB1 / ITA5	Mesodermal Migration	(62)	Migration and Invasion	(63)
FN1	Convergence and extension	(64)	Invasion	(65)
HAS2	Dorsal migration of lateral cells	(66)	Invasion	(67)
MMP14	Convergence and extension	(68)	Invasion	(69)
COX1	Epiboly	(70)	Invasion	(71)
PTGES	Convergence and extension	(72)	Invasion	(73)
SLC39A6	Aterior migration	(74)	EMT	(75)
GNA12 / 13	Convergence and extension	(41)	Migration and Invasion	(76)
OGT	Epiboly	(77)	Migration and Invasion	(78)
CCN1	Cell Movement	(79)	Migration and Invasion	(80)
TRPM7	Convergence and extension	(81)	Migration	(82)
MAPKAPK2	Epiboly	(83)	Migration	(84)
B4GALT1	Convergence and extension	(85)	Invasion	(86)
IER2	Convergence and extension	(87)	Migration	(88)
TIP1	Convergence and extension	(89)	Migration and Invasion	(90)
PAK5	Convergence and extension	(91)	Migration	(92)
MARCKS	Convergence and extension	(93)	Migration and Invasion	(94)

Table S1.

A list of the fifty genes that play essential role in governing both metastasis and

gastrulation progression. The gastrulation defects in Xenopus or zebrafish that are

induced by knockdown of each of these genes, were indicated. The molecular

mechanism in metastasis that are inhibited by knockdown of each of the same genes,

were indicated.

Table S2. A list of the drugs that interfere with epiboly progression in zebrafish
713

Chemical name	Chemical formula	Effect of 10uM	Effect of 50uM
Acitretin	C21H26O3	Delayed	Delayed
Adrenosterone	C19H24O3	Delayed	Delayed
Albendazole	C12H15N3O2S	Severe delayed	Severe delayed
Alfadolone acetate	C23H34O5	Delayed	Delayed
Alfaxalone	C21H32O3	Delayed	Delayed
Alprostadil	C20H34O5	Delayed	Delayed
Altrenogest	C21H26O2	Slightly delayed	Delayed
Ampiroxicam	C20H21N3O7S	Non effect	Delayed
Anethole-trithione	C10H8OS3	Delayed	Delayed
Antimycin A	C28H40N2O9	Delayed	Delayed
Avobenzone	C20H22O3	Delayed	Delayed
Benzoxiquine	C16H11NO2	Non effect	Delayed
Bosentan	C27H29N5O6S	Delayed	Delayed
Butoconazole nitrate	C19H18Cl3N3O3S	Delayed	Toxic lethal
Camptothecine (S,+)	C20H16N2O4	Severe delayed	Severe delayed
Carbenoxolone disodium salt	C34H48Na2O7	Delayed	Toxic lethal
Carmofur	C11H16FN3O3	Slightly delayed	Delayed
Carprofen	C15H12ClNO2	Severe delayed	Toxic lethal
Cefdinir	C14H13N5O5S2	Delayed	Delayed
Celecoxib	C17H14F3N3O2S	Delayed	Delayed
Chlorambucil	C14H19Cl2NO2	Slightly delayed	Delayed
Chlorhexidine	C22H30Cl2N10	Non effect	Toxic lethal
Ciclopirox ethanolamine	C14H24N2O3	Delayed	Severe delayed
Cinoxacin	C12H10N2O5	Delayed	Severe delayed
Clofibrate	C12H15ClO3	Non effect	Severe delayed
Clopidogrel	C16H16ClNO2S	Non effect	Delayed
Clorgyline hydrochloride	C13H16Cl3NO	Delayed	Delayed
Colchicine	C22H25NO6	Non effect	Delayed
Deptropine citrate	C29H35NO8	Delayed	Delayed
Desipramine hydrochloride	C18H23ClN2	Delayed	Delayed
Diclofenac sodium	C14H10Cl2NNaO2	Delayed	Severe delayed
Dicumarol	C19H12O6	Delayed	Severe delayed
Diethylstilbestrol	C18H20O2	Delayed	Toxic lethal
Dimaprit dihydrochloride	C6H17Cl2N3S	Slightly delayed	Delayed
Disulfiram	C10H20N2S4	Delayed	Delayed
Dopamine hydrochloride	C8H12ClNO2	Delayed	Delayed
Eburnamonine (-)	C19H22N2O	Delayed	Delayed
Ethaverine hydrochloride	C24H30ClNO4	Delayed	Delayed
Ethinylestradiol	C20H24O2	Delayed	Severe delayed
Ethopropazine hydrochloride	C19H25ClN2S	Delayed	Delayed
Ethoxyquin	C14H19NO	Non effect	Delayed
Exemestane	C20H24O2	Slightly delayed	Delayed
Ezetimibe	C24H21F2NO3	Slightly delayed	Delayed
Fenbendazole	C15H13N3O2S	Non effect	Delayed
Fenoprofen calcium salt dihydrate	C30H30CaO8	Slightly delayed	Delayed
Fentiazac	C17H12ClNO2S	Toxic lethal	Toxic lethal
Floxuridine	C9H11FN2O5	Delayed	Toxic lethal
Flunixin meglumine	C21H28F3N3O7	Delayed	Toxic lethal
Flutamide	C11H11F3N2O3	Delayed	Toxic lethal
Fluticasone propionate	C25H31F3O5S	Non effect	Delayed
Furosemide	C12H11ClN2O5S	Delayed	Delayed
Gatifloxacin	C19H22FN3O4	Delayed	Delayed
Gemcitabine	C9H11F2N3O4	Delayed	Delayed
Gemfibrozil	C15H22O3	Delayed	Toxic lethal
Gestrinone	C21H24O2	Delayed	Delayed
Haloprogin	C9H4Cl3IO	Delayed	Toxic lethal
Hexachlorophene	C13H6Cl6O2	Delayed	Severe delayed
Hexestrol	C18H22O2	Slightly delayed	Delayed
Ibudilast	C14H18N2O	Non effect	Delayed
Idazoxan hydrochloride	C11H13ClN2O2	Slightly delayed	Delayed
Idazoxan hydrochloride	C11H13ClN2O2	Non effect	Delayed

Idebenone	C19H30O5	Severe delayed	Toxic lethal
Indomethacin	C19H16ClNO4	Non effect	Delayed
Ipriflavone	C18H16O3	Delayed	Severe delayed
Isotretinoin	C20H28O2	Non effect	Severe delayed
Isradipine	C19H21N3O5	Non effect	Delayed
Lansoprazole	C16H14F3N3O2S	Slightly delayed	Delayed
Latanoprost	C26H40O5	Non effect	Delayed
Leflunomide	C12H9F3N2O2	Delayed	Severe delayed
Letrozole	C17H11N5	Non effect	Delayed
Lithocholic acid	C24H40O3	Non effect	Delayed
Lodoxamide	C11H6ClN3O6	Non effect	Delayed
Lofepramine	C26H27ClN2O	Non effect	Delayed
Loratadine	C22H23ClN2O2	Delayed	Delayed
Loxapine succinate	C22H24ClN3O5	Delayed	Delayed
Mebendazole	C16H13N3O3	Severe delayed	Severe delayed
Mebendazole	C22H26N2O2	Non effect	Delayed
Meloxicam	C14H13N3O4S2	Delayed	Toxic lethal
Methiazole	C12H15N3O2S	Delayed	Delayed
Mevastatin	C23H34O5	Non effect	Delayed
MK 801 hydrogen maleate	C20H19NO4	Slightly delayed	Delayed
Nabumetone	C15H16O2	Non effect	Severe delayed
Naftopidil dihydrochloride	C24H30Cl2N2O3	Slightly delayed	Delayed
Nandrolone	C18H26O2	Delayed	Delayed
Naproxen sodium salt	C14H13NaO3	Delayed	Delayed
Niclosamide	C13H8Cl2N2O4	Delayed	Delayed
Nifekalant	C19H27N5O5	Delayed	Delayed
Niflumic acid	C13H9F3N2O2	Delayed	Delayed
Nimesulide	C13H12N2O5S	Non effect	Delayed
Nisoldipine	C20H24N2O6	Delayed	Toxic lethal
Nitazoxanide	C12H9N3O5S	Severe delayed	Severe delayed
Norethindrone	C20H26O2	Non effect	Delayed
Norgestimate	C23H31NO3	Slightly delayed	Delayed
Oxfendazol	C15H13N3O3S	Slightly delayed	Delayed
Oxibendazol	C12H15N3O3	Severe delayed	Severe delayed
Oxymetholone	C21H32O3	Slightly delayed	Delayed
Parbendazole	C13H17N3O2	Severe delayed	Severe delayed
Parthenolide	C15H20O3	Non effect	Delayed
Penciclovir	C10H15N5O3	Non effect	Delayed
Pentobarbital	C11H18N2O3	Non effect	Delayed
Phenazopyridine hydrochloride	C11H12ClN5	Delayed	Toxic lethal
Phenothiazine	C12H9NS	Non effect	Delayed
Phenoxybenzamine hydrochloride	C18H23Cl2NO	Non effect	Delayed
Pizotifen malate	C23H27NO5S	Delayed	Severe delayed
Pramoxine hydrochloride	C17H28ClNO3	Slightly delayed	Delayed
Prilocaine hydrochloride	C13H21ClN2O	Non effect	Delayed
Primidone	C12H14N2O2	Slightly delayed	Delayed
Racecadotril	C21H23NO4S	Slightly delayed	Delayed
Riluzole hydrochloride	C8H6ClF3N2OS	Non effect	Delayed
Ritonavir	C37H48N6O5S2	Non effect	Severe delayed
S(-)Eticlopride hydrochloride	C17H26Cl2N2O3	Delayed	Delayed
Salmeterol	C25H37NO4	Non effect	Delayed
Streptomycin sulfate	C42H84N14O36S3	Non effect	Delayed
Sulconazole nitrate	C18H16Cl3N3O3S	Delayed	Delayed
Tegafur	C8H9FN2O3	Delayed	Delayed
Telmisartan	C33H30N4O2	Severe delayed	Toxic lethal
Tenatoprazole	C16H18N4O3S	Non effect	Delayed
Terbinafine	C21H25N	Non effect	Delayed
Thimerosal	C9H9HgNaO2S	Non effect	Delayed
Thiorphan	C12H15NO3S	Delayed	Delayed
Tolcapone	C14H11NO5	Severe delayed	Severe delayed
Topotecan	C23H23N3O5	Delayed	Delayed
Tracazolate hydrochloride	C16H25ClN4O2	Severe delayed	Delayed
Tribenoside	C29H34O6	Delayed	Delayed
Triclabendazole	C14H9Cl3N2OS	Delayed	Delayed
Triclosan	C12H7Cl3O2	Delayed	Severe delayed

Trioxsalen	C14H12O3	Delayed	Delayed
Troglitazone	C24H27NO5S	Severe delayed	Toxic lethal
Valproic acid	C8H16O2	Non effect	Delayed
Voriconazole	C16H14F3N5O	Non effect	Delayed
Zardaverine	C12H10F2N2O3	Slightly delayed	Delayed
Zuclopenthixol dihydrochloride	C22H27Cl3N2OS	Delayed	Delayed

Table S2. A list of the drugs that interfere with epiboly progression in zebrafish.

Related to Figure 1.

A list of positive "hit" drugs that interfered with epiboly progression. Gastrulation

defects or status of each of the zebrafish embryos that were treated with either 10μM

or 50μM concentrations, are indicated.

Table S3. Effects of pharmacological inhibition of HTR2C on metastatic
dissemination of MDA-MB-231 cells in zebrafish xenografted models
723

		Experiment #1	Experiment #2	Experiment #3	Average of Experiments
Drug: Vehicle Cell: MDA-MB-231	Non-dissemination	0% n1=0/17	0% n2=0/12	6.66% n3=1/15	2.22±3.84%
	Head	58.82% n1=10/17	91.66% n2=11/12	6.66% n3=1/15	72.38±17.15%
	Trunk	52.94% n1=9/17	8.33% n2=1/12	20% n3=2/15	27.09±23.13%
	End-tail	100% n1=17/17	100% n2=12/12	86.66% n3=13/15	95.55±7.69%
Drug: Pizotifen Cell: MDA-MB-231	Non-dissemination	55% n1=11/20	31.57% n2=6/19	45.45 % n3=10/22	44.01±11.77%
	Head	5% n1=1/20	31.57% n2=6/19	18.18% n3=4/22	18.25±13.28%
	Trunk	5% n1=1/20	10.52% n2=2/19	4.45% n3=1/22	6.69±3.32%
	End-tail	45% n1=9/20	57.89% n2=11/19	50% n3=11/22	50.96±6.50%

724
725 Related to Figure 2D.

726 The numbers and frequencies of the fish showing the dissemination patterns in vehicle

727 or Pizotifen-treated group, were indicated. The fish showed both patterns of

728 dissemination were redundantly counted in this analysis.

Table S4. Effects of pharmacological inhibition of HTR2C on metastatic dissemination of Mia-PaCa2 cells in zebrafish xenografted models

		Experiment _#1	Experiment _#2	Average of Experiments
Drug: Vehicle Cell: MIA-PaCa2	Non-dissemination	17.64% n1=3/17	16.66% n2=2/12	17.15+0.69%
	Head	82.35% n1=14/17	66.66% n2=8/12	74.50+11.09%
	Trunk	29.41% n1=5/17	8.33% n2=1/12	18.87+14.90%
	End-tail	70.58% n1=12/17	83.33% n2=10/17	76.96%+9.01
Drug: Pizotifen Cell: MIA-PaCa2	Non-dissemination	40% n1=4/10	52.63% n2=10/19	46.31+8.93%
	Head	20% n1=2/10	10.52% n2=2/19	15.26+6.69%
	Trunk	10% n1=1/10	5.26% n2=1/19	7.63+3.34%
	End-tail	40% n1=4/10	42.05% n2=8/19	41.4+1.48%

Related to Figure S3.

The numbers and frequencies of the fish showing the dissemination patterns in vehicle

or Pizotifen-treated group, were indicated. The fish showed both patterns of

dissemination were redundantly counted in this analysis.

Table S5. Effects of genetic inhibition of HTR2C on metastatic dissemination of MDA-MB-231 cells in zebrafish xenografted models

		Experiment _#1	Experiment _#2	Average of Experiments
shLacZ	Non-dissemination	0% n1=0/10	0% n2=0/10	0%
	Head	60% n1=6/10	100% n2=10/10	80 ±28.28%
	Trunk	30% n1=3/10	10% n2=1/10	20±14.14%
	End-tail	80% n1=8/10	100% n2=10/10	90%±14.14
shHTR2C	Non-dissemination	80% n1=12/15	76.84% n2=14/19	76.84±4.46 %
	Head	6.66% n1=1/15	15.78% n2=3/19	11.22±6.45%
	Trunk	6.66% n1=1/15	5.26% n2=1/19	5.96±0.99%
	End-tail	20% n1=3/15	26.31% n2=5/19	23.15±4.46%

Related to Figure 2E.

The numbers and frequencies of the fish showing the dissemination patterns in the

zebrafish that were inoculated with either shLacZ or shHTR2C MDA-MB-231 cells,

748 were indicated. The fish showed both patterns of dissemination were redundantly

749 counted in this analysis.

750 **Table S6. Effects of HTR2C overexpression on metastatic dissemination of**
751 **MCF7 cells in zebrafish xenografted models**
752

		Experiment_#1	Experiment_#2	Average of Experiments
VC	Non-dissemination	46.15% n1=6/13	40% n2=4/10	43.07±4.35%
	Head	46.15% n1=6/13	20% n2=2/10	33.07±18.49%
	Trunk	0% n1=0/13	0% n2=0/10	0%
	End-tail	53.84% n1=7/13	60% n2=6/10	56.92±4.35
HTR2C	Non-dissemination	0% n1=0/14	0% n2=0/15	0%
	Head	100% n1=14/14	93.33% n2=14/15	96.66±4.71%
	Trunk	64.28% n1=9/14	73.33% n2=11/15	68.80±6.39%
	End-tail	85.71% n1=12/14	93.33% n2=14/15	89.52±5.38%

754 Related to Figure 4E.

755 The numbers and frequencies of the fish showing the dissemination patterns in the

756 zebrafish that were inoculated with MCF7 cells expressing either VC or HTR2C, were

757 indicated. The fish showed both patterns of dissemination were redundantly counted

758 in this analysis.

759
760
References for Table S1.

1.   Kondo M (2007) Bone morphogenetic proteins in the early development
of zebrafish. *FEBS J* 274(12):2960-2967.
2.   Katsuno Y, *et al.* (2008) Bone morphogenetic protein signaling enhances
invasion and bone metastasis of breast cancer cells through Smad
pathway. *Oncogene* 27(49):6322-6333.
3.   Tada M & Smith JC (2000) Xwnt11 is a target of Xenopus Brachyury:
regulation of gastrulation movements via Dishevelled, but not through the
canonical Wnt pathway. *Development* 127:2227-2238.
4.   Vincan E & Barker N (2008) The upstream components of the Wnt
signalling pathway in the dynamic EMT and MET associated with
colorectal cancer progression. *Clin Exp Metastasis* 25(6):657-663.
5.   Yang X, Dormann D, Munsterberg AE, & Weijer CJ (2002) Cell movement
patterns during gastrulation in the chick are controlled by positive and
negative chemotaxis mediated by FGF4 and FGF8. *Dev Cell* 3(3):425-437.

6.   Nomura S, *et al.* (2008) FGF10/FGFR2 signal induces cell migration and
invasion in pancreatic cancer. *Br J Cancer* 99(2):305-313.
7.   Song S, *et al.* (2013) Pou5f1-dependent EGF expression controls E-
cadherin endocytosis, cell adhesion, and zebrafish epiboly movements.
Dev Cell 24(5):486-501.
8.   Lu Z, Jiang G, Blume-Jensen P, & Hunter T (2001) Epidermal growth
factor-induced tumor cell invasion and metastasis initiated by
dephosphorylation and downregulation of focal adhesion kinase. *Mol Cell*
Biol 21(12):4016-4031.
9.   Damm EW & Winklbauer R (2011) PDGF-A controls mesoderm cell
orientation and radial intercalation during Xenopus gastrulation.
Development 138(3):565-575.
10.   Jechlinger M, *et al.* (2006) Autocrine PDGFR signaling promotes mammary
cancer metastasis. *J Clin Invest* 116(6):1561-1570.
11.   Mizoguchi T, Verkade H, Heath JK, Kuroiwa A, & Kikuchi Y (2008)
Sdf1/Cxcr4 signaling controls the dorsal migration of endodermal cells
during zebrafish gastrulation. *Development* 135(15):2521-2529.
12.   Shen HB, Gu ZQ, Jian K, & Qi J (2013) CXCR4-mediated Stat3 activation is
essential for CXCL12-induced cell invasion in bladder cancer. *Tumour Biol*
34(3):1839-1845.
13.   Montero J-A, Kilian B, Chan J, Bayliss PE, & Heisenberg C-P (2003)
Phosphoinositide 3-Kinase Is Required for Process Outgrowth and Cell
Polarization of Gastrulating Mesendodermal Cells. *Current Biology*
13(15):1279-1289.
14.   Wander SA, *et al.* (2013) PI3K/mTOR inhibition can impair tumor
invasion and metastasis in vivo despite a lack of antiproliferative action in
vitro: implications for targeted therapy. *Breast Cancer Res Treat*
138(2):369-381.
15.   Tsai WB, Zhang X, Sharma D, Wu W, & Kinsey WH (2005) Role of Yes
kinase during early zebrafish development. *Dev Biol* 277(1):129-141.
16.   Barraclough J, Hodgkinson C, Hogg A, Dive C, & Welman A (2007)
Increases in c-Yes Expression Level and Activity Promote Motility But Not
Proliferation of Human Colorectal Carcinoma Cells. *Neoplasia* 9(9):745-
IN732.
17.   Sharma D, Holets L, Zhang X, & Kinsey WH (2005) Role of Fyn kinase in
signaling associated with epiboly during zebrafish development. *Dev Biol*
285(2):462-476.
18.   Yadav V & Denning MF (2011) Fyn is induced by Ras/PI3K/Akt signaling
and is required for enhanced invasion/migration. *Mol Carcinog* 50(5):346-
352.
19.   Krens SF, *et al.* (2008) Distinct functions for ERK1 and ERK2 in cell
migration processes during zebrafish gastrulation. *Dev Biol* 319(2):370-
383.
20.   Radtke S, Milanovic M, & Rossé C (2013) ERK2 but not ERK1 mediates
HGF-induced motility in non-small cell lung carcinoma cell lines. *J Cell Sci.*
126(Pt11):2381-2391.
21.   Jopling C, van Geemen D, & den Hertog J (2007) Shp2 knockdown and
Noonan/LEOPARD mutant Shp2-induced gastrulation defects. *PLoS Genet*
3(12):e225.
22.   Wang FM, *et al.* (2005) SHP-2 promoting migration and metastasis of
MCF-7 with loss of E-cadherin, dephosphorylation of FAK and secretion of

MMP-9 induced by IL-1beta in vivo and in vitro. *Breast Cancer Res Treat*
89(1):5-14.

23.   Ip YT & Gridley T (2002) Cell movements during gastrulation: snail
dependent and independent pathways. *Curr Opin Genet Dev* 12(4):423-
429.

24.   Batlle E, Sancho E, Francí C, Domínguez D, & Monfar M (2000) The
transcription factor snail is a repressor of E-cadherin gene expression in
epithelial tumour cells. *Nat Cell Biol* 2(2):84-89.

25.   Shi J, Severson C, Yang J, Wedlich D, & Klymkowsky MW (2011) Snail2
controls mesodermal BMP/Wnt induction of neural crest. *Development*
138(15):3135-3145.

26.   Castanon I & Baylies MK (2002) A Twist in fate: evolutionary comparison
of Twist structure and function. *Gene* 287(1-2):11-22.

27.   Yang J*, et al.* (2004) Twist, a master regulator of morphogenesis, plays an
essential role in tumor metastasis. *Cell* 117(7):927-939.

28.   Fernando RI*, et al.* (2010) The T-box transcription factor Brachyury
promotes epithelial-mesenchymal transition in human tumor cells. *J Clin*
Invest 120(2):533-544.

29.   Vannier C, Mock K, Brabletz T, & Driever W (2013) Zeb1 regulates E-
cadherin and Epcam (epithelial cell adhesion molecule) expression to
control cell behavior in early zebrafish development. *J Biol Chem*
288(26):18643-18659.

30.   Spaderna S*, et al.* (2008) The transcriptional repressor ZEB1 promotes
metastasis and loss of cell polarity in cancer. *Cancer Res* 68(2):537-544.

31.   Sander V, Reversade B, & De Robertis EM (2007) The opposing homeobox
genes Goosecoid and Vent1/2 self-regulate Xenopus patterning. *EMBO J*
26(12):2955-2965.

32.   Hartwell KA*, et al.* (2006) The Spemann organizer gene, Goosecoid,
promotes tumor metastasis. *Proc Natl Acad Sci U S A* 103(50):18969-
18974.

33.   Wilm B, James RG, Schultheiss TM, & Hogan BL (2004) The forkhead
genes, Foxc1 and Foxc2, regulate paraxial versus intermediate mesoderm
cell fate. *Dev Biol* 271(1):176-189.

34.   Mani SA*, et al.* (2007) Mesenchyme Forkhead 1 (FOXC2) plays a key role
in metastasis and is associated with aggressive basal-like breast cancers.
Proc Natl Acad Sci U S A 104(24):10069-10074.

35.   Miyagi C*, et al.* (2004) STAT3 noncell-autonomously controls planar cell
polarity during zebrafish convergence and extension. *J Cell Biol*
166(7):975-981.

36.   Abdulghani J*, et al.* (2008) Stat3 promotes metastatic progression of
prostate cancer. *Am J Pathol* 172(6):1717-1728.

37.   Lachnit M, Kur E, & Driever W (2008) Alterations of the cytoskeleton in all
three embryonic lineages contribute to the epiboly defect of Pou5f1/Oct4
deficient MZspg zebrafish embryos. *Dev Biol* 315(1):1-17.

38.   Dai X*, et al.* (2013) OCT4 regulates epithelial-mesenchymal transition and
its knockdown inhibits colorectal cancer cell migration and invasion.
Oncol Rep 29(1):155-160.

39.   O'Carroll D*, et al.* (2001) The polycomb-group gene Ezh2 is required for
early mouse development. *Mol Cell Biol* 21(13):4330-4336.

40.   Ren G, *et al.* (2012) Polycomb protein EZH2 regulates tumor invasion via
the transcriptional repression of the metastasis suppressor RKIP in breast
and prostate cancer. *Cancer Res* 72(12):3091-3104.

41.   Lin F, *et al.* (2005) Essential roles of G{alpha}12/13 signaling in distinct
cell behaviors driving zebrafish convergence and extension gastrulation
movements. *J Cell Biol* 169(5):777-787.

42.   Chen MW, *et al.* (2010) H3K9 histone methyltransferase G9a promotes
lung cancer invasion and metastasis by silencing the cell adhesion
molecule Ep-CAM. *Cancer Res* 70(20):7830-7840.

43.   van der Lugt NM, *et al.* (1994) Posterior transformation, neurological
abnormalities, and severe hematopoietic defects in mice with a targeted
deletion of the bmi-1 proto-oncogene. *Genes Dev* 8(7):757-769.

44.   Guo BH, *et al.* (2011) Bmi-1 promotes invasion and metastasis, and its
elevated expression is correlated with an advanced stage of breast cancer.
Mol Cancer 10(1):10.

45.   Zhu S, Liu L, Korzh V, Gong Z, & Low BC (2006) RhoA acts downstream of
Wnt5 and Wnt11 to regulate convergence and extension movements by
involving effectors Rho kinase and Diaphanous: use of zebrafish as an in
vivo model for GTPase signaling. *Cell Signal* 18(3):359-372.

46.   Yoshioka K, Nakamori S, & Itoh K (1999) Overexpression of small GTP-
binding protein RhoA promotes invasion of tumor cells. *Cancer Res*
59(8):2004-2010.

47.   Choi SC & Han JK (2002) Xenopus Cdc42 regulates convergent extension
movements during gastrulation through Wnt/Ca2+ signaling pathway.
Dev Biol 244(2):342-357.

48.   Reymond N, *et al.* (2012) Cdc42 promotes transendothelial migration of
cancer cells through beta1 integrin. *J Cell Biol* 199(4):653-668.

49.   Habas R, Dawid IB, & He X (2003) Coactivation of Rac and Rho by
Wnt/Frizzled signaling is required for vertebrate gastrulation. . *Genes Dev*
17(2):295-309.

50.   Vega FM & Ridley AJ (2008) Rho GTPases in cancer cell biology. *FEBS Lett*
582(14):2093-2101.

51.   Marlow F, Topczewski J, Sepich D, & Solnica-Krezel L (2002) Zebrafish
Rho kinase 2 acts downstream of Wnt11 to mediate cell polarity and
effective convergence and extension movements. *Curr Biol* 12(11):876-
884.

52.   Itoh K, *et al.* (1999) An essential part for Rho-associated kinase in the
transcellular invasion of tumor cells. *Nat Med* 5(2):221-225.

53.   Kusakabe M & Nishida E (2004) The polarity-inducing kinase Par-1
controls Xenopus gastrulation in cooperation with 14-3-3 and aPKC.
EMBO J 23(21):4190-4201.

54.   Shi X, Gangadharan B, Brass LF, Ruf W, & Mueller BM (2004) Protease-
activated receptors (PAR1 and PAR2) contribute to tumor cell motility
and metastasis. *Mol Cancer Res* 2(7):395-402.

55.   Gunaratne A, Thai BL, & Di Guglielmo GM (2013) Atypical protein kinase C
phosphorylates Par6 and facilitates transforming growth factor beta-
induced epithelial-to-mesenchymal transition. *Mol Cell Biol* 33(5):874-
886.

56.   Seifert K, Ibrahim H, Stodtmeister T, Winklbauer R, & Niessen CM (2009)
An adhesion-independent, aPKC-dependent function for cadherins in
morphogenetic movements. *J Cell Sci* 122(Pt 14):2514-2523.

57. Yamazaki K, *et al.* (2009) Adenylate cyclase-associated protein 1
overexpressed in pancreatic cancers is involved in cancer cell motility.
Lab Invest 89(4):425-432.
58. Link V, *et al.* (2006) Identification of regulators of germ layer
morphogenesis using proteomics in zebrafish. *J Cell Sci* 119(Pt 10):2073-
2083.
59. Khanna C, *et al.* (2004) The membrane-cytoskeleton linker ezrin is
necessary for osteosarcoma metastasis. *Nat Med* 10(2):182-186.
60. Slanchev K, *et al.* (2009) The epithelial cell adhesion molecule EpCAM is
required for epithelial morphogenesis and integrity during zebrafish
epiboly and skin development. *PLoS Genet* 5(7):e1000563.
61. Ni J, *et al.* (2012) Role of the EpCAM (CD326) in prostate cancer
metastasis and progression. *Cancer Metastasis Rev* 31(3-4):779-791.
62. Skalski M, Alfandari D, & Darribere T (1998) A key function for alphav
containing integrins in mesodermal cell migration during Pleurodeles
waltl gastrulation. *Dev Biol* 195(2):158-173.
63. Felding-Habermann B (2003) Integrin adhesion receptors in tumor
metastasis. *Clin Exp Metastasis* 20(3):203-213.
64. Marsden M & DeSimone DW (2003) Integrin-ECM Interactions Regulate
Cadherin-Dependent Cell Adhesion and Are Required for Convergent
Extension in Xenopus. *Current Biology* 13(14):1182-1191.
65. Malik G, *et al.* (2010) Plasma fibronectin promotes lung metastasis by
contributions to fibrin clots and tumor cell invasion. *Cancer Res*
70(11):4327-4334.
66. Bakkers J, *et al.* (2004) Has2 is required upstream of Rac1 to govern
dorsal migration of lateral cells during zebrafish gastrulation.
Development 131(3):525-537.
67. Kim HR, *et al.* (2004) Hyaluronan facilitates invasion of colon carcinoma
cells in vitro via interaction with CD44. *Cancer Res* 64(13):4569-4576.
68. Coyle RC, Latimer A, & Jessen JR (2008) Membrane-type 1 matrix
metalloproteinase regulates cell migration during zebrafish gastrulation:
evidence for an interaction with non-canonical Wnt signaling. *Exp Cell Res*
314(10):2150-2162.
69. Perentes JY, *et al.* (2011) Cancer cell-associated MT1-MMP promotes
blood vessel invasion and distant metastasis in triple-negative mammary
tumors. *Cancer Res* 71(13):4527-4538.
70. Cha YI, *et al.* (2006) Cyclooxygenase-1-derived PGE2 promotes cell
motility via the G-protein-coupled EP4 receptor during vertebrate
gastrulation. *Genes Dev* 20(1):77-86.
71. Kundu N & Fulton AM (2002) Selective cyclooxygenase (COX)-1 or COX-2
inhibitors control metastatic disease in a murine model of breast cancer.
Cancer Res 62(8):2343-2346.
72. Speirs CK, *et al.* (2010) Prostaglandin Gbetagamma signaling stimulates
gastrulation movements by limiting cell adhesion through Snai1a
stabilization. *Development* 137(8):1327-1337.
73. Wang D & Dubois RN (2006) Prostaglandins and cancer. *Gut* 55(1):115-
122.
74. Yamashita S, *et al.* (2004) Zinc transporter LIVI controls epithelial-
mesenchymal transition in zebrafish gastrula organizer. *Nature*
429(6989):298-302.

75. Lue HW, *et al.* (2011) LIV-1 promotes prostate cancer epithelial-to-
mesenchymal transition and metastasis through HB-EGF shedding and
EGFR-mediated ERK signaling. *PLoS One* 6(11):e27720.
76. Yagi H, *et al.* (2011) A synthetic biology approach reveals a CXCR4-G13-
Rho signaling axis driving transendothelial migration of metastatic breast
cancer cells. *Sci Signal* 4(191):ra60.
77. Webster DM, *et al.* (2009) O-GlcNAc modifications regulate cell survival
and epiboly during zebrafish development. *BMC Dev Biol* 9:28.
78. Lynch TP, *et al.* (2012) Critical role of O-Linked beta-N-acetylglucosamine
transferase in prostate cancer invasion, angiogenesis, and metastasis. *J
Biol Chem* 287(14):11070-11081.
79. Latinkic BV (2003) Xenopus Cyr61 regulates gastrulation movements and
modulates Wnt signalling. *Development* 130(11):2429-2441.
80. Lin J, *et al.* (2012) A novel anti-Cyr61 antibody inhibits breast cancer
growth and metastasis in vivo. *Cancer Immunol Immunother* 61(5):677-
687.
81. Liu W, *et al.* (2011) TRPM7 regulates gastrulation during vertebrate
embryogenesis. *Dev Biol* 350(2):348-357.
82. Middelbeek J, *et al.* (2012) TRPM7 is required for breast tumor cell
metastasis. *Cancer Res* 72(16):4250-4261.
83. Holloway BA, *et al.* (2009) A novel role for MAPKAPK2 in morphogenesis
during zebrafish development. *PLoS Genet* 5(3):e1000413.
84. Kumar B, *et al.* (2010) p38 mitogen-activated protein kinase-driven
MAPKAPK2 regulates invasion of bladder cancer by modulation of MMP-2
and MMP-9 activity. *Cancer Res* 70(2):832-841.
85. Machingo QJ, Fritz A, & Shur BD (2006) A beta1,4-galactosyltransferase is
required for convergent extension movements in zebrafish. *Dev Biol*
297(2):471-482.
86. Zhu X, *et al.* (2005) Elevated beta1,4-galactosyltransferase I in highly
metastatic human lung cancer cells. Identification of E1AF as important
transcription activator. *J Biol Chem* 280(13):12503-12516.
87. Hong SK, Tanegashima K, & Dawid IB (2011) XIer2 is required for
convergent extension movements during Xenopus development. *Int J Dev
Biol* 55(10-12):917-921.
88. Neeb A, *et al.* (2012) The immediate early gene Ier2 promotes tumor cell
motility and metastasis, and predicts poor survival of colorectal cancer
patients. *Oncogene* 31(33):3796-3806.
89. Besser J, Leito JT, van der Meer DL, & Bagowski CP (2007) Tip-1 induces
filopodia growth and is important for gastrulation movements during
zebrafish development. *Dev Growth Differ* 49(3):205-214.
90. Han M, Wang H, Zhang HT, & Han Z (2012) The PDZ protein TIP-1
facilitates cell migration and pulmonary metastasis of human invasive
breast cancer cells in athymic mice. *Biochem Biophys Res Commun*
422(1):139-145.
91. Faure S, *et al.* (2005) Xenopus p21-activated kinase 5 regulates
blastomeres' adhesive properties during convergent extension
movements. *Dev Biol* 277(2):472-492.
92. Gong W, *et al.* (2009) P21-activated kinase 5 is overexpressed during
colorectal cancer progression and regulates colorectal carcinoma cell
adhesion and migration. *Int J Cancer* 125(3):548-555.

93.   Iioka H, Ueno N, & Kinoshita N (2004) Essential role of MARCKS in cortical
1028   actin dynamics during gastrulation movements. *J Cell Biol* 164(2):169-
1029   174.
1030   94.   Rombouts K*, et al.* (2013) Myristoylated Alanine-Rich protein Kinase C
1031   Substrate (MARCKS) expression modulates the metastatic phenotype in
1032   human and murine colon carcinoma in vitro and in vivo. *Cancer Lett*
1033   333(2):244-252.
1034
1035

Table S2. A list of the drugs that interfere with epiboly progression in zebrafish

Chemical name	Chemical formula	Effect of 10uM	Effect of 50uM
Acitretin	C21H26O3	Delayed	Delayed
Adrenosterone	C19H24O3	Delayed	Delayed
Albendazole	C12H15N3O2S	Severe delayed	Severe delayed
Alfadolone acetate	C23H34O5	Delayed	Delayed
Alfaxalone	C21H32O3	Delayed	Delayed
Alprostadil	C20H34O5	Delayed	Delayed
Altrenogest	C21H26O2	Slightly delayed	Delayed
Ampiroxicam	C20H21N3O7S	Non effect	Delayed
Anethole-trithione	C10H8OS3	Delayed	Delayed
Antimycin A	C28H40N2O9	Delayed	Delayed
Avobenzone	C20H22O3	Delayed	Delayed
Benzoxiquine	C16H11NO2	Non effect	Delayed
Bosentan	C27H29N5O6S	Delayed	Delayed
Butoconazole nitrate	C19H18Cl3N3O3S	Delayed	Toxic lethal
Camptothecine (S,+)	C20H16N2O4	Severe delayed	Severe delayed
Carbenoxolone disodium salt	C34H48Na2O7	Delayed	Toxic lethal
Carmofur	C11H16FN3O3	Slightly delayed	Delayed
Carprofen	C15H12ClNO2	Severe delayed	Toxic lethal
Cefdinir	C14H13N5O5S2	Delayed	Delayed
Celecoxib	C17H14F3N3O2S	Delayed	Delayed
Chlorambucil	C14H19Cl2NO2	Slightly delayed	Delayed
Chlorhexidine	C22H30Cl2N10	Non effect	Toxic lethal
Ciclopirox ethanolamine	C14H24N2O3	Delayed	Severe delayed
Cinoxacin	C12H10N2O5	Delayed	Severe delayed
Clofibrate	C12H15ClO3	Non effect	Severe delayed
Clopidogrel	C16H16ClNO2S	Non effect	Delayed
Clorgyline hydrochloride	C13H16Cl3NO	Delayed	Delayed
Colchicine	C22H25NO6	Non effect	Delayed
Deptropine citrate	C29H35NO8	Delayed	Delayed
Desipramine hydrochloride	C18H23ClN2	Delayed	Delayed
Diclofenac sodium	C14H10Cl2NNaO2	Delayed	Severe delayed
Dicumarol	C19H12O6	Delayed	Severe delayed
Diethylstilbestrol	C18H20O2	Delayed	Toxic lethal
Dimaprit dihydrochloride	C6H17Cl2N3S	Slightly delayed	Delayed
Disulfiram	C10H20N2S4	Delayed	Delayed
Dopamine hydrochloride	C8H12ClNO2	Delayed	Delayed
Eburnamonine (-)	C19H22N2O	Delayed	Delayed
Ethaverine hydrochloride	C24H30ClNO4	Delayed	Delayed
Ethinylestradiol	C20H24O2	Delayed	Severe delayed
Ethopropazine hydrochloride	C19H25ClN2S	Delayed	Delayed
Ethoxyquin	C14H19NO	Non effect	Delayed
Exemestane	C20H24O2	Slightly delayed	Delayed
Ezetimibe	C24H21F2NO3	Slightly delayed	Delayed
Fenbendazole	C15H13N3O2S	Non effect	Delayed
Fenoprofen calcium salt dihydrate	C30H30CaO8	Slightly delayed	Delayed
Fentiazac	C17H12ClNO2S	Toxic lethal	Toxic lethal
Floxuridine	C9H11FN2O5	Delayed	Toxic lethal
Flunixin meglumine	C21H28F3N3O7	Delayed	Toxic lethal
Flutamide	C11H11F3N2O3	Delayed	Toxic lethal
Fluticasone propionate	C25H31F3O5S	Non effect	Delayed
Furosemide	C12H11ClN2O5S	Delayed	Delayed
Gatifloxacin	C19H22FN3O4	Delayed	Delayed
Gemcitabine	C9H11F2N3O4	Delayed	Delayed
Gemfibrozil	C15H22O3	Delayed	Toxic lethal
Gestrinone	C21H24O2	Delayed	Delayed
Haloprogin	C9H4Cl3IO	Delayed	Toxic lethal
Hexachlorophene	C13H6Cl6O2	Delayed	Severe delayed
Hexestrol	C18H22O2	Slightly delayed	Delayed
Ibudilast	C14H18N2O	Non effect	Delayed
Idazoxan hydrochloride	C11H13ClN2O2	Slightly delayed	Delayed
Idazoxan hydrochloride	C11H13ClN2O2	Non effect	Delayed
Idebenone	C19H30O5	Severe delayed	Toxic lethal

Indomethacin	C19H16ClNO4	Non effect	Delayed
Ipriflavone	C18H16O3	Delayed	Severe delayed
Isotretinoin	C20H28O2	Non effect	Severe delayed
Isradipine	C19H21N3O5	Non effect	Delayed
Lansoprazole	C16H14F3N3O2S	Slightly delayed	Delayed
Latanoprost	C26H40O5	Non effect	Delayed
Leflunomide	C12H9F3N2O2	Delayed	Severe delayed
Letrozole	C17H11N5	Non effect	Delayed
Lithocholic acid	C24H40O3	Non effect	Delayed
Lodoxamide	C11H6ClN3O6	Non effect	Delayed
Lofepramine	C26H27ClN2O	Non effect	Delayed
Loratadine	C22H23ClN2O2	Delayed	Delayed
Loxapine succinate	C22H24ClN3O5	Delayed	Delayed
Mebendazole	C16H13N3O3	Severe delayed	Severe delayed
Mebendazole	C22H26N2O2	Non effect	Delayed
Meloxicam	C14H13N3O4S2	Delayed	Toxic lethal
Methiazole	C12H15N3O2S	Delayed	Delayed
Mevastatin	C23H34O5	Non effect	Delayed
MK 801 hydrogen maleate	C20H19NO4	Slightly delayed	Delayed
Nabumetone	C15H16O2	Non effect	Severe delayed
Naftopidil dihydrochloride	C24H30Cl2N2O3	Slightly delayed	Delayed
Nandrolone	C18H26O2	Delayed	Delayed
Naproxen sodium salt	C14H13NaO3	Delayed	Delayed
Niclosamide	C13H8Cl2N2O4	Delayed	Delayed
Nifekalant	C19H27N5O5	Delayed	Delayed
Niflumic acid	C13H9F3N2O2	Delayed	Delayed
Nimesulide	C13H12N2O5S	Non effect	Delayed
Nisoldipine	C20H24N2O6	Delayed	Toxic lethal
Nitazoxanide	C12H9N3O5S	Severe delayed	Severe delayed
Norethindrone	C20H26O2	Non effect	Delayed
Norgestimate	C23H31NO3	Slightly delayed	Delayed
Oxfendazol	C15H13N3O3S	Slightly delayed	Delayed
Oxibendazol	C12H15N3O3	Severe delayed	Severe delayed
Oxymetholone	C21H32O3	Slightly delayed	Delayed
Parbendazole	C13H17N3O2	Severe delayed	Severe delayed
Parthenolide	C15H20O3	Non effect	Delayed
Penciclovir	C10H15N5O3	Non effect	Delayed
Pentobarbital	C11H18N2O3	Non effect	Delayed
Phenazopyridine hydrochloride	C11H12ClN5	Delayed	Toxic lethal
Phenothiazine	C12H9NS	Non effect	Delayed
Phenoxybenzamine hydrochloride	C18H23Cl2NO	Non effect	Delayed
Pizotifen malate	C23H27NO5S	Delayed	Severe delayed
Pramoxine hydrochloride	C17H28ClNO3	Slightly delayed	Delayed
Prilocaine hydrochloride	C13H21ClN2O	Non effect	Delayed
Primidone	C12H14N2O2	Slightly delayed	Delayed
Racecadotril	C21H23NO4S	Slightly delayed	Delayed
Riluzole hydrochloride	C8H6ClF3N2OS	Non effect	Delayed
Ritonavir	C37H48N6O5S2	Non effect	Severe delayed
S(-)Eticlopride hydrochloride	C17H26Cl2N2O3	Delayed	Delayed
Salmeterol	C25H37NO4	Non effect	Delayed
Streptomycin sulfate	C42H84N14O36S3	Non effect	Delayed
Sulconazole nitrate	C18H16Cl3N3O3S	Delayed	Delayed
Tegafur	C8H9FN2O3	Delayed	Delayed
Telmisartan	C33H30N4O2	Severe delayed	Toxic lethal
Tenatoprazole	C16H18N4O3S	Non effect	Delayed
Terbinafine	C21H25N	Non effect	Delayed
Thimerosal	C9H9HgNaO2S	Non effect	Delayed
Thiorphan	C12H15NO3S	Delayed	Delayed
Tolcapone	C14H11NO5	Severe delayed	Severe delayed
Topotecan	C23H23N3O5	Delayed	Delayed
Tracazolate hydrochloride	C16H25ClN4O2	Severe delayed	Delayed
Tribenoside	C29H34O6	Delayed	Delayed
Triclabendazole	C14H9Cl3N2OS	Delayed	Delayed
Triclosan	C12H7Cl3O2	Delayed	Severe delayed
Trioxsalen	C14H12O3	Delayed	Delayed

Troglitazone	C24H27NO5S	Severe delayed	Toxic lethal
Valproic acid	C8H16O2	Non effect	Delayed
Voriconazole	C16H14F3N5O	Non effect	Delayed
Zardaverine	C12H10F2N2O3	Slightly delayed	Delayed
Zuclopenthixol dihydrochloride	C22H27Cl3N2OS	Delayed	Delayed

1038

1039 **Table S2. A list of the drugs that interfere with epiboly progression in zebrafish.**

1040 A list of positive "hit" drugs that interfered with epiboly progression. Gastrulation defects

1041 or status of each of the zebrafish embryos that were treated with either 10μM or 50μM

1042 concentrations, are indicated.

1043

1044

References for Table S1.

1. Kondo M (2007) Bone morphogenetic proteins in the early development of
zebrafish. *FEBS J* 274(12):2960-2967.
2. Katsuno Y, *et al.* (2008) Bone morphogenetic protein signaling enhances
invasion and bone metastasis of breast cancer cells through Smad pathway.
Oncogene 27(49):6322-6333.
3. Tada M & Smith JC (2000) Xwnt11 is a target of Xenopus Brachyury:
regulation of gastrulation movements via Dishevelled, but not through the
canonical Wnt pathway. *Development* 127:2227-2238.
4. Vincan E & Barker N (2008) The upstream components of the Wnt signalling
pathway in the dynamic EMT and MET associated with colorectal cancer
progression. *Clin Exp Metastasis* 25(6):657-663.
5. Yang X, Dormann D, Munsterberg AE, & Weijer CJ (2002) Cell movement
patterns during gastrulation in the chick are controlled by positive and
negative chemotaxis mediated by FGF4 and FGF8. *Dev Cell* 3(3):425-437.
6. Nomura S, *et al.* (2008) FGF10/FGFR2 signal induces cell migration and
invasion in pancreatic cancer. *Br J Cancer* 99(2):305-313.
7. Song S, *et al.* (2013) Pou5f1-dependent EGF expression controls E-cadherin
endocytosis, cell adhesion, and zebrafish epiboly movements. *Dev Cell*
24(5):486-501.
8. Lu Z, Jiang G, Blume-Jensen P, & Hunter T (2001) Epidermal growth factor-
induced tumor cell invasion and metastasis initiated by dephosphorylation
and downregulation of focal adhesion kinase. *Mol Cell Biol* 21(12):4016-4031.
9. Damm EW & Winklbauer R (2011) PDGF-A controls mesoderm cell
orientation and radial intercalation during Xenopus gastrulation.
Development 138(3):565-575.
10. Jechlinger M, *et al.* (2006) Autocrine PDGFR signaling promotes mammary
cancer metastasis. *J Clin Invest* 116(6):1561-1570.
11. Mizoguchi T, Verkade H, Heath JK, Kuroiwa A, & Kikuchi Y (2008) Sdf1/Cxcr4
signaling controls the dorsal migration of endodermal cells during zebrafish
gastrulation. *Development* 135(15):2521-2529.
12. Shen HB, Gu ZQ, Jian K, & Qi J (2013) CXCR4-mediated Stat3 activation is
essential for CXCL12-induced cell invasion in bladder cancer. *Tumour Biol*
34(3):1839-1845.

13.   Montero J-A, Kilian B, Chan J, Bayliss PE, & Heisenberg C-P (2003)
Phosphoinositide 3-Kinase Is Required for Process Outgrowth and Cell
Polarization of Gastrulating Mesendodermal Cells. *Current Biology*
13(15):1279-1289.
14.   Wander SA*, et al.* (2013) PI3K/mTOR inhibition can impair tumor invasion
and metastasis in vivo despite a lack of antiproliferative action in vitro:
implications for targeted therapy. *Breast Cancer Res Treat* 138(2):369-381.
15.   Tsai WB, Zhang X, Sharma D, Wu W, & Kinsey WH (2005) Role of Yes kinase
during early zebrafish development. *Dev Biol* 277(1):129-141.
16.   Barraclough J, Hodgkinson C, Hogg A, Dive C, & Welman A (2007) Increases in
c-Yes Expression Level and Activity Promote Motility But Not Proliferation of
Human Colorectal Carcinoma Cells. *Neoplasia* 9(9):745-IN732.
17.   Sharma D, Holets L, Zhang X, & Kinsey WH (2005) Role of Fyn kinase in
signaling associated with epiboly during zebrafish development. *Dev Biol*
285(2):462-476.
18.   Yadav V & Denning MF (2011) Fyn is induced by Ras/PI3K/Akt signaling and
is required for enhanced invasion/migration. *Mol Carcinog* 50(5):346-352.
19.   Krens SF*, et al.* (2008) Distinct functions for ERK1 and ERK2 in cell migration
processes during zebrafish gastrulation. *Dev Biol* 319(2):370-383.
20.   Radtke S, Milanovic M, & Rossé C (2013) ERK2 but not ERK1 mediates HGF-
induced motility in non-small cell lung carcinoma cell lines. *J Cell Sci.*
126(Pt11):2381-2391.
21.   Jopling C, van Geemen D, & den Hertog J (2007) Shp2 knockdown and
Noonan/LEOPARD mutant Shp2-induced gastrulation defects. *PLoS Genet*
3(12):e225.
22.   Wang FM*, et al.* (2005) SHP-2 promoting migration and metastasis of MCF-7
with loss of E-cadherin, dephosphorylation of FAK and secretion of MMP-9
induced by IL-1beta in vivo and in vitro. *Breast Cancer Res Treat* 89(1):5-14.
23.   Ip YT & Gridley T (2002) Cell movements during gastrulation: snail
dependent and independent pathways. *Curr Opin Genet Dev* 12(4):423-429.
24.   Batlle E, Sancho E, Francí C, Domínguez D, & Monfar M (2000) The
transcription factor snail is a repressor of E-cadherin gene expression in
epithelial tumour cells. *Nat Cell Biol* 2(2):84-89.
25.   Shi J, Severson C, Yang J, Wedlich D, & Klymkowsky MW (2011) Snail2
controls mesodermal BMP/Wnt induction of neural crest. *Development*
138(15):3135-3145.
26.   Castanon I & Baylies MK (2002) A Twist in fate: evolutionary comparison of
Twist structure and function. *Gene* 287(1-2):11-22.
27.   Yang J*, et al.* (2004) Twist, a master regulator of morphogenesis, plays an
essential role in tumor metastasis. *Cell* 117(7):927-939.
28.   Fernando RI*, et al.* (2010) The T-box transcription factor Brachyury
promotes epithelial-mesenchymal transition in human tumor cells. *J Clin*
Invest 120(2):533-544.
29.   Vannier C, Mock K, Brabletz T, & Driever W (2013) Zeb1 regulates E-cadherin
and Epcam (epithelial cell adhesion molecule) expression to control cell
behavior in early zebrafish development. *J Biol Chem* 288(26):18643-18659.
30.   Spaderna S*, et al.* (2008) The transcriptional repressor ZEB1 promotes
metastasis and loss of cell polarity in cancer. *Cancer Res* 68(2):537-544.
31.   Sander V, Reversade B, & De Robertis EM (2007) The opposing homeobox
genes Goosecoid and Vent1/2 self-regulate Xenopus patterning. *EMBO J*
26(12):2955-2965.
32.   Hartwell KA*, et al.* (2006) The Spemann organizer gene, Goosecoid, promotes
tumor metastasis. *Proc Natl Acad Sci U S A* 103(50):18969-18974.

33. Wilm B, James RG, Schultheiss TM, & Hogan BL (2004) The forkhead genes,
Foxc1 and Foxc2, regulate paraxial versus intermediate mesoderm cell fate.
Dev Biol 271(1):176-189.
34. Mani SA*, et al.* (2007) Mesenchyme Forkhead 1 (FOXC2) plays a key role in
metastasis and is associated with aggressive basal-like breast cancers. *Proc*
Natl Acad Sci U S A 104(24):10069-10074.
35. Miyagi C*, et al.* (2004) STAT3 noncell-autonomously controls planar cell
polarity during zebrafish convergence and extension. *J Cell Biol* 166(7):975-
981.
36. Abdulghani J*, et al.* (2008) Stat3 promotes metastatic progression of prostate
cancer. *Am J Pathol* 172(6):1717-1728.
37. Lachnit M, Kur E, & Driever W (2008) Alterations of the cytoskeleton in all
three embryonic lineages contribute to the epiboly defect of Pou5f1/Oct4
deficient MZspg zebrafish embryos. *Dev Biol* 315(1):1-17.
38. Dai X*, et al.* (2013) OCT4 regulates epithelial-mesenchymal transition and its
knockdown inhibits colorectal cancer cell migration and invasion. *Oncol Rep*
29(1):155-160.
39. O'Carroll D*, et al.* (2001) The polycomb-group gene Ezh2 is required for early
mouse development. *Mol Cell Biol* 21(13):4330-4336.
40. Ren G*, et al.* (2012) Polycomb protein EZH2 regulates tumor invasion via the
transcriptional repression of the metastasis suppressor RKIP in breast and
prostate cancer. *Cancer Res* 72(12):3091-3104.
41. Lin F*, et al.* (2005) Essential roles of G{alpha}12/13 signaling in distinct cell
behaviors driving zebrafish convergence and extension gastrulation
movements. *J Cell Biol* 169(5):777-787.
42. Chen MW*, et al.* (2010) H3K9 histone methyltransferase G9a promotes lung
cancer invasion and metastasis by silencing the cell adhesion molecule Ep-
CAM. *Cancer Res* 70(20):7830-7840.
43. van der Lugt NM*, et al.* (1994) Posterior transformation, neurological
abnormalities, and severe hematopoietic defects in mice with a targeted
deletion of the bmi-1 proto-oncogene. *Genes Dev* 8(7):757-769.
44. Guo BH*, et al.* (2011) Bmi-1 promotes invasion and metastasis, and its
elevated expression is correlated with an advanced stage of breast cancer.
Mol Cancer 10(1):10.
45. Zhu S, Liu L, Korzh V, Gong Z, & Low BC (2006) RhoA acts downstream of
Wnt5 and Wnt11 to regulate convergence and extension movements by
involving effectors Rho kinase and Diaphanous: use of zebrafish as an in vivo
model for GTPase signaling. *Cell Signal* 18(3):359-372.
46. Yoshioka K, Nakamori S, & Itoh K (1999) Overexpression of small GTP-
binding protein RhoA promotes invasion of tumor cells. *Cancer Res*
59(8):2004-2010.
47. Choi SC & Han JK (2002) Xenopus Cdc42 regulates convergent extension
movements during gastrulation through Wnt/Ca2+ signaling pathway. *Dev*
Biol 244(2):342-357.
48. Reymond N*, et al.* (2012) Cdc42 promotes transendothelial migration of
cancer cells through beta1 integrin. *J Cell Biol* 199(4):653-668.
49. Habas R, Dawid IB, & He X (2003) Coactivation of Rac and Rho by
Wnt/Frizzled signaling is required for vertebrate gastrulation. . *Genes Dev*
17(2):295-309.
50. Vega FM & Ridley AJ (2008) Rho GTPases in cancer cell biology. *FEBS Lett*
582(14):2093-2101.

51. Marlow F, Topczewski J, Sepich D, & Solnica-Krezel L (2002) Zebrafish Rho
kinase 2 acts downstream of Wnt11 to mediate cell polarity and effective
convergence and extension movements. *Curr Biol* 12(11):876-884.
52. Itoh K, *et al.* (1999) An essential part for Rho-associated kinase in the
transcellular invasion of tumor cells. *Nat Med* 5(2):221-225.
53. Kusakabe M & Nishida E (2004) The polarity-inducing kinase Par-1 controls
Xenopus gastrulation in cooperation with 14-3-3 and aPKC. *EMBO J*
23(21):4190-4201.
54. Shi X, Gangadharan B, Brass LF, Ruf W, & Mueller BM (2004) Protease-
activated receptors (PAR1 and PAR2) contribute to tumor cell motility and
metastasis. *Mol Cancer Res* 2(7):395-402.
55. Gunaratne A, Thai BL, & Di Guglielmo GM (2013) Atypical protein kinase C
phosphorylates Par6 and facilitates transforming growth factor beta-induced
epithelial-to-mesenchymal transition. *Mol Cell Biol* 33(5):874-886.
56. Seifert K, Ibrahim H, Stodtmeister T, Winklbauer R, & Niessen CM (2009) An
adhesion-independent, aPKC-dependent function for cadherins in
morphogenetic movements. *J Cell Sci* 122(Pt 14):2514-2523.
57. Yamazaki K, *et al.* (2009) Adenylate cyclase-associated protein 1
overexpressed in pancreatic cancers is involved in cancer cell motility. *Lab
Invest* 89(4):425-432.
58. Link V, *et al.* (2006) Identification of regulators of germ layer morphogenesis
using proteomics in zebrafish. *J Cell Sci* 119(Pt 10):2073-2083.
59. Khanna C, *et al.* (2004) The membrane-cytoskeleton linker ezrin is necessary
for osteosarcoma metastasis. *Nat Med* 10(2):182-186.
60. Slanchev K, *et al.* (2009) The epithelial cell adhesion molecule EpCAM is
required for epithelial morphogenesis and integrity during zebrafish epiboly
and skin development. *PLoS Genet* 5(7):e1000563.
61. Ni J, *et al.* (2012) Role of the EpCAM (CD326) in prostate cancer metastasis
and progression. *Cancer Metastasis Rev* 31(3-4):779-791.
62. Skalski M, Alfandari D, & Darribere T (1998) A key function for alphav
containing integrins in mesodermal cell migration during Pleurodeles waltl
gastrulation. *Dev Biol* 195(2):158-173.
63. Felding-Habermann B (2003) Integrin adhesion receptors in tumor
metastasis. *Clin Exp Metastasis* 20(3):203-213.
64. Marsden M & DeSimone DW (2003) Integrin-ECM Interactions Regulate
Cadherin-Dependent Cell Adhesion and Are Required for Convergent
Extension in Xenopus. *Current Biology* 13(14):1182-1191.
65. Malik G, *et al.* (2010) Plasma fibronectin promotes lung metastasis by
contributions to fibrin clots and tumor cell invasion. *Cancer Res* 70(11):4327-
4334.
66. Bakkers J, *et al.* (2004) Has2 is required upstream of Rac1 to govern dorsal
migration of lateral cells during zebrafish gastrulation. *Development*
131(3):525-537.
67. Kim HR, *et al.* (2004) Hyaluronan facilitates invasion of colon carcinoma cells
in vitro via interaction with CD44. *Cancer Res* 64(13):4569-4576.
68. Coyle RC, Latimer A, & Jessen JR (2008) Membrane-type 1 matrix
metalloproteinase regulates cell migration during zebrafish gastrulation:
evidence for an interaction with non-canonical Wnt signaling. *Exp Cell Res*
314(10):2150-2162.
69. Perentes JY, *et al.* (2011) Cancer cell-associated MT1-MMP promotes blood
vessel invasion and distant metastasis in triple-negative mammary tumors.
Cancer Res 71(13):4527-4538.

70. Cha YI, *et al.* (2006) Cyclooxygenase-1-derived PGE2 promotes cell motility
via the G-protein-coupled EP4 receptor during vertebrate gastrulation. *Genes*
Dev 20(1):77-86.
71. Kundu N & Fulton AM (2002) Selective cyclooxygenase (COX)-1 or COX-2
inhibitors control metastatic disease in a murine model of breast cancer.
Cancer Res 62(8):2343-2346.
72. Speirs CK, *et al.* (2010) Prostaglandin Gbetagamma signaling stimulates
gastrulation movements by limiting cell adhesion through Snai1a
stabilization. *Development* 137(8):1327-1337.
73. Wang D & Dubois RN (2006) Prostaglandins and cancer. *Gut* 55(1):115-122.
74. Yamashita S, *et al.* (2004) Zinc transporter LIVI controls epithelial-
mesenchymal transition in zebrafish gastrula organizer. *Nature*
429(6989):298-302.
75. Lue HW, *et al.* (2011) LIV-1 promotes prostate cancer epithelial-to-
mesenchymal transition and metastasis through HB-EGF shedding and EGFR-
mediated ERK signaling. *PLoS One* 6(11):e27720.
76. Yagi H, *et al.* (2011) A synthetic biology approach reveals a CXCR4-G13-Rho
signaling axis driving transendothelial migration of metastatic breast cancer
cells. *Sci Signal* 4(191):ra60.
77. Webster DM, *et al.* (2009) O-GlcNAc modifications regulate cell survival and
epiboly during zebrafish development. *BMC Dev Biol* 9:28.
78. Lynch TP, *et al.* (2012) Critical role of O-Linked beta-N-acetylglucosamine
transferase in prostate cancer invasion, angiogenesis, and metastasis. *J Biol*
Chem 287(14):11070-11081.
79. Latinkic BV (2003) Xenopus Cyr61 regulates gastrulation movements and
modulates Wnt signalling. *Development* 130(11):2429-2441.
80. Lin J, *et al.* (2012) A novel anti-Cyr61 antibody inhibits breast cancer growth
and metastasis in vivo. *Cancer Immunol Immunother* 61(5):677-687.
81. Liu W, *et al.* (2011) TRPM7 regulates gastrulation during vertebrate
embryogenesis. *Dev Biol* 350(2):348-357.
82. Middelbeek J, *et al.* (2012) TRPM7 is required for breast tumor cell
metastasis. *Cancer Res* 72(16):4250-4261.
83. Holloway BA, *et al.* (2009) A novel role for MAPKAPK2 in morphogenesis
during zebrafish development. *PLoS Genet* 5(3):e1000413.
84. Kumar B, *et al.* (2010) p38 mitogen-activated protein kinase-driven
MAPKAPK2 regulates invasion of bladder cancer by modulation of MMP-2
and MMP-9 activity. *Cancer Res* 70(2):832-841.
85. Machingo QJ, Fritz A, & Shur BD (2006) A beta1,4-galactosyltransferase is
required for convergent extension movements in zebrafish. *Dev Biol*
297(2):471-482.
86. Zhu X, *et al.* (2005) Elevated beta1,4-galactosyltransferase I in highly
metastatic human lung cancer cells. Identification of E1AF as important
transcription activator. *J Biol Chem* 280(13):12503-12516.
87. Hong SK, Tanegashima K, & Dawid IB (2011) XIer2 is required for convergent
extension movements during Xenopus development. *Int J Dev Biol* 55(10-
12):917-921.
88. Neeb A, *et al.* (2012) The immediate early gene Ier2 promotes tumor cell
motility and metastasis, and predicts poor survival of colorectal cancer
patients. *Oncogene* 31(33):3796-3806.
89. Besser J, Leito JT, van der Meer DL, & Bagowski CP (2007) Tip-1 induces
filopodia growth and is important for gastrulation movements during
zebrafish development. *Dev Growth Differ* 49(3):205-214.

90.   Han M, Wang H, Zhang HT, & Han Z (2012) The PDZ protein TIP-1 facilitates
1288   cell migration and pulmonary metastasis of human invasive breast cancer
1289   cells in athymic mice. *Biochem Biophys Res Commun* 422(1):139-145.
1290   91.   Faure S*, et al.* (2005) Xenopus p21-activated kinase 5 regulates blastomeres'
1291   adhesive properties during convergent extension movements. *Dev Biol*
1292   277(2):472-492.
1293   92.   Gong W*, et al.* (2009) P21-activated kinase 5 is overexpressed during
1294   colorectal cancer progression and regulates colorectal carcinoma cell
1295   adhesion and migration. *Int J Cancer* 125(3):548-555.
1296   93.   Iioka H, Ueno N, & Kinoshita N (2004) Essential role of MARCKS in cortical
1297   actin dynamics during gastrulation movements. *J Cell Biol* 164(2):169-174.
1298   94.   Rombouts K*, et al.* (2013) Myristoylated Alanine-Rich protein Kinase C
1299   Substrate (MARCKS) expression modulates the metastatic phenotype in
1300   human and murine colon carcinoma in vitro and in vivo. *Cancer Lett*
1301   333(2):244-252.
1302   95.   Medici D, Hay ED, & B.R. O (2008) Snail and Slug Promote Epithelial-
1303   Mesenchymal Transition through -Catenin–T-Cell Factor-4-dependent Expression
1304   of Transforming Growth Factor-3. Molecular Biology of the Cell 19(11):4875–
1305   4887.

1306

1307

Legend for source data

Figure 2a_HRT2C_source data

Western blot analysis of HRT2C protein levels in non-metastatic human cancer cell

line, MCF7 (breast) and highly metastatic human cancer cell lines, MDA-MB-231

(breast), MDA-MB-435 (melanoma), MIA-PaCa2 (pancreas), SW620 (colon) and

PC3 (prostate).

Figure 2a_GAPDH_source data

Western blot analysis of GAPDH protein levels in non-metastatic human cancer cell

line, MCF7 (breast) and highly metastatic human cancer cell lines, MDA-MB-231

(breast), MDA-MB-435 (melanoma), MIA-PaCa2 (pancreas), SW620 (colon) and

PC3 (prostate).

Figure 3C_E-cadherin in MCF7_ source data

Western blot analysis of HRT2C protein levels in MCF7 cells expressing either the

control vector or HTR2C.

1324

1325   **Figure 3C_EpCAM in MCF7_ source data**

1326   Western blot analysis of EpCAM protein levels in MCF7 cells expressing either the

1327   control vector or HTR2C.

1328

1329   **Figure 3C_Vimentin in MCF7_ source data**

1330   Western blot analysis of Vimentin protein levels in MCF7 cells expressing either the

1331   control vector or HTR2C.

1332

1333   **Figure 3C_N-cadherin in MCF7_ source data**

1334   Western blot analysis of N-cadherin protein levels in MCF7 cells expressing either the

1335   control vector or HTR2C.

1336

1337   **Figure 3C_Zeb1 in MCF7_ source data**

1338   Western blot analysis of Zeb1 protein levels in MCF7 cells expressing either the

1339   control vector or HTR2C.

1340

1341   **Figure 3C_HRT2C in MCF7_ source data**

1342   Western blot analysis of HRT2C protein levels in MCF7 cells expressing either the

1343   control vector or HTR2C.

1344

1345   **Figure 3C_GAPDH in MCF7_source data**

1346   Western blot analysis of GAPDH protein levels in MCF7 cells expressing either the

1347   control vector or HTR2C.

1348

1349   **Figure 3C_E-cadherin in HaCaT_source data**

1350   Western blot analysis of E-cadherin protein levels in HaCaT cells expressing either

1351   the control vector or HTR2C.

1352

1353   **Figure 3C_EpCAM in HaCaT_source data**

1354   Western blot analysis of EpCAM protein levels in HaCaT cells expressing either the

1355   control vector or HTR2C.

1356

1357   **Figure 3C_Vimentin in HaCaT_source data**

1358   Western blot analysis of Vimentin protein levels in HaCaT cells expressing either the

1359   control vector or HTR2C.

1360

1361   **Figure 3C_Zeb1 in HaCaT_source data**

1362   Western blot analysis of zeb1 protein levels in HaCaT cells expressing either the

1363   control vector or HTR2C.

1364

1365   **Figure 3C_HRT2C in HaCaT_source data**

1366   Western blot analysis of HRT2C protein levels in HaCaT cells expressing either the

1367   control vector or HTR2C.

1368

1369   **Figure 3C_GAPDH in HaCaT_source data**

1370   Western blot analysis of GAPDH protein levels in HaCaT cells expressing either the

1371   control vector or HTR2C.

1372

1373   **Figure 5C_E-cadherin_source data**

1374   Western blot analysis of E-cadherin protein levels in whole cell lysate of 4T1 primary

1375   tumors from either vehicle or Pizotifen-treated mice.

1376

1377 **Figure 5C_Zeb1 _source data**

1378 Western blot analysis of Zeb1 protein levels in whole cell lysate of 4T1 primary

1379 tumors from either vehicle or Pizotifen-treated mice.

1380

1381 **Figure 5C_ Phosphorylation of serine-9 in GSK3β_source data**

1382 Western blot analysis of the protein levels of phosphorylation of serine-9 in GSK3β in

1383 whole cell lysate of 4T1 primary tumors from either vehicle or Pizotifen-treated mice.

1384

1385 **Figure 5C_ GSK3β_source data**

1386 Western blot analysis of GSK3β protein levels in whole cell lysate of 4T1 primary

1387 tumors from either vehicle or Pizotifen-treated mice.

1388

1389 **Figure 5C_Luciferase_source data**

1390 Western blot analysis of Luciferase protein levels in whole cell lysate of 4T1 primary

1391 tumors from either vehicle or Pizotifen-treated mice.

1392

1393 **Figure 5C_β-catenin in the nucleus_source data**

1394 Western blot analysis of β-catenin protein levels in the nucleus of 4T1 primary tumors

1395 from either vehicle or Pizotifen-treated mice.

1396

1397 **Figure 5C_Histone H3 in the nucleus_source data**

1398 Western blot analysis of Histone H3 protein levels in the nucleus of 4T1 primary

1399 tumors from either vehicle or Pizotifen-treated mice.

1400

1401 **Figure 5C_β-catenin in the cytoplasm_source data**

1402   Western blot analysis of β-catenin protein levels in the cytoplasm of 4T1 primary

1403   tumors from either vehicle or Pizotifen-treated mice.

1404

1405   **Figure 5C_β-tubulin in the cytoplasm_source data**

1406   Western blot analysis of β-tubulin protein levels in the cytoplasm of 4T1 primary

1407   tumors from either vehicle or Pizotifen-treated mice.

1408

1409   **Figure 5D_E-cadherin_source data**

1410   Western blot analysis of E-cadherin protein levels in either vehicle or Pizotifen-treated

1411   MDA-MB-231 cells or E-cadherin positive cells in Pizotifen-treated MDA-MB-231

1412   cells.

1413

1414   **Figure 5D_EpCAM_source data**

1415   Western blot analysis of EpCAM protein levels in either vehicle or Pizotifen-treated

1416   MDA-MB-231 cells or E-cadherin positive cells in Pizotifen-treated MDA-MB-231

1417   cells.

1418

1419   **Figure 5D_Keratin18 (KRT18)_source data**

1420   Western blot analysis of KRT18 protein levels in either vehicle or Pizotifen-treated

1421   MDA-MB-231 cells or E-cadherin positive cells in Pizotifen-treated MDA-MB-231

1422   cells.

1423

1424   **Figure 5D_Keratin19 (KRT19)_source data**

1425   Western blot analysis of KRT19 protein levels in either vehicle or Pizotifen-treated

1426   MDA-MB-231 cells or E-cadherin positive cells in Pizotifen-treated MDA-MB-231

1427   cells.

1428

1429 **Figure 5D_Vimentin_source data**

1430 Western blot analysis of Vimentin protein levels in either vehicle or Pizotifen-treated

1431 MDA-MB-231 cells or E-cadherin positive cells in Pizotifen-treated MDA-MB-231

1432 cells.

1433

1434 **Figure 5D_MMP1_source data**

1435 Western blot analysis of MMP1 protein levels in either vehicle or Pizotifen-treated

1436 MDA-MB-231 cells or E-cadherin positive cells in Pizotifen-treated MDA-MB-231

1437 cells.

1438

1439 **Figure 5D_MMP3_source data**

1440 Western blot analysis of MMP3 protein levels in either vehicle or Pizotifen-treated

1441 MDA-MB-231 cells or E-cadherin positive cells in Pizotifen-treated MDA-MB-231

1442 cells.

1443

1444 **Figure 5D_S100A4_source data**

1445 Western blot analysis of S100A4 protein levels in either vehicle or Pizotifen-treated

1446 MDA-MB-231 cells or E-cadherin positive cells in Pizotifen-treated MDA-MB-231

1447 cells.

1448

1449 **Figure 5D_Zeb1_source data**

1450 Western blot analysis of Zeb1 protein levels in either vehicle or Pizotifen-treated

1451 MDA-MB-231 cells or E-cadherin positive cells in Pizotifen-treated MDA-MB-231

1452 cells.

1453

1454 **Figure 5D_GAPDH_source data**

1455 Western blot analysis of GAPDH protein levels in either vehicle or Pizotifen-treated

1456 MDA-MB-231 cells or E-cadherin positive cells in Pizotifen-treated MDA-MB-231

1457 cells.

1458

1459 **Figure 5F_β-catenin in the nucleus_source data**

1460 Western blot analysis of β-catenin protein levels in the nuclear of MCF7 cells

1461 expressing either the control vector or HTR2C.

1462

1463 **Figure 5F_Histone H3 in the nucleus _source data**

1464 Western blot analysis of Histone H3 protein levels in the nuclear of MCF7 cells

1465 expressing either the control vector or HTR2C.

1466

1467 **Figure 5F_β-catenin in the cytoplasm_source data**

1468 Western blot analysis of β-catenin protein levels in the cytoplasm of MCF7 cells

1469 expressing either the control vector or HTR2C.

1470

1471 **Figure 5F_β-tubulin in the cytoplasm_source data**

1472 Western blot analysis of β-tubulin protein levels in the cytoplasm of MCF7 cells

1473 expressing either the control vector or HTR2C.

1474

1475 **Figure 5F_Phosphorylation of serine-9 in GSK3β_source data**

1476 Western blot analysis of the protein levels of phosphorylation of serine-9 in GSK3β in

1477 whole cell lysate of MCF7 cells expressing either the control vector or HTR2C.

1478

1479 **Figure 5F_GSK3β_source data**

1480   Western blot analysis of GSK3β protein levels in whole cell lysate of MCF7 cells

1481   expressing either the control vector or HTR2C.

1482

1483   **Figure 5F_GAPDH_source data**

1484   Western blot analysis of GAPDH protein levels in whole cell lysate of MCF7 cells

1485   expressing either the control vector or HTR2C.

1486

1487   **Figure 5H_β-catenin in the nucleus_source data**

1488   Western blot analysis of β-catenin protein levels in the nucleus of either vehicle or

1489   Pizotifen-treated MDA-MB-231 cells or E-cadherin positive cells in Pizotifen-treated

1490   MDA-MB-231 cells.

1491

1492   **Figure 5H_Histone H3 in the nucleus _source data**

1493   Western blot analysis of Histone H3 protein levels in the nucleus of either vehicle or

1494   Pizotifen-treated MDA-MB-231 cells or E-cadherin positive cells in Pizotifen-treated

1495   MDA-MB-231 cells.

1496

1497   **Figure 5H_β-catenin in the cytoplasm_source data**

1498   Western blot analysis of β-catenin protein levels in the cytoplasm of either vehicle or

1499   Pizotifen-treated MDA-MB-231 cells or E-cadherin positive cells in Pizotifen-treated

1500   MDA-MB-231 cells.

1501

1502   **Figure 5H_β-tubulin in the cytoplasm_source data**

1503   Western blot analysis of β-tubulin protein levels in the cytoplasm of either vehicle or

1504   Pizotifen-treated MDA-MB-231 cells or E-cadherin positive cells in Pizotifen-treated

1505   MDA-MB-231 cells.

1506

1507 **Figure 5H_Phosphorylation of serine-9 in GSK3β_source data**

1508 Western blot analysis of the protein levels of phosphorylation of serine-9 in GSK3β in

1509 whole cell lysate of either vehicle or Pizotifen-treated MDA-MB-231 cells or E-

1510 cadherin positive cells in Pizotifen-treated MDA-MB-231 cells.

1511

1512 **Figure 5H_GSK3β_source data**

1513 Western blot analysis of GSK3β protein levels in whole cell lysate of either vehicle or

1514 Pizotifen-treated MDA-MB-231 cells or E-cadherin positive cells in Pizotifen-treated

1515 MDA-MB-231 cells.

1516

1517 **Figure 5H_GAPDH_source data**

1518 Western blot analysis of GSK3β protein levels in whole cell lysate of either vehicle or

1519 Pizotifen-treated MDA-MB-231 cells or E-cadherin positive cells in Pizotifen-treated

1520 MDA-MB-231 cells.

1521

1522 **Figure S1A_ PRMT1_source data**

1523 Western blot analysis of PRMT1 protein levels in non-metastatic human cancer cell

1524 line (MCF7) and highly metastatic human cancer cell lines (MDA-MB-231, MDA-

1525 MB-435, MIA-PaCa2, PC9, HCCLM3, SW620 and PC3).

1526

1527 **Figure S1A_ CYP11A1_source data**

1528 Western blot analysis of CYP11A1 protein levels in non-metastatic human cancer cell

1529 line (MCF7) and highly metastatic human cancer cell lines (MDA-MB-231, MDA-

1530 MB-435, MIA-PaCa2, PC9, HCCLM3, SW620 and PC3).

1531

1532 **Figure S1A_ β-actin_source data**

1533 Western blot analysis of β-actin protein levels in non-metastatic human cancer cell

1534 line (MCF7) and highly metastatic human cancer cell lines (MDA-MB-231, MDA-

1535 MB-435, MIA-PaCa2, PC9, HCCLM3, SW620 and PC3).

1536

1537 **Figure S1A_ PRMT1_source data**

1538 Western blot analysis of PRMT1 protein levels in sub-clones of MBA-MB-231 cells

1539 which were transfected with either a control shRNA targeting LacZ or one of four

1540 independent shRNAs targeting PRMT1 (clone #1 to #4).

1541

1542 **Figure S1A_ β-actin _source data**

1543 Western blot analysis of β-actin protein levels in sub-clones of MBA-MB-231 cells

1544 which were transfected with either a control shRNA targeting LacZ or one of four

1545 independent shRNAs targeting PRMT1 (clone #1 to #4).

1546

1547 **Figure S1A_CYP11A1_source data**

1548 Western blot analysis of CYP11A1 protein levels in sub-clones of MBA-MB-231

1549 cells which were transfected with either a control shRNA targeting LacZ or one of

1550 four independent shRNAs targeting CYP11A1 (clone #1 to #4).

1551

1552 **Figure S1A_ β-actin_source data**

1553 Western blot analysis of β-actin protein levels in sub-clones of MBA-MB-231 cells

1554 which were transfected with either a control shRNA targeting LacZ or one of four

1555 independent shRNAs targeting CYP11A1 (clone #1 to #4).

1556

1557 **Figure S2A_DRD2_source data**

1558   Western blot analysis of DRD2 protein levels in non-metastatic human cancer cell

1559   line, MCF7 (breast) and highly metastatic human cancer cell lines, MDA-MB-231

1560   (breast), MDA-MB-435 (melanoma), MIA-PaCa2 (pancreas), PC3 (prostate) and

1561   SW620 (colon)

1562

1563   **Figure S2A_GAPDH_source data**

1564   Western blot analysis of GAPDH protein levels in non-metastatic human cancer cell

1565   line, MCF7 (breast) and highly metastatic human cancer cell lines, MDA-MB-231

1566   (breast), MDA-MB-435 (melanoma), MIA-PaCa2 (pancreas), PC3 (prostate) and

1567   SW620 (colon)