Zebrafish embryo screen identifies anti-metastasis drugs

Joji Nakayama[1,2,3,4§], Lora Tan[1], Boon Cher Goh[2], Shu Wang[1, 5], Hideki Makinoshima[3,6] and

Zhiyuan Gong[1§]

[1]Department of Biological Science, National University of Singapore, Singapore

[2]Cancer Science Institute of Singapore, National University of Singapore, Singapore

[3]Tsuruoka Metabolomics Laboratory, National Cancer Center, Tsuruoka, Japan

[4]Shonai Regional Industry Promotion Center, Tsuruoka, Japan.

[5]Institute of Bioengineering and Nanotechnology, Singapore

[6]Division of Translational Research, Exploratory Oncology Research and Clinical Trial

Center, National Cancer Center, Kashiwa, Japan.

Keywords: Zebrafish, metastasis, gastrulation, phenotyping screening

[§]Corresponding authors:

Joji Nakayama

Tsuruoka Metabolomics Laboratory, National Cancer Center, Mizukami 246-2, Kakuganji,

Tsuruoka, Yamagata, Japan 975-0052.

email: Zmetastasis@gmail.com

Zhiyuan Gong

Department of Biological Science, National University of Singapore, Singapore 117543.

email: dbsgzy@nus.edu.sg

The authors disclose no potential conflicts of interest.

Total words excluding abstract, material and methods, references, and figure legends: 1390

Number of figures: 2

Number of supplementary figures: 3

Number of supplementary tables: 3

Abstract

Metastasis is responsible for approximately 90% of cancer-associated mortality, but few

models exist that allow for rapid and effective screening of anti-metastasis drugs. Here we

report a novel screening concept utilizing conserved mechanisms between metastasis and

zebrafish gastrulation for identifying anti-metastasis drugs. Molecular mechanisms involved

in metastasis and vertebrate gastrulation are conserved; at least fifty common genes play

essential roles in governing these mechanisms. We hypothesized that small chemicals that

interrupt zebrafish gastrulation might also suppress metastatic progression of cancer cells and

developed a phenotype-based chemical screen to test the hypothesis. The screen used

epiboly, the first morphogenetic movement in gastrulation as a marker, and identified a

serotonin receptor 2C (HTR2C) antagonist as an epiboly-interrupting drug. A mouse model

of metastasis confirmed that pharmacologic and genetic inhibition of HTR2C suppressed

metastasis. Taken together, this screen could offer a rapid and high-throughput platform for

discovery of anti-metastasis drugs.

Introduction

Metastasis, a leading contributor to the morbidity of cancer patients, occurs through multiple steps: invasion, intravasation, extravasation, colonization, and metastatic tumor formation. The physical translocation of cancer cells is an initial step of metastasis and molecular mechanisms of it involve cell motility, the breakdown of local basement membrane, loss of cell polarity, acquisition of stem cell-like properties, and epithelial-mesenchymal transition (EMT) (1, 2). These cell-biological phenomena are also observed during vertebrate gastrulation in that evolutionarily conserved morphogenetic movements of epiboly, internalization, convergence, and extension progress (3). In zebrafish, the first morphogenetic movement, epiboly, is initiated at approximately 4 hours post fertilization (hpf) to move cells from the animal pole to eventually engulf the entire yolk cell by 10 hpf (4, 5). The embryonic cell movements are governed by the molecular mechanisms that are partially shared in metastatic cell dissemination.

At least fifty common genes play essential roles in both metastasis and gastrulation (Supplementary Table 1): Knockdown of these genes in zebrafish induced gastrulation defects; conversely, overexpression of these genes conferred metastatic potential on cancer cells while knockdown of these genes suppressed metastasis. Based on this evidence, we hypothesized that small chemicals that interrupt zebrafish gastrulation might also suppress metastatic progression of cancer cells.

Here we report a novel screening concept utilizing conserved mechanisms between metastasis and zebrafish gastrulation for identifying anti-metastasis drugs. The screen which used epiboly, the first morphogenetic movement in gastrulation as a marker, identified a serotonin receptor 2C (HTR2C) antagonist as an epiboly-interrupting drug, and a mouse model of metastasis confirmed pharmacologic and genetic inhibition of HTR2C suppressed metastasis.

Results

We first conducted experiments of whether hindering the molecular function of protein

arginine methyltransferase 1 (PRMT1) and cytochrome P450 family 11 (CYP11A1), whose

knockdown induced gastrulation defects in zebrafish but whose involvement in metastasis is

unclear might suppress cell motility and invasion of cancer cells (6)(7). Knockdown of

PRMT1 or CYP11A1 suppressed the motility and invasion of metastatic human cancer cells

(Supplementary Figure 1A-C). Furthermore, we conducted an inverse examination of whether

Niclosamide and Vinpocetine which suppress metastatic dissemination of cancer cells, could

interrupt epiboly of zebrafish embryos (8)(9). Zebrafish embryos treated with either

Niclosamide or Vinpocetine showed complete arrest or severe delay in epiboly, respectively

(Supplementary Figure 1D). Thus, epiboly could serve as a marker for a screening assay and

epiboly-interrupting drugs could potentially suppress metastasis of human cancer cells.

We screened 1,280 FDA, EMA or other agencies-approved drugs (Prestwick, Inc) in

zebrafish epiboly inhibition assay. The screening showed that 0.9% (12/1280) of the drugs,

including Actimycin A and Tolcapone, induced severe or complete arrest of embryonic cell

movement when embryos were treated with 10μM concentration. 5.2% (66/1280) of the

drugs, such as Dicumarol, Racecadotril, Pizotifen and S(-) Eticlopride hydrochloride, induced

either delayed epiboly or interrupted epiboly of the embryos. 93.3% (1194/1280) of the drugs

had no effect on epiboly. 0.6% (8/1280) of the drugs induced toxic lethality. Epiboly

progression was affected more severely when embryos were treated with 50 μM

concentration (Figure 1A, Supplementary Figure 1E and Supplementary Table 2). Among the

epiboly-interrupting drugs, Adrenosterone, Zardaverine, Racecadotril and Disulfiram have

already been reported to inhibit metastasis-related molecular mechanisms (10-13). This

evidence supports that epiboly-interrupting drugs have the potential to suppress metastasis of

human cancer cells.

We subjected the 78 epiboly-interrupting drugs that showed a suppressor effect on

epiboly progression at a 10μM concentration to *in vitro* examination of whether the drugs

could suppress cell motility and invasion of MDA-MB-231 cells through a Boyden chamber

assay. Sixteen of the 78 drugs strongly affected cell viability and were excluded. The

remaining 62 drugs were subjected to the assay, and twenty of the 62 drugs inhibited cell

motility and invasion of the cells without affecting cell viability (Figure 1B). Among the 20

drugs, three drugs: Hexachlorophene, Nitazoxanide and Ipriflavone were removed since the

primary targets of the drugs, are either not expressed in mammalian cells or unknown. The

known primary targets of the remaining 17 drugs are reported to be expressed by mammalian

cells (Supplementary Table 3). Among the 17 drugs, Pizotifen and S(-) Eticlopride

hydrochloride suppressed cell motility and invasion of several highly metastatic human

cancer cell lines in a dose-dependent manner (Figure 1C and supplementary Figure 2A).

We selected Pizotifen as a candidate drug for blocking metastasis since its primary

target, serotonin receptor 2C (HTR2C), was highly expressed only in metastatic cell lines

(Figure 1D). Furthermore, HTR2C induced EMT on human epithelial cells and promoted

metastatic dissemination of human cancer cells in a zebrafish xenotransplantation model

(Supplementary Figure 3A-E); conversely, pharmacologic and genetic inhibition of HTR2C

suppressed metastatic dissemination of highly metastatic human cancer cells in the model

(Figure 1E and supplementary Figure 2B&C).

Finally, we validated the metastasis-suppressor effect of Pizotifen in a mouse model

of metastasis(14). Luciferase-expressing 4T1 murine mammary carcinoma cells were

inoculated into the mammary fat pads (MFP) of female BALB/c mice. Bioluminescence

imaging and tumor measurement revealed no anti-tumor effect of Pizotifen on the primary

tumor (Figure 2A&B). After 70 days from the inoculation, bioluminescence imaging detected

light emitted in the lungs, livers and lymph nodes of vehicle-treated mice but not those of

Pizotifen-treated mice (Figure 2B). Vehicle-treated mice formed 5 to 50 metastatic nodules

per lung in all 10 mice analyzed; conversely, Pizotifen-treated mice (n=10) formed 0 to 5

nodules per lung in all 10 mice analyzed (Figure 2C). Histological analyses confirmed that

metastatic lesions in the lungs in all vehicle-treated mice; conversely, they were detected in

only 2 Pizotifen-treated mice and the rest 8 mice showed metastatic colony formations

around the bronchiole of the lung. In addition, 4 of 10 vehicle-treated mice exhibited

metastasis in the liver and the rest showed metastatic colony formation around the portal tract

of the liver. In contrast, none of 10 Pizotifen-treated mice showed liver metastases and only 5

mice showed metastatic colony formation around the portal tract (Figure 2D). These results

indicate that Pizotifen can suppress metastasis progression without affecting primary tumor

growth.

To eliminate potential off-target effects of Pizotifen, we confirmed whether

knockdown of HTR2C would show the same effects. Sub-clones of 4T1 cells that expressed

shRNA targeting either LacZ (shLacZ_4T1) or HTR2C (shHTR2C_4T1) were injected into

the MFP of female BALB/c mice. All of the mice (n=5) that were inoculated with

shLacZ_4T1 cells showed metastases in the lungs; conversely, only one of the mice (n=5)

that were inoculated with shHTR2C_4T1 cells showed metastases in the lungs and the rest of

the mice did not. The mean number of metastatic lesions in the lung significantly decreased

to 10% of those of mice that were inoculated with shLacZ_4T1 cells (Figure 2E&F). These

results indicate that pharmacological and genetic inhibition of HTR2C showed an anti-

metastatic effect in the 4T1 model system.

Discussion

Reducing or eliminating mortality associated with metastatic disease is a key goal of medical

oncology, but few models exist that allow for rapid and effective screening of novel compounds

that target metastasis. We developed a unique screening concept utilizing zebrafish gastrulation

to identify anti-metastatic drugs. Our results clearly confirmed our hypothesis: 25.6% (20/76

drugs) of epiboly-interrupting drugs could also suppress cell motility and invasion of highly

metastatic human cell lines in vitro (Figure 1B). A mouse model of metastasis confirmed that

Pizotifen, which is one of the epiboly-interrupting drugs, could suppress metastasis (Figure

2A-C). Thus, this screen could offer a rapid and high-throughput platform for discovery of

anti-metastasis drugs.

There are at least two advantages to the screen described herein. One is that the screen

can easily be converted to a chemical genetic screening platform. Indeed, we have provided

the first evidence that HTR2C, which is a primary target of Pizotifen, induced EMT and

promoted metastatic dissemination of cancer cells (Supplementary Figure 3A-E). In this

research, 1,280 FDA approval drugs were screened, this is less than a few percent of all of

druggable targets (approximately 100 targets) in the human proteome in the body. If chemical

genetic screening using specific inhibitor libraries were conducted, more genes that contribute

to metastasis and gastrulation could be identified. The second advantage is that the screen

enables one researcher to test 100 drugs in 5 hours with using optical microscopy, drugs, and

zebrafish embryos. That indicates this screen is not only highly efficient, low-cost, and low-

labor but also enables researchers who do not have high throughput screening instruments to

conduct drug screening for anti-metastasis drugs.

Materials and Methods

Zebrafish embryo screening

Each drug was added to a well of a 24-well plate containing approximately 20 zebrafish

embryos per well in either 10 μM or 50 μM final concentration when the embryos reached

the sphere stage. Chemical treatment was initiated at 4 hours post-fertilization (hpf) and

approximately 20 embryos were treated with two different concentrations for each compound

tested. The treatment was ended at 9 hpf when vehicle (DMSO) treated embryos as control

reach 80-90% completion of the epiboly stage. The compounds which induced delay (<50%

epiboly) in epiboly were selected as hit compounds for in vitro testing using highly metastatic

human cancer cell lines. The study protocol was approved by the Institutional Animal Care

and Use Committee of the National University of Singapore (protocol number: R16-1068).

Reagents

FDA, EMA and other agencies approved chemical libraries was purchased from Prestwick

Chemical (Illkirch, France). Pizotifen and S(-) Eticlopride hydrochloride were purchased

from Sigma-Aldrich (St. Louis, MO).

Cell culture and cell viability assay

MCF7, MDA-MB-231, MDA-MB-435, MIA-PaCa2, PC3, SW620, PC9 and HaCaT cells

were obtained from American Type Culture Collection (ATCC, Manassas, VA). Luciferase-

expressing 4T1 (4T1-12B) cells were provided from Dr. Gary Sahagian (Tufts University,

Boston, MA). All culture methods followed the supplier's instruction. Cell viability assay

was performed as previously described (Nakayama et al., 2020).

Plasmid

A DNA fragment coding for HTR2C was amplified by PCR with primers containing

restriction enzyme recognition sequences. The HTR2C coding fragment was amplified from

hsp70l:mCherry-T2A-CreERT2 plasmid (11).

Immunoblotting

Western blotting was performed as described previously (Nakayama et al., 2020). Anti-

PRMT1, anti-CYP11A1, anti-E-cadherin, anti-EpCAM, anti-Vimentin, anti-N-cadherin, and

anti-GAPDH antibodies were purchased from Cell signaling Technology. Anti-HTR2C and

anti-DRD2 antibodies were purchased form Abcam.

Flow cytometry

Cells were stained with FITC-conjugated E-cadherin antibody (Biolegend, San Diego, CA).

Flow cytometry was performed as described (Nakayama et al., 2009) and analyzed with

FlowJo software (TreeStar, Ashland, OR).

shRNA mediated gene knockdown

The short hairpin RNA (shRNA)-expressing lentivirus vectors were

constructed using pLVX-shRNA1 vector (Clontech). PRMT1-shRNA_#3–targeting sequence

is GTGTTCCAGTATCTCTGATTA; PRMT1-shRNA_#4–targeting sequence is

TTGACTCCTACGCACACTTTG. CYP11A1-shRNA_#4–targeting sequence is

GCGATTCATTGATGCCATCTA; CYP11A1-shRNA_#4–targeting sequence is

GAAATCCAACACCTCAGCGAT. Human HTR2C-shRNA–targeting sequence is

TCATGCACCTCTGCGCTATAT. Mouse HTR2C-shRNA–targeting sequence is . LacZ-

shRNA– targeting sequence is CTACACAAATCAGCGATT.

Immunofluorescence

Immunofluorescence microscopy assay was performed by previously described (ref). Goat

anti-mouse and goat anti-rabbit immunoglobulin G (IgG) antibodies conjugated to Alexa

Fluor 488 (Life Technologies) and diluted at 1:100 were used. Nuclei were visualized by the

addition of 2µg/ml of 4', 6-diamidino-2-phenylindole (DAPI) and photographed at 100x

magnification by a fluorescent microscope (Zeiss, Germany).

Boyden chamber cell motility and invasion assay

These assays were performed by previously described (Nakayama et al., 2020). In Boyden

chamber assay, either 3x105 MDA-MB-231, 1x106 MDA-MB-435, or 5x105 PC9 cells were

applied to each well in the upper chamber.

Zebrafish xenotransplantation model

Tg(kdrl:eGFP) zebrafish was provided by Dr. Stainier (Max Planck Institute for Heart and

Lung Research). Embryos that were derived from the line were maintained in E3 medium

containing 200µM 1-phenyl-2-thiourea (PTU). Approximately 100-400 Red fluorescence

protein (RFP)-labelled MBA-MB-231 or MIA-Paca2 cells were injected into the duct of

Cuvier of the zebrafish at 2dpf. In this model, the dissemination patterns were generally

divided into three categories: (i) head dissemination, in which disseminated cells exist in the

vessel of the head part; (ii) trunk dissemination, in which the cells were observed in the

vessel dilating from the trunk to the tail; (iii) end-tail dissemination, in which the cells were

observed in the vessel of the end-tail part

Spontaneous metastasis mouse model

4T1-12B cells (2x104) were injected into the #4 mammary fat pad while the mice were

anesthetized. On day two post inoculation, the mice were randomly assigned to two groups

and one group received once daily intraperitoneal injections of 10mg/kg Pizotifen while the

other group received a vehicle injection. To monitor tumor growth and metastases, mice were

imaged biweekly by IVIS Imaging System (ParkinElmer). The primary tumor was resected

10 days after inoculation. The study protocol (protocol number: BRC IACUC #110612) was

approved by A*STAR (Agency for Science, Technology and Research, Singapore).

Histological Analysis

All OCT embedded primary tumors, lungs, and livers of mice from the spontaneous

metastasis 4T1 model were sectioned on a cryostat. Eight micron sections were taken at five

hundred micron intervals through the entirety of the livers and lungs. Sections were

subsequently stained with hematoxylin and eosin. Metastatic lesions were counted under a

microscope in each section for both lungs and livers.

Statistics

Data were analyzed by Student's t test; $p < 0.05$ was considered significant.

References

1.   Welch DR, Hurst DR. Defining the Hallmarks of Metastasis Cancer research.
2019;79(12):3011-27. .
2.   Chaffer CL, Weinberg RA. A perspective on cancer cell metastasis Science.
2011;331(6024):1559-64.
3.   Solnica-Krezel L. Conserved patterns of cell movements during vertebrate
gastrulation Curr Biol. 2005;15(6):R213-28. .
4.   Latimer A, Jessen JR. Extracellular matrix assembly and organization during
zebrafish gastrulation. Matrix Biol. 2010;29(2):89-96.
5.   Solnica-Krezel L. Gastrulation in zebrafish -- all just about adhesion? Curr Opin
Genet Dev. 2006;16(4):433-41.
6.   Tsai Y, Pan H, Hung C. The predominant protein arginine methyltransferase
PRMT1 is critical for zebrafish convergence and extension during gastrulation. FEBS J.
2011;278(6):905-17.
7.   Hsu J, Liang M, Chen C. Pregnenolone stabilizes microtubules and promotes
zebrafish embryonic cell movement. Nature. 2006;439(7075):480-3.
8.   Sack U, Walther W, Scudiero D. Novel effect of antihelminthic Niclosamide on
S100A4-mediated metastatic progression in colon cancer. J Natl Cancer Inst.
2011;103(13):1018-36.
9.   Huang E, Xue S, Zhang X. Vinpocetine inhibits breast cancer cells growth in vitro
and in vivo. . Apoptosis. 2012;17(10):1120-30.
10.   Nakayama J, Lu J, Makinoshima H, Gong Z. A Novel Zebrafish Model of Metastasis
Identifies the HSD11β1 Inhibitor Adrenosterone as a Suppressor of Epithelial-

Mesenchymal Transition and Metastatic Dissemination Mol Cancer Res.
2020;18(3):477-87.
11.   Kolosionek E, Savai R, Ghofrani HA. Expression and activity of phosphodiesterase
isoforms during epithelial mesenchymal transition: the role of phosphodiesterase 4 Mol
Biol Cell. 2009;20(22):4751-65.
12.   Sasaki T, Kuniyasu H, Luo Y. Serum CD10 is associated with liver metastasis in
colorectal cancer J Surg Res. 2014;192(2):390-4.
13.   Liu P, Kumar IS, Brown S, Kannappan V. Disulfiram targets cancer stem-like cells
and reverses resistance and cross-resistance in acquired paclitaxel-resistant triple-
negative breast cancer cells. Br J Cancer. 2013;109(7):1876-85.
14.   Tao K, Fang M, Alroy J, Sahagian GG. Imagable 4T1 model for the study of late
stage breast cancer BMC Cancer. 2008;9(8):228.
294
295

Acknowledgments

We thank Drs. Joshua Collins (NIH/NIDCR) and Diane Palmieri (NIH/NCI) for helping this

research, Dr. Herrick (Albany medical collage) for providing pCMV-h5TH2C-VSV with us.

This study was funded by grants from National Medical Research Council of Singapore and

Ministry of Education of Singapore.

Author Contributions

Design research; J.N. Conducting experiments; J.N. and L.T. Analyzing data: J.N. Writing

the paper; J.N. and Z.G. Funding Acquisition; Z.G., S.W., B.C.G., H.M. and Supervision;

Z.G.

Declaration of Interests

J.N., L.T., B.C.G., S.W., H.M. and Z.G. declare no conflict of interest.

Figure 1

Figure 1. A chemical screen for identification of epiboly-interrupting drugs.

(A) Representative samples of the embryos that were treated with indicated drugs. (B) Effect

of the epiboly-interrupting drugs on cell motility and invasion of MDA-MB-231 cells

through Boyden chamber assays. (C) Effect of pizotifen on cell motility and invasion of

MDA-MB-231, MDA-MB-435 and PC9 cells through Boyden chamber assays. (D) Western

blot analysis of HTR2C levels in various human cancer cell lines with GAPDH as loading

control. (E) Representative images of dissemination of RFP-labelled MDA-MB-231 (231R)

cells in zebrafish xenotransplantation model. The fish that were inoculated with 231R cells,

were treated with either vehicle or Pizotifen. Arrows indicate disseminated 231R cells. The

mean frequencies of the fish showing head, trunk, or end-tail dissemination were counted

(right). Each value is indicated as the mean±SEM of two independent experiments. Statistical

analysis was determined by Student's t test.

Figure 2

Figure 2. Epiboly-interrupting drugs suppressed metastatic progression in a mouse

metastasis model .

(A) Mean volumes (n=10 per group) of 4T1 primary tumors formed in MFP of either vehicle

or Pizotifen-treated mice at day 10 post-injection. (B) Representative images of primary

tumors on day 10 post-injection (top) and metastatic burden on day 70 post-injection

(bottom). (C) Number of metastatic nodules in the lung of either vehicle or Pizotifen-treated

mice. (D) Representative H&E staining of the lung (top) and liver (bottom) from either

vehicle or Pizotifen-treated mice. (E) The mean number of metastatic lesions in step sections

of the lungs from the mice that were inoculated with either shLacZ or shHTR2C_4T1 cells.

(F) Representative H&E staining of the lung and liver from the mice that were inoculated

with 4T1 cells expressing shRNA targeting for either LacZ or HTR2C. Each value is

indicated as the mean±SEM. Statistical analysis was determined by Student's *t* test.

 Supplementary Figure 1

339
340
341

Supplementary Figure 1. Molecular mechanisms of epiboly in zebrafish overlap with

those in cancer metastasis.

(A) Western blot analysis of PRMT1 (upper) and CYP11A1 (middle) protein levels in non-

metastatic human cancer cell lines, (MCF10, MCF7, MDA-MB-453, Sun159, SKBr3) and

highly metastatic human cancer cell lines, (BT549 and MDA-MB-231); β-actin loading

control is shown (bottom). (B) Knockdown of PRMT1 or CYP11A1 in MDA-MB-231 cells.

MBA-MB-231 cells were transfected with a control shRNA targeting LacZ, and one of four

independent shRNAs targeting PRMT1 (clone #1 to #4) or one of two independent shRNAs

targeting CYP11A1 (clone #1 to #4). Reduced PRMT1 and CYP11A1 expression,

determined by western blot, in sub-cell lines of MDA-MB-231 cells expressing PRMT1

shRNA (clone #3 and #4 or CYP11A1 (clone #2 and #4), compared with controls (parental

cell line MDA-MB-231 and control shRNA cells); β-actin levels shown as a loading control.

(C) Effect of shRNAs targeting either PRMT1 or CYP11A1 on cell motility and invasion of

MBA-MB-231 cells. Parental MDA-MB-231 cells and four sub-cell lines of MDA-MB-231

cells that were transfected with either shRNA targeting either LacZ, two independent

shRNAs targeting PRMT1 (clone #3 and #4) or two independent shRNAs targeting

CYP11A1 (clone #2 and #4), were subjected to Boyden chamber assays. (D) Zebrafish

embryos treated with either vehicle (DMSO), 10μM Niclosamide, or 50μM vinpocetine.

Approximately 20 embryos were treated with either DMSO as a vehicle control, niclosamide,

or vinpocetine. The treatment was started at 4 hpf when all of embryos reached sphere stage

and ended at 9 hpf when control embryos reached 80-90% epiboly stage. Each experiment

was performed at least twice. Statistical analysis was determined by Student's *t* test. (E)

Cumulative results of the chemical screen in which each drug was used at either 10μM (left) or

50μM (right) concentrations. 1,280 FDA, EMA or other agencies-approved drugs were subjected

to this screening. Positive "hit" drugs were those that interrupted epiboly progression.

368
Supplementary Figure 2

Supplementary Figure 2. Epiboly-interrupting drugs suppressed metastatic dissemination

of human cancer cells in a zebrafish xenotransplantation model.

(A) Effect of S (-) Eticlopride hydrochloride on cell motility and invasion of MBA-MB-231,

MDA-MB-435, and PC9 cells. Either vehicle or pizotifen treated cells were subjected to Boyden

chamber assays. Fetal bovine serum (1%v/v) was used as the chemoattractant in both assays.

Each experiment was performed at least twice. Statistical analysis was determined by Student's *t*

test. (B) and (C) Representative images of dissemination of RFP-labelled MIA-PaCa2, RFP-

labelled MDA-MB-231 cells expressing shRNA targeting either LacZ (shLacZ_231R) or HTR2C

(shHTR2C_231R) cells in zebrafish xenotransplantation model. The fish larvae that were

inoculated with RFP-labelled MIA-PaCa2 cells, were treated with either vehicle (top) or the drug

(lower) (B). The fish larvae that were inoculated with either shLacZ_231R (top) or

shHTR2C_231R cells (lower) (C). Arrows indicate disseminated RFP-labelled MIA-PaCa2 or

231R cells. The images were shown in 4x magnification. Scale bar, 100μm. The mean

frequencies of the fish showing head, trunk, or end-tail dissemination were counted (graph on

right). Each value is indicated as the mean ± SEM of two independent experiments. Statistical

analysis was determined by Student's t test.

   Supplementary Figure 3

Supplementary Figure 3. HTR2C induced EMT-mediated metastatic dissemination of

human cancer cells.

(A) The morphologies of the MCF7 and HaCaT cells expressing either the control vector or

HTR2C were revealed by phase contrast microscopy. (B) Immunofluorescence staining of E-

cadherin, EpCAM, Vimentin, and N-cadherin expressions in the MCF7 cells from

Supplementary Figure 3A. (C) Expression of E-cadherin, EpCAM, Vimentin, N-cadherin, and

HTR2C was examined by western-blotting in the MCF7 and HaCaT cells; GAPDH loading

control is shown (bottom). (D) Effect of HTR2C on cell motility and invasion of MCF7 cells.

MCF7 cells were subjected to Boyden chamber assays. Fetal bovine serum (1%v/v) was used as

the chemoattractant in both assays. Each experiment was performed at least twice. (E)

Representative images of dissemination patterns of MCF7 cells expressing either the control

vector (top left) or HTR2C (lower left) in a zebrafish xenotransplantation model. White arrows

head indicate disseminated MCF7 cells. The mean frequencies of the fish showing head, trunk, or

end-tail dissemination tabulated (right). Each value is indicated as the mean ± SEM of two

independent experiments. Statistical analysis was determined by Student's *t* test.

Supplementary Table 1. A list of the genes that are involved between gastrulation and metastasis progression

Genes	Gastrulation Defects	Ref	Effects in Metastasis	Ref
BMP	Convergence and extension	(1)	EMT	(2)
WNT	Convergence and extension	(3)	Migration and Invasion	(4)
FGF	Convergence and extension	(5)	Invision	(6)
EGF	Epiboly	(7)	Migration	(8)
PDGF	Convergence and extension	(9)	EMT	(10)
CXCL12	Migration of endodermal cells	(11)	Migration and Invasion	(12)
CXCR4	Migration of endodermal cells	(11)	Migration and Invasion	(12)
PIK3CA	Convergence and extension	(13)	Migration and Invasion	(14)
YES	Epiboly	(15)	Migration	(16)
FYN	Epiboly	(17)	Migration and Invasion	(18)
MAPK1	Epiboly	(19)	Migration	(20)
SHP2	Convergence and extension	(21)	Migration	(22)
SNAI1	Convergence and extension	(23)	EMT	(24)
SNAI2	Mesoderm & Neural crest formation	(25)	EMT	(24)
TWIST1	Mesoderm formation	(26)	EMT	(27)
TBXT	Convergence and extension	(3)	EMT	(28)
ZEB1	Epiboly	(29)	EMT	(30)
GSC	Mesodermal patterning	(31)	EMT	(32)
FOXC2	Unclear, defects in gastrulation	(33)	EMT	(34)
STAT3	Convergence and extension	(35)	Migration	(36)
POU5F1	Epiboly	(37)	EMT	(38)
EZH2	Unclear, defects in gastrulation	(39)	Invasion	(40)
EHMT2	Defects in Neurogenesis	(41)	Migration and Invasion	(42)
BMI1	Defects in skelton formation	(43)	EMT	(44)
RHOA	Convergence and extension	(45)	Migration and Invasion	(46)
CDC42	Convergence and extension	(47)	Migration and Invasion	(48)
RAC1	Convergence and extension	(49)	Migration and Invasion	(50)
ROCK2	Convergence and extension	(51)	Migration and Invasion	(52)
PAR1	Convergence and extension	(53)	Migration	(54)
PRKCI	Convergence and extension	(53)	EMT	(55)
CAP1	Convergence and extension	(56)	Migration	(57)
EZR	Epiboly	(58)	Migration	(59)
EPCAM	Epiboly	(60)	Migration and Invasion	(61)
ITGB1 / ITA5	Mesodermal Migration	(62)	Migration and Invasion	(63)
FN1	Convergence and extension	(64)	Invasion	(65)
HAS2	Dorsal migration of lateral cells	(66)	Invasion	(67)
MMP14	Convergence and extension	(68)	Invasion	(69)
COX1	Epiboly	(70)	Invasion	(71)
PTGES	Convergence and extension	(72)	Invasion	(73)
SLC39A6	Aterior migration	(74)	EMT	(75)
GNA12 / 13	Convergence and extension	(41)	Migration and Invasion	(76)
OGT	Epiboly	(77)	Migration and Invasion	(78)
CCN1	Cell Movement	(79)	Migration and Invasion	(80)
TRPM7	Convergence and extension	(81)	Migration	(82)
MAPKAPK2	Epiboly	(83)	Migration	(84)
B4GALT1	Convergence and extension	(85)	Invasion	(86)
IER2	Convergence and extension	(87)	Migration	(88)
TIP1	Convergence and extension	(89)	Migration and Invasion	(90)
PAK5	Convergence and extension	(91)	Migration	(92)
MARCKS	Convergence and extension	(93)	Migration and Invasion	(94)

Supplementary Table 1.

A list of the fifty genes that play essential role in governing both metastasis and gastrulation

progression. The gastrulation defects in Xenopus or zebrafish that are induced by knockdown

415   of each of these genes, were indicated. The molecular mechanism in metastasis that are

416   inhibited by knockdown of each of the same genes, were indicated.

417   Supplementary Table 2. A list of the drugs that interfere with epiboly progression in
418   zebrafish

Chemical name	Chemical formula	Effect of 10uM	Effect of 50uM
Acitretin	C21H26O3	Delayed	Delayed
Adrenosterone	C19H24O3	Delayed	Delayed
Albendazole	C12H15N3O2S	Severe delayed	Severe delayed
Alfadolone acetate	C23H34O5	Delayed	Delayed
Alfaxalone	C21H32O3	Delayed	Delayed
Alprostadil	C20H34O5	Delayed	Delayed
Altrenogest	C21H26O2	Slightly delayed	Delayed
Ampiroxicam	C20H21N3O7S	Non effect	Delayed
Anethole-trithione	C10H8OS3	Delayed	Delayed
Antimycin A	C28H40N2O9	Delayed	Delayed
Avobenzone	C20H22O3	Delayed	Delayed
Benzoxiquine	C16H11NO2	Non effect	Delayed
Bosentan	C27H29N5O6S	Delayed	Delayed
Butoconazole nitrate	C19H18Cl3N3O3S	Delayed	Toxic lethal
Camptothecine (S,+)	C20H16N2O4	Severe delayed	Severe delayed
Carbenoxolone disodium salt	C34H48Na2O7	Delayed	Toxic lethal
Carmofur	C11H16FN3O3	Slightly delayed	Delayed
Carprofen	C15H12ClNO2	Severe delayed	Toxic lethal
Cefdinir	C14H13N5O5S2	Delayed	Delayed
Celecoxib	C17H14F3N3O2S	Delayed	Delayed
Chlorambucil	C14H19Cl2NO2	Slightly delayed	Delayed
Chlorhexidine	C22H30Cl2N10	Non effect	Toxic lethal
Ciclopirox ethanolamine	C14H24N2O3	Delayed	Severe delayed
Cinoxacin	C12H10N2O5	Delayed	Severe delayed
Clofibrate	C12H15ClO3	Non effect	Severe delayed
Clopidogrel	C16H16ClNO2S	Non effect	Delayed
Clorgyline hydrochloride	C13H16Cl3NO	Delayed	Delayed
Colchicine	C22H25NO6	Non effect	Delayed
Deptropine citrate	C29H35NO8	Delayed	Delayed
Desipramine hydrochloride	C18H23ClN2	Delayed	Delayed
Diclofenac sodium	C14H10Cl2NNaO2	Delayed	Severe delayed
Dicumarol	C19H12O6	Delayed	Severe delayed
Diethylstilbestrol	C18H20O2	Delayed	Toxic lethal
Dimaprit dihydrochloride	C6H17Cl2N3S	Slightly delayed	Delayed
Disulfiram	C10H20N2S4	Delayed	Delayed
Dopamine hydrochloride	C8H12ClNO2	Delayed	Delayed
Eburnamonine (-)	C19H22N2O	Delayed	Delayed
Ethaverine hydrochloride	C24H30ClNO4	Delayed	Delayed
Ethinylestradiol	C20H24O2	Delayed	Severe delayed
Ethopropazine hydrochloride	C19H25ClN2S	Delayed	Delayed
Ethoxyquin	C14H19NO	Non effect	Delayed
Exemestane	C20H24O2	Slightly delayed	Delayed
Ezetimibe	C24H21F2NO3	Slightly delayed	Delayed
Fenbendazole	C15H13N3O2S	Non effect	Delayed
Fenoprofen calcium salt dihydrate	C30H30CaO8	Slightly delayed	Delayed
Fentiazac	C17H12ClNO2S	Toxic lethal	Toxic lethal
Floxuridine	C9H11FN2O5	Delayed	Toxic lethal
Flunixin meglumine	C21H28F3N3O7	Delayed	Toxic lethal
Flutamide	C11H11F3N2O3	Delayed	Toxic lethal
Fluticasone propionate	C25H31F3O5S	Non effect	Delayed
Furosemide	C12H11ClN2O5S	Delayed	Delayed
Gatifloxacin	C19H22FN3O4	Delayed	Delayed
Gemcitabine	C9H11F2N3O4	Delayed	Delayed
Gemfibrozil	C15H22O3	Delayed	Toxic lethal
Gestrinone	C21H24O2	Delayed	Delayed
Haloprogin	C9H4Cl3IO	Delayed	Toxic lethal
Hexachlorophene	C13H6Cl6O2	Delayed	Severe delayed
Hexestrol	C18H22O2	Slightly delayed	Delayed
Ibudilast	C14H18N2O	Non effect	Delayed

Idazoxan hydrochloride	C11H13ClN2O2	Slightly delayed	Delayed
Idazoxan hydrochloride	C11H13ClN2O2	Non effect	Delayed
Idebenone	C19H30O5	Severe delayed	Toxic lethal
Indomethacin	C19H16ClNO4	Non effect	Delayed
Ipriflavone	C18H16O3	Delayed	Severe delayed
Isotretinoin	C20H28O2	Non effect	Severe delayed
Isradipine	C19H21N3O5	Non effect	Delayed
Lansoprazole	C16H14F3N3O2S	Slightly delayed	Delayed
Latanoprost	C26H40O5	Non effect	Delayed
Leflunomide	C12H9F3N2O2	Delayed	Severe delayed
Letrozole	C17H11N5	Non effect	Delayed
Lithocholic acid	C24H40O3	Non effect	Delayed
Lodoxamide	C11H6ClN3O6	Non effect	Delayed
Lofepramine	C26H27ClN2O	Non effect	Delayed
Loratadine	C22H23ClN2O2	Delayed	Delayed
Loxapine succinate	C22H24ClN3O5	Delayed	Delayed
Mebendazole	C16H13N3O3	Severe delayed	Severe delayed
Mebendazole	C22H26N2O2	Non effect	Delayed
Meloxicam	C14H13N3O4S2	Delayed	Toxic lethal
Methiazole	C12H15N3O2S	Delayed	Delayed
Mevastatin	C23H34O5	Non effect	Delayed
MK 801 hydrogen maleate	C20H19NO4	Slightly delayed	Delayed
Nabumetone	C15H16O2	Non effect	Severe delayed
Naftopidil dihydrochloride	C24H30Cl2N2O3	Slightly delayed	Delayed
Nandrolone	C18H26O2	Delayed	Delayed
Naproxen sodium salt	C14H13NaO3	Delayed	Delayed
Niclosamide	C13H8Cl2N2O4	Delayed	Delayed
Nifekalant	C19H27N5O5	Delayed	Delayed
Niflumic acid	C13H9F3N2O2	Delayed	Delayed
Nimesulide	C13H12N2O5S	Non effect	Delayed
Nisoldipine	C20H24N2O6	Delayed	Toxic lethal
Nitazoxanide	C12H9N3O5S	Severe delayed	Severe delayed
Norethindrone	C20H26O2	Non effect	Delayed
Norgestimate	C23H31NO3	Slightly delayed	Delayed
Oxfendazol	C15H13N3O3S	Slightly delayed	Delayed
Oxibendazol	C12H15N3O3	Severe delayed	Severe delayed
Oxymetholone	C21H32O3	Slightly delayed	Delayed
Parbendazole	C13H17N3O2	Severe delayed	Severe delayed
Parthenolide	C15H20O3	Non effect	Delayed
Penciclovir	C10H15N5O3	Non effect	Delayed
Pentobarbital	C11H18N2O3	Non effect	Delayed
Phenazopyridine hydrochloride	C11H12ClN5	Delayed	Toxic lethal
Phenothiazine	C12H9NS	Non effect	Delayed
Phenoxybenzamine hydrochloride	C18H23Cl2NO	Non effect	Delayed
Pizotifen malate	C23H27NO5S	Delayed	Severe delayed
Pramoxine hydrochloride	C17H28ClNO3	Slightly delayed	Delayed
Prilocaine hydrochloride	C13H21ClN2O	Non effect	Delayed
Primidone	C12H14N2O2	Slightly delayed	Delayed
Racecadotril	C21H23NO4S	Slightly delayed	Delayed
Riluzole hydrochloride	C8H6ClF3N2OS	Non effect	Delayed
Ritonavir	C37H48N6O5S2	Non effect	Severe delayed
S(-)Eticlopride hydrochloride	C17H26Cl2N2O3	Delayed	Delayed
Salmeterol	C25H37NO4	Non effect	Delayed
Streptomycin sulfate	C42H84N14O36S3	Non effect	Delayed
Sulconazole nitrate	C18H16Cl3N3O3S	Delayed	Delayed
Tegafur	C8H9FN2O3	Delayed	Delayed
Telmisartan	C33H30N4O2	Severe delayed	Toxic lethal
Tenatoprazole	C16H18N4O3S	Non effect	Delayed
Terbinafine	C21H25N	Non effect	Delayed
Thimerosal	C9H9HgNaO2S	Non effect	Delayed
Thiorphan	C12H15NO3S	Delayed	Delayed
Tolcapone	C14H11NO5	Severe delayed	Severe delayed
Topotecan	C23H23N3O5	Delayed	Delayed
Tracazolate hydrochloride	C16H25ClN4O2	Severe delayed	Delayed

Tribenoside	C29H34O6	Delayed	Delayed
Triclabendazole	C14H9Cl3N2OS	Delayed	Delayed
Triclosan	C12H7Cl3O2	Delayed	Severe delayed
Trioxsalen	C14H12O3	Delayed	Delayed
Troglitazone	C24H27NO5S	Severe delayed	Toxic lethal
Valproic acid	C8H16O2	Non effect	Delayed
Voriconazole	C16H14F3N5O	Non effect	Delayed
Zardaverine	C12H10F2N2O3	Slightly delayed	Delayed
Zuclopenthixol dihydrochloride	C22H27Cl3N2OS	Delayed	Delayed

Supplementary Table 2. A list of the drugs that interfere with epiboly progression in

zebrafish

A list of positive "hit" drugs that interfered with epiboly progression. Gastrulation defects or

status of each of the zebrafish embryos that were treated with either 10μM or 50μM

concentrations, are indicated.

Supplementary Table 3. Primary targets of the identified drugs
428

The identified drugs	Primary targets of the identified drugs
Hexachlorophene	D-lactate dehydrogenase (D-LDH), not expressed in mammalian cells
Troglitazone	Agonist for Peroxisome proliferator-activated receptor α and γ (PPARα and γ)
Pizotifen malate	5-hydroxytryptamine receptor 2C (HTR2C)
Salmeterol	Adrenergic receptor beta 2 (ADRB2)
Nitazoxanide	Pyruvate ferredoxin oxidoreductase (PFOR), not expressed in mammalian cells
Valproic acid	Histone deacetylases (HDACs)
Dicumarol	NAD(P)H dehydrogenase quinone 1 (NQO1)
Loxapine succinate	Dopamine receptor D2 and D4 (DRD2 and DRD4)
Adrenosterone	Hydroxysteroid (11-beta) dehydrogenase 1 (HSD11β1)
Riluzole hydrochloride	Glutamate R and Voltage-dependent Na+ channel
Naftopidil dihydrochloride	5-hydroxytryptamine receptor 1A (HTR1A) and α1-adrenergic receptor (AR)
S(-)Eticlopride hydrochloride	Dopamine receptor D2 (DRD2)
Racecadotril	Membrane metallo-endopeptidase (MME)
Ipriflavone	Unknow
Flurbiprofen	Cyclooxygenase 1 and 2 (Cox1 and 2)
Zardaverine	Phosphodiesterase III/IV (PDE3/4)
Leflunomide	Dihydroorotate dehydrogenase (DHODH)
Olmesartan	Angiotensin II receptor alpha
Disulfiram	Aldehyde dehydrogenase (ALDH) Dopamine β-hydroxylase (DBH)
Zuclopenthixol dihydrochloride	Dopamine receptors D1 and D2 (DRD1 and 2)

References for supplemental table 1

1.   Kondo M. Bone morphogenetic proteins in the early development of zebrafish. FEBS J. 2007;274(12):2960-7.

2.   Katsuno Y, Hanyu A, Kanda H, Ishikawa Y, Akiyama F, Iwase T, et al. Bone morphogenetic protein signaling enhances invasion and bone metastasis of breast cancer cells through Smad pathway. Oncogene. 2008;27(49):6322-33.

3.   Tada M, Smith JC. Xwnt11 is a target of Xenopus Brachyury: regulation of gastrulation movements via Dishevelled, but not through the canonical Wnt pathway. Development. 2000;127:2227-38.

4.   Vincan E, Barker N. The upstream components of the Wnt signalling pathway in the dynamic EMT and MET associated with colorectal cancer progression. Clin Exp Metastasis. 2008;25(6):657-63.

5.   Yang X, Dormann D, Munsterberg AE, Weijer CJ. Cell movement patterns during gastrulation in the chick are controlled by positive and negative chemotaxis mediated by FGF4 and FGF8. Dev Cell. 2002;3(3):425-37.

6.   Nomura S, Yoshitomi H, Takano S, Shida T, Kobayashi S, Ohtsuka M, et al. FGF10/FGFR2 signal induces cell migration and invasion in pancreatic cancer. Br J Cancer. 2008;99(2):305-13.

7.   Song S, Eckerle S, Onichtchouk D, Marrs JA, Nitschke R, Driever W. Pou5f1-dependent EGF expression controls E-cadherin endocytosis, cell adhesion, and zebrafish epiboly movements. Dev Cell. 2013;24(5):486-501.

8.   Lu Z, Jiang G, Blume-Jensen P, Hunter T. Epidermal growth factor-induced tumor cell invasion and metastasis initiated by dephosphorylation and downregulation of focal adhesion kinase. Mol Cell Biol. 2001;21(12):4016-31.

9.   Damm EW, Winklbauer R. PDGF-A controls mesoderm cell orientation and radial intercalation during Xenopus gastrulation. Development. 2011;138(3):565-75.

10.   Jechlinger M, Sommer A, Moriggl R, Seither P, Kraut N, Capodiecci P, et al. Autocrine PDGFR signaling promotes mammary cancer metastasis. J Clin Invest. 2006;116(6):1561-70.

11.   Mizoguchi T, Verkade H, Heath JK, Kuroiwa A, Kikuchi Y. Sdf1/Cxcr4 signaling controls the dorsal migration of endodermal cells during zebrafish gastrulation. Development. 2008;135(15):2521-9.

12.   Shen HB, Gu ZQ, Jian K, Qi J. CXCR4-mediated Stat3 activation is essential for CXCL12-induced cell invasion in bladder cancer. Tumour Biol. 2013;34(3):1839-45.

13.   Montero J-A, Kilian B, Chan J, Bayliss PE, Heisenberg C-P. Phosphoinositide 3-Kinase Is Required for Process Outgrowth and Cell Polarization of Gastrulating Mesendodermal Cells. Current Biology. 2003;13(15):1279-89.

14.   Wander SA, Zhao D, Besser AH, Hong F, Wei J, Ince TA, et al. PI3K/mTOR inhibition can impair tumor invasion and metastasis in vivo despite a lack of antiproliferative action in vitro: implications for targeted therapy. Breast Cancer Res Treat. 2013;138(2):369-81.

15.   Tsai WB, Zhang X, Sharma D, Wu W, Kinsey WH. Role of Yes kinase during early zebrafish development. Dev Biol. 2005;277(1):129-41.

16.   Barraclough J, Hodgkinson C, Hogg A, Dive C, Welman A. Increases in c-Yes Expression Level and Activity Promote Motility But Not Proliferation of Human Colorectal Carcinoma Cells. Neoplasia. 2007;9(9):745-IN32.

17.   Sharma D, Holets L, Zhang X, Kinsey WH. Role of Fyn kinase in signaling
associated with epiboly during zebrafish development. Dev Biol.
2005;285(2):462-76.
18.   Yadav V, Denning MF. Fyn is induced by Ras/PI3K/Akt signaling and is
required for enhanced invasion/migration. Mol Carcinog. 2011;50(5):346-52.
19.   Krens SF, He S, Lamers GE, Meijer AH, Bakkers J, Schmidt T, et al. Distinct
functions for ERK1 and ERK2 in cell migration processes during zebrafish
gastrulation. Dev Biol. 2008;319(2):370-83.
20.   Radtke S, Milanovic M, Rossé C. ERK2 but not ERK1 mediates HGF-
induced motility in non-small cell lung carcinoma cell lines. J Cell Sci.
2013;126(Pt11):2381-91.
21.   Jopling C, van Geemen D, den Hertog J. Shp2 knockdown and
Noonan/LEOPARD mutant Shp2-induced gastrulation defects. PLoS Genet.
2007;3(12):e225.
22.   Wang FM, Liu HQ, Liu SR, Tang SP, Yang L, Feng GS. SHP-2 promoting
migration and metastasis of MCF-7 with loss of E-cadherin, dephosphorylation of
FAK and secretion of MMP-9 induced by IL-1beta in vivo and in vitro. Breast
Cancer Res Treat. 2005;89(1):5-14.
23.   Ip YT, Gridley T. Cell movements during gastrulation: snail dependent and
independent pathways. Curr Opin Genet Dev. 2002;12(4):423-9.
24.   Batlle E, Sancho E, Francí C, Domínguez D, Monfar M. The transcription
factor snail is a repressor of E-cadherin gene expression in epithelial tumour
cells. Nat Cell Biol. 2000;2(2):84-9.
25.   Shi J, Severson C, Yang J, Wedlich D, Klymkowsky MW. Snail2 controls
mesodermal BMP/Wnt induction of neural crest. Development.
2011;138(15):3135-45.
26.   Castanon I, Baylies MK. A Twist in fate: evolutionary comparison of Twist
structure and function. Gene. 2002;287(1-2):11-22.
27.   Yang J, Mani SA, Donaher JL, Ramaswamy S, Itzykson RA, Come C, et al.
Twist, a master regulator of morphogenesis, plays an essential role in tumor
metastasis. Cell. 2004;117(7):927-39.
28.   Fernando RI, Litzinger M, Trono P, Hamilton DH, Schlom J, Palena C. The
T-box transcription factor Brachyury promotes epithelial-mesenchymal
transition in human tumor cells. J Clin Invest. 2010;120(2):533-44.
29.   Vannier C, Mock K, Brabletz T, Driever W. Zeb1 regulates E-cadherin and
Epcam (epithelial cell adhesion molecule) expression to control cell behavior in
early zebrafish development. J Biol Chem. 2013;288(26):18643-59.
30.   Spaderna S, Schmalhofer O, Wahlbuhl M, Dimmler A, Bauer K, Sultan A, et
al. The transcriptional repressor ZEB1 promotes metastasis and loss of cell
polarity in cancer. Cancer Res. 2008;68(2):537-44.
31.   Sander V, Reversade B, De Robertis EM. The opposing homeobox genes
Goosecoid and Vent1/2 self-regulate Xenopus patterning. EMBO J.
2007;26(12):2955-65.
32.   Hartwell KA, Muir B, Reinhardt F, Carpenter AE, Sgroi DC, Weinberg RA.
The Spemann organizer gene, Goosecoid, promotes tumor metastasis. Proc Natl
Acad Sci U S A. 2006;103(50):18969-74.
33.   Wilm B, James RG, Schultheiss TM, Hogan BL. The forkhead genes, Foxc1
and Foxc2, regulate paraxial versus intermediate mesoderm cell fate. Dev Biol.
2004;271(1):176-89.
34.   Mani SA, Yang J, Brooks M, Schwaninger G, Zhou A, Miura N, et al.
Mesenchyme Forkhead 1 (FOXC2) plays a key role in metastasis and is associated

with aggressive basal-like breast cancers. Proc Natl Acad Sci U S A.
2007;104(24):10069-74.
35.   Miyagi C, Yamashita S, Ohba Y, Yoshizaki H, Matsuda M, Hirano T. STAT3
noncell-autonomously controls planar cell polarity during zebrafish convergence
and extension. J Cell Biol. 2004;166(7):975-81.
36.   Abdulghani J, Gu L, Dagvadorj A, Lutz J, Leiby B, Bonuccelli G, et al. Stat3
promotes metastatic progression of prostate cancer. Am J Pathol.
2008;172(6):1717-28.
37.   Lachnit M, Kur E, Driever W. Alterations of the cytoskeleton in all three
embryonic lineages contribute to the epiboly defect of Pou5f1/Oct4 deficient
MZspg zebrafish embryos. Dev Biol. 2008;315(1):1-17.
38.   Dai X, Ge J, Wang X, Qian X, Zhang C, Li X. OCT4 regulates epithelial-
mesenchymal transition and its knockdown inhibits colorectal cancer cell
migration and invasion. Oncol Rep. 2013;29(1):155-60.
39.   O'Carroll D, Erhardt S, Pagani M, Barton SC, Surani MA, Jenuwein T. The
polycomb-group gene Ezh2 is required for early mouse development. Mol Cell
Biol. 2001;21(13):4330-6.
40.   Ren G, Baritaki S, Marathe H, Feng J, Park S, Beach S, et al. Polycomb
protein EZH2 regulates tumor invasion via the transcriptional repression of the
metastasis suppressor RKIP in breast and prostate cancer. Cancer Res.
2012;72(12):3091-104.
41.   Lin F, Sepich DS, Chen S, Topczewski J, Yin C, Solnica-Krezel L, et al.
Essential roles of G{alpha}12/13 signaling in distinct cell behaviors driving
zebrafish convergence and extension gastrulation movements. J Cell Biol.
2005;169(5):777-87.
42.   Chen MW, Hua KT, Kao HJ, Chi CC, Wei LH, Johansson G, et al. H3K9
histone methyltransferase G9a promotes lung cancer invasion and metastasis by
silencing the cell adhesion molecule Ep-CAM. Cancer Res. 2010;70(20):7830-40.
43.   van der Lugt NM, Domen J, Linders K, van Roon M, Robanus-Maandag E, te
Riele H, et al. Posterior transformation, neurological abnormalities, and severe
hematopoietic defects in mice with a targeted deletion of the bmi-1 proto-
oncogene. Genes Dev. 1994;8(7):757-69.
44.   Guo BH, Feng Y, Zhang R, Xu LH, Li MZ, Kung HF, et al. Bmi-1 promotes
invasion and metastasis, and its elevated expression is correlated with an
advanced stage of breast cancer. Mol Cancer. 2011;10(1):10.
45.   Zhu S, Liu L, Korzh V, Gong Z, Low BC. RhoA acts downstream of Wnt5 and
Wnt11 to regulate convergence and extension movements by involving effectors
Rho kinase and Diaphanous: use of zebrafish as an in vivo model for GTPase
signaling. Cell Signal. 2006;18(3):359-72.
46.   Yoshioka K, Nakamori S, Itoh K. Overexpression of small GTP-binding
protein RhoA promotes invasion of tumor cells. Cancer Res. 1999;59(8):2004-10.
47.   Choi SC, Han JK. Xenopus Cdc42 regulates convergent extension
movements during gastrulation through Wnt/Ca2+ signaling pathway. Dev Biol.
2002;244(2):342-57.
48.   Reymond N, Im JH, Garg R, Vega FM, Borda d'Agua B, Riou P, et al. Cdc42
promotes transendothelial migration of cancer cells through beta1 integrin. J Cell
Biol. 2012;199(4):653-68.
49.   Habas R, Dawid IB, He X. Coactivation of Rac and Rho by Wnt/Frizzled
signaling is required for vertebrate gastrulation. . Genes Dev. 2003;17(2):295-
309.

50. Vega FM, Ridley AJ. Rho GTPases in cancer cell biology. FEBS Lett.
2008;582(14):2093-101.
51. Marlow F, Topczewski J, Sepich D, Solnica-Krezel L. Zebrafish Rho kinase 2
acts downstream of Wnt11 to mediate cell polarity and effective convergence
and extension movements. Curr Biol. 2002;12(11):876-84.
52. Itoh K, Yoshioka K, Akedo H, Uehata M, Ishizaki T, Narumiya S. An
essential part for Rho-associated kinase in the transcellular invasion of tumor
cells. Nat Med. 1999;5(2):221-5.
53. Kusakabe M, Nishida E. The polarity-inducing kinase Par-1 controls
Xenopus gastrulation in cooperation with 14-3-3 and aPKC. EMBO J.
2004;23(21):4190-201.
54. Shi X, Gangadharan B, Brass LF, Ruf W, Mueller BM. Protease-activated
receptors (PAR1 and PAR2) contribute to tumor cell motility and metastasis. Mol
Cancer Res. 2004;2(7):395-402.
55. Gunaratne A, Thai BL, Di Guglielmo GM. Atypical protein kinase C
phosphorylates Par6 and facilitates transforming growth factor beta-induced
epithelial-to-mesenchymal transition. Mol Cell Biol. 2013;33(5):874-86.
56. Seifert K, Ibrahim H, Stodtmeister T, Winklbauer R, Niessen CM. An
adhesion-independent, aPKC-dependent function for cadherins in morphogenetic
movements. J Cell Sci. 2009;122(Pt 14):2514-23.
57. Yamazaki K, Takamura M, Masugi Y, Mori T, Du W, Hibi T, et al. Adenylate
cyclase-associated protein 1 overexpressed in pancreatic cancers is involved in
cancer cell motility. Lab Invest. 2009;89(4):425-32.
58. Link V, Carvalho L, Castanon I, Stockinger P, Shevchenko A, Heisenberg CP.
Identification of regulators of germ layer morphogenesis using proteomics in
zebrafish. J Cell Sci. 2006;119(Pt 10):2073-83.
59. Khanna C, Wan X, Bose S, Cassaday R, Olomu O, Mendoza A, et al. The
membrane-cytoskeleton linker ezrin is necessary for osteosarcoma metastasis.
Nat Med. 2004;10(2):182-6.
60. Slanchev K, Carney TJ, Stemmler MP, Koschorz B, Amsterdam A, Schwarz
H, et al. The epithelial cell adhesion molecule EpCAM is required for epithelial
morphogenesis and integrity during zebrafish epiboly and skin development.
PLoS Genet. 2009;5(7):e1000563.
61. Ni J, Cozzi PJ, Duan W, Shigdar S, Graham PH, John KH, et al. Role of the
EpCAM (CD326) in prostate cancer metastasis and progression. Cancer
Metastasis Rev. 2012;31(3-4):779-91.
62. Skalski M, Alfandari D, Darribere T. A key function for alphav containing
integrins in mesodermal cell migration during Pleurodeles waltl gastrulation.
Dev Biol. 1998;195(2):158-73.
63. Felding-Habermann B. Integrin adhesion receptors in tumor metastasis.
Clin Exp Metastasis. 2003;20(3):203-13.
64. Marsden M, DeSimone DW. Integrin-ECM Interactions Regulate Cadherin-
Dependent Cell Adhesion and Are Required for Convergent Extension in
Xenopus. Current Biology. 2003;13(14):1182-91.
65. Malik G, Knowles LM, Dhir R, Xu S, Yang S, Ruoslahti E, et al. Plasma
fibronectin promotes lung metastasis by contributions to fibrin clots and tumor
cell invasion. Cancer Res. 2010;70(11):4327-34.
66. Bakkers J, Kramer C, Pothof J, Quaedvlieg NE, Spaink HP, Hammerschmidt
M. Has2 is required upstream of Rac1 to govern dorsal migration of lateral cells
during zebrafish gastrulation. Development. 2004;131(3):525-37.

67.   Kim HR, Wheeler MA, Wilson CM, Iida J, Eng D, Simpson MA, et al.
Hyaluronan facilitates invasion of colon carcinoma cells in vitro via interaction
with CD44. Cancer Res. 2004;64(13):4569-76.
68.   Coyle RC, Latimer A, Jessen JR. Membrane-type 1 matrix
metalloproteinase regulates cell migration during zebrafish gastrulation:
evidence for an interaction with non-canonical Wnt signaling. Exp Cell Res.
2008;314(10):2150-62.
69.   Perentes JY, Kirkpatrick ND, Nagano S, Smith EY, Shaver CM, Sgroi D, et al.
Cancer cell-associated MT1-MMP promotes blood vessel invasion and distant
metastasis in triple-negative mammary tumors. Cancer Res. 2011;71(13):4527-
38.
70.   Cha YI, Kim SH, Sepich D, Buchanan FG, Solnica-Krezel L, DuBois RN.
Cyclooxygenase-1-derived PGE2 promotes cell motility via the G-protein-coupled
EP4 receptor during vertebrate gastrulation. Genes Dev. 2006;20(1):77-86.
71.   Kundu N, Fulton AM. Selective cyclooxygenase (COX)-1 or COX-2
inhibitors control metastatic disease in a murine model of breast cancer. Cancer
Res. 2002;62(8):2343-6.
72.   Speirs CK, Jernigan KK, Kim SH, Cha YI, Lin F, Sepich DS, et al.
Prostaglandin Gbetagamma signaling stimulates gastrulation movements by
limiting cell adhesion through Snai1a stabilization. Development.
2010;137(8):1327-37.
73.   Wang D, Dubois RN. Prostaglandins and cancer. Gut. 2006;55(1):115-22.
74.   Yamashita S, Miyagi C, Fukada T, Kagara N, Che YS, Hirano T. Zinc
transporter LIVI controls epithelial-mesenchymal transition in zebrafish gastrula
organizer. Nature. 2004;429(6989):298-302.
75.   Lue HW, Yang X, Wang R, Qian W, Xu RZ, Lyles R, et al. LIV-1 promotes
prostate cancer epithelial-to-mesenchymal transition and metastasis through
HB-EGF shedding and EGFR-mediated ERK signaling. PLoS One.
2011;6(11):e27720.
76.   Yagi H, Tan W, Dillenburg-Pilla P, Armando S, Amornphimoltham P,
Simaan M, et al. A synthetic biology approach reveals a CXCR4-G13-Rho signaling
axis driving transendothelial migration of metastatic breast cancer cells. Sci
Signal. 2011;4(191):ra60.
77.   Webster DM, Teo CF, Sun Y, Wloga D, Gay S, Klonowski KD, et al. O-GlcNAc
modifications regulate cell survival and epiboly during zebrafish development.
BMC Dev Biol. 2009;9:28.
78.   Lynch TP, Ferrer CM, Jackson SR, Shahriari KS, Vosseller K, Reginato MJ.
Critical role of O-Linked beta-N-acetylglucosamine transferase in prostate cancer
invasion, angiogenesis, and metastasis. J Biol Chem. 2012;287(14):11070-81.
79.   Latinkic BV. Xenopus Cyr61 regulates gastrulation movements and
modulates Wnt signalling. Development. 2003;130(11):2429-41.
80.   Lin J, Huo R, Wang L, Zhou Z, Sun Y, Shen B, et al. A novel anti-Cyr61
antibody inhibits breast cancer growth and metastasis in vivo. Cancer Immunol
Immunother. 2012;61(5):677-87.
81.   Liu W, Su LT, Khadka DK, Mezzacappa C, Komiya Y, Sato A, et al. TRPM7
regulates gastrulation during vertebrate embryogenesis. Dev Biol.
2011;350(2):348-57.
82.   Middelbeek J, Kuipers AJ, Henneman L, Visser D, Eidhof I, van Horssen R,
et al. TRPM7 is required for breast tumor cell metastasis. Cancer Res.
2012;72(16):4250-61.

83.   Holloway BA, Gomez de la Torre Canny S, Ye Y, Slusarski DC, Freisinger
CM, Dosch R, et al. A novel role for MAPKAPK2 in morphogenesis during
zebrafish development. PLoS Genet. 2009;5(3):e1000413.
84.   Kumar B, Koul S, Petersen J, Khandrika L, Hwa JS, Meacham RB, et al. p38
mitogen-activated protein kinase-driven MAPKAPK2 regulates invasion of
bladder cancer by modulation of MMP-2 and MMP-9 activity. Cancer Res.
2010;70(2):832-41.
85.   Machingo QJ, Fritz A, Shur BD. A beta1,4-galactosyltransferase is required
for convergent extension movements in zebrafish. Dev Biol. 2006;297(2):471-82.
86.   Zhu X, Jiang J, Shen H, Wang H, Zong H, Li Z, et al. Elevated beta1,4-
galactosyltransferase I in highly metastatic human lung cancer cells.
Identification of E1AF as important transcription activator. J Biol Chem.
2005;280(13):12503-16.
87.   Hong SK, Tanegashima K, Dawid IB. XIer2 is required for convergent
extension movements during Xenopus development. Int J Dev Biol. 2011;55(10-
12):917-21.
88.   Neeb A, Wallbaum S, Novac N, Dukovic-Schulze S, Scholl I, Schreiber C, et
al. The immediate early gene Ier2 promotes tumor cell motility and metastasis,
and predicts poor survival of colorectal cancer patients. Oncogene.
2012;31(33):3796-806.
89.   Besser J, Leito JT, van der Meer DL, Bagowski CP. Tip-1 induces filopodia
growth and is important for gastrulation movements during zebrafish
development. Dev Growth Differ. 2007;49(3):205-14.
90.   Han M, Wang H, Zhang HT, Han Z. The PDZ protein TIP-1 facilitates cell
migration and pulmonary metastasis of human invasive breast cancer cells in
athymic mice. Biochem Biophys Res Commun. 2012;422(1):139-45.
91.   Faure S, Cau J, de Santa Barbara P, Bigou S, Ge Q, Delsert C, et al. Xenopus
p21-activated kinase 5 regulates blastomeres' adhesive properties during
convergent extension movements. Dev Biol. 2005;277(2):472-92.
92.   Gong W, An Z, Wang Y, Pan X, Fang W, Jiang B, et al. P21-activated kinase 5
is overexpressed during colorectal cancer progression and regulates colorectal
carcinoma cell adhesion and migration. Int J Cancer. 2009;125(3):548-55.
93.   Iioka H, Ueno N, Kinoshita N. Essential role of MARCKS in cortical actin
dynamics during gastrulation movements. J Cell Biol. 2004;164(2):169-74.
94.   Rombouts K, Carloni V, Mello T, Omenetti S, Galastri S, Madiai S, et al.
Myristoylated Alanine-Rich protein Kinase C Substrate (MARCKS) expression
modulates the metastatic phenotype in human and murine colon carcinoma in
vitro and in vivo. Cancer Lett. 2013;333(2):244-52.
719
720