Submission type: **Research article**

**Reciprocal recombination reflects sexual reproduction in symbiotic arbuscular**

**mycorrhizal fungi**

Ivan D. Mateus[1]*, Ben Auxier[2], Mam M. S. Ndiaye[1], Joaquim Cruz[1], Soon-Jae Lee[1] & Ian R.

Sanders[1]

[1]*Department of Ecology and Evolution, University of Lausanne, Biophore building, 1015*

*Lausanne, Switzerland*

[2]*Laboratory of Genetics, Wageningen University, Wageningen, the Netherlands.*

* Author for correspondence: ivandario.mateusgonzalez@unil.ch

**Keywords:** Arbuscular mycorrhizal fungi, AMF, haplotypes, recombination, sexual

reproduction

## **Abstract**

Arbuscular mycorrhizal fungi (AMF) are part of the most widespread fungal-plant symbiosis.

They colonize at least 80% of plant species, promote plant growth and plant diversity. These

fungi are multinucleated and contain either one or two haploid nuclear genotypes

(monokaryon and dikaryon) identified by the alleles at a putative mating-type locus. This

taxon has been considered as an ancient asexual scandal because of the lack of observable

sexual structures. Despite identification of a putative mating-type locus and functional

activation of genes related to mating when two isolates co-exist, it remains unknown if AMF

take part in a mainly sexual life cycle.

We used publicly available genome sequences to test if dikaryon genomes display

signatures of sexual reproduction in the form of reciprocal recombination patterns, or if they

display signatures of parasexual reproduction involving gene conversion.

We used short-read and long-read sequence data to identify nucleus genotype-specific

haplotypes within dikaryons and then compared them to orthologous gene sequences from

related monokaryon isolates displaying the same putative MAT-types. We observed that

these genotype-specific haplotypes display reciprocal recombination and not gene

conversion.

These results are consistent with a sexual origin of the dikaryon rather than a parasexual

origin and provides an important step to understand the life cycle of these globally important

symbiotic fungi.

## **Introduction**

Arbuscular mycorrhizal fungi are plant symbionts, forming symbioses with most plant species, promoting plant growth (Harrison, 1997), plant community diversity (van der Heijden *et al.*, 1998; Antunes *et al.*, 2011) and how plants cope with biotic (Thygesen *et al.*, 2004) or abiotic stresses (Augé, 2001). As a consequence, they are widely used in agriculture (Smith & Read, 2008). In a single agricultural field, the presence of at least 17 different genotypes of *Rhizophagus irregularis* displaying spatial genetic structure have been detected (Koch *et al.*, 2004; Croll *et al.*, 2008). Furthermore, these *R. irregularis* isolates display different levels of within isolate genetic diversity (Wyss *et al.*, 2016), which has been reported to produce differential effects on plant growth (Angelard *et al.*, 2010). Understanding how genetic variability is generated in AMF, is important because it could be harnessed to generate genetic variants that could be beneficial for their plant hosts (Sanders, 2010).

AMF are part of the Glomeromycotina subphylum (Spatafora *et al.*, 2016), which fossil records date to at least ~400 Million years ago (Remy *et al.*, 1994). They are coenocytic (without septa separating otherwise adjacent compartments), their hyphae harbor hundreds of nuclei within the same cytoplasm (Marleau *et al.*, 2011) and no single-nucleus state has been recorded. The nuclei of these fungi have been reported as haploid (Lin *et al.*, 2014; Ropars *et al.*, 2016; Kobayashi *et al.*, 2018; Morin *et al.*, 2019) . This group of fungi has been previously considered as an ancient asexual scandal (Judson & Normark, 1996), due to low morphological diversification and the absence of observable sexual structures. However, evidence suggests that sexual reproduction could be possible in AMF since these fungi contain a complete meiosis machinery (Halary *et al.*, 2011). Furthermore, a putative mating-type determining locus (MAT) has been proposed (Ropars *et al.*, 2016), population genetic data suggests the existence of recombination in AMF populations (Croll & Sanders, 2009) and activation of genes related to meiosis has been detected when different isolates of the same species co-exist in plant roots (Mateus *et al.*, 2020, 2021).

Several *R. irregularis* isolates issued from the same geographic location have been reported as monokaryon (A1,B12,C2) and dikaryon (A4, A5, C3 and G1) (Ropars *et al.*, 2016; Chen *et al.*, 2018a; Masclaux *et al.*, 2018; Kokkoris *et al.*, 2021), evidencing that monokaryon and dikaryon isolates co-exist in the same location. Single-nucleus from dikaryons (isolates A4, A5 and SL1) cluster into two genetically different nucleotypes (where nucleotype refers to the genotype of a nucleus), that are associated with the identity of a putative MAT-locus (Chen *et al.*, 2018a). This demonstrates that the presence of two copies of the putative MAT-locus is a reliable marker of the dikaryon state.

Like most fungi, AMF can undergo anastomosis, the fusion of hyphae. Through these connections, bi-directional flow of cytoplasm has been observed between genetically different AMF individuals (Giovannetti *et al.*, 1999). Via anastomosis, the transfer of genetic material between vegetative compatible isolates (parasexuality) has been suggested as a mechanism of maintenance of genetic diversity in the absence of sexual recombination (Bever & Morton, 1999). In fungi, hyphal fusion between different individuals leads to cell death, however, non-self vegetative compatibility has also been observed in AMF when different isolates form perfect hyphal fusions (Croll *et al.*, 2009). In the model species *Aspergillus nidulans* parasexuality involves fusion of two haploid nuclei, mitotic recombination and haploidization of the diploid nuclei by i.e. chromosome loss (Pontecorvo, 1956). In consequence, a genomic signature of parasexuality is gene-conversion where there is a loss of heterozygosity. However, sexual reproduction can also display signatures of gene-conversion but also involves meiotic recombination and the segregation on reciprocal products.

The existence and relevance of sexuality and/or parasexuality for the evolution of AMF remains unknown (Yildirir *et al.*, 2020). It is still unknown whether the transition between a dikaryon and monokaryon life stages involves a sexual event involving meiotic recombination, or a parasexual event.

In the absence of stable transformation methods in AMF (Helber & Requena, 2008),

haplotype analyses of genomic data could be an important resource to identify genomic

signatures of mitotic or meiotic recombination in dikaryon isolates.  The haplotypes from

dikaryon isolates can then be compared to orthologous sequences of related monokaryon

isolates (that share the same MAT-type) and could allow to identify genomic signatures of a

sexual event, involving meiotic recombination, or a parasexual event which results in the

loss of heterozygosity or gene conversion.

AMF genomes display large gene duplication events (Morin *et al.*, 2019), which make it

difficult to call haplotypes. The identification of haplotypes in dikaryon isolates could be

made by: 1) analyzing short-read sequences to identify genome-wide copy number variation.

These analyses consist in obtaining the read depth, or coverage, after mapping the reads to

a genome assembly and identifying changes in coverage across the genome (Yoon *et al.*,

2009). In AMF, a drop in coverage analysis was used to originally identify the putative MAT-

locus (Ropars *et al.*, 2016) highlighting the potential to identify nucleotype-specific

haplotypes in dikaryons with this technique. 2) Analyzing long-read sequencing to identify

haplotypes and, consequently, nucleotype-specific haplotypes in fungal dikaryon isolates (Li

*et al.*, 2019).

Here, we demonstrated that AMF dikaryons display reciprocal recombination by analyzing

nucleotype-specific haplotypes in dikaryon (A5) and monokaryon isolates (A1-C2). In this

study we used publicly available whole genome, single-nucleus short-read sequence data

and long-read bulk-isolate sequence data to identify nucleotype-specific haplotypes in

dikaryon isolates. We identified regions displaying drops in coverage in short-read whole

genome sequence assemblies. In these regions we detected the presence of genes that

have two copies in the dikaryon isolates and one copy in monokaryons. We then confirmed

independently, with short-read genome sequence data from single-nucleus, that in dikaryon

isolates, different nuclei have different alleles and that they are not always associated to the

MAT identity, evidencing a reciprocal recombination genomic signature. Finally, we validated

the analysis, by evaluating long-read genome sequence assemblies and confirmed that A5

dikaryon display genomic signatures of reciprocal recombination, which suggest sexual

reproduction in AMF.

**Materials and Methods**

**Source data**

We used public-available sequence reads, genome assemblies and annotations of isolates

A1, A4, A5, C2 of *R. irregularis* for this study, including data from bulk-isolate and single-

nucleus sequencing (Supplementary Table 1). We used short-read whole genome

assemblies from isolates A1, A4, A5 and C2 (Ropars *et al.*, 2016). We used single-nucleus

genome assemblies from isolates A1, A5 and C2 (Chen *et al.*, 2018a). We also analyzed

long-read genome assemblies of isolates A1, A5 and C2 (Chaturvedi). We downloaded the

sequence reads from the sequence read archive (SRA) using the SRAtoolkit software with

the prefetch and fastq-dump tools (Leinonen *et al.*, 2011).

**Coverage analysis**

We first trimmed the sequence reads using Trim Galore! (Krueger, 2015) with the default

parameters. We then used BWA (Li, 2013) to index the reference genome assemblies and

BWA mem -M (Li, 2013) to map the reads to the reference whole-genome assemblies. We

mapped the reads coming from a given isolate to the reference genome assembly of the

same isolate (i.e. reads A1 mapped to reference A1). We then kept the reads that display a

mapping quality of at least 30. We used the genomecov tool from bedtools (Quinlan & Hall,

2010) to calculate the coverage for each position. We then created a ready-to-use algorithm

that detects genome-wide drop in coverage analysis in whole-genome data (Supplementary

File 1). The algorithm divides the data in portions of 50kb. Then, with a sliding window

approach consisting of windows of 400bp and steps of 100bp, the algorithm searches for

drops in coverage of 0.3-0.6 times lower than the median coverage of the entire genome

and that with a minimum length of 1000bp (Please refer to Supplementary File 1 for the

algorithm specifications implemented in the R programming language). We then further

filtered these drop in coverage regions by keeping only the regions that display an average

of 1.25 coverage difference between the neighboring regions and the drop in coverage

region.

**Gene detection in drops**

We identified all the genes located within genomic regions that presented a drop of

sequencing coverage. We used the 'intersect' command from the BEDTools suite with the

existing gene annotations corresponding to each *R. irregularis* isolate (GTF format) and their

query regions with drops in coverage (BED format) to identify overlapping genes (Quinlan &

Hall, 2010). Genes in scaffolds smaller that 1kb were not considered for further analyses.

*de novo* **single-nucleus assemblies**

We trimmed the raw reads by using TrimGalore-0.6.0 (Krueger, 2015) with default

parameters. After trimming, we performed single-nucleus *de novo* assemblies with SPAdes

v3.14 (Bankevich *et al.*, 2012) with the following parameters: -k 21,33,55,77 --sc --careful --

cov-cutoff auto. The resulted single-nucleus genome assemblies were used for further

analysis. The length, number of contigs and N50 value of the *de novo* assemblies was

evaluated with quast- 5.1.0rc1 with default parameters (Mikheenko *et al.*, 2018).

**Identification of genes in genome assemblies**

To identify the position of gene sequences on the different genome assemblies, we first

extracted a query sequence. We then used the console NCBI+ blast suite  (Camacho *et al.*,

2009) to blast the query against the desired target. In the case of the putative MAT-locus, we

used the homeodomain genes HD2 and HD1-like as query (HD2:KT946661.1, HD1-like:

KU597387 from isolate A1). For further downstream analyses, we extracted the sequences

from the genome assemblies by using the blastdbcmd command from the NCBI+ suite. We

used a reciprocal blast approach to identify the gene sequences corresponding between the

whole genome sequence data and the single-nucleus data. We considered the bests hits by

evaluating the % identity, mismatches, e-value and bitscore.

**Orthology inference**

We used Orthofinder 2.3.11 (Emms & Kelly, 2019) to identify orthologs of genes found inside

the drop in coverage regions within the same isolate. We also identified orthologs in the

long-read assemblies and identified single copy orthologs in isolates A1, C2, the primary

assembly of A5 and the haplotig assembly of A5. We used the orthogroups output from

Orthofinder for the different analyses.

**Synteny plots**

We compared genomic regions by performing synteny plots computed with EasyFig2.2.3

(Sullivan *et al.*, 2011). We provide full Genbank files to compare genomic regions to each

other. The software executes a blast comparison between the regions to determine their

homology.

**Genetic distance between nucleus specific haplotypes**

Coding sequences for the 12 confirmed nucleotype-specific genes were extracted using the

Blast+ command line blastdbcmd tool (Camacho *et al.*, 2009). The sequences were then

aligned with MAFFT (Katoh *et al.*, 2017) using the --auto option. Then, the ape package

(Paradis *et al.*, 2004) of R was used to calculate the pairwise distance between the 4 alleles

(2 from A5, and 1 each from A1 and C2).

**Recombination detection**

We compared the sequences from drop in coverage regions from both nuclei specific

haplotypes of isolate A5 and isolates A1 and C2 to detect if isolate A5 display recombination

events between the two putative parental isolates. After identification of the syntenic region

among the different isolates, we aligned the sequences with MAFFT (Katoh *et al.*, 2017) and

evaluated whether the sequence of one of the nucleotype of isolate A5 was similar to A1 and

the other similar to C2.

**Phylogenetic analyses**

We used MEGA-X (Kumar *et al.*, 2018) for the different phylogenetic reconstructions shown

in the study. We first aligned the data with ClustalW. We then find the best DNA models

describing the relation between the sequences. Finally, we used a maximum likelihood

phylogeny reconstruction with 100 bootstraps to infer the phylogenetic relation among the

samples. In several cases, we were not able to perform maximum likelihood phylogenies

because of the low number of samples to compare, so UPGMA trees were done instead.

Phylogenetic reconstructions of the different orthologous groups on Figure 4 where

produced by the Orthofinder software.

**Results**

**Drop in coverage analysis reveals potential nucleotype-specific haplotypes**

Previously, a drop in coverage analysis of selected regions was used for the identification of

a putative MAT-locus in *R. irregularis* (Ropars *et al.*, 2016). To perform a similar analysis

across the entire genome, we developed a script (Supplementary File 1) that allows us to

identify genome-wide drop in coverage events (for the accessions of raw data and genome

assemblies used in this study see Supplementary Table 1).

We identified drops in coverage in 4 different isolates of *R. irregularis* which are reported to

be dikaryons (A4 and A5) and monokaryons (A1 and C2) (Figure 1, Supplementary Table 2).

The number of coverage drops and genes inside the drop in coverage regions were different

between dikaryons (A4-A5) and monokaryons (A1-C2) (Supplementary Table 3). These

results indicated that dikaryotic isolates displayed more heterozygous regions than the

monokaryons, suggesting that the genes present within the regions showing a drop in

coverage are potential candidates for highly divergent, nucleotype-specific alleles in

dikaryons. Confirming the reliability of our approach, we detected the expected drop in

coverage in the putative MAT-locus region in isolates A4 and A5 but not in isolates A1 and

C2 (Figure 1d).

One cause for a drop in coverage could be copy number variation between the nucleotypes

in a dikaryon. To test for this, we inferred orthologous gene families among the different

isolates to identify if the genes present in the drop in coverage regions displayed more than

one copy in their own genome. We used the gene annotation available for each isolate and

inferred the orthology of all the genes present in each genome. Many orthologous groups

had more than one copy within each isolate (A4: 20%, A5: 18%, A1: 17% and C2: 19%;

Supplementary figure 1, Supplementary Table 4) consistent with the high reported incidence

of paralogs in these fungi (Morin *et al.*, 2019). We further identified orthologous gene families

of genes detected in drop in coverage in isolates A4 and A5 independently. Under the

assumption of a monokaryon-dikaryon genome organization in *R. irregularis*, to avoid the

confounding effect of duplications and reduce the complexity of the dataset, we retained only

the orthologous groups that are present in the drop in coverage regions and that display two

copies in the dikaryon isolates (A4, A5) and a single copy in the monokaryons (A1, C2)

(Figure 2a, Supplementary Table 5). In the regions where a drop in coverage was detected,

we identified 32 orthologous groups that are present with two copies in isolate A4 and only a

single copy in isolates A1 and C2. We also identified 27 orthologous groups in isolate A5

that display two copies. Only two orthologous groups were common between the two

isolates: namely, HD2 and HD1-like which are part of the putative MAT-locus in *R. irregularis*

(Figure 2b).  As reported in Ropars *et al.,* we observed that the two copies of the putative

MAT-locus in the dikaryons were located in different contigs. One copy of HD2 and HD1-like

genes were present in a long contig of the genome assembly, while the second copy was

present in a much shorter contig (Figure 2c). We observed the same pattern for the other

orthologous groups, where the second copy was always present in a second shorter contig

(for several examples see Figure 2d).

To further sort paralogous genes from true orthologs, we performed a synteny analysis to

compare the genomic location of the presumed orthologous genes among isolates. We

identified that 12 out of 32 predicted orthologs in A4 and 16 out of 27 predicted orthologs in

A5, were located in the same genomic location on the different isolates, suggesting that they

should be considered as orthologs (Supplementary Table 5, for examples of inferred

orthologs and paralogs see Supplementary Figure 2).

Hence, in the whole genome assemblies of dikaryon isolates, two divergent alleles were

assembled into different scaffolds; one longer scaffold containing neighboring regions and a

shorter scaffold without the neighboring regions. Given that the nuclei are haploid in the

dikaryon isolates, two possibilities are consistent with this previous fact: The two copies

could be present within the same or in different nuclei.

**Drop in coverage signatures represent nucleotype-specific haplotypes**

To confirm that genes found inside drop in coverage regions are nucleotype-specific, we

used sequencing reads of individual nuclei of dikaryon isolates A4 and A5 (Chen *et al.*,

2018a) to produce *de novo* single-nucleus assemblies. The *de novo* assemblies were very

fragmented and incomplete (Supplmentary figure 3ad, Supplementary table 6) and their

utilization was highly limited. This limitation resulted in the inability to identify some genes

and some complete gene sequences. However, a reciprocal blast approach between the

whole genome assembly and the single-nucleus assemblies allowed us to detect sequences

in the single-nucleus assemblies corresponding to the genes detected in the whole genome

assemblies.

We tested in the dikaryons if the genes identified in the drop in coverage regions were

present in the form of different alleles in different single nuclei by using a reciprocal blast

approach. We confirmed 9 orthologous genes to be nucleotype-specific in isolate A4 and 12

orthologous genes in isolate A5 (Supplementary Table 7). We found that the population of

nuclei clustered in two groups that corresponded to the identity of the putative MAT-locus

contained in each nucleus (Figure 3). This result confirms that nucleotype-specific alleles in

dikaryon isolates can be identified based on genes found in drop in coverage regions and

that are represented by a duplication within the genome assembly.

**Nucleotype-specific alleles from A5 share a more recent evolutionary origin with**

**monokaryon isolates A1 and C2 than among them**

The origin of dikaryon isolates could be investigated through comparisons of monokaryon

isolates that display the same alleles of the putative MAT-locus as those found in the

dikaryons (Isolates A5:MAT-3/MAT-6; A1:MAT-3; C2: MAT-6). A phylogenetic reconstruction

of the putative MAT-locus suggests that MAT-3 from isolates A1 and A5 are more closely

related than MAT-6 from isolates C2 and A5 (Ropars *et al.*, 2016). Furthermore, genome-

wide reduced genome representation phylogenetic reconstructions of several *R. irregularis*

isolates indicated that isolate A5 is more closely related to isolate A1 than to isolate C2

(Wyss *et al.*, 2016; Savary *et al.*, 2018).

To confirm the previous findings, for each previously defined nucleotype-specific gene, we

compared the phylogenetic relationship of the two nucleotype-specific alleles in isolate A5

isolate and in isolates A1, C2 and A4. We observed that for several nucleotype-specific

genes, genes from isolate A1 clustered with one of the alleles of isolate A5, but it was not

always the case in isolate C2 (Figure 4a).

We then analyzed the genetic distance of each of the 12 nucleotype-specific genes

independently between the two alleles of isolate A5 and the homologous allele in isolates A1

and C2. The mean nucleotide distance between the two A5 alleles was 0.147. The mean

distance between A1 and C2 was 0.132. In contrast, the mean of the minimum distance

between an allele of isolate A5 and isolate A1 was 0.013 and between A5 and C2 was 0.043

(Figure 4b). We did not observe any case where the two A5 alleles clustered together,

instead we observed that for all 12 nucleotype-specific gene the two A5 alleles were more

similar to the allele from isolate A1 or C2 (Figure 4c). The mean distances calculated

between alleles in this study are much higher than average distances calculated on the

whole genome between different isolates (Chen *et al.*, 2018b), reflecting our selection

criteria for nucleotype-specific regions. As each of the A5 alleles was closer to A1 or C2,

instead of the two A5 alleles being most similar, this indicates that the alleles of A5 share a

more recent evolutionary origin with these monokaryons than the two alleles within A5.

**Identification of reciprocal recombination between nucleotype-specific haplotypes in**

**isolate A5 using single-nucleus data**

Knowledge about nucleotype-specific haplotypes of dikaryon isolate A5 and their orthologs

in isolates A1 and C2 allowed us to test whether parasexual or sexual genomic signatures

could be identified in dikaryon isolate A5 (Figure 5a). We scanned the different nucleotype-

specific-haplotypes for the detection of recombination events within the two haplotypes of

isolate A5. Comparison of haplotypes issued of the short-read whole genome assemblies

from isolates A1, C2 and the two haplotypes of A5 showed that each nucleotype-specific

sequence from isolate A5 was highly similar to either isolate A1 or C2, but we did not identify

any recombination events within the haplotypes (Supplementary Figure 3).

To further assess the potential for clonal relationships between the two nucleotypes within

A5 and isolates A1 and C2, we compared the nucleotype-specific haplotypes on the single-

nucleus assemblies from isolate A5 and their orthologs on the single-nucleus assemblies of

isolates A1 and C2 (Supplementary Table 8). The difference with the previous analysis is

that with the single-nucleus data, we are able to identify the identity of the putative MAT-type

of the A5 haplotypes. We found that for several nucleotype-specific genes (i.e. OG2995,

OG3981 and OG4715), the sequences from isolates A5 (MAT-3 type) and A1 (MAT-3 type)

clustered together (Supplementary Figure 4, Figure 5b). However, we found that for other

nucleotype-specific genes (OG4925, OG4492 and OG4493), the sequences from isolate A5

(MAT-3 type) clustered with the sequences from isolate C2 which has a MAT-6 type

(Supplementary figure 5, Figure 5c). The alignments on these nucleotype-specific genes

show that A5 nuclei with MAT-3 have similar, but not identical alleles as C2 nuclei (MAT-6

type). In the same way, A5 nuclei with MAT-6 type harbor alleles similar to those of A1 nuclei

(MAT-3 type). These results demonstrate that the A5 isolate harbors nucleotypes with

regions highly similar to isolates A1 and C2, but that the A1-like alleles are not always found

in the same nucleus. The presence of reciprocal recombinant nucleotypes in isolate A5,

involving isolates sharing the same MAT-type, strongly suggest that isolate A5 results from a

sexual and not parasexual event between isolates similar to A1 and C2.

**Confirmation of reciprocal recombination events in isolate A5 using long-read**
**genome assemblies**

We compared single-copy orthologous genes among the A1, C2 assemblies and the phased

assembly of isolate A5, which is divided in a primary assembly and an haplotig assembly.

The long-read assemblies were more complete and contain longer scaffolds than the short

reads assemblies (Supplementary Figure 3bcef). The A5 phased assembly consisted of a

primary assembly of 392 scaffolds and an haplotig assembly on 292 of the primary scaffolds.

The primary assembly was 115Mb long and the haplotig assembly was 19.2Mb (Chaturvedi

*et al.*, 2021).

We identified 1250 single-copy orthologs among isolates A1, C2, the A5 primary assembly

and the A5 haplotig assembly (Supplementary Table 9).  We then focused only in the

orthologs were one of the genes of A5 clustered with either A1 or C2. The majority of the

genes sequences were very similar among all the isolates, however we identified 44

orthologous groups where one of the haplotype genes of A5 cluster with either isolate A1 or

C2 (23 orthogroups A1-A5haplotig and 21 orthogroups A1-A5primary). We observed that for

several contiguous regions, an orthogroup displayed the A5primary gene clustering with A1

and in another orthogroup, contained in the same contiguous region, the A5 primary gene

clustered with the C2 gene (Figure 6), confirming that reciprocal recombination signatures

are found within continuous haplotypes (Supplementary Table 9).

**Discussion**

In this study, we used short and long read published sequencing data to demonstrate that

signatures of reciprocal recombination can be detected in genomes of dikaryon AMF isolates

that display different putative MAT-types. The signatures of reciprocal recombination are

compatible with a sexual reproduction life cycle, rather than a parasexual reproduction

mechanism. These results suggest that after all, AMF are not that scandalous.

In the absence of experimental evidence that isolates with different putative MAT-types are

involved in sexual reproduction, genomic signatures of recombination can help us to

understand whether sexual or parasexual reproduction are involved in the AMF life cycle. On

one hand, parasexual reproduction genomic signatures involves gene-conversion, which

result is the loss of heterozygosity (Forche *et al.*, 2008). On the other hand, reciprocal

recombination associated with a putative sex-determining region (MAT-type) is a genomic

signature of sexual reproduction, although gene-conversion could also happen in sexual

reproduction (Sun *et al.*, 2012). It could be plausible that parasexual recombination could

lead to a reciprocal recombination pattern by fusion of two monokaryons, mitotic reciprocal

recombination, followed by reciprocal chromosome loss. However, the presence of a

complete meiotic machinery, the presence of compatible putative MAT-types, and the

upregulation of genes involved in a putative mating response when two isolates co-exist

(Mateus *et al.*, 2020, 2021), suggest that a plausible hypothesis explaining the results

observed in this study is a reciprocal recombination pattern detected in dikaryons, the result

of sexual reproduction.

We identified few regions that display recombination patterns within continuous haplotypes

in the dikaryon isolate A5. The principal reason, is that we did not evaluate all the different

genes present, but only the genes that display only 1 copy in an haploid phased contig. We

only compared single-copy orthologs among the different assemblies, to avoid the

confounding effect of paralogy on our results. In consequence, the examples of reciprocal

recombination shown in this study, are only a small subset of the total plausible

recombination spots present in their genomes. With the information of nucleotype-specific

haplotypes identified with the drop in coverage analysis, we found that nucleotypes of isolate

A5 were as little 1% diverged from isolate A1 and 4% diverged from isolate C2, suggesting

that isolates sharing the same MAT-type as the monokaryon isolates A1 and C2 are closely

related ancestors from which the dikaryon A5 arose.  Interestingly, with the long-read data,

we identified several loci where the A1 and C2 genes were more closely related among them

than to the alleles of isolate A5. This suggest, that there could be other evolutionary forces

shaping AMF genome evolution. Transposable elements (TE) have been reported to be

responsible of genome duplications, inversions, insertions and deletions (Daboussi & Capy,

2003). In nature, AMF co-exist with different conspecifics and they co-exists aswell with

different host-plants. We could speculate, that gene trajectories within AMF populations

could be also driven by TE-mediated horizontal gene transfer events. However, an important

sampling of AMF genomes is necessary to better understand the genome architecture

features and life-history of the different gene families (Badet & Croll, 2020).

An intriguing question, is how only two coexisting nucleotypes and a well-orchestrated

mechanism such as meiosis could be achieved without the presence of a single-cell stage in

the AMF life cycle.  A genetic bottleneck, where only 1 or 2 nuclei co-occur has never been

detected in AMF. Despite, the last feature, two studies that evaluated nuclear imbalance in

dikaryon isolates show that nuclei ratio among single-spore lines are conserved despite

multiple culturing for years (Robbins *et al.*, 2021) or are stable when exposed to different

host plants (Kokkoris *et al.*, 2021). These studies suggest that a nuclei regulation

mechanism should exist and could be responsible of the stability of the nucleotypes in the

dikaryons.

In this study, we compared haplotypes within the dikaryon A5 to monokaryon isolates (A1

and C2) that contain the same putative MAT-types as the dikaryon A5 to identify the

recombination patterns.

We were able to analyze these signatures on the A5 dikaryon, but not on the dikaryon A4,

because there are no available genomic assemblies of monokaryon isolates displaying the

same putative MAT-type as in Isolate A4.  We hypothesized that MAT-type compatibility

exists between the two different putative MAT-types identified in the dikaryon isolates.

However, to date there is no direct experimental evidence that different putative MAT-types

could be compatible. An indirect evidence of compatibility between different putative MAT-

types was found, when two different isolates harboring different putative MAT-types elicited

a putative fungal mating response (Mateus *et al.*, 2020, 2021). However, Mateus et al., did

not test if  the mating of two co-existing isolates did took place (Mateus *et al.*, 2020). In

consequence, in order to experimentally identify mating and MAT-type compatibility,

crossing experiments between strains harboring different putative MAT-types, including

transcriptome and recombinant progeny analyses should be performed.

Our drop in coverage approach allowed us to identify divergent nucleotype-specific genes in

dikaryon isolates situated in different contigs of the short-read whole genome assemblies.

This approach differs from previous approaches of global intra-isolate divergence

assessment that measured the number of SNPs (Chen *et al.*, 2020) or poly-allelic sites

(Wyss *et al.*, 2016). Although the comparison of both types of measurements gave similar

information (intra-isolate divergence), their comparison should be carefully addressed as

their methodology and the types of sequences compared are different. While SNPs are best

identified in low divergence regions, where reads can be confidently mapped to the same

contig, the sequences with large genetic differences, detected with the drop in coverage

approach, are highly divergent to the point that they are assembled in different contigs in the

same genome assembly. Consequently, genetic divergence should be higher in the regions

with large genetic divergence than when the two alleles are collapsed in the genome

assembly, resulting in sequences displaying several SNPs. Contrary, to the drop in coverage

analysis using short-read data, the phased haplotypes from long-read sequencing, display

genes that differ in several SNP's between the two dikaryon haplotypes.

Inter-nucleus recombination has been previously reported (Chen *et al.*, 2018a), although the

robustness of the analysis has been questioned. The application of strict filtering parameters

such as removal of heterozygous sites in haploid nuclei, duplicated regions of the genome,

and low-coverage depths base calls results in an extreme loss of the signal of recombination

(Auxier & Bazzicalupo, 2019). Although some of these limitations, as coverage depth and

filtering-out heterozygous sites were addressed (Chen *et al.*, 2020), other limitations such as

replicability (recombination events shown in several nuclei sharing the same putative MAT-

type) and issues related to the whole-genome amplification step as the formation of chimeric

sequences (Yilmaz & Singh, 2012), allelic drop-out (Lauri *et al.*, 2013) and SNP miscalling

(Ning *et al.*, 2014) are inherent limitations of the analysis of any single nucleus amplification

data.  Furthermore, the reported evidence of large inter-nuclei recombination (Chen *et al.*,

2018a, 2020) does not fit the observations about a monokaryon – dikaryons organization. In

Chen *et al.,* 2018a, the selected examples of genotypes presented in figure 3 show 4 nuclei

per isolate, which represent four different nuclei for SL1, A4 and three different nuclei on A5.

The same pattern can be observed in Supplementary Figure 5 from Chen *et al.,* 2020, the

examples of genotypes show 7 out of 8 different nuclei in isolate A5, at least 7 different

nuclei in isolate A4 and at least 5 different nuclei in SL1. The presence of repeated inter-

nuclear recombination in dikaryons without prior crossing with other isolates, as observed in

the single-nucleus genotypes shown in Chen et al. (2018a, 2020) would result in

heterokaryons with more than two types of nuclei. But this does not appear to be the case

for *R. irregularis* (Ropars *et al.*, 2016; Chen *et al.*, 2018a; Masclaux *et al.*, 2019; Auxier &

Bazzicalupo, 2019).

Maintaining monokaryon and dikaryon isolates within the same natural population suggests

that both forms are stable over time. Rather than a promiscuous mixing between isolates via

anastomosis, a mechanism of recognition that involves the putative MAT-locus seems to

regulate which isolates can form a dikaryon (Corradi & Brachmann, 2017). The fact that

nucleotype-specific haplotypes from isolate A5 are more closely related to isolates A1 and

C2, and that A5 nucleotypes display recombination, suggests that isolates sharing the same

putative MAT-type as A1 and C2 could be the origin of a recombining dikaryon isolate.

However, we cannot discard that these findings could apply to another step of the AMF life

cycle. It could also be possible that a stable A5 isolate could segregate producing

recombined monokaryons that share the same putative MAT-type as isolates A1 and C2,

that can disperse and then fuse again to form stable dikaryons and complete a life cycle

which involves recombination. It then becomes crucial to experimentally confirm if

monokaryon isolates having different putative MAT-types could generate a dikaryon-like

form and if a dikaryon isolate could segregate into recombining monokaryon isolates.

Understanding the life cycle of AMF could have an enormous impact in the generation of

AMF genetic variability. The generation of diverse AMF monokaryons or dikaryons could be

used to generate variants that enhance plant growth and have an enormous potential in

agriculture (Sanders, 2010).

**<u>Acknowledgments</u>**

We would like to thank Daniel Croll for his feedback on a preliminary version of this

manuscript. We specially thank Bart Nieuwenhuis and his group for the constructive

feedback made on the Biorvix version of this manuscript. This study was funded by the

Swiss National Science Foundation (Grant number: 31003A_162549 to IRS).

## **Author contributions**

IDM designed and supervised all the analyses. MMSN developed the drop in coverage

detection algorithm. JC identified the genes present in the regions displaying drop in

coverage. BA developed the genetic relatedness analysis on the nucleotype-specific

haplotypes. SL provided valuable comments during all the process. IDM, performed all the

bioinformatic analyses, identified the nucleotype-specific genes, performed the phylogenetic

analysis, gene-retrieval from genome assemblies and the recombination detection. IDM and

IRS wrote the manuscript. All the authors provided valuable contributions and modifications

of the manuscript.

## **Data availability**

522 All data analyzed in this manuscript its available in the NCBI repository and the respective

523 accession identifiers could be found in Supplementary table 1. The custom code developed

524 for the identification of drop in coverage regions could be found in Supplementary file 1.

## **Conflict of interest**

526 The authors declare that they have no conflict of interest.

### References

**Angelard C, Colard A, Niculita-Hirzel H, Croll D, Sanders IR**. **2010**. Segregation in a mycorrhizal fungus alters rice growth and symbiosis-specific gene transcription. *Current Biology* **20**: 1216–21.

**Antunes PM, Koch AM, Morton JB, Rillig MC, Klironomos JN**. **2011**. Evidence for functional divergence in arbuscular mycorrhizal fungi from contrasting climatic origins. *The New phytologist* **189**: 507–14.

**Augé RM**. **2001**. Water relations, drought and vesicular-arbuscular mycorrhizal symbiosis. *Mycorrhiza* **11**: 3–42.

**Auxier B, Bazzicalupo A**. **2019**. Comment on 'Single nucleus sequencing reveals evidence of inter-nucleus recombination in arbuscular mycorrhizal fungi'. *eLife* **8**: 1–9.

**Badet T, Croll D**. **2020**. The rise and fall of genes: origins and functions of plant pathogen pangenomes. *Current Opinion in Plant Biology* **56**: 65–73.

**Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, *et al.* 2012**. SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology* **19**: 455–477.

**Bever JD, Morton J**. **1999**. Heritable variation and mechanism of inheritance of spore shape within a population of Scutellosporea pellucida, An arbuscular mycorrhi. *American journal of botany* **86**: 1209–1216.

**Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL**. **2009**. BLAST+: Architecture and applications. *BMC Bioinformatics* **10**: 1–9.

**Chaturvedi A, Cruz Corella J, Robbins C, Loha A, Menin L, Gasilova N, Masclaux FG,**

**Lee S, Sanders IR**. **2021**. The methylome of the model arbuscular mycorrhizal fungus,

Rhizophagus irregularis, shares characteristics with early diverging fungi and Dikarya.

*Communications Biology* **4**: 901.

**Chen ECH, Mathieu S, Hoffrichter A, Ropars J, Dreissig S, Fuchs J, Brachmann A,**

**Corradi N**. **2020**. More Filtering on SNP Calling Does Not Remove Evidence of Inter-

Nucleus Recombination in Dikaryotic Arbuscular Mycorrhizal Fungi. *Frontiers in Plant*

*Science* **11**: 1–9.

**Chen EC, Mathieu S, Hoffrichter A, Sedzielewska-Toro K, Peart M, Pelin A, Ndikumana**

**S, Ropars J, Dreissig S, Fuchs J,** *et al.* **2018a**. Single nucleus sequencing reveals

evidence of inter-nucleus recombination in arbuscular mycorrhizal fungi. *eLife* **7**: 1–17.

**Chen ECH, Morin E, Beaudet D, Noel J, Yildirir G, Ndikumana S, Charron P, St-Onge C,**

**Giorgi J, Krüger M,** *et al.* **2018b**. High intraspecific genome diversity in the model

arbuscular mycorrhizal symbiont Rhizophagus irregularis. *New Phytologist* **220**: 1161–1171.

**Corradi N, Brachmann A**. **2017**. Fungal Mating in the Most Widespread Plant Symbionts?

*Trends in Plant Science* **22**: 175–183.

**Croll D, Giovannetti M, Koch AM, Sbrana C, Ehinger M, Lammers PJ, Sanders IR**. **2009**.

Nonself vegetative fusion and genetic exchange in the arbuscular mycorrhizal fungus

Glomus intraradices. *The New phytologist* **181**: 924–37.

**Croll D, Sanders IR**. **2009**. Recombination in Glomus intraradices, a supposed ancient

asexual arbuscular mycorrhizal fungus. *BMC Evolutionary Biology* **9**: 13.

**Croll D, Wille L, Gamper H a, Mathimaran N, Lammers PJ, Corradi N, Sanders IR**. **2008**.

Genetic diversity and host plant preferences revealed by simple sequence repeat and

mitochondrial markers in a population of the arbuscular mycorrhizal fungus Glomus

intraradices. *The New phytologist* **178**: 672–87.

**Daboussi M-J, Capy P**. **2003**. Transposable Elements in Filamentous Fungi. *Annual*

*Review of Microbiology* **57**: 275–299.

**Emms DM, Kelly S**. **2019**. OrthoFinder: Phylogenetic orthology inference for comparative

genomics. *Genome Biology* **20**: 1–14.

**Forche A, Alby K, Schaefer D, Johnson AD, Berman J, Bennett RJ**. **2008**. The

Parasexual Cycle in Candida albicans Provides an Alternative Pathway to Meiosis for the

Formation of Recombinant Strains (J Heitman, Ed.). *PLoS Biology* **6**: e110.

**Giovannetti M, Azzolini D, Citernesi AS**. **1999**. Anastomosis Formation and Nuclear and

Protoplasmic Exchange in Arbuscular Mycorrhizal Fungi. *Applied and environmental*

*microbiology* **65**: 5571–5575.

**Halary S, Malik S-B, Lildhar L, Slamovits CH, Hijri M, Corradi N**. **2011**. Conserved

meiotic machinery in Glomus spp., a putatively ancient asexual fungal lineage. *Genome*

*biology and evolution* **3**: 950–8.

**Harrison MJ**. **1997**. The arbuscular mycorrhizal symbiosis: An underground association.

*Trends in Plant Science* **2**: 54–60.

**van der Heijden M, Klironomos J, Ursic M, Moutoglis P, Streitwolf-Engel R, Boller T,**

**Wiemken A, Sanders IR**. **1998**. Mycorrhizal fungal diversity determines plant biodiversity,

ecosystem variability and productivity. *Nature* **74**: 69–72.

**Helber N, Requena N**. **2008**. Expression of the fluorescence markers DsRed and GFP

fused to a nuclear localization signal in the arbuscular mycorrhizal fungus Glomus

intraradices. *The New phytologist* **177**: 537–48.

**Judson OP, Normark BB**. **1996**. Ancient asexual scandals. *Trends in Ecology and*

*Evolution* **11**: 41–46.

**Katoh K, Rozewicki J, Yamada KD**. **2017**. MAFFT online service: multiple sequence

alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics*: 1–7.

**Kobayashi Y, Maeda T, Yamaguchi K, Kameoka H, Tanaka S, Ezawa T, Shigenobu S,**

**Kawaguchi M**. **2018**. The genome of Rhizophagus clarus HR1 reveals a common genetic

basis for auxotrophy among arbuscular mycorrhizal fungi. *BMC Genomics* **19**: 1–11.

**Koch AM, Sanders IR, Kuhn G, Fontanillas P, Fumagalli L**. **2004**. High genetic variability

and low local diversity in a population of arbuscular mycorrhizal fungi. *PNAS* **101**: 2369–

2374.

**Kokkoris V, Chagnon P-L, Yildirir G, Clarke K, Goh D, MacLean AM, Dettman J, Stefani**

**F, Corradi N**. **2021**. Host identity influences nuclear dynamics in arbuscular mycorrhizal

fungi. *Current Biology* **31**: 1531-1538.e6.

**Krueger F**. **2015**. Trim galore. *A wrapper tool around Cutadapt and FastQC to consistently*

*apply quality and adapter trimming to FastQ files* **516**: 517.

**Kumar S, Stecher G, Li M, Knyaz C, Tamura K**. **2018**. MEGA X: Molecular evolutionary

genetics analysis across computing platforms. *Molecular Biology and Evolution* **35**: 1547–

1549.

**Lauri A, Lazzari G, Galli C, Lagutina I, Genzini E, Braga F, Mariani P, Williams JL**. **2013**.

Assessment of MDA efficiency for genotyping using cloned embryo biopsies. *Genomics* **101**:

24–9.

**Leinonen R, Sugawara H, Shumway M**. **2011**. The sequence read archive. *Nucleic Acids*

*Research* **39**: 2010–2012.

**Li H**. **2013**. Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM. **00**: 1–3.

**Li F, Upadhyaya NM, Sperschneider J, Matny O, Nguyen-Phuc H, Mago R, Raley C,**

**Miller ME, Silverstein KAT, Henningsen E, *et al.* 2019**. Emergence of the Ug99 lineage of

the wheat stem rust pathogen through somatic hybridisation. *Nature communications* **10**:

5068.

**Lin K, Limpens E, Zhang Z, Ivanov S, Saunders DGO, Mu D, Pang E, Cao H, Cha H, Lin**

**T, *et al.* 2014**. Single nucleus genome sequencing reveals high similarity among nuclei of an

endomycorrhizal fungus. *PLoS genetics* **10**: e1004078.

**Marleau J, Dalpé Y, St-Arnaud M, Hijri M**. **2011**. Spore development and nuclear

inheritance in arbuscular mycorrhizal fungi. *BMC evolutionary biology* **11**: 51.

**Masclaux FG, Wyss T, Mateus-Gonzalez ID, Aletti C, Sanders IR**. **2018**. Variation in allele

frequencies at the bg112 locus reveals unequal inheritance of nuclei in a dikaryotic isolate of

the fungus Rhizophagus irregularis. *Mycorrhiza* **28**: 369–377.

**Masclaux FG, Wyss T, Pagni M, Rosikiewicz P, Sanders IR**. **2019**. Investigating

unexplained genetic variation and its expression in the arbuscular mycorrhizal fungus

Rhizophagus irregularis: A comparison of whole genome and RAD sequencing data. *PLoS*

*ONE* **14**: 1–20.

**Mateus ID, Lee S, Sanders IR, Sanders IR**. **2021**. Co-existence of AMF with different

putative MAT-alleles induces genes homologous to those involved in mating in other fungi :

a reply to Malar et al . *The ISME Journal*.

**Mateus ID, Rojas EC, Savary R, Dupuis C, Masclaux FG, Aletti C, Sanders IR**. **2020**.

Coexistence of genetically different Rhizophagus irregularis isolates induces genes involved

in a putative fungal mating response. *The ISME Journal*.

**Mikheenko A, Prjibelski A, Saveliev V, Antipov D, Gurevich A**. **2018**. Versatile genome

assembly evaluation with QUAST-LG. *Bioinformatics* **34**: i142–i150.

**Morin E, Miyauchi S, San Clemente H, Chen ECH, Pelin A, Providencia I, Ndikumana S,**

**Beaudet D, Hainaut M, Drula E,** ***et al.*** **2019**. Comparative genomics of Rhizophagus

irregularis , R. cerebriforme , R. diaphanus and Gigaspora rosea highlights specific genetic

features in Glomeromycotina. *New Phytologist* **222**: 1584–1598.

**Ning L, Liu G, Li G, Hou Y, Tong Y, He J**. **2014**. Current challenges in the bioinformatics of

single cell genomics. *Frontiers in Oncology* **4 JAN**: 1–7.

**Paradis E, Claude J, Strimmer K**. **2004**. APE: Analyses of phylogenetics and evolution in R

language. *Bioinformatics* **20**: 289–290.

**Pontecorvo G**. **1956**. The Parasexual Cycle in Fungi. *Annual Review of Microbiology* **10**:

393–400.

**Quinlan AR, Hall IM**. **2010**. BEDTools: A flexible suite of utilities for comparing genomic

features. *Bioinformatics* **26**: 841–842.

**Remy W, Taylor TN, Hass H, Kerp H**. **1994**. Four hundred-million-year-old vesicular

arbuscular mycorrhizae. *Proceedings of the National Academy of Sciences of the United*

*States of America* **91**: 11841–11843.

**Robbins C, Cruz Corella J, Aletti C, Seiler R, Mateus ID, Lee S, Masclaux FG, Sanders**

**IR**. **2021**. Generation of disproportionate nuclear genotype proportions in Rhizophagus

irregularis progeny causes allelic imbalance in gene transcription . *New Phytologist*.

**Ropars J, Toro KS, Noel J, Pelin A, Charron P, Farinelli L, Marton T, Krüger M, Fuchs**

**J, Brachmann A,** ***et al.*** **2016**. Evidence for the sexual origin of heterokaryosis in arbuscular

mycorrhizal fungi. *Nature Microbiology* **1**: 16033.

**Sanders IR**. **2010**. Designer mycorrhizas: Using natural genetic variation in AM fungi to

increase plant growth. *ISME Journal* **4**: 1081–1083.

**Savary R, Masclaux FG, Wyss T, Droh G, Cruz Corella J, Machado AP, Morton JB,**

**Sanders IR**. **2018**. A population genomics approach shows widespread geographical

distribution of cryptic genomic forms of the symbiotic fungus Rhizophagus irregularis. *ISME*

*Journal* **12**: 17–30.

**Smith SE, Read DJ**. **2008**. *Mycorrhizal Symbiosis.* London: Elsevier Ltd.

**Spatafora JW, Chang Y, Benny GL, Lazarus K, Smith ME, Berbee ML, Bonito G,**

**Corradi N, Grigoriev I, Gryganskyi A,** *et al.* **2016**. A phylum-level phylogenetic

classification of zygomycete fungi based on genome-scale data. *Mycologia* **108**: 1028–1046.

**Sullivan MJ, Petty NK, Beatson SA**. **2011**. Easyfig: A genome comparison visualizer.

*Bioinformatics* **27**: 1009–1010.

**Sun S, Hsueh YP, Heitman J**. **2012**. Gene conversion occurs within the mating-type locus

of Cryptococcus neoformans during sexual reproduction. *PLoS Genetics* **8**.

**Thygesen K, Larsen J, Bødker L**. **2004**. Arbuscular mycorrhizal fungi reduce development

of pea root-rot caused by Aphanomyces euteiches using oospores as pathogen inoculum.

*European Journal of Plant Pathology* **110**: 411–419.

**Wyss T, Masclaux FG, Rosikiewicz P, Pagni M, Sanders IR**. **2016**. Population genomics

reveals that within-fungus polymorphism is common and maintained in populations of the

mycorrhizal fungus Rhizophagus irregularis. *The ISME Journal* **10**: 2514–2526.

**Yildirir G, Malar C M, Kokkoris V, Corradi N**. **2020**. Parasexual and Sexual Reproduction

in Arbuscular Mycorrhizal Fungi: Room for Both. *Trends in Microbiology* **xx**: 1–3.

**Yilmaz S, Singh AK**. **2012**. Single cell genome sequencing. *Current opinion in*

*biotechnology* **23**: 437–43.

**Yoon S, Xuan Z, Makarov V, Ye K, Sebat J**. **2009**. Sensitive and accurate detection of

copy number variants using read depth of coverage. *Genome Research* **19**: 1586–1592.

**Figure legends**

**Figure 1. Drop in coverage events in isolates A4, A5, A1 and C2. a,** Examples of drop in

coverage events. We plotted the normalized coverage (y) per position (x). Grey rectangles

represent the region detected by the algorithm. The horizontal dashed line represents the

normalized coverage. **b,** Number of regions showing a drop in coverage that were detected

in each isolate. **c,** Number of genes found in the regions showing a drop in coverage in each

isolate. **d,** Summary statistics of regions showing a drop in coverage: i, proportion of total

length of regions showing a drop in coverage and proportion of contigs that contain regions

with a drop in coverage. ii, Histogram representing the lengths of identified regions where a

drop in coverage was detected. iii, Coverage plot on the putative MAT-locus. Drop in

coverage was detected in isolates A4 and A5 but not in A1 and C2.

**Figure 2. Identification of orthologs of genes present in regions showing a drop in**

**coverage. a,** Orthologous groups that display two or more genes in dikaryons and only a

single gene in monokaryons. This analysis was performed independently on isolate A4 and

isolate A5. * orthologous genes containing HD2 and HD1-like genes respectively. **b,** Venn

diagram representing the number of shared orthologous groups within drop in coverage

regions between isolates A4 and A5. Only two orthologous groups were shared between the

isolates, they contain the putative MAT-locus genes HD2 and HD1-like. **c,** Synteny plot

between the two contigs containing the different alleles of the putative MAT-locus of isolates

A4 and A5. **d,** Synteny plot between the two contigs containing different alleles of other

orthologous genes. Please note that the synteny figures are made from the public available

annotations of each genome assembly. Differences in size of open-reading frames (ORF)

among isolates are due to differences in detection of ORF on each isolate and likely could

be the result of the annotation process.

**Figure 3. Single-nucleus sequence data confirms that genes contained in regions**

**where a drop in coverage was observed are nucleotype-specific.** Phylogenetic

reconstruction of single-nucleus for genes found in regions where a drop in coverage was

detected in A4 and A5 isolates. The genes are named by their membership to the

orthologous groups previously defined. Branch support consisting of 100 bootstraps is

shown. When only sequences from three nuclei were included, we performed an UPGMA

hierarchical clustering. **a,** data for nuclei from A4 isolate. **b,** data for nuclei from A5 isolate.

**Figure 4. Nucleotypes from isolate A5 share a more recent evolutionary origin to**

**isolates A1 and C2 than among them. a,** Phylogenetic reconstruction of nucleotype-

specific alleles in isolate A5 and its orthologs in isolates A1, C2 and A4. **c,** Average genetic

distances between the different nucleotype-specific alleles from two alleles from isolates A5

and their homologs in isolates A1 and C2. We show histograms representing the genetic

distance between the two nucleotype-specific alleles of isolate A5 and their homologues in

isolates A1 and C2. For comparisons between A5 and C1 or A1 we used the minimum

distance of the two alleles from A5. **d**, Scenarios of genetic similarity between the two A5 alleles and alleles from A1 and C2.

**Figure 5. Recombination events in nucleotype-specific haplotypes in A5 isolate. a,** Schematic representation of possible outcomes after fusion of two different isolates. Please note the schema illustrates different contigs, separated by blank lines and no different chromosomes. **b,**Phylogenetic relationship of different nucleotype-specific alleles among different nuclei from A1, A5 and C2 isolates. Cases where no recombination was detected. Nuclei having the same MAT-type clustered together. **c**, Phylogenetic relationship of different nucleotype-specific alleles among different nuclei from A1, A5 and C2 isolates. Cases where recombination was detected. Nuclei having the same MAT-type did not clustered together. We performed 100 bootstraps for the branch support of all phylogenetic constructions.

**Figure 6. Recombination events identified in continuous haplotypes issued from the long-read genome assembly of dikaryon isolate A5. a** Schematic representation of recombination events within continuous haplotypes. We highlight single-copy orthologous genes represented by different colors depending if they cluster with a gene of isolate A1 MAT-3 (yellow) or C2 MAT-6 (violet). Genetic recombination is demonstrated when within a continuous haplotype (grey background) a change in clustering pattern between an A5 haplotype and isolate A1 or C2 is observed. We can also observe that for few cases the two haplotypes of A5 clustered together (green). **b** Phylogenetic reconstructions of the different single-copy orthologous groups identified within the same haplotype. We can observe that the clustering pattern change within the same haplotype (i.e. A1-A5primary and C2-A5primary). We performed 100 bootstraps for the branch support of all phylogenetic constructions.

**a**

Schematic representation
Parasexual and Sexual

**A1-like**   **C2-like**
MAT-3       MAT-6

**Parasexual**
loss of heterozygosity in
dikaryon

Nucleotype A    Nucleotype B

MAT-3           MAT-6

**Sexual**
reciprocal
heterozygosity

Nucleotype A    Nucleotype B

MAT-3           MAT-6

**b**

Single-nuclei comparison
no recombination between locus and mating type

**c**

Single-nuclei comparison
recombination between locus and mating type