Cumulative growth and stress responses to the 2018-2019 drought in a European floodplain forest

Florian Schnabel[1,2,], Sarah Purrucker[2], Lara Schmitt[2], Rolf A. Engelmann[1,2], Anja Kahl[2], Ronny Richter[1,2,3], Carolin Seele-Dilbat[2,4], Georgios Skiadaresis[5], Christian Wirth[1,2,6]*

*Corresponding author: E-mail: florian.schnabel@idiv.de; Phone: +49-341-97-38595

Affiliations

[1]German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany

[2]Systematic Botany and Functional Biodiversity, Leipzig University, Leipzig, Germany

[3]Geoinformatics and Remote Sensing, Leipzig University, Leipzig, Germany

[4]Department of Conservation Biology, Helmholtz Centre for Environmental Research (UFZ), Leipzig, Germany

[5]Chair of Silviculture, Institute of Forest Sciences, Freiburg University, Freiburg, Germany

[6]Max-Planck Institute for Biogeochemistry, Jena, Germany

Keywords: climate change, hotter drought, forest ecology, dendroecology, tree rings, growth resistance, δ13C, cavitation resistance, stomatal control, temperate forests

Running head: Tree responses to consecutive drought

Abstract

Droughts increasingly threaten the world's forests and their potential to mitigate climate change. In 2018-2019, Central European forests were hit by two consecutive hotter drought years, an unprecedented phenomenon that is likely to occur more frequently with climate change. Here, we examine tree growth and physiological stress responses (increase in carbon isotope composition; $\Delta\delta^{13}C$) to this consecutive drought based on tree-rings of dominant tree species in a Central European floodplain forest. Tree growth was not reduced for most species in 2018, indicating that water supply in floodplain forests can partly buffer meteorological water deficits. Drought stress responses in 2018 were comparable to former single drought years but the hotter drought in 2018 induced drought legacies in tree growth while former droughts did not. We observed strong decreases in tree growth and increases in $\Delta\delta^{13}C$ across all tree species in 2019, which are likely driven by the cumulative stress both consecutive hotter droughts exerted. Our results show that consecutive hotter droughts pose a novel threat to forests under climate change, even in forest ecosystems with comparably high levels of water supply.

Introduction

The frequency and intensity of droughts and corresponding surges of forest dieback events around the globe are projected to increase in the 21st century (Allen et al., 2010; IPCC, 2014). This critically endangers the world's forests and the variety of ecosystem services they sustain, such as their potential to act as carbon sink (Anderegg et al., 2020) and as a nature-based solution for climate change mitigation (Griscom et al., 2017). Recent drought events, moreover, belong to a new category, so called 'hotter droughts', where low precipitation coincides with heat waves, which creates a positive feedback loop between soil-water depletion through evapotranspiration

and increased surface temperatures through reduced cooling by latent heat production (Allen,

Breshears, & McDowell, 2015; Buras, Rammig, & Zang, 2020). In 2018-2019, Central Europe has

been hit by two consecutive and hotter summer drought events, a phenomenon unprecedented at

least in the last 250 years but likely to occur more frequently with intensifying climate change

(Hari, Rakovec, Markonis, Hanel, & Kumar, 2020). The 2018 hotter drought alone had already

stronger negative effects on European ecosystems than the formerly severest drought event in 2003

(Buras et al., 2020) and induced widespread premature leaf senescence and tree mortality (Schuldt

et al., 2020). An increasing number of studies has shown that droughts can affect tree growth and

hence carbon cycling in forests for years after the actual drought event and that such 'legacy

effects' are widespread in forest ecosystems (e.g. Anderegg et al., 2015; Kannenberg et al., 2019;

Szejner, Belmecheri, Ehleringer, & Monson, 2020). The consecutive hotter drought in 2019 may

thus have critically amplified drought stress as trees were hit that already had emptied carbon

reserves, impaired hydraulic functioning due to embolism and weakened defense systems

(Anderegg et al., 2013; Schuldt et al., 2020) and only access to emptied soil water reserves.

Drought effects on forests can be analyzed retrospectively through analyses of tree-rings, which

are an archive of past growing conditions including climate and water availability (Schweingruber,

1996). In dendroecology the annual growth of trees (i.e. the width of tree rings formed each year)

is a principal indicator of drought effects, which can be analyzed through comparing growth in

drought years with mean growth in a reference period, i.e. years with 'normal' climate conditions

prior to the drought event (with growth reductions indicating drought stress) (Lloret, Keeling, &

Sala, 2011; Schwarz et al., 2020). This growth response to drought can be quantified using the

growth resistance index of Lloret et al. (2011), which may be an especially suitable approach when

rapid impact assessments are needed and no data is available on the post disturbance period. Next

to growth, the carbon isotope ratio of ^{13}C to ^{12}C in wood – called $\delta^{13}C$ – is a widely-used

physiological indicator of a tree`s water status and drought stress (Farquhar, Ehleringer, & Hubick,

1989; Grossiord et al., 2014; Jucker et al., 2017). The $\delta^{13}C$ composition in tissues of C_3 plants like

most trees is a record of the ratio between intercellular and ambient CO_2 concentration during the

time of carbon fixation that is modulated by both, CO_2 assimilation and stomatal aperture

(Farquhar et al., 1989). Under ample water supply and fully open stomata, plants discriminate

against the heavier ^{13}C in favor of the lighter ^{12}C. However, under water shortage, stomatal

conductance is more strongly down-regulated than assimilation, which induces an increase of $\delta^{13}C$

in the wood formed during drought (Farquhar et al., 1989; Grossiord et al., 2014). Thus, drought

stress can be quantified as increase in wood carbon isotope ratio ($\Delta\delta^{13}C$) between drought and

normal years. Hence, growth responses and $\Delta\delta^{13}C$ combined provide a powerful tool to quantify

drought effects on trees.

Tree species vary greatly in their susceptibility to drought due to physiological and morphological

differences. Among other features such as root system architecture (Sánchez-Pérez, Lucot, Bariac,

& Trémolières, 2008), two key factors that might drive tree species reactions to drought are

stomatal control and resistance to cavitation (Choat et al., 2012; Martínez-Vilalta & Garcia-Forner,

2017; McDowell et al., 2008). Stomatal closure in response to water deficits enables plants to

avoid critically low water potentials through transpiration losses and thus hydraulic failure but

species differ largely in their type of stomatal control (Martínez-Vilalta & Garcia-Forner, 2017;

McDowell et al., 2008): Isohydric or water saving species close their stomata fast during water

shortage, while anisohydric or water spending species keep their stomata open and continue to

transpire (Martínez-Vilalta & Garcia-Forner, 2017). Next to stomatal control, xylem resistance to

cavitation is a key determinant of tree responses to drought as embolism decreases water

availability, which leads to desiccation and at extreme levels to tree death (Choat et al., 2012). It

is conceivable that stomatal control and cavitation resistance interact, as a water spending behavior

necessitates a continued water uptake via roots and, all else being equal, carries an increased risk

for xylem cavitation (McDowell et al., 2008). However, whether this translates into water spending

species exhibiting generally higher cavitation resistance and *vice versa* still remains elusive as

some studies found indications for such a correlation (Klein, 2014; Martínez-Vilalta & Garcia-

Forner, 2017) while others did not (Kröber, Zhang, Ehmig, & Bruelheide, 2014). We expect water

saving species to show earlier growth and $\Delta\delta^{13}C$ responses, while water spending species may face

high cavitation risks during severe and prolonged drought conditions characterized by very low

soil moisture availability. Hence, for understanding and generalizing the effects of consecutive

droughts on forests, tree species should be examined that differ in such traits.

The high tree species richness of floodplain forests (Ward, Tockner, & Schiemer, 1999) makes

them ideally suited for comparative studies of tree species reactions to consecutive droughts as

they are one of the few systems where coexisting mature trees spanning an entire gradient of

hydraulic behaviors can be found. Floodplain forests rank among the most rapidly disappearing

ecosystems due to land conversion and drainage (Leuschner & Ellenberg, 2017; Mikac et al., 2018)

and novel climatic conditions – like prolonged droughts – may amplify this trend through changing

the hydrological regimes on which these forests depend. For instance, sinking groundwater levels

may increase tree growth sensitivity to drought and susceptibility to drought-induced dieback

(Mikac et al., 2018; Skiadaresis, Schwarz, & Bauhus, 2019) and this might bring these forests,

which are among the most dynamic, productive and diverse Central European habitats (Kowalska

et al., 2020; Tockner & Stanford, 2002), closer to a tipping point. On the contrary, the higher water

availability in floodplain forests may buffer drought effects to a certain extent as trees might have

access to groundwater in addition to precipitation-derived moisture (Heklau, Jetschke, Bruelheide,

Seidler, & Haider, 2019). Hence, it is conceivable that if drought effects on growth and $\Delta\delta^{13}C$

were observed in floodplain trees, other forest ecosystems might experience even stronger effects.

Here, we focus on the effect of the two consecutive drought years 2018-2019 characterized by

extremely hot and dry conditions (Fig. 1), as well as their cumulative effects, on tree growth and

$\Delta\delta^{13}C$ as physiological stress response. To this end, we reconstruct the stress exerted by this

unprecedented event and compare it to past (single) drought events based on tree-ring records from

the dominant tree species – *Quercus robur*, *Acer pseudoplatanus* and *Fraxinus excelsior* – in the

Leipzig floodplain forest, one of the few remaining and thus highly protected floodplain forests in

Central Europe (BMU & BfN, 2021; Günther-Diringer et al., 2021). We sampled trees in two

environmental strata representing topographic differences in distance to groundwater. We expect

the results for the hypotheses proposed below to be more pronounced in the drier stratum.

Specifically, we tested the following hypothesis:

H1 Drought stress responses in 2018 – measured as growth reduction and a positive $\Delta\delta^{13}C$ – are

comparable to stress responses during former drought years.

H2 The consecutive drought years 2018-2019 lead to a drastic growth reduction and a further

increase in $\Delta\delta^{13}C$ in 2019.

H3 Water spending species respond faster to drought stress (already in 2018), while water saving

species react later but show stronger reactions to the consecutive drought in 2019.

Fig. 1 Annual standardized water balance of precipitation minus potential evapotranspiration (A, January-

December) and mean growing season temperature (B, April-September) per year from 1979-2019 in the

Leipzig floodplain forest. The water balance was calculated as Standardized Precipitation

Evapotranspiration Index (SPEI; Vicente-Serrano, Beguería, & López-Moreno, 2010). Points are colored

according to their value with deeper red indicating increasing drought and temperature. The horizontal line

in (A) represents the long-term mean, negative values indicate water deficits and positive values water

surpluses. SPEI values below -1 and above 1 can be considered exceptionally dry and wet, respectively.

See Figs. S1-S2 for additional SPEI lengths, climatic and hydrological variables that we used to identify

drought events.

Methods

Study site

In this study, we used data collected from a Central European floodplain forest ecosystem located in the northwest of the city of Leipzig, Germany. The Leipzig floodplain forest is one of the few remaining and thus highly protected floodplain forests in Central Europe (BMU & BfN, 2021; Günther-Diringer et al., 2021) and lies in the transition zone between maritime and continental climate characterized by warm summers, with an annual mean temperature of 9.6 °C and an annual precipitation sum of 525 mm (1979-2019; DWD, Station Leipzig/Halle). Its main rivers Weiße Elster, Luppe, Pleiße and Parthe formed the floodplain landscape, but their course and thus the floodplain forest itself has been strongly influenced by human interventions over the last centuries (Gutte, 2011). The straightening of rivers as well as dike and canal constructions strongly influenced the hydrological regime of the floodplain forest, which today does not experience regular flooding anymore (Haase & Gläser, 2009). The floodplain soils originated from an accumulation of alluvial sediments, such as gravel, sand and loam, as result of several glacial periods (Haase & Gläser, 2009). These are nowadays covered by an alluvial clay layer with a thickness between 1-4 m, rich in nutrients and with a high pH (around 6-7) (Gutte, 2011; Haase & Gläser, 2009). The principal soil available to trees is thus a loamy Vega, with partly gleyed conditions, above gravel and sand filled with groundwater.

Tree species

The contemporary floodplain forest ecosystem can be characterized as *Ficario-Ulmetum* Knapp ex Medwecka-Kornas 1952 with oak, elm and ash being the dominant tree species (Härdtle et al., 2020). The absence of flooding, however, resulted in an on-going gradual shift to an oak-hornbeam

forest (*Galium-carpinetum stachyetosum*) and allowed other tree species (especially maple), which

are intolerant to flooding, to become dominant. Moreover, elm (*Ulmus minor*) largely disappeared

from the tree canopy due to the Dutch Elm Disease since the 1960s. Nowadays, the dominant tree

species of Leipzig's floodplain forest are *Quercus robur L.* (hereafter oak), *Acer pseudoplatanus*

L. (hereafter maple) and *Fraxinus excelsior L.* (hereafter ash) (Haase & Gläser, 2009; Richter, Reu,

Wirth, Doktor, & Vohland, 2016), on which we focus in the present study. These three species

feature contrasting adaptations to drought in terms of stomatal control and cavitation resistance,

which allowed us to explore a range of species response strategies to consecutive drought stress.

In terms of stomatal control, former studies classified oak (Cocozza et al., 2020; Thomsen,

Reisdorff, Gröngröft, Jensen, & Eschenbach, 2020) and maple (Köcher, Gebauer, Horna, &

Leuschner, 2009; Lemoine, Peltier, & Marigo, 2001; Leuschner, Wedde, & Lübbe, 2019) as rather

water saving (isohydric), while ash was shown to follow a water spending (anisohydric) strategy

(Köcher et al., 2009; Lemoine et al., 2001; Leuschner et al., 2019). Consistent with this view,

studies focussing on sap flow measurements on mature trees, which can be considered as proxy

for tree transpiration and hence stomatal aperture, reported a significant downregulation of sap

flow with decreasing soil water availability for maple but not for ash (Brinkmann, Eugster,

Zweifel, Buchmann, & Kahmen, 2016; Hölscher, Koch, Korn, & Leuschner, 2005; Köcher et al.,

2009), indicating water saving versus water spending modes, respectively. Importantly, we view

stomatal control here as a gradient and not as a dichotomy between water saving or water spending

behaviour (but see Martínez-Vilalta & Garcia-Forner, 2017), with classifications depending

strongly on the compared species. However, quantitative data on traits indicative of stomatal

control like sap flow is scarce (particularly for *Q. robur*) and we are not aware of any study that

compared mature individuals of all herein examined species under similar site conditions. We

therefore assembled high-resolution sap flow and soil moisture data recorded during the 2018

drought in the Leipzig floodplain forest (Leipzig Canopy Crane facility) to provide a quantitative

comparison of species-specific sensitivity to decreasing soil moisture under severe drought (Fig.

2a-b). Consistent with the classification above, oak and maple significantly downregulated sap

flow under drought conditions (indicating water saving behaviour) while ash maintained similar

sap flow rates (indicating water spending behaviour) (Fig. 2b). In terms of species resistance to

cavitation, we relied on published values of the water potential at which 50% of xylem conductivity

is lost due to cavitation (Ψ_{50}, Choat et al., 2012), the most common measure of embolism resistance

in trees (Choat et al., 2012). This comparison indicates a similar cavitation resistance in oak (-2.8

MPa) and ash (-2.8 MPa) while maple is less resistant (-1.6 MPa).

Fig. 2 Soil moisture development (a) and sap flux density (J_s) regulation of oak, maple and ash (b) during the 2018 hotter drought in the Leipzig floodplain forest. The observed soil moisture development (a) was used to delineate two periods with contrasting soil moisture conditions, a moist and dry period, respectively. During the dry period soil moisture levels approached 0.24 m^3/m^3 (red horizontal line), the permanent wilting point of vegetation on clay soils (Weil & Brady, 2017). Boxplots (b) show daily maxima in J_s during the two month period with moist soil (mid-Mai to mid-July) and during the period with dry soil at later stages during the 2018 drought (mid-July to mid-September). A statistically significant downregulation of J_s under dry compared to moist conditions is indicated by asterisks over the respective species' boxplot ('***' $p < 0.001$; '**' $p < 0.01$; '*' $p < 0.05$). See Methods S1 for details on the sap flow and soil moisture measurements and analyses.

Drought year identification

The definition and identification of drought is central to the analysis of drought effects. Here, we define drought as period with water deficits compared to normal conditions, where 'normal' can be quantified as a percentile of the long-term mean of meteorological or hydrological variables (Schwarz et al., 2020; van Loon et al., 2016). Following suggestions by Schwarz et al. (2020) we selected drought years based on climatic and hydrological information alone without considering tree growth reductions to avoid a biased selection that could for example result in the exclusion of drought years without reduced growth. We used the Standardized Precipitation Evapotranspiration Index (SPEI; Vicente-Serrano et al., 2010) and river discharge data to identify drought years. The SPEI is a commonly used drought index (Hari et al., 2020; Schwarz et al., 2020; Skiadaresis et al., 2019) based on the standardized monthly water balance of precipitation minus potential evapotranspiration. It can quantify drought severity according to a droughts intensity and duration and can be calculated at different time scales (e.g. 1-12 months) (Vicente-Serrano et al., 2010). Here, we used three different SPEI lengths that represent the climatic water balance of the main vegetation period (SPEI for 3 months, Mai-July), the full vegetation period (SPEI for 6 months, April-September) and the full year (SPEI for 12 months, January-December) for each year and with a 40-year reference period (1979-2019; Fig. S1). SPEI series were calculated with the SPEI package (Beguería & Vicente-Serrano, 2017) in R from monthly precipitation (mm) and potential evapotranspiration (mm) data derived from the weather station located closest to the study sites (DWD, Station Leipzig/Halle, ID 2932; Fig. S1).

We classified years with SPEI values ≤ -1 as drought years, years with SPEI values ≥ 1 as

particularly wet and years with values between -1 and 1 as 'normal' (McKee, Doesken, & Kleist,

1993). To take into account the hydrological regime of the floodplain forest, which is in addition

to local precipitation strongly influenced by its rivers, we compared the SPEI derived classification

to river discharge calculated for the same periods as the SPEIs (Fig. S2). We considered only years

without particularly high discharge as drought years. Focusing on a 20-year period before the

2018-2019 consecutive drought, we selected 2005, 2009 and 2017 as reference years with normal

climatic conditions, while single drought years – in contrast to the 2018-2019 consecutive drought

– were 2003, 2006 and 2015 (Fig. 1; Figs. S1-S2; hereafter 'single drought years'). We did not

consider a longer period to minimize the effect of past forest management and ground water

fluctuation related influences on tree growth. Both, the drought in 2018 and the one in 2019 were

the most severe droughts in the last 40 years (i.e. they had the lowest SPEI values), but 2018 had

slightly lower SPEI values and was especially characterized by an extreme heat wave during the

vegetation period.

Tree selection and increment core extraction

We selected trees for extracting wood increment cores from permanent forest research plots of the

"Lebendige Luppe" (living Luppe river) project (Scholz et al., 2018), which cover a gradient in

topographic distances to the groundwater level (Fig. S3). The project features three distinct strata

of distance to groundwater: dry (> 2m), intermediate (1-2m) and moist (≤ 1m) plots, with 20 plots

per stratum each 0.25ha in size. Plots were not flooded since 1973 due to flood control measures

(dikes, river-straightening etc.), except for winter 2011 and summer 2013, when the area of Leipzig

experienced extreme flood events. We chose to sample trees on dry and moist plots to cover both

ends of the gradient of hydrological site conditions within the floodplain forest (Fig. S3). Across

these plots, we extracted tree increment cores from at least 40 tree individuals per species (20 trees

per stratum) from each of the three dominant tree species oak, maple and ash, amounting to 120

sampled trees. From each tree, we extracted one increment core at a height of 80cm with a Ø 5mm

increment corer (Suunto, Sweden) in January-February 2020, i.e. in the winter after tree ring

formation of the second consecutive drought year 2019 was completed. Trees with diameters at

breast height (dbh) > 20cm were selected according to their dominance, past management history

and health status. Competition for light is a central determinant of tree growth and δ^{13}C that might

complicate the detection of drought effects (Grossiord et al., 2014). We therefore sampled only

dominant and co-dominant individuals, i.e. trees belonging to category 1-2 according to the

classification of Kraft (1884), that were no direct competitors and further excluded plots that

showed signs of forest management in recent years. We further selected only healthy appearing

trees, excluding those ash trees visually affected by 'ash dieback' (*Hymenoscyphus fraxineus*) and

those maple trees visually affected by the 'sooty bark disease' (*Cryptostroma corticale*). Both

fungal pathogens had caused widespread tree damages and diebacks in the Leipzig floodplain

forest during the 2018-2019 consecutive drought and especially very few ash trees were

completely unaffected (Wirth et al., 2021). We used the classification key of Lenz, Straßer,

Baumann, and Baie (2012) for ash dieback infestation and sampled only trees showing no to only

little signs of infestation (level 0-2 of infestation levels 0-5) based on annual infestation records

for four years prior to sampling, while we sampled maple trees that showed no visual sign of sooty

bark disease. Importantly, our sample is thus representative for the most vital individuals of the

entire population. Since the number of trees fulfilling these strict criteria was too low within the

plot area, we sampled also oak and maple trees in the direct vicinity of the plots.

Tree growth analysis

Tree cores were dried at 70 °C for at least 3 days and then clamped in wooden alignment strips. For surface preparation, we used a core microtome (WSL, Switzerland) (Gärtner & Nievergelt, 2010) to enhance visibility of tree ring boundaries. Tree ring-width was measured with a LINTAB 6 measuring table and the TSAPWin Professional 4.64 program © 2002-2009 Frank Rinn / RINNTECH with an accuracy of 1/1000 mm. The measured sequences were cross-dated against a species-specific master chronology developed in former works for the same area as well as against each other using COFECHA (Grissino-Mayer, 2001). This allowed us to identify missing rings, which were more often found in maple trees and in the consecutive drought years 2018-2019. Years without growth were included as zero for the respective year. Sequences that could not be dated unequivocally were excluded from further analysis. The final number of trees included for growth analysis was 114 trees instead of the planned 120, including 40 oak, 32 maple and 42 ash trees from 11 moist and 15 dry plots. Mean series length was 109 years for oak, 79 years for maple and 94 years for ash trees (Table S1).

Tree ring-width provides an integrated record of past growth conditions as influenced by environmental factors including but not limited to climate and shows an inherent decrease in ring-width with increasing tree size (Schweingruber, 1996). As we focus here on climatic influences on growth, we removed age-related trends from the raw tree ring-width chronologies via a Negative Exponential curve (Fritts, 1976), which provided the best compromise between removing long-term age trends and preserving decadal variability in growth using the package dplR (Bunn, 2008; Bunn et al., 2020). We assessed the climatic sensitivity of tree growth through computing

bootstrapped Pearson's correlation functions between species-specific chronologies and monthly

climatic variables (Fig. S4), using the package treeclim (Zang & Biondi, 2015). Species-specific

chronologies are shown in Fig. S5.

We quantified the growth response of trees to the consecutive drought years 2018 and 2019 as well

as to single drought years (hereafter 'growth response') for each individual tree using the growth

resistance index proposed by Lloret et al. (2011) as:

$Growth\ response = \frac{Dr_{growth}}{PreDr_{growth}} - 1$ (1)

where Dr_{growth} is a tree's detrended growth in drought year(s) and $PreDr_{growth}$ is a tree's detrended

growth in the reference period characterized by normal climatic conditions. The growth response

index is thus standardized around zero with positive values meaning higher and negative values

lower growth during drought year(s) compared to reference years. We calculated the growth

response for 2018, 2019 and for single drought years (using their mean growth response in all

analysis). Recent decades experienced an unprecedented surge in temperatures and drought events

(Buras et al., 2020; Schuldt et al., 2020), making the use of a continuous reference period of several

years before drought events that is not influenced by drought itself difficult. We therefore used the

mean growth in three years (2005, 2009, 2017) that were characterized by normal climatic

conditions and not proceeded by a drought year (see above) to calculate $PreDr_{growth}$. We used

several years to calculate $PreDr_{growth}$ and the growth response in single drought years as this should

reduce the influence of outliers caused by individual tree reactions to other factors than climate

(e.g. changes in competitive interactions, waterlogging). However, as the choice of growth data

and a reference period can strongly influence results of the index used (Schwarz et al., 2020), we

tested for the robustness of here reported relationships. We examined growth responses in detail

based on raw and detrended ring-width and further compared growth responses calculated with a

one-year pre-period (2017 was the only climatically normal year before the 2018-2019 consecutive

drought; Fig. S1) to growth responses calculated with the mean reference period detailed above.

We quantified drought legacies in tree growth as observed growth minus predicted growth as

expected based on the climatic conditions in the year after the drought event (Anderegg et al.,

2015; Kannenberg et al., 2019). Tree growth in the last 40 years (1979-2019) was predicted using

tree-specific regressions between detrended tree ring-width and the most important drought-related

climate variable for each species (the climate variable, which showed the strongest correlations

with growth; Fig. S4): These were temperature in August ($r = 0.51$) for oak, SPEI 12 of July ($r =$

0.64) for maple and SPEI6 of July for ash ($r = 0.63$). We used the climate variable with the highest

growth-climate correlation, since the ability to quantify drought legacy effects depends on an

informative (i.e. significant) null model (but see Anderegg et al., 2015). Reported drought legacy

effects thus quantify the deviation of observed growth from expected growth based on climate in

year 1 after single drought years (2003, 2006, 2015) and after the hotter drought in 2018, that is,

in the years 2004, 2007, 2016 and in the consecutive drought year 2019. Climatic conditions in

year 1 after all single drought years were neither exceptionally dry nor wet (Fig. 1), providing a

suitable baseline against which the legacy effects of consecutive drought can be compared.

Carbon isotope analysis

The stable carbon isotope composition (δ^{13}C) in wood of the same cores was measured after completing tree ring-width measurements. The tree rings of the herein analysed consecutive drought years 2018-2019, of single drought years (2003, 2006, 2015) and of reference years (2005, 2009, 2017) were separated and their wood tissue homogenized. Some trees, especially maple ones, did not form tree rings during the 2018-2019 consecutive drought, likely due to intense drought stress. As their δ^{13}C could thus not be analysed, we excluded these trees from our isotope analysis (6 maple and 1 oak tree). The homogenized material of the tree rings in reference years (2005, 2009 and 2017) and single drought years (2003, 2006 and 2015), was pooled by mixing equal shares of the material from each of the three years. The isotope analysis was done at the BGC stable isotope laboratory of the Max Planck Institute for Biogeochemistry in Jena, Germany. The results were expressed as isotopic ratio δ^{13}C, calculated with the equation of Farquhar et al. (1989) as follows:

$$\delta^{13}C = \left(\frac{\delta^{13}C\ (sample)}{\delta^{13}C(standard)} - 1\right) * 1000‰ \tag{2}$$

where $\delta^{13}C$ (sample) and $\delta^{13}C$ (standard) are the abundance ratios between ^{13}C and ^{12}C of the given sample and Vienna PeeDee Belemnite international standard (VPDB). Isotope ratios were expressed in δ-notation in per mil units (‰). We calculated the increase in δ^{13}C from reference to drought years for each individual tree as indicator of a tree's physiological stress response to drought as:

$$\Delta\delta^{13}C = Dr_{\delta^{13}C} - PreDr_{\delta^{13}C} \tag{3}$$

where $Dr_{\delta^{13}C}$ is the isotope composition in drought year(s) and $PreDr_{\delta^{13}C}$ the isotope composition

in the reference years (see e.g. Grossiord et al., 2014). Positive values of $\Delta\delta^{13}C$ thus indicate higher

and negative values lower stress during drought year(s) compared to reference years. Drought and

reference years used to calculate $\Delta\delta^{13}C$ were the same as in the growth response analysis.

Statistical analysis

We used linear mixed-effect models (LMMs) to understand the effects of consecutive drought

years on tree growth and $\Delta\delta^{13}C$ in comparison to single drought years (using their mean growth

response and $\Delta\delta^{13}C$ in all analysis). We were further interested in understanding how these effects

were modulated by changes in a trees distance to groundwater. We fitted species-specific LMMs

for analysing the growth response and $\Delta\delta^{13}C$ with the packages lme4 (Bates, Mächler, Bolker, &

Walker, 2015) and lmerTest (Kuznetsova, Brockhoff, & Christensen, 2017) and a significance

level of α=0.05. Drought event (single droughts, 2018, 2019), groundwater level (dry > 2m, moist

≤ 1m) and their interaction were modelled as fixed effects, while tree identities nested within plots

were used as nested random effects to account for differences between plots and for repeated

measurements on each individual tree. We selected the most parsimonious model structure via

backward model selection, first adjusting the random (likelihood ratio tests) and then the fixed

effect model structure (F-tests), using the step function in lmerTest. The most parsimonious LMM

structure consistently retained only a fixed effect of drought event and tree identity nested within

plots as random effects, indicating that water table did not significantly (P>0.05) influence

observed relationships. Only for the growth response LMM of oak we found a significant

interaction (P=0.0413) between drought event and groundwater level, which however disappeared

when using non-detrended growth data or a one-year reference period. Therefore, to report only

the most robust relationships, we present all final LMMs with drought event as the only fixed

effect. Final LMMs (Table S2) were fit using restricted maximum likelihood estimation (REML)

and marginal means and confidence intervals (95%) were predicted with the ggeffects package

(Lüdecke, 2018). We used post-hoc pairwise comparisons with adjusted p-values for multiple

comparisons (Tukey's Honest Significant Difference) to compare differences between drought

events using the emmeans package (Lenth, 2020) (Table S3). Model assumptions (normality,

independence and homogeneity of variance) where visually checked through examining model

residuals and through quantile-quantile plots. Drought legacy effects were analysed using the same

modelling procedure (see Table S2). All analyses were conducted in R version 4.0.3 (R Core Team,

2020).

Results

We found pronounced responses to drought stress in terms of tree growth and $\Delta\delta^{13}C$ across the

examined tree species, with strongest stress responses in the second of two consecutive drought

years (2019). The mean growth response to single drought years (2003, 2006 and 2015) before the

2018-2019 consecutive drought ranged around zero for oak and maple, while growth in ash tended

to be reduced (Fig. 3a-c). This indicates a similar tree growth in single drought years and in

climatically 'normal' years for oak and maple but not for ash. Growth of oak and maple even

tended to be higher in 2018 compared to normal years (mean growth response above zero). The

hotter drought in 2018 did not induce growth responses in oak and maple that differed significantly

from single drought years (P>0.1 for both species) but ash experienced a significantly stronger

growth reduction (t=-2.94, P=0.004; Fig. 3c; Table S2). In 2019, the second consecutive and

extreme drought year, the growth reduction in all species was significantly stronger than in single

drought years (oak t=-2.00, P=0.049; maple t=-2.74, P=0.008; ash t=-7.22, P<0.001; Fig. 3a-c;

Table S2) and in comparison to 2018 (Table S3). Observed growth responses were largely

insensitive to the type of growth data (raw or detrended) and reference period (one-year or pooled

years) used (Figs. S6-S7). We used species-specific models but provide evidence for significant

differences between species in Fig. S8. Distance to groundwater had an overall small influence on

the growth response of the examined species (non-significant effect of groundwater level for maple

and ash). Only for oak we found indications for a smaller growth response on moist plots in 2019

(significant interaction of drought year and groundwater level, P=0.041). Among the three

analysed species, ash showed the highest growth sensitivity to drought (especially to SPEIs

indicating summer drought) while oak was the least sensitive (Fig. S4).

We did not find drought legacy effects in tree growth after single drought years, that is, observed

tree growth in year 1 after these droughts was not significantly lower than growth predicted based

on climate (Fig. 4a-c; Table S2). For ash, observed growth even tended to be higher than predicted

(Fig. 4c). In contrast, the hotter drought year 2018 induced substantial legacy effects in oak and

ash but not in maple (Fig. 4a-c), that is, observed growth in the second consecutive drought year

2019 was significantly lower than expected based on climate. Legacy effects in 2018 were

significantly different from legacy effects in single drought years for oak and ash but not for maple

(oak t=-3.05, P=0.004; ash t=-4.93, P<0.001; maple t=-0.89, P=0.380; Fig. 4a-c; Table S2).

Fig. 3 Growth response and increase in the carbon isotope ratio ($\Delta\delta^{13}C$) in wood of oak, maple and ash in drought years. The figure shows the growth response (upper panels) and $\Delta\delta^{13}C$ (lower panels) in the consecutive hotter drought years 2018 and 2019 compared to the mean growth response and $\Delta\delta^{13}C$ in single drought years (2003, 2006 and 2015). Zero corresponds to a comparable growth and $\delta^{13}C$ in drought and climatically normal years. Negative growth response values indicate growth reductions while positive $\Delta\delta^{13}C$ values indicate stress increases during drought compared to normal years. The growth response and $\Delta\delta^{13}C$ were calculated with Eq. 1 and Eq. 3, respectively. Black points show estimated marginal means and error bars the 95% confidence intervals of linear mixed-effect model fits, with non-overlapping confidence intervals indicating signfEcant differences. Coloured points show the growth response and $\Delta\delta^{13}C$ values per

tree and species (oak n=40, n=39; maple n=32, n=26; ash n=42, n=42) and are jittered to enhance visibility.

The tree ring-widths have been detrended with a negative exponential function. Statistically significant

differences in the growth response and $\Delta\delta^{13}$C between the years 2018 and 2019 compared to single drought

years are indicated by asterisks over the respective year ('***' p <0.001; '**' p <0.01; '*' 'p <0.05).

Fig. 4 Drought legacy effects in growth of oak, maple and ash in the year following drought events. The

figure shows drought legacy effects induced by the hotter drought year 2018 compared to mean legacy

effects induced by single drought years (2003, 2006 and 2015). Legacy effects were quantified as observed

minus predicted (detrended) tree ring-width based on climate in year 1 after the drought event. Zero

corresponds to growth as expected based on climate conditions, while negative values indicate drought

legacies in form of lower than expected post-drought growth. Black points show estimated marginal means

and error bars the 95% confidence intervals of linear mixed-effect model fits, with non-overlapping

confidence intervals indicating signficant differences. Coloured points show legacy effects per tree and

species (oak n=40; maple n=32; ash n=42) and are jittered to enhance visibility. Statistically significant

differences in legacy effects between 2018 compared to single drought years are indicated by asterisks

('***' p <0.001; '**' p <0.01; '* 'p <0.05).

We found positive $\Delta\delta^{13}C$ values across all species and drought years, indicating tree physiological

stress responses to drought irrespective of drought type (single or consecutive; Fig. 3d-f).

However, the magnitude of $\Delta\delta^{13}C$ increases varied strongly between drought years and species.

For oak and maple, $\Delta\delta^{13}C$ was not significantly enhanced in 2018 compared to single drought

years (P=0.85 and P=0.79), while ash had significantly higher $\Delta\delta^{13}C$ values (t=2.85, P=0.006; Fig.

3f; Table S2). Across all species, we found a strong increase in $\Delta\delta^{13}C$ in 2019 compared to single

drought years (oak t=3.93, P<0.001; maple t=2.80, P=0.007; ash t=14.80, P<0.001; Fig. 3d-f; Table

S2) and in comparison to 2018 (Table S3). The $\Delta\delta^{13}C$ increase was strongest for ash. Distance to

groundwater had no significant influence on $\Delta\delta^{13}C$ for all examined species. Together these results

indicate that drought stress in 2018 was, except for ash, comparable to stress in former single

drought years, while the second consecutive drought year 2019 induced the strongest growth

reductions and increases in $\Delta\delta^{13}C$ across all species.

Discussion

Using tree growth reductions and increases $\Delta\delta^{13}C$ as indicators of drought stress, we report a strong

increase in drought-related stress in the second of two consecutive hotter drought years across all

examined species. Drought responses were strikingly consistent for both indicators (growth

response and $\Delta\delta^{13}C$; Fig. 3), but the timing and magnitude of responses were species-specific: Oak

showed the overall smallest stress response followed closely by maple with the strongest response

in ash. The 2019 drought, although an extreme drought as well, was meteorologically less severe

than the preceding drought year 2018, as indicated by less negative SPEI values and lower growing

season temperature (Fig. 1). This indicates that the cumulative drought effect exerted by both years

was likely the principal driver of the stress increase in 2019, which is consistent with our result

that the 2018 drought – in contrast to preceding drought years – induced significant legacies in tree

growth (Fig. 4). The 2018 hotter drought was the severest drought so far recorded in Central

Europe (Buras et al., 2020; Hari et al., 2020; Schuldt et al., 2020), but, as predicted, we found

physiological stress increases ($\Delta\delta^{13}C$) to be comparable to former single drought years and tree

growth to be largely within the range of climatically 'normal' years. Hence, the comparably high

water availability in floodplain forests may partly buffer tree stress responses to single but not to

consecutive drought years.

Our conclusion that the effects of single drought years were buffered to some extent contrasts with

the dramatic drought effects reported across European forests in 2018 that suffered widespread

defoliation, xylem hydraulic failure and mortality (Buras et al., 2020; Schuldt et al., 2020) but is

consistent with other floodplain forest studies. For instance, the exceptionally high gross primary

production during the warm spring in 2018 was found to compensate for losses later that year due

to drought in a Czech floodplain forest (Kowalska et al., 2020). Similarly, tree growth recovered

within two years after the 1976 drought for all herein analyzed floodplain forest species, which

was attributed to the drought buffering effect of water availability in floodplain forests (Heklau et

al., 2019). Nonetheless, we found physiological stress increases ($\Delta\delta^{13}C$) in 2018 while tree growth

in most species did not react. This confirms the view of clearer drought signals in $\Delta\delta^{13}C$ compared

to tree ring-width, potentially due to tree growth being maintained from carbon reserves even under

low soil water availability (Jucker et al., 2017).

This picture changed dramatically in 2019. As hypothesized, we observed the strongest stress

responses in the second consecutive drought year, likely driven by the cumulative effect of both

drought years. Drought legacy effects (Anderegg et al., 2015) were found to be widespread in

forests and to affect tree growth and $\Delta\delta^{13}C$ 1-5 years after the actual drought event (Anderegg et

al., 2013; Anderegg et al., 2015; Gazol et al., 2020; Kannenberg et al., 2019; Lloret et al., 2011;

Szejner et al., 2020). We observed significant drought legacy effects in tree growth after the 2018

hotter drought but not after former single drought years. Hence, in a system where drought legacy

effects have not been observed previously, the hotter drought in 2018 was severe enough to induce

such legacies. Trees were thus likely still suffering from 2018 when the second drought year 2019

struck. It should be considered that these drought years (2003, 2006, 2015) were – before the 2018-

2019 drought – themselves considered as some of the severest droughts in Central Europe with

strongly decreased tree growth, increased $\delta^{13}C$ and increased tree mortality being reported at other

sites (Allen et al., 2015; Büntgen et al., 2021). Their comparably low effect thus supports our view

of high water availability in floodplain forests partly buffering tree stress responses and

simultaneously underlines the unprecedented nature of 2018-2019. Results on drought legacy

effects from former studies are only partially applicable to our case, as these focused on legacy

effects in post drought periods in which trees where already (partially) recovering. In such cases it

is difficult to disentangle drought legacy effects from other changes in a trees environment that

may affect tree growth and $\Delta\delta^{13}C$ (Gazol et al., 2020). In contrast, we focus here on two

consecutive hotter drought years, unprecedented in their severity at least since 250 years (Hari et

al., 2020), which left the trees no time to recover. The few studies that studied prolonged droughts,

moreover, did not examine the cumulative built-up of drought effects from year-to-year as they

used either mean tree growth across drought years or growth in the last year of drought to calculate

growth responses to drought (Schwarz et al., 2020). In comparison, the strong reactions we report

for 2019 should be mainly attributable to legacy effects of 2018, although drought timing may

have also played a role as we discuss below. Other changes in the trees' environment like reduced

competition for light are unlikely within a single year. In addition, forest management can be

excluded as potential cause as we did not sample trees in stands that experienced recent

interventions. Similarly, assessments of the 2018 hotter drought found early signs of strong

drought legacy effects (Buras et al., 2020; Schuldt et al., 2020).

Several physiological mechanisms could explain drought legacy effects (Anderegg et al., 2015)

and thereby cumulative drought stress. Drought-induced xylem cavitation may impair growth and

transpiration (and thus effect $\Delta\delta^{13}C$) (McDowell et al., 2008; Schuldt et al., 2020). Under

consecutive drought, this damage persists, while vulnerability to cavitation may continue to

increase under successive drought stress (Anderegg et al., 2013). In the second drought year, less

nonstructural carbohydrates (NSC) reserves were likely left for xylem repair, growth and

especially for keeping up the trees' defense system, which increases their susceptibility to pests

and pathogens (Anderegg et al., 2013; Hartmann & Trumbore, 2016; McDowell et al., 2008;

Schuldt et al., 2020). Large parts of the here examined stands suffered severe damages and

mortality through sooty bark disease, ash dieback and secondary beetle infestations in 2019, which

points to depleted NSC stores causing a collapse of the trees' immune system. We studied only

the most vital tree individuals of the population thus largely excluding disease effects from our

sample. However, as the majority of ash trees in the forest were affected to some degree (Wirth et

al., 2021), we cannot completely disentangle whether the species intrinsic traits, incipient ash-

dieback or their interaction caused the strong stress response in this species. Finally, drought

induces shifts in carbon allocation in favor of the canopy and root system at the expense of radial

growth (Kannenberg et al., 2019) and, when photosynthesis is insufficient to meet demands, NSC

reserves are utilized to maintain autotrophic respiration, growth and tissue repair (Hartmann

& Trumbore, 2016; Richardson et al., 2013). This enriches the reserve pool and tissues built from

it in ^{13}C as the isotopically lighter ^{12}C is turned over faster than ^{13}C. Both processes may have

further contributed to the strong decrease in tree-ring growth and increase in $\Delta\delta^{13}C$ in 2019 in

addition to fractionation through stomata closure. Interestingly, if the strong $\Delta\delta^{13}C$ increases in

2019 were indeed (partly) triggered by NSC dynamics, we could exclude that the drought strongly

impaired phloem transport and hence carbon translocation in the here examined trees. That the

latter could happen has been shown for example in drought experiments with Norway spruce

(Hartmann, Ziegler, & Trumbore, 2013).

In addition to physiological drought legacies, different meteorological and abiotic conditions may

have contributed to the strong stress responses observed in 2019. The increasing frequency of

droughts and exceptionally hot years, especially since 2015 (Schuldt et al., 2020) (Fig. 1), may

have successively reduced the ability of trees to withstand drought (Anderegg et al., 2013), leaving

them weakened when the 2018-2019 drought struck (Buras et al., 2020). The second hotter drought

year 2019 started already with severe soil moisture shortages as the low winter and spring

precipitation in 2018-2019 was not enough to refill soil water reservoirs (UFZ Drought Monitor /

Helmholtz Centre for Environmental Research). The drought legacy effects we found in tree

growth therefore likely resulted from both, physiological and abiotic drought legacies. Next to

drought duration and intensity drought timing may influence tree radial growth (Schwarz et al.,

2020). We observed variable timings of climatic drought onset, with single drought years being

characterized by both spring and summer droughts, 2018 by summer drought (onset in Mai) and

2019 by drought during spring and summer (onset in February) (Fig. S9). Studies examining intra-

annual radial growth at high temporal resolution show that maple, ash and *Q. petraea* continue to

grow until August (if not affected by drought) (Brinkmann et al., 2016; Dietrich, Delzon, Hoch,

Kahmen, & Battipaglia, 2018), which, together with our own observation of strong growth-climate

correlations in spring and summer months (Fig. S4), points at all species being effected by drought

during their growing phase. Nonetheless, drought effects on growth and $\Delta\delta^{13}C$ are likely strongest

if the timing of drought is such that both early and latewood development are affected (Schwarz

et al., 2020). That the drought in 2019 affected the entire growing season while the drought in 2018

did not may therefore – in addition to legacy effects of 2018 – have contributed to the strong stress

responses we report. Our sampling sites cover the whole gradient of groundwater conditions in the

examined floodplain forest but interestingly we found only small effects of groundwater level. The

reasons remain speculative. Differences in distances to the groundwater level may have been too

small to induce strong effects on tree performance or, alternatively, more intense rooting on dry

plots may have compensated for lower water availability (Skiadaresis et al., 2019). We did not

observe a temporal trend in groundwater levels (neither decrease nor increase) over the study

period and decreases in response to the 2018-2019 drought were small (Fig. S10). Temporal

changes in groundwater level are thus unlikely to have had major influences on observed

responses. Finally, trees in floodplain forests take up water from upper (unsaturated) soil horizons

that are fed by precipitation and through capillary rise from the groundwater level (Sánchez-Pérez

et al., 2008; Singer et al., 2014). Under severe drought this capillary rise may have been interrupted

at all distances from the groundwater and the loamy Vega soils of our study site reached moisture

levels close to the permanent wilting point of vegetation for clay soils (Fig. 2a).

The magnitude and timing of drought stress responses were species-specific, which may be related

to differences in species hydraulic traits. Oak and ash feature similar cavitation resistance but

contrasting stomatal control which may explain the stronger drought stress response observed in

ash compared to oak. We report a highly significant downregulation of sap flow with decreasing

soil moisture during the 2018 drought in oak and maple but no downregulation in ash (Fig. 2),

indicating rather water saving and water spending strategies, respectively. Its water saving strategy

may have helped oak to avoid xylem cavitation during peak drought periods, while the water

spending strategy of ash carried a higher cavitation risk (Martínez-Vilalta & Garcia-Forner, 2017;

McDowell et al., 2008), as both species feature similar levels of cavitation resistance (both -2.8

MPa; Choat et al., 2012). Moreover, a water spending strategy necessitates continued water uptake

via roots (McDowell et al., 2008), which may be an especially risky strategy on severely dried out

clay soils. Interestingly, the water saving behavior we report here for oak (*Q. robur*) contrasts with

observations of water spending behavior (no downregulation of sap flow) in *Q. petraea* (Dietrich

et al., 2018; Leuzinger, Zotz, Asshoff, & Körner, 2005), the second 'common' oak species in

Central Europe. This difference may be explained by the higher drought adaptation of *Q. petraea*

compared to *Q. robur* (Nardini & Tyree, 1999), as evidenced also by its higher cavitation resistance

(Ψ_{50} of -3.5 vs -2.8 MPa; Choat et al., 2012).

Oak and maple showed similar $\Delta\delta^{13}C$ responses in all drought years consistent with their similar

stomatal control. In contrast, ash showed a stronger response particularly in 2019. On first sight,

this may come as a surprise as one may expect lower $\Delta\delta^{13}C$ increases (which are related to stomatal

closure) in a water spending compared to water saving species. However, potentially high

hydraulic damages in ash during the severe 2018 drought would necessitate a high mobilization of

NSC reserves for damage repair. As discussed above, this would enrich the reserve pool and tissues

built from it in $\Delta\delta^{13}C$ and could explain the strong $\Delta\delta^{13}C$ increases we observe in the second

consecutive drought year 2019. The overall intermediate drought reaction of maple, which is often

considered drought sensitive (Leuschner et al., 2019), may be related to its higher vulnerability to

cavitation and/or its water saving behavior that may have prevented severe damages to a certain

extent. Moreover, the reaction of maple may also be influenced by its less exposed crown position

(maple trees were rather co-dominant) which can reduce irradiance and water pressure deficits

(Montgomery, Reich, & Palik, 2010). Other traits may have influenced the responses observed but

establishing species-specific differences remains challenging. For instance, ash was reported to

have most root biomass concentrated to shallow soil layers while oak occupied deeper layers in

another riparian hardwood forests (Sánchez-Pérez et al., 2008). However, other studies reported

rather deep rooting in ash and an intruding ability to plastically shift its water uptake to deeper soil

layers (Brinkmann, Eugster, Buchmann, & Kahmen, 2019; Meißner, Köhler, Schwendenmann, &

Hölscher, 2012). Despite compelling progress in functional trait research (Kattge et al., 2020),

harmonized assessments of key hydraulic traits, particularly root and stomatal control related ones,

remain scarce but are urgently needed. Finally, here reported drought effects may be influenced

through the naturally high tree species richness of floodplain forests (Ward et al., 1999), as diverse

tree communities with dissimilar hydraulic traits may outperform species poor communities

through complementarity in water use (Sánchez-Pérez et al., 2008; Schnabel et al., 2019).

The response of forests to the increasing frequency and intensity of droughts (IPCC, 2014) will

affect a variety of ecosystem services and will determine if forests act as carbon sink or source in

the 21st century. Here, we show that the cumulative effect of two consecutive hotter drought years

(2018-2019) induces strong drought-related stress responses in a Central European floodplain

forest. Our retrospective analysis based on tree rings allowed us a robust comparison of the stress

responses observed in 2018-2019 compared to responses in former severe drought years (2003,

2006, 2015) on the same tree individuals. Stress responses in 2019 were far stronger than in any

other examined drought year, indicating that consecutive hotter drought years exert a novel stress

for the examined trees. Comparisons of living and dead trees affected by drought show that radial

growth reductions are widespread before tree mortality and that sudden changes in tree growth

often precede mortality caused by tree hydraulic failure (Cailleret et al., 2017; Obladen et al.,

2021). Nonetheless, linking responses in growth and $\Delta\delta^{13}C$ to tree mortality and carbon ecosystem

fluxes remains challenging (Kannenberg et al., 2019) as the drought effects we report cannot be

directly upscaled to whole forest ecosystems nor be used to predict future mortality risks. Against

this background it is important to consider that we found partly buffered tree stress responses,

presumably because floodplain trees are fed by groundwater in addition to precipitation, and

examined only the most vital tree individuals of the population. Our results thus show a 'best-case

scenario' and more severe tree responses could be expected if entire tree populations or other forest

ecosystems were examined, as indicated by wide-spread tree mortality observed in assessments of

the 2018 drought (Buras et al., 2020; Schuldt et al., 2020) and at our own site (Wirth et al., 2021).

Further, it remains unknown how the here observed responses will affect tree recovery after and

resilience (Lloret et al., 2011; Schwarz et al., 2020) to (future) droughts. The strong drought legacy

effects we observe in the second drought year (2019), the reported persistence of legacy effects for

years (Anderegg et al., 2015) and successive declines in drought resistance under consecutive

drought (Anderegg et al., 2013; Lloret et al., 2011) are worrisome. On the other hand, a species

like oak that combines a high tolerance to drought and flood (Scharnweber, Manthey, & Wilmking,

2013), may remain resilient, underlining its importance for floodplain forests. Consecutive hotter

droughts are projected to become more frequent (Hari et al., 2020). Our ability to forecast forest

responses to climate change hinges on our ability to understand what drives tree responses to this

novel stress.

Acknowledgements

We thank our colleagues of the ‚Lebendige Luppe' (living Luppe river) project for establishing

the plot network that allowed us to draw representative conclusions and Andreas Sickert

(Stadtforst) und Andreas Padberg (Sachsenforst) for the possibility to extract tree increment cores.

The Lebendige Luppe project is funded by the German Federal Agency for Nature Conservation

(BfN) in cooperation with the German Federal Ministry for the Environment, Nature Conservation

and Nuclear Safety (BMU). We further acknowledge the help of the BGC stable isotope laboratory

of the Max Planck Institute for Biogeochemistry in Jena, Germany. F.S. was supported by the

International Research Training Group TreeDì funded by the Deutsche Forschungsgemeinschaft

(DFG, German Research Foundation) – 319936945/GRK2324.

Author contributions

C.W., F.S., R.A.E., A.K., R.R., C.S.-D. and G.S. conceived the idea and developed the concept of

the study. S.P., L.S., A.K., F.S., R.A.E., R.R. and C.S.-D. collected data. F.S., S.P, L.S., R.R. and

G.S. analyzed data. F.S., C.S.-D. and G.S. created figures. F.S. wrote the manuscript with

contributions from S.P. and L.S. All authors discussed the results and contributed substantially to

revisions.

Data availability

Tree growth and δ¹³C data supporting the findings of this study are available on the iDiv Data

repository: https://idata.idiv.de (Note: data will be released publicly upon publication). Climate

data is available at: https://cdc.dwd.de/portal/.

References

Allen, C. D., Breshears, D. D., & McDowell, N. G. (2015). On underestimation of global

vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene.

Ecosphere, *6*(8). https://doi.org/10.1890/ES15-00203.1

Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., . . .

Cobb, N. (2010). A global overview of drought and heat-induced tree mortality reveals

emerging climate change risks for forests. *Forest Ecology and Management*, *259*(4), 660–684.

https://doi.org/10.1016/j.foreco.2009.09.001

Anderegg, W. R. L., Plavcová, L., Anderegg, L. D. L., Hacke, U. G., Berry, J. A., & Field, C. B.

(2013). Drought's legacy: Multiyear hydraulic deterioration underlies widespread aspen forest

die-off and portends increased future risk. *Global Change Biology*, *19*(4), 1188–1196.

https://doi.org/10.1111/gcb.12100

Anderegg, W. R. L., Schwalm, C., Biondi, F., Camarero, J. J., Koch, G., Litvak, M., . . .

Pacala, S. (2015). Pervasive drought legacies in forest ecosystems and their implications for

carbon cycle models. *Science*, *349*(6247), 528–532.

Anderegg, W. R. L., Trugman, A. T., Badgley, G., Anderson, C. M., Bartuska, A., Ciais, P., . . .

Randerson, J. T. (2020). Climate-driven risks to the climate mitigation potential of forests.

Science, *368*(6497). https://doi.org/10.1126/science.aaz7005

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models

Using lme4. *Journal of Statistical Software*, *67*(1). https://doi.org/10.18637/jss.v067.i01

Beguería, S., & Vicente-Serrano, S. M. (2017). SPEI: Calculation of the Standardised

Precipitation-Evapotranspiration Index. [Computer software]. Retrieved from

https://CRAN.R-project.org/package=SPEI

BMU & BfN (2021). *Auenzustandsbericht. Flussauen in Deutschland*. Berlin, Germany.

Brinkmann, N., Eugster, W., Buchmann, N., & Kahmen, A. (2019). Species-specific differences

in water uptake depth of mature temperate trees vary with water availability in the soil. *Plant*

Biology (Stuttgart, Germany), *21*(1), 71–81. https://doi.org/10.1111/plb.12907

Brinkmann, N., Eugster, W., Zweifel, R., Buchmann, N., & Kahmen, A. (2016). Temperate tree

species show identical response in tree water deficit but different sensitivities in sap flow to

summer soil drying. *Tree Physiology*, *36*(12), 1508–1519.

https://doi.org/10.1093/treephys/tpw062

Bunn, A. G. (2008). A dendrochronology program library in R (dplR). *Dendrochronologia*,

26(2), 115–124. https://doi.org/10.1016/j.dendro.2008.01.002

Bunn, A. G., Korpela, M., Biondi, F., Campelo, F., Mérian, P., Qeadan, F., & Zang, C. (2020).

dplR: Dendrochronology Program Library in R. Retrieved from https://CRAN.R-

project.org/package=dplR

Büntgen, U., Urban, O., Krusic, P. J., Rybníček, M., Kolář, T., Kyncl, T., . . . Trnka, M. (2021).

Recent European drought extremes beyond Common Era background variability. *Nature*

Geoscience. Advance online publication. https://doi.org/10.1038/s41561-021-00698-0

Buras, A., Rammig, A., & Zang, C. S. (2020). Quantifying impacts of the 2018 drought on

European ecosystems in comparison to 2003. *Biogeosciences*, *17*(6), 1655–1672.

https://doi.org/10.5194/bg-17-1655-2020

Cailleret, M., Jansen, S., Robert, E. M. R., DeSoto, L., Aakala, T., Antos, J. A., . . . Martínez-

Vilalta, J. (2017). A synthesis of radial growth patterns preceding tree mortality. *Global*

Change Biology, *23*(4), 1675–1690. https://doi.org/10.1111/gcb.13535

Choat, B., Jansen, S., Brodribb, T. J., Cochard, H., Delzon, S., Bhaskar, R., . . . Zanne, A. E.

(2012). Global convergence in the vulnerability of forests to drought. *Nature*, *491*(7426),

752–755. https://doi.org/10.1038/nature11688

Cocozza, C., Paoletti, E., Mrak, T., Zavadlav, S., Levanič, T., Kraigher, H., . . . Hoshika, Y.

(2020). Isotopic and Water Relation Responses to Ozone and Water Stress in Seedlings of

Three Oak Species with Different Adaptation Strategies. *Forests*, *11*(8), 864.

https://doi.org/10.3390/f11080864

Dietrich, L., Delzon, S., Hoch, G., Kahmen, A., & Battipaglia, G. (2018). No role for xylem

embolism or carbohydrate shortage in temperate trees during the severe 2015 drought.

Journal of Ecology, *197*(4), 1142. https://doi.org/10.1111/1365-2745.13051

Farquhar, G. D., Ehleringer, J. R., & Hubick, K. T. (1989). Carbon Isotope Discrimination and

Photosynthesis. *Annual Review of Plant Physiology and Plant Molecular Biology*, *40*(1), 503–

537. https://doi.org/10.1146/annurev.pp.40.060189.002443

Fritts, H. (1976). *Tree Rings and Climate*. London, UK: Academic Press. Retrieved from

http://site.ebrary.com/lib/alltitles/docDetail.action?docID=10685739

Gärtner, H., & Nievergelt, D. (2010). The core-microtome: A new tool for surface preparation on

cores and time series analysis of varying cell parameters. *Dendrochronologia*, *28*(2), 85–92.

https://doi.org/10.1016/j.dendro.2009.09.002

Gazol, A., Camarero, J. J., Sangüesa-Barreda, G., Serra-Maluquer, X., Sánchez-Salguero, R.,

Coll, L., & Casals, P. (2020). Tree Species Are Differently Impacted by Cumulative Drought

Stress and Present Higher Growth Synchrony in Dry Places. *Frontiers in Forests and Global*

Change, *3*. https://doi.org/10.3389/ffgc.2020.573346

Griscom, B. W., Adams, J., Ellis, P. W., Houghton, R. A., Lomax, G., Miteva, D. A., . . .

Fargione, J. (2017). Natural climate solutions. *Proceedings of the National Academy of*

Sciences of the United States of America, *114*(44), 11645–11650.

https://doi.org/10.1073/pnas.1710465114

Grissino-Mayer, H. D. (2001). Evaluating Crossdating Accuracy: A Manual and Tutorial for the

Computer Program COFECHA. *Tree-Ring Research*. Retrieved from

http://hdl.handle.net/10150/251654

Grossiord, C., Granier, A., Ratcliffe, S., Bouriaud, O., Bruelheide, H., Checko, E., . . .

Gessler, A. (2014). Tree diversity does not always improve resistance of forest ecosystems to

drought. *Proceedings of the National Academy of Sciences of the United States of America*,

111(41), 14812–14815. https://doi.org/10.1073/pnas.1411970111

785 Günther-Diringer, D., Berner, K., Koenzen, U., Kurth, A., Modrak, P., Ackermann, W., . . .

786   Heyden, J. (2021). Methodische Grundlagen zum Auenzustandsbericht 2021: Erfassung,

787   Bilanzierung und Bewertung von Flussauen. *BfN-Skripten*, *591*.

788 Gutte, P. (2011). Das Querco-Ulmetum minoris ISSLER 1942, der Stieleichen-Ulmen-

789   Hartholzwald, in der Elster-Luppe-Aue bei Leipzig. *Mauritiana*, *22*, 213–242.

790 Haase, D., & Gläser, J. (2009). Determinants of floodplain forest development illustrated by the

791   example of the floodplain forest in the District of Leipzig. *Forest Ecology and Management*,

792   *258*(5), 887–894. https://doi.org/10.1016/j.foreco.2009.03.025

793 Härdtle, W., Bergmeier, E., Fichtner, A., Heinken, T., Hölzel, N., Remy, D., . . . Dierschke, H.

794   (2020). Pflanzengesellschaft des Jahres 2021: Hartholz-Auenwald (Ficario-Ulmetum).

795   *Tuexenia*, *40*, 373–399.

796 Hari, V., Rakovec, O., Markonis, Y., Hanel, M., & Kumar, R. (2020). Increased future

797   occurrences of the exceptional 2018-2019 Central European drought under global warming.

798   *Scientific Reports*, *10*(1), 12207. https://doi.org/10.1038/s41598-020-68872-9

799 Hartmann, H., & Trumbore, S. (2016). Understanding the roles of nonstructural carbohydrates in

800   forest trees - from what we can measure to what we want to know. *New Phytologist*, *211*(2),

801   386–403. https://doi.org/10.1111/nph.13955

802 Hartmann, H., Ziegler, W., & Trumbore, S. (2013). Lethal drought leads to reduction in

803   nonstructural carbohydrates in N orway spruce tree roots but not in the canopy. *Functional*

804   *Ecology*, *27*(2), 413–427. https://doi.org/10.1111/1365-2435.12046

805 Heklau, H., Jetschke, G., Bruelheide, H., Seidler, G., & Haider, S. (2019). Species-specific

806   responses of wood growth to flooding and climate in floodplain forests in Central Germany.

807   *IForest - Biogeosciences and Forestry*, *12*(3), 226–236. https://doi.org/10.3832/ifor2845-012

Hölscher, D., Koch, O., Korn, S., & Leuschner, C. (2005). Sap flux of five co-occurring tree

species in a temperate broad-leaved forest during seasonal soil drought. *Trees*, *19*(6), 628–

637. https://doi.org/10.1007/s00468-005-0426-3

Field, C. B., Barros, V. R., Dokken, D. J., Mach, K. J., Mastrandrea, M. D., Bilir, T. E., . . .

White, L. L. (Eds.) (2014). *Climate change 2014: Impacts, Adaptation, and Vulnerability.*

Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth

Assessment Report of the Intergovernmental Panel on Climate Change. New York, NY:

Cambridge University Press.

Jucker, T., Grossiord, C., Bonal, D., Bouriaud, O., Gessler, A., & Coomes, D. A. (2017).

Detecting the fingerprint of drought across Europe's forests: do carbon isotope ratios and stem

growth rates tell similar stories? *Forest Ecosystems*, *4*(1), 706.

https://doi.org/10.1186/s40663-017-0111-1

Kannenberg, S. A., Novick, K. A., Alexander, M. R., Maxwell, J. T., Moore, D. J. P.,

Phillips, R. P., & Anderegg, W. R. L. (2019). Linking drought legacy effects across scales:

From leaves to tree rings to ecosystems. *Global Change Biology*, *25*(9), 2978–2992.

https://doi.org/10.1111/gcb.14710

Kattge, J., Bönisch, G., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., . . . Wirth, C. (2020).

Try plant trait database - enhanced coverage and open access. *Global Change Biology*, *26*(1),

119–188. https://doi.org/10.1111/gcb.14904

Klein, T. (2014). The variability of stomatal sensitivity to leaf water potential across tree species

indicates a continuum between isohydric and anisohydric behaviours. *Functional Ecology*,

28(6), 1313–1320. https://doi.org/10.1111/1365-2435.12289

Köcher, P., Gebauer, T., Horna, V., & Leuschner, C. (2009). Leaf water status and stem xylem

flux in relation to soil drought in five temperate broad-leaved tree species with contrasting

water use strategies. *Annals of Forest Science*, *66*(1), 101.

https://doi.org/10.1051/forest/2008076

Kowalska, N., Šigut, L., Stojanović, M., Fischer, M., Kyselova, I., & Pavelka, M. (2020).

Analysis of floodplain forest sensitivity to drought. *Philosophical Transactions of the Royal*

Society of London. Series B, Biological Sciences, *375*(1810), 20190518.

https://doi.org/10.1098/rstb.2019.0518

Kraft, G. (1884). *Beiträge zur Lehre von den Durchforstungen, Schlagstellungen und*

Lichtungshieben. Hannover, Germany: Klindworth's Verlag.

Kröber, W., Zhang, S., Ehmig, M., & Bruelheide, H. (2014). Linking xylem hydraulic

conductivity and vulnerability to the leaf economics spectrum--a cross-species study of 39

evergreen and deciduous broadleaved subtropical tree species. *PLoS ONE*, *9*(11), e109211.

https://doi.org/10.1371/journal.pone.0109211

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in

Linear Mixed Effects Models. *Journal of Statistical Software*, *82*(13).

https://doi.org/10.18637/jss.v082.i13

Lemoine, D., Peltier, J.-P., & Marigo, G. (2001). Comparative studies of the water relations and

the hydraulic characteristics in Fraxinus excelsior, Acer pseudoplatanus and A. opalus trees

under soil water contrasted conditions. *Annals of Forest Science*, *58*(7), 723–731.

https://doi.org/10.1051/forest:2001159

Lenth, R. (2020). emmeans: Estimated Marginal Means, aka Least-Squares Means. [Computer

software]. Retrieved from https://CRAN.R-project.org/package=emmeans

Lenz, H., Straßer, L., Baumann, M., & Baie, U. (2012). Boniturschlüssel zur Einstufung der

Vitalität von Alteschen. *AFZ-DerWald*, *3*, 18–19.

Leuschner, C., & Ellenberg, H. (2017). *Vegetation ecology of Central Europe: Vegetation*

Ecology of Central Europe, Volume I (L. Sutcliffe, Trans.). *Life sciences*. Cham, Switzerland:

Springer.

Leuschner, C., Wedde, P., & Lübbe, T. (2019). The relation between pressure–volume curve

traits and stomatal regulation of water potential in five temperate broadleaf tree species.

Annals of Forest Science, *76*(2). https://doi.org/10.1007/s13595-019-0838-7

Leuzinger, S., Zotz, G., Asshoff, R., & Körner, C. (2005). Responses of deciduous forest trees to

severe drought in Central Europe. *Tree Physiology*, *25*(6), 641–650.

https://doi.org/10.1093/treephys/25.6.641

Lloret, F., Keeling, E. G., & Sala, A. (2011). Components of tree resilience: Effects of

successive low-growth episodes in old ponderosa pine forests. *Oikos*, *120*(12), 1909–1920.

https://doi.org/10.1111/j.1600-0706.2011.19372.x

Lüdecke, D. (2018). ggeffects: Tidy Data Frames of Marginal Effects from Regression Models.

Journal of Open Source Software, *3*(26), 772. https://doi.org/10.21105/joss.00772

Martínez-Vilalta, J., & Garcia-Forner, N. (2017). Water potential regulation, stomatal behaviour

and hydraulic transport under drought: Deconstructing the iso/anisohydric concept. *Plant,*

Cell and Environment, *40*(6), 962–976. https://doi.org/10.1111/pce.12846

McDowell, N., Pockman, W. T., Allen, C. D., Breshears, D. D., Cobb, N., Kolb, T., . . .

Yepez, E. A. (2008). Mechanisms of plant survival and mortality during drought: Why do

some plants survive while others succumb to drought? *New Phytologist*, *178*(4), 719–739.

https://doi.org/10.1111/j.1469-8137.2008.02436.x

McKee, T. B., Doesken, N. J., & Kleist, J. (1993). The relationship of drought frequency and

duration to time scales. In *Proceedings of the 8th Conference on Applied Climatology*.

Meißner, M., Köhler, M., Schwendenmann, L., & Hölscher, D. (2012). Partitioning of soil water

among canopy trees during a soil desiccation period in a temperate mixed forest.

Biogeosciences, *9*(8), 3465–3474. https://doi.org/10.5194/bg-9-3465-2012

Mikac, S., Žmegač, A., Trlin, D., Paulić, V., Oršanić, M., & Anić, I. (2018). Drought-induced

shift in tree response to climate in floodplain forests of Southeastern Europe. *Scientific*

Reports, *8*(1), 16495. https://doi.org/10.1038/s41598-018-34875-w

Montgomery, R. A., Reich, P. B., & Palik, B. J. (2010). Untangling positive and negative biotic

interactions: Views from above and below ground in a forest ecosystem. *Ecology*, *91*(12),

3641–3655. https://doi.org/10.1890/09-1663.1

Nardini, A., & Tyree, M. T. (1999). Root and shoot hydraulic conductance of seven Quercus

species. *Annals of Forest Science*, *56*(5), 371–377. https://doi.org/10.1051/forest:19990502

Obladen, N., Dechering, P., Skiadaresis, G., Tegel, W., Keßler, J., Höllerl, S., . . . Seim, A.

(2021). Tree mortality of European beech and Norway spruce induced by 2018-2019 hot

droughts in central Germany. *Agricultural and Forest Meteorology*, *307*, 108482.

https://doi.org/10.1016/j.agrformet.2021.108482

R Core Team (2020). A language and environment for statistical computing [Computer

software]. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from

https://www.R-project.org/

Richardson, A. D., Carbone, M. S., Keenan, T. F., Czimczik, C. I., Hollinger, D. Y.,

Murakami, P., . . . Xu, X. (2013). Seasonal dynamics and age of stemwood nonstructural

carbohydrates in temperate forest trees. *The New Phytologist*, *197*(3), 850–861.

https://doi.org/10.1111/nph.12042

Richter, R., Reu, B., Wirth, C., Doktor, D., & Vohland, M. (2016). The use of airborne

hyperspectral data for tree species classification in a species-rich Central European forest

area. *International Journal of Applied Earth Observation and Geoinformation*, *52*, 464–474.

https://doi.org/10.1016/j.jag.2016.07.018

Sánchez-Pérez, J. M., Lucot, E., Bariac, T., & Trémolières, M. (2008). Water uptake by trees in a

riparian hardwood forest (Rhine floodplain, France). *Hydrological Processes*, *22*(3), 366–375.

https://doi.org/10.1002/hyp.6604

Scharnweber, T., Manthey, M., & Wilmking, M. (2013). Differential radial growth patterns

between beech (Fagus sylvatica L.) and oak (Quercus robur L.) on periodically waterlogged

soils. *Tree Physiology*, *33*(4), 425–437. https://doi.org/10.1093/treephys/tpt020

Schnabel, F., Schwarz, J. A., Dănescu, A., Fichtner, A., Nock, C. A., Bauhus, J., & Potvin, C.

(2019). Drivers of productivity and its temporal stability in a tropical tree diversity

experiment. *Global Change Biology*, *25*(12), 4257-4272. https://doi.org/10.1111/gcb.14792

Scholz, M., Seele, C., Engelmann, R. A., Hartmann, H., Heinrich, J. Henle, K.,

Herkelrath, A., . . . Riedel, J. (2018). Das Projekt Lebendige Luppe – ein Beitrag zur

Renaturierung der Leipziger Nord-West-Aue. *Auenmagazin*, *14*(14), 14–21.

Schuldt, B., Buras, A., Arend, M., Vitasse, Y., Beierkuhnlein, C., Damm, A., . . . Kahmen, A.

(2020). A first assessment of the impact of the extreme 2018 summer drought on Central

European forests. *Basic and Applied Ecology*, *45*, 86–103.

https://doi.org/10.1016/j.baae.2020.04.003

Schwarz, J., Skiadaresis, G., Kohler, M., Kunz, J., Schnabel, F., Vitali, V., & Bauhus, J. (2020).

Quantifying Growth Responses of Trees to Drought—a Critique of Commonly Used

Resilience Indices and Recommendations for Future Studies. *Current Forestry Reports*, *6*,

185–200. https://doi.org/10.1007/s40725-020-00119-2

Schweingruber, F. H. (1996). *Tree rings and environment: dendroecology*. Berne, Switzerland:

Paul Haupt AG Bern.

Singer, M. B., Sargeant, C. I., Piégay, H., Riquier, J., Wilson, R. J. S., & Evans, C. M. (2014).

Floodplain ecohydrology: Climatic, anthropogenic, and local physical controls on partitioning

of water sources to riparian trees. *Water Resources Research*, *50*(5), 4490–4513.

https://doi.org/10.1002/2014WR015581

Skiadaresis, G., Schwarz, J. A., & Bauhus, J. (2019). Groundwater Extraction in Floodplain

Forests Reduces Radial Growth and Increases Summer Drought Sensitivity of Pedunculate

Oak Trees (Quercus robur L.). *Frontiers in Forests and Global Change*, *2*, 267.

https://doi.org/10.3389/ffgc.2019.00005

Szejner, P., Belmecheri, S., Ehleringer, J. R., & Monson, R. K. (2020). Recent increases in

drought frequency cause observed multi-year drought legacies in the tree rings of semi-arid

forests. *Oecologia*, *192*(1), 241–259. https://doi.org/10.1007/s00442-019-04550-6

Thomsen, S., Reisdorff, C., Gröngröft, A., Jensen, K., & Eschenbach, A. (2020).

"Responsiveness of mature oak trees (Quercus robur L.) to soil water dynamics and

meteorological constraints in urban environments". *Urban Ecosystems*, *23*(1), 173–186.

https://doi.org/10.1007/s11252-019-00908-z

Tockner, K., & Stanford, J. A. (2002). Riverine flood plains: present state and future trends.

Environmental Conservation, *29*(3), 308–330. https://doi.org/10.1017/S037689290200022X

UFZ Drought Monitor / Helmholtz Centre for Environmental Research. Retrieved from

https://www.ufz.de/index.php?en=37937

Van Loon, A. F., Stahl, K., Di Baldassarre, G., Clark, J., Rangecroft, S., Wanders, N., . . . van

Lanen, H. A. J. (2016). Drought in a human-modified world: reframing drought definitions,

understanding, and analysis approaches. *Hydrology and Earth System Sciences*, *20*(9), 3631–

3650. https://doi.org/10.5194/hess-20-3631-2016

Vicente-Serrano, S. M., Beguería, S., & López-Moreno, J. I. (2010). A Multiscalar Drought

Index Sensitive to Global Warming: The Standardized Precipitation Evapotranspiration Index.

Journal of Climate, *23*(7), 1696–1718. https://doi.org/10.1175/2009JCLI2909.1

Ward, J. V., Tockner, K., & Schiemer, F. (1999). Biodiversity of floodplain river ecosystems:

ecotones and connectivity1. *Regulated Rivers: Research & Management*, *15*(1-3), 125–139.

https://doi.org/10.1002/(SICI)1099-1646(199901/06)15:1/3<125::AID-RRR523>3.0.CO;2-E

Weil, R. R., & Brady, N. C. (2017). *Nature and properties of soils* (15th edition): Pearson.

Wirth, C., Engelmann, R. A., Haack, N., Hartmann, H., Richter, R., Schnabel, F., . . . Seele-

Dilbat, C. (2021). Naturschutz und Klimawandel im Leipziger Auwald: Ein

Biodiversitätshotspot an der Belastungsgrenze. *Biologie in Unserer Zeit*, *51*(1), 55–65.

https://doi.org/10.11576/biuz-4107

Zang, C., & Biondi, F. (2015). treeclim: an R package for the numerical calibration of proxy-

climate relationships. *Ecography*, *38*(4), 431–436. https://doi.org/10.1111/ecog.01335

962