TITLE: Fine-tuned repression of Drp1 driven mitochondrial fission primes a

'stem/progenitor-like state' to accelerate neoplastic transformation

AUTHORS: Spurlock B[1], Parker D[1], Basu MK[2], Hjelmeland A[3], G Sajina[3], Liu S[1], Siegal

GP[2], Gunter A[1], Moran A[1] and Mitra K[1]*

* corresponding author (kasturi@uab.edu)

AFFILIATION:

1. Department of Genetics, University of Alabama at Birmingham,

10   Birmingham, Alabama 35294

11   **2.** Department of Pathology, University of Alabama at Birmingham,

12   Birmingham, Alabama 35294

13   **3.** Department of Cell Development and Integrative Biology, University of Alabama

14   at Birmingham,

4. Department of Biology, University of Alabama at Birmingham.

Running title:

Mitochondria based priming of 'stem/progenitor-like state' towards driving

transformation

Keywords:

Drp1, Mitochondrial fission, stem/progenitor-like state, neoplastic transformation,

keratinocytes, scRNA-seq, microscopy

ABSTRACT:

The opposing processes of mitochondrial fission and fusion are emerging as crucial regulators of stemness. Gene knockout of the master regulator of mitochondrial fission, Drp1, prevents neoplastic transformation. However, stem/progenitor cells maintaining repressed mitochondrial fission are primed for self-renewal and proliferation. Here, we demonstrate that only fine-tuned repression of Drp1 establishes a 'stem/progenitor-like state' towards supporting carcinogen driven neoplastic transformation of keratinocytes, while more complete Drp1 repression prevents it. Only fine-tuned Drp1 repression maintains small networks of fused mitochondria to sustain a unique gene-expression profile with elevated stem/progenitor cell functional markers (Krt15, Sox2 etc) and their regulators (Cyclin E). Cells with such a mitochondria-primed state are slow cycling, susceptible to transformation, and when enriched by mild carcinogen exposure sustains elevated self-renewal/proliferation to form less differentiated tumors. Therefore, our data for the first time highlights a 'goldilocks' level of Drp1 repression that supports stem/progenitor state dependent neoplastic transformation.

INTRODUCTION

Stem cells exhibit plasticity as the state of stemness is modulated by various intrinsic and extrinsic factors (Magee, Piskounova et al., 2012)(Folmes, Dzeja et al., 2012). Involvement of mitochondria in regulation of the stem cell state is complex (Lisowski, Kannan et al., 2018)(Khacho & Slack, 2017, Zhang, Menzies et al., 2018). The master regulators of mitochondrial fission and fusion processes are emerging as crucial regulators of both embryonic and adult stem cells (Khacho & Slack, 2017, Lisowski et al., 2018)(Spurlock, Tullet et al., 2020). The Dynamin Related Protein 1 (Drp1) is the master regulator of mitochondrial fission that breaks larger mitochondria into smaller elements (Friedman & Nunnari, 2014, Kageyama, Zhang et al., 2011). The effects of Drp1 driven mitochondrial fission is opposed by fusion between mitochondria driven by the Mitofusins and Opa1(Chen & Chan, 2017)(Schrepfer & Scorrano, 2016). Therefore, Drp1 repression allows unopposed mitochondrial fusion to sustain a hyperfused mitochondrial state. Enhanced mitochondrial fusion sustains stemness of certain adult stem cells of neural, germ line and mammary lineages (Khacho, Clark et al., 2016, Senos Demarco, Uyemura et al., 2019, Wu, Chen et al., 2019). Particularly, repression of Drp1 supports stemness and repression of Mitofusin or Opa1 inactivates stemness in adult mouse neural lineage (Iwata, Casimir et al., 2020, Khacho & Slack, 2017). The other extreme state, i.e. unopposed mitochondrial fission, is critical for achieving pluripotency of stem cells during reprogramming (Prieto, Leon et al., 2016). However, sustained mitochondrial fission reduces pluripotency of stem cells (Zhong, Cui et al.,

2019). Thus, it can be speculated that a balance of timely fission and fusion of

mitochondria may be critical for maintaining stem cell states.

In various tumors, the bulk tumor cell populations are maintained by the adult

neoplastic stem cells (also called tumor initiating cells) as they self-renew, proliferate

and differentiate (Magee, Piskounova et al., 2012). Drp1 has been linked to tumor

formation in various cancer types (Nagdas, Kashatus et al., 2019, Serasinghe, Wieder

et al., 2015, Tanwar, Parker et al., 2016, Tsuyoshi, Orisaka et al., 2020, Xie, Wu et al.,

2015). Drp1 activation sustains neoplastic stem cells at least in the astrocytic lineage

(Xie et al., 2015), and genetic ablation of Drp1 prevents neoplastic transformation

(Serasinghe et al., 2015). In contrast, an elevated mitochondrial fusion state, sustained

by mitofusin, can drive immortalization of neoplastic stem cells to support tumorigenesis

in a Drosophila model (Bonnay, Veloso et al., 2020). We found that select neoplastic

ovarian epithelial stem cells (marked by Aldh) have lower levels of mitochondrial fission

and can convert to a state of elevated mitochondrial fission when self-renewal and

proliferation is activated. Importantly, the cells with minimum mitochondrial fission are

found within a neoplastic stem cell sub-population that has >10 folds self-renewal and

proliferation ability compared to the other neoplastic stem cell subpopulation (Spurlock

et al., 2019)(Spurlock et al., 2021b). Thus, we proposed, that a repressed mitochondrial

fission state may prime neoplastic stem cells towards maximizing their potential of self-

renewal and proliferation. Similar priming was demonstrated in a normal hematopoietic

stem cell subpopulation that maintains Drp1 repression (Liang, Arif et al., 2020). Here,

we investigated the causative role of Drp1 repression in mitochondria driven priming of

a stem/progenitor-like like state in skin cells, given skin stem cells serve as an excellent

model for adult stem cells (Fuchs, 2016). We established a transformed skin

keratinocyte model that sustains abundant self-renewing/proliferating cells with

attenuated Drp1 activity. Using genetic perturbation strategies and single cell analyses

of the parental and the transformed keratinocyte model we demonstrate that only fine-

tuned repression of Drp1 establishes a 'stem/progenitor-like state' towards supporting

neoplastic transformation, whereas more complete Drp1 repression prevents it. Our

data for the first time highlights a 'goldilocks' level of Drp1 repression that supports

stem/progenitor-like state dependent neoplastic transformation.

RESULTS AND DISCUSSION

Establishment of a mild carcinogen transformed keratinocyte model that

maintains abundant self-renewing/proliferating cells and attenuated Drp1 activity.

Chemical carcinogens like 2,3,7,8-tetrachlorodibenzodioxin (TCDD) is widely

used for transforming skin tissue and the keratinocyte HaCaT cells that models key

aspects of skin regeneration (Wincent, Bengtsson et al., 2012)(Hao, Lee et al.,

2012)(Ray & Swanson, 2004)(Der Vartanian, Quetin et al., 2019)(Boelsma, Verhoeven

et al., 1999, Jung, Jung et al., 2016, Schoop, Mirancea et al., 1999). The HaCaT cells

carry non-functional p53 (St John, Sauter et al., 2000), which provides the appropriate

cellular context given Drp1 inactivation increases cell proliferation in the absence of

active p53 (Mitra, Rikhy et al., 2012, Mitra, Wunder et al., 2009). We used both mild and

strong dose of TCDD to uncover any possibility of mitochondria based priming of

stemness during TCDD driven neoplastic transformation. TCDD driven carcinogenesis

is caused by activation and upregulation of the aryl hydrocarbon receptor (AHR)

(Mulero-Navarro & Fernandez-Salguero, 2016) (Leclerc, Staats Pires et al., 2021).

Exposure to milder (T-1nM) and stronger (T-10nM) doses of TCDD causes a

comparable increase in cell proliferation of HaCaT cells in a standard transformation

protocol (Fig. 1A). The stable transformed colonies of TCDD-1nM (TF-1) and TCDD-

10nM (TF-10) maintain comparable upregulation of AHR, even in the absence of TCDD

(Fig. 1B). Unlike the parental HaCaT cells (Parental), both transformed cells are able to

form subcutaneous xenograft tumors, confirming their transformation status

(Supplementary. Fig. 1A). Interestingly, pathological evaluation of H&E stained tumor

sections revealed that the TF-1 cells gave rise to malignant tumors harboring less

differentiated (primitive) squamous cells with large nuclei (blue) and little visible

cytoplasm (pink) (Fig. 1C, left). Whereas the TF-10 cells formed tumors harboring

differentiated stratified squamous epithelium with cells having expansion of cytoplasm

(Fig. 1C, right); the clear gaps represent artifacts secondary to cryo-sectioning.

Notably, among the major molecules regulating mitochondrial fission and fusion

(Drp1, Mfn1, Mfn2, Opa1), Drp1 exhibited the most prominent differences (Fig. 1D).

Particularly, the TF1 population has markedly lower levels of the cell cycle regulated

activating phosphorylation of Drp1 (pDrp1-S616), while both the transformed

populations have higher levels of Drp1 protein compared to the Parental line (Fig. 1D).

The TF-1 population also has lower levels of other tested mitochondrial markers

(Tom20, Cytochrome C and Fis1), compared to the TF-10 population (Supplementary

Fig. 1B). Next, to assess the steady state contribution of mitochondrial fission and

fusion on mitochondrial shape we quantified mitochondrial [Fission] and [Fusion5]

metrics in individual live cells stably expressing the fluorescent mitochondrial reporter,

mitoPSmO (Spurlock et al., 2019)(Spurlock et al., 2021b) (Fig. 1E). In comparison to the

TF-1 and Parental populations, the TF-10 population exhibits reduced levels of the

[Fusion5] metric and slightly elevated level of the [Fission] metric (Fig. 1F). Notably,

although the TF-1 population has higher median [Fusion5], lower range of [Fusion5] is

also detected in this population (Fig. 1F). Since the [Fusion5] metric reflects

mitochondrial length in each cell (Spurlock et al., 2019), this data suggests a wider

range of mitochondrial length in the TF-1 population, which could be potentially brought

about by regulating Drp1 driven mitochondrial fission (Fig. 1D).

The TF-1 HaCaT population has significantly higher *in vitro* cell proliferation rate,

compared to the Parental and the TF-10 population (Fig 1G). Furthermore, the TF-1

population form markedly larger spheroids in conditions that support self-renewal and

proliferation, in comparison to those formed by the Parental or the TF-10 population

(Fig. 1H). More importantly, ELDA statistics applied on spheroid formation assay for

determination of *in vitro* stem cell frequency (Hu & Smyth, 2009) demonstrated that the

TF-1 population has double the abundance of self-renewing/proliferating cells than the

TF-10 population, while their abundance is one order higher in both transformed

populations compared to the Parental (Fig. 1I).

The above data confirms the derivation of a transformed Drp1-attenuated-TF-1

keratinocyte model that maintains greater self-renewal/proliferative ability and forms

less differentiated tumors compared to those transformed by a 10-fold higher TCDD

dose. Thus, we conclude that the Drp1-attenuated-TF-1 keratinocyte population is

enriched in neoplastic stem/progenitor cells.

Transformed Drp1-attenuated-TF-1 keratinocyte population maintains an

expanded slow-cycling sub-population with elevated stem/progenitor cell

markers

To define the neoplastic stem cells of our newly derived transformed Drp1-

attenuated-TF-1 population, we performed scRNA-seq (Luecken & Theis, 2019). Cell

clustering in a UMAP plot shows 6 clusters within Parental, TF-1 and TF-10 populations

(Fig. 2A, Clusters 0-5). While Cluster 4 is reduced in both transformed populations in

comparison to the Parental, the TF-1 population exhibits a marked expansion of Cluster

3 and reduction of Cluster 5 (Fig. 2A, Supplementary. Fig.2A, left panel). These results

also hold true with the lowest cluster resolution (Supplementary. Fig. 2B. arrows). We

identified the top most candidate genes to mark each cluster (color coded arrows in

Supplementary Fig. 2C, Supplementary. Data Table 1). Indeed, the neoplastic

stem/progenitor cell enriched Drp1-att TF-1 population shows >3-fold upregulation of

the epidermal stem cell marker Krt15 (Giroux, Lento et al., 2017, Gonzales & Fuchs,

2017, Purba, Haslam et al., 2014) and of the neoplastic stem cell markers for skin

carcinomas, SOX2 and SOX4 (Boumahdi, Driessens et al., 2014, Foronda, Martinez et

al., 2014) (Fig. 2B, arrow). Since the markers increase across all TF-1 clusters, with

maximum levels in Cluster 3, we conclude that the TF-1 population exhibits upregulation

of the stem/progenitor markers independent of their cluster distribution. On the other

hand, the other stem/progenitor cell markers, like Krt5, Krt14, Krt19 (Gonzales & Fuchs,

2017)(Karantza, 2011) are also enriched by ~2-fold in Cluster 3, but comparably in all

the three cell populations, as exemplified by Krt5 (Fig. 2B, arrow, Supplementary Data

Table 1). Notably, among the 3 cell populations, TF-1 harbors markedly fewer cells in

Cluster 3 with elevation of the differentiating cell marker Krt13 (Karantza, 2011). Cluster

5, reduced in TF-1 population, is marked by 1.5 to >2- fold reduced expression of 12 of

the 13mtDNA genes in all the 3 cell populations (Supplementary Fig. 2D, arrow). Cluster

4, reduced in both transformed populations, is marked by >8-fold upregulation of OASL,

a gene involved in the interferon pathway, in all the three cell populations

(Supplementary Fig. 2E).

Since flow cytometry on PI stained cells revealed modest differences in cell cycle

distribution between the Parental and transformed populations (Supplementary. Fig. 2A,

right panel), we asked if the scRNA-seq derived clusters represent distinct cell cycle

phases. We performed the standard gene set enrichment analyses (GSEA) of the

overall cluster marker profiles (Subramanian, Tamayo et al., 2005), where some

statistically significant REACTOME pathways (q value <0.01) were renamed based on

their leading-edge genes (Supplementary. Data Table 2). GSEA showed upregulation of

genes in "Mitosis and Relevant Cytoskeletal Proteins" in Cluster 2 and that of genes in

"DNA repair / replication and Histones" in Cluster 0 (Fig. 2C); Cluster 1 has

downregulation of "Histones and Ubiquitin related" genes. Indeed, the mitotic Cyclin B

and its partner kinase CDK1, as well as Cyclin A are markedly higher in Cluster 2-G2-M,

while PCNA, which peaks early in S phase (Maga & Hubscher, 2003, Zerjatke, Gak et

al., 2017) is highest in Cluster 0-S-Histone-hi in all 3 cell populations (Fig. 2D); the S

phase cyclins E1/E2 and their partner CDK2 were not detected in our sc-RNAseq data

set. Surprisingly, Krt15 marked Cluster 3 has dramatic downregulation of genes in

"Overall Cell Cycle" as well as the above categories representing distinct cell cycle

phases (Fig. 2C), suggesting their cell cycling is slow. Cluster 3-SloCycl-Krt15-hi

maintains extreme low levels of major cyclins in all three cell populations, most

prominently in TF-1 (Fig. 2D). PCNA, which is dramatically reduced in quiescent cells

(Maga & Hubscher, 2003, Zerjatke, Gak et al., 2017), is markedly reduced in the

SloCycl-Krt15-hi cluster (and others) only in the TF-1 population (Fig. 2D). SloCycl-

Krt15-hi has upregulation of genes involved in "Keratin" and "Signaling" pathways (Fig.

2C). Notably, the Notch pathway target, Hes1, which actively maintains quiescence

(Sang, Coller et al., 2008)(Moriyama, Durham et al., 2008), is upregulated across the

clusters in the TF-1 population (Supplementary Fig. 2E). These data strongly suggest

that the enrichment of the SloCycl-Krt15-hi cluster in the TF-1 population happens due

to active maintenance of quiescence, rendering them slow cycling. Cluster 4 does not

relate to cell cycle and shows upregulation of genes related to "Cytokine" (interferons)

pathway (Fig. 2C), many of which are suppressed by AHR activation (Di Meglio, Duarte

et al., 2014) (Supplementary. Data Table 1). This finding is consistent with our findings

in Fig. 1B and Fig. 2A. The downregulated mtDNA genes in Cluster 5 were identified as

"TCA cycle" and "Complex 1 biogenesis" in our overrepresentation analyses (* in Fig.

2C), but not in GSEA. Towards understanding the entry into the slow cycling state of the

Krt15hi-Cluster 3, we performed trajectory analyses using Slingshot algorithm that

identifies clusters related to each other based on their gene expression (Street *et al*,

2018). The output trajectory was the following: "G2-M" to "S-Histones-hi" to "Cytokine"

to "Histones/Ub-lo" bifurcating into "SloCycl-Krt15-hi" or "mtDNA-lo" (Fig. 2C, open

arrows, Supplementary. Fig. 2F). This implies that the cells in the TF-1 population may

preferentially reside in the SloCycl-Krt15-hi cluster and not in the mtDNA-lo cluster, thus

expanding the former and reducing the latter (Fig. 2A).

To confirm the cell cycle pathway results from GSEA, we quantified cell cycle

scores and distribution in each cluster in cell population using a Seurat algorithm that

employs several validated predictive cell cycle genes (Tirosh *et al*, 2016). This showed

that the TF-1 population has lower S score and comparable G2-M score with the

Parental and the TF-10 populations (Fig. 2E). Indeed, Cluster 2 is abundant in G2-M

cells, Cluster 0 (and 1) in S and Krt15 marked Cluster 3 is in G0/G1 (expected for

quiescent/slow-cycling cells) (Fig. 2F,G). This data taken with the lower levels of

Histone transcripts in Cluster 1 than in Cluster 0 (* in Fig. Supplementary 2C) suggest

that Cluster 0 may be in early S phase while Cluster 1 in late S phase. We noted that

the expanded 3-SloCycl-Krt15-hi cluster in the TF-1 population has significantly more

cells in G0/G1 and less cells in S than the other two populations (Fig. 2F,G, arrows),

which is opposite to the 5-mtDNA-lo cluster that is reduced in the TF-1 population (Fig.

2F,G, arrows). This G1 to S reciprocity between Cluster 3 and 5 (quantified from the

distributions) is >5-fold in the TF-1 population with respect to the Parental and TF-10

population (Fig. 2G, inset). This suggests that altered cell cycle regulation in the TF-1

population may allow them to preferentially reside in the 3-SloCycl-Krt15-hi cluster and

not in the 5-mtDNA-lo cluster. Such a regulation may involve attenuation of CDK1

driven Drp1(S616) phosphorylation (Fig. 1D) due to marked reduction of CyclinB/CDK1

particularly in the TF-1 population in late S phase (Fig. 2D), where decision for

expansion of the slow cycling Krt15-hi cluster is likely made (Fig. 2C, Suppl Fig. 2F).

Thus, our scRNA-seq characterization reveals that our newly derived

transformed Drp1-attenuated-TF-1 keratinocyte population, enriched with self-renewing

cells, maintain an expanded sub-population of slow cycling cells expressing high levels

of stem/progenitor markers (lineage specific: Krt15; general: Sox2 and Sox4), likely due

to altered cell cycle regulation. Our data is consistent with the Krt15+ epidermal stem

cells, which generate various epithelial cell types or contribute to epidermal carcinoma,

being primarily slow cycling (Giroux et al., 2017, Gonzales & Fuchs, 2017, Purba et al.,

2014)(Morris, Liu et al., 2004). Moreover, the higher *in vitro* cell proliferation rate of the

Drp1-attenuated-TF-1 keratinocytes (Fig. 1G) could result from the higher clonogenic

capacity of the Krt15-hi cells (Morris et al., 2004)(Seykora & Cotsarelis, 2011).

As opposed to more complete Drp1 repression, fine-tuned Drp1 repression

sustains a [CyclinE^{hi}-Sox2^{hi}-Krt15^{hi}] state and accelerates carcinogen driven

keratinocyte transformation

Repression of Drp1 driven mitochondrial fission elevates stem/progenitor cell

markers in mouse embryonic fibroblasts (MEFs), and in cells of neural and ovarian

epithelial lineage (Iwata et al., 2020, Parker et al., 2015). In the epidermal lineage, we

observed marked increase in in Sox2 protein levels (but no other embryonic stem cell

markers) in the Drp1 attenuated TF-1 population in comparison to the Parental and TF-

10 keratinocyte population (Fig. 3A, Supplementary Fig. 3A), corroborating the scRNA-

seq data (Fig. 2B). Krt15 protein levels are elevated in both transformed populations in

comparison to the Parental (Fig. 3A), unlike the cluster specific differences in transcript

levels (Fig. 2B). Immunostaining based co-expression analyses of Sox2 and Krt15 in

individual cells revealed that only a subpopulation of Drp1-attenuated-TF-1 population

has elevation of both Krt15 and Sox2 (Fig. 3B, Supplementary. Fig. 3B). Similar

analyses showed that the highest levels of Sox2 in the TF-1 population is sustained in

cells with an intermediate level of Drp1 protein levels (Fig. 3C) and lower pDrp1(S616)

levels (indicative of lower Drp1 activity) (Fig. 3D). Therefore, we investigated whether

repression of Drp1 in the Parental cells may cause the elevation of Sox2 and/or Krt15

observed in the TF-1 population derived from the Parental cells. Therefore, we knocked

down Drp1 expression in Parental keratinocytes with two validated Drp1 shRNAs

(Parker et al., 2015, Tanwar et al., 2016), one weaker (W) and the other 5 folds stronger

(S) almost mimicking Drp1 genetic ablation (Fig. 3E). Surprisingly, while weaker Drp1

knockdown elevates Krt15 levels, the stronger Drp1 knockdown fails to do so (Fig. 3E).

On the other hand, both weak and strong Drp1 knockdown caused marked elevation of

Sox2 in a graded fashion (Fig. 3E), consistent with Drp1 ablated MEFs (Parker et al.,

2015) and neural stem cells (Iwata et al., 2020). Thus, unlike the stronger Drp1

knockdown, the weaker Drp1 knockdown maintains detectable Drp1 protein, and

uniquely sustains a Sox2-hi/Krt15-hi sub-population characteristic of the naturally Drp1

attenuated TF-1 population enriched in neoplastic stem cells.

Cell cycle status is a crucial determinant of stemness (Orford & Scadden, 2008,

Otsuki & Brand, 2018, Pauklin & Vallier, 2013). Sox2 expression is under the influence

of various G1 cyclins (Liu, Michowski et al., 2019). Cyclin E, which is indispensable for

S phase entry from quiescence and also for neoplastic transformation (Geng, Yu et al.,

2003, Siu, Rosner et al., 2012), is critical for sustaining Sox2 expression (Liu et al.,

2019). Interestingly, among the major cyclins, Cyclin E1 levels are particularly higher in

the Sox2 enriched Drp1 attenuated TF-1 population, while Cyclin D is particularly higher

in the TF-10 population, when compared to the Parental, (corroborating scRNA-seq

data, Fig. 2D) (Fig. 3F). Moreover, Cyclin E1 is preferentially accumulated in the

nucleus in the TF-1 population, which is required to impact gene expression of Sox2

(and others) (Fig. 3G,H). The level of pCyclinE(T-62) reflects the level of active Cyclin E

that drives transcription and remains susceptible to degradation (Siu et al., 2012).

Immunostaining based co-expression analyses showed that in the TF-1 population,

elevated Sox2 is sustained in cells with lower pCyclinE(T-62) than in the Parental

population that has overall lower Sox2 (Fig. 3I). Moreover, cycloheximide chase assay,

to assess degradation rate of proteins, revealed no Cyclin E degradation in TF-1 cells in

comparison to marked Cyckin E degradation in the Parental cells; both populations

maintain Tom20 (mitochondrial marker) within the 2 hour assay time frame

(Supplementary. Fig. 3C). Thus, our data is consistent with elevated Sox2 in the TF-1

population being sustained by elevated Cyclin E levels achieved by attenuation of its

degradation kinetics. Given mitochondrial regulation of Cyclin E regulates Cyclin E

degradation (Mandal, Freije et al., 2010, Parker et al., 2015), we tested if repression of

Drp1 driven mitochondrial fission in the Parental cells can elevate Cyclin E levels.

Indeed, Cyclin E is dramatically accumulated in the nucleus in a graded manner with

Drp1 knockdown efficacy (Fig. 3J,K), similar to Sox2 (Fig. 3E), suggesting Drp1

repression elevates nuclear Cyclin E and Sox2 levels similar to the naturally Drp1

attenuated TF-1 population.

Given, Sox2 and Cyclin E are important for neoplastic transformation in various

cancer types (Hwang & Clurman, 2005, Schaefer & Lengerke, 2020, Teixeira & Reed,

2017), and Krt15+ cells give rise to skin carcinoma (Li, Park et al., 2013, Seykora &

Cotsarelis, 2011), we asked if repression of Drp1 can modulate the process of

carcinogen driven neoplastic transformation of keratinocytes. Thus, we exposed control

and Drp1 knockdown Parental keratinocytes to 1 and 10nM TCDD following standard

TCDD driven transformation protocol. Interestingly, the weaker Drp1 knockdown, which

maintains Krt15-hi/Sox2-hi status and elevated nuclear Cyclin E, got transformed by

10nM TCDD even earlier than the control (Fig. 3L); 1 nM TCDD did not cause any

transformation at this time point. This happened in spite of their lower proliferation rate

than the control in the earlier time point. On the other hand, the stronger Drp1

knockdown, which also maintained slower cell proliferation rate and a Krt15-lo/Sox2-hi

status with elevated nuclear Cyclin E, is not transformed in the same time window (Fig.

3L).

We conclude that the neoplastic stem/progenitor cell enriched Drp1-attenuated

TF-1 population maintains a [CyclinE^{hi}-Sox2^{hi}-Krt15^{hi}] sub-population. Fine-tuned Drp1

repression in the Parental cells primes the 'stem/progenitor-like [CyclinE^{hi}-Sox2^{hi}-

Krt15^{hi}] state', making them susceptible to neoplastic transformation, which is

suppressed by more complete Drp1 repression. It is tempting to speculate that inability

to sustain of this primed state may prevent neoplastic transformation in Drp1 ablated

MEFs (Serasinghe et al., 2015).

Fine-tuned Drp1 repression maintains smaller fused mitochondria, and sustains transcriptomic profile of the neoplastic stem cell enriched TF-1 population, which is suppressed by more complete Drp1 repression

Towards understanding how the 'stem/progenitor-like state' is primed by fine-

tuned (incomplete) Drp1 repression, we performed detailed comparison of mitochondrial

shape and over all gene expression between the Parental keratinocytes expressing

control, weak or strong Drp1 shRNAs. Quantification of mitochondrial fission and fusion

metrics contributing to mitochondrial shape in live cells confirmed that the stronger

(more complete) Drp1 knockdown maintains ~77% cells in the uniformly hyperfused

state i.e. [Fission] metric <0.2 and [Fusion5] metric >80 (Spurlock et al., 2019), while the

control population maintains ~55% cells in this state (Fig. 4A, blue box, Supplementary.

Fig. 3D). Interestingly, the weaker (incomplete) Drp1 knockdown maintains only ~40%

cells in the uniformly hyperfused state (Fig. 4A, blue box), while enriching a sub-

population of cells (~28%) in the [Fission] range of 0.2-0.4 and [Fusion5] range of 40-80

(Fig. 4A, red box), reflecting the mitochondrial fragments in the midst of extensive

tubular mitochondria (Supplementary Fig. 3D). We also found that the cells in the

enriched sub-population (Fission: 0.2-0.4 and Fusion5: 40-80) have lower mitochondrial

number and total mitochondrial length (Fig. 4B), signifying decrease in mitochondrial

content. However, cells with hyperfused mitochondria in the weaker Drp1 knockdown

population (Fission<0.2; Fusion5>80) have distinctly higher mitochondrial number but

not mitochondrial length (Fig. 4B). To confirm that such an increase in mitochondrial

number is due to a greater number of smaller networks of fused mitochondria, we

quantified the percent length of the longest mitochondrial elements (Fusion1). Indeed,

less than 7% of cells have >80% of the mitochondrial length as hyperfused in the

weaker Drp1 knockdown, compared to ~40% in the stronger Drp1 knockdown (Fig. 4C,

top). More importantly, the enriched population of cells in the weaker Drp1 knockdown

with fission range of 0.2-0.4 have only up to 40% of the mitochondrial lengths in fused

networks (Fig. 4C, bottom). We detected cells with similar mitochondrial fission and

fusion features particularly in (FACS sorted) mitochondria primed ovarian neoplastic

stem cell population that has 10-fold higher ability of self-renewal and proliferation

(Spurlock et al., 2021b). We conclude that the remnant Drp1 driven mitochondrial

fission after fine-tuned Drp1 repression enriches a subpopulation of cells maintaining

smaller fused mitochondria (by preventing their hyperfusion) with reduced mitochondrial

content and also distinct mitochondrial puncta, towards sustaining a [CyclinE^{hi}-Sox2^{hi}-

Krt15^{hi}] state. The functional significance of such previously unappreciated distinctions

in mitochondrial shape may involve Drp1 driven regulation of mitophagy of damaged

mitochondria (Twig & Shirihai, 2011)

To probe the contribution of gene expression of any particular cell population in

sustaining a stem/progenitor-like state after fine-tuned Drp1 repression, we performed

scRNA-seq of the control and Drp1 knockdown Parental populations. Analyses of the

overall gene expression (within the limits of gene coverage) confirmed that the weaker

Drp1 knockdown, which maintains higher Drp1 transcript levels, modulates a smaller

gene set to a smaller degree compared to the stronger Drp1 knockdown that maintains

lower Drp1 transcript levels (Supplementary. Fig. 4A, B). However, graded Drp1

knockdown has both 'similar' (graded upregulation or downregulation) and 'opposite'

effects (upregulation in one and downregulation in the other) (Supplementary Fig. 4C).

The opposite effect is dominated by Krt15, Krt13, Krt4, Krt5 transcripts being

upregulated in the weaker Drp1 knockdown and downregulated in the stronger Drp1

knockdown (color coded * in Fig. 4D, Supplementary Fig. 4C), corroborating our

findings on Krt15 protein levels (Fig. 3E). On the other hand, genes for various

mitochondrial proteins, involved in mitochondrial energetics, redox or biogenesis, were

upregulated particularly in the stronger Drp1 knockdown (Fig. 4E upper panel,

Supplementary Fig. 4C, bold). This is consistent with our observation of decreased

mitochondrial content in the weaker Drp1 knockdown (Fig. 4B). Synthesis of the

mitochondrial proteins can be potentially sustained by elevated ribosomal genes and

genes involved in protein synthesis in the stronger Drp1 knockdown (Fig. 4E, lower

panel). This finding is consistent with observations linking Drp1 repression to ribosomal

genes and/or protein synthesis (Tanwar et al., 2016)(Zhao Q et al., 2021)(Favaro,

Romanello et al., 2019). We speculate that repression of Drp1 beyond an optimal level

may initiate compensatory changes through the retrograde pathway to trigger

mitochondrial biogenesis (Ryan & Hoogenraad, 2007).

Cell clustering in a UMAP plot (Supplementary. Fig. 4D, Supplementary Data

Table 3) revealed remarkable correspondence to the conserved cluster markers of the

Parental/transformed data set (Fig.2, Supplementary. Fig. 4D, E), allowing assignment

of cluster identities of G2-M, S-early, S-late, mtDNA-lo, SloCycl-Krt15-hi and Cytokine to

the control and Drp1 knockdown data set. Strikingly, the weaker Drp1-knockdown

population exhibits expansion of the SloCycl-Krt15-hi cluster and reduction of the

Cytokine and mtDNA-lo clusters observed in the stem cell enriched Drp1-attenuated-

TF-1 population, albeit to a lesser degree (Fig. 4F). This similarity in cluster organization

suggests that fine-tuned Drp1 repression in Parental population establishes a

stem/progenitor cell-like gene expression profile to poise them for accelerated

neoplastic transformation (Fig. 3L). On the other hand, the stronger Drp1 knockdown

markedly reduced the abundance of the SloCycl-Krt15-hi cluster and exhibits overall

opposite trend of cluster profile compared to the weaker Drp1 knockdown and TF-1

population (Fig. 4F). Krt15 and various other marker Keratin genes are dramatically

suppressed across the clusters in the stronger Drp1 knockdown and over all elevated in

the weaker Drp1 knockdown (Supplementary Fig. 4F), confirming and expanding our

findings on Krt15 protein levels (Fig. 3E). Moreover, analyses of cell cycle distribution

(as in Fig. 2) revealed higher G2-M score, reduced S score and perturbed G1 to S

reciprocity between SloCycl-Krt15-hi and the mtDNA-lo clusters particularly in the

stronger Drp1 knockdown (Fig. 4G, Supplementary. Fig. 4G). The dramatic reduction of

the SloCycl-Krt15-hi cluster in the stronger Drp1 knockdown cells could be due to their

inability of exiting the elevated CyclinB1/CDK1 driven G2-M regulation in this cluster

(Fig. 4H, left). Also, the stronger Drp1 knockdown maintained marked elevation of the

growth factor cyclin, Cyclin D2 and reduction of the S phase marker, PCNA, across

clusters (Fig. 4H, right); Cyclin E1 and Sox2 transcripts were not detected in this data

set. Therefore, our data demonstrates that particularly fine-tuned Drp1 repression

sustains a gene expression profile characteristic of the neoplastic stem cell enriched

Drp1$^-$attenuated-[CyclinE^{hi}-Sox2^{hi}-Krt15^{hi}]-TF-1 population, which is prevented by more

complete Drp1 repression that induces cell cycle disturbances.

Taken together, detailed molecular and mitochondrial shape analyses of our

newly derived transformed TF-1 and its Parental keratinocyte model suggest, for the

first time, that a fine-tuned "goldilocks" level of Drp1 repression maintains small

networks of fused mitochondria to prime a slow cycling 'stem/progenitor-like [CyclinE^{hi}-

Sox2^{hi}-Krt15^{hi}] state' towards accelerating neoplastic transformation. Cyclin E, which

maintains expression of stem cell markers, is indispensable for exiting cell cycle

quiescence and also for neoplastic transformation (Hwang & Clurman, 2005, Schaefer

& Lengerke, 2020, Teixeira & Reed, 2017)(Liu et al., 2019). We speculate that our

defined CyclinE^{hi} state, enriched by fine-tuned Drp1 repression, may reduce depth of

cell cycle quiescence (van Velthoven & Rando, 2019) towards rapidly transitioning to a

self-renewing/proliferation state to support neoplastic transformation. Such transition

may involve further activation of mitochondrial fission induced by signaling pathways

supporting stemness which would be prevented with more complete Drp1 repression

(as shown here) or total ablation of Drp1 (Kashatus, Nascimento et al., 2015,

Serasinghe et al., 2015)(Spurlock et al., 2019). This conceptualization helps explain

how regulated reduction and increase of Drp1 activity can potentially maintain adult

stem cell properties (Iwata et al., 2020, Khacho & Slack, 2017, Parker et al., 2015, Xie

et al., 2015). Further *in vivo* validation of our findings would reveal if and how bodily

stresses causing mild repression of Drp1 driven mitochondrial fission, can possibly

sustain the mitochondria primed 'stem/progenitor-like state' in tissues making them

susceptible to neoplastic transformation. Enhanced mitochondrial fusion, proposed as a

required event of neoplastic transformation of stem cells (Bonnay et al., 2020), may also

involve such mitochondria driven priming.

MATERIALS AND METHODS

Materials

Biochemicals were obtained from Fisher Biochemicals or Sigma. Other materials: Drp1

shRNA (Dharmacon); Flouromount G (SouthernBiotech); FuGENE® 6 Transfection

Reagent (Promega); Luminata Forte Western HRP substrate (Millipore); 4%

paraformaldehyde aqueous solution and Triton X-100 (FisherScientific); Labtek

chambers (Nalgene Nunc International); DMEM (GIBCO); Geltrex (GIBCO); Primary

antibodies were against: Drp1 (western blotting, 1:1000; immunofluorescence, 1:100;

BD Biosciences); Cyclin E1 (HE12) (western blotting, 1:1000; immunofluorescence

1:100; Cell Signaling); actin AB-5 (1:10000; BD Biociences), Sox2 (western blotting,

1:500; immunofluorescence, 1:100; BD Bioscience), HSP-60 (1:200; BD Bioscience);

Cytokeratin 15 (western blotting, 1:2000; immunofluorescence, 1:200; Abcam); Oct4

(immunofluorescence, 1:50; Cell Signaling); Nanog (immunofluorescence, 1:50; Cell

Signaling). Secondary antibodies were from Jackson ImmunoResearch Laboratories.

Athymic nude mice were purchased from Jackson laboratories.

Cell culture methods

HaCaT cells were cultured on plastic dishes in Dulbecco's Modified Eagle's Medium

(DMEM) with high glucose (4.5 g/L), sodium pyruvate (1 mM), L-Glutamine (4 mM),

Penicillin (100 U/mL), Streptomycin (100 µg/mL), and 10% FBS using standard

techniques. Transformed keratinocytes were derived from the Parental HaCaTs by

selecting multiple clones after treatment with the noted dose of TCDD with

replenishment with fresh TCDD every 2-3 days. The derived lines were treated with two

courses of BM-Cyclin treatment to eliminate any mycoplasma infection.

Transfections were preformed using Fugene 6 Transfection Reagent modifying

manufacturer's protocols. Lines stably expressing Drp1-shRNAs or PSmO were

selected using puromycin treatment (2 μg/mL) with regular media changes for 2 weeks

until resistant colonies grew. The stable Drp1 knockdown lines were treated with two

courses of BM-Cyclin treatment to eliminate any mycoplasma infection.

Standard cell proliferation assay was performed as described in (Parker et al.,

2015). The extreme limiting dilution analysis (ELDA) of spheroid formation ability was

performed by seeding 1, 10, 100, and 1000 cells per well with 24 wells per dose in a 96

well UltraLow Attachment plate and TIC-supporting media (DMEM:F12 supplemented

with human growth factors as published elsewhere (Spurlock et al., 2019)). Wells were

Supplementaryemented with media equaling 10% of the total volume every second day.

On Day 5, the total number of wells for each dose containing nascent spheroids was

tabulated. The online ELDA tool was used to analyze the results (Hu & Smyth, 2009).

<u>Immunoblotting and Immunofluorescence</u>

Immunoblotting was performed using standard techniques. Whole cell lysates were run

on 10% polyacrylamide gels and transferred to PVDF membranes followed by probing

with appropriate primary and secondary antibodies.

Immunofluorecence was performed as described previously (Parker et al., 2015)

on cells seeded in LabTek 8-well glass bottom or slide chambers. Briefly, cells were

fixed in freshly prepared 4% aqueous paraformaldehyde supplemented with 4 %w/v

sucrose and permeablized in freshly prepared 0.1 % Triton X-100 prior to staining.

Immunostained cells were mounted in Fluoromount with DNA stain Hoechst 33342 (10

µg/mL).

Confocal microscopy, image processing and analysis

Confocal microscopy was performed on a Zeiss LSM700 microscope equipped with 405

nm, 488 nm, 555 nm, and 639 nm lasers, using the proprietary Zen Black (2012)

software. Confocal micrographs were acquired with optimized laser powers and

appropriate filters to minimize crosstalk, cross-excitation and bleaching. Live cells were

imaged using a temperature and CO_2 controlled chamber set to 37 C and 5% CO_2.

Image processing and analysis of relative protein abundance and localization

were performed using Zen Black and Zen Blue software to obtain background corrected

mean fluorescence intensities within defined regions of interest drawn on maximum

intensity z-projections of optical slices. Nuclear and cytosolic regions were demarcated

based on DNA stain Hoechst 33342 and the immunostain of the molecule of interest,

respectively. The [Fission] and [Fusion1/5] metrics using MitoGraph v2.1 software run

on 3D stacks of confocal optical slices acquired from live cells expressing mito-PSmO

or stained with Mitotracker Green, as described in (Spurlock et al., 2019). [Fission] is

total mitochondria number / total mitochondrial length; [Fusion5] is (sum of top 5

mitochondrial length / total mitochondrial length) X 100. Microsoft Excel and IBM SPSS

Statistics 23 were used to perform background corrections, bivariate analyses and

significance testing.

Tumor Forming Assay in mice and histochemistry

5 million transformed HaCaT cells were injected in subcutaneously in the flank of

athymic nude mice to form xenograft tumors. Prior to injection, cells were trypsinized

and washed once with sterile PBS, and finally resuspended to 25,000 cells/μL in room

temperature PBS with Geltrex. Tumors arising within 6 to 8 months were harvested for

histochemical analyses. The harvested tumors were frozen unfixed in TissueTek OCT

compound on dry-ice. A Cryostat was used to obtain 5 μm slices, which were then fixed

using freshly made cold 4% paraformaldehyde and then stained with H&E using

standard techniques.

Single Cell RNAseq and data analyses

Trypsinized single cells were washed in PBS (without ca++ or Mg++) with 0.04% BSA

and tested for >90% viability. Single cell analysis was performed on a 10xGenomics

platform according to 3' v3.1 NextGem Dual Index manual. The 3'-biased cDNA libraries

were constructed through the following steps: cDNA fragmentation, end repair & A-

tailing, and size selection by SPRIselect beads, adaptor ligation, and sample index PCR

amplification, and then SPRIselect beads size selection again. The final constructed 3'-

biased single cell libraries were sequenced by Illumina Nextseq500 machine, targeting

total reads per cell for 20,000 at minimum, and the sequencing cycles consisted of 28bp

for read 1, 90 bp for read 2, and 10 bp for i7 index, and 10bp for i5 index.

Count matrices were generated from the single-cell raw fastq files using 10x

genomics *cellranger* software (v.4.0) using hg38 reference genome provided by 10x

Genomics. We filtered the data to only include cells expressing at least 2000 detectable

genes (using "nFeature_RNA" filter); this also filtered out dead cells with >10% mt-DNA

gene expression in our data set. Our data coverage included approximately ~ 3000

genes in average of 5000 cells in each population. The resulting count matrices were

analyzed by Seurat package (v3.9.9) with the standard workflow (Butler *et al*, 2018;

Stuart *et al*, 2019). After applying the (*nFeature_RNA* > 2000), the expression data were

then normalized using the *NormalizeData* function in Seurat and variable featuresets

were identified using the *FindVariableFeatures* function in Seurat. Depending on the

type of comparison, specific sets of samples were combined using the *IntegrateData*

function. The integrated datasets were scaled and cells were clustered with up to 20

dimensions (dim=1:20). We performed clustering using several resolution parameters

(0.1-0.8) and visually selected optimal resolution for specific datasets. The clusters were

visualized using UMAP with up to 20 dimensions. Marker genes for each cluster were

calculated with the *FindAllMarkers* function and statistical significance was calculated by

Wilcoxon Rank Sum test. Differential expression of the cluster markers was carried out

using the FindMarkers function of Seurat. For this, we only used the normalized data

before integration (assay 'RNA') using the Wilcoxon rank sum test, as suggested by

Seurat documentation. For cell-cycle scoring, we use the CellCycleScoring function of

Seurat, which provides scores of G2M and S phase scores and assigns cell-cycle

phases based on G2-M and S scores, or in G1/G0 when both scores are low (Tirosh *et*

al, 2016). The cell-lineage determination and trajectory calculation were carried out

using the *Slingshot* (Street *et al*, 2018) package with Seurat clusters and default

parameters. The algorithm was allowed to automatically identify the start and the end

clusters. GSEA analyses were performed with ranked log fold change (LFC) of markers

(>0.1, p<0.01) using MsigDB as described in (Tanwar et al., 2016). Overrepresentation

analyses were performed with genes with negative LFC in Cluster 5 using the

Reactome PA package of R v 3.6.0.

Statistical analyses:

Drawing of plots and statistical analyses were performed using Excel, SPSS or R

package as mentioned in appropriate section above.

ACKNOWLEDGEMENTS:

BS, KM, AM are supported by NIH award (# R33ES025662). We acknowledge Dezhi

Wang from the Pathology Core Research Laboratory for preparation of H&E stained

slides, UAB Flow Cytometry and Single Cell Services Core (supported by AR048311,

AI027667) and the Dr. Mike Crowley from the Heflin Genomics Core for sc-RNAseq.

AUTHOR CONTRIBUTION

BS designed and performed experiments and analyzed data along with AM and AG; DP

generated the transformed lines and performed basic characterization; ABH supervised

the xenograft experiments performed by BS and SG; GPS performed the pathological

evaluation of tumor sections; SL provided consultation for the sc-RNAseq and prepared

the scRNAseq library; MKB provided consultation for the sc-RNAsq analyses and

performed the analyses; KM conceived the study, helped in experimental design,

analyzed data and wrote the manuscript with BS.

DECLARATION OF INTEREST

The authors declare no competing interests.

 FIGURE LEGENDS:

Figure 1. Establishment of a mild carcinogen transformed keratinocyte model that

maintains abundant self-renewing/proliferating cells and attenuated Drp1 activity.

A) Cell proliferation assay (quantified by crystal violet staining) of HaCaT cells after

exposure to TCDD (1 or 10 nM) over 4 and 16 days, with Toluene (10nM) as vehicle

control. **B)** Immunoblot analyses of AHR and actin (loading control) in transformed

HaCaT populations, TF-1 and TF-10, in the absence of TCDD. **C)** Representative

micrographs (40X) of H&E stained sections of tumor xenografts formed by TF-1 and TF-

10 HaCaT cells; zoom of boxed region shown in the bottom panels. **D)** Immunoblot

analyses of Drp1, pDrp1(S616), Mfn1, Mfn2, Opa1 and actin (loading control) in TF-1

and TF-10 HaCaT cells. **E)** Maximum intensity projection of confocal optical sections of

representative Parental, TF-1 and TF-10 HaCaT cells expressing mitoPSmO. **F)**

Quantification of mitochondrial [Fission] and [Fusion5] metrics from micrographs

represented in (E). **G)** Quantification of cell proliferation assay (as in A) of Parental, TF-

1 and TF-10 HaCaT cells over 7 days in the absence of TCDD. **H)** Representative

micrograph showing spheroids formed when 10^4 Parental, TF-1 or TF-10 HaCaT cells

were maintained in low attachment plate in presence of stem cell medium. **I)**

Quantification of spheroid forming frequency of Parental, TF-1 or TF-10 HaCaT cells

using ELDA. * signifies p value of <0.05 in T test (A,F) and ELDA (I); scale bar depicts

10 μm (D) and 100 μm (H).

Figure. 2. Transformed Drp1-attenuated-TF-1 keratinocyte population maintains

an expanded slow-cycling sub-population with elevated stem/progenitor cell

markers. A) UMAP plot of scRNA-seq derived clusters (0-5) of Parental, TF-1 and TF-

10 HaCaT cells (top); pie chart of the percentage distribution of the clusters (bottom, p

value from ChiSquare test); color coded arrows point the clusters with unique reciprocal

abundance in TF-1 cells. **B)** Violin plots depicting expression of Krt13, Krt15, Krt5, Sox2

and Sox4 in the clusters identified in (A) in the Parental, TF-1 and TF-10 HaCaT cells;

color coded arrow point to Cluster 3 marked by Krts. **C)** Normalized Enrichment Scores

(NES) of functional pathways identified by GSEA analyses of marker genes of Clusters

0-4 (Supplementary data Table 2) and by overrepresentation analyses for Cluster 5 (*

represents categories including mtDNA genes); functional categories defined by

majority of the leading-edge genes are in parentheses, while clusters are arranged

based on trajectory analyses (open arrows) (see Supplementary Fig. 2F). **D)** Violin plots

depicting expression of cyclinA2, cyclinB1, CDK1, cyclinD2 and PCNA in the clusters

identified in (A) in the Parental, TF-1 and TF-10 HaCaT cells; the functional category of

genes identified in (C) is included in the cluster identity. **E)** Distribution of G2-M and S

scores obtained by gene expression analyses of cell cycle predictive genes in the

Parental, TF-1 and TF-10 HaCaT cells. **F)** Feature plot of G2-M and S scores obtained

from (E). **G)** Bar plot showing fraction of cells in G2-M, S and G0/G1, computed from (E)

in the functionally categorized clusters in the Parental, TF-1 and TF-10 HaCaT cells;

inset shows G1 to S reciprocity of the clusters (color-coded arrows) with unique

reciprocal abundance in TF-1 cells in (A).

Figure 3. As opposed to more complete Drp1 repression, fine-tuned Drp1

repression sustains a [CyclinE^{hi}-Sox2^{hi}-Krt15^{hi}] state and accelerates carcinogen

driven keratinocyte transformation. A) Immunoblot analyses of Sox2, Krt15 and actin

(loading control) in Parental and TF-1 and TF-10 HaCaT cells. **B)** Dot plot of [Sox2] and

[Krt15] levels obtained from confocal micrographs of co-immunostained Parental and

TF-1 HaCaT cells; black lines, marking 4 quadrants, represent levels of Sox2 and Krt15

determined from normalized signal in each population (see Supplementary Fig. 3B). **C)**

Dot plot of [Sox2] and [Drp1] levels obtained from confocal micrographs of co-

immunostained Parental and TF-1 HaCaT cells; dotted lines mark the levels of [Drp1] in

cells expressing higher [Sox2] levels. **D)** Dot plot of [Sox2] and [pDrp1(S616)] levels

obtained from confocal micrographs of co-immunostained Parental and TF-1 HaCaT

cells; dotted lines mark the levels of [pDrp1(S616)] in cells expressing higher [Sox2]

levels. **E)** Immunoblot analyses of Drp1, Krt15, Sox2 and actin (loading control) in

Parental HaCaT cells expressing non-targeted shRNA (Cnt), weaker (W) or stronger (S)

Drp1 shRNA. **F)** Immunoblot analyses of Cyclins E1, E2, A2, B1, D2 and actin (loading

control) in Parental and stable TF-1 and TF-10 HaCaT cells. **G)** Representative confocal

micrographs of Cyclin E1 immunostaining in Parental and TF-1 HaCaT cells; N depict

the nucleus in each cell identified by DNA stain (not shown). **H)** Box plot showing

quantification of nuclear to cytosolic ratio of Cyclin E1 levels quantified from experiment

described in (G). **I)** Dot plot of [Sox2] and [CyclinE1(pT62)] levels obtained from

confocal micrographs of co-immunostained Parental and TF-1 HaCaT cells; dotted lines

mark the levels of [CyclinE1(pT62)] in cells expressing higher [Sox2] levels. **J)**

Representative images of Cyclin E1 immunostaining in Parental HaCat cells expressing

non-targeted shRNA (Cnt), weaker (W) or stronger (S) Drp1 shRNA. **K)** Box plot

showing quantification of nuclear to cytosolic ratio of Cyclin E1 levels quantified from

experiment described in (J). **L)** Cell proliferation assay of Parental HaCaT cells

expressing non-targeted shRNA (Cnt), weaker (W) or stronger (S) Drp1 shRNA after

exposure to TCDD (1nM or 10nM) at an early and a late time point in the neoplastic

transformation protocol; 'Toluene' is the vehicle control for TCDD and 'Control' is with

no chemical. * signifies p value of <0.05 in KW (G) and T test (L); scale bar depicts 20

μm.

Figure 4. Fine-tuned Drp1 repression maintains smaller fused mitochondrial

elements, and sustains transcriptomic profile similar to the neoplastic stem cell

enriched TF-1 population, which is suppressed by more complete Drp1

repression. A) Dot plot of [Fission] and [Fusion5] obtained from confocal micrographs

of Mitotracker green stained live Parental HaCaT cells expressing non-targeted shRNA

as control (Cnt), weaker (W) or stronger (S) Drp1 shRNA; numbers denote the

percentage of cells in the adjacent boxes with p value from ChiSquare test;

representative images in Supplementary Fig. 3D **B)** Dot plot of [Fission] or [Fusion5] of

single cells (from A) with their total mitochondrial number and total mitochondrial length;

lines represent the ranges of [Fission] and [Fusion5] to demarcate cell subpopulations in

boxes in (A). **C)** Bar graph signifying percentage of cells from (A) with >80 [Fusion1]

metric (top); dot plot of [Fission] and [Fusion1] of cells from (A) (bottom). **D)** Barplots

showing positive and negative log fold change values of top 20 genes in Parental

HaCaT cells expressing weaker (W) (Top) or stronger (S) (bottom) shRNA with respect

to the cells expressing non-targeting shRNA (Cnt); 0 signifies no change and color

coded * denote relevant Krt genes. **E)** Barplot showing ratio of absolute expression

values of mitochondrial genes (top) and ribosomal or protein translation genes (bottom)

in the Parental HaCat cells expressing the stronger (S) or the weaker (W) Drp1 shRNA

with respect to those expressing non-targeting shRNA (Cnt); 1 signifies no change. **F)**

Line plot showing % population of Parental cells expressing the strong (S) or weak (W)

Drp1 shRNA in the named scRNA-seq derived cell clusters, with respect to those

expressing non-targeting shRNA (Cnt). This is compared to the same in the TF-1 cells

with respect to its Parental population; 1 signifies no change from respective control. **G)**

Distribution of G2-M and S scores obtained by gene expression analyses of cell cycle

predictive genes in the Parental HaCaT cells expressing non-targeted shRNA (Cnt),

weaker Drp1-shRNA (W) or stronger shRNA (S). **H)** Violin plots depicting expression of

cyclinB1, CDK1 (left panel) and cyclinD2, PCNA (right panel) in the named scRNA-seq

derived cell clusters in the Parental HaCaT cells expressing non-targeted shRNA (Cnt),

weaker Drp1-shRNA (W) or stronger shRNA (S).

REFERENCES

Boelsma E, Verhoeven MC, Ponec M (1999) Reconstruction of a human skin equivalent using a spontaneously transformed keratinocyte cell line (HaCaT). J Invest Dermatol 112: 489-98

Bonnay F, Veloso A, Steinmann V, Kocher T, Abdusselamoglu MD, Bajaj S, Rivelles E, Landskron L, Esterbauer H, Zinzen RP, Knoblich JA (2020) Oxidative Metabolism Drives Immortalization of Neural Stem Cells during Tumorigenesis. Cell 182: 1490-1507 e19

Boumahdi S, Driessens G, Lapouge G, Rorive S, Nassar D, Le Mercier M, Delatte B, Caauwe A, Lenglez S, Nkusi E, Brohee S, Salmon I, Dubois C, del Marmol V, Fuks F, Beck B, Blanpain C (2014) SOX2 controls tumour initiation and cancer stem-cell functions in squamous-cell carcinoma. Nature 511: 246-50

Butler A, Hoffman P, Smibert P, Papalexi E & Satija R (2018) Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nature Biotechnology* 36: 411–420

Chen H, Chan DC (2017) Mitochondrial Dynamics in Regulating the Unique Phenotypes of Cancer and Stem Cells. Cell Metab 26: 39-48

Der Vartanian A, Quetin M, Michineau S, Aurade F, Hayashi S, Dubois C, Rocancourt

D, Drayton-Libotte B, Szegedi A, Buckingham M, Conway SJ, Gervais M, Relaix F

(2019) PAX3 Confers Functional Heterogeneity in Skeletal Muscle Stem Cell

Responses to Environmental Stress. Cell Stem Cell 24: 958-973 e9

Di Meglio P, Duarte JH, Ahlfors H, Owens ND, Li Y, Villanova F, Tosi I, Hirota K, Nestle

FO, Mrowietz U, Gilchrist MJ, Stockinger B (2014) Activation of the aryl hydrocarbon

receptor dampens the severity of inflammatory skin conditions. Immunity 40: 989-1001

Favaro G, Romanello V, Varanita T, Andrea Desbats M, Morbidoni V, Tezze C, Albiero

M, Canato M, Gherardi G, De Stefani D, Mammucari C, Blaauw B, Boncompagni S,

Protasi F, Reggiani C, Scorrano L, Salviati L, Sandri M (2019) DRP1-mediated

mitochondrial shape controls calcium homeostasis and muscle mass. Nat Commun 10:

2576

Folmes CD, Dzeja PP, Nelson TJ, Terzic A (2012) Metabolic plasticity in stem cell

homeostasis and differentiation. Cell Stem Cell 11: 596-606

Foronda M, Martinez P, Schoeftner S, Gomez-Lopez G, Schneider R, Flores JM,

Pisano DG, Blasco MA (2014) Sox4 links tumor suppression to accelerated aging in

mice by modulating stem cell activation. Cell Rep 8: 487-500

Friedman JR, Nunnari J (2014) Mitochondrial form and function. Nature 505: 335-43

Fuchs E (2016) Epithelial Skin Biology: Three Decades of Developmental Biology, a

Hundred Questions Answered and a Thousand New Ones to Address. Curr Top Dev

Biol 116: 357-74

Geng Y, Yu Q, Sicinska E, Das M, Schneider JE, Bhattacharya S, Rideout WM,

Bronson RT, Gardner H, Sicinski P (2003) Cyclin E ablation in the mouse. Cell 114:

431-43

Giroux V, Lento AA, Islam M, Pitarresi JR, Kharbanda A, Hamilton KE, Whelan KA,

Long A, Rhoades B, Tang Q, Nakagawa H, Lengner CJ, Bass AJ, Wileyto EP, Klein-

Szanto AJ, Wang TC, Rustgi AK (2017) Long-lived keratin 15+ esophageal progenitor

cells contribute to homeostasis and regeneration. J Clin Invest 127: 2378-2391

Gonzales KAU, Fuchs E (2017) Skin and Its Regenerative Powers: An Alliance between

Stem Cells and Their Niche. Dev Cell 43: 387-401

Hao N, Lee KL, Furness SG, Bosdotter C, Poellinger L, Whitelaw ML (2012)

Xenobiotics and loss of cell adhesion drive distinct transcriptional outcomes by aryl

hydrocarbon receptor signaling. Mol Pharmacol 82: 1082-93

Hoppins S (2014) The regulation of mitochondrial dynamics. Curr Opin Cell Biol 29: 46-

52

Hu Y, Smyth GK (2009) ELDA: extreme limiting dilution analysis for comparing depleted

and enriched populations in stem cell and other assays. J Immunol Methods 347: 70-8

Hwang HC, Clurman BE (2005) Cyclin E in normal and neoplastic cell cycles.

Oncogene 24: 2776-86

Iwata R, Casimir P, Vanderhaeghen P (2020) Mitochondrial dynamics in postmitotic

cells regulate neurogenesis. Science 369: 858-862

Jung MH, Jung SM, Shin HS (2016) Co-stimulation of HaCaT keratinization with

mechanical stress and air-exposure using a novel 3D culture device. Sci Rep 6: 33889

Kageyama Y, Zhang Z, Sesaki H (2011) Mitochondrial division: molecular machinery

and physiological functions. Curr Opin Cell Biol 23: 427-34

Karantza V (2011) Keratins in health and cancer: more than mere epithelial cell

markers. Oncogene 30: 127-38

Kashatus JA, Nascimento A, Myers LJ, Sher A, Byrne FL, Hoehn KL, Counter CM,

Kashatus DF (2015) Erk2 phosphorylation of Drp1 promotes mitochondrial fission and

MAPK-driven tumor growth. Mol Cell 57: 537-51

Khacho M, Clark A, Svoboda DS, Azzi J, MacLaurin JG, Meghaizel C, Sesaki H,

Lagace DC, Germain M, Harper ME, Park DS, Slack RS (2016) Mitochondrial Dynamics

Impacts Stem Cell Identity and Fate Decisions by Regulating a Nuclear Transcriptional

Program. Cell Stem Cell 19: 232-247

Khacho M, Slack RS (2017) Mitochondrial activity in the regulation of stem cell self-

renewal and differentiation. Curr Opin Cell Biol 49: 1-8

Leclerc D, Staats Pires AC, Guillemin GJ, Gilot D (2021) Detrimental activation of AhR

pathway in cancer: an overview of therapeutic strategies. Curr Opin Immunol 70: 15-26

Li S, Park H, Trempus CS, Gordon D, Liu Y, Cotsarelis G, Morris RJ (2013) A keratin 15

containing stem cell population from the hair follicle contributes to squamous papilloma

development in the mouse. Mol Carcinog 52: 751-9

Liang R, Arif T, Kalmykova S, Kasianov A, Lin M, Menon V, Qiu J, Bernitz JM, Moore K,

Lin F, Benson DL, Tzavaras N, Mahajan M, Papatsenko D, Ghaffari S (2020)

Restraining Lysosomal Activity Preserves Hematopoietic Stem Cell Quiescence and

Potency. Cell Stem Cell 26: 359-376 e7

Lisowski P, Kannan P, Mlody B, Prigione A (2018) Mitochondria and the dynamic

control of stem cell homeostasis. EMBO Rep 19

Liu L, Michowski W, Kolodziejczyk A, Sicinski P (2019) The cell cycle in stem cell

proliferation, pluripotency and differentiation. Nat Cell Biol 21: 1060-1067

Luecken MD, Theis FJ (2019) Current best practices in single-cell RNA-seq analysis: a

tutorial. Mol Syst Biol 15: e8746

Maga G, Hubscher U (2003) Proliferating cell nuclear antigen (PCNA): a dancer with

many partners. J Cell Sci 116: 3051-60

Magee JA, Piskounova E, Morrison SJ (2012) Cancer stem cells: impact, heterogeneity,

and uncertainty. Cancer Cell 21: 283-96

Mandal S, Freije WA, Guptan P, Banerjee U (2010) Metabolic control of G1-S transition:

cyclin E degradation by p53-induced activation of the ubiquitin-proteasome system. J

Cell Biol 188: 473-9

Mitra K, Lippincott-Schwartz J (2010) Analysis of mitochondrial dynamics and functions

using imaging approaches. Curr Protoc Cell Biol Chapter 4: Unit 4 25 1-21

Mitra K, Rikhy R, Lilly M, Lippincott-Schwartz J (2012) DRP1-dependent mitochondrial

fission initiates follicle cell differentiation during Drosophila oogenesis. J Cell Biol 197:

487-97

Mitra K, Wunder C, Roysam B, Lin G, Lippincott-Schwartz J (2009) A hyperfused

mitochondrial state achieved at G1-S regulates cyclin E buildup and entry into S phase.

Proc Natl Acad Sci U S A 106: 11960-5

Moriyama M, Durham AD, Moriyama H, Hasegawa K, Nishikawa S, Radtke F, Osawa M

(2008) Multiple roles of Notch signaling in the regulation of epidermal development. Dev

Cell 14: 594-604

Morris RJ, Liu Y, Marles L, Yang Z, Trempus C, Li S, Lin JS, Sawicki JA, Cotsarelis G

(2004) Capturing and profiling adult hair follicle stem cells. Nat Biotechnol 22: 411-7

Mulero-Navarro S, Fernandez-Salguero PM (2016) New Trends in Aryl Hydrocarbon

Receptor Biology. Front Cell Dev Biol 4: 45

Nagdas S, Kashatus JA, Nascimento A, Hussain SS, Trainor RE, Pollock SR, Adair SJ,

Michaels AD, Sesaki H, Stelow EB, Bauer TW, Kashatus DF (2019) Drp1 Promotes

KRas-Driven Metabolic Changes to Drive Pancreatic Tumor Growth. Cell Rep 28: 1845-

1859 e5

Orford KW, Scadden DT (2008) Deconstructing stem cell self-renewal: genetic insights

into cell-cycle regulation. Nat Rev Genet 9: 115-28

Otsuki L, Brand AH (2018) Cell cycle heterogeneity directs the timing of neural stem cell

activation from quiescence. Science 360: 99-102

Parker DJ, Iyer A, Shah S, Moran A, Hjelmeland AB, Basu MK, Liu R, Mitra K (2015) A

new mitochondrial pool of cyclin E, regulated by Drp1, is linked to cell-density-

dependent cell proliferation. J Cell Sci 128: 4171-82

Pauklin S, Vallier L (2013) The cell-cycle state of stem cells determines cell fate

propensity. Cell 155: 135-47

Prieto J, Leon M, Ponsoda X, Sendra R, Bort R, Ferrer-Lorente R, Raya A, Lopez-

Garcia C, Torres J (2016) Early ERK1/2 activation promotes DRP1-dependent

mitochondrial fission necessary for cell reprogramming. Nat Commun 7: 11124

Purba TS, Haslam IS, Poblet E, Jimenez F, Gandarillas A, Izeta A, Paus R (2014)

Human epithelial hair follicle stem cells and their progeny: current state of knowledge,

the widening gap in translational research and future challenges. Bioessays 36: 513-25

Ray SS, Swanson HI (2004) Dioxin-induced immortalization of normal human

keratinocytes and silencing of p53 and p16INK4a. J Biol Chem 279: 27187-93

Ryan MT, Hoogenraad NJ (2007) Mitochondrial-nuclear communications. Annu Rev

Biochem 76: 701-22

Sang L, Coller HA, Roberts JM (2008) Control of the reversibility of cellular quiescence

by the transcriptional repressor HES1. Science 321: 1095-100

Schaefer T, Lengerke C (2020) SOX2 protein biochemistry in stemness,

reprogramming, and cancer: the PI3K/AKT/SOX2 axis and beyond. Oncogene 39: 278-

292

Schoop VM, Mirancea N, Fusenig NE (1999) Epidermal organization and differentiation

of HaCaT keratinocytes in organotypic coculture with human dermal fibroblasts. J Invest

Dermatol 112: 343-53

Schrepfer E, Scorrano L (2016) Mitofusins, from Mitochondria to Metabolism. Mol Cell

61: 683-694

Senos Demarco R, Uyemura BS, D'Alterio C, Jones DL (2019) Mitochondrial fusion

regulates lipid homeostasis and stem cell maintenance in the Drosophila testis. Nat Cell

Biol 21: 710-720

Serasinghe MN, Wieder SY, Renault TT, Elkholi R, Asciolla JJ, Yao JL, Jabado O,

Hoehn K, Kageyama Y, Sesaki H, Chipuk JE (2015) Mitochondrial division is requisite

to RAS-induced transformation and targeted by oncogenic MAPK pathway inhibitors.

Mol Cell 57: 521-36

Seykora JT, Cotsarelis G (2011) Keratin 15-positive stem cells give rise to basal cell

carcinomas in irradiated Ptch1(+/-) mice. Cancer Cell 19: 5-6

Siegle JM, Basin A, Sastre-Perona A, Yonekubo Y, Brown J, Sennett R, Rendl M,

Tsirigos A, Carucci JA, Schober M (2014) SOX2 is a cancer-specific regulator of tumour

initiating potential in cutaneous squamous cell carcinoma. Nat Commun 5: 4511

Simpson CL, Tokito MK, Uppala R, Sarkar MK, Gudjonsson JE, Holzbaur ELF (2021)

NIX initiates mitochondrial fragmentation via DRP1 to drive epidermal differentiation.

Cell Rep 34: 108689

Siu KT, Rosner MR, Minella AC (2012) An integrated view of cyclin E function and

regulation. Cell Cycle 11: 57-64

Spurlock B, Gupta P, Basu MK, Mukherjee A, Hjelmeland AB, Darley-Usmar V, Parker

D, Foxall ME, Mitra K (2019) New quantitative approach reveals heterogeneity in

mitochondrial structure-function relations in tumor initiating cells. J Cell Sci

Spurlock B, Tullet J, Hartman JLt, Mitra K (2020) Interplay of mitochondrial fission-

fusion with cell cycle regulation: Possible impacts on stem cell and organismal aging.

Exp Gerontol 135: 110919

Spurlock B and Mitra K (2021a) Methods Molecular Biology, Vol. 2275, Volkmar

Weissig and Marvin Edeas (Eds): Mitochondrial Medicine , 978-1-0716-1261-3,

482923_2_En, (Chapter 27).

Spurlock B, Hanumanthu VS and Mitra K (2021b). Strategy of Isolating 'Primed' Tumor

Initiating Cells Based on Mitochondrial Transmembrane Potential. Bio-protocol online,

inPress.

St John LS, Sauter ER, Herlyn M, Litwin S, Adler-Storthz K (2000) Endogenous p53

gene status predicts the response of human squamous cell carcinomas to wild-type

p53. Cancer Gene Ther 7: 749-56

Street K, Risso D, Fletcher RB, Das D, Ngai J, Yosef N, Purdom E & Dudoit S (2018)

Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics. *BMC*

Genomics 19: 477

Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, Hao Y,

Stoeckius M, Smibert P & Satija R (2019) Comprehensive Integration of Single-Cell

Data. *Cell* 177: 1888-1902.e21

Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich

A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP (2005) Gene set enrichment

analysis: a knowledge-based approach for interpreting genome-wide expression

profiles. Proc Natl Acad Sci U S A 102: 15545-50

Tirosh I, Izar B, Prakadan SM, Wadsworth MH, Treacy D, Trombetta JJ, Rotem A,

Rodman C, Lian C, Murphy G, *et al* (2016) Dissecting the multicellular ecosystem of

metastatic melanoma by single-cell RNA-seq. *Science* 352: 189–196

Tanwar DK, Parker DJ, Gupta P, Spurlock B, Alvarez RD, Basu MK, Mitra K (2016)

Crosstalk between the mitochondrial fission protein, Drp1, and the cell cycle is identified

across various cancer types and can impact survival of epithelial ovarian cancer

patients. Oncotarget 7: 60021-60037

Teixeira LK, Reed SI (2017) Cyclin E Deregulation and Genomic Instability. Adv Exp

Med Biol 1042: 527-547

Tsuyoshi H, Orisaka M, Fujita Y, Asare-Werehene M, Tsang BK, Yoshida Y (2020)

Prognostic impact of Dynamin related protein 1 (Drp1) in epithelial ovarian cancer. BMC

Cancer 20: 467

van Velthoven CTJ, Rando TA (2019) Stem Cell Quiescence: Dynamism, Restraint, and

Cellular Idling. Cell Stem Cell 24: 213-225

Twig G, Shirihai OS (2011) The interplay between mitochondrial dynamics and

mitophagy. Antioxid Redox Signal 14: 1939-51

Vantaggiato C, Castelli M, Giovarelli M, Orso G, Bassi MT, Clementi E, De Palma C

(2019) The Fine Tuning of Drp1-Dependent Mitochondrial Remodeling and Autophagy

Controls Neuronal Differentiation. Front Cell Neurosci 13: 120

Wincent E, Bengtsson J, Mohammadi Bardbori A, Alsberg T, Luecke S, Rannug U,

Rannug A (2012) Inhibition of cytochrome P4501-dependent clearance of the

endogenous agonist FICZ as a mechanism for activation of the aryl hydrocarbon

receptor. Proc Natl Acad Sci U S A 109: 4479-84

Wu MJ, Chen YS, Kim MR, Chang CC, Gampala S, Zhang Y, Wang Y, Chang CY,

Yang JY, Chang CJ (2019) Epithelial-Mesenchymal Transition Directs Stem Cell

Polarity via Regulation of Mitofusin. Cell Metab 29: 993-1002 e6

Xie Q, Wu Q, Horbinski CM, Flavahan WA, Yang K, Zhou W, Dombrowski SM, Huang

Z, Fang X, Shi Y, Ferguson AN, Kashatus DF, Bao S, Rich JN (2015) Mitochondrial

control by DRP1 in brain tumor initiating cells. Nat Neurosci 18: 501-10

Zerjatke T, Gak IA, Kirova D, Fuhrmann M, Daniel K, Gonciarz M, Muller D, Glauche I,

Mansfeld J (2017) Quantitative Cell Cycle Analysis Based on an Endogenous All-in-One

Reporter for Cell Tracking and Classification. Cell Rep 19: 1953-1966

Zhang H, Menzies KJ, Auwerx J (2018) The role of mitochondria in stem cell fate and

aging. Development 145

1000

1001   Zhao Q, Yan S, Lu J, Parker DJ, Wu H, Sun Q, Crossman DK, Liu S, Wang Q, Sesaki

1002   H, Mitra K, Liu K, Jiao K (2021) Mitochondrial Fission Regulates Transcription of

1003   Ribosomal Protein Genes in Embryonic Hearts

1004   bioRxiv 2021.02.10.430627; doi: https://doi.org/10.1101/2021.02.10.430627

1005

1006   Zhong X, Cui P, Cai Y, Wang L, He X, Long P, Lu K, Yan R, Zhang Y, Pan X, Zhao X, Li

1007   W, Zhang H, Zhou Q, Gao P (2019) Mitochondrial Dynamics Is Critical for the Full

1008   Pluripotency and Embryonic Developmental Potential of Pluripotent Stem Cells. Cell

1009   Metab 29: 979-992 e4

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

 SUPPLEMENTARY FIGURE LEGENDS:

Supplementary Figure 1. A) Quantification of incidence and volume of xenograft

tumors formed by TF-1 and TF-10 HaCaT cells; # indicates presence of a distant tumor.

B) Immunoblot analyses of Cytochrome C, Fis1, Tom20 and actin (loading control) in

Parental and TF-1 and TF-10 HaCaT cells.

Supplementary Figure 2. A) Bar plot showing the percentage distribution of the

clusters (0-5) in each of the cell populations identified by scRNA-seq analyses of

Parental, TF-1 and TF-10 HaCat cells (left); numbers show cell cycle distribution from

flowcytometry analyses of PI stained cells in the corresponding population (right). **B)**

UMAP plot of scRNA-seq derived clusters (0-5) of Parental, TF-1 and TF-10 HaCaT

cells obtained with minimum cluster resolution; color coded arrows point the major

change in cluster distribution in transformed cells. **C)** Heat map of statistically significant

marker genes (LFC ≥1.5 folds, p-adjusted <0.0001) for the color-coded clusters (0-5) of

Parental, TF-1 and TF-10 HaCaT cells; color coded arrows point to the top most marker

gene in the 3 clusters that are have different distribution between the Parental and

transformed populations; color index shows log fold change; * marks the Histone gene

cluster. **D)** Violin plots depicting expression of the mt-DNA genes in the named scRNA-

seq derived cell clusters in the Parental, TF-1 and TF-10 HaCaT cells. **E)** Violin plots

depicting expression of HES1 and OASL genes in the named scRNA-seq derived cell

clusters in the Parental, TF-1 and TF-10 HaCaT cells. **F)** PCA plot of trajectory analyses

of the scRNA-seq derived cell clusters in the Parental, TF-1 and TF-10 HaCaT cells.

Supplementary Figure 3. A) Representative confocal micrographs of immunostained

Parental and TF-1 HaCaT cells showing endogenous expression of Oct4, Nanog and

Sox2; blue depicting nuclear staining with Hoechst. **B)** Dot plot of [Sox2] and [Krt15]

obtained from confocal micrographs of co-immunostained Parental and TF-1 HaCaT

cells; [Sox2] levels normalized to the minimum in the respective population (left) and

[Krt15] levels normalized to the minimum in the respective population (right); black lines

demarcate levels of Sox2 and Krt15 as determined from the normalized signal of the

other. **C)** Representative confocal micrographs of Parental and TF-1 HaCaT cells

showing endogenous expression of Tom20 and Cyclin E1 after 2 hours incubation with

DMSO or Cychloheximide, with the pre-incubation time point (0h) as control. **D)**

Representative maximum intensity projections (MIP) of confocal optical sections

(micrograph) and the corresponding MitoGraph generated binary images (binary) for

quantifying [Fission] and [Fusion1/5] metrics of the Parental HaCaT cells expressing

non-targeted shRNA (Cnt), strong (S) or weak (W) Drp1 shRNA; the cells represent

populations in colored boxes in Fig. 4A while the number depicts their abundance in

percentage. Scale bar depicts 100 μm (A, C) or 15 μm (D).

Supplementary Figure 4. A) Bar plot showing the knockdown efficiency of strong (S)

or weak (W) Drp1 shRNA expressed as ratio of Drp1 transcript levels in the knockdown

HaCaT and that of the cells expressing the non-targeted shRNA control. **B)** Bar plot

showing log fold change of overall gene expression of Parental HaCaT cells expressing

strong (S) or weak (W) Drp1 shRNA with respect to those expressing the non-targeted

shRNA (Cnt). **C)** Plots showing ratio of the expression of covered genes between the

Parental HaCaT cells expressing strong (S) or weak (W) Drp1 shRNA; "Similar effects"

showing ratio of increase or decrease in the (S) Drp1 shRNA over the (W) Drp1 shRNA

(back arrows in left panel), while opposite effects showing opposite trends in genes as

labelled in the arrows (back arrows right panel); grey arrows point to the top 20 genes of

each category. **D)** UMAP plot of scRNA-seq based clusters (0-5) of Parental HaCaT

cells expressing non-targeted shRNA (Cnt), strong (S) or weak (W) Drp1 shRNA; arrow

pointing the SloCyc-Krt15-hi cluster (note color coding is not same as the one presented

in Fig. 2). **E)** Bar plot showing percentage of statistically significant common genes

(Upregulated or Downregulated) between color coded clusters obtained with the

scRNA-seq data set of parental and transformed HaCaT cells (Fig. 2) and that of

parental expressing control or Drp1 shRNAs (Fig. 4); same color coding is maintained

between the two data sets. **F)** Violin plots depicting expression of Krt13, Krt15, Krt4,

Krt5, Krt6A, Krt7 genes in the named scRNA-seq based cell clusters in the Parental

HaCaT cells expressing non-targeted shRNA (Cnt), strong (S) or weak (W) Drp1

shRNA. **G)** Bar plot showing fraction of cells in G2-M, S and G0/G1, computed from

main Fig. 4G, in the functionally categorized clusters in the Parental cells expressing

non-targeted shRNA (Cnt), strong (S) or weak (W) Drp1 shRNA.

1081

A

Cell proliferation (au)

Toluene
TCDD 1 nM
TCDD 10 nM

Exposure: 4d 16d 4d 16d 4d 16d

B

Parental TF-1 TF-10

AHR

Actin

C

TF-1   TF-10

Tumor mass

Stratified-like squamous epithelium

Tumor-host (connective tissue) interface

Squamous cell carcinoma

D

Parental TF-1 TF-10

pDrp1 (S616)

Drp1

Mfn1

Mfn2

OPA1

Actin

E

Parental   TF-1   TF-10

F

Parental TF-1 TF-10

[Fission]

[Fusion5]

G

Cell Proliferation (au)

Parental
TF-1
TF-10

Days

H

Parental   TF-1   TF-10

I

Cell Type	Stem/progenitor cell frequency
Parental	1 in 3352
TF-1	1 in 376
TF-10	1 in 603

Figure. 1

Figure. 2

Figure. 3

Figure. 4

A

Cells Injected	Tumor incidence	Tumor volume
Parental	0/9	N.A.
TF-1	#1/9	188.1 #363.1
TF-10	3/9	287.9 ± 173

B

Parental   TF-1   TF-10

Cytochrome C
Fis1
TOM20
Actin

Extended Fig. 1

Extended Fig. 2

Extended Fig. 3

Extended Fig. 4