Odor-evoked Increases in Spiking Variability in the Olfactory Bulb

Cheng Ly[†,*] ; Andrea K. Barreiro[#] ; Shree Hari Gautam[‡] ; Woodrow L. Shew[‡]

†*Department of Statistical Sciences and Operations Research, Virginia Commonwealth University,*

Richmond, VA 23284 U.S.A.

*Correspondences should be addressed to Cheng Ly: CLy@vcu.edu

Department of Mathematics, Southern Methodist University, Dallas, TX 75275 U.S.A.

E-mail: abarreiro@smu.edu

‡*Department of Physics, University of Arkansas, Fayetteville, AR 72701 U.S.A.*

E-mail: shgautam@uark.edu; woodrowshew@gmail.com

Abstract

At the onset of sensory stimulation, the variability and co-variability of spiking activity is widely reported to decrease, especially in cortex. Considering the potential benefits of such decreased variability for coding, it has been suggested that this could be a general principle governing all sensory systems. Here we show that this is not so. We recorded neurons in olfactory bulb (OB) of anesthetized rats and found increased variability and co-variability of spiking at the onset of olfactory stimulation. Using models and analysis, we predicted that these increases arise due to network interactions within OB, without increasing variability of input from the nose. We tested and confirmed this prediction using optogenetic stimulation of OB in awake animals. Our results establish increases in spiking variability at stimulus onset as a viable alternative coding strategy to the more commonly observed decreases in variability in many cortical systems.

Introduction

The variability of spiking in response to sensory input can significantly impact the coding efficacy of downstream neurons. Numerous experimental observations in diverse sensory cortical regions have established that the onset of sensory input causes a drop in the average spiking variability (Churchland et al., 2010). Similarly, the spiking co-variability among simultaneously recorded pairs of neurons decreases in the evoked state (Doiron et al., 2016). There have been many theoretical studies detailing how neuron populations can have both increases in firing rate and decreases in (co-)variability at stimulus onset. The most prominent mechanistic explanations are succinctly

summarized in Doiron et al. (2016), and include changes in noise levels via (pre-)synaptic trans-mission (Churchland et al., 2010; Ponce-Alvarez et al., 2013; Rosenbaum et al., 2013), altering transfer function of inputs to outputs (de la Rocha et al., 2007; Barreiro et al., 2010, 2012), and increases in correlated inhibition leading to cancellation of correlated inputs (Renart et al., 2010; Litwin-Kumar et al., 2011; Middleton et al., 2012; Tetzlaff et al., 2012; Ly et al., 2012).

Does stimulus-induced reduction of variability and co-variability generalize across all sensory neural systems? Here we first sought to answer this question in the olfactory bulb (**OB**), which processes sensory signals generated by olfactory receptor neurons (**ORN**) in the nose and passes this sensory information on to piriform cortex. Notably, the OB was not discussed in Churchland et al. (2010) nor in any of the experiments cited in Doiron et al. (2016) despite its crucial role in olfaction. We recorded single unit activity from anesthetized rats during precisely controlled ol-factory stimulation. We were surprised to find that stimulus onset caused an increase in variability and co-variability of spiking among mitral cells (**MC**s) in OB.

Stimulus-induced increases in spiking variability and co-variability are not widely reported and are understudied (but see Kazama and Wilson (2009); Wanner and Friedrich (2020)). Moreover, the mechanisms governing spiking variability and co-variability in OB are not well understood. Thus, our next goals were to elucidate mechanisms of variability in the OB circuit, and more specifically, to clarify why the OB variability increases rather than decreases. We focus on two possible sources of variability. First, the input to OB is generated by olfactory receptor neurons, thus variability of ORN activity is one likely source of OB variability. It has been observed that ORN activity increases with stimulus intensity (Scott et al., 2007; Furudono et al., 2013), but the extent to which the variability (across trials or across glomeruli) of ORN input changes at stimulus onset is unclear. In *Drosophila*, changes in ORN inputs that occur with stimulus onset modulate spiking of projection neurons, the analog of MCs (Kazama and Wilson, 2009). A second important factor in spiking variability of recurrent networks, in general, is inhibition. Thus, we also explored the role of granule cells (**GC**, an important inhibitory neuron in OB) in OB variability. GC activity is known to affect spiking variability and co-variability in OB (Galán et al., 2006; Giridhar et al., 2011).

To this end, we develop and analyzed computational models of the OB. We use a relatively simple firing rate model that contains many known biophysical details of the OB, including having 3 important cell types (MC, GC, and periglomerular cells (**PGC**)), sub-network architecture, mem-brane time-constants, and synaptic dynamics (based on the Cleland OB models (Li and Cleland, 2013, 2017)). Our model enables a detailed computational approach for determining coupling strengths and ORN inputs that best match our data. We found for a given ORN variance regime that uncorrelated GC inhibition to MC must be strong, excitation to GC is weaker, and lateral inhi-bition providing shared input to multiple MCs must be weak. When accounting for three key spike statistics — time-varying firing rate, spike count variance and covariance — we found that the total

error between model and data is minimized when the ORN input noise is fixed across spontaneous and evoked states. Thus our model predicts that ORN input noise does not need to increase in the evoked state to observe evoked increases in OB spiking variability.

We tested this prediction using previous experiments in awake mice (Bolding and Franks, 2018a,b). In these experiments, the optogenetic stimulation of OB acted as a surrogate input to OB, replacing ORN input, but with no increase in ORN variability at stimulus onset. In agreement with our model prediction, OB activity exhibited an increase in firing variability at the onset of optogenetic stimulation.

Taken together, experimental data and our model and theory establish increased variability at stimulus onset as a viable strategy for sensory systems and demonstrate how this occurs in OB.

Results

We performed *in vivo* multi-electrode array recordings of the OB in the mitral cell layer of anesthetized rats (see **Materials and Methods**) to capture both spontaneous and odor-evoked spiking activity of populations of putative MCs. This yielded a large number of cells and simultaneously recorded pairs of cells with heterogeneous trial-averaged spiking activity. Figure 1 summarizes the trial-averaged spiking, showing the population firing rate (peri-stimulus time histogram, **PSTH** in Fig. 1**A**), spike count variance (Fig. 1**B**) and spike count covariance (Fig. 1**C**) over 100 ms time windows. The choice of 100 ms time window is an intermediate value between the shorter time scales (membrane time constants, AMPA, GABA$_A$, etc.) and longer ones (NMDA, calcium and other ionic currents). Details about the trial-averaged spiking statistics on a cell-by-cell, or pair-by-pair basis are shown in Figure 1**D**–**F**, where the time-averaged spike count is shown in each state (spontaneous and evoked). The vast majority of cells and pairs have larger stimulus-evoked values than spontaneous.

Consistent with many sensory and motor neurons, the odor-evoked firing rates are larger than the spontaneous firing rates on average. However, note that the spike count variability (population-averaged variance, Fig. 1**B**, and Fano factor, defined as variance over mean, Fig. 1**G**) *increases* in the odor-evoked state, in contrast to many cortical regions where spiking variability decreases with stimuli (Churchland et al., 2010). Similarly, we see on average an *increase* in spike count co-variability (population-averaged covariance, Fig. 1**C**, and Pearson's correlation, Fig. 1**H**) in the evoked state compared to the spontaneous state, again in contrast to many cortical regions (Doiron et al., 2016). Another way to measure population Fano factor is via the slope of regression line (with intercept at the origin) for a cell's variance against mean spike count; similarly correlation can also be measured by slope of regression line of covariance of pairs against product of standard deviations. By these measures we again see evoked increases of these statistics (Fig. 1**I, J**).

Figure 1: **Spiking statistics from experiments. A)–C)** Spiking statistics in spontaneous ($t < 0$) and odor-evoked states ($t > 0$); the firing rate (**A**), spike count variance (**B**) and covariance (**C**). **D)–F)** Trial- and time-averaged spike count statistics of individual cells/pairs in gray (black squares = population average). **D)** Mean spike count increases with odor (vertical log-scale), 918 cells (82%) increase, time-averaged in each state. **E)** Spike count variance increases with odor (vertical log-scale), 908 cells (81.1%) increase. **F)** Spike count covariance increases with odor, 10194 pairs (57.7%) increase, some with very large evoked covariance. **G)–J)** Scaled measures of variability shown for completeness. **G)** Fano factor of spike counts (variance/mean) also increases with odor. **H)** Pearson's Correlation increases with odor. **I)** Spike count variance versus mean in spontaneous (black) and evoked (red) states, time-averaged in each state. Lines are regression of best least squares fit to data with 0 intercept, another way to measure Fano factor. Two red lines: the solid line (slope of 1.83, $R^2 = 0.660$) is regression for all data, lower line (red dashed, slope of 1.52, $R^2 = 0.657$) is regression on a subset of data (evoked mean count ≤ largest spontaneous mean count) to crudely account for larger mean counts in the evoked state (see mean-matching in Churchland et al. (2010)). Spontaneous slope is 1.21, $R^2 = 0.870$. **J)** Spike count covariance versus product of standard deviations of spike counts. The slope of the regression line is another measure of Pearson's correlation ρ; inset is zoomed in. Spontaneous slope: 0.14, $R^2 = 0.385$; evoked slope: 0.51, $R^2 = 0.750$; evoked slope with cut-off: 0.27, $R^2 = 0.565$. **K)** Spike statistics from simultaneously recorded cells in the anterior piriform cortex (**aPC**) are consistent with other data: evoked decreases in Fano factor (Churchland et al., 2010; Miura et al., 2012) and correlation (Miura et al., 2012). Spike counts were binned in 100 ms half-overlapping windows, except for **K** where 200 ms was used. Data averaged over 10 trials and over cell population (1120 cells in **A, B, D, E, G**; 17674 pairs in **C, F, H**; 194 cells and 7395 pairs in **K**). Shaded gray/tan regions show relative population heterogeneity: $\mu \pm 0.2$std (standard deviation across the population/pairs after trial-averaging; 0.2 scale for visualization).

Churchland et al. (2010) accounted for the increase in mean activity in the evoked state to attribute a
decrease in Fano factor to the spike count variance by a 'mean-matching method'. Here we account
for increases in mean spike count in a more crude and conservative way by performing linear
regression for a subset of data, removing all cells/pairs that have evoked mean count larger than
the largest spontaneous mean count and regressing on the resulting subset: we still observe evoked
increases in Fano factor and correlation (red dotted lines in Fig. 1**I, J**). Although these scaled
measures of variability increase with stimuli (see Fig. 1**G–J**), we focus on capturing the three
unscaled statistics (PSTH, variance and covariance of spike counts) because they each directly

effect common measures of neural coding (e.g. the Fisher information) whereas it is less clear how scaled statistics such as Fano factor and correlation relate to neural coding (Kohn et al., 2016).

Finally, Figure 1K shows that simultaneously recorded cells from anterior piriform cortex (**aPC**) are consistent with previous reports. Our data shows odor-evoked increases in the population firing rate (left), and odor-evoked decreases in Fano factor (middle, Churchland et al. (2010); Miura et al. (2012)) and correlation (right, Miura et al. (2012)).

Firing rate model

Dissecting the detailed effects of neural attributes, such as ORN inputs and OB coupling strengths, on MC spike statistic modulation is difficult experimentally, especially *in vivo*. We are currently unaware of any experiments that that simultaneously consider ORN activity and its affect on MC spiking (although Scott et al. (2007) considered how ORN activity affects activity in superficial OB which is considerably different than the MC layer). Therefore we develop a simple computational model that allowed us to observe the affects of different circuit manipulations.

While realistic biophysical models can capture a lot of known electrophysiology of the mammalian OB circuit (see multi-compartment OB models by Cleland and colleagues (Li and Cleland, 2013, 2017)), they are computationally expensive to simulate, thus making our thorough theoretically study intractable. We also wish to capture the time-varying statistics over a lengthy time interval of 4 seconds with 1 ms resolution, which will require many realizations in order to resolve the first and second-order firing statistics. Furthermore, we will perform a robust search in a large region of parameter space. Therefore we developed a firing rate model that retains key components of the Cleland models (Li and Cleland, 2013, 2017), including the the membrane time-constants of three important cell types (MC, GC, PGC), and the synaptic dynamics (GABA$_A$, NMDA, AMPA and their associated equations, see **Materials and Methods**).

With a firing rate model for each type of cell, we coupled these into a minimal sub-network of 7 cells with 2 representative glomeruli (Fig. 2A) each with a PGC and MC cell. We also include small noise correlations that model common input from ORNs to MC (Kazama and Wilson, 2009), as well as small correlated inputs to other OB cell types (see **Materials and Methods** for details). There are too many parameters in the circuit to feasibly consider, so we set the coupling strengths to be symmetric between the 2 glomeruli resulting in each MC being statistically identical, and we focus on lateral inhibition from GC and recurrent MC excitation to GC.

Our focus on lateral inhibition was informed by OB anatomy, and experimental and theoretical studies demonstrating the importance of the MC–GC network in controlling spiking co-variability. The long lateral GC dendrites are known to span multiple glomeruli and impinge on correspond MCs, thus it is reasonable to assume GC strongly impacts MC spike covariance. Giridhar et al. (2011) showed in slice recordings of OB that lateral (GC) inhibition can alter spike co-variability

differently depending on if it is common or independent to MC. Lateral GC inhibition is also known to modulate MC synchrony (Galán et al., 2006). Moreover, many theoretical studies have shown the importance of correlated inhibition for modulating spiking co-variability in general circuits (Renart et al., 2010; Tetzlaff et al., 2012; Middleton et al., 2012; Ly et al., 2012). To address the role of common and independent GC inhibition, we include 3 GCs with 2 providing independent inhibition to each MC and 1 GC providing common inhibition to both MCs.

Matching experimental data with model

To test whether odor-evoked increases in variability stems from changes in the variability of inputs ($\sigma(t)$) from ORN), we considered three ways to modulate the $\sigma(t)$: i) relatively small and fixed σ (**Smaller** σ), ii) σ that increases in the evoked state, except for GC cells (σ **Vary in Time**), iii) large fixed σ (**Larger** σ), see Figure 2**B**, bottom. In the 3 input noise regimes, the smaller and larger σ values match the spontaneous and evoked values of time-varying $\sigma(t)$. For all parameters, the same time-varying $\mu(t)$ is used (Fig. 2**B** top, see **Appendix: Input statistics give reasonable model results**).

For each noise regime, we performed a robust search through 3D parameter space to find the coupling strengths (w_{MG}, w_{GM}, w_{Gc}, see Fig. 2**A**) that match the time-varying firing rate (peri-stimulus time histogram, or **PSTH**) within a certain tolerance. Specifically, a parameter set is considered 'Close to Data' if: $\int_{-2}^{2} W_1(t)|r(t) - r_{data}(t)| \, dt \leq 1.4$, where $W_1(t)$ weights the uncertainty of the data (Fig. 6**A**). This tolerance results in good model fits to the data PSTH (Fig. 2**D**), and also yields hundreds of parameters for each of the 3 input noise regimes (see Fig. 2**C**).

The criteria of matching the data PSTH alone (see **Materials and Methods**) drastically reduces the size of parameter space: out of 10,000 randomly distributed points, only 361 are close to the data for small and fixed input noise (Fig. 2**Ci**), 295 for time-varying input noise (Fig. 2**Cii**), and 129 for large fixed input noise (Fig. 2**Ciii**). Interestingly, the region of parameter space that results in good fits to the data PSTH looks similar for all 3 input noise regimes. The strength of synaptic input of independent GC inhibition to MC (w_{MG}) is relatively large, but tends to decrease as the common GC inhibition (w_{Gc}) increases in strength (lower left panels of Fig. 2**C**). Overall the MC excitation to GCs (w_{GM}) is weaker, but there are both AMPA and NMDA synaptic currents compared to just GABA$_A$ for GC inhibition.

Capturing MC spike variance and covariance

When we consider all 3 statistics from our data, the region of parameter space (Fig. 2**C**) of viable models for the PSTH should significantly shrink to reveal a more specific structure (Barreiro et al., 2017). To determine parameters that match the spike count variance and covariance, we only

Figure 2: **Feasible region in parameter space to match firing rate (PSTH) from data. A)** Left: sub-network of 7 cells highlighting the 3 coupling strengths we systematically vary: w_{MG}, w_{GM}, and w_{Gc}. **A)** Right: transfer functions used in our model, obtained numerically from Cleland OB models (Li and Cleland, 2013, 2017) (see **Appendix: Details of the transfer funtion calculation** Figure A3C). **B)** Model of odor input from ORN (see Eq. (4)), having time-varying mean $\mu(t)$ (top) and 3 ways to assess variance of ORN inputs (bottom): i) fixed but small σ (left), ii) step increase from spontaneous to evoked state (σ **Vary in Time**, except for GC), and iii) large fixed σ. The smaller and larger σ match the spontaneous and evoked values when σ varies in time. **C)** The black dots shows the region in parameter space that is within tolerance of the PSTH from data, with a 3D view (upper right) and all 2D projections. Repeated for the 3 input noise regimes. Out of 10,000 parameter sets, there are 361, 295, and 129 (respectively for **Ci, Cii, Ciii**) that are within tolerance. **D)** Blue curves show trial-averaged MC firing rate within tolerance of data (black). The gray dashed curves show the model extremes that have the worst matches to data.

considered coupling strengths that are close to the data PSTH (black dots in Fig. 2C) because our goal is for a model to simultaneously match all spike statistics.

Figure 3: **Capturing second order spike statistics with the model.** Models that best match all three time-varying statistics (PSTH, spike count variance, and covariance) result in different levels of error depending on input statistics. **A)** Model results (blue) of spike count variance that best match data (black) from among set of coupling strengths that well-match the PSTH (Fig. 2**C, D**) for all 3 input noise regimes. The model extremes (gray dash) show the worst matches to the spike count variance among models that well-match the PSTH. **B)** Results for spike count covariance, same format as **A**. **C)** Summary of errors between model and data: small fixed σ (gray) result in overall best performance, followed by time-varying σ (tan) and large fixed σ (blue). Box plots for 10 best parameter sets: the shaded rectangle captures two middle quartile (25%–75%), median is solid line through shaded rectangle, and whiskers contain the most extreme points. Total Error and Error of Variance have the same ordering of input noise regimes; for matching the covariance alone, the time-varying σ performs best. **D)** Small region of parameter space corresponding to blue curves in **A**–**B**. **Di** shows the 3D region, **Dii** shows the 2D projections of the same points. There is some variation within a noise regime but the average values among the 10 best are very similar, see Table 1.

We assess how well the models match our data by:

$$\textbf{Total Error} \quad = \quad \textbf{Error of PSTH} + \textbf{Error of Variance} + \textbf{Error of Covariance}$$

$$\textbf{Error of PSTH} \quad = \quad \int_{-2}^{2} W_1(t)\big|r(t) - r_{data}(t)\big|\, dt \Big/ \|r_{data}(t)\| \tag{1}$$

$$\textbf{Error of Variance} \quad = \quad \int_{-2}^{2} W_2(t)\big|V(t) - V_{data}(t)\big|\, dt \Big/ \|V_{data}(t)\| \tag{2}$$

$$\textbf{Error of Covariance} \quad = \quad \int_{-2}^{2} W_3(t)\big|C(t) - C_{data}(t)\big|\, dt \Big/ \|C_{data}(t)\| \tag{3}$$

where the weights $W_{1,2,3}(t)$ account for the (population) uncertainty of the experimental data (see Fig. 6**A**), and we have normalized each integral by the magnitude of the corresponding quantity from the data $\|X_{data}(t)\| := \sqrt{\int_{-2}^{2} \left(W_j(t)X_{data}(t)\right)^2 \, dt}$.

The results are shown in Figure 3**A, B** for each of the input noise regimes. The 10 best model fits with the lowest **Total Error** are the blue curves; the parameter(s) with smallest error for both the variance and covariance are contained in the set of 10 smallest **Total Error**, for all input noise regimes. We see that the small input noise regime results in the best fit to the variance (Fig. 3**Ai**) and reasonably good fits to the covariance (Fig. 3**Bi**), but the time-varying noise fits the covariance best (Fig. 3**Bii**). The large fixed input noise regime performs the worst (Fig. 3**Aiii, Biii**). Even within the parameter sets that capture the data PSTH well, the variance and covariance can be quite different from the data (gray-dashed curves correspond to the largest error).

The errors between the model with the 10 best parameters and experimental data are shown in boxplots in Figure 3C for each of the input noise regimes. For both **Total Error** and **Error of Variance**, the small fixed input noise performs best (black), followed by time-varying input noise (tan), and large fixed input noise (blue). But for **Error of Covariance**, the time-varying input noise performs best, followed by small fixed input noise.

From Figure 3**D**, we see in all ORN input noise regimes that independent GC inhibition to distinct MC has to be large, while MC excitation is modest, and common inhibition to both MC has to be relatively weak. Although there is some variation in coupling strengths across the different noise regimes, the averages are remarkably similar (Table 1).

Table 1: The average coupling strengths among the 10 best parameter sets in Figure 3D.

Noise regime	Ind. GC Inh. (w_{MG})	Shared GC Inh. (w_{Gc})	MC Excit. (w_{GM})
Smaller σ	6.67	1.57	3.79
σ Vary in Time	7.35	1.85	3.31
Larger σ	7.46	1.64	3.63

Small fixed ORN input noise performs best

With a specific set of three ORN input noise values, we thus far have performed a robust search through the parameter space of coupling strengths to determine the profile of input noise that minimizes error with various statistics of our data. Next we addressed how robust these observations are by considering a range of input noise values; however, reproducing everything in the prior sections would require impractical amounts of computation. To circumvent this, we use a previously developed mathematical reduction (Barreiro and Ly, 2017a) (also see **Materials and Methods: Steady-sate approximation of statistics of firing rate model**) to efficiently approximate the steady-state spiking statistics.

We used the reduction to compute statistics for a set of networks, in which we allowed the evoked input noise variance (σ_{large}) to take a range of linearly spaced values, i.e., $\hat{\sigma}_{large} = (j - 1)/10 * \sigma_{small} + (1 - (j-1)/10) * \sigma_{large}$, $j = 1, 2, \ldots, 10$ from left to right; see insets in Fig. 4). For a fair comparison of the 3 input noise regimes, we use a large random sample of the coupling strengths that maintain the same 3D structure (Fig. 6**B**), and calculated the spike statistics to get a large set of error values.

Figure 4 confirms that our results are robust. Testing a total of 10 triple input noise sets, we find that small fixed noise yields the lowest Total Error, followed by time-varying input noise and large fixed noise (Fig. 4**A**). The results for Error of Variance similarly hold (Fig. 4**B**). Finally, if matching the spike count covariance is desired, the time-varying input noise gives the smallest error (Fig. 4**C**).

Although our firing rate network model has only 7 cells and identical marginal MC spiking statistics, it is still sophisticated enough to exhibit odor-evoked decreases in variability (see **Appendix: Firing rate model can exhibit evoked decreases in variability**, Fig. A2). That is, even qualitatively matching our experimental data (i.e., increases in mean and variability at stimulus onset) requires parameter tuning in this model.

Model prediction confirmed with awake mice

A natural prediction from our modeling results is that the ORN input noise does not have to increase in the evoked state to observe odor-evoked increases in MC spiking variability. Perhaps the OB circuit itself can shape the increases in variability rather than inheriting increased variability from ORN. We test this model prediction with a publicly available dataset (Bolding and Franks, 2018b) originally collected for Bolding and Franks (2018a). The Franks lab performed direct optical MC stimulation with 1 s light pulses in awake and freely breathing mice (Fig. 7 in (Bolding and Franks, 2018a)). By circumventing the normal sensory pathway with direct MC stimulation, the ORN input statistics (i.e., variance) presumably remains similar during light-evoked MC spiking.

Figure 4: **Probing other ORN input noise values.** Testing other ORN input noise values to show robustness of prior results for matching time-varying data statistics, but using a steady-state approximation for computational feasibility. Each triplet of input noise values has a small fixed σ (black), time-varying with a step (tan), and large fixed σ (blue); like before, the time-varying values are set to the small values in the spontaneous state and large values in the evoked state. Results for Total Error **(A)**, Error of Variance **(B)**, and Error of Covariance **(C)** are all consistent with what we observed before (Fig. 3C), with the differences converging as the noise values approach each other. We test 10 total triplets of input noise sets, using a random sample of coupling strengths (Fig. 6C) that cover the best parameter values (Fig. 3D), and a random set of scaling factors (Fig. 6C) to approximate spike statistics in 100 ms windows. The whiskers and inter-quartiles in the box plots capture the variation of errors over the random samples of coupling strengths and scaling factors.

²³⁴ Hence our model prediction would bear out if the spike count variance and covariance increase in ²³⁵ the evoked state.

²³⁶   Figure 5 indeed shows evoked increases in OB spike count variance (Fig. 5B) and covariance ²³⁷ for several light intensities (same ones used in Bolding and Franks (2018a)). The OB spike co-²³⁸ variance clearly increases after light stimulation (Fig. 5C), although the increases is brief for only ²³⁹ about 300 ms. The increase in spike variance is less noticeable by visual inspection (Fig. 5B), but ²⁴⁰ two-sample t-tests assuming unequal variance shows that the increase from spontaneous to evoked ²⁴¹ is statistically significant with $p-$values of $0.0296, 5.5 \times 10^{-4}, 1.2 \times 10^{-4}$ for the three respec-²⁴² tive light intensities (5, 10, 20 mW/mm^2). We specifically determined that the mean spontaneous ²⁴³ values are less than the mean evoked values.

Figure 5: **Data from the Franks lab shows evoked increases in MC spiking variability** Using publicly available data (Bolding and Franks, 2018b), we verify our model prediction that ORN input noise does not have to increase to observe evoked increases in MC spiking variance and covariance. We use the same 3 light intensities in Bolding and Franks (2018a). **A)** shows the population average PSTH, **B)** the spike variance, and **C)** the spike count covariance. The PSTH increases in the evoked state, and the spike count covariance clearly increases for a brief period after light stimulation. Evoked increases of spike count variance is harder to see, but two-sample t-test assuming unequal variance shows statistically significant increases for all light intensities ($p-\text{value} = 0.0296, 5.5 \times 10^{-4}, 1.2 \times 10^{-4}$, respectively). Spike counts were calculated in $100\,\text{ms}$ half-overlapping windows; there were 77 total cells and 812 total pairs, with 10 trials for each light intensity. Shaded regions show relative population heterogeneity after trial-averaging: $\mu \pm 0.05\text{std}$ (standard deviation across population/pairs after trial-averaging).

Discussion

Using both experimental data from multi-unit recordings of rat OB and a biophysically-inspired model, we studied the mechanisms by which the olfactory system exhibits stimulus-induced increases in spiking variability and co-variability in putative MCs. This phenomenon is relatively understudied, in contrast to the stimulus-induced decreases in (co-)variability that have been widely reported in many cortical and motor regions (Churchland et al., 2010; Doiron et al., 2016). Although there are reports of odor-evoked increases in (co-)variability of OB cells in insects and fish (Kazama and Wilson, 2009; Wanner and Friedrich, 2020), these systems differ from the mammalian system in important ways. Thus, the mechanisms for how the mammalian ORN–OB system accomplishes a stimulus-induced increase in spike variability is unknown. We have described in detail how both presynaptic ORN inputs and coupling within the OB are structured to capture the time-varying statistics of our data. We also verify our model prediction with data from awake mice (Bolding and Franks, 2018a,b). Thus evoked increases in spiking variability in the OB seems to be a core principle in many species.

Recent recordings in the OB of developing zebrafish show time-varying statistics of correlation that are similar to what we observed in rat: the correlation of activity measured by calcium imaging increased suddenly in the evoked state, followed by a slow decay to a level that remained above the spontaneous values (Wanner and Friedrich, 2020). These authors explained correlation of MC cells by analyzing detailed connectivity structures of interneurons and MC using electron microscopy, neglecting GC because they have an insignificant role at this stage of zebrafish development. Kazama and Wilson (2009) also showed that the correlation of projection neurons (**PN**, analogous to MC) in *Drosophila* OB increased in the evoked state compared to spontaneous. This was observed for pairs PN from the same or different glomeruli, and independent of whether ORN inputs were shared or not. They attributed the increase in PN spike correlation primarily to common inputs from ORN rather than lateral connections in the glomeruli layer. We focussed on both ORN inputs and MC–GC coupling strengths because they likely have significant affects on MC spike dynamics in mammals (Cang and Isaacson, 2003; Galán et al., 2006; Giridhar et al., 2011).

Our model prediction that evoked increases in MC spike count variability can occur without increases in the noise of ORN input (i.e., attribute it to the OB circuit) was verified with another dataset (Bolding and Franks, 2018a,b). To our knowledge, whether ORN spiking variance actual stays constant in natural conditions has not been directly verified. There have been many ORN recordings in mammals (Spors et al., 2006), *Drosophila* (Nagel and Wilson, 2011; Kazama and Wilson, 2009) and *Locust* (Raman et al., 2010), but the trial-to-trial variability of ORN spiking in both spontaneous and evoked states has not been reported or is not readily available. Duchamp-Viret et al. (2005) characterized the noisy spiking of ORN in tracheotomized and breathing rats but only in the spontaneous state. The data in Figure 1 of Nagel and Wilson (2011) has shaded

gray regions to show trial-variability of ORN spiking, and indeed by visual inspection the trial-variability appears to be constant in spontaneous and evoked states for several different odors (but not all) in *Drosophila*, consistent with our model prediction.

ORN responses are known to have odor-evoked increases in spatial-temporal variability, for example in locusts (Raman et al., 2010) and mammals (Spors et al., 2006). Calcium imaging of ORN inputs to OB in anesthetized breathing rats *in vivo* has shown that across glomeruli, the calcium activity with the same odor we use (ethyl butyrate) can vary dramatically compared to the spontaneous state (Spors et al., 2006). Although our OB model consists of only 2 glomeruli and is far from a realistic spatial model, these data appear to refute our model prediction that ORN input variance have similar values in both states. However, Spors et al. (2006) showed that higher odor concentrations may result in less variability across glomeruli (their Fig. 2B). Furthermore, the important effects of breath cycles and active sniffing (Wachowiak, 2011) are not in our data because rats in our experiments were anesthetized. Active sniffing and breathing may result in larger evoked ORN input variance because some glomeruli are more locked to the breathing cycle while others responded tonically (Spors et al., 2006). We are unaware of any experiments that refute or validate our model predictions about coupling strengths within OB, but relatively strong GC inhibition is well-known in the mammalian OB (Cang and Isaacson, 2003; Schoppa and Urban, 2003; Boyd et al., 2012; Markopoulos et al., 2012; Burton and Urban, 2015). Nevertheless, we hope our model predictions about OB synaptic strengths will be insightful in further mechanistic studies of OB spiking modulation.

We conclude by listing several factors which could in principle affect our results, but which we were not able to consider here. We did not account for different odors in our data, and the OB model did not explicitly model feedback from piriform or other cortical areas even though there is evidence of odor-specific feedback from piriform that can decorrelate OB responses (Otazu et al., 2015). Different odors elicit different spatiotemporal OB responses that would quantitatively change our results, and whether our results would qualitatively hold is an open question, but at the very least, evoked increases in spike variability in our data is consistent with others (Wanner and Friedrich (2020); Kazama and Wilson (2009), and Bolding and Franks (2018a) as verified by us). Also, we did not investigate how changes in the transfer functions (Fig. 2A) would alter the population spike count statistics, even though it is known to shape spike statistics (de la Rocha et al., 2007; Tetzlaff et al., 2012; Barreiro and Ly, 2017b, 2018). We use a network model that does not incorporate heterogeneity of MCs or other cells, so the model cannot simultaneously match the 3 statistics we focus on *as well as* the Fano factor and spike count correlation. Fano factor and correlation in our data are not simply scaled by the population mean and variances (as they would be in a homogeneous network model) because of the large degree of heterogeneity. All cells/pairs do not simply increase or decrease uniformly in the evoked state (Fig. 1D–F), and the modulation of (co-)variability could strongly depend on the underlying function or selectivity of individual

cells/pairs (Ruff and Cohen, 2014; Moreno-Bote et al., 2014). Finally, limiting our investigation to a single odor means that we did not explicitly consider information or coding: much theoretical work has been devoted to whether relatively larger co-variability is better (da Silveira and Berry II, 2014) or worse (Ecker et al., 2011) for coding, with a recent study showing this also depends on population size (Bartolo et al., 2020). This intersects with the question of heterogeneity, because the relationship between collective population statistics (i.e., not just population-averaged statistics) and efficient/accurate neural coding is complicated (Kohn et al., 2016). We hope to consider each of these important issues in future work.

Methods and Materials

See https://github.com/chengly70/OB for MATLAB code implementing all models in this paper, as well as code to display the Franks' lab data.

Firing rate model details

Our firing rate model contains many features of a multi-compartment biophysical OB model (Li and Cleland, 2013, 2017), including the synaptic variables, transfer functions, and membrane time constants. But this model is also simple enough to enable fast simulation times to assess how various attributes modulate firing statistics; see Kanashiro et al. (2017) for similar use of firing rate models. The voltage, intrinsic currents and all associated dynamic variables are modeled with a single state variable x_j that is a phenomenological representation of the cell's activity, which is mapped to a firing rate via a monotonically increasing function F_j.

$$\tau_j \frac{dx_j}{dt} = -x_j + \underbrace{\mu_j(t) + \sigma_j(t)\xi_j(t)}_{\text{Odor Inputs}} + \sum_k w_{j,k} G_k(r_k(t), t) \tag{4}$$

$$r_j(t) = F_j(x_j) \tag{5}$$

Noise terms $\xi_j(t)$ represent inputs, are white in time (i.e. $\langle \xi_j(t) \rangle = 0$, $\langle \xi_j(t)\xi_j(t') \rangle = \delta(t - t')$), and are pairwise correlated (Galán et al., 2006; Marella and Ermentrout, 2010; Litwin-Kumar et al., 2011). Input noise terms are correlated between an MC and PGC pair within a glomerulus because they have input connections from the same ORN cells ($\langle \xi_j(t)\xi_k(t) \rangle = 0.1$), correlated between the two MCs because of common ORN inputs (Kazama and Wilson, 2009) ($\langle \xi_j(t)\xi_k(t) \rangle = 0.05$), and between the GCs ($\langle \xi_j(t)\xi_k(t) \rangle = 0.05$) because they are known to synchronize (Li and Cleland, 2017; Marella and Ermentrout, 2010). The correlation values are small (0.05 or 0.1), with the largest value between PGC and MC (0.1) corresponding to common ORN inputs (Kazama and Wilson, 2009). All other pairs of cells have no input correlation.

The transfer functions $F_j(\cdot)$ (Fig. 2**A**) we use is numerically calculated from our implementation of the Cleland models (Li and Cleland, 2013, 2017). For simplicity, we choose static functions F_j for each cell type that is similar to several computed transfer functions, see **Appendix: Details of the transfer function calculation** Figure A3C. Any inputs outside of the ranges in Figure 2**A** are capped, so inputs below the input range give 0 firing and inputs larger than the maximum input range result in the maximum firing rate.

We use the synapse equations from the Cleland models but replace the presynaptic voltage with a function of firing rate $f(r(t))$ since firing rates are what drive average synaptic activity:

$$G_k(t) = \alpha \int_{-\infty}^t f_k(r_k(t')) \exp\left(-\beta(t-t') - \alpha \int_{t'}^t f_k(r_k(s))\,ds\right) dt' \tag{6}$$

Here the strengths of the synapse $w_{j,k}$ is a signed-quantity: < 0 for GABA$_A$ and > 0 for both AMPA and NMDA; but all plots of coupling strengths (w_{MG}, w_{Gc}) show the absolute value. The parameters α, β are obtained directly from the biophysical model (Li and Cleland, 2013, 2017) and vary with synapse type (i.e., α_{NMDA}). Here we choose: $f_k(r_k(t')) = r_k(t')/\bar{r}$ where $\bar{r} = \max r$. The synaptic filter $(s(t) := \frac{G_k}{w_{j,k}})$ satisfies the synapse equations used in the biophysical model (20) but with $a_{on}(v_{pre})$ replaced with $f(r(t))$. The steady state of G_k is:

$$\lim_{t\to\infty} G_k(t) = \frac{\alpha f_k(r_k)}{\beta + \alpha f_k(r_k)} = \frac{\alpha r_k/\bar{r}}{\beta + \alpha r_k/\bar{r}} \tag{7}$$

The spike count variance/covariance is modeled by the variance/covariance of the firing rate integrated over a time window T. That is, we define the random variable

$$R_j(t) = \int_{t-T/2}^{t+T/2} r_j(t')\,dt' \tag{8}$$

and use $Var(R_j)$ and $Cov(R_j, R_k)$ as proxies for the spike count variance and covariance respectively. For this study, we focused on a time window of $T = 100$ ms.

Each network was simulated using the explicit Euler-Maruyama method for stochastic differential equations with a resolution of 1 ms, for a simulation time of 4 seconds (2 seconds before stimulus onset and 2 seconds afterwards). For each set of parameter values, we simulated 100,000 realizations in order to accurately resolve the mean value of second-order statistics such as spike count covariance.

Coupling strength parameter space and error calculations

For each ORN noise regime, we calculate model PSTH $r(t)$ for 10,000 points that fill the 3D volume of parameter space generated by Halton sequence sampling (Kocis and Whiten, 1997). To account for uncertainty about a population-averaged statistic, we weight the difference between the model and data PSTH $r_{data}(t)$ with $W_1(t)$, which is a monotonically decreasing function of the standard deviation (across cells) of the data PSTH (Fig. 6A). Error calculations for the spike count variance and covariance are computed analogously.

Steady-state approximation of statistics of firing rate model

We derived a steady-state approximation of the complete set of 1st and 2nd order statistics of a general firing rate model (i.e., equations (4)–(5)), with a weak coupling assumption in another paper (Barreiro and Ly, 2017a). The approximation consists of a set of transcendental equations that are to be solved self-consistently; this procedure is more computationally efficient than performing Monte Carlo simulations and averaging over realizations. Here we augment that method to allow for larger coupling strengths that are necessary for capturing the spike statistics of our experimental data as follows. We use a new set of mean parameter values that is shifted from μ_j, resulting in a new operating point where the network coupling is effectively weak. The shift in the μ_j parameters is accomplished in a principled manner and is different depending on the coupling strengths $w_{j,k}$ and μ_j's. We replace μ_j with:

$$\tilde{\mu}_j = \mu_j - \mu_j^1 \tag{9}$$

where μ_j^1 is the solution to:

$$\mu_j^1 = \mu_j + \sum_k w_{j,k} G_k(F_k(\mu_k^1)) \tag{10}$$

Then we find a solution to the following set of equations:

$$\tilde{\mu}(j) = \tilde{\mu}_j + \sum_k w_{j,k} \mathcal{E}_1(k) \tag{11}$$

$$\sigma^2(j)\tau_j = \frac{\sigma_j^2}{2} + \sigma_j \sum_k w_{j,k} \mathcal{N}_{\mathcal{F}}(j,k) + \frac{1}{2} \sum_k w_{j,k}^2 \mathcal{V}(k) + \sum_{k \neq l} w_{j,k} w_{j,l} \mathcal{C}_{\mathcal{V}}(k,l) \tag{12}$$

$$Cov(j,k)\frac{\tau_j+\tau_k}{2} = \frac{1}{2}c_{j,k}\sigma_j\sigma_k + \frac{1}{2}\sigma_j \sum_l w_{k,l}\mathcal{N}_{\mathcal{F}}(j,l)$$

$$+\frac{1}{2}\sigma_k \sum_l w_{j,l}\mathcal{N}_{\mathcal{F}}(k,l) + \frac{1}{2}\sum_{l_1,l_2} w_{j,l_1} w_{k,l_2} \mathcal{C}_{\mathcal{V}}(l_1,l_2). \tag{13}$$

where the definitions of $\mathcal{E}_1(k), \mathcal{N}_{\mathcal{F}}(j,k), \mathcal{V}(k), \mathcal{C}_{\mathcal{V}}(j,k)$ are in Table 2.

Figure 6: **Details for error calculations. A)** Time-varying weights for each of the spike statistics. We choose a monotonically decreasing function of the standard deviation of the data (across population / pairs). The weight of the PSTH is $W_1(t) = 1 + 0.5 \tanh\left((std(t) - 10)/5\right)$, the spike count variance is $W_2(T) = 1 + 0.5 \tanh\left((std(T) - 1)/5\right)$, covariance is: $W_3(T) = 1 + 0.5 \tanh\left((std(T) - 0.2)/5\right)$. **B)** Black dots are random samples of coupling strength parameters preserving relationships of parameters that best match time-varying statistics of data for the 3 input noise regimes (10 best parameters in Fig. 3D are in red, tan, and blue), used for results in Fig. 4. **C)** Scaling factors from simulations (red, tan and blue circles, again from 10 best parameters) relating instantaneous spike count variance and covariance to spike statistics in 100 ms time window. Horizontal axis shows ratio of spike count variance in instantaneous window (dt) to 100 ms time window, vertical shows same ratio for covariance. Each parameter set has 2 ratios: time-averaged ratios in spontaneous state, and time-averaged ratios in the evoked state. We use a single large random sample of ratios (black dots are same for all 3 panels) preserving the dependencies between the variance and covariance ratios; used for results in Fig. 4. Code to generate all random samples on GitHub.

389   The final step in our approximation is to shift the mean back: $\mu(j) = \tilde{\mu}(j) + \mu_j^1$ and solve for

Table 2: Definitions of variables for the steady-state approximation of statistics (11)–(13). Whenever $j = k$ in the double integrals (e.g., in $\mathcal{N}_\mathcal{F}, \mathcal{S}$), the bivariate normal distribution $\varrho_{j,k}$ is replaced with the standard normal distribution ϱ_1. Note that order of the arguments matters in $\mathcal{N}_\mathcal{F}$: $\mathcal{N}_\mathcal{F}(j,k) \neq \mathcal{N}_\mathcal{F}(k,j)$ in general; all of these quantities depend on $\tilde{\mu}(\cdot), \sigma(\cdot)$. For notational convenience, we define $H_k(x) := G_k(F_k(x))$.

Abbreviation	Definition
$\mathcal{E}_1(k)$	$\displaystyle\int H_k(\sigma(k)y + \tilde{\mu}(k))\varrho_1(y)\,dy$
$\mathcal{E}_2(k)$	$\displaystyle\int H_k^2(\sigma(k)y + \tilde{\mu}(k))\,\varrho_1(y)\,dy$
$\mathcal{V}(k)$	$\displaystyle\int H_k^2(\sigma(k)y + \tilde{\mu}(k))\,\varrho_1(y)\,dy - \left(\int H_k(\sigma(k)y + \tilde{\mu}(k))\,\varrho_1(y)\,dy\right)^2$
	$= \mathcal{E}_2(k) - [\mathcal{E}_1(k)]^2$
$\mathcal{N}_\mathcal{F}(j,k)$	$\displaystyle\iint H_k(\sigma(k)y_1 + \tilde{\mu}(k))\frac{y_2}{\sqrt{2}}\varrho_{j,k}(y_1, y_2)\,dy_1 dy_2,\,\text{if } j \neq k$
	$\displaystyle\int H_j(\sigma(j)y + \tilde{\mu}(j))\frac{y}{\sqrt{2}}\varrho_1(y)\,dy,\,\text{if } j = k$
$\mathcal{S}(j,k)$	$\displaystyle\iint H_j(\sigma(j)y_1 + \tilde{\mu}(j))H_k(\sigma(k)y_2 + \tilde{\mu}(k))\varrho_{j,k}(y_1, y_2)\,dy_1 dy_2$
$\mathcal{C}_\mathcal{V}(j,k)$	$\mathcal{S}(j,k) - \mathcal{E}_1(j)\mathcal{E}_1(k)$

the firing statistics:

$$\mu(j) = \tilde{\mu}(j) + \mu_j^1 \tag{14}$$

$$r_j = \int F_j(\sigma(j) + \mu(j)x)\varrho_1(x)\,dx \tag{15}$$

$$Cov(r_j, r_k) = \iint F_j(\sigma(j) + \mu(j)x_1)F_k(\sigma(k) + \mu(k)x_2)\varrho_{j,k}(x_1, x_2)\,dx_1 dx_2 - r_j r_k \tag{16}$$

Where ϱ_1 is the standard normal distribution and $\varrho_{j,k}$ is a standard normal bivariate distribution with correlation $\frac{Cov(j,k)}{\sigma(j)\sigma(k)}$ from Eqs. (12)–(13).

We can compare the firing rate r_j (Eq. (15)) directly to the corresponding value from our experimental data, but the approximate variance and covariance (Eq. (16)) are computed for instantaneous time rather than the 100 ms time windows. Unlike firing rate, second-order statistics can depend on the time window used for measurement in nontrivial ways (Gabbiani and Koch, 1998). To model spiking activity in 100 ms windows, we hypothesized that the relationship between statistics measured at instantaneous time and statistics measured in 100 ms bins could be modeled by a scaling factor. We used results from Monte Carlo simulations of the firing rate model (Eqs. (4)–(8)) to find this scaling factor as described below.

We computed the ratio between variances and covariances from the steady-state approximation (variances and covariances, in either the spontaneous or evoked state), and the corresponding statistic in each 100 ms bin from the 30 best parameter simulations (10 each for small σ, varying

σ, and large σ). For example, $Var(\mathcal{R}_j)/Var(R_j)$ gives us a variance ratio for cell j, where R_j are spike counts in 100 ms windows (Eq. (8)) and $\mathcal{R}_j$ are spike counts in 1 ms windows ($\mathcal{R}_j = r_j dt$, see Eq. (5)). Similarly $Cov(\mathcal{R}_j, \mathcal{R}_k)/Cov(R_j, R_k)$ gives us a covariance ratio for cells j, k. This yielded a range of values in the vicinity of 2±0.5, with variation between simulations, across time, and across statistics (variance vs. covariance, see Fig. 6C). However, we did notice a correlation between ratios for variance and covariance involving the same cells; therefore we grouped statistics into pairs, e.g. $Var(\mathcal{R}_j)/Var(R_j), Cov(\mathcal{R}_j, \mathcal{R}_k)/Cov(R_j, R_k))$. We show these paired ratios as black dots in each panel of Figure 6C; superimposed we show the time-averaged values during spontaneous and evoked activity in red, tan and blue, respectively.

These panels demonstrate several points about the ratio between instantaneous and 100 ms time bin statistics: first, for a given cell or cell pair the ratios for variance and covariance are correlated; second, the ratios vary even within a single simulation; third, all of the ratios have the same order of magnitude and lie within 25% of an average value of 2. Thus, when we measure error for the networks for which only the instantaneous rates have been computed (i.e. Figure 4), we will actually use a large number of scaling factor pairs to produce a distribution of error values. Each is plausible, given the uncertainty in our "fudge factor". Since the ratios are fairly similar, we use the same random sample of 500 ratios (black dots) for all noise regimes; see 'get_RatiosRand.m' at https://github.com/chengly70/OB.

Finally, we generate a random set of 1000 coupling strengths for each noise regime (black dots in Fig. 6B) as follows. For a given noise regime, we take the 10 best sets (red, tan or blue in Fig. 6**Bi**–**Biii**), calculate the mean of (w_{GM}, w_{MG}, w_{Gc}), denoted by $\vec{\mu}$ and their 3×3 covariance matrix $\mathbf{Cw}$. We then calculate a Cholesky decomposition $\mathbf{Cw} = \mathbf{LL}'$ ($\mathbf{L}$ is a 3×3 lower triangular matrix), and generate 990 multivariate normal random variables:

$$\vec{\mu} + 0.7\mathbf{L}\vec{Z}$$

where $\vec{Z}$ is an instance of a 3×1 uncorrelated standard normal. The samples have the same mean (w_{GM}, w_{MG}, w_{Gc}) (up to finite sampling error) and similar covariance (compare black dots with red, tan and blue in Fig. 6**B**). The 990 random samples are combined with the 10 best coupling strengths for a total of 1000 sets; see 'get_GsRand.m' at https://github.com/chengly70/OB.

So for each noise regime (i.e., each error bar in Fig. 4, there is a large set of errors between experimental data and model that stem from both the random sample of coupling strengths (1000 sets, black dots) and random ratios (500 sets, black dots). With a steady-state approximation, we must augment the error calculation; we use the same weights as before (see Fig. 6**A**) except we only consider a 1 s period in the spontaneous state $(-2 \le t \le -1)$ and a 1 s period in the evoked state $(1 \le t \le 2)$.

Electrophysiological recordings

All procedures were carried out in accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health and approved by University of Arkansas Institutional Animal Care and Use Committee (protocol #14049). Data were collected from 11 adult male rats (240-427 g; *Rattus Norvegicus*, Sprague-Dawley outbred, Harlan Laboratories, TX, USA) housed in an environment of controlled humidity (60%) and temperature (23°C) with 12 h light-dark cycles. The experiments were performed in the light phase.

Surgical preparations. Anesthesia was induced with isoflurane inhalation and maintained with urethane (1.5 g/kg body weight (**bw**) dissolved in saline, intraperitoneal injection (**ip**)). Dexamethasone (2 mg/kg bw, ip) and atropine sulphate (0.4 mg/kg bw, ip) were administered before performing surgical procedures. Throughout surgery and electrophysiological recordings, core body temperature was maintained at 37°C with a thermostatically controlled heating pad. To isolate the effects of olfactory stimulation from breath-related effects, we performed a double tracheotomy surgery as described previously (Gautam and Verhagen, 2012). A Teflon tube (OD 2.1 mm, upper tracheotomy tube) was inserted 10 mm into the nasopharynx through the rostral end of the tracheal cut. Another Teflon tube (OD 2.3 mm, lower tracheotomy tube) was inserted into the caudal end of the tracheal cut to allow breathing, with the breath bypassing the nasal cavity. Both tubes were fixed and sealed to the tissues using surgical thread. Local anesthetic (2% Lidocaine) was applied at all pressure points and incisions. Subsequently, a craniotomy was performed on the dorsal surface of the skull over the right olfactory bulb (2 mm x 2 mm, centered 8.5 mm rostral to bregma and 1.5 mm lateral from midline).

Olfactory Stimulation. A Teflon tube was inserted into the right nostril and the left nostril was sealed by suturing. Odorized air was delivered for 1 s in duration at 1 minute intervals. The odorant was Ethyl Butyrate (**EB**, saturated vapor). We note that the full experimental data set included additional odors and retronasal stimulation, but here we consider only orthonasal stimulation with EB.

Electrophysiology. A 32-channel microelectrode array (**MEA**, A4x2tet, NeuroNexus, MI, USA) was inserted 400 μm deep from dorsal surface of OB targeting tufted and mitral cell populations. The MEA probe consisted of 4 shanks (diameter: 15 μm, inter-shank spacing: 200 μm), each with eight iridium recording sites arranged in two tetrode groups near the shank tip (inter-tetrode spacing: 150 μm, within tetrode spacing 25 μm). Simultaneous with the OB recordings, we recorded from a second MEA placed in anterior piriform cortex. Voltage was measured with respect to an AgCl ground pellet placed in the saline-soaked gel foams covering the exposed brain surface around the inserted MEAs. Voltages were digitized with 30 kHz sample rate (Cereplex + Cerebus, Blackrock Microsystems, UT, USA). Recordings were band-pass filtered between 300 and

3000 Hz and semiautomatic spike sorting was performed using Klustakwik software, which is well suited to the type of electrode arrays used here (Rossant et al., 2016).

Acknowledgments

We thank Brent Doiron for enlightening conversations about this research. We thank the Southern Methodist University (SMU) Center for Research Computing for providing computational resources.

References

Balu, R., P. Larimer, and B. W. Strowbridge (2004). Phasic stimuli evoke precisely timed spikes in intermittently discharging mitral cells. Journal of neurophysiology 92(2), 743–753.

Barreiro, A., S. Gautam, W. Shew, and C. Ly (2017). A theoretical framework for analyzing coupled neuronal networks: Application to the olfactory system. PLoS Computational Biology 13, e1005780.

Barreiro, A. and C. Ly (2017a). Practical approximation method for firing-rate models of coupled neural networks with correlated inputs. Physical Review E 96, 022413.

Barreiro, A. and C. Ly (2017b). When do correlations increase with firing rates in recurrent networks? PLoS Computational Biology 13, e1005506.

Barreiro, A. and C. Ly (2018). Investigating the correlation-firing rate relationship in heterogeneous recurrent networks. Journal of Mathematical Neuroscience 8, 8.

Barreiro, A., E. Thilo, and E. Shea-Brown (2012). A-current and type i/type ii transition determine collective spiking from common input. Journal of Neurophysiology 108(6), 1631.

Barreiro, A. K., E. Shea-Brown, and E. Thilo (2010). Time scales of spike-train correlation for neural oscillators with common drive. Physical Review E 81, 011916.

Bartolo, R., R. C. Saunders, A. R. Mitz, and B. B. Averbeck (2020). Information-limiting correlations in large neural populations. Journal of Neuroscience 40(8), 1668–1678.

Bolding, K. A. and K. M. Franks (2018a). Recurrent cortical circuits implement concentration-invariant odor coding. Science 361(6407).

Bolding, K. A. and K. M. Franks (2018b). Simultaneous extracellular recordings from mice olfactory bulb (OB) and piriform cortex (PCx) and respiration data in response to odor stimuli and optogenetic stimulation of OB. CRCNS.org Date accessed: January 2021(https://dx.doi.org/10.6080/K00C4SZB).

Boyd, A. M., J. F. Sturgill, C. Poo, and J. S. Isaacson (2012). Cortical feedback control of olfactory bulb circuits. Neuron 76(6), 1161–1174.

Burton, S. D. and N. N. Urban (2015). Rapid feedforward inhibition and asynchronous excitation regulate granule cell activity in the mammalian main olfactory bulb. The Journal of Neuroscience 35(42), 14103–14122.

Cang, J. and J. S. Isaacson (2003). In vivo whole-cell recording of odor-evoked synaptic transmission in the rat olfactory bulb. Journal of Neuroscience 23(10), 4108–4116.

Churchland, M., B. Yu, et al. (2010). Stimulus onset quenches neural variability: a widespread cortical phenomenon. Nature Neuroscience 13, 369–378.

da Silveira, R. A. and M. J. Berry II (2014). High-fidelity coding with correlated neurons. PLoS Computational Biology 10(11), e1003970.

de la Rocha, J., B. Doiron, E. Shea-Brown, K. Josić, and A. Reyes (2007). Correlation between neural spike trains increases with firing rate. Nature 448, 802–806.

Desmaisons, D., J.-D. Vincent, and P.-M. Lledo (1999). Control of action potential timing by intrinsic subthreshold oscillations in olfactory bulb output neurons. Journal of Neuroscience 19(24), 10727–10737.

Doiron, B., A. Litwin-Kumar, R. Rosenbaum, G. Ocker, and K. Josić (2016). The mechanics of state-dependent neural correlations. Nature Neuroscience 19(3), 383–393.

Duchamp-Viret, P., L. Kostal, M. Chaput, P. Lánsky, and J.-P. Rospars (2005). Patterns of spontaneous activity in single rat olfactory receptor neurons are different in normally breathing and tracheotomized animals. Journal of Neurobiology 65(2), 97–114.

Ecker, A., P. Berens, A. Tolias, and M. Bethge (2011). The effect of noise correlations in populations of diversely tuned neurons. The Journal of Neuroscience 31(40), 14272–14283.

Furudono, Y., G. Cruz, and G. Lowe (2013). Glomerular input patterns in the mouse olfactory bulb evoked by retronasal odor stimuli. BMC neuroscience 14(1), 45.

Gabbiani, F. and C. Koch (1998). Principles of spike train analysis. In C. Koch and I. Segev (Eds.), Methods in Neuronal Modeling: From Ions to Networks, pp. 313–360. MIT Press.

Galán, R., N. Fourcaud-Trocmé, B. Ermentrout, and N. Urban (2006). Correlation-induced synchronization of oscillations in olfactory bulb neurons. The Journal of Neuroscience 26, 3646–3655.

Gautam, S. H. and J. V. Verhagen (2012). Retronasal odor representations in the dorsal olfactory bulb of rats. The Journal of Neuroscience 32(23), 7949–7959.

Giridhar, S., B. Doiron, and N. N. Urban (2011). Timescale-dependent shaping of correlation by olfactory bulb lateral inhibition. Proceedings of the National Academy of Sciences 108(14), 5843–5848.

Kanashiro, T., G. K. Ocker, M. R. Cohen, and B. Doiron (2017). Attentional modulation of neuronal variability in circuit models of cortex. Elife 6, e23978.

Kazama, H. and R. I. Wilson (2009). Origins of correlated activity in an olfactory circuit. Nature neuroscience 12(9), 1136.

Kocis, L. and W. J. Whiten (1997). Computational investigations of low-discrepancy sequences. ACM Transactions on Mathematical Software (TOMS) 23(2), 266–294.

Kohn, A., R. Coen-Cagli, I. Kanitscheider, and A. Pouget (2016). Correlations and neuronal population information. Annual review of neuroscience 39(0).

Li, G. and T. A. Cleland (2013). A two-layer biophysical model of cholinergic neuromodulation in olfactory bulb. Journal of Neuroscience 33(7), 3037–3058.

Li, G. and T. A. Cleland (2017). A coupled-oscillator model of olfactory bulb gamma oscillations. PLoS computational biology 13(11), e1005760.

Litwin-Kumar, A., A. Oswald, N. Urban, and B. Doiron (2011). Balanced synaptic input shapes the correlation between neural spike trains. PLoS Computational Biology 7, e1002305.

Ly, C., J. Middleton, and B. Doiron (2012). Cellular and circuit mechanisms maintain low spike covariability and enhance population coding in somatosensory cortex. Frontiers in Computational Neuroscience 6, 1–26.

Marella, S. and B. Ermentrout (2010). Amplification of asynchronous inhibition-mediated synchronization by feedback in recurrent networks. PLoS Computational Biology 6(2), e1000679.

Markopoulos, F., D. Rokni, D. H. Gire, and V. N. Murthy (2012). Functional properties of cortical feedback projections to the olfactory bulb. Neuron 76(6), 1175–1188.

McQuiston, A. and L. Katz (2001). Electrophysiology of interneurons in the glomerular layer of the rat olfactory bulb. Journal of neurophysiology 86(4), 1899–1907.

Middleton, J., C. Omar, and B. D. D. Simons (2012). Neural correlation is stimulus modulated by feedforward inhibitory circuitry. The Journal of Neuroscience 32, 506–518.

Miura, K., Z. Mainen, and N. Uchida (2012). Odor representations in olfactory cortex: distributed rate coding and decorrelated population activity. Neuron 74, 1087–1098.

Moreno-Bote, R., J. Beck, I. Kanitscheider, X. Pitkow, P. Latham, and A. Pouget (2014). Information-limiting correlations. Nature neuroscience 17(10), 1410–1417.

Nagel, K. I. and R. I. Wilson (2011). Biophysical mechanisms underlying olfactory receptor neuron dynamics. Nature neuroscience 14(2), 208–216.

Otazu, G. H., H. Chae, M. B. Davis, and D. F. Albeanu (2015). Cortical feedback decorrelates olfactory bulb output in awake mice. Neuron 86(6), 1461–1477.

Ponce-Alvarez, A., A. Thiele, T. Albright, G. Stoner, and G. Deco (2013). Stimulus-dependent variability and noise correlations in cortical mt neurons. Proceedings of the National Academy of Sciences 110, 13162–13167.

Rall, W., G. M. Shepherd, T. S. Reese, and M. W. Brightman (1966). Dendrodendritic synaptic pathway for inhibition in the olfactory bulb. Experimental neurology 14(1), 44–56.

Raman, B., J. Joseph, J. Tang, and M. Stopfer (2010). Temporally diverse firing patterns in olfactory receptor neurons underlie spatiotemporal neural codes for odors. Journal of Neuroscience 30(6), 1994–2006.

Renart, A., J. de la Rocha, P. Bartho, L. Hollender, N. Parga, A. Reyes, and K. Harris (2010). The asynchronous state in cortical circuits. Science 327, 587–590.

Rosenbaum, R., J. E. Rubin, and B. Doiron (2013). Short-term synaptic depression and stochastic vesicle dynamics reduce and shape neuronal correlations. Journal of neurophysiology 109(2), 475–484.

Rossant, C., S. Kadir, D. Goodman, J. Schulman, M. Hunter, A. Saleem, A. Grosmark, M. Belluscio, G. Denfield, A. Ecker, A. Tolias, S. Solomon, G. Buzsáki, M. Carandini, and K. Harris (2016). Spike sorting for large, dense electrode arrays. Nature neuroscience 19(4), 634–641.

Ruff, D. A. and M. R. Cohen (2014). Attention can either increase or decrease spike count correlations in visual cortex. Nature neuroscience 17(11), 1591–1597.

Schoppa, N. E. and N. N. Urban (2003). Dendritic processing within olfactory bulb circuits. Trends in neurosciences 26(9), 501–506.

Scott, J. W., H. P. Acevedo, L. Sherrill, and M. Phan (2007). Responses of the rat olfactory epithelium to retronasal air flow. Journal of neurophysiology 97(3), 1941–1950.

Spors, H., M. Wachowiak, L. B. Cohen, and R. W. Friedrich (2006). Temporal dynamics and latency patterns of receptor neuron input to the olfactory bulb. Journal of Neuroscience 26(4), 1247–1259.

Tetzlaff, T., M. Helias, G. Einevoll, and M. Diesmann (2012). Decorrelation of neural-network activity by inhibitory feedback. PLoS Computational Biology 8, e1002596.

Wachowiak, M. (2011). All in a sniff: olfaction as a model for active sensing. Neuron 71(6), 962–973.

Wanner, A. A. and R. W. Friedrich (2020). Whitening of odor representations by the wiring diagram of the olfactory bulb. Nature Neuroscience 23(3), 433–442.

Wellis, D. P. and J. W. Scott (1990). Intracellular responses of identified rat olfactory bulb interneurons to electrical and odor stimulation. Journal of Neurophysiology 64(3), 932–947.

Xiong, W. and W. R. Chen (2002). Dynamic gating of spike propagation in the mitral cell lateral dendrites. Neuron 34(1), 115–126.

Appendix

Input statistics give reasonable model results

For all firing rate models, we fixed the time-varying mean of ORN inputs $\mu(t)$, insuring MC received stronger inputs than PGC and also including odor-evoked increases for GC (Fig. 2B, top). GC do not receive direct odor input, but we justify increasing $\mu(t)$ to account for unmodeled excitation from other glomeruli to adequately capture odor-evoked increases in GC firing, but we never allowed the σ for GC to be time-varying.

The resulting firing rates depend on specific parameters, but we checked that the set of coupling strengths (Fig. 3D) for the models with 3 different sets of noise input resulted in MC spontaneous

firing rates being larger than GC and slightly larger than PGC while in the evoked state the MC firing is slightly smaller than PGC but still larger than GC (see Fig. A1**A**). Although our data only has MC recordings, *in vivo* anesthetized recordings of freely breathing rats by the Isaacson lab showed larger spontaneous and evoked firing for MC than GC on average (Cang and Isaacson, 2003). While firing in that study depends on odor concentration among other factors, a visual comparison of their Figures 6 and 10 shows larger evoked firing by MC than GC.

The firing rate relationships in Fig. A1**A** are consistent with the populations averages in the large and realistic coupled OB model in Li and Cleland (2017), where PGC and MC fire with similar rates that are higher than GC. Furthermore, Wellis and Scott (1990) showed intracellular voltage traces with current injection for both GC and PGC with varying behaviors from cell to cell (like Cang and Isaacson (2003)), but PGC visually appear to fire more spikes (their Figs. 8 and 9) than GC (their Fig. 5), even when GC exhibited strong responses to odor stimulation; they did not report comparisons of firing rates.

Figure A1: **Spike statistics of other cell types in model.** Firing rate model results for 3 input noise regimes, averaging over the 10 best sets of coupling strengths (Fig. 3**D** in main text). These model results are consistent with experiments, see **Input statistics give reasonable model results**. **A)** Firing rate for 3 different cell types in the firing rate model, shaded error bars representing standard deviation across 10 best parameter sets. **B)** Spike count variance for 3 different cell types, showing 3 different input noise regimes. **C)** Spike count covariance for 3 pairs of cell types, again combining 3 input noise regimes. Standard deviations (shaded regions) are omitted in **B)** and **C)** because they are small.

Firing rate model can exhibit evoked decreases in variability

Figure A2: **Firing rate model can exhibit odor-evoked decreases in variability. A)** Parameters where covariance decreases with stimulus and mean spike count increases (magenta dots) from the entire set of 10,000 points and for all 3 input noise regimes. Results vary depending on input noise regime, but both the independent and common GC inhibition must be relatively strong while there does not appear to be a dependence on MC excitation. **B)** Covariance values when evoked is less than spontaneous (magenta), plotted with data (black); the blue squares denote averages of the model results. **C)** With much larger GC inhibition than before, stimulus-induced decreases in variance and increases in mean spiking (magenta dots) can occur. For **C)** and **D)** we test 1000 points with the inhibition values twice as large: $10 \leq w_{MG} \leq 20$, $5 \leq w_{Gc} \leq 10$, and a subset of excitation values $4 \leq w_{GM} \leq 8$. **Ci)** Small fixed σ easily results in evoked decreases in variance. For time-varying σ (**Cii**) and large fixed σ (**Ciii**), the inhibition has to be relatively large for evoked decreases in variance. In all of the (w_{MG}, w_{Gc}) plots, the upper right corner is gray because when total GC inhibition is excessive, the mean spike counts decrease in the evoked state. **D)** All variance values plotted with data (same format as **B)**).

Although the firing rate model we use is relatively simple, we show here that the dynamic range

is vast. In particular, circuit parameters could be manipulated to yield a decrease in variability and covariability with stimulus onset.

Using Monte Carlo simulations, we calculated the steady-state spike count covariance in the spontaneous and evoked states over the entire 10,000 point parameter space of coupling strengths and identified those parameter combinations for which both variance and covariance decreased in the evoked state (Fig. A2**A**, magenta points). We see that the coupling strength values are slightly different for the 3 input noise regimes, with small fixed σ having the most such points, following by large fixed σ, then time-varying σ (Fig. A2**i–iii**). Notice that in all cases, both the independent and common GC inhibition have to be relatively strong while there does not appear to be any dependence on MC excitation to GCs. The actual steady state values are shown in Figure A2**B**, plotted with the average values (blue squares) and experimental data (black curve).

The model can also exhibit stimulus-evoked decreases in the spike count variance, but to robustly show this in all 3 input noise regimes, we had to consider much larger inhibition strength values: $10 \leq w_{MG} \leq 20$, $5 \leq w_{Gc} \leq 10$, and $4 \leq w_{GM} \leq 8$ (Fig. A2**C**). There does not appear to be a dependence on the MC excitation (w_{GM}) but there is a dependence on total inhibition (w_{MG} and w_{Gc}) in that both have to be large (Fig. A2**C**). In the (w_{Gc}, w_{MG}) plots (Fig. A2**Ci, Cii, Ciii**, lower left sub-panels), we see there is a region of gray points in the upper-right corner corresponding to larger w_{Gc} and w_{MG} values where the inhibition is so strong that the mean spike counts do not increase in the evoked state. Notice that in all cases, the common GC inhibition w_{Gc} spans the range because all that matters for the MC spike variance is the total GC inhibition, in contrast to the covariance where w_{Gc} has a prominent effect that cannot be counteracted with independent inhibition. Figure A2**D** shows the steady-state spike count variance values in spontaneous and evoked (magenta) along with average values (blue squares) and the experimental data (black curve).

Figure A2 demonstrates that even qualitatively matching our experimental data (i.e., increases in mean and variability at stimulus onset) requires parameter tuning in this model.

Details of the transfer function calculation

To calculate he transfer functions used in our main firing rate model, we implemented and simulated a large-scale model of OB cells. Our model implementation of all three multi-compartment cell types (MC, PGC, GC) are based entirely on models developed by the Cleland Lab (Li and Cleland, 2013, 2017), with coupling between different cell types mediated by dendrodendritic interactions (Rall et al., 1966). For example, synaptic coupling between MC and GC only occurs at part of the lateral dendrite of MC and part of the dendrite/spine of GC; MC and PGC are coupled only at the MC tuft and PGC dendrite/spine.

The main parameter we vary here and all cells to get the transfer function is the applied current:

I_{app} (see Eq. (17)–(19)). Each cell model contains different compartments, each having intrinsic ionic currents, including sodium, persistent sodium, potassium delayed rectifier, fast-inactivating A-current, slow-inactivating transient potassium current, L-type calcium current and a calcium-dependent potassium current. All parameters, compartments, intrinsic ionic currents and their gating variables are the same as in (Li and Cleland, 2013, 2017) unless otherwise stated.

Mitral cell model.

$$\tau_m \frac{\partial V_m}{\partial t} = \lambda_m^2 \frac{\partial^2 V_m}{\partial x^2} + I_{app} - \sum_{\text{intrinsic}} I_{ion} - g_{MP}(t)(V_m - \mathcal{E}_I) - g_{MG}(t)(V_m - \mathcal{E}_I) \tag{17}$$

where x represents spatial location assumed to be 1-dimensional with no flux boundary conditions, and with four compartments: tuft, apical dendrite, soma, and lateral dendrite.

The complete list of intrinsic ionic currents are: I_{Na} (sodium), I_{NaP} (persistent sodium), I_{DR} potassium delayed rectifier, I_A (fast-inactivating-) and I_{KS} (slow-inactivating transient potassium current), I_{CaL} (L-type calcium current), I_{KCa} (calcium-dependent potassium current). Not all compartments contain all of these ionic currents, see Li and Cleland (2013, 2017) or our code on GitHub.

Granule cell model.

$$\tau_g \frac{\partial V_g}{\partial t} = \lambda_g^2 \frac{\partial^2 V_g}{\partial x^2} + I_{app} - \sum_{\text{intrinsic}} I_{ion} - g_{GM}(t)(V_g - \mathcal{E}_E) - g_{GG}(t)(V_g - \mathcal{E}_I) \tag{18}$$

where x represents spatial location assumed to be 1-dimensional with no flux boundary conditions, and with two compartments: dendrite/spine, and soma.

Periglomerular cell model.

$$\tau_p \frac{\partial V_p}{\partial t} = \lambda_p^2 \frac{\partial^2 V_p}{\partial x^2} + I_{app} - \sum_{\text{intrinsic}} I_{ion} - g_{PM}(t)(V_p - \mathcal{E}_E) \tag{19}$$

where x represents spatial location assumed to be 1-dimensional with no flux boundary conditions, and with two compartments: dendrite/spine, and soma.

We use standard synaptic connections, with all presynaptic GC and PGC providing GABA$_A$ inputs, and all MC providing both AMPA and NMDA – all AMPA and NMDA connection strengths are the same for any given pre/postsynaptic pair, with $g_X(t) = w_X s_X(t)$ where w_X is the strength and all s_X evolve according to:

$$\frac{ds}{dt} = \alpha a_{on}(v_{pre}(t))(1 - s) - \beta s \tag{20}$$

Here $a_{on}(v_{pre}) = 1/\left(1 + \exp\left(-(v_{pre} - v_t)/v_s\right)\right)$. We adopt the Li and Cleland (2013, 2017) convention of graded inhibitory synapses; GABA$_A$ synapses activate via a sigmoidal function with a lower threshold of $v_t = -40$ mV and scale $v_s = 2$, while AMPA and NMDA synapes have $v_t = 0$ mV and $v_s = 0.2$. The α and β values are: $\alpha_{GABA}^{-1} = 1.25$ ms, $\beta_{GABA}^{-1} = 18$ ms, $\alpha_{AMPA}^{-1} = 1$ ms, $\beta_{AMPA}^{-1} = 5.5$ ms, $\alpha_{NMDA}^{-1} = 52$ ms, $\beta_{NMDA}^{-1} = 343$ ms.

In Fig. A3**Ai** we see that with an abrupt step of current injection from 0 to $50\,\mu$A/cm^2, the uncoupled PGCs exhibit transient spikes but no sustained spiking, with similar voltage profile as the model in Li and Cleland (2013) (their Fig. 4) and in slice recordings (McQuiston and Katz, 2001). The uncoupled MCs exhibit sub-threshold oscillations as observed in experiments (Desmaisons et al., 1999; Balu et al., 2004), and the number of spikes in a burst increases with current step size (Balu et al., 2004); see Fig. A3**Aii**, again like the model in Li and Cleland (2013). Figure A3**D** shows example voltage trajectories in a coupled network where PGC and GC provide GABA$_A$ synaptic inputs and MC provides both AMPA and NMDA inputs with same weight to the appropriate compartments. Voltages within the same cell vary slightly depending on compartment and spatial location. The inset of MC voltage (Fig. A3**D** bottom) shows relatively fast spike propagation, on the order of a few milliseconds (Xiong and Chen, 2002).

Calculating transfer functions

Firing rate models generally use the transfer functions of individual cell models that are uncoupled. However, since there is no sustained firing in the uncoupled PGC model (Fig. A3**Ai**) with even large values of injected current, the PGC transfer function is not viable. Thus, to incorporate the affects of PGC, we have to instead use transfer functions calculated from reciprocally coupled cells.

We considered two reciprocally coupled networks: i) 1 GC and 1 MC, and ii) 1 PGC and 1 MC; in each network we varied the constant current infection values with fixed synaptic coupling strengths $w_{GM} = w_{MG} = w_{PM} = w_{MP}$ that are dimensionless; recall $g_X(t) = w_X s_X(t)$ where s_X evolves according to Eq. (20) and g_X are in the voltage equations for all 3 cell types. The firing rates were calculated with 50 s of biological time (removing initial 1 s of transients); the average of the two MC firing rates is used.

We calculated several transfer functions (Fig.A3**C**) with fixed synaptic coupling strengths $w_{GM} = w_{MG} = w_{PM} = w_{MP}$, using values of 60, 80, 100, 120, and 140. Figure A3**C** shows the resulting curves for each cell type and for the synaptic coupling strengths. Since the input (or constant current injection values) are scaled and the curves appear relatively similar, we use the $w = 100$ curves.

Figure A3: **Multi-compartment biophysical OB model used for the firing rate model.** Small network of Cleland OB models (Li and Cleland, 2013, 2017). **A**) The firing rate response of uncoupled cells capture some known behavior observed in experiments. i) From rest (no current applied), an abrupt step of current injection to $50\,\mu A/cm^2$ results in transient spikes but no sustained firing, consistent with Li and Cleland (2013); McQuiston and Katz (2001). ii) Uncoupled MC show prominent sub-threshold oscillations (Desmaisons et al., 1999; Balu et al., 2004) and an increase in number of spikes per burst as applied current values increase, consistent with experiments in (Balu et al., 2004). **B**) Voltage traces of the coupled network show random spontaneous and evoked firing; inset of MC at bottom shows relatively fast propagation within a cell, on the order of a few milliseconds (Xiong and Chen, 2002). The results in **A**) and **B**) are not surprising because Li and Cleland (2013, 2017) showed these. **C**) Numerically computed transfer functions with several coupling strengths. Since the shape of the curves are qualitatively similar, we set the curves to be the solid thick curves ($w = 100$) in the firing rate model.