Combinatorial patterns of gene expression changes contribute to variable expressivity of the developmental delay-associated 16p12.1 deletion

Matthew Jensen,[1,2,7] Anastasia Tyryshkina,[1,3,7] Lucilla Pizzo,[1] Corrine Smolen,[1,2] Maitreya Das,[1] Emily Huber,[1] Arjun Krishnan,[4,5] and Santhosh Girirajan[1,2,3,6]*

1. Department of Biochemistry and Molecular Biology, Pennsylvania State University, University Park, PA 16802, USA
2. Bioinformatics and Genomics Program, Huck Institute of the Life Sciences, Pennsylvania State University, University Park, PA 16802, USA
3. Neuroscience Program, Huck Institute of the Life Sciences, Pennsylvania State University, University Park, PA 16802, USA
4. Department of Computational Mathematics, Science and Engineering, Michigan State University, East Lansing, MI 48824, USA
5. Department of Biochemistry and Molecular Biology, Michigan State University, East Lansing, MI 48824, USA
6. Department of Anthropology, Pennsylvania State University, University Park, PA 16802, USA
7. These authors contributed equally to this work

* Correspondence to:
Santhosh Girirajan
205A Life Sciences Building
Pennsylvania State University
University Park, PA 16802
E-mail: sxg47@psu.edu
Twitter: @girirajan16

ABSTRACT

Recent studies have suggested that individual variants do not sufficiently explain the variable expressivity of phenotypes observed in complex disorders. For example, the 16p12.1 deletion is associated with developmental delay and neuropsychiatric features in affected individuals, but is inherited in >90% of cases from a mildly-affected parent. While children with the deletion are more likely to carry additional "second-hit" variants than their parents, the mechanisms for how these variants contribute to phenotypic variability are unknown. We performed detailed clinical assessments, whole-genome sequencing, and RNA sequencing of lymphoblastoid cell lines for 32 individuals in five large families with multiple members carrying the 16p12.1 deletion. We found that the deletion dysregulates multiple autism and brain development genes such as *FOXP1*, *ANK3*, and *MEF2*. Carrier children also showed expression changes that were inherited as well as *de novo* compared with their parents, which matched with 39/47 observed developmental phenotypes. We identified significant enrichments for 13/25 classes of "second-hit" variants in genes with expression changes, where 7/25 variant classes were only enriched when inherited from the non-carrier parent, including missense SNVs and large deletions. In 11 instances, including for *ZEB2* and *SYNJ1*, gene expression was synergistically altered by both the deletion and inherited "second-hits" in carrier children. Finally, brain-specific interaction network analysis showed strong connectivity between genes carrying "second-hits" and genes with transcriptome alterations, including differential expression, alternative splicing, and allele-specific expression. Our study shows that family-based assessments of transcriptome data are highly relevant towards understanding the genetic mechanisms associated with complex disorders.

INTRODUCTION

Complex genetic disorders, such as autism, intellectual disability/developmental delay (ID), epilepsy, and schizophrenia, have been attributed to rare copy-number variants (CNVs), or deletions and duplications encompassing multiple genes, as well as single nucleotide variants (SNVs)[1,3,5,7]. Despite advances in high-throughput sequencing methods and quantitative assessments of large cohorts, individual variants implicated for these disorders do not sufficiently explain the variable expressivity of clinical features often observed in affected individuals[9,11,13]. An example is the 520-kbp deletion at chromosome 16p12.1 [MIM: 136570], which was originally described in children with developmental delay[14,15] but was subsequently found to confer increased risk for schizophrenia[16], epilepsy[17], and cognitive defects in control populations[18]. Unlike syndromic CNVs such as Smith-Magenis syndrome [MIM: 182290] that primarily occur *de novo*[19], the 16p12.1 deletion is inherited in over 90% of affected children from carrier parents who manifest subclinical or mild cognitive and neuropsychiatric features[2,14]. In fact, we recently found that children with the deletion were more likely to carry an additional burden of rare CNVs[15] and deleterious variants in genes intolerant to variation[2], or "second-hit" variants, compared to their carrier parents, making the deletion an ideal model for assessing the combined effects of multiple variants towards variable clinical outcomes.

While dissecting the pathogenicity of complex disorders has been challenging, cohort and family-based studies that integrate multiple variants with different effect sizes or functional outcomes have provided insights into how the genetic architecture contributes to changes in penetrance, severity, and complexity of phenotypes. In particular, analysis of gene expression patterns in human cells allows for dissecting the direct and indirect effects of genomic variants towards biological functions in complex disorders. *For example*, Merla and colleagues assessed gene expression in skin fibroblasts and lymphoblastoid cell lines (LCLs) from individuals with the 7q11.23 deletion, associated with Williams syndrome [MIM: 194050], and found that several genes adjacent to the deletion region were also downregulated compared to controls[20]. Similarly, expression changes due to the autism-associated 16p11.2 deletion [MIM: 611913] correlated with changes in head circumference phenotypes[21] and converged on several neurodevelopmental pathways, including synaptic function and chromatin modification[22]. Other studies have used expression data to identify pathogenic variants potentially missed by genome sequencing studies[23–25]. *For example*, Frésard and colleagues identified novel causal variants for 6/80

individuals with rare undiagnosed diseases through paired analysis of whole-blood

transcriptomes and genomes[23]. Additionally, several recent studies have used family-based

approaches to study the effects of rare inherited variants towards gene expression. *For example,*

Pala and colleagues found that rare inherited variants in both coding and non-coding regions

increased the likelihood of gene expression changes among 61 families in the bottlenecked

Sardinia population, indicating the importance of such variants towards disease risk[26]. While

these studies have shown the utility of assessing transcriptomic consequences of individual

causal variants, they were focused on either control populations *or* relatively invariable

disorders, and did not examine the simultaneous effects of multiple variants with different effect

sizes towards changes in gene expression within major biological pathways.

Here, we integrated whole-genome sequencing and transcriptome data of LCLs from 32

individuals in five large 16p12.1 deletion families who manifested variable phenotypes, in order

to investigate how the combined effects of the deletion and "second-hits" perturb transcriptional

networks and biological functions. We found that the 16p12.1 deletion disrupts expression of

genes involved in neuronal and developmental functions, such as signal transduction and cell

proliferation, as well as genes preferentially expressed in the fetal and adult brain. We further

identified significant contributions of several classes of rare "second-hit" coding and non-coding

variants towards changes in gene expression among carrier children compared with their parents,

especially when the variants were inherited from the noncarrier parent. In fact, we found 11

instances of genes in carrier children whose expression was synergistically altered by the

combined effects of the 16p12.1 deletion and "second-hit" variants inherited from the noncarrier

parent. Although a relatively small sample size precluded global analyses between these

expression changes and developmental phenotypes, we found that specific expression changes

contributed towards distinct clinical features of affected children through disruption of biological

functions related to neurodevelopment. Our results suggest that the 16p12.1 deletion and

"second-hit" variants jointly disrupt the developmental transcriptome through shared pathways to

contribute towards developmental phenotypes, emphasizing the importance of family-based

transcriptome studies for complex disorders.

MATERIALS AND METHODS

Patient recruitment and clinical phenotype analysis

We obtained clinical data and whole-blood DNA from 32 individuals in five families with the 16p12.1 deletion. Carrier children and their family members provided informed consent according to a protocol reviewed and approved by the Pennsylvania State University Institutional Review Board (IRB #STUDY00000278). Among the recruited individuals were 10 children with the deletion ("carrier children"), six sets of carrier and noncarrier parents, three sets of carrier and noncarrier grandparents, and four noncarrier siblings (**Figure S1; Table S1**). Affected children and family members were identified as carriers of the deletion through prior clinical diagnostic tests, which we confirmed using SNP microarray analysis[2]. Phenotypic information from the five families was collected through self-reported questionnaires, medical records, and interviews with family members. These data represent comprehensive medical history of affected children and their family members, including neuropsychiatric and developmental features (including cognitive, behavioral, and psychiatric diagnoses), anthropomorphic measures, abnormalities across multiple organ systems (nervous, craniofacial, musculoskeletal, cardiac, hearing/vision, digestive, and urinary systems), and family history of medical or psychiatric disorders.

DNA extraction, whole-genome sequencing, and variant identification

We identified a full range of rare deleterious variants from whole-genome sequencing and SNP microarray for each of the 32 family members in our cohort. Genomic DNA was extracted from peripheral blood using QIAamp DNA Blood Maxi extraction kit (Qiagen, Hilden, Germany) and treated with RNAse. DNA levels were then quantified using Quant-iT™ PicoGreen™ dsDNA assay methods (Thermo Fisher Scientific, Waltham, MA, USA), and sample integrity was assessed in agarose gel. After constructing Illumina TruSeq DNA PCR-free libraries (San Diego, CA, USA), whole-genome sequencing was performed on each sample by Macrogen Labs (Rockville, MD, USA) using an Illumina HiSeq X sequencer to obtain an average coverage of 34.5X. Raw sequencing data were processed for quality control using Trimmomatic[27] with leading:5, trailing:5, and slidingwindow:4:20 parameters, aligned to the human hg19 reference genome using BWA[28] v.0.7.13, and sorted and indexed using Samtools[29] v.1.9.

The GATK Best Practices pipeline v.3.8 (HaplotypeCaller) and v.4.0.11

(GenotypeGVCFs)[30] was used to identify SNVs and small indels from WGS data. In short,

duplicate reads were marked and removed using PicardTools, and after calibration of base-pair

quality scores, GATK HaplotypeCaller was used to identify variants in each sample. Variant

calls were then pooled for joint genotyping and calibration of variant quality scores. Custom-

built pipelines using Annovar[31] v.2016Feb01 applied a total of 430 annotation classes for variant

function, population frequency, conservation, genomic region, and predicted pathogenicity.

Variants were filtered based on the following quality metrics[32]: QUAL >50, read depth >8, allele

balance between 0.25-0.75 (or >0.9 for homozygous variants), and quality depth (QUAL/reads

with alternate allele) <1.5. Rare variants were defined as variants with frequency ≤0.001 in the

gnomAD v.2.1.1 genome database[33], and present in <10 samples in our in-house WGS cohort of

125 families (335 individuals) with rare CNVs. We finally classified rare SNVs and small indels

for downstream analysis as follows: Rare missense and loss-of-function (including frameshift

and stopgain) variants within protein coding regions, as well as variants in the 5' and 3' UTR or

within 1 kbp of the transcription start (TSS) or end sites (upstream and downstream), were

classified based on their RefSeq-defined genomic locations in Annovar (**Figure S2**). Splice-site

variants were identified based on MutationTaster annotations[34] for disease-causing ("D") or

disease-causing automatic ("A") variants. Rare non-coding regulatory variants within 50 kbp of

TSS for protein-coding genes were classified according to ChromHMM chromatin state

segmentation data for GM12878 lymphoblastoid cells[35], available from the ENCODE Project,

into promoters (chromosome states 1-3), enhancers (states 4-7), or silencers (state 12). With the

exception of loss-of-function and splice-site variants, all coding and non-coding variants were

filtered for CADD Phred-like pathogenicity scores[4] ≥10. Inheritance patterns of these variants

were determined using in-house pipelines.

Copy-number variants and structural variants were identified using a combination of

WGS data for all samples and SNP microarray data for 25/32 samples, previously described in [2].

To identify variants from microarray data, Illumina Omni 2.5 BeadChip microarray experiments

were performed for each sample at either the HudsonAlpha Institute for Biotechnology

(Huntsville, AL, USA; n=18), Yale Center for Genome Analysis (New Haven, CT, USA; n=5),

or the Department of Genome Sciences at the University of Washington (Seattle, WA, USA;

n=2). CNV calls for each sample were generated using PennCNV[36] v.1.0.3, and were filtered for

≥50 kbp in length and ≥5 target probes. CNVs and SVs were also detected from aligned WGS

data using a combination of four pipelines: CNVNator[37] v.0.4.1 (bin size of 200), DELLY[38]

v.0.8.2, LUMPY-sv[39] v.0.2.13 with Smoove v.0.2.5, and Manta[40] v.1.6.0. In both WGS and

microarray-derived datasets, adjacent CNVs were merged if they overlapped or had a gap <20%

of CNV length and <50 kbp. We then integrated the CNV and SV calls from each of the datasets

as follows: For smaller CNVs and SVs <50 kbp, any duplication or deletion called by at least

two of the four WGS-based callers were considered for downstream analysis, with the minimum

intersected regions defining the new breakpoints. For larger CNVs and SVs >50 kbp, the union

of CNVNator and PennCNV calls were considered for downstream analysis. Integrated calls

were based on 50% reciprocal overlap among the callers. After integration, variants were filtered

for presence in <10 individuals in our in-house WGS cohort, as determined by 50% reciprocal

overlap. Finally, RefSeq gene-coding regions spanned by SVs were categorized as follows:

encapsulating variants which span the entire gene, interstitial variants that are contained within a

gene, and 5' and 3' UTR variants that overhang the gene on either end (**Figure S2**). Inheritance

patterns of CNV and SV calls were determined if calls in the child and parent had >50%

reciprocal overlap.

Short tandem repeats were called from aligned WGS data with GangSTR[41] v.2.4, using

the GangSTR hg19 reference file v.13.1. The calls were filtered and analyzed using three tools

from the STR analysis toolkit TRTools[42]. First, dumpSTR was used to filter for quality of calls

using the following parameters: read depth >20, read depth <1000, QUAL >0.9, spanbound only

(calls that are spanned by reads), and filter bad confidence intervals (filtered calls whose

maximum likelihood estimates were not within the confidence interval). The reads were then

merged with mergeSTR, and basic statistics were calculated using statSTR. In addition to the

dumpSTR filters, we applied the following filters to our call set: >95% of samples called for the

STR location, variance at location >0, and overlap of the STR location with a RefSeq defined

protein-coding gene. STR expansions were defined as any call for which the deviation of the

repeat length was greater than the mean length plus three standard deviations among all

individuals in our cohort. Finally, STR variants were categorized according to their genomic

location, including exonic, intronic, 5' or 3' UTR, upstream, and downstream (**Figure S2**).

Inheritance patterns of STRs were determined by matching the number of repeats in the child to

their parents.

Generation of lymphoblastoid cell lines and RNA-sequencing

Peripheral blood samples for all 32 individuals in our cohort were submitted to the Coriell Institute for Medical Research (Camden, NJ, USA) for generation of lymphoblastoid cell lines through Epstein-Barr virus transformation of B-lymphocytes (**Table S2**). After receiving the LCL samples, cells were grown at 5% CO_2 and a concentration of 1X 106 cells/mL under L-glutamine-supplemented RPMI 1640 medium (11875-119, Thermo Fisher Scientific) containing 15% fetal bovine serum (35-010-CV, Corning Life Sciences, Tewksbury, MA, USA) and Cytiva HyClone™ Penicillin Streptomycin solution (SV30010, Thermo Fisher Scientific). Total RNA was isolated from three biological replicates of P6-P7 cells per sample using TRIzol Reagent (Thermo Fisher Scientific) and PureLink RNA Mini Kit (12183018A, Thermo Fisher Scientific), and subsequently treated with DNA-free DNA Removal Kit (AM1906, Thermo Fisher Scientific). RNA integrity number scores (RIN) were assessed using Agilent Bioanalyzer 2100 (**Figure S3A**), and replicates with RIN scores >8.5 were sequenced. Paired-end 50 bp libraries for each replicate were generated using Illumina TruSeq Stranded mRNA kit, and were sequenced using Illumina NovaSeq at the Penn State College of Medicine Genome Sciences Facility (Hershey, PA, USA). Two sequencing runs of 48 replicates were performed, with the biological replicates of each sample split among the two runs to mitigate batch effects, to generate a total of 43.5 million reads/replicate.

Quantification of gene expression and coverage of disease genes

Sequenced RNA reads were filtered using Trimmomatic[27] v.0.36 to remove reads <30 bp long. Following the GTEx Consortium RNA-Seq pipeline[6], the filtered reads were aligned to the human genome (GENCODE v.19) using STAR[43] v.2.4.2a, and sorted and indexed using Samtools[29] v.1.9. Duplicates reads were marked with PicardTools v.2.9.0. We assessed the quality of the aligned reads with transcript integrity scores[44], which moderately correlated (r=0.38, p=1.0×10^{-4}, Pearson correlation) with the RIN scores for each sample (**Figure S3A**). Isoform counts for GENCODE 19 genes were quantified using RSEM[45] v.1.2.22. A collapsed gene coordinate GTF file was generated using the GENCODE 19 gene coordinates and the GTEX collapse_annotation script. Gene-level counts and transcripts per million read (TPM) values were quantified using RNASeQC[46] v.1.1.8, using strict mode and the collapsed gene coordinates.

After filtering for transcripts where all three replicates of at least one sample showed >0.2

TPM, we obtained a total of 24,340 expressed transcripts across our cohort, representing 43.3%

of all GENCODE transcripts. We further compared our set of expressed LCL genes to disease

gene databases[12,47–50] and genes expressed in the adult brain from GTEx consortium RNA-Seq

data[6]. We defined expressed genes in GTEx tissues if they showed >0.5 TPM in 80% of samples

for a particular tissue. The expressed LCL genes covered >70% of each of these gene sets,

including 83% of genes expressed in GTEx brain tissues (**Figure S4A-B**). These data are in

concordance with gene expression data from GTEx, where gene expression values in LCLs and

brain tissues showed an average Spearman correlation of 0.84 (**Figure S4C**). These findings

indicate that our LCL data would be able to identify changes in expression patterns for most

genes related to neurodevelopmental disease.

Differential expression and outlier expression analysis

We performed differential expression analysis between all 16p12.1 deletion carriers and

noncarriers, as well as between parents and offspring across the five families, using edgeR[51]

v.3.30.0 on gene-level counts to create generalized linear models and perform quasi-likelihood

F-tests with Benjamini-Hochberg correction. For testing differences between all deletion carriers

(n=19) and noncarriers (n=13), we included family as a covariate in the linear model, used

default filtering for low-expressed genes, and removed genes with sex-specific differences in

GTEx LCL samples as well as genes on the X and Y chromosomes (due to unequal sex ratios in

deletion carriers and noncarriers). To control for expression outliers, we iteratively identified sets

of differentially expressed genes, defined using an FDR<0.05 threshold (Benjamini-Hochberg

correction), between deletion carriers and noncarriers after removing one sample at a time. We

then took the intersection of the resulting 32 sets of differentially expressed genes, and obtained

a total of 1,569 transcripts differentially expressed in the deletion carriers (**Table S3**). We also

performed differential expression analysis using PQLseq v.1.2 to account for gene expression

similarity due to relatedness[52]. We first generated input files from unfiltered WGS SNV data

using PLINK[53] v.1.9, and used GEMMA v.0.98.3, which calculates kinship between two

individuals based on genotype similarity[54], to generate a kinship matrix for our cohort. This

matrix was used as input for PQLseq along with gene-level counts from RNA-Seq data, after

removing the same sex-specific genes as for the edgeR analysis.

We next performed family-based analysis on 13 separate trios identified across the five

families (nine carrier children compared to parents and four carrier parents compared to

grandparents), which are listed in **Table S1**. For example, we separately analyzed two trios in

family GL_001 (**Figure S1**). For comparison, we analyzed an additional four trios with

noncarrier children compared to carrier and noncarrier parents (**Figure S5C**). For each trio, we

first used an edgeR workflow without covariates to identify differentially expressed genes

between the offspring and carrier parent (|logFC|>0.5, FDR<0.05, Benjamini-Hochberg

correction), and separately assessed expression changes between the offspring and noncarrier

parent. Genes with low expression (expressed in <25% of all replicates) and sex-specific genes

were removed from edgeR analysis. We then overlapped the two sets of differentially expressed

genes to classify expression changes by their inheritance patterns as follows: "*de novo*" if the

gene was differentially expressed in the offspring compared with both parents; "inherited from

the carrier parent" if the gene was only differentially expressed compared with the noncarrier

parent; and "inherited from the noncarrier parent" if the gene was only differentially expressed

compared with the carrier parent (**Figure S5A**).

To identify genes with outlier expression in our cohort, we calculated z-scores of gene

expression values in each individual for 14,212 protein coding genes expressed in the LCL

samples. We normalized the expression values in each person by calculating the median TPM

expression across the three replicates for each gene, transformed the values using $\log_2(x + 1)$,

and calculated z-scores for each log-transformed TPM compared with all samples in our cohort.

As principal component analysis (PCA) showed clustering of samples by family (**Figure S6A**),

we used PEER[55] v.1.0 to correct the z-scores using one PEER principal component (**Figure**

S6B). After correction, we further assessed for clustering of samples and replicates, and found

strong Spearman correlations among replicates derived from the same sample (**Figure S3B**). We

defined outlier genes as any gene with |z-score| >2 (**Figure S6C**), in line with recent studies

utilizing outlier expression values[23].

Enrichment analysis for biological function, brain expression, and disease relevance

Enrichment analysis for sets of differentially expressed genes was performed using goseq[56]

v.3.12, which tests for overrepresentation of gene categories in RNA-seq data. Goseq controls

for selection bias in RNA-seq datasets by modeling the distribution of transcript lengths of

differentially expressed genes. We assessed for enriched biological processes using the Gene

Ontology database[57], as well as genes expressed in specific adult brain tissues from GTEx[6] and

developing brain tissues from the BrainSpan Atlas[58]. We defined preferentially-expressed genes

in GTEx and BrainSpan tissues as expression >2 standard deviations higher than the median

expression across all tissues for that gene. We further assessed for enrichment of differentially

expressed gene sets for candidate neurodevelopmental disease genes (DBD Gene Database)[47], as

well as specific gene sets for autism (SFARI Gene database)[48], intellectual disability (DDD and

DDG2P databases)[49,50], and schizophrenia[12]. Finally, we annotated sets of genes with altered

expression for two common measures of intolerance to variation, RVIS[8] and pLI[10], and used

genes considered to be intolerant to variation (RVIS $<20^{th}$ percentile or pLI score >0.9) for

downstream analysis.

Weighted gene correlation network analysis

We performed weighted gene correlation network analysis (WGCNA) to identify modules of

genes that were co-expressed among samples in our cohort[59]. We used the R package tximport[60]

to import RSEM-derived gene expression counts, filtered genes for >10 counts/replicate in at

least one sample, and used DESeq2[61] to generate variance-stabilized expression counts for each

gene. To detect co-expression patterns specific to deletion carriers, we used ComBat[62] within sva

v.3.12 to perform batch correction with family as a covariate. We detected 35 co-expression

modules in our samples using WGCNA[59] v.1.69, with the following parameters: power threshold

= 8, signed hybrid network, unsigned topological overlap matrix, bi-weight mid-correlation,

module size = 30-30000, and merge cut height = 0.25. Two modules showed strong sex-specific

gene expression and were excluded from further analysis. The average gene expression values in

each module were compared between carriers and noncarriers using two-tailed t-tests, and genes

in each of these modules were tested for enrichment of Gene Ontology terms using goseq.

Integration of gene expression and genomic variant data

We calculated the effect size of different classes of rare "second-hit" variants towards gene

expression changes, stratified by sample type and inheritance patterns. We compared 25 classes

of rare variants identified from WGS data (**Figure S2**) towards differentially expressed genes in

family trios as well as outlier expression genes in all individuals. For all comparisons, we

calculated odds ratios and 95% confidence intervals for each variant class towards changes in

expression using Fisher's exact and Wald tests, respectively (**Table S4**). We note that we

considered each variant class independently, so that dysregulated genes with multiple types of

disrupting variants were counted within multiple variant classes. For the differential expression

analysis, we first assessed variants in the 13 trios with carrier offspring for genes with

differential expression (**Table S1**), and then determined the effects of variants in carrier children

(n=9 trios) inherited from carrier or noncarrier parents towards expression changes that were

inherited from the same parent. For the outlier expression analysis, we first assessed variants for

outlier expression genes in all individuals. We then stratified these data by sample type (carrier

child, carrier parent, and noncarrier parent), and compared variants in carrier children that were

inherited from their carrier or noncarrier parent. To identify synergistic effects between the

16p12.1 deletion and "second-hit" variants, we identified a subset of genes with outlier

expression in deletion carriers that were also differentially expressed in the global comparison of

carriers and noncarriers, and then identified those genes which also had "second-hit" variants

inherited from the noncarrier parent.

Alternative splicing analysis

To assess alternative splicing events from RNA-sequencing data, we used DESeq2[61] to detect

differential expression of isoforms. After importing isoform-level expression counts from RSEM

using tximport[60], we filtered for genes with >2 counts across all samples, and performed pairwise

comparisons between carrier offspring and their parents in the 13 trios listed in **Table S1**, plus

the four trios with noncarrier children for comparisons. We then repeated the DESeq2 analysis

for gene expression counts, and only included differentially expressed isoforms within genes that

did not show overall differential expression, to specifically account for isoform changes due to

alternative splicing. Similar to the family-based differential expression analysis, we assigned

inheritance patterns to each alternative splicing event observed in offspring based on the pairwise

comparisons to each parent. We further compared alternate isoforms identified by DESeq2 to

those in GTEx LCL data[6] to identify unique isoforms in our cohort. Finally, we integrated these

data with 12 classes of putative splice-site disrupting variants identified from WGS data, and

calculated odds ratios as described above.

Allele-specific expression analysis

We used the phASER[63] v.1.1.1 pipeline to identify allele-specific expression events in our cohort. We first used whatshap[64] v.0.18 to perform read-backed and pedigree-informed phasing of our WGS samples, and then merged the three replicate BAM files of aligned RNA-Seq reads for each sample together using Samtools, We then used phASER, which uses phased WGS data to infer phasing of RNA-Seq samples, to phase the RNA-Seq alignments and to count the number of reads per haplotype block. We ran phASER with the parameters --mapq 255 and --baseq 10, and used the recommended blacklist to remove HLA genes. Finally, we quantified log-fold changes for allelic counts in each protein-coding gene with >10 read counts using phASER Gene AE, and identified ASE for genes with FDR>0.05 using binomial tests with Benjamini-Hochberg correction. For each identified ASE event, we examined the over-expressed haplotype for presence of a deleterious rare coding variant identified from WGS, which would potentially indicate pathogenic effects of the ASE event. Finally, we determined inheritance patterns of ASE genes based on the presence of ASE in parents of offspring.

eQTL discovery and analysis

We used QTLTools[65] v.1.2 to identify eQTLs in our cohort. Because we had three replicates per participant, we first calculated the median TPM values for all transcripts in an individual. Genes were filtered for >0.1 median TPM in more than 50% of samples. Principal components for gene expression (from RNA-Seq data) and genotype (from whatshap-phased WGS data) were then computed using QTLtools. The top three genotype and the top two gene expression principal components were used as covariates for the linear model, in addition to three explicit covariates (family, sex, and carrier status). QTLtools cis-permutation tests (n=1000 replicates) were then used to discover eGenes and associated variants in our samples. We performed multiple testing correction with the QTLtools script runFDR_cis.R. Finally, we annotated significant eQTL variants (FDR<0.05) associated with protein-coding genes for presence in GTEx LCL data, genomic location, population frequency, and biological functions using the WGS Annovar-based pipeline[31].

Brain-specific network analysis

We assessed the connectivity patterns of genes with "second-hit" variants and changes in expression in the context of a brain-specific interaction network. This network was previously built using a Bayesian classifier trained on hundreds of gene co-expression, protein-protein interaction, and regulatory-sequence datasets, in order to predict the likelihood of interactions between any two pairs of brain-expressed genes[66,67]. To create a network containing only the highest probability predicted gene interactions, we extracted all pairs of genes with weighted probabilities >2.0, representing the top ~0.5% of pairwise interactions (217,975,718 pairs of genes). We then calculated the weighted shortest path lengths for all pairs of genes in the network, using the inverse of the probabilities as weights for each edge. Finally, we created sub-networks that contained genes with "second-hit" protein-coding variants (LOF, missense, splice-site, exonic STR, or encapsulated deletion or duplication) or expression changes (differential expression, outlier expression, alternative splicing, ASE, or eQTL minor allele) for each carrier offspring from the 13 trios (**Table S1**). For each trio, we calculated the average shortest paths between all pairs of genes with expression changes and genes with "second-hit" coding variants, and then compared these distances to average shortest paths calculated from 100 permuted network replicates, where genes were randomly reassigned to different nodes. Network analysis was performed using the NetworkX package in Python[68].

Statistical analysis

All genomic and statistical analyses were conducted using either Python v.3.7.3, with packages numpy[69] v.1.16.2, scipy[70] v.1.1.0, and pandas[71] v.1.0.0, or using R v.3.5.1 (R Foundation for Statistical Computing, Vienna, Austria). Details of all statistical tests, including summary statistics, test statistics, p-values, and confidence intervals, are provided in **Table S4**.

RESULTS

The 16p12.1 deletion leads to pervasive disruption of genes involved in neurodevelopment

We performed RNA-sequencing on LCL samples from 19 deletion carriers and 13 noncarriers from five large families with multiple affected members (**Figure 1; Figure S1; Table S1**), and identified 1,569 transcripts that were differentially expressed (FDR<0.05) in deletion carriers compared with noncarriers (**Figures 2A-B; Table S3**). Application of additional corrections for relatedness among the samples[52] (see Methods) yielded 1,044 differentially expressed transcripts, of which 840 (80.5%) were also identified in the main analysis (**Figure S7A; Table S3**). We first confirmed that each of the seven protein-coding genes in the deletion region were downregulated in deletion carriers (**Figure 2C**). Interestingly, 21 protein-coding genes adjacent to the 16p12.1 region (between chromosomal bands 16p11.2 and 16p12.3) also showed differential gene expression in carriers, 18 of which were under-expressed in the deletion carriers. *For example*, two genes within the autism-associated 16p11.2 region, *SEZ6L2* and *DOC2A*, were downregulated in carriers of the 16p12.1 deletion. As none of the carriers harbored an atypical deletion, it is possible that these adjacent genes could be affected by haploinsufficiency of regulatory elements located within the deletion region. In fact, three downregulated genes adjacent to the deletion, *DNAH3*, *OTOA*, and *NPIPB4*, exhibited chromatin interactions with enhancer elements within the deletion region, detected using published Hi-C data of LCL samples[72] (**Figure 2C**).

We found that differentially expressed genes in deletion carriers were enriched (FDR<0.05) for multiple biological functions, including biological adhesion and cell proliferation regulation for relatedness-corrected genes, and signal transduction and locomotion for genes without relatedness correction (**Table S3**). Additionally, we observed an enrichment (FDR=0.015) for candidate autism genes[48], including *FOXP1*, *CUL7*, *ANK3*, and *EP300*, among the differentially-expressed genes (**Figure 2B; Table S3**). Weighted-gene correlation network analysis further identified several modules of genes with significant expression changes in deletion carriers (p<0.05, two-tailed t-test), including downregulated genes enriched for cell signaling and adhesion, and upregulated genes enriched for neurogenesis, nervous system development, and MAPK and Notch signaling (**Figure S8; Table S5**). Differentially expressed genes in deletion carriers were further enriched (FDR<0.05) for genes preferentially expressed in the hippocampus and basal ganglia of the adult brain[6] (**Figure S7B; Table S3**), as well as in the

striatum, thalamus, and frontal cortex during late fetal and adolescent/young adulthood

timepoints[58], which are critical transition periods for expression of neurodevelopmental genes[73–

75] (**Figure 2D; Table S3**). Overall, our data suggest that the 16p12.1 deletion leads to pervasive

transcriptomic changes across multiple biological and neuronal processes in the developing

brain.

Family-specific patterns of gene expression changes influence developmental phenotypes

We next investigated how gene expression patterns segregated within 13 complete trios with

carrier offspring extracted from the five families, including carrier children compared to their

parents as well as carrier parents compared with grandparents (**Table S1**). For each trio, we

identified differentially expressed genes for offspring-carrier parent and offspring-noncarrier

parent pairs (see Methods), and overlapped the two sets of differentially expressed genes to

categorize expression changes based on their inheritance pattern (**Figure S5A; Tables S6, S7**).

We found no differences (p=0.735, two-tailed paired Mann-Whitney test) in the proportion of

differentially expressed genes inherited from the carrier (avg. 2,223 genes/offspring) or

noncarrier parent (avg. 1,908 genes/offspring; **Figure 3A**). This suggests that "second-hit"

variants from the noncarrier parent contribute equally to gene expression changes, and therefore

to disease pathogenicity, as the combined effects of the deletion and any "second-hit" variants

from the carrier parent, an observation that corresponds with our recent findings of increased

burden of "second-hits" transmitted to the child from noncarrier parents[2] (**Figure S5B**).

Interestingly, we also observed an average of 1,192 genes/offspring that were differentially

expressed compared with both parents (**Figure 3A**), such as *SHANK2*, *FOXP1*, and *CACNA1D*.

These expression changes potentially represent effects of *de novo* variants or combinatorial

effects of variants inherited from both parents, which could explain the increased phenotypic

severity observed in the carrier children. However, the trends in expression patterns widely

varied across families, which in some cases could be explained by family history of

neuropsychiatric disease (**Figure S5C**). For example, we found that children within family

GL_004, whose parents were unaffected or presented with mild depression, had the lowest

number of gene expression changes among any carrier children in the cohort. Meanwhile,

children in families GL_001 and GL_052, whose carrier parents manifested multiple overt

cognitive and neuropsychiatric features, had higher proportions of inherited expression changes

from their carrier parents than their noncarrier parents (**Figure S5C**).

We next assessed the dysregulated biological functions in each trio (**Figure S9; Table**

S7), and found that *de novo* or inherited differentially expressed genes in carrier children were

enriched for biological processes (FDR<0.05) that could be related to 39 out of 47 (83.0%)

specific developmental phenotypes observed in the affected children (**Figure 3B**). For example,

inherited gene expression changes for carrier child P1C_01 in family GL_001 were enriched for

nervous system development, neurotransmitter metabolism, neuron projection, and synaptic

transmission functions, while their *de novo* expression changes were enriched for genes involved

in skeletal and muscular development. The inherited changes in neuronal genes could contribute

to the intellectual disability/developmental delay (ID/DD) and speech delay phenotypes observed

in the child, as both parents also had several psychiatric features, while the *de novo* changes in

developmental genes could be related to hypotonia, growth delay, and craniofacial features

uniquely observed in the child. Overall, these results suggest that expression changes of

neurodevelopmental-related genes could account for phenotypic differences among carriers of

the 16p12.1 deletion.

"Second-hit" variants and the 16p12.1 deletion show synergistic effects towards gene

expression

We next investigated whether changes in gene expression could be attributed to "second-hits", or

rare genetic modifiers elsewhere in the genome. Rare variants disrupting protein-coding regions

and nearby regulatory elements have been previously linked to gene expression changes in both

control populations[26,76–78] and disease cohorts, where causal genes may be missed by DNA

sequencing methods[21,23–25]. We hypothesized that "second-hits" by themselves or in combination

with the deletion could contribute to the observed gene expression changes in affected children.

We therefore identified 25 classes of rare gene-disruptive "second-hit" variants from WGS data

for each individual (**Figure S2; Tables S8, S9**), including SNVs and indels in coding and non-

coding regulatory regions (UTRs, introns, and putative promoter, enhancer, and silencer

elements within 50 kbp of a gene) with Phred-like CADD scores[4] >10, and CNVs and short

tandem repeats (STRs) that spanned gene-coding regions. We then calculated the likelihood that

these "second-hit" variants are associated with changes in expression of a proximal gene, as

determined by either differential expression analysis between carrier offspring and their parents

in the 13 trios, or outlier expression analysis among all individuals in the cohort[21,76] (see

Methods). While family-based differential expression analysis detects all expression changes

between affected children and their parents[79], including those due to the downstream effects of

the deletion, outlier analysis more robustly identifies specific effects of "second-hits" towards

larger changes in expression, including synergistic effects in combination with the deletion.

Overall, we observed an average of 285 outlier genes (|z-score| >2) per individual, including

candidate neurodevelopmental genes[47] such as *CTNNB1*, *FOXG1*, *DISC1*, and *ZNF804A* (**Figure**

S6; Tables S6, S10). We found that 10.8% of outlier genes (avg. 31/286 per person) and 11.2%

of differentially expressed genes (avg. 310/2,774 per carrier offspring) were potentially disrupted

by a rare coding or non-coding variant (**Table S9**). Altered expression of genes without such

variants could be due to several factors, such as common variants, DNA methylation events,

downstream effects of other dysregulated genes, or environmental factors.

517   In agreement with previous studies[26,76], we found that genes with outlier expression were

enriched ($p<0.05$, Fisher's exact test) for 6/25 classes of rare variants that directly affected gene-

coding regions, including loss-of-function (LOF), missense, and splice-site SNVs, 5' UTR

overhanging deletions, and 5' UTR overhanging and gene-encapsulating duplications (**Figure**

4A). Similarly, we found that 10/25 variant classes were significantly associated with

differentially expressed genes in carrier offspring for the 13 trios ($p<0.05$, Fisher's exact test),

including coding missense SNVs, duplications overhanging the 5' UTR, and encapsulated

deletions (**Figure S10A**). We further found that outlier genes had significantly higher burdens of

rare variants in aggregate ($p=1.01\times10^{-3}$, one-tailed t-test) and for 7/25 individual classes ($p<0.05$)

compared with non-outlier genes (**Figures S11A-B**). Interestingly, we also found that outlier

genes that were intolerant to variation (pLI score >0.9 or RVIS percentile <20; $p<3.26\times10^{-4}$, two-

tailed t-test) or preferentially expressed in the brain ($p=0.011$) had a higher burden of nearby rare

variants than other outlier genes (**Figure S11C**). We next assessed the effect size of "second-

hits" towards outlier gene expression among carriers and noncarriers of the deletion, and found

significant enrichments ($p<0.05$, Fisher's exact test) of LOF variants, 5' UTR overhanging

duplications, and 5' UTR-disrupting SNVs towards outlier expression in carrier children but not

in carrier parents (**Figure 4B**). We observed similar findings among differentially expressed

genes, where missense SNVs, upstream SNVs, encapsulated deletions, and 5' UTR overhanging

duplications were only likely to alter gene expression in carrier children (**Figure S10B**). Notably,

LOF variants and 5' UTR overhanging duplications were also enriched for outlier expression in

noncarrier parents, suggesting that these classes of "second-hit" variants were more likely to be

deleterious in carrier children when inherited from noncarrier parents (**Figure 4B**). In fact, we

observed five classes of "second-hit" variants in carrier children, including gene-encapsulating

and 3' UTR overhanging deletions, 5' UTR overhanging duplications, and missense and

upstream SNVs, that had significant effects towards outlier expression (p<0.05, Fisher's exact

test) when inherited from the noncarrier parent but not from the carrier parent (**Figure 4C**).

Similarly, we found that LOF variants and SNVs in silencer regions correlated with differential

gene expression in carrier children only when inherited from the noncarrier parent (p<0.05,

Fisher's exact test) (**Figure S10C**). For example, a carrier child in family GL_007, who

exhibited hypotonia and muscle weakness, inherited a deleterious variant from their noncarrier

parent in the 5' UTR of *EIF2AK1* [MIM: 613635], associated with motor dysfunction[80], that

potentially led to downregulation of that gene (**Figure 4D**). Overall, our findings showed that

distinct classes of "second-hit" variants differentially contribute towards changes in gene

expression when inherited in a complex manner from either the carrier or noncarrier parent.

We next investigated whether "second-hit" variants showed synergistic effects towards

expression changes in genes also dysregulated by the 16p12.1 deletion. We found 11 instances of

genes, such as *ANK3, DOCK10,* and *SLC26A1*, that were differentially expressed in all deletion

carriers, showed outlier expression in an individual deletion carrier, and had a nearby variant

(two coding and nine non-coding) inherited from the noncarrier parent (**Figure 4E; Figure S12;**

Table S11). For example, two carrier children in family GL_005 inherited an intronic variant

within *ZEB2* [MIM: 605802] from their noncarrier parent, whose altered dosage is associated

with Mowat-Wilson syndrome[81,82] [MIM: 235730]. While both the 16p12.1 deletion and the

"second-hit" variant individually corresponded with increased *ZEB2* expression, the presence of

both variants in the carrier children resulted in even stronger overexpression of the gene

compared to those with either individual variant (**Figure 4E**). Overexpression of *ZEB2* could

contribute to the Mowat-Wilson like features observed in the carrier child P1C_05, including

ID/DD, seizures, hypotonia, and digestive abnormalities. Similarly, a carrier child in GL_005

inherited a rare variant in a promoter region upstream of the mRNA splicing-associated[83] gene

SNRNP27 from their noncarrier parent. *SNRNP27* is over-expressed in deletion carriers but

under-expressed in both the carrier child and the noncarrier parent, highlighting a case where a

"second-hit" variant reverses an expression change caused by the deletion (**Figure 4E**).

Furthermore, a carrier child in GL_004 shared an intronic variant with two noncarrier relatives in

the gene *SYNJ1* [MIM: 604297], which is associated with synaptic transmission[84] and is under-

expressed in carriers of the deletion. While other individuals with the same variant had normal

SYNJ1 expression, the carrier child exhibited under-expression of the gene compared to both

carriers and noncarriers of the deletion, suggesting that the variant may alter *SYNJ1* expression

only in the presence of the deletion (**Figure 4E**). While it is possible that other variants

elsewhere in the genome could also influence expression levels of these genes, these examples

highlight putative synergistic effects between the 16p12.1 deletion and "second-hit" variants

towards gene expression, where the "second-hit" variants may amplify or reduce the effects of

the CNV.

A broad range of transcriptomic alterations contribute to phenotypic variability of the

16p12.1 deletion

To identify a complete spectrum of gene expression alterations in each individual, we next

evaluated alternative gene splicing, allele-specific expression (ASE), and expression quantitative

trait loci (eQTL) among individuals in our cohort. We first identified an average of 3,267

alternative isoforms present in carrier offspring of the 13 trios compared to their parents (**Tables**

S6, S12), including for several neurodevelopmental-associated genes[47] such as *KANSL1*,

SHANK2, and *SYNGAP1*. After categorizing splicing events by inheritance, we found no

differences between splicing events inherited from carriers (average=1,307) or noncarriers

(average=1,392; p=0.635, two-tailed paired Mann-Whitney test), with fewer changes occurring

de novo (average=568; p=2.44×10^{-4}; **Figure S13A**). We next found enrichments (p<0.05,

Fisher's exact test) for alternative splicing in genes disrupted by "second-hit" splice-site,

intronic, or missense SNVs, interstitial and 3' UTR overhanging deletions, and intronic STRs

(**Figure S13B; Table S9**). We also found that intronic SNVs were more likely to disrupt splicing

in carrier children if they were inherited from the noncarrier parent (p=0.033, Fisher's exact test)

than the carrier parent (p=0.132, **Figure 5A**), while missense SNVs were more likely to lead to

alternative splicing when present in carrier children (p=0.034) than in carrier parents (p=0.168)

(**Figure S13C**). For example, a deleterious splice-site variant in the transcriptional activator

TADA2A [MIM: 602276] led to an alternate isoform (TADA2A-003) in multiple family members of GL_007 that was not observed in GTEx LCL data (**Figure 5B**). TADA2A is a candidate gene within the schizophrenia-associated 17q12 deletion[85] [MIM: 614527], and five out of six family members with the splicing variant have schizophrenia-like clinical features (i.e. hallucinations or delusions), including four deletion carriers and one noncarrier child.

Next, we identified an average of 285 genes with ASE per individual in our cohort (**Tables S6, S13**, including for the neurodevelopmental-associated genes[47] DNMT3A, NSUN2, and HDAC8. ASE events in offspring of the 13 trios were more likely to occur de novo than inherited (p=2.44×10^{-4}, two-tailed paired Mann-Whitney test), in contrast to differential expression and alternative splicing events (**Figure S14A**). Genes with ASE have previously been shown to have a higher burden of nearby rare deleterious variants[76], and the pathogenicity of a gene with ASE increases with the presence of a deleterious variant on the overexpressed allele[78]. In our cohort, we found five ASE events in carrier children that led to overexpression of a deleterious "second-hit" coding variant (**Figure S14B**). For example, two carrier children with autism in family GL_007 showed overexpression of a deleterious "second-hit" missense variant in the candidate autism gene[86] CARD11 [MIM: 607210], which was inherited from their noncarrier parent (**Figure 5C**).

We further performed eQTL discovery analysis to identify variants statistically correlated with expression changes in our cohort, agnostic to variant pathogenicity or population frequency. We identified 21 eQTLs which modulated the expression of 23 eGenes (**Figure S15A; Table S14**), of which 19 were not present in GTEx LCL data, representing unique loci in our cohort. Carrier children showed a trend (p=0.107, two-tailed Mann-Whitney test) towards carrying a higher number of minor eQTLs alleles (average=4.3/person) than their carrier parents (average=3.2/person) (**Figure S15B; Table S6**). Furthermore, several eGenes had biological functions related to neuronal processes[87–89] (**Table S14**), including SERPINF1, BEGAIN, and ARFGEF2. For example, we identified a relatively rare eQTL (allele frequency = 0.015) for overexpression of ARHGAP39 [MIM: 615880], a key regulator of neurogenesis and dendrite morphology associated with learning and memory[90] (**Figure 5D**). The eQTL minor allele, located in a transcription factor binding cluster, was only observed in a carrier child and their noncarrier parent within GL_007, who both presented with neuropsychiatric phenotypes.

To assess the joint contributions of each type of expression change among the individuals

in our cohort, we assessed correlations between the numbers of gene expression changes

assessed in our study by inheritance pattern (**Figure 5E**). We observed nine significant positive

correlations ($p<0.05$, Pearson correlation) between pairs of gene expression changes in each

person, which often shared the same inheritance patterns. For example, the number of genes with

differential expression strongly correlated with the number of genes with alternative splicing

when inherited from the carrier parent ($r=0.93$, $p=4.41\times10^{-6}$) or non-carrier parent ($r=0.91$,

$p=1.37\times10^{-5}$). Together, the correlations between transcriptomic alterations suggest that different

types of gene expression changes could be inherited together, potentially due to the same

"second-hit" variants disrupting expression of similar genes and biological pathways, as is

observed for signals in genome-wide association studies[91].

639   Genes with "second-hit" variants and expression changes show strong connectivity in a

640   brain-specific network

Finally, to determine whether associations between transcriptomic changes and "second-hit"

variants in LCL samples were also relevant in the nervous system, we assessed connectivity

patterns of genes with "second-hit" variants and altered gene expression using a brain-specific

interaction network[66,67]. We generated individual networks for carrier offspring in the 13 trios,

and calculated shortest distances between genes with protein-coding "second-hit" variants and

genes with observed expression changes in each offspring (see Methods). We found that the

average shortest distances between genes with "second-hits" and expression changes were

significantly smaller in 7/13 offspring than those derived from permuted networks ($p<0.05$, one-

tailed z-test with $n=100$ permutations). In fact, networks for offspring in aggregate had

significantly smaller shortest distances than the shortest distances from the sets of permuted

networks ($p=4.88\times10^{-4}$, two-tailed paired Mann-Whitney test) (**Figure 5F**). These data indicate

that "second-hit" variants closely interact with genes with expression changes in the brain,

suggesting a potential mechanism for how gene expression changes that underlie developmental

phenotypes can be influenced by "second-hit" variants in the genome.

DISCUSSION

We previously described a two-hit model for the 16p12.1 deletion, where the presence of both the deletion and "second-hit" variants determine the phenotypic trajectory of affected children[2,15]. Here, we propose a mechanism for how the deletion and "second-hits" jointly interact to alter clinical phenotypes by way of the transcriptome. We found that the 16p12.1 deletion itself disrupts the expression of genes across the genome through direct effects, such as chromatin interactions, and through indirect effects, such as downstream genetic interactions (**Figure 6**). For example, chromatin interactions were observed between regions within the 16p12.1 deletion and flanking genes such as *DNAH3*, and 1,493 genes outside of chromosome 16 were also dysregulated in deletion carriers. Similarly, we found that "second-hits" disrupt gene expression through both direct and indirect mechanisms. Genes with nearby "second-hit" variants were more likely to exhibit outlier expression and alternative splicing, and genes with "second-hits" were more closely connected to genes with expression changes in a brain-specific network than random sets of genes in permuted networks. In fact, we observed 11 examples of combined effects of the deletion and "second-hit" variants towards expression in our cohort, including the candidate neurodevelopmental genes *SYNJ1* and *ZEB2*. These synergistic effects towards gene expression are similar to those previously observed for eQTLs[92] and HLA alleles[93], except that these effects are potentially due to the combined effects of rare deleterious variants. These results suggest that the 16p12.1 deletion and the "second-hit" variants jointly mold the shape of the transcriptome, resulting in strong dysregulation of developmental genes and contributing to neuropsychiatric features in 16p12.1 deletion carriers.

Results from our study align with previous studies, which found that rare variants of different classes have varying effect sizes towards gene expression[26,76]. Our study extends this paradigm by identifying classes of rare "second-hit" variants whose contributions to gene expression changes differ by inheritance pattern. We found that high-effect variants, such as duplications and loss-of-function variants, always cause expression changes regardless of parent-of-origin, and splice-site variants almost always lead to changes in isoform expression. In contrast, medium-effect variants, including missense, intronic, and upstream SNVs, were more strongly associated with gene expression changes when inherited from the noncarrier parent than the carrier parent. These findings indicate that the noncarrier parents are more likely to pass gene expression-altering "second-hit" variants down to their carrier children, providing an explanation

for why children with the deletion have more severe phenotypic manifestations than their carrier

parents (**Figure 6**). Future family-based transcriptome studies with larger sample sizes may be

able to pinpoint specific rare variants within dysregulated genes that are associated with distinct

phenotypes in the carrier children.

We also identified putative biological and developmental pathways disrupted by both the

deletion and "second-hit" variants. For example, we found that genes differentially expressed by

the deletion were preferentially expressed in multiple brain tissues during development, and were

enriched for core signaling and developmental pathways. In fact, knockdown of individual

homologs of 16p12.1 genes in *Drosophila melanogaster* models showed neuronal phenotypes

and transcriptome disruptions, suggesting that the individual effects of multiple genes in the

deletion sensitize the genome for neuropsychiatric phenotypes[94]. Interestingly, we found several

examples of biological functions and mechanisms that were simultaneously dysregulated by both

the deletion and "second-hit" variants in the carrier children. For example, most carrier children

inherited differential expression in genes enriched for nervous system development, cell

adhesion, signaling, and locomotion from both their carrier and noncarrier parents. These results

provide insights into how the deletion and "second-hit" variants synergistically dysregulate

genes and pathways related to development, ultimately contributing towards a wide range of

developmental phenotypes observed in children with the deletion.

Some limitations can be noted in the context of our study. *First*, we investigated gene

expression changes within patient-derived LCL samples, which have reduced relevance for brain

expression. However, over 80% of genes expressed in GTEx brain samples, as well as over 70%

of neurodevelopmental disease genes, were expressed in our LCL samples (**Figures S4A-B**).

Nevertheless, repeating the study in tissues that are implicated in neurodevelopmental disorders,

potentially using patient-derived reprogrammed neuronal progenitor cells, would verify the

associations between variants, expression changes, and clinical features. *Second*, we have a

relatively small cohort of 32 individuals within five families. It would be useful to determine

whether the identified associations are strengthened in a larger cohort. Phenotypically more

diverse cohorts would also allow for performing additional correlations between gene expression

changes and specific clinical features, such as whether more outlier genes are present among

families with stronger histories of neuropsychiatric disease. Overall, our work supports a model

for complex disorders, where combinatorial effects of multiple variants with different effect sizes

affect expression of genes in developmental pathways, which further influence the expressivity

of clinical features. These results exemplify that family-based transcriptome studies, similar to

family-based genome studies, can help explain changes in phenotypes from parents to children

and between siblings, especially in complex disorders with a high degree of intra- and inter-

familial variability.

SUPPLEMENTAL DATA

Supplemental Data include 15 figures and 14 tables in Excel format.

ACKNOWLEDGEMENTS

We are grateful to all members of the five families who participated in this study. We also thank Sherryann Wert, Melissa Berkowitz, and Tara Schmidlen (Coriell Institute) for their assistance in the generation of stable LCL lines; Craig Praul and Yuka Imamura (Penn State Genomics Core Facility) for assistance with designing the RNA-sequencing strategy; Casey Brown (UPenn), Istvan Albert (Penn State), and Qunhua Li (Penn State) for assistance with statistical analysis of transcriptome data; and Jesse Gillis (CSHL), Dajiang Liu (Penn State), and members of the Girirajan lab for useful discussions and critical reading of the manuscript. We appreciate access to data from the Genotype-Tissue Expression (GTEx) Project, which was supported by the Common Fund of the Office of the Director of the National Institutes of Health, and by NCI, NHGRI, NHLBI, NIDA, NIMH, and NINDS. The data used for the analyses described in this manuscript were obtained from the GTEx Portal in January 2020. This work was supported by NIH R01-GM121907 and resources from the Huck Institutes of the Life Sciences to S.G. M.J. was supported by NIH T32-GM102057. A.T. was supported by NIH T32-LM012415. L.P. was supported by Fulbright Commission Uruguay-ANII. A.K. was supported by NIH R35-GM128765. The funding bodies had no role in data collection, analysis, or interpretation.

WEB RESOURCES

BrainSpan Atlas, https://www.brainspan.org/

CADD, https://cadd.gs.washington.edu/

ChromHMM, https://www.encodeproject.org/software/chromhmm/

DBD Gene Database, https://dbd.geisingeradmi.org/

DDD, https://www.ddduk.org/

DDG2P, https://decipher.sanger.ac.uk/ddd/ddgenes

Gene Ontology, http://geneontology.org/

gnomAD/pLI, https://gnomad.broadinstitute.org/

GTEx, https://www.gtexportal.org/home/

MutationTaster, http://www.mutationtaster.org/

OMIM, https://www.omim.org/

RVIS, http://genic-intolerance.org/

SFARI Gene, https://gene.sfari.org/

DATA AND CODE AVAILABILITY

Patient-derived LCL samples generated in this study are available at the NIGMS Human Genetic

Cell Repository at the Coriell Institute (https://www.coriell.org/1/NIGMS). Whole genome

sequencing and RNA-sequencing data generated in this study will be available at the NCBI

dbGaP database (https://www.ncbi.nlm.nih.gov/gap/). Accession numbers for LCL samples and

sequencing data are provided in **Table S2**. All code generated for this project, including

pipelines for running bioinformatic software and custom analysis scripts, is available at

https://github.com/girirajanlab/16p12.1_rnaseq_analysis and

https://github.com/girirajanlab/16p12.1_WGS_analysis.

REFERENCES

1. Eichler, E.E. (2019). Genetic Variation, Comparative Genomics, and the Diagnosis of Disease. N. Engl. J. Med. *381*, 64–74.

2. Pizzo, L., Jensen, M., Polyak, A., Rosenfeld, J.A., Mannik, K., Krishnan, A., McCready, E., Pichon, O., Le Caignec, C., Van Dijck, A., et al. (2019). Rare variants in the genetic background modulate cognitive and developmental phenotypes in individuals carrying disease-associated variants. Genet. Med. *21*, 816–825.

3. Berkovic, S.F., Scheffer, I.E., Petrou, S., Delanty, N., Dixon-Salazar, T.J., Dlugos, D.J., Helbig, I., Frankel, W.N., Goldstein, D.B., Heinzen, E.L., et al. (2015). A roadmap for precision medicine in the epilepsies. Lancet Neurol. *14*, 1219–1228.

4. Kircher, M., Witten, D.M., Jain, P., O'roak, B.J., Cooper, G.M., and Shendure, J. (2014). A general framework for estimating the relative pathogenicity of human genetic variants. Nat. Genet. *46*, 310–315.

5. Schaaf, C.P., Betancur, C., Yuen, R.K.C., Parr, J.R., Skuse, D.H., Gallagher, L., Bernier, R.A., Buchanan, J.A., Buxbaum, J.D., Chen, C.A., et al. (2020). A framework for an evidence-based gene list relevant to autism spectrum disorder. Nat. Rev. Genet. *21*, 367–376.

6. Ardlie, K.G., DeLuca, D.S., Segrè, A. V., Sullivan, T.J., Young, T.R., Gelfand, E.T., Trowbridge, C.A., Maller, J.B., Tukiainen, T., Lek, M., et al. (2015). The Genotype-Tissue Expression (GTEx) pilot analysis: Multitissue gene regulation in humans. Science *348*, 648–660.

7. Kavanagh, D.H., Tansey, K.E., O'Donovan, M.C., and Owen, M.J. (2015). Schizophrenia genetics: Emerging themes for a complex disorder. Mol. Psychiatry *20*, 72–76.

8. Petrovski, S., Wang, Q., Heinzen, E.L., Allen, A.S., and Goldstein, D.B. (2013). Genic Intolerance to Functional Variation and the Interpretation of Personal Genomes. PLoS Genet. *9*, e1003709.

9. Manolio, T.A., Collins, F.S., Cox, N.J., Goldstein, D.B., Hindorff, L.A., Hunter, D.J., McCarthy, M.I., Ramos, E.M., Cardon, L.R., Chakravarti, A., et al. (2009). Finding the missing heritability of complex diseases. Nature *461*, 747–753.

10. Lek, M., Karczewski, K.J., Minikel, E. V., Samocha, K.E., Banks, E., Fennell, T., O'Donnell-Luria, A.H., Ware, J.S., Hill, A.J., Cummings, B.B., et al. (2016). Analysis of protein-coding genetic variation in 60,706 humans. Nature *536*, 285–291.

11. Boyle, E.A., Li, Y.I., and Pritchard, J.K. (2017). An Expanded View of Complex Traits: From Polygenic to Omnigenic. Cell *169*, 1177–1186.

12. Purcell, S.M., Moran, J.L., Fromer, M., Ruderfer, D., Solovieff, N., Roussos, P., O'Dushlaine, C., Chambert, K., Bergen, S.E., Kähler, A., et al. (2014). A polygenic burden of rare disruptive mutations in schizophrenia. Nature *506*, 185–190.

13. Coe, B.P., Girirajan, S., and Eichler, E.E. (2012). A genetic model for neurodevelopmental disease. Curr. Opin. Neurobiol. *22*, 829–836.

14. Girirajan, S., Rosenfeld, J.A., Cooper, G.M., Antonacci, F., Siswara, P., Itsara, A., Vives, L.,

Walsh, T., McCarthy, S.E., Baker, C., et al. (2010). A recurrent 16p12.1 microdeletion supports a two-hit model for severe developmental delay. Nat. Genet. *42*, 203–209.

15. Girirajan, S., Rosenfeld, J.A., Coe, B.P., Parikh, S., Friedman, N., Goldstein, A., Filipink, R.A., McConnell, J.S., Angle, B., Meschino, W.S., et al. (2012). Phenotypic Heterogeneity of Genomic Disorders and Rare Copy-Number Variants. N. Engl. J. Med. *367*, 1321–1331.

16. Rees, E., Kendall, K., Pardiñas, A.F., Legge, S.E., Pocklington, A., Escott-Price, V., MacCabe, J.H., Collier, D.A., Holmans, P., O'Donovan, M.C., et al. (2016). Analysis of intellectual disability copy number variants for association with schizophrenia. JAMA Psychiatry *73*, 963–969.

17. Mefford, H.C., Muhle, H., Ostertag, P., von Spiczak, S., Buysse, K., Baker, C., Franke, A., Malafosse, A., Genton, P., Thomas, P., et al. (2010). Genome-wide copy number variation in epilepsy: novel susceptibility loci in idiopathic generalized and focal epilepsies. PLoS Genet. *6*, e1000962.

18. Stefansson, H., Meyer-Lindenberg, A., Steinberg, S., Magnusdottir, B., Morgen, K., Arnarsdottir, S., Bjornsdottir, G., Walters, G.B., Jonsdottir, G.A., Doyle, O.M., et al. (2014). CNVs conferring risk of autism or schizophrenia affect cognition in controls. Nature *505*, 361–366.

19. Girirajan, S., and Eichler, E.E. (2010). Phenotypic variability and genetic susceptibility to genomic disorders. Hum. Mol. Genet. *19*, R176-87.

20. Merla, G., Howald, C., Henrichsen, C.N., Lyle, R., Wyss, C., Zabot, M.T., Antonarakis, S.E., and Reymond, A. (2006). Submicroscopic deletion in patients with Williams-Beuren syndrome influences expression levels of the nonhemizygous flanking genes. Am. J. Hum. Genet. *79*, 332–341.

21. Luo, R., Sanders, S.J., Tian, Y., Voineagu, I., Huang, N., Chu, S.H., Klei, L., Cai, C., Ou, J., Lowe, J.K., et al. (2012). Genome-wide transcriptome profiling reveals the functional impact of rare de novo and recurrent CNVs in autism spectrum disorders. Am. J. Hum. Genet. *91*, 38–55.

22. Blumenthal, I., Ragavendran, A., Erdin, S., Klei, L., Sugathan, A., Guide, J.R., Manavalan, P., Zhou, J.Q., Wheeler, V.C., Levin, J.Z., et al. (2014). Transcriptional consequences of 16p11.2 deletion and duplication in mouse cortex and multiplex autism families. Am. J. Hum. Genet. *94*, 870–883.

23. Frésard, L., Smail, C., Ferraro, N.M., Teran, N.A., Li, X., Smith, K.S., Bonner, D., Kernohan, K.D., Marwaha, S., Zappala, Z., et al. (2019). Identification of rare-disease genes using blood transcriptome sequencing and large control cohorts. Nat. Med. *25*, 911–919.

24. Gonorazky, H.D., Naumenko, S., Ramani, A.K., Nelakuditi, V., Mashouri, P., Wang, P., Kao, D., Ohri, K., Viththiyapaskaran, S., Tarnopolsky, M.A., et al. (2019). Expanding the Boundaries of RNA Sequencing as a Diagnostic Tool for Rare Mendelian Disease. Am. J. Hum. Genet. *104*, 466–483.

25. Mohammadi, P., Castel, S.E., Cummings, B.B., Einson, J., Sousa, C., Hoffman, P., Donkervoort, S., Jiang, Z., Mohassel, P., Reghan Foley, A., et al. (2019). Genetic regulatory variation in populations informs transcriptome analysis in rare disease. Science *366*, 351–356.

26. Pala, M., Zappala, Z., Marongiu, M., Li, X., Davis, J.R., Cusano, R., Crobu, F., Kukurba,
K.R., Gloudemans, M.J., Reinier, F., et al. (2017). Population- and individual-specific regulatory
variation in Sardinia. Nat. Genet. *49*, 700–707.

27. Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: A flexible trimmer for
Illumina sequence data. Bioinformatics *30*, 2114–2120.

28. Li, H., and Durbin, R. (2010). Fast and accurate long-read alignment with Burrows-Wheeler
transform. Bioinformatics *26*, 589–595.

29. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis,
G., and Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics
25, 2078–2079.

30. Poplin, R., Ruano-Rubio, V., DePristo, M.A., Fennell, T.J., Carneiro, M.O., Auwera, G.A.
Van der, Kling, D.E., Gauthier, L.D., Levy-Moonshine, A., Roazen, D., et al. (2017). Scaling
accurate genetic variant discovery to tens of thousands of samples. BioRxiv 201178.

31. Wang, K., Li, M., and Hakonarson, H. (2010). ANNOVAR: Functional annotation of genetic
variants from high-throughput sequencing data. Nucleic Acids Res. *38*, e164.

32. Krumm, N., Turner, T.N., Baker, C., Vives, L., Mohajeri, K., Witherspoon, K., Raja, A.,
Coe, B.P., Stessman, H.A., He, Z.X., et al. (2015). Excess of rare, inherited truncating mutations
in autism. Nat. Genet. *47*, 582–588.

33. Karczewski, K.J., Francioli, L.C., Tiao, G., Cummings, B.B., Alföldi, J., Wang, Q., Collins,
R.L., Laricchia, K.M., Ganna, A., Birnbaum, D.P., et al. (2020). The mutational constraint
spectrum quantified from variation in 141,456 humans. Nature *581*, 434–443.

34. Schwarz, J.M., Cooper, D.N., Schuelke, M., and Seelow, D. (2014). MutationTaster2:
Mutation prediction for the deep-sequencing age. Nat. Methods *11*, 361–362.

35. Ernst, J., Kheradpour, P., Mikkelsen, T.S., Shoresh, N., Ward, L.D., Epstein, C.B., Zhang,
X., Wang, L., Issner, R., Coyne, M., et al. (2011). Mapping and analysis of chromatin state
dynamics in nine human cell types. Nature *473*, 43–49.

36. Wang, K., Li, M., Hadley, D., Liu, R., Glessner, J., Grant, S.F.A., Hakonarson, H., and
Bucan, M. (2007). PennCNV: An integrated hidden Markov model designed for high-resolution
copy number variation detection in whole-genome SNP genotyping data. Genome Res. *17*,
1665–1674.

37. Abyzov, A., Urban, A.E., Snyder, M., and Gerstein, M. (2011). CNVnator: An approach to
discover, genotype, and characterize typical and atypical CNVs from family and population
genome sequencing. Genome Res. *21*, 974–984.

38. Rausch, T., Zichner, T., Schlattl, A., Stütz, A.M., Benes, V., and Korbel, J.O. (2012).
DELLY: Structural variant discovery by integrated paired-end and split-read analysis.
Bioinformatics *28*, 333–339.

39. Layer, R.M., Chiang, C., Quinlan, A.R., and Hall, I.M. (2014). LUMPY: A probabilistic
framework for structural variant discovery. Genome Biol. *15*, R84.

40. Chen, X., Schulz-Trieglaff, O., Shaw, R., Barnes, B., Schlesinger, F., Källberg, M., Cox,
A.J., Kruglyak, S., and Saunders, C.T. (2016). Manta: Rapid detection of structural variants and
indels for germline and cancer sequencing applications. Bioinformatics *32*, 1220–1222.

41. Mousavi, N., Shleizer-Burko, S., Yanicky, R., and Gymrek, M. (2019). Profiling the genome-
wide landscape of tandem repeat expansions. Nucleic Acids Res. *47*, e90.

42. Mousavi, N., Margoliash, J., Pusarla, N., Saini, S., Yanicky, R., and Gymrek, M. (2020).
TRTools: a toolkit for genome-wide analysis of tandem repeats. Bioinformatics ePub.

43. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,
M., and Gingeras, T.R. (2013). STAR: Ultrafast universal RNA-seq aligner. Bioinformatics *29*,
15–21.

44. Wang, L., Nie, J., Sicotte, H., Li, Y., Eckel-Passow, J.E., Dasari, S., Vedell, P.T., Barman,
P., Wang, L., Weinshiboum, R., et al. (2016). Measure transcript integrity using RNA-seq data.
BMC Bioinformatics *17*, 58.

45. Li, B., and Dewey, C.N. (2011). RSEM: Accurate transcript quantification from RNA-Seq
data with or without a reference genome. BMC Bioinformatics *12*, 323.

46. Deluca, D.S., Levin, J.Z., Sivachenko, A., Fennell, T., Nazaire, M.D., Williams, C., Reich,
M., Winckler, W., and Getz, G. (2012). RNA-SeQC: RNA-seq metrics for quality control and
process optimization. Bioinformatics *28*, 1530–1532.

47. Gonzalez-Mantilla, A.J., Moreno-De-Luca, A., Ledbetter, D.H., and Martin, C.L. (2016). A
cross-disorder method to identify novel candidate genes for developmental brain disorders.
JAMA Psychiatry *73*, 275–283.

48. Abrahams, B.S., Arking, D.E., Campbell, D.B., Mefford, H.C., Morrow, E.M., Weiss, L.A.,
Menashe, I., Wadkins, T., Banerjee-Basu, S., and Packer, A. (2013). SFARI Gene 2.0: A
community-driven knowledgebase for the autism spectrum disorders (ASDs). Mol. Autism *4*, 36.

49. Firth, H. V., Richards, S.M., Bevan, A.P., Clayton, S., Corpas, M., Rajan, D., Vooren, S.
Van, Moreau, Y., Pettett, R.M., and Carter, N.P. (2009). DECIPHER: Database of Chromosomal
Imbalance and Phenotype in Humans Using Ensembl Resources. Am. J. Hum. Genet. *84*, 524–
533.

50. Wright, C.F., Fitzgerald, T.W., Jones, W.D., Clayton, S., McRae, J.F., Van Kogelenberg, M.,
King, D.A., Ambridge, K., Barrett, D.M., Bayzetinova, T., et al. (2015). Genetic diagnosis of
developmental disorders in the DDD study: A scalable analysis of genome-wide research data.
Lancet *385*, 1305–1314.

51. Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2009). edgeR: A Bioconductor package
for differential expression analysis of digital gene expression data. Bioinformatics *26*, 139–140.

52. Sun, S., Zhu, J., Mozaffari, S., Ober, C., Chen, M., and Zhou, X. (2019). Heritability
estimation and differential analysis of count data with generalized linear mixed models in
genomic sequencing studies. Bioinformatics *35*, 487–496.

53. Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A.R., Bender, D., Maller, J.,
Sklar, P., De Bakker, P.I.W., Daly, M.J., et al. (2007). PLINK: A tool set for whole-genome

association and population-based linkage analyses. Am. J. Hum. Genet. *81*, 559–575.

54. Zhou, X., and Stephens, M. (2012). Genome-wide efficient mixed-model analysis for
association studies. Nat. Genet. *44*, 821–824.

55. Stegle, O., Parts, L., Piipari, M., Winn, J., and Durbin, R. (2012). Using probabilistic
estimation of expression residuals (PEER) to obtain increased power and interpretability of gene
expression analyses. Nat. Protoc. *7*, 500–507.

56. Young, M.D., Wakefield, M.J., Smyth, G.K., and Oshlack, A. (2010). Gene ontology
analysis for RNA-seq: accounting for selection bias. Genome Biol. *11*, R14.

57. The Gene Ontology Consortium (2019). The Gene Ontology Resource: 20 years and still
GOing strong. Nucleic Acids Res. *47*, D330–D338.

58. Miller, J.A., Ding, S.-L., Sunkin, S.M., Smith, K.A., Ng, L., Szafer, A., Ebbert, A., Riley,
Z.L., Royall, J.J., Aiona, K., et al. (2014). Transcriptional landscape of the prenatal human brain.
Nature *508*, 199–206.

59. Langfelder, P., and Horvath, S. (2008). WGCNA: An R package for weighted correlation
network analysis. BMC Bioinformatics *9*, 559.

60. Soneson, C., Love, M.I., and Robinson, M.D. (2016). Differential analyses for RNA-seq:
Transcript-level estimates improve gene-level inferences. F1000Res. *4*, 1521.

61. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550.

62. Johnson, W.E., Li, C., and Rabinovic, A. (2007). Adjusting batch effects in microarray
expression data using empirical Bayes methods. Biostatistics *8*, 118–127.

63. Castel, S.E., Mohammadi, P., Chung, W.K., Shen, Y., and Lappalainen, T. (2016). Rare
variant phasing and haplotypic expression from RNA sequencing with phASER. Nat. Commun.
7, 12817.

64. Martin, M., Patterson, M., Garg, S., O Fischer, S., Pisanti, N., Klau, G., Schöenhuth, A., and
Marschall, T. (2016). WhatsHap: fast and accurate read-based phasing. BioRxiv 085050.

65. Delaneau, O., Ongen, H., Brown, A.A., Fort, A., Panousis, N.I., and Dermitzakis, E.T.
(2017). A complete tool set for molecular QTL discovery and analysis. Nat. Commun. *8*, 15452.

66. Greene, C.S., Krishnan, A., Wong, A.K., Ricciotti, E., Zelaya, R.A., Himmelstein, D.S.,
Zhang, R., Hartmann, B.M., Zaslavsky, E., Sealfon, S.C., et al. (2015). Understanding
multicellular function and disease with human tissue-specific networks. Nat. Genet. *47*, 569–576.

67. Krishnan, A., Zhang, R., Yao, V., Theesfeld, C.L., Wong, A.K., Tadych, A., Volfovsky, N.,
Packer, A., Lash, A., and Troyanskaya, O.G. (2016). Genome-wide prediction and functional
characterization of the genetic basis of autism spectrum disorder. Nat. Neurosci. *19*, 1454–1462.

68. Hagberg, A.A., Schult, D.A., and Swart, P.J. (2008). Exploring network structure, dynamics,
and function using NetworkX. In 7th Python in Science Conference (SciPy 2008), pp. 11–15.

69. Harris, C.R., Millman, K.J., van der Walt, S.J., Gommers, R., Virtanen, P., Cournapeau, D.,

Wieser, E., Taylor, J., Berg, S., Smith, N.J., et al. (2020). Array programming with NumPy. Nature *585*, 357–362.

70. Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., et al. (2020). SciPy 1.0: fundamental algorithms for scientific computing in Python. Nat. Methods *17*, 261–272.

71. McKinney, W. (2010). Data Structures for Statistical Computing in Python. In Proceedings of the 9th Python in Science Conference, pp. 56–61.

72. Rao, S.S.P., Huntley, M.H., Durand, N.C., Stamenova, E.K., Bochkov, I.D., Robinson, J.T., Sanborn, A.L., Machol, I., Omer, A.D., Lander, E.S., et al. (2014). A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping. Cell *159*, 1665–1680.

73. Werling, D.M., Pochareddy, S., Choi, J., An, J.Y., Sheppard, B., Peng, M., Li, Z., Dastmalchi, C., Santpere, G., Sousa, A.M.M., et al. (2020). Whole-Genome and RNA Sequencing Reveal Variation and Transcriptomic Coordination in the Developing Human Prefrontal Cortex. Cell Rep. *31*, 107489.

74. Courchesne, E., Gazestani, V.H., and Lewis, N.E. (2020). Prenatal Origins of ASD: The When, What, and How of ASD Development. Trends Neurosci. *43*, 326–342.

75. Skene, N.G., Roy, M., and Grant, S.G. (2017). A genomic lifespan program that reorganises the young adult brain is targeted in schizophrenia. Elife *6*, e17915.

76. Li, X., Kim, Y., Tsang, E.K., Davis, J.R., Damani, F.N., Chiang, C., Hess, G.T., Zappala, Z., Strober, B.J., Scott, A.J., et al. (2017). The impact of rare variation on gene expression across tissues. Nature *550*, 239–243.

77. Zhao, J., Akinsanmi, I., Arafat, D., Cradick, T.J., Lee, C.M., Banskota, S., Marigorta, U.M., Bao, G., and Gibson, G. (2016). A Burden of Rare Variants Associated with Extremes of Gene Expression in Human Peripheral Blood. Am. J. Hum. Genet. *98*, 299–309.

78. Castel, S.E., Cervera, A., Mohammadi, P., Aguet, F., Reverter, F., Wolman, A., Guigo, R., Iossifov, I., Vasileva, A., and Lappalainen, T. (2018). Modified penetrance of coding variants by cis-regulatory variation contributes to disease risk. Nat. Genet. *50*, 1327–1334.

79. Ballouz, S., Dörfel, M., Crow, M., Crain, J., Faivre, L., Keegan, C.E., Kitsiou-Tzeli, S., Tzetis, M., Lyon, G.J., and Gillis, J. (2017). Not by systems alone: Replicability assessment of disease expression signals. BioRxiv 128439.

80. Mao, D., Reuter, C.M., Ruzhnikov, M.R.Z., Beck, A.E., Farrow, E.G., Emrick, L.T., Rosenfeld, J.A., Mackenzie, K.M., Robak, L., Wheeler, M.T., et al. (2020). De novo EIF2AK1 and EIF2AK2 Variants Are Associated with Developmental Delay, Leukoencephalopathy, and Neurologic Decompensation. Am. J. Hum. Genet. *106*, 570–583.

81. Yuan, H., Zhang, L., Chen, M., Zhu, J., Meng, Z., and Liang, L. (2015). A de novo triplication on 2q22.3 including the entire ZEB2 gene associated with global developmental delay, multiple congenital anomalies and behavioral abnormalities. Mol. Cytogenet. *8*, 99.

82. Baxter, A.L., Vivian, J.L., Hagelstrom, R.T., Hossain, W., Golden, W.L., Wassman, E.R., Vanzo, R.J., and Butler, M.G. (2017). A Novel Partial Duplication of ZEB2 and Review of

997 ZEB2 Involvement in Mowat-Wilson Syndrome. Mol. Syndromol. *8*, 211–218.

998 83. Zahler, A.M., Rogel, L.E., Glover, M.L., Yitiz, S., Ragle, J.M., and Katzman, S. (2018).
999 SNRP-27, the C. Elegans homolog of the tri-snRNP 27K protein, has a role in 5′ splice site
1000 positioning in the spliceosome. RNA *24*, 1314–1325.

1001 84. Saito, T., Guan, F., Papolos, D.F., Lau, S., Klein, M., Fann, C.S.J., and Lachman, H.M.
1002 (2001). Mutation analysis of SYNJ1: A possible candidate gene for chromosome 21q22-linked
1003 bipolar disorder. Mol. Psychiatry *6*, 387–395.

1004 85. Mitchel, M.W., Moreno-De-Luca, D., Myers, S.M., Finucane, B., Ledbetter, D.H., and
1005 Martin, C.L. (2016). 17q12 Recurrent Deletion Syndrome. In GeneReviews®, M.P. Adam, H.H.
1006 Ardinger, R.A. Pagon, S.E. Wallace, L.J. Bean, K. Stephens, and A. Amemiya, eds. (University
1007 of Washington, Seattle).

1008 86. Lim, E.T., Uddin, M., De Rubeis, S., Chan, Y., Kamumbu, A.S., Zhang, X., D'Gama, A.M.,
1009 Kim, S.N., Hill, R.S., Goldberg, A.P., et al. (2017). Rates, distribution and implications of
1010 postzygotic mosaic mutations in autism spectrum disorder. Nat. Neurosci. *20*, 1217–1224.

1011 87. Ramírez-Castillejo, C., Sánchez-Sánchez, F., Andreu-Agulló, C., Ferrón, S.R., Aroca-
1012 Aguilar, J.D., Sánchez, P., Mira, H., Escribano, J., and Fariñas, I. (2006). Pigment epithelium-
1013 derived factor is a niche signal for neural stem cell renewal. Nat. Neurosci. *9*, 331–339.

1014 88. Yao, I., Iida, J., Nishimura, W., and Hata, Y. (2002). Synaptic and nuclear localization of
1015 brain-enriched guanylate kinase-associated protein. J. Neurosci. *22*, 5354–5364.

1016 89. De Wit, M.C.Y., De Coo, I.F.M., Halley, D.J.J., Lequin, M.H., and Mancini, G.M.S. (2009).
1017 Movement disorder and neuronal migration disorder due to ARFGEF2 mutation. Neurogenetics
1018 *10*, 333–336.

1019 90. Nowak, F. V. (2018). Porf-2 = arhgap39 = vilse: A pivotal role in neurodevelopment,
1020 learning and memory. ENeuro *5*, e0082-18.2018.

1021 91. Dickson, S.P., Wang, K., Krantz, I., Hakonarson, H., and Goldstein, D.B. (2010). Rare
1022 Variants Create Synthetic Genome-Wide Associations. PLoS Biol. *8*, e1000294.

1023 92. Powell, J.E., Henders, A.K., McRae, A.F., Kim, J., Hemani, G., Martin, N.G., Dermitzakis,
1024 E.T., Gibson, G., Montgomery, G.W., and Visscher, P.M. (2013). Congruence of Additive and
1025 Non-Additive Effects on Gene Expression Estimated from Pedigree and SNP Data. PLoS Genet.
1026 *9*, e1003502.

1027 93. Lenz, T.L., Deutsch, A.J., Han, B., Hu, X., Okada, Y., Eyre, S., Knapp, M., Zhernakova, A.,
1028 Huizinga, T.W.J., Abecasis, G., et al. (2015). Widespread non-additive and interaction effects
1029 within HLA loci modulate the risk of autoimmune diseases. Nat. Genet. *47*, 1085–1090.

1030 94. Pizzo, L., Lasser, M., Yusuff, T., Jensen, M., Ingraham, P., Huber, E., Singh, M.D.,
1031 Monahan, C., Iyer, J., Desai, I., et al. (2021). Functional assessment of the "two-hit" model for
1032 neurodevelopmental defects in Drosophila and X. laevis. BioRxiv 2020.09.14.295923.

1033 95. Wang, Y., Song, F., Zhang, B., Zhang, L., Xu, J., Kuang, D., Li, D., Choudhary, M.N.K., Li,
1034 Y., Hu, M., et al. (2018). The 3D Genome Browser: A web-based browser for visualizing 3D
1035 genome organization and long-range chromatin interactions. Genome Biol. *19*, 151.

FIGURE TITLES AND LEGENDS

Figure 1. Overview of experimental design. We performed whole genome sequencing, RNA sequencing, and clinical phenotyping on five large families (32 total individuals) with the 16p12.1 deletion, indicated with red asterisks in the pedigrees. Children (green) and adults (blue) in the pedigrees are shaded by phenotypic severity, with white indicating no clinical features, lighter shades indicating mild features, darker shades indicating severe features, and grey indicating no phenotypic data available, as detailed in **Figure S1** and **Table S1**. We then performed multiple analyses to assess the role of the deletion and rare "second-hit" variants towards the observed transcriptomic changes and developmental phenotypes, including differential expression between carriers and noncarriers of the deletion, differential expression between parents and carrier offspring in 13 trios from the five families, outlier gene expression among all individuals, identification of additional transcriptomic alterations such as alternative splicing and allele-specific expression, and gene interaction patterns in the context of a brain-specific network.

Figure 2. Differential expression of genes between carriers and noncarriers of the 16p12.1 deletion. (A) Volcano plot showing downstream (non-16p12.1 deletion) genes differentially expressed (FDR<0.05) between carriers (n=19) and noncarriers (n=13) of the deletion. Red circles indicate genes preferentially expressed in GTEx cerebral cortex tissues. **(B)** Scatter plot showing all genes differentially expressed between carriers and noncarriers of the deletion by chromosome, excluding genes on sex chromosomes. Genes are colored by FDR of differential expression. Labeled genes indicate candidate autism genes with differential expression. **(C)** Expression changes and chromatin connectivity of genes within the 16p12.1 region. The top plot shows pairwise chromatin interactions within the 3.5 Mbp 16p12.1 region, with red lines representing stronger Hi-C intensity, while the bottom plot shows $\log_2$-fold change of expression in deletion carriers of genes within and adjacent to the 16p12.1 deletion. The Hi-C data is from previously reported Hi-C experiments of LCL samples[72], and the heatmap was generated using the 3D Genome Browser[95]. **(D)** Line plot shows enrichment ($\log_{10}$ FDR) of differentially expressed genes in deletion carriers for genes preferentially expressed in six select BrainSpan tissues across 11 developmental timepoints.

Figure 3. Differential expression of genes between offspring and carrier and noncarrier parents. (A) Boxplot shows the proportion of differentially expressed genes in carrier offspring of 13 trios (**Table S1**) that either occurred *de novo* or were inherited from their carrier or noncarrier parents (*p<0.05, two-tailed paired Mann-Whitney test). Boxplot indicates median (center line), 25th and 75th percentiles (bounds of box), and minimum and maximum (whiskers). **(B)** Table shows observed clinical features in eight carrier children with overt developmental phenotypes, as well as enrichments (FDR<0.05) of differentially expressed genes in each carrier child for biological functions related to each clinical feature. Cells are colored according to the inheritance pattern of differentially expressed genes for each enriched biological process.

Figure 4. Enrichment of "second-hit" variants near genes with outlier expression. (A) Forest plot shows enrichment (*p<0.05, Fisher's exact test) of genes with outlier expression in all individuals in the cohort (n=32) for rare proximal coding and non-coding variants, including single-nucleotide variants (SNVs) and insertions/deletions (indels) with CADD scores >10[4], structural variants (SVs), and short tandem repeats (STRs). **(B)** Forest plot shows classes of "second-hit" variants with significant enrichment (*p<0.05, Fisher's exact test) towards genes with outlier expression in carrier children (n=10), carrier parents (n=6), or noncarrier parents (n=6). **(C)** Forest plot shows classes of "second-hit" variants with significant enrichment (*p<0.05, Fisher's exact test) towards genes with outlier expression in carrier children (n=9) that are inherited from either carrier or noncarrier parents. All forest plots show log-odds ratios (dots) and 95% confidence intervals (whiskers). **(D)** Scatter plot shows expression values (transcripts per million, or TPM) for *EIF2AK1* in LCL replicates for all individuals in our cohort. Samples in blue have outlier expression of *EIF2AK1* (z-score <-2) and carry a deleterious "second-hit" variant in the 5' UTR of the gene. **(E)** Scatter plots show expression values (TPM) of genes with synergistic effects due to the 16p12.1 deletion and inherited "second-hit" variants. Blue circles indicate expression values for samples from carrier children and family members with the rare "second-hit" variants, orange circles indicate expression values for samples from family members without the "second-hit" variant, and green circles indicate expression values of samples from other deletion carriers and noncarriers in the cohort. Black lines denote mean gene expression for LCL replicates of each individual.

Figure 5. Alternative splicing, allele specific expression, eQTL, and network analysis. (A) Forest plot shows classes of rare variants with significant enrichment (*p<0.05, Fisher's exact test) towards genes with alternative splicing in carrier children (n=9) that are inherited from either carrier or noncarrier parents. Forest plot shows log-odds ratios (dots) and 95% confidence intervals (whiskers). **(B)** Scatter plot shows isoform usage percentage for the isoform *TADA2A-003* in LCL replicates for individuals in family GL_007. Samples in blue carry a "second-hit" splice-site variant in *TADA2A* and exhibit a higher frequency of the alternative isoform. **(C)** Scatter plot shows allele frequencies for the autism-associated gene *CARD11* in carrier child P2C_07, noncarrier parent F2NC_07, and carrier parent M2C_07 in GL_007. Blue circles indicate allele frequency for haplotypes carrying a "second-hit" coding variant disrupting *CARD11*. **(D)** Scatter plot shows z-scores for expression values of *ARHGAP39* for all individuals with available genotypes for the gene. Two individuals who carry the minor allele for the *ARHGAP39* eQTL (blue dots) have higher expression of the gene than the rest of the cohort (orange dots). **(E)** Plot shows correlations among the numbers of gene expression alterations in carrier offspring for the 13 trios assessed in our study. The color and size of the circles are proportional to the correlation coefficients between classes of gene expression changes, where blue indicates a positive correlation and red indicates a negative correlation. Asterisks denote significant correlations (p<0.05, Pearson correlation). **(F)** Boxplot shows the average shortest distances for carrier offspring in the 13 trios between pairs of genes with "second-hit" coding variants and genes with identified expression changes in a brain-specific interaction network. Genes with expression changes were more strongly connected to genes with "second-hit" variants in the brain-specific network than the average distances for genes within 100 permuted brain-specific networks per sample (p=4.88×10^{-4}, two-tailed paired Mann-Whitney test). Boxplot indicates median (center line), 25th and 75th percentiles (bounds of box), and minimum and maximum (whiskers).

Figure 6. Genetic and transcriptomic mechanisms for phenotypic variability in 16p12.1 deletion families. Affected children inherit the 16p12.1 deletion (red) and a smaller number of rare "second-hit" variants (green) from a carrier parent, and a larger number of "second-hit" variants from the noncarrier parent (blue). Altered expression of genes due to these "second-hit" variants affects nearby downstream connected genes in an interaction network (grey), causing

1127 additional transcriptomic perturbation. Because of this, carrier children have numerous gene

1128 expression changes compared with their carrier parents, including genes showing synergistic

1129 effects of the deletion and "second-hit" variants (orange), potentially accounting for more severe

1130 developmental phenotypes observed in the children.

Figure 1

Figure 2

A Genes differentially expressed by the 16p12.1 deletion

B Genes differentially expressed in deletion carriers by chromosome

C Gene expression changes at chromosome 16p12.1

D Enrichment of DEGs in the developing brain

Figure 3

A

Family-based diff. expression

p=8.06e-3
p=7.32e-4   p=0.735

Percentage of all genes with differential expression

0.8, 0.6, 0.4, 0.2, 0.0

De novo · Inh. from carrier · Inh. from noncarrier

B

Biological function enrichments for DEGs of carrier childen

Carrier child	P1C_01							P1C_04	P1C_05		P2C_05	P1C_07		P2C_07			P3C_07		P2C_52
Clinical phenotype	Craniofacial features	Micro/macrocephaly	Neuropsych. features	Hypotonia	Brain abnormalities	Kidney/urinary defects	Musculoskeletal defects	Neuropsych. features	Neuropsych. features	Epilepsy	Neuropsych. features	Neuropsych. features	Epilepsy	Neuropsych. features	Hypotonia	Musculoskeletal features	Neuropsych. features	Musculoskeletal features	Neuropsych. features
Nervous system dev.		red	red		red						blue	red	red	red			red		red
Neurotransmitter metab.			orange									green	green	green					blue
Neuron projection					green							blue	blue	blue					blue
Synapse organization											blue						red		
Synaptic transmission			green							orange		blue	blue				green		blue
Axon formation												orange	orange	blue					
Axon outgrowth			green		green				blue	blue									
Skeletal development	blue						blue									red		green	
Kidney development						blue													
Muscle development				blue	blue										red	red			

De novo [blue]   Inh. carrier [green]   Inh. noncarrier [orange]   Inh. and *de novo* [red]

Figure 4

A Effects of "second-hits" towards outlier expression

B Effects of "second-hits" by sample type

C Effects of "second-hits" by inheritance

D Example of outlier expression

E Synergistic effects of 16p12.1 deletion and "second-hit" variants towards expression

Figure 5

Figure 6

Carrier parent

"Second-hit" variants

16p12.1 del

"Second-hit" variants

Noncarrier parent

"Second-hit" variants

"Second-hit" variants

Carrier child

"Second-hit" variants

16p12.1 del

"Second-hit" variants