Daily torpor reduces the energetic consequences of habitat

selection for a widespread bat

Jesse M. Alston[1,2,3,5], Michael E. Dillon[1], Douglas A. Keinath[4], Ian M.

Abernethy[3], and Jacob R. Goheen[1]

[1] Program in Ecology, Department of Zoology and Physiology, University of

Wyoming, Laramie, Wyoming 82071 USA

[2] Wyoming Natural Diversity Database, University of Wyoming, Laramie, Wyoming

82071 USA

[3] Center for Advanced Systems Understanding (CASUS), Görlitz, Sachsen 02826

DEU

[4] Wyoming Ecological Services Field Office, United States Fish and Wildlife Service,

Cheyenne, Wyoming 82009 USA

[5] Corresponding author: jalston@uwyo.edu

Abstract

Homeothermy requires increased metabolic rates as temperatures decline below the thermoneutral zone, so homeotherms typically select microhabitats within or near their thermoneutral zones during periods of inactivity. However, many mammals and birds are heterotherms that relax internal controls on body temperature when maintaining a high, stable body temperature is energetically costly. Such heterotherms should be less tied to microhabitats near their thermoneutral zones, and because heterotherms spend more time in torpor and expend less energy at colder temperatures, heterotherms may even select microhabitats in which temperatures are well below their thermoneutral zones. We studied how temperature and daily torpor influence selection of diurnal roosts by a heterothermic bat (*Myotis thysanodes*). We (1) quantified the relationship between ambient temperature and daily duration of torpor, (2) simulated daily energy expenditure over a range of microhabitat (roost) temperatures, and (3) quantified the influence of roost temperature on roost selection. While warm roosts substantially reduced energy expenditure of simulated homeothermic bats, heterothermic bats modulated their use of torpor to maintain a constant level of energy expenditure over the course of a day. Daily torpor expanded the range of energetically economical microhabitats, such that roost selection was independent of roost temperature. Our work adds to a growing literature documenting functions of torpor beyond its historical conceptualization as a last-resort measure to save energy during extended or acute energetic stress.

Introduction

The thermal environments in which organisms live strongly influence metabolic rates (Huey and Stevenson 1979, Brown et al. 2004, Pörtner and Knust 2007). Among homeotherms—which regulate body temperature internally within a narrow range to optimize physiological processes—metabolic heat production is tightly regulated in response to variation in temperature in the surrounding environment (i.e., ambient temperature; Lowell and Spiegelman 2000). Controlling body temperature thus requires increased energy expenditure by homeotherms when ambient temperatures depart from the thermoneutral zone (i.e., the range of ambient temperatures in which homeotherms can regulate body temperature with minimal metabolic effort; McNab 2002). Because survival and reproduction require that energy intake equal or exceed energy expenditure, operating in ambient temperatures outside the thermoneutral zone can reduce fitness over time (Angilletta et al. 2010, Boyles et al. 2011).

Although the influence of ambient temperature on metabolism in homeotherms is understood relatively well, many animals are heterotherms that can temporarily or partially allow body temperature to track ambient temperature (Withers et al. 2016). Heterothermy is common among mammals and birds (Geiser, 2004; Geiser and Ruf, 1995; McKechnie and Mzilikazi, 2011; Ruf and Geiser, 2015) and can reduce energy expenditure during both hot and cold periods (Stawski and Geiser 2012, Boyles et al. 2016, Nowack et al. 2017, Reher and Dausmann 2021). As ambient temperatures depart the thermoneutral zone, heterotherms can relax internal controls on metabolism; this physiological response allows body temperature to track ambient temperature and reduce or altogether eliminate the energetic costs of maintaining stable body temperatures outside the thermoneutral zone (Levesque et al. 2016). Heterotherms often

achieve this by entering daily torpor, a short-term hypometabolic state of inactivity (Ruf

and Geiser 2015).

Heterotherms use daily torpor more as ambient temperatures decline below the

thermoneutral zone (Chruszcz and Barclay, 2002; Geiser and Broome, 1993;

Rambaldini and Brigham, 2008; Solick and Barclay, 2006), but it is unclear how this

tendency translates to differences in energy expenditure across differences in

temperature. For a given period of time, total energy expenditure for heterotherms

depends on (1) the duration and frequency of bouts of torpor, (2) ambient temperatures,

and (3) the difference in metabolic rates between torpor and homeothermy at a given

ambient temperature. Energy expenditure might increase as ambient temperatures fall

below the thermoneutral zone: even though heterotherms save energy by using torpor,

declines in energy expenditure from using torpor more when it is cold do not fully

compensate for the increased energetic costs of maintaining homeothermy in colder

ambient temperatures (Fig. 1B). In this scenario, periodic bouts of torpor dampen but

do not completely offset increases in energy expenditure during periods of

homeothermy at cold ambient temperatures. Alternatively, it is possible that energy

expenditure by heterotherms is stable through a wide range of ambient temperatures

because energy savings from using progressively more torpor at progressively colder

ambient temperatures closely matches increases in energy expenditure from

maintaining homeothermy at colder ambient temperatures (Fig. 1C). Finally, as ambient

temperatures decline, the energetic savings from torpor could more than offset the

increased energy expenditure necessary to maintain homeothermy (Fig. 1D).

Such relationships between ambient temperature and energy expenditure have

cascading repercussions for other aspects of an animal's life. For example, ambient

temperature often influences habitat selection by animals seeking to minimize energy

expenditure (e.g., Huey 1991, Freitas et al. 2016, Sarmento et al. 2019, Alston et al.

2020). Homeotherms have relatively fixed relationships between ambient temperature

and metabolic rate, and thus often consistently select habitats to maintain optimal body

temperatures with little metabolic effort (e.g., Poole et al. 2016, Courbin et al. 2017,

Sarmento et al. 2019). In contrast, looser relationships between ambient temperature

and metabolic rate for heterotherms may allow heterotherms to select habitats with less

regard to ambient temperature, or even to prefer habitats that might be colder than

optimal for homeotherms. For example, heterothermic Australian owlet-nightjars

(*Aegotheles cristatus*) preferentially roost in colder, less thermally stable tree cavities,

whereas homeothermic cavity-nesting birds typically select warmer, more thermally

stable tree cavities (Doucette et al. 2011). Empirical data on habitat selection by

heterotherms is rare, however, particularly for free-ranging animals.

Uncertainty surrounding the form and strength of relationships between ambient

temperature and energy expenditure limit our understanding of temperature-driven

habitat selection by heterotherms. For an animal attempting to minimize energy

expenditure during periods of inactivity, each of the hypothetical relationships between

energy expenditure and ambient temperature in Fig. 1 would result in a different pattern

of habitat selection. A heterotherm exhibiting the relationship shown by the red (B) line

in Fig. 1 should select warm microhabitats to save energy, similar to homeotherms. A

heterotherm exhibiting the relationship shown by the grey (C) line in Fig. 1 should not

select microhabitats based on their thermal characteristics. This pattern of habitat

selection would also diverge from the pattern followed by homeotherms. A heterotherm

exhibiting the relationship shown by the blue (D) line in Fig. 1 should select cool

microhabitats to save energy, opposite of the pattern followed by homeotherms.

Empirical tests of the influence of ambient temperature on energy expenditure are thus

needed to understand how ambient temperature drives habitat selection for heterotherms.

We sought to understand how ambient temperature influences energy expenditure, and how energy expenditure in turn influences habitat selection, in a bat that is widely distributed throughout western North America (fringed myotis, *Myotis thysanodes*). Like other bats inhabiting temperate latitudes, fringed myotis are heterotherms that are believed to select diurnal roosts to minimize energy expenditure during diurnal periods of inactivity (Sedgeley 2001, Willis and Brigham 2005, Ruczyński 2006). At temperate latitudes, temperature within roosts can vary substantially throughout the day and year, and ambient temperature influences the amount of time bats spend in torpor each day. Like other heterotherms, bats spend more time in torpor when it is cold than when it is hot (Chruszcz and Barclay 2002, Solick and Barclay 2006, Rambaldini and Brigham 2008). We hypothesized that differences in energy expenditure at roosts of varying temperatures drive patterns of roost selection (i.e., bats select roosts that minimize energy expenditure). Specifically, we weighed evidence for four competing sets of predictions (Fig. 2):

Prediction Set 1: Bats select warm roosts regardless of ambient temperature. In this scenario, energy expenditure during the day should be higher in cool roosts than in warm roosts (Fig. 2.1A) because the energetic benefits from being warmer when bats are maintaining homeothermy outweigh the energetic costs of spending less time in torpor. If this is the case, bats should select roosts that are warmer compared to available structures on the landscape (Fig. 2.1B); this pattern of selection should be consistent regardless of ambient temperature during the day (Fig. 2.1C).

Prediction Set 2: Bats select cool roosts regardless of ambient temperature. In this scenario, energy expenditure during the day should be higher in warm roosts than

in cool roosts (Fig. 2.2A) because the energetic benefits from spending more time in

torpor outweigh the energetic costs of being colder when bats are maintaining

homeothermy. If this is the case, bats should select roosts that are cooler compared to

available structures on the landscape (Fig. 2.2B); this pattern of selection should be

consistent regardless of ambient temperature during the day (Fig. 2.2C).

Prediction Set 3: Bats do not alter roost selection as ambient temperatures

change. In this scenario, energy expenditure during the day is roughly equal across

roosts of all temperatures (Fig. 2.3A). This could occur if bats modulate use of torpor

such that roost temperatures over the course of a day have little influence on overall

energy expenditure. In this case, bats should select roosts that are similar in temperature

to available structures on the landscape (Fig. 2.3B), and this pattern of selection should

be consistent regardless of ambient temperature during the day (Fig. 2.3C).

Prediction Set 4: Bats select cool roosts on cool days and warm roosts on warm

days (shifting roost selection). In this scenario, energy expenditure is lower in cool

roosts than in warm roosts on cool days, lower in warm roosts than in cool roosts on

warm days, and consistently higher in roosts at intermediate ambient temperatures (Fig.

2.4A). This may arise because of threshold effects from a non-linear relationship

between ambient temperature and torpor use. Namely, a threshold may exist above

which homeothermy requires relatively little energy even as bats spend little time in

torpor, but below which bats save a substantial amount of energy by using torpor. Near

the threshold, however, bats may use relatively little torpor even as maintaining

homeothermy is relatively energetically costly. In this case, bats should select roosts

that are roughly the same temperature on average as available structures on the

landscape (though the distribution may be bimodal; Fig. 2.4B), and temperatures in

roosts should be positively correlated with ambient temperature (Fig. 2.4C).

Methods

Study Area and Species. We conducted our study during the summers of 2017 and 2018 on Jewel Cave National Monument (43˚ 45' N, 103˚ 45' W) and surrounding areas of Black Hills National Forest in South Dakota, USA. Our study area is described in Alston et al. (2019). Mean monthly summer high temperatures range between $22 - 27$˚C and mean monthly summer precipitation ranges between $60 - 80$ mm (Western Regional Climate Center 2018). Open ponderosa pine (*Pinus ponderosa*) forests dominate, with Rocky Mountain juniper (*Juniperus scopulorum*) and quaking aspen (*Populus tremuloides*) occurring locally. Forests are actively managed to prevent wildfire, and those managed by the US Forest Service and private landowners also undergo intensive logging. Forests form a mosaic with northern mixed-grass prairie where a large stand-replacing fire occurred in in 2000. A large network of caves lie underground, and the landscape exhibits substantial topographic relief in the form of intersecting canyon systems and rock outcrops.

Fringed myotis roost in caves, mines, rock crevices, tree cavities, and under the sloughing bark of dead trees, and forage in forest canopy and riparian areas (O'Farrell and Studier 1980). We chose to focus on males because sex ratios of bats in the Black Hills are heavily (>90%) male-biased (a common pattern in high-elevation areas; Barclay, 1991; Cryan et al. 2000; Senior et al. 2005), because male *M. thysanodes* usually roost solitarily (O'Farrell and Studier 1980), and because male bats maintain consistent patterns of torpor use throughout the reproductive season (unlike females, which alter patterns of torpor use at different stages of reproduction; Chruszcz and Barclay, 2002; Dzal and Brigham, 2013; Johnson and Lacki, 2014).

Capture and VHF Telemetry. We used mist nets to capture bats over permanent and semi-permanent water sources (e.g., springs, stock tanks, and stock ponds). From

June through August of 2017 and 2018, we netted 20 and 49 nights, respectively, at 15

water sources. We opened mist nets at civil sunset and closed them after five hours or

during inclement weather.

We affixed temperature-sensitive VHF transmitters (LB-2XT model .28/.33 g –

Holohil Systems Ltd., Carp, ON, Canada) between the scapulae of adult male fringed

myotis with latex surgical adhesive (Osto-Bond, Montreal Ostomy, Montreal, QC,

Canada). The transmitters measure and transmit data on skin temperature—an accurate

proxy for body temperature—of bats, enabling researchers to delineate bouts of torpor

(Barclay et al. 1996, Chruszcz and Barclay 2002, Stawski and Geiser 2010). All

transmitters weighed <5% of the mass of the bat (Aldridge and Brigham 1988). We

tracked bats to roosts each day transmitters were active, and installed VHF data loggers

(SRX800-D1 – Lotek Wireless Inc., Newmarket, ON, Canada) that collected and

recorded data transmitted by the VHF transmitters. All protocols were approved by the

University of Wyoming and National Park Service Animal Care and Use Committees

and met guidelines approved by the American Society of Mammalogists for research

on wild mammals (Sikes and the Animal Care and Use Committee of the American

Society of Mammalogists 2016).

Energetic Modelling. To quantify torpor use, we delineated bouts of torpor

from data logger readings that captured full days (i.e., from roost entry in the morning

to roost exit in the evening) of skin temperature data from individual bats. This was a

fraction of total days in which we located roosts, because bats typically were not located

until after they entered roosts. We defined torpor as beginning when skin temperature

dropped below the lowest skin temperature of bats maintaining homeothermy during a

day and ending when skin temperature began a steep rise that led to bats re-entering

homeothermy or leaving a roost (as recommended by Barclay et al. 2001; Appendix

S1: Fig. S1). Because fat reserves and body mass can substantially alter the amount of time spent in torpor (Wojciechowski et al. 2007, Stawski and Geiser 2010, Vuarin et al. 2013), we also controlled for the body mass of each individual at time of capture on torpor duration. We then used the modelling software 'Stan' (Carpenter et al. 2017) via the R package 'brms' (v2.13.0; Bürkner 2017) to build a linear Bayesian hierarchical model to quantify the influence of ambient temperature and body mass on torpor duration while accounting for non-independence among data points collected from the same individual. The model included 3 chains run for 13,000 iterations (1,000 iterations of warm-up and 12,000 iterations of sampling). We assessed chain convergence using the Gelman-Rubin diagnostic ($\hat{R}$) and precision of parameter estimation using effective sample size. $\hat{R} < 1.01$ and effective sample sizes $> 10,000$ represent acceptable convergence and parameter precision (Gelman et al. 2013, Kruschke 2015). We used leave-one-out cross validation to check model fit using the R packages 'loo' (v2.2.0; Vehtari et al. 2017) and 'bayesplot' (v1.7.2; Gabry et al. 2019) to visually assess the cross-validated probability integral transform.

To quantify energy expenditure in bats, we combined published estimates of metabolic rates of fringed myotis as a function of temperature (Studier and O'Farrell 1976) and the linear model of the influence of ambient temperature on torpor use to simulate the influence of roost temperature on energy expenditure. Specifically, we simulated minute-by-minute energy expenditure by bats in each used roost between 0445 hrs and 2100 hrs (typical entry and exit times for bats in our study) on each day over the duration of our study period. We modeled torpor use as a function of decision rules that reflect torpor use observed over the course of our study (raw data presented in Appendix S1: Table S1). Specifically, we assumed that bats entered torpor immediately upon entering roosts, exited torpor after an interval determined by roost

temperature, and remained in homeothermy for the rest of the time spent in the roost

except for a shorter bout of torpor in the evening. We further assumed that bats would

use 86.9% of the duration of daily torpor in the morning and 13.1% in the afternoon

unless the afternoon bout of torpor would be less than 30 minutes in duration, in which

case 100% of the day's torpor would occur in the morning period. We also assumed

that the mean duration of torpor that we observed would be used in the baseline

"average" roost, with the duration of torpor in warmer and cooler roosts determined by

the slope of the modeled relationship between ambient temperature and torpor use

described in the above paragraph. To account for uncertainty in our estimate of the slope

of the relationship between ambient temperature and daily torpor use, for each roost on

each day we randomly drew a different slope estimate for this relationship from the

posterior distribution of slope estimates from the model described in the prior

paragraph.

Roost Characterization. To characterize rock roost structures, we collected data

for 31 roosts and 62 randomly sampled available (i.e., unused by bats in our study)

roosts. Hereafter, we distinguish between 'used roosts' and available but unused

'available roosts'; we use the term 'roost structure' when we refer to both used and

available roosts simultaneously. We identified available rock roosts in two ways: at

each used roost, we 1) located the nearest rock crevice large enough to hold a bat, and

2) generated a paired point in a random cardinal direction a random distance between

100 – 300 m away, then located the nearest rock crevice large enough to hold a bat.

To characterize tree roost structures, we collected data for 9 used roosts and 36

randomly sampled available roosts. We identified available tree roosts in two ways: at

each used roost, we 1) located the nearest snag and selected the nearest cavity large

enough to hold a bat, and 2) generated a paired point in a randomly determined distance

between 100 – 300 m away, in a randomly-determined (cardinal) direction, then located

the nearest tree cavity large enough to hold a bat. For each available point, we placed

data loggers in two locations: one in a cavity in the trunk and one underneath sloughing

bark. We defined available roost trees as any dead tree with a visible defect (e.g.,

sloughing bark or cavities) sufficiently large to hold a bat. This description fit every

tree in which we found a bat roosting.

In Summer 2018, we monitored temperatures within both used and available

roosts using data loggers (Model MX2201; Onset Computer Corporation, Bourne, MA,

USA). The first data loggers were deployed on 17 July 2018, and the last data logger

was removed on 8 October 2018. This period of time includes the full range of daily

high temperatures occurring during the active season for bats at our study site. During

data logger deployment and opportunistically thereafter, we checked roost structures

for the presence of bats. We sometimes found bats in used roosts, but we never found

bats in available roosts. When we found bats in used roosts, we waited to deploy data

loggers until there was no bat within the roost.

To quantify the thermal characteristics of each roost structure, we calculated the

mean temperature within each roost structure for periods between 0445 and 2100 hrs,

which corresponds with the period in which a bat is likely to be within a roost

(Appendix S1: Table S1). To control for potential confounding variables, we also

calculated the timing of the peak temperature in all roost structures (because if two roost

structures have the same mean temperature but peak in temperature at different times,

the roost structure with the later peak will have cooler temperatures in the morning

when bats use torpor most), and the standard deviation of temperature during the day

(because stability in roost temperature can be an important factor in roost selection;

Sedgeley, 2001). To quantify the timing of the daily temperature peak, we located the

peak temperature in each roost structure for each day and calculated the mean time of day at which this occurred over our study period. To quantify thermal stability in roost structures, we calculated the standard deviation of temperatures between 0445 and 2100 hrs in each roost structure for each day and calculated the mean daily standard deviation over our study period. To ensure consistency, we only calculated these values for the period between July 28 and September 31 (a period in which all data loggers were actively logging temperatures, and in which average daily high temperatures correspond with the range a bat might be exposed to during the active season in our study area).

We used the R statistical software environment (R Core Team 2020) to quantify differences between used and available roosts. To determine whether bats select cooler roosts than those available, we used the modelling software 'Stan' (Carpenter et al. 2017) via the R package 'brms' (v2.13.0; Bürkner 2017) to build a binomial-family Bayesian model to quantify the influence of mean temperature within roost structures, the timing of daily peaks in temperature within roost structures, and the standard deviation of temperatures within roost structures on roost selection. The model included 3 chains run for 13,000 iterations (1,000 iterations of warm-up and 12,000 iterations of sampling). We assessed chain convergence using $\hat{R}$ and precision of parameter estimation using effective sample size. We checked predictive performance with receiver operating curve analysis using the R package 'pROC' (v1.16.2; Robin et al. 2011) and used the R package 'bayesplot' (v1.7.2; Gabry et al. 2019) to visually assess binned residual plots.

Results

We tracked 46 bats to 107 roosts (93 in rock crevices and 14 in trees) and collected 27

full days of skin temperature data from 7 bats. Data from 16 data loggers within roost

structures (3 used rock, 12 available rock, 1 available tree) could not be collected

because they were not relocated or were dislodged from roost structures. We thus

excluded these data from analyses, leaving a total of 122 (78 rock, 44 tree) data loggers

that collected data on temperatures within roost structures.

Use of torpor stabilized daily energy expenditure across the range of roost

temperatures observed during our telemetry study. In our model of the effect of ambient

temperature on daily torpor duration, 95% credible intervals for the effect of mean

ambient temperature over the course of the day on daily torpor duration did not cross 0

(parameter estimate: -37.4 min; 95% credible intervals: -64.0 – -12.6 min), indicating

that bats spent ca. 37 minutes less in torpor per day for each additional 1˚C in daily

mean ambient temperature between 0445 hrs and 2100 hrs (Fig. S2). Assessment of the

cross-validated probability integral transform indicated that model fit was adequate.

When incorporated into our simulation of bat energy expenditure over the course of a

typical day, this estimate of the relationship between ambient temperature and torpor

use led to similar estimates of energy expenditure across temperatures within used

roosts (Fig. 3; blue points). Daily energy expenditure was roughly equivalent in all

roosts. Our estimates for energy expenditure using observed bat behavior were always

substantially lower and less variable than our estimates for energy expenditure if bats

had remained in homeothermy all day (Fig. 3; red points). Bats that remain in

homeothermy would expend substantially more energy in cool roosts than warm roosts.

Overall, temperatures in both rock and tree roost structures were similar, though

roost structures in trees were slightly cooler and less stable than roost structures in

rocks. We therefore pooled rock and tree roost structures in roost selection analyses, but we report descriptive statistics for each type of roost structure in Appendix S1.

Despite substantial variation in temperatures among roost structures, we found little evidence that the thermal characteristics of used roosts differed from those of available roosts (Fig. 4). In our model of roost selection, 95% credible intervals for the effect of mean ambient temperature over the course of the day on roost selection did not cross 0 (parameter estimate: 0.30; 95% credible intervals: 0.04 – 0.58), indicating that bats were more likely to roost in warm roost structures than cool ones. However, predictive performance was poor (AUC: 0.650), and overall, used roosts (20.1˚C) had similar mean temperatures as available roosts (19.4˚C; Fig. 4A). Bats also did not differentiate between roost structures with temperatures peaking late in the day versus roost structures with temperatures peaking early in the day (Fig 4B). In our model of roost selection, 95% credible intervals for the effect of the timing of daily peaks in temperature on roost selection crossed 0 (parameter estimate: -0.10; 95% credible intervals: -0.34 – 0.14). Overall, used roosts (1408 hrs) peaked in temperature at similar times as available roosts (1434 hrs). Bats also did not differentiate between roosts with stable temperatures and those with more variable temperatures (Fig. 4C). In our model of roost selection, 95% credible intervals for the effect of standard deviation in roost temperature over the course of the day on roost selection crossed 0 (parameter estimate: -0.20; 95% credible intervals: -0.47 – 0.06) Overall, there was no difference in the standard deviation of temperatures of used roosts (7.0˚C) and available roosts (7.0˚C). Finally, there was also no relationship between ambient temperature on a given day and mean temperatures within roosts used on that day ($R^2 = 0.03$; $p = 0.132$; Fig. 5).

Discussion

The thermal environments in which animals operate strongly influence physiological

processes, and can thereby pose substantial challenges in variable environments. How

animals overcome these challenges is a central question in animal ecology. Attempts to

address this question have focused largely on poikilotherms and homeotherms. Because

heterotherms are neither as strongly tied to narrow ranges of body temperature as

homeotherms nor as subject to ambient temperatures as poikilotherms, the relationships

between temperature and habitat selection for heterotherms should differ fundamentally

from those of either homeotherms or poikilotherms. Specifically, whereas

homeotherms select microhabitats near the thermoneutral zone during periods of

inactivity, heterotherms should have less incentive to do so.

We sought to better understand how variation in ambient temperature influences

use of daily torpor and habitat selection for heterotherms, using a species of bat as a

model system. Simulations of energy expenditure at varying roost temperatures

revealed that bats can modulate use of torpor to maintain constant energy expenditure

over the course of a day over a wide range of temperatures within roosts. As a result,

roost selection was not driven by temperatures within roosts. Our results provide

evidence for Prediction Set 3 (no selection) in our introduction (Fig. 2).

The energetic savings associated with torpor—particularly at cooler

temperatures—likely result in habitat selection that differs substantially from habitat

selection by homeotherms. For example, we showed that use of daily torpor can reduce

the energetic costs of inhabiting roosts that are colder than optimal for homeotherms. If

bats were strict homeotherms, the energetic costs of inhabiting cool roosts would have

been substantially higher (Fig. 3), which would likely result in bats selecting warm

roosts. In contrast, heterothermic bats face little pressure to select warm habitats, even

on relatively cool days. Daily torpor does not simply loosen the thermal constraints

facing homeotherms at temperatures below the thermoneutral zone—it can entirely

mitigate them. Additional studies of the relationships between temperature, torpor use,

and habitat selection would be valuable for establishing the generality of this finding

for other heterothermic species.

Individual traits (e.g., sex, age, and reproductive condition) can alter the

energetic costs and benefits of using torpor for heterotherms, thereby driving

divergence from the pattern demonstrated in this study. For example, roost selection by

bats varies by sex, age, and reproductive condition (Elmore et al. 2004, Hein et al.

2008). While male bats in our study did not select roosts with specific thermal

characteristics, female bats seem to use less torpor and prefer warmer roosts than males

while pregnant or raising young, and females typically aggregate in social maternity

colonies rather than roosting solitarily (Hamilton and Barclay 1994, Kerth et al. 2001,

Ruczyński 2006). Compared to males, then, roost selection by females will likely be

governed more strongly by roost temperature (though social thermoregulation via

huddling can influence temperatures within roosts more than a roost's physical and

environmental characteristics; Pretzlaff et al. 2010; Willis and Brigham, 2007). Further

research on the roles of sex, age, and reproductive condition on torpor use in

heterotherms (and thus habitat selection by heterotherms) is likely to reveal important

context for our findings.

Climate warming increases energy expenditure for many animals, including

both poikilotherms (Pörtner and Knust 2007, Dillon et al. 2010) and homeotherms

(Humphries et al. 2002, Şekercioğlu et al. 2012). However, the degree to which climate

warming will impact heterotherms is poorly understood, largely due to a lack of data

on relationships between ambient temperature, torpor use, and thermolability that is

needed to accurately model the influence of ambient temperature on heterotherm

metabolism (Levesque et al. 2016). Our results indicate that temperature-dependent use of torpor may stabilize energy expenditure, and thus buffer against the energetic costs associated with variable ambient temperatures. However, most of the energetic savings from heterothermy arise during periods of cold. Increased temperatures due to climate change may thus reduce the relative energetic benefits of heterothermy compared to homeothermy, as homeotherms experience fewer and milder periods of cold.

In conclusion, we showed that a heterothermic bat selected neither warm nor cool roosts, because bats can modulate torpor use to stabilize energy expenditure over the course of a day. Unlike homeotherms, bats face little pressure to select warm habitats to avoid heat loss during periods of inactivity—when maintaining a high, stable body temperature becomes energetically costly, bats can calibrate the duration of torpor such that energy expenditure stays constant through a wide range of ambient temperature. Although such fine-tuning of torpor use to stabilize daily energy expenditure is intuitive, it has not been demonstrated in previous studies to the best of our knowledge.

Acknowledgements

Many thanks to L. Boodoo, C. McFarland, E. Greene, B. Tabor, and B. Phillips for help with fieldwork; L. Shoemaker, D. Laughlin, J. Rick, and D. Goodhouse for helpful comments on pre-submission versions of this manuscript; and P. Ortegon, D. Licht, M. Wiles, D. Austin, B. Phillips, E. Thomas, and A. Stover for their logistical support of this project. Research funding was provided by the National Park Service, the Department of Zoology and Physiology at the University of Wyoming, the Berry Ecology Center, the American Society of Mammalogists, Prairie Biotic Research, Inc., and the Wyoming Chapter of The Wildlife Society. The findings and conclusions

in this article are those of the authors and do not necessarily represent the views of the

U.S. Fish and Wildlife Service or the National Park Service. We conducted field

research on the traditional lands of the Lakȟóta, Sahnish, Tsitsistas, Hinono'eino,

K'oigu, and Ná'ishą peoples. The Lakȟóta people know this land as Ȟe Sápa and

Pahá Sápa, which was taken by the United States in the Agreement of 1877 in

violation of the 1868 Fort Laramie Treaty.

Data Availability

Data and code used in analyses for this paper will be archived on *Zenodo* upon

acceptance of this manuscript and are available upon request by editors or reviewers.

Author Contributions

JA, JG, and MD conceived and designed the study; JA, DK, and IA obtained funding

for the study; JA collected and analyzed the data and led writing of the manuscript.

All authors contributed to manuscript drafts and gave final approval for publication.

References

Aldridge, H.D.J.N., and R.M. Brigham. 1988. Load carrying and maneuverability in

an insectivorous bat: a test of the 5% "rule" of radio-telemetry. Journal of

Mammalogy 69:379–382.

Alston, J.M., I.M. Abernethy, D.A. Keinath, and J.R. Goheen. 2019. Roost selection

by male northern long-eared bats (*Myotis septentrionalis*) in a managed fire-

adapted forest. Forest Ecology and Management 446:251–256.

Alston, J.M., M.J. Joyce, J.A. Merkle, and R.A. Moen. 2020. Temperature shapes

469   movement and habitat selection by a heat-sensitive ungulate. Landscape

470   Ecology:790048.

471   Angilletta, M.J., B.S. Cooper, M.S. Schuler, and J.G. Boyles. 2010. The evolution of

472   thermal physiology in endotherms. Frontiers in Bioscience E2:861–881.

473   Barclay, R.M., C.L. Lausen, and L. Hollis. 2001. What's hot and what's not: defining

474   torpor in free-ranging birds and mammals. Canadian Journal of Zoology

475   79:1885–1890.

476   Barclay, R.M.R. 1991. Population structure of temperate zone insectivorous bats in

477   relation to foraging behaviour and energy demand. Journal of Animal Ecology

478   60:165–178.

479   Barclay, R.M.R., M.C. Kalcounis, L.H. Crampton, C. Stefan, M.J. Vonhof, L.

480   Wilkinson, and R.M. Brigham. 1996. Can external radiotransmitters be used to

481   assess body temperature and torpor in bats? Journal of Mammalogy 77:1102–

482   1106.

483   Boyles, J.G., L.P. McGuire, E. Boyles, J.P. Reimer, C.A.C. Brooks, R.W. Rutherford,

484   T.A. Rutherford, J.O. Whitaker, and G.F. McCracken. 2016. Physiological and

485   behavioral adaptations in bats living at high latitudes. Physiology & Behavior

486   165:322–327.

487   Boyles, J.G., F. Seebacher, B. Smit, and A.E. McKechnie. 2011. Adaptive

488   thermoregulation in endotherms may alter responses to climate change.

489   Integrative and Comparative Biology 51:676–690.

490   Brown, J.H., J.F. Gillooly, A.P. Allen, V.M. Savage, and G.B. West. 2004. Toward a

491   metabolic theory of ecology. Ecology:1771–1789.

Bürkner, P.-C. 2017. brms: an R package for Bayesian multilevel models using Stan.

Journal of Statistical Software 80:1–28.

Carpenter, B., A. Gelman, M.D. Hoffman, D. Lee, B. Goodrich, M. Betancourt, M.

Brubaker, J. Guo, P. Li, and A. Riddell. 2017. Stan: a probabilistic

programming language. Journal of Statistical Software 76.

Chruszcz, B.J., and R.M.R. Barclay. 2002. Thermoregulatory ecology of a solitary

bat, *Myotis evotis*, roosting in rock crevices. Functional Ecology 16:18–26.

Courbin, N., C. Dussault, A. Veillette, M.-A. Giroux, and S.D. Côté. 2017. Coping

with strong variations in winter severity: plastic habitat selection of deer at

high density. Behavioral Ecology 28:1037–1046.

Cryan, P.M., M.A. Bogan, and J.S. Altenbach. 2000. Effect of elevation on

distribution of female bats in the Black Hills, South Dakota. Journal of

Mammalogy 81:719–725.

Cunningham, S.J., J.L. Gardner, and R.O. Martin. 2021. Opportunity costs and the

response of birds and mammals to climate warming. Frontiers in Ecology and

the Environment, early view.

Dillon, M.E., G. Wang, and R.B. Huey. 2010. Global metabolic impacts of recent

climate warming. Nature 467:704–706.

Doucette, L.I., R.M. Brigham, C.R. Pavey, and F. Geiser. 2011. Roost type influences

torpor use by Australian owlet-nightjars. Naturwissenschaften 98:845.

Dzal, Y.A., and R.M. Brigham. 2013. The tradeoff between torpor use and

reproduction in little brown bats (*Myotis lucifugus*). Journal of Comparative

Physiology B 183:279–288.

Elmore, L.W., D.A. Miller, and F.J. Vilella. 2004. Selection of diurnal roosts by red

bats (*Lasiurus borealis*) in an intensively managed pine forest in Mississippi.

Forest Ecology and Management 199:11–20.

Freitas, C., E.M. Olsen, H. Knutsen, J. Albretsen, and E. Moland. 2016. Temperature-

associated habitat selection in a cold-water marine fish. Journal of Animal

Ecology 85:628–637.

Gabry, J., D. Simpson, A. Vehtari, M. Betancourt, and A. Gelman. 2019.

Visualization in Bayesian workflow. Journal of the Royal Statistical Society:

Series A (Statistics in Society) 182:389–402.

Geiser, F. 2004. Metabolic rate and body temperature reduction during hibernation

and daily torpor. Annual Review of Physiology 66:239–274.

Geiser, F., and L.S. Broome. 1993. The effect of temperature on the pattern of torpor

in a marsupial hibernator. Journal of Comparative Physiology B 163:133–137.

Geiser, F., and T. Ruf. 1995. Hibernation versus daily torpor in mammals and birds:

physiological variables and classification of torpor patterns. Physiological

Zoology 68:935–966.

Gelman, A., J.B. Carlin, H.S. Stern, D. B. Dunson, A. Vehtari, and D.B. Rubin. 2013.

Bayesian Data Analysis. Third. CRC Press, Boca Raton, FL, USA.

Hamilton, I.M., and R.M.R. Barclay. 1994. Patterns of daily torpor and day-roost

selection by male and female big brown bats (*Eptesicus fuscus*). Canadian

Journal of Zoology 72:744–749.

Hein, C.D., S.B. Castleberry, and K.V. Miller. 2008. Sex-specific summer roost-site

selection by Seminole bats in response to landscape-level forest management.

Journal of Mammalogy 89:964–972.

Huey, R.B. 1991. Physiological consequences of habitat selection. American

Naturalist 137:S91–S115.

Huey, R.B., and R.D. Stevenson. 1979. Integrating thermal physiology and ecology of

ectotherms: a discussion of approaches. Integrative and Comparative Biology

19:357–366.

Humphries, M.M., D.W. Thomas, and J.R. Speakman. 2002. Climate-mediated

energetic constraints on the distribution of hibernating mammals. Nature

418:313–316.

Johnson, J.S., and M.J. Lacki. 2014. Effects of reproductive condition, roost

microclimate, and weather patterns on summer torpor use by a vespertilionid

bat. Ecology and Evolution 4:157–166.

Kerth, G., K. Weissmann, and B. König. 2001. Day roost selection in female

Bechstein's bats (*Myotis bechsteinii*): a field experiment to determine the

influence of roost temperature. Oecologia 126:1–9.

Kruschke, J. 2015. Doing Bayesian analysis. Second. Academic Press, Waltham, MA,

USA.

Levesque, D.L., J. Nowack, and C. Stawski. 2016. Modelling mammalian energetics:

the heterothermy problem. Climate Change Responses 3:7.

Lowell, B.B., and B.M. Spiegelman. 2000. Towards a molecular understanding of

adaptive thermogenesis. Nature 404:652–660.

McKechnie, A.E., and N. Mzilikazi. 2011. Heterothermy in Afrotropical mammals

and birds: a review. Integrative and Comparative Biology 51:349–363.

McNab, B.K. 2002. The physiological ecology of vertebrates: a view from energetics.

Cornell University Press, Ithaca, NY, USA.

Nowack, J., C. Stawski, and F. Geiser. 2017. More functions of torpor and their roles

in a changing world. Journal of Comparative Physiology B 187:889–897.

O'Farrell, M. J., and E. H. Studier. 1980. Myotis thysanodes. Mammalian Species:1–

5.

Poole, K.G., R. Serrouya, I.E. Teske, and K. Podrasky. 2016. Rocky Mountain

bighorn sheep (*Ovis canadensis canadensis*) winter habitat selection and

seasonal movements in an area of active coal mining. Canadian Journal of

Zoology 94:733–745.

Pörtner, H.O., and R. Knust. 2007. Climate change affects marine fishes through the

oxygen limitation of thermal tolerance. Science 315:95–97.

Pretzlaff, I., G. Kerth, and K.H. Dausmann. 2010. Communally breeding bats use

physiological and behavioural adjustments to optimise daily energy

expenditure. Naturwissenschaften 97:353–363.

R Core Team. 2020. R: a language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

Rambaldini, D.A., and R.M. Brigham. 2008. Torpor use by free-ranging pallid bats

(*Antrozous pallidus*) at the northern extent of their range. Journal of

Mammalogy 89:933–941.

Reher, S., and K.H. Dausmann. 2021. Tropical bats counter heat by combining torpor

with adaptive hyperthermia. Proceedings of the Royal Society B: Biological

Sciences 288:20202059.

Robin, X., N. Turck, A. Hainard, N. Tiberti, F. Lisacek, J.-C. Sanchez, and M. Müller.

2011. pROC: an open-source package for R and S+ to analyze and compare

ROC curves. BMC Bioinformatics 12:77.

Ruczyński, I. 2006. Influence of temperature on maternity roost selection by noctule

bats (*Nyctalus noctula*) and Leisler's bats (*N. leisleri*) in Białowieża Primeval

Forest, Poland. Canadian Journal of Zoology 84:900–907.

Ruf, T., and F. Geiser. 2015. Daily torpor and hibernation in birds and mammals.

Biological Reviews 90:891–926.

Sarmento, W., M. Biel, and J. Berger. 2019. Seeking snow and breathing hard –

behavioral tactics in high elevation mammals to combat warming

temperatures. PLOS ONE 14.

Sedgeley, J.A. 2001. Quality of cavity microclimate as a factor influencing selection

of maternity roosts by a tree-dwelling bat, *Chalinolobus tuberculatus*, in New

Zealand. Journal of Applied Ecology 38:425–438.

Şekercioğlu, Ç.H., R.B. Primack, and J. Wormworth. 2012. The effects of climate

change on tropical birds. Biological Conservation 148:1–18.

Senior, P., R.K. Butlin, and J.D. Altringham. 2005. Sex and segregation in temperate

bats. Proceedings of the Royal Society B: Biological Sciences 272:2467–2473.

Sikes, R.S., and Animal Care and Use Committee of the American Society of

Mammalogists. 2016. 2016 guidelines of the American Society of

Mammalogists for the use of wild mammals in research and education. Journal

of Mammalogy 97:663–688.

Solick, D.I., and R.M.R. Barclay. 2006. Thermoregulation and roosting behaviour of

reproductive and nonreproductive female western long-eared bats (*Myotis*

evotis) in the Rocky Mountains of Alberta. Canadian Journal of Zoology

84:589–599.

Stawski, C., and F. Geiser. 2010. Fat and fed: frequent use of summer torpor in a

subtropical bat. Naturwissenschaften 97:29–35.

Stawski, C., and F. Geiser. 2012. Will temperature effects or phenotypic plasticity

613   determine the thermal response of a heterothermic tropical bat to climate

614   change? PLOS ONE 7:e40278.

615   Studier, E.H., and M.J. O'Farrell. 1976. Biology of Myotis thysanodes and M.

616   lucifugus (Chiroptera: Vespertilionidae)—III. Metabolism, heart rate,

617   breathing rate, evaporative water loss and general energetics. Comparative

618   Biochemistry and Physiology Part A: Physiology 54:423–432.

619   Vehtari, A., A. Gelman, and J. Gabry. 2017. Practical Bayesian model evaluation

620   using leave-one-out cross-validation and WAIC. Statistics and Computing

621   27:1413–1432.

622   Vuarin, P., M. Dammhahn, and P.-Y. Henry. 2013. Individual flexibility in energy

623   saving: body size and condition constrain torpor use. Functional Ecology

624   27:793–799.

625   Western Regional Climate Center. 2018. NCDC 1981-2010 monthly normals: Custer,

626   SD.

627   Willis, C.K.R., and R.M. Brigham. 2005. Physiological and ecological aspects of

628   roost selection by reproductive female hoary bats (Lasiurus cinereus). Journal

629   of Mammalogy 86:85–94.

630   Willis, C.K.R., and R.M. Brigham. 2007. Social thermoregulation exerts more

631   influence than microclimate on forest roost preferences by a cavity-dwelling

632   bat. Behavioral Ecology and Sociobiology 62:97–108.

633   Withers, P. C., C. E. Cooper, S. K. Maloney, F. Bozinovic, and A.P. Cruz-Neto. 2016.

634   Ecological and environmental physiology of mammals. First edition. Oxford

635   University Press, Oxford, U.K.

636   Wojciechowski, M.S., M. Jefimow, and E. Tęgowska. 2007. Environmental

637   conditions, rather than season, determine torpor use and temperature selection

638   in large mouse-eared bats (*Myotis myotis*). Comparative Biochemistry and

639   Physiology Part A: Molecular & Integrative Physiology 147:828–840

Figure Captions

Fig. 1. Heuristic diagram outlining the potential energetic benefit to an individual bat

of using periodic bouts of daily torpor rather than remaining in homeothermy at

ambient temperatures below the thermoneutral zone. This diagram is similar to a

classic Scholander curve except for one detail: while a Scholander curve illustrates

metabolic rate or energy expenditure at a constant ambient temperature and

physiological state (i.e., either homeothermy or torpor) in laboratory conditions, this

diagram illustrates energy expenditure when ambient temperature and physiological

state vary through time as they do in field conditions. Specifically, this diagram

assumes that (1) bats use more torpor when it is cold than when it is warm, (2)

ambient temperatures vary over the course of the day, and (3) ambient temperatures

below the thermoneutral zone are more prevalent than ambient temperatures above the

thermoneutral zone (see Cunningham et al. 2021). Each hypothetical relationship

would result in a different pattern of roost selection by animals seeking to minimize

energy expenditure during periods of inactivity. The black (A) line represents energy

expenditure over a day while maintaining homeothermy 100% of the time (i.e., never

using torpor). The red (B), grey (C), and blue (D) lines indicate energy expenditure

over a day while using some amount of torpor. For all three relationships, torpor

provides energy savings (i.e., the difference between the black and other lines), and

these savings are most pronounced at colder ambient temperatures. (B) For bats that

use torpor, energy expenditure *increases* at colder ambient temperatures because

while some energy is saved from employing torpor, maintaining homeothermy is

more costly at colder than at warmer ambient temperatures. A bat exhibiting this

relationship should select *warm* roosts to reduce energy use. (C) For bats that use

torpor, energy expenditure *is stable* across a wide range of ambient temperatures

because the energy saved from employing torpor matches (and thus offsets) the

increase in energy expended to maintain homeothermy at colder temperatures. A bat

exhibiting this relationship should not benefit from selecting either warm or cool

roosts, and should thus select neither warm nor cool roosts. (D) For bats that use

torpor, energy expenditure *decreases* at colder ambient temperatures because

relatively more energy is saved from using torpor even as maintaining homeothermy

is more costly at colder than at warmer ambient temperatures. A bat exhibiting this

relationship should select *cool* roosts to reduce energy use.

Fig. 2. Four competing sets of predictions of roost selection by a heterothermic bat.

Each column represents one of four sets of predictions, and each row represents a

statistical relationship consistent with the predictions. In column 1, energy

expenditure over the course of a day is higher in cool roosts than in warm roosts (1A).

In response, bats select warm roosts to minimize energy expenditure during the day

(1B). In this scenario, there should be no directional relationship between ambient

temperature and roost temperature (i.e., bats always select warm roosts regardless of

ambient temperature; 1C). In column 2, energy expenditure over the course of a day is

higher in warm roosts than in cool roosts (2A). In response, bats select cool roosts to

minimize energy expenditure during the day (2B). In this scenario, there should be no

directional relationship between ambient temperature and roost temperature (i.e., bats

always select cool roosts regardless of ambient temperature; 2C). In column 3, energy

expenditure over the course of a day is constant across roosts of all temperatures

(because bats can adaptively use torpor so that roost temperatures over the course of a

day have little influence on overall energy expenditure; 3A). Because energy

expenditure is consistent across roosts of all temperatures, bats do not select roosts

due to roost temperature (3B). In this scenario, there is no relationship between

ambient temperature and roost temperature (i.e., bats never select roosts due to

temperatures within roosts, regardless of ambient temperature; 3C). In column 4,

energy expenditure peaks at intermediate roost temperatures where bats use relatively

little torpor but the costs of maintaining homeothermy are relatively high (4A). In

response, bats select cool roosts on cool days and warm roosts on warm days (4B)

because torpor saves more energy in cool roosts than in warm roosts. In this scenario,

the relationship between ambient temperature and roost temperature should be

positive (i.e., bats select warmer roosts on warmer days; 4C).

Fig. 3. Results of our simulation of daily energy expenditure by fringed myotis over

the range of temperatures observed in used roosts. Each point represents one day. The

red points represent estimated daily energy expenditure if bats never used torpor. The

blue points represent our estimate of energy expenditure over the course of a day if

part of the day is spent in torpor (with daily duration of torpor a function of daily

ambient temperature as observed in our study). The lines represent loess regressions

of the relationship between roost temperature and daily energy expenditure. Estimates

of daily energy expenditure incorporating observed bat behavior are steady across all

roost temperatures observed during our study. The blue points in this figure

correspond with Row A in Fig. 2, and are most closely matched by Fig 2.3A.

Fig. 4. Kernel density plots comparing thermal characteristics within used and available roost structures: mean temperature (A), time of day at peak temperature (B), and the standard deviation of temperature (C). Blue distributions represent used roosts, while orange distributions represent available roosts. These plots illustrate the results of our binomial model of roost selection. Used roosts were slightly warmer on average than available roosts, but their distributions largely overlapped (A). Temperatures peaked slightly earlier in used roosts than available roosts, but this was a function of temperatures in warmer roosts tending to peak earlier in the day ($r = -0.19$ for the relationship between mean temperature within roost structures and time of day at peak temperature) and their distributions largely overlap (B). The standard deviation in temperatures within used roosts is very similar to the standard deviation in temperatures within available roosts, although bats did not use the few roost structures with very high standard deviations (C). Panel A in this figure corresponds with Row B in Fig. 2, and is most closely matched by Fig. 2.3B.

Fig. 5. Scatter plot of the relationship between ambient temperature on a given day and mean temperature within used roosts. Each point is based on observed data, and represents a roost used for one day; some roosts (n = 14) were used on multiple days and are thus represented by multiple points on this plot. The line represents the regression line for this relationship and the grey band represents 95% confidence intervals. Ambient temperature on a given day did not influence whether bats used warm or cool roosts ($p = 0.06$; $R^2 = 0.04$). This figure corresponds with Row C in Fig. 2, and is most closely matched by Fig. 2.3C.

735   **Fig. 1**

737 **Fig. 2**

739 **Fig. 3**

741   **Fig. 4**

743 **Fig. 5**

Appendix S1: Supplementary Data

Descriptive Statistics for Rock vs. Tree Roost Structures

During the day, rock crevices averaged 20.1˚C (range: 16.5˚ – 24.2˚C) while tree roost structures averaged 18.8˚C (range: 16.1˚ – 25.5˚C). Daily maximum temperatures within rock crevices averaged 26.1˚C (range: 17.9˚ – 40.8˚C), while daily maximum temperatures within tree roost structures averaged 28.3˚C (range: 21.0˚ – 52.1˚C). Temperatures within rock crevices peaked at 1441 hrs on average (range = 1005 – 1742 hrs), while temperatures within tree roost structures peaked at 1357 hrs on average (range = 1056 – 1659 hrs). Ambient temperature strongly influenced temperatures within roost structures. Temperatures within rock crevices at each hour (in ˚C) followed the equation $7.67 + 0.73*$ambient temperature ($R^2 = 0.54$), while temperatures within tree roost structures at each hour followed the equation $1.63 + 1.00*$ambient temperature ($R^2 = 0.63$).

Temperatures within used rock crevices averaged 20.5˚C (range: 16.8˚ – 23.3˚C) while temperatures within available rock crevices averaged 19.9˚C (range: 16.5˚ – 24.2˚C). Temperatures within used tree roosts averaged 18.6˚C (range: 17.4˚ – 20.4˚C) while temperatures within available tree cavities averaged 19.2˚C (range: 16.1˚ – 25.5˚C) and temperatures within available spaces under sloughing bark averaged 18.4˚C (range: 16.1˚ – 21.0˚C).

Temperatures within used rock crevices peaked on average at 1414 hrs (range: 1105 – 1719 hrs), while temperatures within available rock crevices peaked on average at 1458 hrs (range: 1005 – 1742 hrs). Temperatures within used tree roosts peaked on average at 1447 hrs (range: 1125 – 1659 hrs), while temperatures within available tree cavities peaked on average at 1410 hrs (range: 1120 – 1608 hrs) and

769   temperatures within available spaces under sloughing bark peaked on average at 1349

770   hrs (range: 1056 – 1608 hrs).

771   The standard deviation of temperatures within used rock crevices was 6.7˚C

772   (range: 4.3˚ – 10.0˚C), while the standard deviation of temperatures within available

773   rock crevices was 6.2˚C (range: 3.2˚ - 11.0˚C). The standard deviation of temperatures

774   within used tree roosts was 7.7˚C (range: 6.7˚ - 9.1˚C), while the standard deviation of

775   temperatures within available tree cavities was 8.7˚C (range: 5.9˚ - 16.4˚C) and within

776   available spaces under sloughing bark was 7.7˚C (range: 6.5˚ - 11.0˚C).

777   There was no difference in ambient temperature between days where rock

778   crevices were used and days where tree roost structures were used (Mann-Whitney U

779     $= 299$; $p = 0.968$).

Table S1. Information on torpor use by bats tracked during our study, including an ID number for each individual, the dates for which we have data, the mass of bats at time of capture, the timing of torpor entry and exit for morning and afternoon bouts of torpor, the duration of periods of periods of torpor in both mornings and afternoons, and the total duration of torpor across the day.

Bat ID	Date	Mass (grams)	AM Torpor Start Time (hrs)	AM Torpor End Time (hrs)	Duration of AM Torpor (mins)	PM Torpor Start Time (hrs)	PM Torpor End Time (hrs)	Duration of PM Torpor (mins)	Total Torpor Duration (mins)
172_063	8/5/2017	6.02	517	1456	579	2013	2055	42	621
172_063	8/6/2017	6.02	451	1210	439	1910	2037	87	526
172_063	8/7/2017	6.02	2245	1557	1032	1840	2044	124	1156
172_904	6/28/2018	6.75	425	733	188	1825	2057	125	313
172_904	6/29/2018	6.75	419	1037	378	1603	2114	277	655
172_904	7/3/2018	6.75	525	944	259	1834	2029	115	374
172_904	7/4/2018	6.75	412	1446	634	1709	2122	253	887
172_904	7/5/2018	6.75	424	1458	597	1930	2043	73	670
172_904	7/6/2018	6.75	511	1016	305	-	-	0	305
172_904	7/7/2018	6.75	438	818	220	-	-	0	220
172_692	7/13/2018	6.92	445	830	225	1936	2043	67	292
172_692	7/14/2018	6.92	435	815	220	-	-	0	220
172_632	7/20/2018	8.04	426	1102	396	1916	2041	85	481
172_753	7/27/2018	8.16	133	2045	1152	-	-	0	1152
172_753	7/28/2018	8.16	2300	2031	1291	-	-	0	1291
172_453	8/4/2018	7.1	449	959	310	1915	2039	84	394
172_784	8/4/2018	7.53	442	1028	346	1951	2023	32	378
172_453	8/5/2018	7.1	459	1156	417	1613	2028	255	672

172_784	8/5/2018	7.53	445	1100	375	1852	2019	87	462
172_453	8/6/2018	7.1	441	916	275	1823	2034	131	406
172_784	8/6/2018	7.53	449	1003	314	-	-	0	314
172_453	8/7/2018	7.1	444	1041	357	-	-	0	357
172_784	8/7/2018	7.53	502	850	228	-	-	0	228
172_063	8/8/2018	6.02	2335	1427	892	1737	2009	152	1044
172_453	8/8/2018	7.1	451	839	228	-	-	0	228
172_784	8/8/2018	7.53	439	852	253	-	-	0	253
172_453	8/10/2018	7.1	456	843	227	-	-	0	227

784   **Fig. S1.** An example of raw skin temperature data that we used to delineate bouts of torpor. Periods of time in red blocks represent periods of

785   activity (flying, foraging, etc.), periods of time in blue blocks represent periods of torpor, and periods in white represent periods of homeothermy

786   or transition between torpor and homeothermy/activity. To delineate bouts of torpor, we used the definition suggested in Barclay et al. (2001).

787
788

789   **Fig. S2**. Scatter plot illustrating the conditional effect of daily mean ambient temperature on the total duration of bouts of torpor during the day.

790   Each point is based on observed data and represents one day. The line represents the regression line for this relationship and the grey band

791   represents 95% credible intervals around this line. Credible intervals for this conditional effect did not cross zero (parameter estimate: -37.4 min;

792   95% credible intervals: -64.0 – -12.6 min), indicating that bats spent ca. 37 minutes less in torpor per day for each additional 1˚C in daily mean

793   ambient temperature between 0445 hrs and 2100 hrs.