1 **Longer study length, standardized sampling techniques, and broader geographic scope**
2 **leads to higher likelihood of detecting stable abundance patterns in long term black-legged**
3 **tick studies**
4
5
6

7 Rowan Christie[1*], Kaitlin Stack Whitney[1,2*], Julia Perrone[3], Christine A. Bahlai[4]
8
[1] Thomas H Gosnell School of Life Sciences, Rochester Institute of Technology, Rochester, NY,
USA
[2] Science, Technology & Society Department, Rochester Institute of Technology, Rochester, NY,
USA
[3] School of Information, Kent State University, Kent, OH, USA
[4] Department of Biological Sciences and Environmental Science and Design Research Initiative,
Kent State University, Kent, OH, USA
16
17 *co-first author status
18
19 Corresponding author:
20 Kaitlin Stack Whitney[1,2]
21 7 Lomb Memorial Drive, Rochester, NY, 14623, USA
22 Email address: kxwsbi@rit.edu

23
24
25

Abstract:

Background.
Understanding how study design and monitoring strategies shape inference within, and synthesis across, studies is critical across biological disciplines. Many biological and field studies are short term and limited in scope. Monitoring studies are critical for informing public health about potential vectors of concern, such as *Ixodes scapularis* (black-legged ticks). Black-legged ticks are a taxon of ecological and human health concern due to their status as primary vectors of *Borrelia burgdorferi*, which causes Lyme disease. However, variation in black-legged tick monitoring, and gaps in data, are currently considered major barriers to understanding population trends and in turn, predicting Lyme disease risk. To understand how variable methodology in black-legged tick studies may influence which population patterns researchers find, we conducted a data synthesis experiment.

Materials and Methods. We searched for publicly available black-legged tick abundance datasets that had at least 9 years of data, using keywords about ticks in internet search engines, literature databases, data repositories and public health websites. Our analysis included 289 datasets from 7 surveys from locations throughout the US, ranging in length from 9 to 24 years. We used a moving window analysis, which is a kind of non-random resampling approach, to investigate the temporal stability of black-legged tick population trajectories across the US. We then used t-tests to assess differences in stability time across different study parameters.

Results. All of our sampled datasets required 4 or more years to reach stability. We also found that several study factors can have an impact on the likelihood of a study reaching stability and of data resulting in misleading results if the study does not reach stability. Specifically, datasets collected via dragging reached stability significantly faster than data collected via opportunistic sampling. Datasets that sampled larva reached stability significantly later than those that sampled adults or nymphs. Additionally, datasets collected at the broadest spatial scale (county) reached stability fastest.

Conclusion. We used 289 datasets from 7 long term black-legged tick studies to conduct a non-random data resampling experiment, revealing that sampling design does shape inferences in black-legged tick population trajectories and how many years it takes to find stable patterns. Specifically, our results show the importance of study length, sampling technique, life stage, and geographic scope in understanding black-legged tick populations, in the absence of standardized surveillance methods. Current public health efforts based on existing black-legged tick datasets must take monitoring study parameters into account, to better understand if and how to use monitoring data to inform decision making. We also recommend that potential forecasting initiatives consider these parameters when projecting black-legged tick population trends.

Introduction

Introduction
Understanding how study design and monitoring strategies shape inference within, and synthesis
across, studies is critical across biological sciences. This is especially important as climatic
conditions and ecosystems change in stark and unpredictable ways, requiring scientists to tease
apart drivers of population and community trends from the past to understand possible futures
(Bahlai et al., 2021). Many biological and field studies are short term and limited in scope,
leading to concerns about how to interpret the many short-term studies in comparison with the
relatively fewer long term ecological studies (Wauchope et al., 2019; Bahlai et al., 2021). Shorter
term studies increase the likelihood of finding misleading trends, potentially leading to
misinformed management approaches (White and Bahlai, 2021). Natural variability in a stable
population might lead a 2-3 year study to appear to capture a strong upward trend in abundance
for a given taxon, simply as a result of natural population cycling (Fournier et al., 2019).
79
Monitoring studies are critical for informing public health about potential vectors of concern.
Surveys indicate that the control and management of vector borne diseases is of immediate
concern for public health in the United States (Hill et al., 2009). Public health officials must
know the geographical range of infectious vectors and their level of abundance to understand
potential disease risk (Townson et al., 2005). However, standardizing results from multiple
studies on vectors can be problematic. Novel sampling approaches and methodologies are
required to effectively monitor variable vector populations (Sougoufara et al., 2020). Reliable
inference about population trends from single studies, or synthesizing across them, must account
for variation in monitoring methods.
89
A critical vector of concern for public health in the US is ticks. They are common vectors that
can transmit multiple diseases to humans and are responsible for vectoring the greatest number
of human diseases in the US (Eisen et al., 2017). Increased outdoor activity, especially during the
ongoing COVID-19 pandemic, may be exposing more people to potential tick-borne illnesses
(HHS 2020). This pattern matches broader tick trends. Globally, both the known range of tick
species and the reported cases of tick-vectored illnesses are increasing (Paddock et al., 2016).
96
Ticks and tick-vectored diseases are commonly monitored using dragging, flagging, and carbon
dioxide baited traps (Falco and Fish, 1992). Dragging is a method where a person pulls a
horizontal pole behind them across the ground and ticks latch onto the cloth attached to the pole
(Salomon et al., 2020). Another common tick survey method is passive surveillance, where
individuals send in ticks they have found on themselves to public health departments, but do not
deliberately search for ticks (Ripoche et al., 2018). While these are common, there is a wide
variety of methods reported across studies. For example, one method of surveying ticks has been
collecting them from hunter-killed deer in the midwestern US (Egizi et al., 2018; Lee et al.,
2013). Additionally, some studies measure density instead of abundance (Kahl et al., 2002), use
distance-based sampling (Gray et al., 1992), or are modeled projections of where ticks are likely
to be (Baldwin et al., 2022), based on varying assumptions about climate and landscape factors
(Estrada-Peña et al., 2013). The lack of a shared and standardized sampling method may be
preventing effective understanding of changes in abundance and distribution of ticks synthesized
across surveys (Dennis et al., 1998; Eisen et al., 2016).
111

112 The lack of a standardized sampling method may also impact the ability to accurately monitor
113 tick-vectored disease. Recent research has identified the lack of standardized data collection, and
114 subsequent data sharing, as impeding researchers' understanding of tick-transmitted pathogens
115 (Estrada-Peña et al., 2021).
116
This is critical, as Lyme disease is a common and increasing illness impacting US residents.
Recent research estimates that more than 400,000 people annually have been diagnosed in the
US between 2010 and 2018 (Kugeler et al., 2021). The primary vectors of *Borrelia burgdorferi*,
bacteria that cause Lyme disease, are Black-legged ticks (Des Vignes and Fish, 1997). Black-
legged ticks are common throughout the US - and both widespread and abundant. Established
black-legged tick populations are most concentrated in upper north-central, northeastern, and
west-coast states (Eisen et al. 2016). Black-legged ticks transmit seven human pathogens of
concern, including *B. burgdorferi*, and both the tick and pathogen transmission have increased
over the past two decades in the US (Eisen and Eisen 2018). In addition, Black-legged ticks are
spreading rapidly, found in more than twice the number of counties in the United States
compared to twenty years ago (Eisen et al. 2016). Systematic surveillance of ticks and tick-borne
pathogens is important for understanding Lyme disease risk (Eisen and Paddock 2021). Variation
in black-legged tick monitoring, and gaps in data, are currently considered major barriers to
predicting Lyme disease risk (Kugeler and Eisen 2020). Understanding tick prevalence and
potential Lyme disease risk can help shift the burden of tick vectored disease management from
individual prevention to states or other public health agencies (Eisen 2020).
133
To understand how variable methodology in black-legged tick studies may influence which
population patterns researchers find, we conducted a data synthesis experiment. To do this, we
used a novel nonrandom sampling approach (Bahlai et al., 2021) to analyze publicly available
black-legged tick datasets from long term monitoring studies to understand the impacts of study
parameters on inferences about black-legged tick abundance and trajectory. Specifically, we
examined study length, sampling technique, life stage, geographic scope, and sampling metrics.
We anticipated that studies using dragging would have less variable findings than other sampling
techniques, due to existing research showing that dragging is more consistent for some life stages
of black-legged tick collection (Falco and Fish, 1992). We also anticipated that the tick life stage
monitored could influence expected trends. Nymphs are most likely to spread *Borrelia*
burgdorferi (Rulison et al., 2013), and may have different abundance patterns than adults or
larvae. Results from this analysis can help researchers contextualize existing monitoring data
and the potential opportunities and challenges when extrapolating from the past to the future.
This data experiment may also help researchers better understand how to design black-legged
tick monitoring studies to avoid misleading inferences, potentially leading to improved
surveillance of a public health risk.
150
Materials & Methods
Data searching
We searched for publicly available black-legged tick abundance dataset that had at least 9 years
of data, using keywords about black-legged ticks in internet search engines, literature databases,
data repositories and public health websites. Specifically, we searched combinations of *tick* and
scapularis in Data Dryad, Data One, Google Datasets, the Long Term Ecological Research data
portal, and the National Ecological Observation Network data portal. We then used Google

search portals to look up individual state department of health websites and state datasets where
black-legged ticks are common for additional sets of survey data. Our complete search was done
using the PRISMA workflow for systematic literature screening, review and reporting (Moher et
al., 2009; Appendix 1).
162
Black-legged tick datasets
Our search yielded 289 datasets from 7 surveys from locations in New Jersey, New York, Iowa,
Connecticut, Massachusetts, ranging in length from 9 years to 24 years. We extracted and scored
information about each dataset in terms of the life stage sampled, sampling techniques used,
geographic scale of the study, and location (Table 1). The surveys varied in the life stages
included (Table 1). Black-legged ticks go through four life stages: egg, larva, nymph, and adult
(CDC 2011). Our search yielded surveys that collected data on adults only, nymphs only, adults
and nymphs, or all life stages other than egg. These included datasets that studied adults (oldest
life stage, n=63 datasets), nymphs (middle life stage, n=68 datasets), and larvae (youngest life
stage, n=8 datasets). Additional datasets (n=150) did not specify the life stage collected and were
not included in subsequent analyses about life stage. The search results also included datasets
that either only included abundance - or also included analysis of the percentage of ticks infected
with *B. burgdorferi*. Across the surveys, we found most measured the abundance of black-legged
ticks (n=175 datasets) and some tested which percentage had *B. burgdorferi* (n=114 datasets).
Surveys also varied in their geographic scope (Table 1). Three surveys were collected and
reported at the county scale, while two were reported by town, and two at smaller spatial scales
(grid and state forest). The studies included datasets summarized by grid (n=24 datasets), state
forest (n=6 datasets), town (n=186 datasets), and county (n=73 datasets). The surveys found in
our search also included two sampling techniques, dragging (n=90 datasets) and opportunistic
surveying of ticks found on individuals (n=198 datasets) (Table 1). One additional dataset
instead recorded "bites found on a person" (n=1), instead of ticks, and was not included in
subsequent analyses.
185
Moving window analysis
To analyze the stability of the abundance trends of these datasets, we used the 'broken window'
algorithm (Bahlai et al. 2021) using RStudio and R statistical software (RStudio Team, 2020; R
Core Team, 2020). This non-random resampling approach uses subsamples (i.e. '*windows*') of
the time series data to gain insights into patterns of how data behaves in arbitrarily selected time
periods (Bahlai et al. 2021). Non-random resampling of existing monitoring data is a powerful
and underused tool to understand trends (White and Bahlai 2021).
193
This algorithm defines the linear slope associated with the longest time series available as the
'true' slope (as it is calculated using the largest possible sample size) and performs calculations
relative to this proxy for comparison. The algorithm uses a yearly interval for its calculations,
and in the case of this study, the metric used as the dependent variable was some unit of
abundance or activity of ticks, as defined by the trapping method. Prior to subsetting within the
algorithm, each raw dataset is subjected to normalization by converting it to a unitless Z score,
allowing methods producing data on dramatically different response scales to be directly
compared. We then ran a '*moving window*' function in the algorithm, which iterates through all
subsets in the data greater than 2 years, and subjects each 'window' to a linear model, which

calculates the slope statistic (change of standardized density over change in year), standard error
of this slope, p-value, and R^2.
Then we ran the '*stability time*' function in the same package to calculate the number of years to
reach stability (Bahlai et al., 2021). The number of years to reach stability is calculated in the
function by using the summary statistics previously calculated for each dataset to compute the
proportion correct for each window within the standard deviation of the longest slope, and
returns the number of windows whose slope are within said standard deviation (95% CI). We
also ran the '*proportion_wrong'* function in the same package (Bahlai et al., 2021), to calculate
the proportion of windows that did not reach stability with a p-value that was not within 0.05 of
the p-value of the true slope (slope of the longest series). The 'significantly wrong' windows
would likely lead a scientist to a spurious conclusion that does not agree with the overall trend.
We calculated both the proportion of 'significantly wrong' windows before stability time and
overall, for each dataset.
216
Statistical analyses
Following the use of the 'broken windows' algorithm, we assessed if and how study parameters
affected years to stability time, the overall proportion wrong, and the proportion wrong before
stability was reached for each dataset. We did this by comparing the means between different
groups using independent samples t-tests in base R using RStudio and R statistical software
(RStudio Team, 2020; R Core Team, 2020). Using this method, we compared results from
datasets that varied by life stage studied, the geographic scope of the study, sampling techniques
used, and the study response variables. For example, we tested whether the mean stability time
for datasets collected using dragging sampling methods was significantly different from the
mean stability time for datasets collected using opportunistic sampling methods. We also used
Pearson product moment correlation tests in base R using RStudio and R statistical software
(RStudio Team, 2020; R Core Team, 2020). These tests were used to assess potential
correlations between study length and stability time, overall proportion wrong, and the overall
proportion wrong compared to proportion wrong before stability was reached. The dataset, R
code, test results, and resulting figures are also provided in the Supplemental Materials.
232
Results
Study length
Using the 'broken window' algorithm, we found that across all datasets, years to stability time
ranged from 4 to 23 years (Figure 1). The proportion of datasets that reached stability by 4 years
is very low; by 10 years, the majority of datasets had reached stability (Figure 1). We found that
study length was positively correlated with years to stability (Pearson product moment
correlation test, r^2 = 0.9, p-value <0.001) (Figure 2). From our analysis, we also found that the
proportion of windows significantly wrong before datasets reached stability ranged from 0 to
100% (Figure 3). We also found that the association between proportion significantly wrong
before stability time and overall proportion wrong were not strongly correlated (t = -1.2, df =
571.46, p-value = 0.23, Figure 3).
244
Sampling technique
We found that datasets with dragging techniques reached stability time faster than datasets
collected using opportunistic, passive sampling methods (t = -8.53, df = 236.23, p-value <0.001)
(Figure 4A). The median number of years to reach stability was 7 for dragging techniques and 12

for opportunistic techniques. However, we did not find a significant difference when we
compared the proportion significantly wrong before stability by sampling technique (t = 0.08, df
= 155.58, p-value = 0.93, Figure 4B). Datasets with dragging sampling techniques in our data
experiment also exhibited a smaller range of years to stability, compared to datasets with
opportunistic passive sampling methods.
254
Life stage
Datasets measuring black-legged tick larvae reached stability slower than datasets measuring
adult or nymph abundance data (adults and larvae, t = -5.96, df = 10.11, p-value <0.001; nymphs
and larvae, t = -5.52, df = 10.54, p-value <0.001, Figure 5A). We found no difference in the time
to stability between datasets measuring adults or nymph blacklegged ticks (t = -0.64, df =
126.84, p-value = 0.52). The median time for datasets measuring larvae to reach stability was
11.5 years, compared to 7 years for adults and nymphs.
262
The mean proportion wrong before stability for adults was 0.35, compared to nymphs (0.16) and
larvae (0.28). Datasets measuring nymph tick life stages had a significantly lower proportion of
windows wrong before stability time compared to adults (t = 2.90, df = 113.64, p-value = 0.004,
Figure 5B). We found no difference in the proportion wrong before stability time between larvae
and nymphs (t = -0.75, df = 7.89, p-value = 0.47) and larvae and adults (t = 0.44, df = 8.77, p-
value = 0.67)
269
Geographic scope
Blacklegged tick datasets from studies reporting data from counties reached stability in fewer
years than datasets reported at smaller spatial scales (town, state forest, or grid) (county and grid,
t = -17.21, df = 33.87, p-value <0.001; county and state forest, t = -5.55, df = 5.22, p-value =
0.002; county and town, t = -17.03, df = 243.09, p-value <0.001; Figure 6A). The median
number years to stability for datasets of county scope was 7 years, compared to 11.5 years for
grid and 12 years for town and state forest. We found no difference in years to stability
comparing across datasets reported from towns, state forests, or grids (town and state forest, t =
0.47, df = 6.12, p-value = 0.65; town and grid, t = 0.22, df = 86.18, p-value = 0.82; state forest
and grid, t = -0.38, df = 6.03, p-value = 0.72; Figure 6A).
280
Datasets from studies reporting data from state forests had a significantly lower proportion of
wrong windows, as measured by the proportion wrong before stability time, compared to
datasets reported from counties and towns (state forest and county, t = 5.24, df = 74.29, p-value
<0.001; state forest and town, t = 7.42, df = 46.44, p-value = p-value <0.001; Figure 6B). Studies
reported from grids did not have a higher proportion of windows wrong before stability time (t =
-1.91, df = 25.10, p-value = 0.068). The mean proportion significantly wrong for data at the scale
of state forest is 0.077, compared to 0.35 of county datasets, 0.31 of town datasets, and 0.23 of
grid datasets. The proportion of wrong windows did not significantly vary between county, town,
and grid studies (county and town, t = 0.72, df = 115.03, p-value = 0.48; county and grid, t=
1.33, df = 43.07, p-value = 0.19; town and grid, t = 1.00, df = 28.52, p-value = 0.32; Figure 6B).
291
Sampling metric
We found no difference in the number of years until a dataset reached stability time comparing

datasets measuring the abundance of black-legged ticks and datasets that tested for *B.*
burgdorferi infection (t = -1.29, df = 283.9, p-value = 0.20; Figure 7A). The datasets also did not
significantly vary in the proportion of wrong windows, as measured by the proportion wrong
before stability time (t = -1.18, df = 232.98, p-value = 0.24; Figure 7B).
298
Discussion
Overall, we found several aspects of how blacklegged ticks are monitored and data reported may
have an impact on the likelihood of a study reflecting longer-term population trends. This is
supported by our findings about datasets reaching stability time, as well the proportion of
significantly wrong windows, varying by study methods. Specifically, datasets using dragging,
counting nymph or adult life stages, and reported by county, reached stability faster than other
datasets. These findings support our hypothesis that the dragging method would be associated
with more consistent patterns than other sampling methods. We also predicted that datasets
collected on different blacklegged tick life stages may vary in stability time; however, we did not
anticipate that larvae would reach stability time slower than other life stages. Our results also
indicate the importance of long term monitoring, as we found that the longer datasets were
correlated with more years to reach stability and many datasets with higher proportions of
misleading windows before stability time. As none of the datasets assessed reached stability time
before 4 years, our analyses indicate that short term studies may be insufficient to explain long
term dynamics in blacklegged ticks.
314
Our findings are consistent with previous research supporting the importance of long term
ecological monitoring for understanding populations in field biology. Several other studies
provide support for using long term studies to understand patterns (White, 2019; Cusser et al.,
2020; Cusser et al. 2021). For example, White found that at least 72% of the vertebrate
populations they studied required at least 10 years of study to detect a significant trend in
abundance, while Cusser et al. determined that it would take at least 16-19 years to detect a
significant effect from no-till management on crop yield and soil moisture, emphasizing the
importance of long term studies. Similarly, we predicted and found that the dragging sampling
technique would produce more consistent abundance trends than datasets associated with
opportunistic surveying. This was expected given research indicating that dragging can be a
consistent way to monitor black-legged ticks (Falco and Fish, 1992). However, other research
has found that data produced using different tick sampling methods is similar (Rulison et al.,
2013). Yet even more recent research attempting to validate dragging with other tick species in
Sweden concluded that dragging was not recommended for tick adults or larvae, due to low
repeatability and agreement (Kjellander et al., 2021). This may be due to the lack of
standardization within dragging cloths or other within-method variability between surveys
(Newman et al., 2019).
332
We anticipated, and found, that datasets studying different life stages of ticks would vary in how
many years it took to reach stability time. Specifically, time to stability was longer for datasets
monitoring larvae, the youngest life stage, compared to adults and nymphs. Additionally,
datasets monitoring adult black-legged ticks had higher proportions of significantly wrong
windows. This finding is critical for researchers trying to glean insights into broad scale black-
legged tick trends across disparate study methods. This suggests that datasets that focus on adults
will be more likely to lead to misleading results if the study has not reached stability, which

supports longer-term studies. This finding is critical for understanding how to apply black-legged
tick monitoring data to potential Lyme disease risk, as different life stages may pose different
levels of exposure risk to humans. Nymph, not adult, abundance has been shown to be the
primary factor in determining risk of exposure (Falco et al., 1999). Due to this potential variation
in transmission by life stage, the recent US Centers for Disease Control and Prevention (CDC)
report on black-legged tick surveillance provides distinct guidance on collecting and interpreting
data from each life stage (CDC 2021). Our results support this distinction. However, life stage is
distinct from other categorizations of tick life history thought to impact abundance and disease
transmission, such as 'physiological age,' a measure that groups ticks based on environmental
stress events (Pool et al., 2017). There may be important variation among ticks of the same life
stage, based on their physiological age, that surveyors are not assessing in the field, as this may
require the addition of molecular biology approaches (Estrada-Peña et al., 2021). And even
though we found no difference in our results when comparing the number of years to reach
stability and proportion wrong before stability in datasets that measured black-legged tick
abundance and the percent with *Borrelia burgdorferi*, this may be due to additional variation
within studies measuring pathogen prevalence within ticks. Some studies test pools of ticks for
infection to reduce cost and labor, and variation in the number of ticks processed can impact the
perceived estimation of prevalence (Estrada-Peña et al., 2021).
358
Our results indicate that sampling a larger geographic area leads to more consistent patterns in
black-legged tick population dynamics than studies conducted on smaller areas. This is
supported by previous work finding high levels of fine-scale spatial distribution of black-legged
ticks (Dumas et al., 2022, Mathews-Martin et al., 2020). Fortunately, county-level information is
a common geographic scale for documenting the presence and abundance of blacklegged ticks
and Lyme disease (e.g. Fleshman et al., 2021). Given that ecological monitoring requires
tradeoffs in effort, cost, and insight (Bennett et al., 2018), this finding may indicate that studies
at larger spatial scales may be able to give indication of population trajectories faster than studies
at smaller scales.
368
Our findings also support the collection of long term tick datasets. Even with the improved
insights from longer monitoring studies, managers and policymakers will still need to make
decisions without complete information. Determining what is 'enough' data for evidence-based
decision making is a challenge (Canessa et al., 2015). As the lack of shared, standardized
sampling methods may hamper the ability to monitor changes in abundance and distribution of
ticks (Dennis et al., 1998; Eisen et al., 2016), understanding patterns and differences between
datasets may help. Our results can help inform if and how black-legged tick monitoring data can
be reasonably interpreted based on study parameters to make informed, rigorous claims to
support management decisions that can help reduce Lyme disease exposure and risk. Moreover,
we further demonstrate the value of resampling existing data to give insights into analyzing
existing data and designing future monitoring programs (White and Bahlai, 2021).
380
Understanding population trajectories is also critical for using ecological data for ecological
forecasting. Ecological forecasting is considered the next critical frontier in broad scale
environmental science (Dietze et al., 2018). Ecological forecasting initiatives are increasing, with
an NSF-funded Research Coordination Network (NSF, 2019), conferences (EFI, 2019; EFI RCN
and NEON, 2020), and ongoing ecological forecast challenges (EFI RCN NEON, 2020). With

study organisms such as black-legged ticks, forecasting could be used to proactively protect
human health. As standardized data has been identified as critical for understanding tick-
transmitted pathogens (Estrada-Peña et al., 2021), future work should examine how these study
parameters, identified as critical through retrospective data resampling, shape forecasted patterns
and simulations. This is especially important for forecasting disease risk, as data gaps and
variation in monitoring data are already considered major challenges (Kugeler and Eisen 2020).
Our results highlight the need to better understand and explicitly incorporate that variability into
the interpretation of black-legged tick data.
394
Conclusions
We used 289 datasets from 7 long term black-legged tick studies to conduct a moving windows
data resampling experiment, revealing that sampling design does shape inferences in black-
legged tick population trajectories and the number of years to reveal stable patterns. Specifically,
our results show the importance of study length, sampling technique, life stage, and geographic
scope in understanding black-legged tick populations, in the absence of standardized surveillance
methods. Current public health efforts based on existing black-legged tick datasets must take
monitoring study parameters into account, to better understand if and how to use monitoring data
to inform decisioning. We also advocate that potential future forecasting initiatives consider
these parameters when projecting future black-legged tick population trends. Understanding how
study design and monitoring shapes synthesis is critical across many biological disciplines.
406
Acknowledgments

RMC was supported by the National Institute of General Medical Sciences of the National
Institutes of Health under Award Number R25GM122672. CAB, JP, and KSW were supported
by the Office of Advanced Cyberinfrastructure in the National Science Foundation under Award
Number #1838807. The content is solely the responsibility of the authors and does not
necessarily represent the official views of the National Institutes of Health or the National
Science Foundation.

References

Bahlai CA, White ER, Perrone JD, Cusser S, Stack Whitney K. 2021. The broken window: An algorithm for quantifying and characterizing misleading trajectories in ecological processes. Ecological Informatics, 64, 101336. https://doi.org/10.1016/j.ecoinf.2021.101336

Baldwin H, Landesman WJ, Borgmann-Winter B, Allen D. 2022. A geographic information system approach to map tick exposure risk at a scale for public health intervention. Journal of Medical Entomology 59(1) 162-172. https://doi.org/10.1093/jme/tjab169

Bennett JR, Maxwell SL, Martin AE, Chadés I, Fahrig L, Gilbert B. 2018. When to monitor and when to act: Value of information theory for multiple management units and limited budgets. J. Appl. Ecol. 55, 2102–2113. doi: 10.1111/1365-2664.13132

Centers for Disease Control and Prevention (CDC). 2021. Surveillance for Ixodes scapularis and pathogens found in this tick species in the United States. Report by the Centers for Disease Control and Prevention, National Center for Emerging and Zoonotic Infectious Diseases (NCEZID), Division of Vector-Borne Diseases (DVBD). Page last updated April 21, 2022. Available at <https://www.cdc.gov/ticks/resources/TickSurveillance_Iscapularis-P.pdf> (accessed 15 November 2021)

Centers for Disease Control and Prevention (CDC). 2011. Lifecycle of Blacklegged ticks. Available at https://www.cdc.gov/lyme/transmission/blacklegged.html (accessed May 23, 2022)

Canessa S, Guillera-Arroita G, Lahoz-Monfort JJ, Southwell DM, Armstrong DP, Chadés I, Lacy RC, Converse SJ. 2015. When do we need more data? A primer on calculating the value of information for applied ecologists. Methods Ecol. Evol. 6, 1219–1228. doi: 10.1111/2041-210X.12423

Cusser S, Helms IV J, Bahlai C, Hadded NM. 2021. How long do population level field experiments need to be? A meta-analysis across the 40-year old LTER network. Ecology Letters https://doi.org/10.1111/ele.13710

Cusser S, Bahlai C, Swinton SM, Robertson GP, Haddad NM. 2020. Long□term research avoids spurious and misleading trends in sustainability attributes of no□till. Glob Change Biol. 26: 3715– 3725. https://doi.org/10.1111/gcb.15080

Dennis DT, Nekomoto TS, Victor JC, Paul WS, Piesman J. 1998. Reported distribution of Ixodes scapularis and Ixodes pacificus (Acari: Ixodidae) in the United States. Journal of Medical Entomology 35(5):629-38. doi: 10.1093/jmedent/35.5.629. PMID: 9775584.

Des Vignes F, Fish D. 1997. Transmission of the agent of human granulocytic ehrlichiosis by host-seeking Ixodus scapularis (Acari:Ixodidae) in southern New York state. Journal of Medical Entomology, 34(4), 379–382. https://doi.org/10.1093/jmedent/34.4.379

Dietze, MC, Fox A, Beck-Johnson LM, Betancourt JL, Hooten MB, Jarnevich CS, Keitt TH, Kenney MA, Laney CM, Larsen LG, Loescher HW, Lunch CK, Pijanowski BC, Randerson JT, Read EK, Tredennick AT, Vargas R, Weathers KC, White EP. 2018. Iterative near-term ecological forecasting. Proceedings of the National Academy of Sciences 115 (7) 1424-1432; DOI: 10.1073/pnas.1710231115

Dumas A, Bouchard C, Lindsay LR, Ogden NH, Leighton PA. 2022. Fine-scale determinants of the spatiotemporal distribution of Ixodes scapularis in Quebec (Canada). Ticks and Tick Borne Diseases. Jan;13(1):101833. doi: 10.1016/j.ttbdis.2021.101833. Epub 2021 Sep 23. PMID: 34600416.

469
470 Ecological Forecasting Initiative (EFI). 2019. EFI 2019 Conference May 13-15, 2019. Available at
471 <https://ecoforecast.org/efi2019/> (accessed 15 November 2021)
472
473 Ecological Forecasting Initiative (EFI) and National Ecological Observation Network (NEON). 2020.
474 NSF-sponsored EFI-Research Coordination Network and NEON. Dates: May 12-13, 2020. Available at
475 <https://ecoforecast.org/efi-rcn-2020-conference/> (accessed 15 November 2021)
476
477 Ecological Forecasting Initiative (EFI) Research Coordination Network (RCN) and National Ecological
478 Observation Network (NEON). 2020. EFI RCN NEON Ecological Forecast Challenge. Available at
479 <https://ecoforecast.org/efi-rcn-forecast-challenges/> (accessed 15 November 2021)
480
481 Egizi A, Roegner V, Faraji A, Healy SP, Schulze TL, Jordan RA. 2018. A historical snapshot of Ixodes
482 scapularis-borne pathogens in New Jersey ticks reflects a changing disease landscape. Ticks and Tick-
483 borne Diseases 9 318-426.
484
485 Eisen L. 2020. Stemming the Rising Tide of Human-Biting Ticks and Tickborne Diseases, United States.
486 Emerging Infectious Disease 26(4) 641-647. doi: 10.3201/eid2604.191629
487
488 Eisen RJ and Eisen L. 2018. The blacklegged tick, Ixodes scapularis: an increasing public health concern.
489 Trends in Parasitology 34 (4): 295-309. doi: 10.1016/j.pt.2017.12.006
490
491 Eisen RJ and Paddock CD. 2021. Tick and tickborne pathogen surveillance as a public health tool in the
492 United States. Journal of Medical Entomology 58(4): 1490-1502.
493
494 Eisen RJ, Eisen L, Beard CB. 2016. County-Scale Distribution of Ixodes scapularis and Ixodes pacificus
495 (Acari: Ixodidae) in the Continental United States. J Med Entomol. 2016;53(2):349-386.
496 doi:10.1093/jme/tjv237
497
498 Eisen RJ, Kugeler KJ, Eisen L, Beard CB, Paddock CD. 2017. Tick-Borne Zoonoses in the United States:
499 Persistent and Emerging Threats to Human Health. ILAR J. 2017;58(3):319-335. doi:10.1093/ilar/ilx005
500
501 Estrada-Peña A, Cevidanes A, Sprong H, Millán J. 2021. Pitfalls in Tick and Tick-Borne Pathogens
502 Research - Some Recommendations and a Call for Data Sharing. Pathogens. 2021;10(6):712.
503 doi:10.3390/pathogens10060712
504
505 Estrada-Peña A, Gray JS, Kahl O, Lane RS, Nijhof AM. 2013. Research on the ecology of ticks and tick-
506 borne pathogens--methodological principles and caveats. Front Cell Infect Microbiol. 2013;3:29.
507 doi:10.3389/fcimb.2013.00029
508
509 Falco RC, Fish D. 1992. A comparison of methods for sampling the deer tick, Ixodes dammini, in a Lyme
510 disease endemic area. Experimental & applied acarology, 14(2), 165–173.
511 https://doi.org/10.1007/BF01219108
512
513 Falco RC, McKenna DF, Daniels TJ, Nadelman RB, Nowakowski J, Fish D, Wormser GP. 1999.
514 Temporal relation between Ixodes scapularis abundance and risk for Lyme disease associated with
515 erythema migrans. Am J Epidemiol. 15;149(8):771-6. doi: 10.1093/oxfordjournals.aje.a009886. PMID:
516 10206627.
517
518 Fleshman A, Graham CB, Maes SE, Foster E, Eisen RJ. 2021. Reported county-level distribution of
519 Lyme disease spirochetes, Borrelia burgdorferi sensu stricto and Borrelia mayonii (Spirochaetales:

520   Spirochaetaceae), in Host-Seeking Ixodes scapularis and Ixodes pacificus Ticks (Acari: Ixodidae) in the
521   Contiguous United States. Journal of Medical Entomology 58 (3) 2021: 1219 - 1233.
522
523   Fournier AM, White ER, Heard SB. 2019. Site-selection bias and apparent population declines in long-
524   term studies. Conserv. Biol. 33, 1–10. doi: 10.1111/cobi.13371
525
526   Gray JS, Kahl O, Janetzki C, Stein J. 1992. Studies on the ecology of Lyme disease in a deer forest in
527   County Galway, Ireland. J Med Entomol. Nov;29(6):915-20. doi: 10.1093/jmedent/29.6.915. PMID:
528   1460628. https://doi.org/10.1093/jmedent/29.6.915
529
530   Hill, C. A., Whitford, F., and MacDonald, J. F. 2009. Assessing and responding to public health
531   entomology needs in Indiana. American Entomol. 55: 114-121
532
533   Kahl O., Gern L., Eisen L., Lane R. S. (2002). Ecological research on Borrelia burgdorferi sensu lato:
534   terminology and some methodological pitfalls, in Lyme Borreliosis: Biology, Epidemiology and Control,
535   eds Gray J. S., Kahl O., Lane R. S., Stanek G. (New York, NY: CABI Publishing;), 29–46.
536
537   Kjellander, P.L., Aronsson, M., Bergvall, U.A. et al. 2021. Validating a common tick survey method:
538   cloth-dragging and line transects. Experimental and Applied Acarology 83, 131–146.
539   https://doi.org/10.1007/s10493-020-00565-4
540
541   Kugeler KJ, Eisen RJ. 2020. Challenges in Predicting Lyme Disease Risk. JAMA Network Open.
542   2020;3(3):e200328. doi:10.1001/jamanetworkopen.2020.0328
543
544   Kugeler KJ, Schwartz AM, Delorey MJ, Mead PS, Hinckley AF. 2021. Estimating the Frequency of
545   Lyme Disease Diagnoses, United States, 2010–2018. Emerg Infect Dis. 27(2):616-619.
546   https://doi.org/10.3201/eid2702.202731
547
548   Lee X, Hardy K, Johnson DP, Paskewitz SM. 2013. Hunter-killed deer surveillance to assess changes in
549   the prevalence and distribution of Ixodes scapularis (Acari: Ixodidae) in Wisconsin. Journal of Medical
550   Entomology 50(3): 632-639.
551
552   Mathews-Martin L, Namèche M, Vourc'h G, Gasser S, Lebert I, Poux V, Barry S, Bord S, Jachacz J,
553   Chalvet-Monfray K, Bourdoiseau G, Pamies S, Sepulveda D, Chambon-Rouvier S, Rene-Martellet M.
554   2020. Questing tick abundance in urban and peri-urban parks in the French city of Lyon. Parasites
555   Vectors 13, 576. https://doi.org/10.1186/s13071-020-04451-1
556
557   Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group. 2009. Preferred Reporting Items for
558   Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097.
559   doi:10.1371/journal.pmed1000097
560
561   National Science Foundation (NSF). 2019. NEON RCN: The Ecological Forecasting Initiative RCN:
562   Using NEON-enabled near-term forecasting to synthesize our understanding of predictability across
563   ecological systems and scales. NSF Division of Environmental Biology Award Abstract #1926388.
564   Available at <https://www.nsf.gov/awardsearch/showAward?AWD_ID=1926388> (accessed 15
565   November 2021)
566
567   Newman BC, Sutton WB, Wang Y, Schweitzer CJ, Moncayo AC, Miller BT. 2019. A standardized
568   method for the construction of a tick drag/flag sampling approach and evaluation of sampling efficacy.
569   Exp Appl Acarol 79, 433–446 2019. https://doi.org/10.1007/s10493-019-00429-6
570

Paddock CD, Lane RS, Staples JE, Labruna MB. 2016. Changing paradigms for tick-borne diseases in the
Americas. In: Mack A, editor. Global health impacts of vector-borne diseases: workshop summary.
National Academies Press; 2016. pp. 221–257. URL:
https://nap.nationalacademies.org/login.php?record_id=21792&page=https%3A%2F%2Fnap.nationalaca
demies.org%2Fdownload%2F21792 (accessed 25 May 2022)

Pool JR, Petronglo JR, Falco RC, Daniels TJ. 2017. Energy usage of known-age blacklegged ticks (Acari:
Ixodidae): What is the best method for determining physiological age? Journal of Medical Entomology
2017;54:949–956. doi: 10.1093/jme/tjx042.

R Core Team. 2020. R: A language and environment for statistical   computing. R Foundation for
Statistical Computing, Vienna, Austria. URL http://www.R-project.org/. (accessed 15 November 2021)

RStudio Team. 2020. RStudio: Integrated Development for R. RStudio, PBC, Boston, MA URL
http://www.rstudio.com/. (accessed 15 November 2021)

Ripoche M, Gasmi S, Adam-Poupart A, Koffi JK, Lindsay LR, Ludwig A, Milord F, Ogden NH,
Thivierge K, Leighton PA. 2018. Passive Tick Surveillance Provides an Accurate Early Signal of
Emerging Lyme Disease Risk and Human Cases in Southern Canada. J Med Entomol. 2018 Jun
28;55(4):1016-1026. doi: 10.1093/jme/tjy030.

Rulison EL, Kuczaj I, Pang G, Hickling GJ, Tsao JI, Ginsberg HS. 2013. Flagging versus dragging as
sampling methods for nymphalIxodes scapularis(Acari: Ixodidae). Journal of Vector Ecology, 38(1), 163-
167. doi:10.1111/j.1948-7134.2013.12022.x

Salomon J, Hamer SA, Swei A. 2020. A Beginner's Guide to Collecting Questing Hard Ticks (Acari:
Ixodidae): A Standardized Tick Dragging Protocol. Journal of Insect Science Volume 20, Issue 6, 11.
https://doi.org/10.1093/jisesa/ieaa073

Sougoufara S, Ottih EC, Tripet, F. 2020. The need for new vector control approaches targeting outdoor
biting anopheline malaria vector communities. Parasites Vectors 13, 295. https://doi.org/10.1186/s13071-
020-04170-7

Townson H, Nathan MB, Zaim M, Guillet P, Manga L, Bos R, Kindhauser M. 2005. Exploiting the
potential of vector control for disease prevention. Bulletin of the World Health Organization 83(12):942-
7. Epub 2006 Jan 30. PMID: 16462987; PMCID: PMC2626501.

United States Department of Health and Human Services. 2020. "Tick-Borne Disease Working Group
2020 Report to Congress 2nd report." URL https://www.hhs.gov/sites/default/files/tbdwg-2020-report_to-
ongress-final.pdf (accessed 27 May 2022)

Wauchope HS, Amano T, Sutherland WJ, Johnston A. 2019. When can we trust population trends? A
method for quantifying the effects of sampling interval and duration. Methods in Ecology and Evolution
10: 2067–2078. doi:10.1111/2041-210X.13302

White ER. 2019. Minimum Time Required to Detect Population Trends: The Need for Long-Term
Monitoring Programs, BioScience, Volume 69, Issue 1, January 2019, Pages 40–46,
https://doi.org/10.1093/biosci/biy144

White ER, Bahlai CA. 2021. Experimenting With the Past to Improve Environmental Monitoring.
Frontiers in Ecology and Evolution 8 (391). DOI=10.3389/fevo.2020.572979

Supplemental Materials
The data for this study are located here -
Rowan Christie, Kaitlin Stack Whitney, Julia Perrone, & Christie Bahlai. (2022). Ixodes
Scapularis monitoring data compiled from 7 studies [Data set]. Zenodo.
https://doi.org/10.5281/zenodo.6540831
627
628 The code for this study is located here:
629 Rowan Christie. (2022). SMCCoder/ixodes_scapularis_research: Ixodes scapularis research code
630 for PeerJ submission (v1.1). Zenodo. https://doi.org/10.5281/zenodo.6820316

631 **Appendix 1. PRISMA workflow for finding and screening black-legged tick studies**
632
633 Screening criteria: We searched for datasets that were at least 9 years and measured black-legged
634 tick density or abundance using any sampling method, documenting the process and results using
635 the PRISMA flow diagram (Moher et al., 2009).
636
637

Date Accessed	Database	Search terms	# of results	# of results without duplicates	# of results after screening
6/26/2019	LTER data portal	scapularis, tick	12		1
6/26/2019	Data One	scapularis, tick	19	18	2
6/26/2019	Data Dryad repository	scapularis, tick	9	7	0
6/26/2019	NEON	tick	2	2	0
6/26/2019	Google Datasets repository	scapularis	75	70	4
6/28/2019	NJ Department of Health	Accessed vector borne surveillance reports	0		0
6/28/2019	CT Department of Health	tick	2,920		1
6/28/2019	PA Department of Health	tick	54		0
6/28/2019	Google	pennsylvania tick datasets	447,000		1
6/28/2019	DE Department of Health	tick	61		0
6/28/2019	Google	delaware tick datasets	18		0
6/28/2019	RI Department of Health	tick	23		0

Tables

Dataset name	Data repository	Location	Geographic scope	Life stage	Sampling technique
Summaries of Tick Testing	The Connecticut Agricultural Experiment Station	Connecticut, US	Town	Unspecified	Found on a person
Incidence of Ticks and Tick Bites at Harvard Forest since 2006	LTER Data Portal	Harvard Forest, Massachusetts, US	State forest	Unspecified	Found on a person
The growing importance of lone star ticks in a Lyme disease endemic county: Passive tick surveillance in Monmouth County, NJ, 2006 – 2016	NCBI	Monmouth County, New Jersey, US	Town	Adults, nymphs	Found on a person
Deer Tick Surveillance: Adults (Oct to Dec) excluding Powassan virus: Beginning 2008	NY Department of Health	New York, US	County	Adults	Dragging
Deer Tick Surveillance: Nymphs (May to Sept) excluding Powassan virus: Beginning 2008	NY Department of Health	New York, US	County	Nymphs	Dragging
Tick-borne disease risk in a forest food web	Data Dryad	Cary Forest, New York, US	Grid	Adults, larvae, nymphs	Dragging
Range expansion and increasing Borrelia burgdorferi infection of the tick Ixodes scapularis (Acari: Ixodidae) in Iowa, 1990-2013	Data Dryad	Iowa, US	County	Adults, nymphs	Found on a person

Table 1. Location, geographic scope, life stage, and sampling technique of black-legged deer tick studies found that met our study criteria (Appendix 1). Datasets from these studies were included in our analyses.

Figures

Figure 1. The fraction of datasets that take *y* years to reach stability.

Number of years for datasets within our study to reach stability. Majority of datasets (n = 289) required 10 years to reach stability, and none reached stability before 4 years.

Figure 2. Comparison of study length, years to reach stability and the number of datasets.

Scatter plot comparing study length with the number of years to reach stability. Point size represents the number of raw datasets (n=289). Overall, the number of years to reach stability increased with study length.

Figure 3. Overall proportion wrong compared to proportion wrong before stability.

Whisker plot comparing the overall proportion wrong and proportion wrong before reaching stability and showing no significant differences (t = -1.2, df = 571.46, p-value = 0.23). Each dot represents the results from a dataset (n=289) analyzed.

Figure 4. Comparison of stability time and proportion significantly wrong before stability by sampling techniques.

A) Years to reach stability by sampling technique, dragging (n=90) and public survey (n=198). Letters above boxes denote significant differences (t=-8.53, df=236.23, p-value < 0.001).

B) Proportion significantly wrong before reaching stability across sampling technique, showing no significant difference between techniques (t=0.08, df = 155.58, p-value = 0.93).

Figure 5. Comparison of stability time and proportion significantly wrong before stability by life stage.

A) Years to reach stability by life stage, larvae (n=8), nymphs (n=68), and adult (n=63). Letters above boxes denote significant differences between larvae and adults (t = -5.96, df = 10.11, p-value <0.001) and larvae and nymphs (t = -5.56, df = 10.33, p-value <0.001), while there is no significant difference between nymphs and adults (t = -0.63, df = 128.99, p-value = 0.53). B) Proportion significantly wrong before reaching stability across life stage. Letters above boxes denote significant differences between nymphs and adults (t = 2.99, df = 112.55, p-value = 0.003). There is no significant differences between nymphs and larvae (t = -0.78, df = 7.85, p-value = 0.46) or adults and larvae (t = 0.44, df = 8.77, p-value = 0.67).

Figure 6. Comparison of stability time and proportion significantly wrong before stability by geographic scope.

A) Years to reach stability by geographic scope, grid (n=24), state forest (n=6), town (n=186) and county (n=73). Letters above boxes denote significant differences between county and town level studies (t = -17.03, df = 243.09, p-value <0.001); county and state forest level studies (t = -5.55, df = 5.22, p-value = 0.002); county and grid level studies (t = -17.21, df = 33.87, p-value <0.001). There is no significant between town and state forest level studies (t = 0.47, df = 6.12, p-value = 0.65); town and grid level studies (t = 0.22, df = 86.18, p-value = 0.8); state forest and grid level studies (t = -0.38, df = 6.03, p-value = 0.72). B) Proportion significantly wrong before reaching stability across geographic scope. Letters above boxes denote significant differences between state forest and county level studies (t = 5.24, df = 74.29, p-value <0.001); state forest and town level studies (t = 7.42, df = 46.44, p-value <0.001). There is no significant difference between county and town level studies (t = 0.72, df = 115.03, p-value =

0.48); county and grid level studies (t = 1.33, df = 43.07, p-value = 0.19); town and grid level studies (t = 1.00, df = 28.52, p-value = 0.32); state forest and grid level studies (t = -1.91, df = 25.10, p-value = 0.07).

Figure 7. Comparison of stability time and proportion significantly wrong before stability by study response variable.

A) Years to reach stability by study response variable, deer tick abundance (n=175) and *B. burgdorferi* presence in deer ticks (n=114) and showing no significant difference between study response variables (t = -1.29, df = 283.9, p-value = 0.20). B) Proportion significantly wrong before reaching stability across study response variables and showing no significant difference between study response variables (t = -1.18, df = 232.98, p-value = 0.24).