Mito-nuclear variation

in transgenerational and reproductive effects

of amino acid and lipid nutrition

Adam J. Dobson[a,b]*, Luisa Kumpitsch[b], Lucas Langer[b],

Emmely Voigt[b], Damian K. Dowling[c], Klaus Reinhardt[b]*

a - Institute of Molecular Cell & Systems Biology, College of Medical Veterinary &

Life Sciences, Davidson Building, University of Glasgow, G12 8QQ, UK.

b - Applied Zoology, Faculty of Biology, Technische Universität Dresden, 01069

Dresden, Germany.

c - School of Biological Sciences, Monash University, Clayton, Victoria 3800,

Australia.

* correspondence: adam.dobson@glasgow.ac.uk, klaus.reinhardt@tu-dresden.de

Abstract

Animals vary genetically in responses to dietary change. Both mitochondrial and nuclear genomes contribute to this variation, but the role of combinatorial "mito-nuclear" genetic variation is understudied. We do not know whether specific nutrients modify patterns of mito-nuclear variation, nor whether putative epigenetic mechanisms play a role. Here, we show that enriching dietary essential amino acids or lipids modifies patterns of mito-nuclear variation in *Drosophila* life-history, including transgenerational effects of lipids. Systematically evaluating alternative statistical models revealed that diet-mito-nuclear interactions were a leading driver of phenotypic variation. Mito-nuclear genotype repeatably predicted phenotypic impacts of nutritional changes, but genotypes bearing naturally co-occurring pairs of mitochondria and nuclei did not necessarily outperform novel pairings, suggesting that nutrition-dependent phenotypes cannot easily be optimised by matching mitochondria to coincident nuclear genotypes. These results enhance understanding of how nutrition and genetics sculpt phenotype, with potential implications for human mitochondrial transfer therapies.

  Introduction

Nutrients and a genetic code are the fundaments of cellular life. Consequently,

variation in nutrition and genotype are primary determinants of variation in health and

biological fitness. Nutritional and genetic variation can also interact, meaning that

different genotypes can respond distinctly to the same nutritional change (Dobson et

al., 2015; Heymsfield and Wadden, 2017; Jumbo-Lucioni et al., 2010; Liao et al.,

2009; Mulvey et al., 2017; Rikke et al., 2010). Biomedically, the interactive effects of

diet and human genetic variation have led to interest in personalised nutrition

(Ordovas et al., 2018), in which diets would be targeted individually, to treat maladies

such as obesity and late-life morbidity (Fontana et al., 2010; Heymsfield and

Wadden, 2017; Simpson and Raubenheimer, 2012). Biologically, unravelling genetic

variation in how fitness is affected by nutrition will be a major part of bridging the

genotype-phenotype gap.

Why do responses to diet vary? Ultimately, cellular metabolic function is the output of

the coordinate activity of enzymes encoded by both the mitochondrial and nuclear

genome. Both genomes exhibit sequence variation, yet the consequences of

mitochondrial genome variation have often been overlooked. Effects of diet do

indeed vary amongst mitochondrial haplotypes, altering metabolism and life-history

(Aw et al., 2018; Bevers et al., 2019; Camus et al., 2020; Drummond et al., 2019;

Mulvey et al., 2017; Nagarajan-Radha et al., 2019; Towarnicki and Ballard, 2018,

2017; Zhu et al., 2014). When both mitochondrial and nuclear genomes vary, we

expect "mito-nuclear" variation, in which effects of variants in one can be either

buffered or augmented by variants in the other. In animals, this has been observed

for fecundity (Mossman et al., 2016a), fertility (Camus et al., 2015), lifespan (Camus

et al., 2015; Vaught et al., 2020), gene expression (Camus et al., 2015; Mossman et

al., 2019, 2016b), and epigenomic modifications (Grunau et al., 2018). Novel mito-

nuclear pairings can even change the course of nuclear genomic evolution (Healy

and Burton, 2020), and lead to reproductive isolation (Hill, 2019; Ma et al., 2016).

The intimacy of mito-nuclear coupling leads us to expect coadaptation (Dowling et

al., 2008), and consequently to predict that pairs from the same population

(sympatric) should function optimally. Some studies have supported this prediction

(Baris et al., 2017; Chang et al., 2016; Ellison and Burton, 2006; Ma et al., 2016;

Meiklejohn et al., 2013), but others have not (Dowling et al., 2010; Mossman et al.,

2016b; Vaught et al., 2020). Despite genetic variation in responses to dietary

variation, and the emerging importance of mito-nuclear interactions in overall genetic

variation, there has been relatively little work to connect the three, by asking whether

the phenotypic impacts of dietary alterations are subject to mito-nuclear interactions.

Trailblazing work in *Drosophila melanogaster* has shown diet-mito-nuclear (DMN)

variation in development time, lifespan and fecundity (Camus et al., 2020; Montooth

et al., 2019; Mossman et al., 2016a; Rand et al., 2018; Zhu et al., 2014), but specific

causal nutrients and cellular mechanisms remain to be established.

Perhaps diet's most immediate fitness effect is regulation of reproduction. Parents

can gain fitness by increasing offspring quantity, or by promoting offspring quality

(Lack, 1947; Smith et al., 1989; Stearns, 1992). Offspring quality and quantity will

depend on fertility, fecundity, and development (viability) of those offspring, each of

which are influenced by diet. Equivalent changes in these traits can be elicited in

distinct taxa by equivalent dietary changes: for example, in numerous studies of

insects and mammals, amino acid nutrition has proven consistently important for

fertility, fecundity and offspring development (Bong et al., 2014; Crean and Senior,

2019; Dong et al., 2016; Fanson et al., 2012; Grandison et al., 2009; Jensen et al.,

2015; Lee et al., 2008; Leitão-Gonçalves et al., 2017; Liang and Zhang, 2006; Ma et

al., 2020; Maklakov et al., 2008; McCracken et al., 2020; Mirth et al., 2018; Skorupa

et al., 2008; Solon-Biet et al., 2015; Svajgr et al., 1972; Tufarelli et al., 2015; Winship

et al., 2018; Wong et al., 2014a; Zanco et al., 2020); whereas high-lipid or high-sugar

diets are obesogenic (Dobson et al., 2019; Hariri and Thibault, 2010; Jang et al.,

2018; Rivera et al., 2015; Saud et al., 2015; Wong et al., 2014b; Woodcock et al.,

2015; Zhao et al., 2020). However, importantly, reproduction is not only regulated by

the individual's current diet: large nutritional fluctuations can induce enduring

physiological and molecular changes, which can transmit across generations,

influencing quality of offspring and even grand-offspring (Barker and Osmond, 1986;

Barker and Thornburg, 2013; Deas et al., 2019; Duque-Guimarães and Ozanne,

2017; Holland et al., 2016; Li et al., 2018; Liang and Zhang, 2006; Öst et al., 2014;

Rivera et al., 2015; Stefana et al., 2017; Wei et al., 2014; Winship et al., 2018). Thus,

diet can influence reproduction via lasting, programmed effects on offspring quality,

which point to epigenetic reprogramming, whilst acute effects of diet on both

offspring quality and quantity point to metabolic changes. But do these effects vary

amongst genotypes? If so, is variation shaped by mitochondrial genomes, nuclear

genomes, or by mito-nuclear genotype?

Understanding health impacts of mito-nuclear interactions is not just a current topic

in biology. It is also pertinent biomedically, because dietary intervention is a

preeminent treatment for human mitochondrial disease (Gorman et al., 2016), and

also following the advent of mitochondrial replacement therapies (Craven et al.,

2010; Tachibana et al., 2018; Wolf et al., 2015). These procedures have been

licensed in the United Kingdom and a few other countries (Cohen et al., 2020). They

aim to treat mitochondrial diseases by placing embryonic nuclei into the cytoplasm,

and therefore mitochondrial context, of a disease-free donor. Some argue that the

conservation of mitochondrial function and the body of evidence for mito-nuclear

interactions suggest that undesired consequences may ensue if the donor

mitochondria function poorly with the nuclear genome (Dobler et al., 2018; Dunham-

Snary and Ballinger, 2015; Reinhardt et al., 2013); although the UK Human Fertility

and Embryology Authority concluded that risks remain "purely theoretical" (HFEA,

2016). To solve this debate it will be important to identify which traits to evaluate in

patients, whether costs or benefits may be diet-dependent, and whether

transgenerational variation may ensue.

Here, we identify specific nutrients sufficient to drive diet-mito-nuclear variation in

Drosophila life-history, and show that specific mito-nuclear genotypes can exhibit

long-lasting, transgenerational responses to transient dietary changes. We bred a

panel of flies with replicated and fully-factorial mito-nuclear variation, and show that

essential amino acids and lipids can elicit novel patterns of variation in fecundity and

offspring performance (development time). We then show for the first time that the

capacity for effects of parental diet on offspring performance are genetically variable,

and mito-nuclear in this case. To elucidate overall variation in fitness, we integrate

our analyses of offspring quality and quantity, showing mito-nuclear variation in how

diet influences reproductive investment; and also that "mis-matched" pairs of

mitochondrial and nuclear genomes do not necessarily perform worse than matched

pairs. Finally, we evaluate performance of alternative models of our extensive

datasets, in which interactions between diet, mitochondrial and nuclear haplotype

were systematically included or excluded. We find that models which include mito-

nuclear or diet-mito-nuclear interactions are favoured overwhelmingly over models

which assume no interaction, suggesting that models which do not account for these

terms are not adequate to explain variance in fitness and health.

 Results

We used a set of *D. melanogaster* lines comprising replicated, fully-factorial

combinations of varied mitochondrial and nuclear genomes (Figure 1a), derived from

heterogeneous populations originally isolated in Australia and Benin (formerly

Dahomey). For brevity, we term Australian and Beninese genotypes *A* and *B*,

respectively. The populations were introgressed either reciprocally (*AB* = *A*

mitochondria / *B* nuclei; *BA* = *B* mitochondria / *A* nuclei), or to themselves to control

for introgression and drift (i.e. *AA* = *A* mitochondria / *A* nuclei; *BB* = *B* mitochondria /

B nuclei). In each generation, a fixed number of females per line were mated to a

fixed number of males from the ancestral population. Each combination was

replicated in triplicate, generating twelve lines altogether (e.g. *AA1, AA2, AA3, BA1,*

BA2, etc). The lines were created to retain equivalent standing nuclear variation from

the ancestral populations in each of the two mitochondrial backgrounds.

Introgressions had been iterated for 74 generations at the time of our first

experiment, and so the percentage of nuclear genome that remained unreplaced by

introgression in the *AB* and *BA* genotypes was expected to be negligibly small

(~5.3e-21%). Ancestors of all lines were cleared of cytoplasmic *Wolbachia*

endosymbionts long prior to these experiments (Vaught et al., 2020), and so we

expect cytoplasmic genetic complements to comprise only mitochondria. The *A* and

B nuclear genomes do not appear to select for differential mitochondrial haplotypes

in this design (Vaught et al., 2020), and we expect the rigorous backcrossing regime

to minimize differential segregation of nuclear genomic variants over *A* and *B*

mitochondria: this experimental design should therefore produce fully-factorial mito-

nuclear variation.

Figure 1

Figure 1. Study design. A. Genetically distinct and heterogeneous populations of

flies were originally isolated from from Benin (formerly Dahomey), and New South

Wales, Australia. Mito-nuclear variation was generated by reciprocally introgressing

each line either into its own respective cytoplasmic background, or into the other

line's cytoplasmic background. Three biologically independent replicate lines were

established per mito-nuclear combination. 74 iterations of introgression were

completed before experiments. **B.** Fly lines were cultured on rearing diet and then

fed experimental diets in adulthood. The heatmap shows estimated content of

different macronutrients, bars at the top indicating caloric content.

EAA and lipid nutrition are sufficient to drive diet-mito-nuclear variation in life-history

To identify nutrients which could drive DMN interactions, we added nutrients to a sugar-yeast diet (Bass et al., 2007), as previously (Dobson et al., 2018; Emran et al., 2014; Grandison et al., 2009). To ensure that potential novelty effects were evenly distributed amongst conditions, the baseline diet was distinct from the medium on which the lines were maintained and developed. For experimental diets, we added one of two nutrients to the baseline, selected for their important and conserved health impacts. Essential amino acids (EAAs) are metabolised by mitochondria (Mariño et al., 2014), and their specific enrichment is sufficient to recapitulate fecundity effects of adding yeast (Emran et al., 2014; Grandison et al., 2009). We also manipulated dietary lipid, because mitochondria metabolise fatty acids, and high-fat animal diets can model Western human disease (Hariri and Thibault, 2010; Heymsfield and Wadden, 2017). Lipid source can have large physiological effects (Brankatschk et al., 2018), and flies are largely vegan in nature (Knittelfelder et al., 2020). Therefore, we supplied a plant-based lipid which set in agar (i.e. in contrast to oils), by adding margarine to the diet (15% w/v, after (Woodcock et al., 2015)). Whilst this design did not control calories, it enabled us to test whether specific, ecologically-relevant nutrients can cause diet-mito-nuclear variation. Estimated nutrient contents (Lesperance and Broderick, 2020) are presented in Figure 1b.

Previous work showed that manipulating complex nutrient sources elicits life-history variation in specific mito-nuclear genotypes, including in fecundity (egg laying), and

offspring development (Camus et al., 2020; Montooth et al., 2019; Mossman et al.,

2016a; Rand et al., 2018; Zhu et al., 2014). We tested whether EAAs and lipid were

sufficient to cause these effects. Starting three days after adult emergence, the

control, EAA-enriched and lipid-enriched foods were fed to flies of all lines for seven

days. We restricted treatment to adults, to avoid developmental variation (Figure 2a).

This experiment was repeated twice. In both experiments, EAAs increased fecundity,

and lipids repressed fecundity. Fecundity was correlated between the two

experiments, indicating technical repeatability (Figure 2b). Biologically, two important

trends were evident. First, genetic replicates consistently grouped together, with all

three replicates of each mito-nuclear genotype tending to sit within one standard

deviation of each other (Figure S1), despite having been separated for >70

generations of introgression. This suggested a highly stereotyped effect of mito-

nuclear genotype on fecundity, which overcame any genetic drift in the fly lines.

Therefore, for simplification, we replotted the data without distinguishing

experimental or genetic replicates (Figure 2c). This outlined the second important

trend, that mito-nuclear variation was shaped by diet, in agreement with a GLMM

analysis indicating a three-way interaction (p=0.009, Table S1). To visualise pairwise

statistical differences between every genotype, we used *post-hoc* estimated marginal

means (EMMs (Searle et al., 1980)) analysis. These EMMs represented the GLMM

output per experimental condition, and so their correlation to the raw data indicated a

good model fit (Figure 2c). Pairwise comparison of EMMs was consistent with our

interpretation that overall patterns of among-genotype variation in fecundity were

diet-dependent (Table S2). These results revealed distinct patterns of mito-nuclear

variation on different diets, and that impacts of consuming a certain diet depend on

mito-nuclear genotype.

Animal fitness is a function of both quantity and quality of offspring, and DMN

variation in offspring development has already been documented. Having shown that

EAAs and lipid were sufficient to cause DMN variation in egg laying, we

hypothesised that variation in offspring quality may additionally manifest in fertility of

those eggs. To test this, after parents had been fed experimental diets for a week as

previously, we allowed eggs laid over 24h to develop in the same experimental

media. We scored time to adult and pupal emergence, along with sex in adults. We

take proportion of eggs emerging as a metric of parental reproductive success, and

time to development as a metric of relative offspring fitness. In every mito-nuclear

genotype, food enrichment with EAAs or lipid either reduced the proportion

developing, extended development time, or both. However, the magnitude of these

effects was highly genotype-dependent (Figure 2d). For explicit analysis, we fit Cox

mixed-effects (CME) survival models. The DMN interaction was statistically

significant for emergence of both males and females to adulthood (each sex

$p < 0.0005$, Table S3, Figure S2). We did not detect robust sex-specificity

(Supplementary Text, Figure S2), and results were broadly repeatable within the

triplicates of each fly genotype (Figure S2). To confirm genotype-specific effects, we

applied *post-hoc* EMM tests to the CME model. This analysis aggregated time to

development and proportion developing into a single quantitative output, because

the EMMs were a posterior view of our CME analysis, which integrates time to

development and proportion developing. In this framework, high EMMs correspond

to faster and/or more frequent development, and vice-versa (Figure 2e). These

EMMs confirmed that EAA or lipid enrichment disadvantaged specific genotypes

(Figure 2e). The *AA* flies bore a particular cost of reduced survival on EAA food. The

differential ranking of EMMs across genotypes on different diets indicated that the

pattern of mito-nuclear genetic variation was dependent on EAA and lipid nutrition.

Indeed, as with fecundity, there was little variation within each triplicate per mito-

nuclear genotype (except that the detrimental effect of EAAs on development of *AA*

flies was pushed to lethality in the *AA3* line, and *BB3* was somewhat more lipid-

resistant than *BB1* and *BB2* (Figure S2)), suggesting that the developmental

response to nutrition is highly stereotyped by mito-nuclear genotype. A parallel

analysis of these same individuals' time to pupation revealed congruent patterns

(Figure S3, Figure S4, Table S4). Altogether, these results show that either lipid or

EAA enrichment are sufficient to potently modify developmental outputs of mito-

nuclear genetic variation.

Figure 2

Figure 2. Diet-mito-nuclear interactions modulate *Drosophila* fecundity. A.

Experimental design. The 12 mito-nuclear lines were reared from egg to adult on

rearing food, and allocated at random to experimental media 6-48h after eclosion, at

a density of five of each sex per vial. After seven days, flies laid eggs on fresh food

for 24 hours. **B.** Intra-line correlations in impact of diet-mito-nuclear variation on egg

laying. Each point shows mean egg laying per line per diet in each of two replicate

experiments, with the replicates of each haplotype grouped by dashed lines. Means

were correlated between experiments (Pearson's r = 0.87, p = 7.6e-12). **C.** Impact of

diet on fecundity is determined by mito-nuclear variation. Data are pooled across the

experimental and genetic replicates presented in panel B. Boxplots show medians,

first and third quartiles and 5th and 95th percentiles. Connected points to right of

each box show *post-hoc* comparisons of means (by estimated marginal means with

95% confidence intervals, calculated from a generalised linear mixed effects model,

showing exponent of EMMs in order to fit to original data scale). Translucent points

to left of each box show raw data. Complementary statistical analysis presented in

Table S1 and Table S2. **D.** Kaplan-Meier plots of development (egg-to-adult) after

parents fed and offspring developed on experimental media. Mito-nuclear genotypes

are indicated in top-left of each panel. **E.** Estimated marginal means (EMMs) with

95% confidence intervals summarising survival analysis (Cox mixed effects) of data

from D. Higher values correspond to faster development and/or increased fertility,

lower values correspond to slower development and/or reduced viability.

Accompanying statistical analysis presented in Table S3.

Persistent life-history effects of dietary EAAs and lipid in specific mito-nuclear genotypes

DMN interactions could occur via numerous mechanisms, but no candidate has yet been confirmed. Dietary variation induces not only immediate changes to metabolism, but also long-lasting epigenetic changes (Barker and Osmond, 1986; Barker and Thornburg, 2013; Deas et al., 2019; Dobson et al., 2017; Duque-Guimarães and Ozanne, 2017; Holland et al., 2016; Li et al., 2018; Liang and Zhang, 2006; Öst et al., 2014; Rivera et al., 2015; Stefana et al., 2017; Wei et al., 2014; Winship et al., 2018). Mitochondria metabolise substrates for epigenetic modifications (Eisenberg et al., 2014; Mariño et al., 2014; Trefely et al., 2020), suggesting that nutritional regulation of epigenetically-encoded traits may vary amongst mito-nuclear genotypes. We asked whether EAAs and lipid might drive DMN interactions for life-history via long-term reprogramming. Whilst such effects are consistent with epigenetic modifications, we conservatively avoid that term for our data because we did not measure epigenetic marks directly. We tested for long-lasting effects of transient EAA and lipid enrichment, both in the generation that experienced the nutritional manipulation, and in their offspring.

We conducted a second fecundity experiment, with a week of adult-onset feeding on experimental diets as previously, but followed by a diet switch back to standardised rearing medium. Fecundity was measured at the end of experimental diet feeding, and 24h after switching back to standardised medium, yielding vial-matched egg counts pre- and post- switch (Figure 3a). The overall effect of experimental diets was

equivalent before and after the switch (EAAs elevated fecundity, lipid repressed

fecundity), confirming that effects of experimental diets persisted for the 24h feeding

on standardised diet (Figure 3b). Before the switch, on experimental diets, a DMN

interaction was once again evident (Figure 3c, Table S5, Figure S5a), and pooling

these data with preceding data (Figure 2) yielded the same qualitative result (Figure

S6, Table S6), confirming that DMN effects were consistent across experiments.

A GLMM analysis confirmed that, after switching back to standardised diet, the main

effects of prior EAA or lipid feeding persisted (Figure 3d, Table S7, Figure S5b).

However, the DMN interaction did not persist (Figure 3d, Table S7, Figure S5b).

These findings suggested that EAA and lipid modify the manifestation of mito-

nuclear variation in fecundity via transient effects; on top of longer-term impacts

which do not depend on mito-nuclear genotype. If so, recovery of fecundity after diet

switch should exhibit a DMN interaction. We estimated recovery rate as the ratio of

eggs laid per vial after/before the switch, with greater ratios indicating greater

plasticity. Recovery from EAA or lipid feeding was indeed variable amongst mito-

nuclear genotypes (Figure 3e; GLMM p=2.91E-09, Table S8; Figure S5c).

Altogether, these analyses revealed two layers of variation, with genotype-

independent main effects of dietary EAAs and lipid which persist after feeding, and

additional DMN variation whilst feeding on the diets, but not after. This suggested

that EAAs and lipid can have lasting fecundity effects which generalise amongst

mito-nuclear genotypes, but did not exclude the possibility of longer-term,

transgenerational diet-mito-nuclear impacts on offspring.

Diet can modulate fitness by effects that span one or more generations (Camilleri-

Carter et al., 2019; Deas et al., 2019; Öst et al., 2014; Rivera et al., 2015). We are

not aware of a prior demonstration of genetic variation in such transgenerational

reprogramming, nor of whether it is subject specifically to mito-nuclear variation.

Since we had already observed that feeding EAAs and lipid to both parents and

offspring modified the manifestation of mito-nuclear genetic variation in development,

we speculated that some of that variation could be due to differential legacies of

parental diet. If so, we expected that feeding EAAs or lipid to parents alone would be

sufficient to modify patterns of mito-nuclear variation in offspring development, even

when all offspring developed in a standardized medium.

We restricted diet manipulations to adults for one week, before egg-laying on

standardised developmental medium, and allowing those eggs to develop on that

standardised medium. This design differs from the preceding study of developmental

effects (Figure 2d-e), because diet was only manipulated in the parental generation.

Therefore any diet-mito-nuclear effect (or lower-order diet effect) detected in this

experiment is transgenerational. We scored time to adult emergence (along with sex

and pupation), and again, results were consistent within each triplicate fly genotype

(Figure S2), and so data were plotted after pooling all three replicates per genotype.

Plotting these data suggested that, indeed, parental nutrition had genotype-specific

effects on development (Figure 3f). This was confirmed by CME analysis (Table S9,

males p=0.003, female p=0.03), without strong evidence of sex bias (Supplementary

Text, Figure S2). EMM analysis confirmed genotype-specific impacts (Figure 3g):

Specifically, *AA* flies developed on average a full day faster when parents were fed

lipid-enriched food (Figure 3f). A parallel analysis of these same individuals' time to

pupation revealed congruent patterns (Figure S3, Figure S4, Table S10). Whilst

some variation may be due to selection, our experimental design largely excludes

this possibility, because the only nutritional variation that individuals under study

experienced was *in utero* when mothers fed on experimental diets, before post-

embryonic development in standardised medium; and variation in rearing density

was also accounted for statistically. Furthermore, since the major effect we observe

is an acceleration of development in one condition (*AA* flies fed lipid), selection

would have to be otherwise ubiquitous but relaxed specifically for this condition. The

results therefore indicate that specific mito-nuclear genotypes are susceptible to

transgenerational effects of parental nutrition, in this case lipid.

Notably, the transgenerational effects we observed in development did not correlate

patterns when both parents and offspring were fed the diets. Our studies of

transgenerational and intergenerational effects of EAAs and lipid were conducted at

the same time (Figure 2d-e flies developed from eggs plotted in Figure 3c, Figure 3f-

g from eggs in Figure 3d), validating this comparison. Altogether, this discord

suggests that nutrients can induce transgenerational effects in specific mito-nuclear

genotypes, but this reprogramming in combination with ongoing nutrient enrichment

in offspring leads to additional changes to patterns of phenotypic variation.

Figure 3

Figure 3. Performance is determined by interplay of mito-nuclear genotype and persistent intragenerational and transgenerational effects of diet. A. Experimental design. 12 Mito-nuclear lines were reared from egg to adult on rearing food, and allocated at random to experimental media 6-48h after eclosion, at a density of five of each sex per vial. After seven days, flies laid eggs on fresh food for 24 hours. Females were then reassigned back to standardised rearing medium, and eggs were laid for a further 24h. **B.** Correspondence between egg laying rate per vial, before and after switch from experimental diets. Egg laying rates before and after switch across all diets were correlated (Pearson's product-moment correlation, $r=0.65$, $p<2.2e-16$). Line of equivalence shown in grey. **C.** Egg laying rate before dietary switch is determined by mito-nuclear variation (*vis* Fig 2C). Boxplots show medians, first and third quartiles and 5th and 95th percentiles. Connected points to right of boxes show *post-hoc* comparisons of means (by estimated marginal means with 95% confidence intervals, calculated from a generalised linear mixed effects model). Translucent points to left of each box show raw data. Complementary statistical analysis presented in Table S5. **D.** Diet-mito-nuclear interactions do not persist 24h after switch from experimental media. Statistical analysis presented in Table S7. **E.** Mito-nuclear interactions govern recovery of fecundity after switch from experimental media. Statistical analysis presented in Table S8. **F.** Kaplan-Meier plots of development to adulthood of eggs laid on developmental media, after parents had previously fed on experimental media. Mito-nuclear genotypes are indicated in top-left of each panel. **G.** Estimated marginal means (EMMs) with 95% confidence intervals summarising survival analysis (Cox mixed effects) of data from F. Higher values correspond to faster development and/or increased fertility, lower values

415   correspond to slower development and/or reduced viability. Accompanying statistical

416   analysis presented in Table S9.

Mito-nuclear (mis-)matching does not predict differential life-history responses to nutrition

Together, offspring quantity and quality define fitness (Lack, 1947; Stearns, 1992). Our analyses suggested that both fecundity (quantity) and fertility (quality) are regulated by EAAs and lipid, but specific mito-nuclear genotypes are differentially receptive or resistant to these effects. To date, the widely-held prediction of a disadvantage to "mis-matched" mito-nuclear pairings, has received equivocal support (Baris et al., 2017; Chang et al., 2016; Ellison and Burton, 2006; Ma et al., 2016; Meiklejohn et al., 2013). Our datasets provided an opportunity to ask whether this prediction applied in terms of differential responses to specific nutrients; whilst taking an integrative view of fitness, encompassing both offspring quality and quantity, and putatively epigenetic effects. To this end, we integrated outputs of our previous analyses by plotted EMMs from against one another, indicating effects of diet and mito-nuclear genotype on offspring quality and quantity, and the relationship between them. We did this separately for data from parents and offspring both fed experimental diets (flies in Figure 2e, which developed from eggs in Figure 3c), and from parent-restricted feeding (flies in Figure 3g, which developed from eggs in Figure 3d).

First, we examined impacts of EAA feeding. Visual inspection indicated that EAA enrichment enhanced offspring quantity, but compromised offspring quality in flies of all genotypes except *BB* (Figure 4a). This was surprising, because we had assumed

that parents should discriminate diets that maximise their own fitness, and therefore

that parental egg-laying would correlate offspring development indices. Instead, the

apparent tradeoff indicates either that (1) EAAs signal diet quality to parents on

normal diets when yeast is the source of most nutrients (including protein), but our

manipulations rendered this quality signal misleading (Zanco et al., 2020); or that (2)

diet is subject to parent-offspring conflict (Trivers, 1974), with parents maximising

their fitness by producing more eggs on high EAA-diets, despite the cost to individual

offspring.

Were costs of EAAs genotype-specific? The quality-quantity tradeoff induced by EAA

feeding was exaggerated dramatically in *AA* flies, whose eggs developed

exceptionally poorly on EAA-enriched food. However, this cost of EAAs was rescued

in both *BA* and *AB* genotypes, indicating that neither *A* mitochondria nor *A* nuclei

alone were sufficient for *AA*'s diet-dependent phenotype, and therefore that these

effects are mito-nuclear (Figure 4a). This outlined a second surprising finding, in

which a sympatric mito-nuclear pairing endured a higher cost of dietary EAAs than

allopatric pairings (Figure 4a). Thus, unexpected, costly, and definitively mito-nuclear

patterns of variation can emerge on specific diets. However, no clear effects of past

EAA feeding on offspring quality or quantity were evident (Figure 4b). Together,

these results indicated that EAAs interact with mito-nuclear genotype during feeding,

but not thereafter. This suggests that the reproductive DMN variation we observed

upon EAA enrichment is an acute effect of diet, and therefore likely metabolic and

not epigenetic.

Next we examined the impact of lipid enrichment. Lipid mostly compromised both

quality and quantity, but one genotype stood out: *AA* flies were most resistant to the

reduction in offspring quality induced by high-lipid diet, and their development index

on lipid-enriched food was in fact better than on EAA-enriched food (Figure 4a). By

this metric, *AA* flies' performance was substantially better than the *AB* or *BA* flies,

confirming that neither *A* mitochondria nor *A* nuclei alone were sufficient to explain

AA's response to diet. This again suggested that genotypes which exhibit notable

responses to diet do not necessarily bear allopatric combinations of mitochondria

and nuclear genomes. Furthermore, the finding that the *AA* genotype was beneficial

in one condition (EAA) despite costs in another (lipid) reveals that it was not simply

sick as would be expected with a poor-functioning mito-nuclear pairing, but instead

exhibits its own genotype-specific pattern of response to diet.

Finally, we evaluated the impact on offspring of parent-restricted lipid enrichment. All

genotypes bore costs, but impacts were highly genotype-specific. Again, the *AA* flies

stood out. Specifically, *AA* flies bore a cost of parental lipid feeding on offspring

quantity, but appeared to enjoy a compensatory benefit from enhanced offspring

quality. In fact, these flies exhibited the highest offspring quality (development index)

observed in any condition in our study. These benefits were not evident in either the

AB or *BA* flies, confirming again that neither *A* mitochondria nor *A* nuclei alone are

sufficient to explain diet-dependent phenotype: the observed effects are mito-

nuclear.

Altogether, these results confirm diet-mito-nuclear shaping of fitness. They show that

diet can induce phenotypic change in specific mito-nuclear genotypes, with neither *A*

mitochondria nor *A* nuclei accounting for the *AA* flies' differential response to diet.

This suggests that mito-nuclear "matching" is not necessarily optimal. In wild

populations, the cost of EAA susceptibility may be balanced by lipid resistance, in

which case these flies were not sick in all circumstances, but rather exhibit a private

pattern of genotype-specific variation, which is evident only on specific diets.

Figure 4

Figure 4. Mito-nuclear interactions dictate dietary modulation of offspring

quality versus quantity. Panels show relationships per diet and per genotype (data

from Figures 2-3). **A** offspring development index and eggs laid, when both parents

and offspring were fed experimental diets, and **B** development index before switch

and release of egg laying after diet, when parents were switched from experimental

to standard diets. Points show EMMs±95% confidence intervals, and grey lines

indicate grand means.

Diet-mito-nuclear effects cause substantial biological variation

Finally, we sought to test the importance of the DMN interactions that we identified

for overall variation in our dataset, for two reasons. First, because the DMN patterns

that we identified may be of previously-unrecognised importance in nature. Second,

because mitochondrial transfer therapy (Craven et al., 2010; Tachibana et al., 2018;

Wolf et al., 2015) presents the possibility of a medical procedure giving patients

novel combinations of mitochondrial and nuclear genomes, which our results

suggest may lead to unpredictable responses to nutritional variation. Therefore, we

sought effect sizes of DMN terms, complementing our preceding analyses by

outlining whether DMN effects were substantial, or biologically trivial despite

statistical significance.

Since *de facto* effect sizes for interaction terms are not easily interpreted, we used

optimization procedures to compare simple models of the data against those

including DMN interactions (Akaike, 1974). Calculating Akaike Weights reveals

performance amongst a set of models, in terms of the relative probability of being the

best description of the data (Wagenmakers and Farrell, 2004). We applied this

procedure to all our previously-presented models (Figures 2-3, Figure S5), and

simpler alternatives. Cases where this analysis favoured including DMN interactions

argued for the importance of this novel tier of variation in health, which mitochondrial

transfer will influence in ways that we cannot currently predict.

Akaike Weights suggested that models accounting for DMN interactions performed

best described all datasets but one, and that purely additive models (assuming no

interactions) were deficient by comparison (Figure 5).

For fecundity data (Figures 2-3), we fit a structured series of models, systematically

including or eliminating all possible interactions amongst diet, mitochondrial and

nuclear genotype. Akaike Weights favoured including DMN interactions (Figure 5a-

d): for example, a model of data from Figure 2c including DMN interactions was 61.7

times more likely to be the best model than another with no interactions (Figure 5a);

and approximately twice more likely than the next-best, which retained only the

diet:nuclear interaction (Figure 5a). Models including DMN terms were also the best

descriptor of fecundity before switching from experimental media (data from Figure

3c), and the release of fecundity after switching (data from Figure 3e). The exception

to the emergent rule was the case of fecundity after switching from experimental

media (Figure 3d), in which the favoured model included a mito-nuclear term, but not

a DMN term. This exception demonstrated that the most complex models were not

favoured automatically, but therefore encouraging confidence in other cases,

suggesting that DMN interactions do indeed explain substantial variation.

For development data (Figures 2e, Figure 3f, Figure S4-S6), including terms for

offspring sex added an additional order of complexity (i.e. a potential four-way

interaction), rendering an assessment of all possible combinations hard to interpret.

Therefore, we calculated Akaike Weights for models including either only first-degree

terms (a purely additive model, without interaction terms), second-degree terms (all

pairwise combinations), third-degree terms (all three-way interactions), or fourth-

degree terms (all four-way interactions, but only when sex was known, in models of

adult emergence). We did this both for development to pupa (Figure 5e-f) and to

adult (Figure 5g-h), and when diet was manipulated either for both parents and

offspring (Figures 5e and 5g), or solely for parents (Figures 5f and 5h). In each case,

models including all possible interaction terms performed best, often by several

orders of magnitude (note logarithmic scale on figures). This analysis indicated that

DMN interactions exert a substantial influence on developmental variation.

In summary, our modelling indicates that key health and fitness metrics - fecundity

and development - are subject to significant interactions amongst mito-nuclear

genotype and diet, and that those interactions account for significant variation in our

data, which reflects the real-world mosaic of diet-mito-nuclear variation.

Figure 5

Figure 5. Models including interactions amongst mitochondrial and nuclear

genotypes and diet treatment are favoured over alternatives. Barplots indicate

Akaike weights analyses of sets of alternative models of data from Figures 2-3.

Columns to left of each panel indicate terms in each model. Higher values indicate a

better-supported model. The probability that one describes data better than another

can be calculated by the ratio of their respective Akaike Weights. Plots compare

models of data of fecundity of varied mito-nuclear genotypes fed experimental diets

(**A,** data from Figure 2; and **B**, data from Figure 3c), after feeding on experimental

diets (**C,** data from Figure 3d), and the ratio of fecundity before and after switching

from experimental to standardised diets (**D**, data from Figure 3e). Plots E-H compare

models of development time and rate, when models included either no interactions

between predictive terms, or various levels of interactions as indicated, for

development to pupa when experimental diets were fed to either parents and

offspring (**E**, data from Figure S3) or parents alone (**F**, data from Figure S3), and

development to adult when experimental diets were fed to either parents and

offspring (**G**, data from Figure 2) or parents alone (**H**, data from Figure 3).

Discussion

Diet is a major source of biological variation. Evidence is now mounting that mito-

nuclear interactions are also a substantial source of variation, with implications in

areas as diverse as mitochondrial disease, ageing, evolution of sex, and speciation

(Ballard and Melvin, 2010; Đorđević et al., 2017; Gemmell et al., 2004; Gershoni et

al., 2009; Havird et al., 2015; Hill, 2017; Innocenti et al., 2011; Latorre-Pellicer et al.,

2019; Milot et al., 2017; Reinhardt et al., 2013; Wallace and Chalkia, 2013). Our

work complements previous studies showing that diet and mito-nuclear variation

interact to produce complex phenotypes, with diet enhancing or reducing the

consequences of mito-nuclear interactions. We now reveal specific nutrients which

are sufficient to drive these effects. The nutrients we identify are of particular

interest, because EAAs are currently attracting a great deal of attention for their

extensive regulation of swathes of life-history and health traits, whilst lipid

consumption is associated with the pandemic of metabolic disease in humans

(Simpson and Raubenheimer, 2012). We now show that responses these nutrients

are exaggerated in some mito-nuclear genotypes, and buffered in others. An

additional, and perhaps more important novel finding, is that transgenerational

effects of these nutrients can emerge in specific mito-nuclear genotypes. These

interactions appear to explain substantial variance in reproductive traits and life-

history. The Darwinian view that reproduction subjugates all other processes

suggests that many further traits may be similarly regulated.

Transient dietary alterations and metabolic disease can drive persistent molecular

and phenotypic change, within and across generations (Dobson et al., 2017; Duque-

Guimarães and Ozanne, 2017; Holland et al., 2016; Stefana et al., 2017). Our study

now shows that specific mito-nuclear genotypes are susceptible to a

transgenerational effect of lipid nutrition. To our knowledge, this is the first

demonstration of any form of genetic variation in transgenerational effects of diet.

Currently, we do not know how this transgenerational effect of lipid is transmitted, but

it may be due to altered epigenetic marks, nutrient provision from mother to

offspring, or microbiota. It will be interesting in future work to ask whether mito-

nuclear genotype shapes further aspects of non-genetic inheritance, and identify

mechanisms. Epigenomic profiling will likely prove insightful. Lipids and mito-nuclear

genotype may interact via mitochondrial metabolic processes such as beta-oxidation,

or by affecting rates of mitochondrial fusion and fission (Senyilmaz et al., 2015). It

will also be interesting to ask whether the transgenerational effects we have

identified in F_1 descendants extend further, to subsequent generations. These

findings indicate that detailed pedigrees containing both mitochondrial and nuclear

information may be required to predict transgenerational effects of nutrition.

A significant outcome of our study is that impacts of diet on offspring quality and

quantity do not necessarily correlate, and their relationship is subject to mito-nuclear

variation. We were surprised that EAA enrichment did not enhance offspring

development, because parental preference for laying eggs on this food led us to

expect that EAAs would promote offspring anabolism. This discrepancy may indicate

that EAAs function as signals of food quality as well as metabolites. This dual role

could drive deleterious outcomes when EAA levels are not representative of the

composition of food that would be found in the yeasts which flies are thought to

consume in nature. Alternatively, the discrepancy may prove an example of parent-

offspring conflict (Trivers, 1974), in which EAAs promote parental fecundity, despite

the detriment of reduced investment into individual offspring. How parent-offspring

conflict is regulated at the interface of diet and genetic variation in general is not

well-studied.

To our knowledge, our study is the first investigation of diet-mito-nuclear interactions

to manipulate specific nutrient classes, rather than a complex ingredient such as

yeast. Our finding that impacts of dietary lipid depend on mito-nuclear genotype may

be relevant to understanding variation in impacts of high-fat human diets. The high-

EAA diets that we used have parallels to high-protein anabolic diets used to increase

yields of livestock and human muscle mass. Our findings that EAAs generally

decrease offspring quality may give pause for thought in use of these diets.

Furthermore, the finding of mito-nuclear variation in the response to EAAs

enrichment demonstrates that certain individuals may bear particularly strong costs

of eating such diets. At present, we do not know the mechanistic basis of the diet-

mito-nuclear interactions that we have uncovered, but the transgenerational effects

hint at epigenetics, consistent with preceding findings that epigenetic marks respond

to genetic variation in the cytoplasm (Bellizzi et al., 2012; Grunau et al., 2018; Vivian

et al., 2017; Yan et al., 2011).

Our findings have implications for understanding variation that could emerge when

novel mito-nuclear pairings come to be, such as after mitochondrial replacement

therapy. Our results do not necessarily support the proposal to "match" mito-nuclear

pairs as a positive predictor of therapeutic success (Dowling et al., 2010; Mossman

et al., 2016a; Vaught et al., 2020), because a naturally co-occurring mito-nuclear pair

(*AA*) responded uniquely to diet, suggesting that costs and benefits are not

necessarily straightforward functions of mito-nuclear matching or mis-matching.

Thus, comparing performance of matched versus unmatched groups may not be

sufficient to evaluate risks of mitochondrial transfer therapy. Our analysis suggests

that mitochondrial transfer may be variously deleterious, beneficial, or have

unforeseen costs and benefits. Dietary recommendations may promote patient

health. Our data also lend kudos to the suggestion of transgenerational effects of

mitochondrial replacement therapy, especially when diet varies. A change of

nomenclature may ultimately be called for, since "matched" and "mis-matched" carry

intrinsic value judgments, but it seems that whether mitochondria and nuclei are of

shared origin does not necessarily predict cost or benefit. We suggest that deep

knowledge of patient and donor nuclear and mitochondrial genotype will likely be

required to predict a healthy match.

Our results have wider implications for the conduct of nutritional and genetic

research. Fly nutrition studies have drawn scrutiny for dietary variation between

studies, attributed to methodological reporting (Lesperance and Broderick, 2020) and

stochastic differences in ingredients (principally yeast) over time (Piper et al., 2014),

which may limit repeatability. Similar issues can ensue after inconsistent

standardisation of genetic background (Burnett et al., 2011). The mito-nuclear field

as a whole suggests that different outcomes are to be expected when mitochondrial

and nuclear haplotypes are not carefully controlled by backcrossing, and our results

indicate that variation between studies may be amplified when diet is manipulated.

Even stochastic variation in diet may interact with mito-nuclear variation, making the

case for careful reporting, and perhaps the use of chemically-defined media (Piper et

al., 2014). Accounting for these sources of variation will likely improve repeatability.

In summary, we have shown that diet, mitochondrial and nuclear haplotype have

complex interactive effects, each capable of modifying the impact of change in

another. Dietary lipid and EAAs are both implicated separately in these interactions,

with relevance not just for proximate reproductive output, but also for lasting

transgenerational effects of diet which impact relative offspring fitness. These

varying, mito-nuclear impacts of diet on offspring quantity and quality appear to be

an important determinant of individual fitness and health.

                          Materials & methods

Diets

Development medium contained 1.4% agar and 4.5% brewer's yeast (both

Gewürzmühle Brecht, Germany), 10% cornmeal and 11.1% sucrose (both Mühle

Milzitz, Germany) (all w/v), 0.45% propionic acid and 3% nipagin (v/v).

Experimental media build on published protocols (Bass et al., 2007; Dobson et al.,

2018; Emran et al., 2014). These media contained a final concentration of 10%

brewer's yeast, 5% sucrose, 1.5% agar (w/v), 3% nipagin and 0.3% propionic acid

(v/v). EAAs were purchased as powder (Sigma), and supplemented by dissolving

into a 6.66x solution in ddH_2O pH 4.5 (final media concentrations: L-arginine 0.43 g/l,

L-histidine 0.21 g/l, L-isoleucine 0.34 g/l, L-leucine 0.48 g/l, L-lysine 0.52 g/l, L-

methionine 0.1 g/l, L-phenylalanine 0.26 g/l, L-threonine 0.37 g/l, L-tryptophan 0.09

g/l, L-valine 0.4 g/l). Margarine (*Ja! Pflanzenmargarine* from Rewe Supermarkets,

Germany; 720 kcal / 100g; 80/100g fat from 23/100g saturated fatty acids, 40/100g

monounsaturated fatty acids, 17/100g polyunsaturated fatty acids) was briefly melted

then mixed thoroughly into the food (15% w/v). Final nutrient contents of rearing and

control media were estimated using the *Drosophila* diet content calculator

(Lesperance and Broderick, 2020), with additional protein, lipid and caloric content

after nutrient supplements calculated according to margarine nutrient content report,

and by assuming caloric equity between EAAs and protein at a caloric value of 4

calories/g (USDA). Vials contained ~5ml of food, and were stored at 4°C for up to 1

week before use.

Flies

D. melanogaster fly lines were established as described in Figure 1 and maintained

at 25°C on development medium throughout their history prior to our experiments.

The ancestral Australian population was isolated in Coffs Harbour, NSW, Australia

(Dowling et al., 2014). The Benin population is the widely-used *Dahomey* population,

isolated in the 1970s in Dahomey (now Benin). Ancestral lines had been cured of the

cytoplasmic endosymbiont *Wolbachia* 66 generations previously by tetracycline

treatment prior to the experiment. For each fly line, 45 females of the desired

mitochondrial background were crossed to 45 males of the desired nuclear

background per generation. Iterating this process over many generations led to

introgression of the desired nuclear background (from males) into each mitochondrial

background. For experiments, flies were collected upon eclosion to adulthood and

fed fresh developmental medium, before being pooled, split and assigned at random

to experimental medium in groups of 5 males and 5 females. Experimental flies were

maintained at 25°C, and transferred to fresh media every 48-72h for one week. Flies

were transferred to fresh medium 24h before egg laying experiments. For

development experiments, eggs were incubated at 25°C and pupation and eclosion

were scored daily. Eclosing adults were lightly CO_2-anaesthetised before counting

and sexing.

Analysis

Data were analysed in R 3.6.1. Markdowns of all analyses are provided as

supplementary material. Fit of fecundity data to a negative binomial distribution was

determined with firdistrplus::descdist and firdistrplus::fitdist. Generalised linear mixed

models of the form

$$y \sim diet * mitochondria * nuclear + (genotype)$$

were fit with lme4::glmer.nb (egg counts, negative binomial distribution) or

lme4::glmer (binomial of egg counts before and after diet switch); in which *diet*

(control/EAA/lipid), *mitochondria* (A/B) and *nuclear* (A/B) were fixed factors,

genotype was a random factor denoting the full replicated genotype (e.g. *AA1, AA2,*

AA3, AB1, AB2, BA1, etc). Where relevant (Figure 2), experimental replicate was

also included as an additional random factor. Dispersion was assessed with

blmeco::dispersion_glmer, and model singularity was tested with lme4::isSingular.

Anova tests (type-3) were conducted with car::Anova, and *post-hoc* analyses were

applied with the functions emmeans::joint_tests, emmeans::pairs, emmeans::emmip

(Searle et al., 1980). AIC was determined with stats::AIC, Akaike weights with

MuMIn::dredge. Four vials were excluded from the diet-switching experiment to

enable model fitting, on the basis of extreme values (minimum and maximum

observations in the experiment) or deviation from a QQ plot before diet switch.

These data are retained, with note, in the data associated to this paper.

Censoring (i.e. failure to develop) was inferred per vial when n. emerging adults was

less than n. eggs (noting assignment of zero censors when adult counts were

greater than eggs in some vials, due to measurement error). A 50:50 sex ratio was

assumed amongst censors. Development to adult was modelled by fitting Cox mixed

effects models of the form

$$y \sim diet * mitocondria * nuclear * sex + eggs + (\frac{genotype}{vial})$$

with coxme::coxme. *Diet, mitochondria, nuclear* and *genotype* terms were as in

models of egg laying, and *vial* was an additional random effect nested in *genotype*.

Eggs coded number of eggs laid in the vial in which the individual developed, to

account for variation in rearing density. Pupal sex was unknown, so the sex term

was excluded from models of pupal development. Anova and *post-hoc* EMM tests

were conducted as per fecundity analyses, with additional stratification by sex when

applying emmeans::joint_tests.

Figures were assembled in Adobe Illustrator. Mitochondria graphics were recoloured

from files freely distributed under an open commons licence. The heatmap of nutrient

content was plotted in R with the superheat library.

789   **Acknowledgments**

790   45

We thank Steven Parratt and Colin Selman for valuable comments on drafts of the

manuscript, and Luke Holman for advice on data analysis. Christin Froschauer and

Ralph Dobler provided invaluable advice in setting up experiments. This work was

supported by funds to KR from the Deutsche Forschungsgemeinschaft (Excellence

Initiative to TU Dresden). AJD was supported by a Dresden Fellowship funded by the

Excellence Initiative of the German Federal and State Governments, a UKRI Future

Leaders Fellowship (MR/S033939/1) and a University of Glasgow Lord Kelvin Adam

Smith Fellowship.

800   **Declaration of interests**

The authors declare no competing interests.

References

Akaike H. 1974. A new look at the statistical model identification. *Ieee T Automat*

Contr **19**:716–723. doi:10.1109/tac.1974.1100705

Aw WC, Towarnicki SG, Melvin RG, Youngson NA, Garvin MR, Hu Y, Nielsen S,

Thomas T, Pickford R, Bustamante S, Vila-Sanjurjo A, Smyth GK, Ballard JWO.

2018. Genotype to phenotype: Diet-by-mitochondrial DNA haplotype interactions

drive metabolic flexibility and organismal fitness. *Plos Genet* **14**:e1007735.

doi:10.1371/journal.pgen.1007735

Ballard JWO, Melvin RG. 2010. Linking the mitochondrial genotype to the organismal

phenotype. *Mol Ecol* **19**:1523–1539. doi:10.1111/j.1365-294x.2010.04594.x

Baris TZ, Wagner DN, Dayan DI, Du X, Blier PU, Pichaud N, Oleksiak MF, Crawford

DL. 2017. Evolved genetic and phenotypic differences due to mitochondrial-

nuclear interactions. *Plos Genet* **13**:e1006517. doi:10.1371/journal.pgen.1006517

Barker DJP, Osmond C. 1986. Infant mortality, childhood nutrition, and ischaemic

heart disease in England and Wales. *Lancet* **327**:1077–1081. doi:10.1016/s0140-

6736(86)91340-1

Barker DJP, Thornburg KL. 2013. The Obstetric Origins of Health for a Lifetime. *Clin*

Obstet Gynecol **56**:511–519. doi:10.1097/grf.0b013e31829cb9ca

Bass TM, Grandison RC, Wong R, Martinez P, Partridge L, Piper MDW. 2007.

Optimization of Dietary Restriction Protocols in Drosophila. *Journals Gerontology*

Ser **62**:1071–1081. doi:10.1093/gerona/62.10.1071

Bellizzi D, DAquila P, Giordano M, Montesanto A, Passarino G. 2012. Global DNA

methylation levels are modulated by mitochondrial DNA variants. *Epigenomics-uk*

4:17–27. doi:10.2217/epi.11.109

Bevers RPJ, Litovchenko M, Kapopoulou A, Braman VS, Robinson MR, Auwerx J,

Hollis B, Deplancke B. 2019. Mitochondrial haplotypes affect metabolic

phenotypes in the Drosophila Genetic Reference Panel. *Nat Metabolism* **1**:1226–

1242. doi:10.1038/s42255-019-0147-3

Bong L-J, Neoh K-B, Lee C-Y, Jaal Z. 2014. Effect of Diet Quality on Survival and

Reproduction of Adult Paederus fuscipes (Coleoptera: Staphylinidae). *J Med*

Entomol **51**:752–759. doi:10.1603/me13145

Brankatschk M, Gutmann T, Knittelfelder O, Palladini A, Prince E, Grzybek M,

Brankatschk B, Shevchenko A, Coskun Ü, Eaton S. 2018. A Temperature-

Dependent Switch in Feeding Preference Improves Drosophila Development and

Survival in the Cold. *Developmental Cell* **46**:781-793.e4.

doi:10.1016/j.devcel.2018.05.028

Burnett C, Valentini S, Cabreiro F, Goss M, Somogyvári M, Piper MD, Hoddinott M,

Sutphin GL, Leko V, McElwee JJ, Vazquez-Manrique RP, Orfila A-M, Ackerman

D, Au C, Vinti G, Riesen M, Howard K, Neri C, Bedalov A, Kaeberlein M, Sőti C,

Partridge L, Gems D. 2011. Absence of effects of Sir2 overexpression on lifespan

in C. elegans and Drosophila. *Nature* **477**:482–485. doi:10.1038/nature10296

Camilleri-Carter T-L, Dowling DK, Robker RL, Piper MDW. 2019. Transgenerational

Obesity and Healthy Aging in Drosophila. *The Journals of Gerontology: Series A*.

doi:10.1093/gerona/glz154

Camus MF, O'Leary M, Reuter M, Lane N. 2020. Impact of mitonuclear interactions

on life-history responses to diet. *Philosophical Transactions Royal Soc B*

375:20190416. doi:10.1098/rstb.2019.0416

Camus MF, Wolf JBW, Morrow EH, Dowling DK. 2015. Single Nucleotides in the

mtDNA Sequence Modify Mitochondrial Molecular Function and Are Associated

with Sex-Specific Effects on Fertility and Aging. *Curr Biol* **25**:2717–2722.

doi:10.1016/j.cub.2015.09.012

Chang C-C, Rodriguez J, Ross J. 2016. Mitochondrial–Nuclear Epistasis Impacts

Fitness and Mitochondrial Physiology of Interpopulation Caenorhabditis briggsae

Hybrids. *G3 Genes Genomes Genetics* **6**:209–219. doi:10.1534/g3.115.022970

Cohen IG, Adashi EY, Gerke S, Palacios-González C, Ravitsky V. 2020. The

Regulation of Mitochondrial Replacement Techniques Around the World. *Annu*

Rev Genom Hum G **21**:1–22. doi:10.1146/annurev-genom-111119-101815

Craven L, Tuppen HA, Greggains GD, Harbottle SJ, Murphy JL, Cree LM, Murdoch

AP, Chinnery PF, Taylor RW, Lightowlers RN, Herbert M, Turnbull DM. 2010.

Pronuclear transfer in human embryos to prevent transmission of mitochondrial

DNA disease. *Nature* **465**:82–85. doi:10.1038/nature08958

Crean AJ, Senior AM. 2019. High-fat diets reduce male reproductive success in

animal models: A systematic review and meta-analysis. *Obesity Reviews* **20**:921–

933. doi:10.1111/obr.12827

Deas JB, Blondel L, Extavour CG. 2019. Ancestral and offspring nutrition interact to

affect life-history traits in Drosophila melanogaster. *Proceedings of the Royal*

Society B **286**:20182778. doi:10.1098/rspb.2018.2778

Dobler R, Dowling DK, Morrow EH, Reinhardt K. 2018. A systematic review and

meta-analysis reveals pervasive effects of germline mitochondrial replacement on

components of health. *Hum Reprod Update* **24**:519–534.

doi:10.1093/humupd/dmy018

Dobson AJ, Boulton-McDonald R, Houchou L, Svermova T, Ren Z, Subrini J,

Vazquez-Prada M, Hoti M, Rodriguez-Lopez M, Ibrahim R, Gregoriou A,

Gkantiragas A, Bähler J, Ezcurra M, Alic N. 2019. Longevity is determined by ETS

transcription factors in multiple tissues and diverse species. *PLOS Genetics*

15:e1008212. doi:10.1371/journal.pgen.1008212

Dobson AJ, Chaston JM, Newell PD, Donahue L, Hermann SL, Sannino DR,

Westmiller S, Wong AC, Clark AG, Lazzaro BP, Douglas AE. 2015. Host genetic

determinants of microbiota-dependent nutrition revealed by genome-wide analysis

of Drosophila melanogaster. *Nature communications* **6**:6312.

doi:10.1038/ncomms7312

Dobson AJ, Ezcurra M, Flanagan CE, Summerfield AC, Piper MDW, Gems D, Alic N.

2017. Nutritional Programming of Lifespan by FOXO Inhibition on Sugar-Rich

Diets. *Cell Reports* **18**:299–306. doi:10.1016/j.celrep.2016.12.029

Dobson AJ, He X, Blanc E, Bolukbasi E, Feseha Y, Yang M, Piper MD. 2018.

Tissue-specific transcriptome profiling of Drosophila reveals roles for GATA

transcription factors in longevity by dietary restriction. *npj Aging and Mechanisms*

of Disease **4**. doi:10.1038/s41514-018-0024-4

Dong H-J, Wu D, Xu S-Y, Li Q, Fang Z-F, Che L-Q, Wu C-M, Xu X-Y, Lin Y. 2016.

Effect of dietary supplementation with amino acids on boar sperm quality and

fertility. *Anim Reprod Sci* **172**:182–189. doi:10.1016/j.anireprosci.2016.08.003

Đorđević M, Stojković B, Savković U, Immonen E, Tucić N, Lazarević J, Arnqvist G.

2017. Sex-specific mitonuclear epistasis and the evolution of mitochondrial

bioenergetics, ageing, and life history in seed beetles. *Evolution* **71**:274–288.

doi:10.1111/evo.13109

Dowling DK, Friberg U, Lindell J. 2008. Evolutionary implications of non-neutral

mitochondrial genetic variation. *Trends Ecol Evol* **23**:546–554.

doi:10.1016/j.tree.2008.05.011

Dowling DK, Meerupati T, Arnqvist G. 2010. Cytonuclear Interactions and the

Economics of Mating in Seed Beetles. *Am Nat* **176**:131–140. doi:10.1086/653671

Dowling DK, Williams BR, Garcia-Gonzalez F. 2014. Maternal sexual interactions

affect offspring survival and ageing. *J Evolution Biol* **27**:88–97.

doi:10.1111/jeb.12276

Drummond E, Short E, Clancy D. 2019. Mitonuclear gene X environment effects on

lifespan and health: How common, how big? *Mitochondrion* **49**:12–18.

doi:10.1016/j.mito.2019.06.009

Dunham-Snary KJ, Ballinger SW. 2015. Mitochondrial-nuclear DNA mismatch

matters. *Science* **349**:1449–1450. doi:10.1126/science.aac5271

Duque-Guimarães D, Ozanne S. 2017. Early nutrition and ageing: can we intervene?

Biogerontology **18**:893–900. doi:10.1007/s10522-017-9691-y

Eisenberg T, Schroeder S, Andryushkova A, Pendl T, Küttner V, Bhukel A, Mariño G,

Pietrocola F, Harger A, Zimmermann A, Moustafa T, Sprenger A, Jany E, Büttner

S, Carmona-Gutierrez D, Ruckenstuhl C, Ring J, Reichelt W, Schimmel K, Leeb

T, Moser C, Schatz S, Kamolz L-P, Magnes C, Sinner F, Sedej S, Fröhlich K-U,

Juhasz G, Pieber TR, Dengjel J, Sigrist SJ, Kroemer G, Madeo F. 2014.

Nucleocytosolic Depletion of the Energy Metabolite Acetyl-Coenzyme A

Stimulates Autophagy and Prolongs Lifespan. *Cell Metab* **19**:431–444.

doi:10.1016/j.cmet.2014.02.010

Ellison CK, Burton RS. 2006. Disruption of mitochondrial function in interpopulation

hybrids of Tigriopus californicus. *Evolution* **60**:1382–1391. doi:10.1554/06-210.1

Emran S, Yang M, He X, Zandveld J, Piper MD. 2014. Target of rapamycin signalling

mediates the lifespan-extending effects of dietary restriction by essential amino

acid alteration. *Aging* **6**:390–8.

Fanson BG, Fanson KV, Taylor PW. 2012. Cost of reproduction in the Queensland

fruit fly: Y-model versus lethal protein hypothesis. *Proc Royal Soc B Biological Sci*

279:4893–4900. doi:10.1098/rspb.2012.2033

Fontana L, Partridge L, Longo VD. 2010. Extending Healthy Life Span—From Yeast

to Humans. *Science* **328**:321–326. doi:10.1126/science.1172539

Gemmell NJ, Metcalf VJ, Allendorf FW. 2004. Mother's curse: the effect of mtDNA

on individual fitness and population viability. *Trends Ecol Evol* **19**:238–244.

doi:10.1016/j.tree.2004.02.002

Gershoni M, Templeton AR, Mishmar D. 2009. Mitochondrial bioenergetics as a

major motive force of speciation. *Bioessays* **31**:642–650.

doi:10.1002/bies.200800139

Gorman GS, Chinnery PF, DiMauro S, Hirano M, Koga Y, McFarland R,

Suomalainen A, Thorburn DR, Zeviani M, Turnbull DM. 2016. Mitochondrial

diseases. *Nat Rev Dis Primers* **2**:16080. doi:10.1038/nrdp.2016.80

Grandison RC, Piper MD, Partridge L. 2009. Amino-acid imbalance explains

extension of lifespan by dietary restriction in Drosophila. *Nature* **462**:1061–1064.

doi:10.1038/nature08619

Grunau C, Voigt S, Dobler R, Dowling DK, Reinhardt K. 2018. The Cytoplasm

Affects the Epigenome in Drosophila melanogaster. *Epigenomes* **2**:17.

doi:10.3390/epigenomes2030017

Hariri N, Thibault L. 2010. High-fat diet-induced obesity in animal models. *Nutr Res*

Rev **23**:270–99. doi:10.1017/s0954422410000168

Havird JC, Hall MD, Dowling DK. 2015. The evolution of sex: A new hypothesis

based on mitochondrial mutational erosion. *Bioessays* **37**:951–958.

doi:10.1002/bies.201500057

Healy TM, Burton RS. 2020. Strong selective effects of mitochondrial DNA on the

nuclear genome. *P Natl Acad Sci Usa* **117**:6616–6621.

doi:10.1073/pnas.1910141117

Heymsfield SB, Wadden TA. 2017. Mechanisms, Pathophysiology, and Management

of Obesity. *New Engl J Med* **376**:254–266. doi:10.1056/nejmra1514009

Hill GE. 2019. Reconciling the mitonuclear compatibility species concept with

rampant mitochondrial introgression. *Integr Comp Biol* **59**:912–924.

doi:10.1093/icb/icz019

Hill GE. 2017. The mitonuclear compatibility species concept. *Auk* **134**:393–409.

doi:10.1642/auk-16-201.1

Holland ML, Lowe R, Caton PW, Gemma C, Carbajosa G, Danson AF, Carpenter

AA, Loche E, Ozanne SE, Rakyan VK. 2016. Early-life nutrition modulates the

epigenetic state of specific rDNA genetic variants in mice. *Science (New York,*

NY) **353**:495–8. doi:10.1126/science.aaf7040

Human Fertility and Embryology Authority. 2016. Scientific review of the safety and

efficacy of methods to avoid mitochondrial disease through assisted conception:

2016 update.

Innocenti P, Morrow EH, Dowling DK. 2011. Experimental Evidence Supports a Sex-

Specific Selective Sieve in Mitochondrial Genome Evolution. *Science* **332**:845–

848. doi:10.1126/science.1201157

Jang C, Hui S, Lu W, Cowan AJ, Morscher RJ, Lee G, Liu W, Tesz GJ, Birnbaum

MJ, Rabinowitz JD. 2018. The Small Intestine Converts Dietary Fructose into

Glucose and Organic Acids. *Cell Metab* **27**:351-361.e3.

doi:10.1016/j.cmet.2017.12.016

Jensen K, McClure C, Priest NK, Hunt J. 2015. Sex-specific effects of protein and

carbohydrate intake on reproduction but not lifespan in Drosophila melanogaster.

Aging Cell **14**:605–615. doi:10.1111/acel.12333

Jumbo-Lucioni P, Ayroles JF, Chambers M, Jordan KW, Leips J, Mackay TF, Luca

M. 2010. Systems genetics analysis of body weight and energy metabolism traits

in Drosophila melanogaster. *BMC Genomics* **11**:297. doi:10.1186/1471-2164-11-

297

Knittelfelder O, Prince E, Sales S, Fritzsche E, Wöhner T, Brankatschk M,

Shevchenko A. 2020. Sterols as dietary markers for Drosophila melanogaster.

Biochimica Et Biophysica Acta Bba - Mol Cell Biology Lipids **1865**:158683.

doi:10.1016/j.bbalip.2020.158683

Lack D. 1947. The significance of clutch size in birds. *Ibis* 302–352.

Latorre-Pellicer A, Lechuga-Vieco AV, Johnston IG, Hämäläinen RH, Pellico J,

Justo-Méndez R, Fernández-Toro JM, Clavería C, Guaras A, Sierra R, Llop J,

Torres M, Criado LM, Suomalainen A, Jones NS, Ruíz-Cabello J, Enríquez JA.

2019. Regulation of Mother-to-Offspring Transmission of mtDNA Heteroplasmy.

Cell Metab **30**:1120-1130.e5. doi:10.1016/j.cmet.2019.09.007

Lee K, Simpson SJ, Clissold FJ, Brooks R, Ballard WJ, Taylor PW, Soran N,

Raubenheimer D. 2008. Lifespan and reproduction in Drosophila: New insights

from nutritional geometry. *Proceedings of the National Academy of Sciences*

105:2498–2503. doi:10.1073/pnas.0710787105

Leitão-Gonçalves R, Carvalho-Santos Z, Francisco A, Fioreze G, Anjos M, Baltazar

C, Elias A, Itskov PM, Piper MD, Ribeiro C. 2017. Commensal bacteria and

essential amino acids control food choice behavior and reproduction. *PLOS*

Biology **15**:e2000862. doi:10.1371/journal.pbio.2000862

Lesperance DNA, Broderick NA. 2020. Meta-analysis of Diets Used in Drosophila

Microbiome Research and Introduction of the Drosophila Dietary Composition

Calculator (DDCC). *G3 Genes Genomes Genetics* **10**:g3.401235.2020.

doi:10.1534/g3.120.401235

Li X, Shi X, Hou Y, Cao X, Gong L, Wang H, Li Jiayu, Li Jibin, Wu C, Xiao D, Qi H,

Xiao X. 2018. Paternal hyperglycemia induces transgenerational inheritance of

susceptibility to hepatic steatosis in rats involving altered methylation on Pparα

promoter. *Biochimica Et Biophysica Acta Mol Basis Dis* **1865**:147–160.

doi:10.1016/j.bbadis.2018.10.040

Liang H, Zhang Z. 2006. Food restriction affects reproduction and survival of F1 and

F2 offspring of Rat-like hamster (Cricetulus triton). *Physiol Behav* **87**:607–613.

doi:10.1016/j.physbeh.2005.12.006

Liao C-Y, Rikke BA, Johnson TE, Diaz V, Nelson JF. 2009. Genetic variation in the

murine lifespan response to dietary restriction: from life extension to life

shortening. *Aging Cell* **9**:92–5. doi:10.1111/j.1474-9726.2009.00533.x

Ma C, Mirth CK, Hall MD, Piper MDW. 2020. Amino acid quality modifies the

quantitative availability of protein for reproduction in Drosophila melanogaster. *J*

Insect Physiol 104050. doi:10.1016/j.jinsphys.2020.104050

Ma H, Gutierrez NM, Morey R, Dyken CV, Kang E, Hayama T, Lee Y, Li Y, Tippner-

Hedges R, Wolf DP, Laurent LC, Mitalipov S. 2016. Incompatibility between

Nuclear and Mitochondrial Genomes Contributes to an Interspecies Reproductive

Barrier. *Cell Metab* **24**:283–94. doi:10.1016/j.cmet.2016.06.012

Maklakov AA, Lummaa V. 2013. Evolution of sex differences in lifespan and aging:

Causes and constraints. *BioEssays* **35**:717–724. doi:10.1002/bies.201300021

Maklakov AA, Simpson SJ, Zajitschek F, Hall MD, Dessmann J, Clissold F,

Raubenheimer D, Bonduriansky R, Brooks RC. 2008. Sex-Specific Fitness Effects

of Nutrient Intake on Reproduction and Lifespan. *Curr Biol* **18**:1062–1066.

doi:10.1016/j.cub.2008.06.059

Mariño G, Pietrocola F, Eisenberg T, Kong Y, Malik SA, Andryushkova A, Schroeder

S, Pendl T, Harger A, Niso-Santano M, Zamzami N, Scoazec M, Durand S, Enot

DP, Fernández ÁF, Martins I, Kepp O, Senovilla L, Bauvy C, Morselli E, Vacchelli

E, Bennetzen M, Magnes C, Sinner F, Pieber T, López-Otín C, Maiuri MC,

Codogno P, Andersen JS, Hill JA, Madeo F, Kroemer G. 2014. Regulation of

Autophagy by Cytosolic Acetyl-Coenzyme A. *Mol Cell* **53**:710–725.

doi:10.1016/j.molcel.2014.01.016

McCracken AW, Adams G, Hartshorne L, Tatar M, Simons MJP. 2020. The hidden

costs of dietary restriction: Implications for its evolutionary and mechanistic

origins. *Sci Adv* **6**:eaay3047. doi:10.1126/sciadv.aay3047

Meiklejohn CD, Holmbeck MA, Siddiq MA, Abt DN, Rand DM, Montooth KL. 2013.

An Incompatibility between a Mitochondrial tRNA and Its Nuclear-Encoded tRNA

Synthetase Compromises Development and Fitness in Drosophila. *Plos Genet*

9:e1003238. doi:10.1371/journal.pgen.1003238

Milot E, Moreau C, Gagnon A, Cohen AA, Brais B, Labuda D. 2017. Mother's curse

neutralizes natural selection against a human genetic disease over three

centuries. *Nat Ecol Evol* **1**:1400–1406. doi:10.1038/s41559-017-0276-6

Mirth CK, Alves AN, Piper MD. 2018. Turning Food Into Eggs: insights from

nutritional biology and developmental physiology of Drosophila. *Curr Opin Insect*

Sci **31**:49–57. doi:10.1016/j.cois.2018.08.006

Montooth KL, Dhawanjewar AS, Meiklejohn CD. 2019. Temperature-sensitive

reproduction and the physiological and evolutionary potential for Mother's Curse.

Integr Comp Biol **59**:890–899. doi:10.1093/icb/icz091

Mossman JA, Biancani LM, Rand DM. 2019. Mitochondrial genomic variation drives

differential nuclear gene expression in discrete regions of Drosophila gene and

protein interaction networks. *Bmc Genomics* **20**:691. doi:10.1186/s12864-019-

6061-y

Mossman JA, Biancani LM, Zhu C-T, Rand DM. 2016a. Mitonuclear Epistasis for

Development Time and Its Modification by Diet in Drosophila. *Genetics* **203**:463–

484. doi:10.1534/genetics.116.187286

Mossman JA, Tross JG, Li N, Wu Z, Rand DM. 2016b. Mitochondrial-Nuclear

Interactions Mediate Sex-Specific Transcriptional Profiles in Drosophila. *Genetics*

204:genetics.116.192328. doi:10.1534/genetics.116.192328

Mulvey L, Sands WA, Salin K, Carr AE, Selman C. 2017. Disentangling the effect of

dietary restriction on mitochondrial function using recombinant inbred mice.

Molecular and cellular endocrinology **455**:41–53. doi:10.1016/j.mce.2016.09.001

Nagarajan-Radha V, Aitkenhead I, Clancy DJ, Chown SL, Dowling DK. 2020. Sex-

specific effects of mitochondrial haplotype on metabolic rate in Drosophila

melanogaster support predictions of the Mother's Curse hypothesis. *Philosophical*

Transactions Royal Soc B **375**:20190178. doi:10.1098/rstb.2019.0178

Nagarajan-Radha V, Rapkin J, Hunt J, Dowling DK. 2019. Interactions Between

Mitochondrial Haplotype and Dietary Macronutrient Ratios Confer Sex-Specific

Effects on Longevity in Drosophila melanogaster. *The Journals of Gerontology:*

Series A 1573–1581. doi:10.1093/gerona/glz104

Ordovas JM, Ferguson LR, Tai ES, Mathers JC. 2018. Personalised nutrition and

health. *Bmj Clin Res Ed* **361**:bmj.k2173. doi:10.1136/bmj.k2173

Öst A, Lempradl A, Casas E, Weigert M, Tiko T, Deniz M, Pantano L, Boenisch U,

Itskov PM, Stoeckius M, Ruf M, Rajewsky N, Reuter G, Iovino N, Ribeiro C,

Alenius M, Heyne S, Vavouri T, Pospisilik AJ. 2014. Paternal Diet Defines

Offspring Chromatin State and Intergenerational Obesity. *Cell* **159**.

doi:10.1016/j.cell.2014.11.005

Piper MD, Blanc E, Leitão-Gonçalves R, Yang M, He X, Linford NJ, Hoddinott MP,

Hopfen C, Soultoukis GA, Niemeyer C, Kerr F, Pletcher SD, Ribeiro C, Partridge

L. 2014. A holidic medium for Drosophila melanogaster. *Nature Methods* **11**:100–

105. doi:10.1038/nmeth.2731

Rand DM, Mossman JA, Zhu L, Biancani LM, Ge JY. 2018. Mitonuclear epistasis,

genotype-by-environment interactions, and personalized genomics of complex

traits in Drosophila: Mitonuclear G x G x E. *Iubmb Life* **70**:1275–1288.

doi:10.1002/iub.1954

Reinhardt K, Dowling DK, Morrow EH. 2013. Mitochondrial Replacement, Evolution,

and the Clinic. *Science* **341**:1345–1346. doi:10.1126/science.1237146

Rikke BA, Liao C-Y, McQueen MB, Nelson JF, Johnson TE. 2010. Genetic

dissection of dietary restriction in mice supports the metabolic efficiency model of

life extension. *Exp Gerontol* **45**:691–701. doi:10.1016/j.exger.2010.04.008

Rivera HM, Kievit P, Kirigiti MA, Bauman LA, Baquero K, Blundell P, Dean TA,

Valleau JC, Takahashi DL, Frazee T, Douville L, Majer J, Smith MS, Grove KL,

Sullivan EL. 2015. Maternal high-fat diet and obesity impact palatable food intake

and dopamine signaling in nonhuman primate offspring. *Obes Silver Spring Md*

23:2157–64. doi:10.1002/oby.21306

Saud S, Summerfield AC, Alic N. 2015. Ablation of insulin-producing cells prevents

obesity but not premature mortality caused by a high-sugar diet in Drosophila.

Proceedings of the Royal Society of London B: Biological Sciences

282:20141720. doi:10.1098/rspb.2014.1720

Searle SR, Speed FM, Milliken GA. 1980. Population Marginal Means in the Linear

Model: An Alternative to Least Squares Means. *Am Statistician* **34**:216–221.

doi:10.1080/00031305.1980.10483031

Senyilmaz D, Virtue S, Xu X, Tan CY, Griffin JL, Miller AK, Vidal-Puig A, Teleman

AA. 2015. Regulation of mitochondrial morphology and function by stearoylation

of TFR1. *Nature* **525**:124–128. doi:10.1038/nature14601

Simpson SJ, Raubenheimer D. 2012. The Nature of Nutrition.

doi:10.1515/9781400842803

Skorupa DA, Dervisefendic A, Zwiener J, Pletcher SD. 2008. Dietary composition

specifies consumption, obesity, and lifespan in Drosophila melanogaster. *Aging*

Cell **7**:478–490. doi:10.1111/j.1474-9726.2008.00400.x

Smith HG, Kallander H, Nilsson J-A. 1989. The Trade-Off Between Offspring

Number and Quality in the Great Tit Parus major. *J Animal Ecol* **58**:383.

doi:10.2307/4837

Solon-Biet SM, Walters KA, Simanainen UK, McMahon AC, Ruohonen K, Ballard

JO, Raubenheimer D, Handelsman DJ, Couteur DG, Simpson SJ. 2015.

Macronutrient balance, reproductive function, and lifespan in aging mice.

Proceedings of the National Academy of Sciences **112**:3481–3486.

doi:10.1073/pnas.1422041112

Stearns SC. 1992. The evolution of life histories. Oxford University Press.

Stefana IM, Driscoll PC, Obata F, Pengelly A, Newell CL, MacRae JI, Gould AP.

2017. Developmental diet regulates Drosophila lifespan via lipid autotoxins.

Nature Communications **8**:1384. doi:10.1038/s41467-017-01740-9

Svajgr AJ, Hammell DL, DeGeeter MJ, Hays VW, Cromwell GL, Dutt RH. 1972.

Reproductive performance of sows on a protein-restricted diet. *Reproduction*

30:455–458. doi:10.1530/jrf.0300455

Tachibana M, Kuno T, Yaegashi N. 2018. Mitochondrial replacement therapy and

assisted reproductive technology: A paradigm shift toward treatment of genetic

diseases in gametes or in early embryos. *Reproductive Medicine Biology* **17**:421–

433. doi:10.1002/rmb2.12230

Towarnicki SG, Ballard JWO. 2018. Mitotype Interacts With Diet to Influence

Longevity, Fitness, and Mitochondrial Functions in Adult Female Drosophila.

Frontiers in Genetics **9**:593. doi:10.3389/fgene.2018.00593

Towarnicki SG, Ballard JWO. 2017. Drosophila mitotypes determine developmental

time in a diet and temperature dependent manner. *Journal of Insect Physiology*

100. doi:10.1016/j.jinsphys.2017.06.002

Trefely S, Lovell CD, Snyder NW, Wellen KE. 2020. Compartmentalized acyl-CoA

metabolism and roles in chromatin regulation. *Mol Metab* **38**:100941.

doi:10.1016/j.molmet.2020.01.005

Trivers RL. 1974. Parent-Offspring Conflict. *Am Zool* **14**:249–264.

doi:10.1093/icb/14.1.249

Tufarelli V, Lacalandra G, Laudadio V. 2015. Reproductive and Metabolic

Responses of Early-lactating Dairy Cows Fed Different Dietary Protein Sources.

Reprod Domest Anim **50**:735–739. doi:10.1111/rda.12566

Vaught RC, Voigt S, Dobler R, Clancy DJ, Reinhardt K, Dowling DK. 2020.

Interactions between cytoplasmic and nuclear genomes confer sex-specific

effects on lifespan in Drosophila melanogaster. *J Evolution Biol* **33**:694–713.

doi:10.1111/jeb.13605

Vivian CJ, Brinker AE, Graw S, Koestler DC, Legendre C, Gooden GC, Salhia B,

Welch DR. 2017. Mitochondrial Genomic Backgrounds Affect Nuclear DNA

Methylation and Gene Expression. *Cancer Res* **77**:6202–6214. doi:10.1158/0008-

5472.can-17-1473

Wagenmakers E-J, Farrell S. 2004. AIC model selection using Akaike weights.

Psychon B Rev **11**:192–196. doi:10.3758/bf03206482

Wallace DC, Chalkia D. 2013. Mitochondrial DNA Genetics and the Heteroplasmy

Conundrum in Evolution and Disease. *Csh Perspect Biol* **5**:a021220.

doi:10.1101/cshperspect.a021220

Wei Y, Yang C-R, Wei Y-P, Zhao Z-A, Hou Y, Schatten H, Sun Q-Y. 2014. Paternally

induced transgenerational inheritance of susceptibility to diabetes in mammals. *P*

Natl Acad Sci Usa **111**:1873–8. doi:10.1073/pnas.1321195111

Winship AL, Gazzard SE, Cullen-McEwen LA, Bertram JF, Hutt KJ. 2018. Maternal

low-protein diet programmes low ovarian reserve in offspring. *Reproduction*

156:299–311. doi:10.1530/rep-18-0247

Wolf DP, Mitalipov N, Mitalipov S. 2015. Mitochondrial replacement therapy in

reproductive medicine. *Trends Mol Med* **21**:68–76.

doi:10.1016/j.molmed.2014.12.001

Wong AC-N, Dobson AJ, Douglas AE. 2014a. Gut microbiota dictates the metabolic

response of Drosophila to diet. *J Exp Biology* **217**:1894–1901.

doi:10.1242/jeb.101725

Woodcock KJ, Kierdorf K, Pouchelon CA, Vivancos V, Dionne MS, Geissmann F.

2015. Macrophage-Derived upd3 Cytokine Causes Impaired Glucose

Homeostasis and Reduced Lifespan in Drosophila Fed a Lipid-Rich Diet.

Immunity **42**:133–144. doi:10.1016/j.immuni.2014.12.023

Yan H, Yan Z, Ma Q, Jiao F, Huang S, Zeng F, Zeng Y. 2011. Association between

mitochondrial DNA haplotype compatibility and increased efficiency of bovine

intersubspecies cloning. *J Genet Genomics* **38**:21–28.

doi:10.1016/j.jcg.2010.12.003

Zanco B, Mirth CK, Sgrò CM, Piper MDW. 2020. A dietary sterol trade off determines

lifespan responses to dietary restriction in Drosophila melanogaster. *Biorxiv*

2020.08.21.260489. doi:10.1101/2020.08.21.260489

Zhao S, Jang C, Liu J, Uehara K, Gilbert M, Izzo L, Zeng X, Trefely S, Fernandez S,

Carrer A, Miller KD, Schug ZT, Snyder NW, Gade TP, Titchenell PM, Rabinowitz

JD, Wellen KE. 2020. Dietary fructose feeds hepatic lipogenesis via microbiota-

derived acetate. *Nature* **579**:586–591. doi:10.1038/s41586-020-2101-7

Zhu C-T, Ingelmo P, Rand DM. 2014. G×G×E for Lifespan in Drosophila:

Mitochondrial, Nuclear, and Dietary Interactions that Modify Longevity. *PLoS*

Genetics **10**. doi:10.1371/journal.pgen.1004354

1188

Table S1. GLMM of fecundity across diets and mito-nuclear haplotypes

Term	Chisq	Df	$Pr(>\chi^2)$
(Intercept)	4342.751	1	<2.20E-16
mito	8.3384	1	0.0039
nuclear	13.6946	1	0.0002
diet	314.9242	2	<2.20E-16
mito:nuclear	6.5883	1	0.0103
mito:diet	10.2151	2	0.0061
nuclear:diet	0.2024	2	0.9036
mito:nuclear:diet	9.3347	2	0.0094

Table S2. Pairwise comparisons of fecundity across diets and mito-nuclear haplotypes

diet	contrast	estimate	SE	Z ratio	P value
	AA vs BA	0.264859	0.0917	2.888	0.0203
	AA vs AB	0.32267	0.0872	3.701	0.0012
	AA vs BB	0.262022	0.0871	3.008	0.014
sya	BA vs AB	0.057811	0.0922	0.627	0.9233
	BA vs BB	-0.002838	0.092	-0.031	1
	AB vs BB	-0.060649	0.0876	-0.692	0.9001
	AA vs BA	0.04355	0.0918	0.474	0.9648
EAA	AA vs AB	0.362769	0.0865	4.193	0.0002
	AA vs BB	0.363417	0.0866	4.199	0.0002

BA vs AB	0.319219	0.0922	3.464	0.003
BA vs BB	0.319866	0.092	3.475	0.0029
AB vs BB	0.000647	0.087	0.007	1
AA vs BA	-0.09705	0.097	-1.001	0.7491
AA vs AB	0.365225	0.0932	3.918	0.0005
AA vs BB	0.42652	0.0935	4.56	<.0001
mar — BA vs AB	0.462275	0.0982	4.706	<.0001
BA vs BB	0.523571	0.0986	5.309	<.0001
AB vs BB	0.061296	0.0951	0.645	0.9174

Table S3. Development on experimental diets: Cox Mixed-Effects model of adult emergence, sex-stratified posterior analysis by Estimated Marginal Means

Sex	Term	Df	F ratio	P
	mito	1	15.517	0.0001
	nuclear	1	13.541	0.0002
	diet	2	322.17	<.0001
female	mito:nuclear	1	20.686	<.0001
	mito:diet	2	25.789	<.0001
	nuclear:diet	2	16.855	<.0001
	mito:nuclear:diet	2	25.733	<.0001
male	mito	1	10.307	0.0013
	nuclear	1	0.223	0.6371

diet	2	226.121	<.0001
mito:nuclear	1	16.511	<.0001
mito:diet	2	8.529	0.0002
nuclear:diet	2	15.201	<.0001
mito:nuclear:diet	2	7.778	0.0004

Table S4. Development on experimental diets: Cox Mixed-Effects model of egg-pupa survival

Term	Chisq	Df	$Pr(>\chi^2)$
mito	1	129.732	<2.20E-16
nuclear	1	148.282	<2.20E-16
diet	2	146.228	<2.20E-16
eggs	1	12.377	0.0004
mito:nuclear	1	66.987	2.73E-16
mito:diet	2	74.55	<2.20E-16
nuclear:diet	2	90.713	<2.20E-16
mito:nuclear:diet	2	35.714	1.76E-08

Table S5. Fecundity before switch from experimental diets: GLMM of egg laying across mito-nuclear haplotypes, after feeding on experimental diets

Term	Chisq	Df	$Pr(>\chi^2)$
(Intercept)	1748.5684	1	<2.20E-16
mito	1.1888	1	0.2756
nuclear	2.2325	1	0.1351

diet	127.1386	2	<2.20E-16
mito:nuclear	5.845	1	0.0156
mito:diet	9.7233	2	0.0077
nuclear:diet	10.8029	2	0.0045
mito:nuclear:diet	9.1987	2	0.0101

Table S6. GLMM of fecundity across diets and mito-nuclear haplotypes – data pooled from Figure 2 and Figure 3c

Term	χ^2	Df	$Pr(>\chi^2)$
(Intercept)	4338.6592	1	<2.20E-16
mito	1.7911	1	0.1808
nuclear	9.8394	1	0.0017
diet	500.2988	2	<2.20E-16
mito:nuclear	1.8161	1	0.1779
mito:diet	14.5664	2	0.0006
nuclear:diet	2.829	2	0.2430
mito:nuclear:diet	17.8267	2	0.0001

Table S7. Fecundity after switch from experimental diets: GLMM of egg laying across mito-nuclear haplotypes, after feeding on experimental diets

Term	χ^2	Df	$Pr(>\chi^2)$
(Intercept)	1341.3688	1	<2.20E-16
mito	2.6895	1	0.1010
nuclear	0.0347	1	0.8522

Term	χ^2	Df	Pr(>χ^2)
diet	36.59	2	1.13E-08
mito:nuclear	3.7278	1	0.0535
mito:diet	0.3195	2	0.8523
nuclear:diet	1.7084	2	0.4256
mito:nuclear:diet	0.5948	2	0.7427

Table S8. Rescue of fecundity after switch from experimental diets: GLMM of egg laying across mito-nuclear haplotypes: after feeding, relative to during feeding (binomial)

Term	χ^2	Df	Pr(>χ^2)
(Intercept)	44.7143	1	2.28E-11
mito	1.4802	1	0.2237
nuclear	4.4348	1	0.0352
diet	139.1165	2	<2.20E-16
mito:nuclear	0.1714	1	0.6788
mito:diet	51.4892	2	6.60E-12
nuclear:diet	17.9358	2	0.0001
mito:nuclear:diet	39.3128	2	2.91E-09

Table S9. Fertility & development after parental feeding on experimental diets: sex-stratified posterior (EMM) analysis of adult emergence

Sex	Term	Df	F ratio	P
female	mito	1	10.243	0.0014
	nuclear	1	1.313	0.2518

	Term	Df	Chisq	Pr
	diet	2	0.345	0.7083
	mito:nuclear	1	0.218	0.6407
	mito:diet	2	4.961	0.007
	nuclear:diet	2	0.777	0.4599
	mito:nuclear:diet	2	3.477	0.0309
	mito	1	9.262	0.0023
	nuclear	1	1.331	0.2486
	diet	2	0.396	0.6731
male	mito:nuclear	1	0.07	0.7913
	mito:diet	2	3.454	0.0316
	nuclear:diet	2	1.545	0.2133
	mito:nuclear:diet	2	5.862	0.0028

Table S10. Development after parental feeding on experimental diets: Cox Mixed-Effects model of egg-pupa survival

Term	Chisq	Df	Pr(>χ^2)
mito	1	3.8079	0.0510
nuclear	1	13.3143	0.0003
diet	2	10.8094	0.0045
eggs	1	31.5515	1.94E-08
mito:nuclear	1	3.8928	0.0485

mito:diet	2	5.5291	0.0630
nuclear:diet	2	8.4317	0.0148
mito:nuclear:diet	2	1.717	0.4238

Table S11. Fertility & development on experimental diets: Cox Mixed-Effects model of adult emergence

Term	Chisq	Df	$Pr(>\chi^2)$
mito	1	75.1042	<2.20E-16
nuclear	1	65.8859	4.78E-16
diet	2	86.1542	<2.20E-16
sex	1	12.9677	0.0003
eggs	1	1.7022	0.1920
mito:nuclear	1	55.7929	8.05E-14
mito:diet	2	86.1953	<2.20E-16
nuclear:diet	2	68.6659	1.23E-15
mito:sex	1	14.1785	0.0002
nuclear:sex	1	14.9357	0.0001
diet:sex	2	12.8281	0.0016
mito:nuclear:diet	2	51.4656	6.67E-12
mito:nuclear:sex	1	13.8949	0.0002
mito:diet:sex	2	25.9358	2.33E-06
nuclear:diet:sex	2	11.5055	0.0032
mito:nuclear:diet:sex	2	16.6248	0.0002

Table S12. Fertility & development after parental feeding on experimental diets: Cox Mixed-Effects model of adult emergence

Term	Chisq	Df	Pr(>χ^2)
mito	1	0.3711	0.5423
nuclear	1	1.9887	0.1585
diet	2	10.3938	0.0055
sex	1	0.1665	0.6832
eggs	1	7.2316	0.0072
mito:nuclear	1	0.2117	0.6454
mito:diet	2	14.3708	0.0008
nuclear:diet	2	6.0394	0.0488
mito:sex	1	1.517	0.2181
nuclear:sex	1	1.4268	0.2323
diet:sex	2	2.9565	0.2280
mito:nuclear:diet	2	6.9531	0.0309
mito:nuclear:sex	1	0.565	0.4522
mito:diet:sex	2	0.2748	0.8716
nuclear:diet:sex	2	1.0484	0.5920
mito:nuclear:diet:sex	2	6.6992	0.0351

Figure S1

Figure S1. Diet-mito-nuclear interactions modulate *Drosophila* fecundity.

Genetic correlations in impact of mito-nuclear variation on egg laying, and impact of diet on egg laying. Each point shows mean egg laying per line per diet in each of two replicate experiments. Bars show standard error. Accompanying statistical analysis presented in Table S1 and Table S2.

Figure S2

Figure S2. Mito-nuclear variation in dietary regulation of adult development: impacts per genetic replicate. Kaplan-Meier plots of development after feeding on experimental diets as specified in key. Mito-nuclear genotypes are indicated per panel. Data are presented pooled across each replicate genotype in Figure 4. Accompanying statistical analysis presented in Table S3.

Figure S3. Mito-nuclear variation in dietary regulation of pupation. (**A**) Kaplan-Meier plots of pupation after feeding on experimental diets as specified in key. Mito-nuclear genotypes are indicated in top-left of each panel. (**B**). Estimated marginal means (EMMs) with 95% confidence intervals summarising survival analysis (Cox mixed effects) of data from A. Higher values correspond to faster development and/or increased fertility, lower values correspond to slower development and/or reduced viability. Data are presented per genetic replicate in Figure S4. Accompanying statistical analysis presented in Table S4.

Figure S4

Figure S4. Mito-nuclear variation in dietary regulation of pupation: impacts per genetic replicate. Kaplan-Meier plots of pupation after feeding on experimental diets as specified in key. Mito-nuclear genotypes are indicated per panel. Data are presented pooled across each replicate genotype in Figure S5. Accompanying statistical analysis presented in Table S4 and Table S10

Figure S5

Figure S5. Mito-nuclear variation in fecundity response to switching from experimental diets to a standardised diet: impacts per genetic replicate.
Fecundity measures are shown per genotype as specified on bottom x axis. Boxplots show medians, first and third quartiles and 5th and 95th percentiles. Points show individual data. **A**. Fecundity before diet switch. **B.** Fecundity after diet switch. **C.** Ratio of fecundity before and after diet switch. Data are presented pooled across each replicate genotype in Figure 3. Accompanying statistical analysis presented in Tables S5, S7 and S8.

Figure S6

Figure S6. Diet-mito-nuclear interactions modulate *Drosophila* fecundity -

emergent pattern from three replicate experiments. Fecundity is shown per

genotype as specified on bottom x axis. Boxplots show medians, first and third

quartiles and 5th and 95th percentiles. Connected points to right of each box show

post-hoc comparisons of means (by estimated marginal means with 95% confidence

intervals, calculated from a generalised linear mixed effects model, showing

exponent of EMMs in order to fit to original data scale). Translucent points to left of

each box show raw data. Accompanying statistical analysis presented in Table S6.

Supplementary Text

Much recent research in *Drosophila* and other organisms has been directed to characterising the consequences transmission of mitochondria from mothers only, which restricts mito-nuclear coevolution to females (Gemmell et al., 2004; Maklakov and Lummaa, 2013; Montooth et al., 2019; Nagarajan-Radha et al., 2020). This is thought to cause a "mother's curse" (Gemmell et al., 2004), in which males bear their mothers' mitochondria despite potential mismatch to variants inherited from fathers. In our study of development we manipulated diet and mito-nuclear genotype, and recorded sex of emerging adult offspring, giving four potential predictive factors. Our main analyses are presented stratified by sex. ANOVA tests of the complex diet:mito:nuclear:sex interaction were nominally supportive of sex biases in DMN interactions, but did not point solely to males. We detected significant sex:diet:mito:nuclear interactions, but even with our large sample sizes we are cautious in interpreting such complex interaction terms. F-ratios for DMN interactions were greater in males when only parents were fed experimental diets (Table S12), but were greater in females when both F0+F1 were fed experimental diets (Table S11), suggesting that sex dimorphisms manifest as different magnitudes of the same interaction. Visual inspection of the data led to a similar conclusion (Figure S2). We therefore interpret that any statistical differences reflected relatively modest differences in the present data, suggesting that mothers curse may exert a relatively benign influence in this case.