1   **Hypothalamic perineuronal nets are regulated by sex and dietary interventions**

2   Nan Zhang[1,2,3], Zili Yan[1], Hailan Liu[1], Meng Yu[1], Yang He[1], Hesong Liu[1], Chen Liang[1], Longlong Tu[1], Lina

3   Wang[1], Na Yin[1], Junying Han[1], Nikolas Scarcelli[1], Yongjie Yang[1], Chunmei Wang[1], Tianshu Zeng[2,3], Lu-Lu

4   Chen[2,3], Yong Xu[1, 4*]

5   [1]Children's Nutrition Research Center, Department of Pediatrics, Baylor College of Medicine, Houston,

6   TX, 77030; [2]Department of Endocrinology, Union Hospital, Tongji Medical College, Huazhong University

7   of Science and Technology, Wuhan, China; [3]Hubei Provincial Clinical Research Center for Diabetes and

8   Metabolic Disorder, Wuhan, China; [4]Department of Molecular and Cellular Biology, Baylor College of

9   Medicine, Houston, TX, 77030

10   *To whom correspondence should be addressed:

11   Yong Xu, PhD, MD; 1100 Bates Street #8066, Houston, TX 77030, USA, Mail stop code: MCB320, E-mail:

12   yongx@bcm.edu, Ph: (713) 798 7199, Fax: (713) 798 7187

13

14   **LEAD AUTHOR: YONG XU**

15

17

18

19

ABSTRACT

Perineuronal nets (PNNs) are widely present in the hypothalamus, and are thought to provide physical protection and ion buffering for neurons, and regulate their synaptic plasticity and intracellular signaling. Recent evidence indicates that PNNs in the mediobasal hypothalamus plays an important role in the regulation of glucose homeostasis. However, whether and how hypothalamic PNNs are regulated are not fully understood. In the present study, we examined whether PNNs in various hypothalamic regions in mice can be regulated by sex, gonadal hormones, dietary interventions, or their interactions. We demonstrated that gonadal hormones are required to maintain normal PNNs in the arcuate nucleus of hypothalamus in both male and female mice. In addition, PNNs in the terete hypothalamic nucleus display a sexual dimorphism with females higher than males, and high-fat diet feeding increases terete PNNs only in female mice but not in male mice. On the other hand, PNNs in other hypothalamic regions are not influenced by sex, gonadal hormones or dietary interventions. In summary, we demonstrated that hypothalamic PNNs are regulated in a region-specific manner and these results provide a framework to further investigate the potential functions of PNNs in regulating energy/glucose homeostasis at the interplay of sex, gonadal hormones and diets.

Introduction

Obesity is now recognized as a serious global health problem due to its increasing prevalence and comorbidities, e.g. the metabolic syndrome. The WHO reported that over 650 million adults worldwide were obese in 2016 and 40 million children under the age of 5 were overweight or obese in 2018. In U.S., the prevalence of adult obesity was 42.4% in 2017~2018 according to the Centers for Disease Control and Prevention. The etiology of human obesity is not fully understood, and effective treatments for obesity and associated metabolic disorders are limited. Recent studies revealed genetic and epigenetic basis for variations in human body mass index (BMI) (1-3), and strikingly the majority of BMI-associated genetic variants affect genes that are enriched in the brain (3,4). In particular, the brain hypothalamus receives metabolic and/or hormonal signals reflecting the body's nutritional status and coordinates neuroendocrine and behavioral responses to maintain body weight balance (5-7). Indeed, various genetic variants associated with human obesity have been demonstrated to cause energy and/or glucose dysregulations through impairing functions of neurons and/or neurocircuits within the hypothalamus (8).

Perineuronal nets (PNNs) in the brain are condensed glycosaminoglycan-rich extracellular matrix structures (9). PNNs typically enmesh neurons in defined circuits, and are thought to provide physical protection and ion buffering for neurons, and regulate their synaptic plasticity and intracellular signaling (9). PNNs abundantly exist in the forebrain regions, e.g. the cortex and the hippocampus, and the PNNs levels in these regions can be regulated by animals' experience, and have been implicated in various neurobiological disorders such as schizophrenia, bipolar disorder, Alzheimer's disease and addictions. Recent evidence indicates that chemical disruption of PNNs in the mediobasal hypothalamus significantly blunts the glucose-lowering effects of central action of fibroblast growth factor 1 (FGF1) in

obese Zucker diabetic fatty (ZDF) rats (10). Thus, hypothalamic PNNs may also play important roles in

the regulation of energy and/or glucose homeostasis. Indeed, abundant PNNs are present in multiple

hypothalamic regions (11,12), but whether and how these hypothalamic PNNs are regulated are not fully

understood.

In the present study, we sought to examine whether PNNs in various hypothalamic regions can be

regulated by dietary interventions in mice. Since many hypothalamic regions or neural populations are

sexually dimorphic and/or regulated by gonadal hormones (13,14), we also explored the potential effects

of sex and/or gonadal hormones on PNN levels. Our results revealed that sex, gonadal hormones and/or

dietary interventions can regulate PNNs in a region-specific manner and provide a framework to further

investigate the potential functions of PNNs in regulating energy/glucose homeostasis at the interplay of

sex, gonadal hormones and diets.

METHODS

Study Animals

C57BL6j male and female mice were group-housed respectively until 14 weeks of age in a temperature-

controlled room under a 12h:12h light-dark cycle with ad lib access to regular chow diet (5V5R-Advanced

Protocol PicoLab Select Rodent 50 IF/6F, PicoLab) and water. Forty nine mice are divided into 16 groups

by four factors (1) sex: male or female groups; (2) surgeries: receiving sham or castration

(CAST)/ovariectomy (OVX) surgeries at 14 weeks of age (see below); (3) dietary interventions: chronic

feeding with chow or a high-fat diet (HFD, D12492i Rodent Diet With 60 kcal% Fat, Research Diets, Inc.),

(4) nutritional status: fed or fasted at the time of perfuse (see below). Care of all animals and procedures

were approved by the Baylor College of Medicine Institutional Animal Care and Use Committee.

Surgery

At the age of 14 weeks, either bilateral gonadectomy (CAST for males and OVX for females) or sham

surgeries were executed in these mice. The mice were anesthetized with inhaled 2% isoflurane. For OVX

in females, the dorsolateral incisions (2 cm) were made, followed by blunted separation of the

subcutaneous fat, and incision (1.5cm) the abdominal wall to expose the reproductive tract. The ovaries

were isolated and excised. The reproductive tract was returned to the peritoneum and the incisions on

the abdominal wall were sutured using sterile Vicryl thread (size 6); the skin incisions were sutured by

sterile Prolene thread. For sham surgeries in female mice, the procedures were the same, except that

the ovaries were kept intact. For castration in males, a central vertical incision (3 cm) was made, followed

by blunted separation of any subcutaneous fat to locate the vas deferens. The vas deferens were ligated

with Vicryl; the testes were cut and taken away from the fat pad. The skin incisions were sutured by

sterile Prolene thread. For sham surgeries in male mice, the procedures were the same, except that the

testes were kept intact.

Food intake, body weight, glucose and body composition

One week after the surgeries, mice were randomly divided into two groups to be either regular chow or

HFD described above ad libitum for four weeks. Food intake, body weight and fed glucose were

measured before the surgeries and once each week after the surgery for five weeks. The glucose was

measured using the glucometer via the tail vein; to avoid confounding effects from circadian clock and

feeding behavior on the glucose level, we always measured blood glucose in the morning of the day and

after a 2-hour short fasting to ensure the empty stomach. Body composition (fat mass and lean mass)

was examined with the Bruker minispec mq10 MRS system before the surgeries and 5 weeks afterwards.

Perfusion and WFA staining

Five weeks after the surgeries, mice were further divided into fasted and ad libitum groups before the

perfusion. For the fasted group, mice were fasted overnight and then perfused in the next morning; for

the ad libitum group, both food and water were provided ad libitum at all time before the perfusion. For

the perfusion, mice were anaesthetized with inhaled isoflurane and perfused with saline followed by 10%

formalin. The brain sections were cut at 25 μm and collected into five consecutive series. One series is

used for the following staining and quantification. Free-floating tissue sections were incubated overnight

at room temperature for 22 hours using a 1:500 dilution of biotin-labelled Wisteria floribunda agglutinin

(WFA) (L1516, Sigma-Aldrich) in PBS + 0.25% Triton X-100. The sections were then washed and incubated

for 2 hours at room temperature in 1:500 Streptavidin protein 488 (21832, Invitrogen). Images were

captured using a fluorescence microscope with 1s exposure time and no change of contrast for all

pictures to visualize various hypothalamic nuclei, including the arcuate hypothalamic nucleus (ARH), the

terete hypothalamic nucleus (TE) , the paraventricular hypothalamic nucleus (PVH), lateral

hypothalamus (LH), the anterior hypothalamus (AH), the ventromedial hypothalamic nucleus (VMH), the

dorsomedial hypothalamic nucleus (DMH), and the perifornical area of the anterior hypothalamus

(PeFAH). Fluorescence intensity of PNNs for each image was quantified using an established method by

the macro plugin "Perineuronal net Intensity Program for the Standardization and Quantification of ECM

Analysis" (PIPSQUEAK AI v5.3.9, Rewire Neuro, Inc.) (15).

Statistics

Statistical analysis was done using SPSS (SPSS 22.0.0.0, IBM SPSS Statistics) and GraphPad Prism

(GraphPad Prism 8.4.2, Graphpad software, LLC). Multivariate linear regression in SPSS was used to

investigate the influence of sex, GDX and chronic dietary intervention on various metabolic parameters

(cumulative food intake, body weight change, blood glucose change, fat change and lean change). If sex

or GDX was found to have a significant effect on any parameter, the p value for the interaction of sex

and GDX was tested. If any two factors were found to have significant effects, the p value for the

interaction of these two factors was tested.

For the PNN fluorescence intensity, in addition to the sex, GDX and chronic diet intervention, we

introduced one more factor (nutritional status) as mice were perfused either after an overnight fasting

or ad libitum. Similarly, we first used the multivariate linear regression analyses in SPSS to examine

effects of these 4 factors on PNNs fluorescence intensity in each hypothalamic region. If sex or GDX was

found to have a significant effect on any parameter, we then used the two-way ANOVA to examine the

interaction of sex and GDX. In case that sex had a significant effect, we used the GraphPad Prism to

further examine effects of dietary intervention and nutritional status in each sex separately. P < 0.05 was

considered statistically significant.

Study approval

Care of all animals and procedures were approved by the Baylor College of Medicine Institutional Animal

Care and Use Committee.

Results

Distribution of WFA-labelled PNNs in the mouse hypothalamus

We first surveyed the distribution of PNNs in the mouse hypothalamus using WFA immunostaining. Abundant WFA-labelled PNNs were detected in multiple hypothalamic regions, including the LH, the ARH, the VMH and the TE (Figure 1). In addition, we also observed modest WFA immunoreactivity in the AH, the PVH and the DMH (Figure 1). Consistent with a previous report (12), WFA-labelled PNNs were also observed in a newly identified hypothalamic area, namely the PeFAH (Figure 1A).

Effects of sex, gonadal hormones and dietary interventions on metabolic parameters

Given the essential roles of the hypothalamus in regulating energy homeostasis, we sought to examine potential effects of dietary interventions on the level of PNNs in the hypothalamus. In order to explore potential influence by sex and/or gonadal hormones, we included both male and female mice with or without gonadectomy (GDX) in these analyses. Briefly, male and female C57Bl6j mice underwent sham or GDX (castration for males and ovariectomy for females) at the 14 weeks of age, followed by 5-week chronic feeding with either a regular chow diet or HFD. As expected, the multivariate linear regression analysis revealed that compared to chow-fed groups, HFD feeding had profound effects on multiple metabolic parameters, including significantly increasing calorie intake, blood glucose, body weight gain and fat mass gain (Table 1). Notably, fat mass gain was significantly smaller in females than in males (Table 1). We also noted that lean mass gain was significantly larger in females than in males, and that GDX significantly reduced lean mass gain; in addition, there was a significant interaction between sex and GDX on this parameter (Table 1). In summary, multiple metabolic parameters in mice were influenced by sex, gonadal hormones, the dietary interventions, and/or their interactions, as reported by others (16,17).

Effects of gonadal hormones on PNN in the ARH

At the end of 5-week feeding, mice were then perfused either ad libitum or after an overnight fasting. We then used these fixed brain samples for WFA immunostaining and then quantified the level of WFA-labelled PNNs in each hypothalamic region in various conditions. In the ARH, the WFA fluorescence intensity was not significantly affected by chronic HFD feeding (vs. chow), or by the acute fasting (vs. ad libitum), as revealed by the multivariate linear regression analysis (Table 2). Interestingly, we noted that GDX significantly reduced WFA fluorescence intensity in the ARH compared to the sham group (Table 2 and Figure 2A-2E). A detailed analysis at different rostral-caudal levels revealed that GDX mainly reduced WFA in the rostral to medial ARH (-1.94 to -1.58 mm to Bregma) but the caudal ARH was not affected (Figure 2F). Because female and male mice experienced different GDX surgeries (OVX for females and CAST for males), we used a two-way ANOVA analysis to further analyze the interaction between sex and GDX. We found that the inhibitory effect of GDX was independent of sex; in other words, both OVX in females and CAST in males significantly reduced WFA fluorescence intensity in the ARH (Table 2 and Figure 2G). We also examined effects of dietary innervations on ARH PNNs in sham or in GDX groups, respectively, but did not detect any significant effect of by either chronic HFD feeding or by acute fasting (data not shown). Thus, these results indicate that endogenous gonadal hormones in both males and females are required to maintain normal PNNs in the ARH.

Effects of sex and HFD feeding on PNN in the TE

In the TE, female mice showed significantly higher WFA fluorescence intensity compared to males, as revealed by the multivariate linear regression analysis (Table 3 and Figure 3A-3E), and this sex difference existed in a rostral subdivision (-1.70 mm to Bregma) and a medial-caudal subdivision (-2.18 to -2.30 mm

to Bregma) of the TE (Figure 3F). Using a two-way ANOVA analysis, we further found that the significant

sex difference only existed in sham mice, but not in GDX mice (Figure 3G). Thus, these results indicate

that gonad-intact female mice have higher PNNs in the TE than gonad-intact males.

Given the clear influence by sex, we reanalyzed the effects of HFD feeding in each sex individually and

used a two-way ANOVA analysis followed by Sidak tests to further explore potential interactions

between chronic HFD feeding and acute fasting. In female mice fed ad libitum, chronic HFD feeding

significantly enhanced WFA fluorescence intensity in the TE compared to chow-fed females, but this

effect was not observed after an overnight fasting (Figure 4A-4E). This HFD-induced increases existed in

a rostral-medial subdivision (-1.70 to -1.94 mm to Bregma) of the TE (Figure 4F). On the other hand, in

male mice, neither chronic HFD feeding nor acute fasting induced any changes in WFA fluorescence

intensity in the TE (Figure 4G). Thus, these results indicate that chronic HFD feeding can increase PNNs

in the TE in female mice but not in male mice.

Other hypothalamic regions

We also analyzed effects of sex, gonadal hormones, and/or dietary interventions on WFA fluorescence

intensity in all other hypothalamic regions. The multivariate linear regression analysis revealed that WFA

fluorescence intensity in the PVH, the PeFAH, the LH, the AH, the VMH or the DMH, was not significantly

altered by these conditions (Table 4).

Discussions

One interesting observation of our study is that endogenous gonadal hormones are required to maintain normal PNNs in the ARH in both male and female mice. The ARH in the hypothalamus has been long believed to contain the first order neurons that respond to the peripheral signals, including leptin (18-20), insulin (21-23), ghrelin (24), asprosin (25) and estrogens (26-28). The ARH neurons include those expressing pro-opiomelanocortin (POMC) and those co-releasing agouti-related peptide (AgRP), neuropeptide Y (NPY), and GABA (referred to as AgRP neurons). POMC neurons synthesize and secret an anorexigenic peptide, α-melanocyte-stimulating hormone (α-MSH) which prevents overeating and body weight gain (29,30). On the other hand, AgRP neurons are orexigenic (31), and activation of AgRP neurons promotes eating even when mice are satiated (32,33). Recent evidence also indicates that non-AgRP GABAergic neurons in the ARH provide a redundant orexigenic mechanism to drive feeding and body weight gain (34). Notably, the majority of PNNs in the ARH enmeshes AgRP neurons and GABAergic neurons, with only a small portion enmeshing POMC neurons (11). Interestingly, reduced PNNs in the ARH is found in ob/ob mice deficient in the leptin gene and in ZDF rats with a mutated leptin receptor gene (10,11), highlighting an important role of leptin signaling in maintaining the normal PNNs in the ARH. Further, the central administration of FGF1 in ZDF rats can enhance PNNs in the ARH, associated with a prolonged glucose-lowering benefit (10). Importantly, chemical disruption of PNNs significantly shortens this glucose-lowering effect of FGF1 (10). Thus, PNNs in the ARH play an essential role in regulating glucose balance.

Here we demonstrated that PNNs in the ARH are significantly reduced by removal of the ovaries in female mice, suggesting that PNNs are also regulated by ovarian hormones. Supporting this notion, co-administration of both ovarian hormones (estrogen and progesterone), which simulates pregnancy, can

trigger PNN formation in the medial preoptic area, a hypothalamic region normally devoid of PNNs in

non-pregnant females (35). While the role of progesterone in energy and glucose homeostasis is not

clear (14), estrogen is well established to provide many metabolic benefits, including suppressing food

intake, increasing energy expenditure, reducing body weight gain, enhancing insulin sensitivity, and

lowering blood glucose (16,36-41). Both estrogen receptor α (ERα) and estrogen receptor β (ERβ) are

implicated in mediating these metabolic effects (42-44). Notably, abundant ERα is expressed by POMC

neurons in the ARH (45-47), to a less extent by AgRP neurons (48,49), while ERβ expression in the ARH

is minimal (50). We recently showed that intact ERα signals are required for the glucose-sensing

functions of ARH neurons (51). Further, estrogen can enhance glutamatergic synapses onto POMC

neurons (52), and increase their excitability (53,54). Importantly, we previously showed that selective

deletion of ERα from POMC-lineage neurons in female mice results in hyperphagia and obesity (26), and

attenuates estrogen-induced anorexia (55). Thus, these findings, together with our current observations

that PNNs are reduced by ovariectomy, suggest that PNNs in the ARH may mediate, at least a portion of,

estrogenic actions on energy and glucose homeostasis, and future investigations are warranted to test

this possibility. Interestingly, we also observed similar reductions of PNNs in the ARH of male mice

depleted with endogenous testosterone. Consistently, it has been reported that testosterone

administration can stimulate PNN formation in the forebrain of song birds (56). However, it is worth

noting that testosterone can be converted into estrogen by aromatase (57). Thus, it remains to be tested

whether the testosterone-induced PNN expression is mediated by testosterone per se or by converted

estrogen.

Another hypothalamic region where PNNs are regulated is the TE, where female mice showed higher

PNNs than males. Importantly, this sex difference was only observed in gonad-intact mice, but was

blunted in mice with gonads surgically removed. These results suggest that PNNs in the TE may be also

regulated by gonadal hormones. Supporting this possibility, estrogen has been shown to increase

oxytocin binding in the TE in rats (58), indicating potential actions of estrogen in this region. It has to be

pointed out that, unlike other hypothalamic regions (e.g. the ARH, VMH, etc.), the TE has received very

little attention with no more than 5 publications regarding this nucleus available in the PubMed. Early

neuroanatomic studies reported that cell bodies of cholecystokinin neurons and calcitonin gene-related

peptide (CGRP) neurons are found in the TE (59,60). In addition, the TE harbors nerve fibers that are

immunoreactive for NPY, α-MSH, β-endorphin, corticortropin-releasing hormone, galanin, and

substance P (59,60), suggesting that the TE neurons receive a wide range of innervations and inputs from

other neural populations. Interestingly, we found that in female mice only, PNNs in the TE was

upregulated by chronic HFD feeding, while such effect was not observed after an overnight fasting. It is

worth noting that among many hypothalamic regions expressing PNNs, the TE is the only region where

PNNs can be regulated by dietary interventions. Considering that the TE receives inputs from NPY and

α-MSH, both of which are implicated in the regulation of feeding and body weight, we suggest that TE

neurons and the PNNs enmeshing these neurons may play an under-appreciated role in the context of

energy homeostasis. This possibility warrants future investigations.

In summary, here we systemically assessed PNNs in multiple hypothalamic regions and explored

potential regulations by sex, gonadal hormones, dietary interventions, and their interactions. We

acknowledge that the limitation of the current study is its descriptive nature. However, we want to point

out that our results filled a gap of knowledge in the relevant field and provided a framework to develop

new hypotheses and design future studies to determine the physiological functions of hypothalamic

PNNs and to explore the regulatory mechanisms. For example, the strong regulations by gonadal

hormones on PNNs in the ARH suggest that gonadal hormones may act through PNNs in the ARH to

regulate energy and/or glucose homeostasis. Further, given that PNNs in the ARH are required to

mediates metabolic benefits of FGF1 (10), the efficacy of FGF1 may need to be examined in animals

depleted of gonadal hormones, which would provide important pre-clinical data for potential

applications of FGF1 or its analogs in aged men with testosterone deficiency or in post-menopausal

women. In addition, the dietary regulations on PNNs in the TE suggest that this understudied

hypothalamic nucleus may play a role in the regulation of feeding and body weight balance. Importantly,

since these dietary regulations only existed in female mice but not in male mice, future studies need to

include at least female animals, if not both. Only including male animals (a common practice in many

laboratories) might miss potential discoveries about PNNs in the TE or TE neurons themselves.

Author Contributions: NZ was involved in experimental design and most of procedures, data acquisition

and analyses, and writing the manuscript. ZY assisted in data analysis. HL, MY, YH, HL, CL, LT, LW, NY, JH,

NS, YY and CW assisted in in surgical procedures and production of study mice. LLC and TZ are involved

in the study design and manuscript preparation. YX is the guarantors of this work and, as such, had full

access to all the data in the study and take responsibility for the integrity of the data and the accuracy

of the data analysis.

Acknowledgements: The authors are grateful for the technical assistance provided by QW Zhang, SY

Huang, DH Liu, B Yuan and YR Wang. The authors are grateful for the consultations provided by Dr.

Michael Schwartz and Dr. Kimberly Alonge at University of Washington. The investigators were

supported by grants from the NIH (K01DK119471 to CW), USDA/CRIS (51000-064-01S to YX) and the

American Heart Association (20POST35120600 to YH).

Data availability statement: All data generated or analyzed during this study are included in this

published article.

References

1. Farooqi IS, O'Rahilly S. The Genetics of Obesity in Humans. 2000.

2. Wahl S, Drong A, Lehne B, Loh M, Scott WR, Kunze S, Tsai PC, Ried JS, Zhang W, Yang Y, Tan S, Fiorito G, Franke L, Guarrera S, Kasela S, Kriebel J, Richmond RC, Adamo M, Afzal U, Ala-Korpela M, Albetti B, Ammerpohl O, Apperley JF, Beekman M, Bertazzi PA, Black SL, Blancher C, Bonder MJ, Brosch M, Carstensen-Kirberg M, de Craen AJ, de Lusignan S, Dehghan A, Elkalaawy M, Fischer K, Franco OH, Gaunt TR, Hampe J, Hashemi M, Isaacs A, Jenkinson A, Jha S, Kato N, Krogh V, Laffan M, Meisinger C, Meitinger T, Mok ZY, Motta V, Ng HK, Nikolakopoulou Z, Nteliopoulos G, Panico S, Pervjakova N, Prokisch H, Rathmann W, Roden M, Rota F, Rozario MA, Sandling JK, Schafmayer C, Schramm K, Siebert R, Slagboom PE, Soininen P, Stolk L, Strauch K, Tai ES, Tarantini L, Thorand B, Tigchelaar EF, Tumino R, Uiterlinden AG, van Duijn C, van Meurs JB, Vineis P, Wickremasinghe AR, Wijmenga C, Yang TP, Yuan W, Zhernakova A, Batterham RL, Smith GD, Deloukas P, Heijmans BT, Herder C, Hofman A, Lindgren CM, Milani L, van der Harst P, Peters A, Illig T, Relton CL, Waldenberger M, Jarvelin MR, Bollati V, Soong R, Spector TD, Scott J, McCarthy MI, Elliott P, Bell JT, Matullo G, Gieger C, Kooner JS, Grallert H, Chambers JC. Epigenome-wide association study of body mass index, and the adverse outcomes of adiposity. *Nature*. 2017;541(7635):81-86.

3. Locke AE, Kahali B, Berndt SI, Justice AE, Pers TH, Day FR, Powell C, Vedantam S, Buchkovich ML, Yang J, Croteau-Chonka DC, Esko T, Fall T, Ferreira T, Gustafsson S, Kutalik Z, Luan J, Magi R, Randall JC, Winkler TW, Wood AR, Workalemahu T, Faul JD, Smith JA, Zhao JH, Zhao W, Chen J, Fehrmann R, Hedman AK, Karjalainen J, Schmidt EM, Absher D, Amin N, Anderson D, Beekman M, Bolton JL, Bragg-Gresham JL, Buyske S, Demirkan A, Deng G, Ehret GB, Feenstra B, Feitosa MF, Fischer K, Goel A, Gong J, Jackson AU, Kanoni S, Kleber ME, Kristiansson K, Lim U, Lotay V, Mangino M, Leach IM, Medina-Gomez C, Medland SE, Nalls MA, Palmer CD, Pasko D, Pechlivanis S, Peters MJ, Prokopenko I, Shungin D, Stancakova A, Strawbridge RJ, Sung YJ, Tanaka T, Teumer A, Trompet S, van der Laan SW, van Setten J, Van Vliet-Ostaptchouk JV, Wang Z, Yengo L, Zhang W, Isaacs A, Albrecht E, Arnlov J, Arscott GM, Attwood AP, Bandinelli S, Barrett A, Bas IN, Bellis C, Bennett AJ, Berne C, Blagieva R, Bluher M, Bohringer S, Bonnycastle LL, Bottcher Y, Boyd HA, Bruinenberg M, Caspersen IH, Chen YI, Clarke R, Daw EW, de Craen AJM, Delgado G, Dimitriou M, Doney ASF, Eklund N, Estrada K, Eury E, Folkersen L, Fraser RM, Garcia ME, Geller F, Giedraitis V, Gigante B, Go AS, Golay A, Goodall AH, Gordon SD, Gorski M, Grabe HJ, Grallert H, Grammer TB, Grassler J, Gronberg H, Groves CJ, Gusto G, Haessler J, Hall P, Haller T, Hallmans G, Hartman CA, Hassinen M, Hayward C, Heard-Costa NL, Helmer Q, Hengstenberg C, Holmen O, Hottenga JJ, James AL, Jeff JM, Johansson A, Jolley J, Juliusdottir T, Kinnunen L, Koenig W, Koskenvuo M, Kratzer W, Laitinen J, Lamina C, Leander K, Lee NR, Lichtner P, Lind L, Lindstrom J, Lo KS, Lobbens S, Lorbeer R, Lu Y, Mach F, Magnusson PKE, Mahajan A, McArdle WL, McLachlan S, Menni C, Merger S, Mihailov E, Milani L, Moayyeri A, Monda KL, Morken MA, Mulas A, Muller G, Muller-Nurasyid M, Musk AW, Nagaraja R, Nothen MM, Nolte IM, Pilz S, Rayner NW, Renstrom F, Rettig R, Ried JS, Ripke S, Robertson NR, Rose LM, Sanna S, Scharnagl H, Scholtens S, Schumacher FR, Scott WR, Seufferlein T, Shi J, Smith AV, Smolonska J, Stanton AV, Steinthorsdottir V, Stirrups K, Stringham HM, Sundstrom J, Swertz MA, Swift AJ, Syvanen AC, Tan ST, Tayo BO, Thorand B, Thorleifsson G, Tyrer JP, Uh HW, Vandenput L, Verhulst FC, Vermeulen SH, Verweij N, Vonk JM, Waite LL, Warren HR, Waterworth D, Weedon MN, Wilkens LR, Willenborg C, Wilsgaard T, Wojczynski MK, Wong A, Wright AF, Zhang Q, Brennan EP, Choi M, Dastani Z, Drong AW, Eriksson P, Franco-Cereceda A, Gadin JR, Gharavi AG, Goddard ME, Handsaker RE, Huang J, Karpe F, Kathiresan S, Keildson S, Kiryluk K, Kubo M, Lee JY, Liang L, Lifton RP, Ma B, McCarroll SA, McKnight AJ, Min JL, Moffatt MF, Montgomery GW, Murabito JM, Nicholson

G, Nyholt DR, Okada Y, Perry JRB, Dorajoo R, Reinmaa E, Salem RM, Sandholm N, Scott RA, Stolk L, Takahashi A, van 't Hooft FM, Vinkhuyzen AAE, Westra HJ, Zheng W, Zondervan KT, Heath AC, Arveiler D, Bakker SJL, Beilby J, Bergman RN, Blangero J, Bovet P, Campbell H, Caulfield MJ, Cesana G, Chakravarti A, Chasman DI, Chines PS, Collins FS, Crawford DC, Cupples LA, Cusi D, Danesh J, de Faire U, den Ruijter HM, Dominiczak AF, Erbel R, Erdmann J, Eriksson JG, Farrall M, Felix SB, Ferrannini E, Ferrieres J, Ford I, Forouhi NG, Forrester T, Franco OH, Gansevoort RT, Gejman PV, Gieger C, Gottesman O, Gudnason V, Gyllensten U, Hall AS, Harris TB, Hattersley AT, Hicks AA, Hindorff LA, Hingorani AD, Hofman A, Homuth G, Hovingh GK, Humphries SE, Hunt SC, Hypponen E, Illig T, Jacobs KB, Jarvelin MR, Jockel KH, Johansen B, Jousilahti P, Jukema JW, Jula AM, Kaprio J, Kastelein JJP, Keinanen-Kiukaanniemi SM, Kiemeney LA, Knekt P, Kooner JS, Kooperberg C, Kovacs P, Kraja AT, Kumari M, Kuusisto J, Lakka TA, Langenberg C, Marchand LL, Lehtimaki T, Lyssenko V, Mannisto S, Marette A, Matise TC, McKenzie CA, McKnight B, Moll FL, Morris AD, Morris AP, Murray JC, Nelis M, Ohlsson C, Oldehinkel AJ, Ong KK, Madden PAF, Pasterkamp G, Peden JF, Peters A, Postma DS, Pramstaller PP, Price JF, Qi L, Raitakari OT, Rankinen T, Rao DC, Rice TK, Ridker PM, Rioux JD, Ritchie MD, Rudan I, Salomaa V, Samani NJ, Saramies J, Sarzynski MA, Schunkert H, Schwarz PEH, Sever P, Shuldiner AR, Sinisalo J, Stolk RP, Strauch K, Tonjes A, Tregouet DA, Tremblay A, Tremoli E, Virtamo J, Vohl MC, Volker U, Waeber G, Willemsen G, Witteman JC, Zillikens MC, Adair LS, Amouyel P, Asselbergs FW, Assimes TL, Bochud M, Boehm BO, Boerwinkle E, Bornstein SR, Bottinger EP, Bouchard C, Cauchi S, Chambers JC, Chanock SJ, Cooper RS, de Bakker PIW, Dedoussis G, Ferrucci L, Franks PW, Froguel P, Groop LC, Haiman CA, Hamsten A, Hui J, Hunter DJ, Hveem K, Kaplan RC, Kivimaki M, Kuh D, Laakso M, Liu Y, Martin NG, Marz W, Melbye M, Metspalu A, Moebus S, Munroe PB, Njolstad I, Oostra BA, Palmer CNA, Pedersen NL, Perola M, Perusse L, Peters U, Power C, Quertermous T, Rauramaa R, Rivadeneira F, Saaristo TE, Saleheen D, Sattar N, Schadt EE, Schlessinger D, Slagboom PE, Snieder H, Spector TD, Thorsteinsdottir U, Stumvoll M, Tuomilehto J, Uitterlinden AG, Uusitupa M, van der Harst P, Walker M, Wallaschofski H, Wareham NJ, Watkins H, Weir DR, Wichmann HE, Wilson JF, Zanen P, Borecki IB, Deloukas P, Fox CS, Heid IM, O'Connell JR, Strachan DP, Stefansson K, van Duijn CM, Abecasis GR, Franke L, Frayling TM, McCarthy MI, Visscher PM, Scherag A, Willer CJ, Boehnke M, Mohlke KL, Lindgren CM, Beckmann JS, Barroso I, North KE, Ingelsson E, Hirschhorn JN, Loos RJF, Speliotes EK. Genetic studies of body mass index yield new insights for obesity biology. *Nature*. 2015;518(7538):197-206.

4. Turcot V, Lu Y, Highland HM, Schurmann C, Justice AE, Fine RS, Bradfield JP, Esko T, Giri A, Graff M, Guo X, Hendricks AE, Karaderi T, Lempradl A, Locke AE, Mahajan A, Marouli E, Sivapalaratnam S, Young KL, Alfred T, Feitosa MF, Masca NGD, Manning AK, Medina-Gomez C, Mudgal P, Ng MCY, Reiner AP, Vedantam S, Willems SM, Winkler TW, Abecasis G, Aben KK, Alam DS, Alharthi SE, Allison M, Amouyel P, Asselbergs FW, Auer PL, Balkau B, Bang LE, Barroso I, Bastarache L, Benn M, Bergmann S, Bielak LF, Blüher M, Boehnke M, Boeing H, Boerwinkle E, Böger CA, Bork-Jensen J, Bots ML, Bottinger EP, Bowden DW, Brandslund I, Breen G, Brilliant MH, Broer L, Brumat M, Burt AA, Butterworth AS, Campbell PT, Cappellani S, Carey DJ, Catamo E, Caulfield MJ, Chambers JC, Chasman DI, Chen YI, Chowdhury R, Christensen C, Chu AY, Cocca M, Collins FS, Cook JP, Corley J, Corominas Galbany J, Cox AJ, Crosslin DS, Cuellar-Partida G, D'Eustacchio A, Danesh J, Davies G, Bakker PIW, Groot MCH, Mutsert R, Deary IJ, Dedoussis G, Demerath EW, Heijer M, Hollander AI, Ruijter HM, Dennis JG, Denny JC, Di Angelantonio E, Drenos F, Du M, Dubé MP, Dunning AM, Easton DF, Edwards TL, Ellinghaus D, Ellinor PT, Elliott P, Evangelou E, Farmaki AE, Farooqi IS, Faul JD, Fauser S, Feng S, Ferrannini E, Ferrieres J, Florez JC, Ford I, Fornage M, Franco OH, Franke A, Franks PW, Friedrich N, Frikke-Schmidt R, Galesloot TE, Gan W, Gandin I, Gasparini P, Gibson J, Giedraitis V, Gjesing AP, Gordon-Larsen P, Gorski M, Grabe HJ, Grant SFA, Grarup N, Griffiths

HL, Grove ML, Gudnason V, Gustafsson S, Haessler J, Hakonarson H, Hammerschlag AR, Hansen T, Harris KM, Harris TB, Hattersley AT, Have CT, Hayward C, He L, Heard-Costa NL, Heath AC, Heid IM, Helgeland Ø, Hernesniemi J, Hewitt AW, Holmen OL, Hovingh GK, Howson JMM, Hu Y, Huang PL, Huffman JE, Ikram MA, Ingelsson E, Jackson AU, Jansson JH, Jarvik GP, Jensen GB, Jia Y, Johansson S, Jørgensen ME, Jørgensen T, Jukema JW, Kahali B, Kahn RS, Kähönen M, Kamstrup PR, Kanoni S, Kaprio J, Karaleftheri M, Kardia SLR, Karpe F, Kathiresan S, Kee F, Kiemeney LA, Kim E, Kitajima H, Komulainen P, Kooner JS, Kooperberg C, Korhonen T, Kovacs P, Kuivaniemi H, Kutalik Z, Kuulasmaa K, Kuusisto J, Laakso M, Lakka TA, Lamparter D, Lange EM, Lange LA, Langenberg C, Larson EB, Lee NR, Lehtimäki T, Lewis CE, Li H, Li J, Li-Gao R, Lin H, Lin KH, Lin LA, Lin X, Lind L, Lindström J, Linneberg A, Liu CT, Liu DJ, Liu Y, Lo KS, Lophatananon A, Lotery AJ, Loukola A, Luan J, Lubitz SA, Lyytikäinen LP, Männistö S, Marenne G, Mazul AL, McCarthy MI, McKean-Cowdin R, Medland SE, Meidtner K, Milani L, Mistry V, Mitchell P, Mohlke KL, Moilanen L, Moitry M, Montgomery GW, Mook-Kanamori DO, Moore C, Mori TA, Morris AD, Morris AP, Müller-Nurasyid M, Munroe PB, Nalls MA, Narisu N, Nelson CP, Neville M, Nielsen SF, Nikus K, Njølstad PR, Nordestgaard BG, Nyholt DR, O'Connel JR, O'Donoghue ML, Olde Loohuis LM, Ophoff RA, Owen KR, Packard CJ, Padmanabhan S, Palmer CNA, Palmer ND, Pasterkamp G, Patel AP, Pattie A, Pedersen O, Peissig PL, Peloso GM, Pennell CE, Perola M, Perry JA, Perry JRB, Pers TH, Person TN, Peters A, Petersen ERB, Peyser PA, Pirie A, Polasek O, Polderman TJ, Puolijoki H, Raitakari OT, Rasheed A, Rauramaa R, Reilly DF, Renström F, Rheinberger M, Ridker PM, Rioux JD, Rivas MA, Roberts DJ, Robertson NR, Robino A, Rolandsson O, Rudan I, Ruth KS, Saleheen D, Salomaa V, Samani NJ, Sapkota Y, Sattar N, Schoen RE, Schreiner PJ, Schulze MB, Scott RA, Segura-Lepe MP, Shah SH, Sheu WH, Sim X, Slater AJ, Small KS, Smith AV, Southam L, Spector TD, Speliotes EK, Starr JM, Stefansson K, Steinthorsdottir V, Stirrups KE, Strauch K, Stringham HM, Stumvoll M, Sun L, Surendran P, Swift AJ, Tada H, Tansey KE, Tardif JC, Taylor KD, Teumer A, Thompson DJ, Thorleifsson G, Thorsteinsdottir U, Thuesen BH, Tönjes A, Tromp G, Trompet S, Tsafantakis E, Tuomilehto J, Tybjaerg-Hansen A, Tyrer JP, Uher R, Uitterlinden AG, Uusitupa M, Laan SW, Duijn CM, Leeuwen N, van Setten J, Vanhala M, Varbo A, Varga TV, Varma R, Velez Edwards DR, Vermeulen SH, Veronesi G, Vestergaard H, Vitart V, Vogt TF, Völker U, Vuckovic D, Wagenknecht LE, Walker M, Wallentin L, Wang F, Wang CA, Wang S, Wang Y, Ware EB, Wareham NJ, Warren HR, Waterworth DM, Wessel J, White HD, Willer CJ, Wilson JG, Witte DR, Wood AR, Wu Y, Yaghootkar H, Yao J, Yao P, Yerges-Armstrong LM, Young R, Zeggini E, Zhan X, Zhang W, Zhao JH, Zhao W, Zhao W, Zhou W, Zondervan KT, Rotter JI, Pospisilik JA, Rivadeneira F, Borecki IB, Deloukas P, Frayling TM, Lettre G, North KE, Lindgren CM, Hirschhorn JN, Loos RJF. Protein-altering variants associated with body mass index implicate pathways that control energy intake and expenditure in obesity. *Nat Genet*. 2018;50(1):26-41.

5.   Hetherington AW, Ranson SW. Hypothalamic lesions and adiposity in the rat. *The Anatomical Record*. 1940;78:149-172.

6.   Elmquist JK, Elias CF, Saper CB. From lesions to leptin: hypothalamic control of food intake and body weight. *Neuron*. 1999;22(2):221-232.

7.   Morton GJ, Cummings DE, Baskin DG, Barsh GS, Schwartz MW. Central nervous system control of food intake and body weight. *Nature*. 2006;443(7109):289-295.

8.   Yang Y, Xu Y. The central melanocortin system and human obesity. *J Mol Cell Biol*. 2021;12(10):785-797.

9.   Testa D, Prochiantz A, Di Nardo AA. Perineuronal nets in brain physiology and disease. *Semin Cell Dev Biol*. 2019;89:125-135.

10.   Alonge KM, Mirzadeh Z, Scarlett JM, Logsdon AF, Brown JM, Cabrales E, Chan CK, Kaiyala KJ, Bentsen MA, Banks WA, Guttman M, Wight TN, Morton GJ, Schwartz MW. Hypothalamic

perineuronal net assembly is required for sustained diabetes remission induced by fibroblast
growth factor 1 in rats. *Nat Metab*. 2020;2(10):1025-1033.

11. Mirzadeh Z, Alonge KM, Cabrales E, Herranz-Perez V, Scarlett JM, Brown JM, Hassouna R,
Matsen ME, Nguyen HT, Garcia-Verdugo JM, Zeltser LM, Schwartz MW. Perineuronal Net
Formation during the Critical Period for Neuronal Maturation in the Hypothalamic Arcuate
Nucleus. *Nat Metab*. 2019;1(2):212-221.

12. Horii-Hayashi N, Sasagawa T, Hashimoto T, Kaneko T, Takeuchi K, Nishi M. A newly identified
mouse hypothalamic area having bidirectional neural connections with the lateral septum: the
perifornical area of the anterior hypothalamus rich in chondroitin sulfate proteoglycans. *Eur J
Neurosci*. 2015;42(6):2322-2334.

13. Kammel LG, Correa SM. Selective sexual differentiation of neurone populations may contribute
to sex-specific outputs of the ventromedial nucleus of the hypothalamus. *J Neuroendocrinol*.
2020;32(1):e12801.

14. Wang C, Xu Y. Mechanisms for Sex Differences in Energy Homeostasis. *J Mol Endocrinol*.
2019;62(2):R129-R143.

15. Slaker ML, Harkness JH, Sorg BA. A standardized and automated method of perineuronal net
analysis using Wisteria floribunda agglutinin staining intensity. *IBRO Rep*. 2016;1:54-60.

16. Riant E, Waget A, Cogo H, Arnal JF, Burcelin R, Gourdy P. Estrogens protect against high-fat
diet-induced insulin resistance and glucose intolerance in mice. *Endocrinology*.
2009;150(5):2109-2117.

17. Moverare-Skrtic S, Venken K, Andersson N, Lindberg MK, Svensson J, Swanson C,
Vanderschueren D, Oscarsson J, Gustafsson JA, Ohlsson C. Dihydrotestosterone treatment results
in obesity and altered lipid metabolism in orchidectomized mice. *Obesity (Silver Spring)*.
2006;14(4):662-672.

18. Balthasar N, Coppari R, McMinn J, Liu SM, Lee CE, Tang V, Kenny CD, McGovern RA, Chua
SC, Jr., Elmquist JK, Lowell BB. Leptin receptor signaling in POMC neurons is required for
normal body weight homeostasis. *Neuron*. 2004;42(6):983-991.

19. Caron A, Dungan Lemko HM, Castorena CM, Fujikawa T, Lee S, Lord CC, Ahmed N, Lee CE,
Holland WL, Liu C, Elmquist JK. POMC neurons expressing leptin receptors coordinate metabolic
responses to fasting via suppression of leptin levels. *Elife*. 2018;7.

20. Xu J, Bartolome CL, Low CS, Yi X, Chien CH, Wang P, Kong D. Genetic identification of leptin
neural circuits in energy and glucose homeostases. *Nature*. 2018;556(7702):505-509.

21. Qiu J, Zhang C, Borgquist A, Nestor CC, Smith AW, Bosch MA, Ku S, Wagner EJ, Ronnekleiv
OK, Kelly MJ. Insulin excites anorexigenic proopiomelanocortin neurons via activation of
canonical transient receptor potential channels. *Cell Metab*. 2014;19(4):682-693.

22. Konner AC, Janoschek R, Plum L, Jordan SD, Rother E, Ma X, Xu C, Enriori P, Hampel B, Barsh
GS, Kahn CR, Cowley MA, Ashcroft FM, Bruning JC. Insulin action in AgRP-expressing neurons
is required for suppression of hepatic glucose production. *Cell Metab*. 2007;5(6):438-449.

23. Hill JW, Elias CF, Fukuda M, Williams KW, Berglund ED, Holland WL, Cho YR, Chuang JC,
Xu Y, Choi M, Lauzon D, Lee CE, Coppari R, Richardson JA, Zigman JM, Chua S, Scherer PE,
Lowell BB, Bruning JC, Elmquist JK. Direct insulin and leptin action on pro-opiomelanocortin
neurons is required for normal glucose homeostasis and fertility. *Cell Metab*. 2010;11(4):286-297.

24. Wang Q, Liu C, Uchida A, Chuang JC, Walker A, Liu T, Osborne-Lawrence S, Mason BL, Mosher
C, Berglund ED, Elmquist JK, Zigman JM. Arcuate AgRP neurons mediate orexigenic and
glucoregulatory actions of ghrelin. *Mol Metab*. 2014;3(1):64-72.

25. Duerrschmid C, He Y, Wang C, Li C, Bournat JC, Romere C, Saha PK, Lee ME, Phillips KJ, Jain
M, Jia P, Zhao Z, Farias M, Wu Q, Milewicz DM, Sutton VR, Moore DD, Butte NF, Krashes MJ,
Xu Y, Chopra AR. Asprosin is a centrally acting orexigenic hormone. *Nat Med*. 2017;23(12):1444-
1453.

26. Xu Y, Nedungadi TP, Zhu L, Sobhani N, Irani BG, Davis KE, Zhang X, Zou F, Gent LM, Hahner
LD, Khan SA, Elias CF, Elmquist JK, Clegg DJ. Distinct hypothalamic neurons mediate estrogenic
effects on energy homeostasis and reproduction. *Cell Metab*. 2011;14(4):453-465.
27. Herber CB, Krause WC, Wang L, Bayrer JR, Li A, Schmitz M, Fields A, Ford B, Zhang Z, Reid
MS, Nomura DK, Nissenson RA, Correa SM, Ingraham HA. Estrogen signaling in arcuate Kiss1
neurons suppresses a sex-dependent female circuit promoting dense strong bones. *Nat Commun*.
2019;10(1):163.
28. Roepke TA, Malyala A, Bosch MA, Kelly MJ, Ronnekleiv OK. Estrogen regulation of genes
important for K+ channel signaling in the arcuate nucleus. *Endocrinology*. 2007;148(10):4937-
4951.
29. Xu AW, Kaelin CB, Morton GJ, Ogimoto K, Stanhope K, Graham J, Baskin DG, Havel P,
Schwartz MW, Barsh GS. Effects of hypothalamic neurodegeneration on energy balance. *PLoS
Biol*. 2005;3(12):e415.
30. Yaswen L, Diehl N, Brennan MB, Hochgeschwender U. Obesity in the mouse model of pro-
opiomelanocortin deficiency responds to peripheral melanocortin. *Nat Med*. 1999;5(9):1066-1070.
31. Wu Q, Boyle MP, Palmiter RD. Loss of GABAergic signaling by AgRP neurons to the
parabrachial nucleus leads to starvation. *Cell*. 2009;137(7):1225-1234.
32. Krashes MJ, Koda S, Ye C, Rogan SC, Adams AC, Cusher DS, Maratos-Flier E, Roth BL, Lowell
BB. Rapid, reversible activation of AgRP neurons drives feeding behavior in mice. *J Clin Invest*.
2011.
33. Aponte Y, Atasoy D, Sternson SM. AGRP neurons are sufficient to orchestrate feeding behavior
rapidly and without training. *Nat Neurosci*. 2011;14(3):351-355.
34. Zhu C, Jiang Z, Xu Y, Cai ZL, Jiang Q, Xue M, Arenkiel BR, Wu Q, Shu G, Tong Q. Profound
and redundant functions of arcuate neurons in obesity development. *Nat Metab*. 2020;2(8):763-
774.
35. Uriarte N, Ferreno M, Mendez D, Nogueira J. Reorganization of perineuronal nets in the medial
Preoptic Area during the reproductive cycle in female rats. *Sci Rep*. 2020;10(1):5479.
36. Blaustein JD, Wade GN. Ovarian influences on the meal patterns of female rats. *Physiol Behav*.
1976;17(2):201-208.
37. Drewett RF. Sexual behaviour and sexual motivation in the female rat. *Nature*.
1973;242(5398):476-477.
38. Geary N, Asarian L, Korach KS, Pfaff DW, Ogawa S. Deficits in E2-dependent control of feeding,
weight gain, and cholecystokinin satiation in ER-alpha null mice. *Endocrinology*.
2001;142(11):4751-4757.
39. Wallen WJ, Belanger MP, Wittnich C. Sex hormones and the selective estrogen receptor
modulator tamoxifen modulate weekly body weights and food intakes in adolescent and adult rats.
J Nutr. 2001;131(9):2351-2357.
40. Paladini CA, Roeper J. Generating bursts (and pauses) in the dopamine midbrain neurons.
Neuroscience. 2014;282:109-121.
41. Rogers NH, Perfield JW, 2nd, Strissel KJ, Obin MS, Greenberg AS. Reduced energy expenditure
and increased inflammation are early events in the development of ovariectomy-induced obesity.
Endocrinology. 2009.
42. Okura T, Koda M, Ando F, Niino N, Ohta S, Shimokata H. Association of polymorphisms in the
estrogen receptor alpha gene with body fat distribution. *Int J Obes Relat Metab Disord*.
2003;27(9):1020-1027.
43. Heine PA, Taylor JA, Iwamoto GA, Lubahn DB, Cooke PS. Increased adipose tissue in male and
female estrogen receptor-alpha knockout mice. *Proc Natl Acad Sci U S A*. 2000;97(23):12729-
12734.

44. Foryst-Ludwig A, Clemenz M, Hohmann S, Hartge M, Sprang C, Frost N, Krikov M, Bhanot S,
Barros R, Morani A, Gustafsson JA, Unger T, Kintscher U. Metabolic actions of estrogen receptor
beta (ERbeta) are mediated by a negative cross-talk with PPARgamma. *PLoS Genet*.
2008;4(6):e1000108.

45. Miller MM, Tousignant P, Yang U, Pedvis S, Billiar RB. Effects of age and long-term ovariectomy
on the estrogen-receptor containing subpopulations of beta-endorphin-immunoreactive neurons in
the arcuate nucleus of female C57BL/6J mice. *Neuroendocrinology*. 1995;61(5):542-551.

46. Attia E. Anorexia nervosa: current status and future directions. *Annu Rev Med*. 2010;61:425-435.

47. de Souza FS, Nasif S, Lopez-Leal R, Levi DH, Low MJ, Rubinsten M. The estrogen receptor alpha
colocalizes with proopiomelanocortin in hypothalamic neurons and binds to a conserved motif
present in the neuron-specific enhancer nPE2. *Eur J Pharmacol*. 2011;660(1):181-187.

48. Roepke TA, Qiu J, Smith AW, Ronnekleiv OK, Kelly MJ. Fasting and 17beta-estradiol
differentially modulate the M-current in neuropeptide Y neurons. *J Neurosci*. 2011;31(33):11825-
11835.

49. Olofsson LE, Pierce AA, Xu AW. Functional requirement of AgRP and NPY neurons in ovarian
cycle-dependent regulation of food intake. *Proc Natl Acad Sci U S A*. 2009;106(37):15932-15937.

50. Merchenthaler I, Lane MV, Numan S, Dellovade TL. Distribution of estrogen receptor alpha and
beta in the mouse central nervous system: in vivo autoradiographic and immunocytochemical
analyses. *J Comp Neurol*. 2004;473(2):270-291.

51. Yu K, He Y, Hyseni I, Pei Z, Yang Y, Xu P, Cai X, Liu H, Qu N, Yu M, Liang C, Yang T, Wang
J, Gourdy P, Arnal JF, Lenfant F, Xu Y, Wang C. 17beta-estradiol promotes acute refeeding in
hungry mice via membrane-initiated ERalpha signaling. *Mol Metab*. 2020;42:101053.

52. Gao Q, Mezei G, Nie Y, Rao Y, Choi CS, Bechmann I, Leranth C, Toran-Allerand D, Priest CA,
Roberts JL, Gao XB, Mobbs C, Shulman GI, Diano S, Horvath TL. Anorectic estrogen mimics
leptin's effect on the rewiring of melanocortin cells and Stat3 signaling in obese animals. *Nat Med*.
2007;13(1):89-94.

53. Malyala A, Zhang C, Bryant DN, Kelly MJ, Ronnekleiv OK. PI3K signaling effects in
hypothalamic neurons mediated by estrogen. *J Comp Neurol*. 2008;506(6):895-911.

54. Saito K, Cao X, He Y, Xu Y. Progress in the molecular understanding of central regulation of body
weight by estrogens. *Obesity (Silver Spring)*. 2015;23(5):919-926.

55. Zhu L, Xu P, Cao X, Yang Y, Hinton AO, Jr., Xia Y, Saito K, Yan X, Zou F, Ding H, Wang C,
Yan C, Saha P, Khan SA, Zhao J, Fukuda M, Tong Q, Clegg DJ, Chan L, Xu Y. The ERalpha-
PI3K Cascade in Proopiomelanocortin Progenitor Neurons Regulates Feeding and Glucose
Balance in Female Mice. *Endocrinology*. 2015;156(12):4474-4491.

56. Cornez G, Shevchouk OT, Ghorbanpoor S, Ball GF, Cornil CA, Balthazart J. Testosterone
stimulates perineuronal nets development around parvalbumin cells in the adult canary brain in
parallel with song crystallization. *Horm Behav*. 2020;119:104643.

57. Henderson KM, Swanston IA. Androgen aromatization by luteinized bovine granulosa cells in
tissue culture. *J Reprod Fertil*. 1978;52(1):131-134.

58. Kremarik P, Freund-Mercier MJ, Stoeckel ME. Estrogen-sensitive oxytocin binding sites are
differently regulated by progesterone in the telencephalon and the hypothalamus of the rat. *J
Neuroendocrinol*. 1995;7(4):281-289.

59. Lantos TA, Gorcs TJ, Palkovits M. Immunohistochemical localization of calcitonin gene-related
peptide in the terete nucleus of the rat hypothalamus. *Neurobiology (Bp)*. 1996;4(1-2):73-84.

60. Lantos TA, Gorcs TJ, Palkovits M. Immunohistochemical mapping of neuropeptides in the
premamillary region of the hypothalamus in rats. *Brain Res Brain Res Rev*. 1995;20(2):209-249.

598
599

Tables

Table 1. Effects of sex, GDX and diet on various metabolic parameters.

	B	p value	95.0% Confidence Interval	
			Lower Bound	Upper Bound
Cumulative intake (calorie)				
Female vs Male[#]	-13.234	0.298	-38.542	12.075
GDX vs Sham[##]	7.461	0.554	-17.798	32.721
High fat diet vs Regular chow[###]	56.660	p < 0.001***	31.306	82.013
Body weight change (g)				
Female vs Male[#]	-0.448	0.620	-2.255	1.359
GDX vs Sham[##]	0.532	0.556	-1.275	2.339
High fat diet vs Regular chow[###]	5.389	p < 0.001***	3.582	7.196
Blood glucose change (mg/dL)				
Female vs Male[#]	13.141	0.128	-3.915	30.198
GDX vs Sham[##]	1.143	0.893	-15.913	18.200
High fat diet vs Regular chow[###]	37.838	p < 0.001***	20.782	54.894
Fat change (g)				
Female vs Male[#]	-2.052	0.012*	-3.633	-0.471
GDX vs Sham[##]	1.495	0.063	-0.085	3.076
P for the interaction of Sex and GDX		0.451		
High fat diet vs Regular Chow[###]	5.539	p < 0.001***	3.958	7.120
P for the interaction of Sex and Diet		0.226		
Lean change (g)				
Female vs Male[#]	1.405	p < 0.001***	0.828	1.983
GDX vs Sham[##]	-0.805	0.007**	-1.383	-0.228
P for the interaction of Sex and GDX		p < 0.001***		
High fat diet vs Regular chow[###]	-0.107	0.710	-0.684	0.470

Here we use male mice as reference. ## Here we use sham mice as reference. ### Here we use regular chow as reference.

Table 2. Effects of sex, GDX, diet and fasting on PNNs in the ARH.

	B	p value	95.0% Confidence Interval	
			Lower Bound	Upper Bound
Female vs Male#	3216.040	0.710	-14091.082	20523.162
GDX vs Sham##	-28525.814	0.002**	-45832.935	-11218.692
High fat diet vs Regular chow###	5478.986	0.527	-11858.261	22816.234
Fast vs Ab libitum####	1070.635	0.902	-16266.612	18407.883

Here we use male mice as reference. ## Here we use sham mice as reference. ### Here we use

regular chow as reference. ####Here we use fed mice as reference.

613 **Table 3. Effects of sex, GDX, diet and nutritional status on PNNs in the TE.**

	B	p value	95.0% Confidence Interval	
			Lower Bound	Upper Bound
Female vs Male[#]	37265.520	0.004**	12278.584	62252.456
GDX vs Sham[##]	8878.365	0.478	-16108.572	33865.301
High fat diet vs Regular chow[###]	3443.149	0.783	-21587.281	28473.579
Fast vs Ab libitum[####]	-2328.269	0.852	-27358.698	22702.161

614 # Here we use male mice as reference. ## Here we use sham mice as reference. ### Here we use regular

615 chow as reference. ####Here we use fed mice as reference.

618 **Table 4. Effects of sex, GDX, diet and nutritional status on PNNs in the PVH, PeFAH, LH, AH, VMH and**

619 **DMH.**

	B	p value	95.0% Confidence Interval	
			Lower Bound	Upper Bound
PVH				
Female vs Male[#]	1093.487	0.785	-6950.901	9137.874
GDX vs Sham[##]	1801.150	0.649	-6123.285	9725.586
High fat diet vs Regular chow[###]	7111.050	0.082	-948.109	15170.209
Fast vs Ab libitum[####]	1437.498	0.721	-6621.661	9496.657
PeFAH				
Female vs Male[#]	10127.575	0.447	-16501.677	36756.827
GDX vs Sham[##]	12003.295	0.361	-14228.883	38235.473
High fat diet vs Regular chow[###]	12968.496	0.332	-13709.655	39646.646
Fast vs Ab libitum[####]	3879.818	0.771	-22798.333	30557.969
LH				
Female vs Male[#]	48723.960	0.118	-12839.004	110286.924
GDX vs Sham[##]	25737.089	0.397	-34907.897	86382.074
High fat diet vs Regular chow[###]	-22043.519	0.475	-83719.530	39632.491
Fast vs Ab libitum[####]	-17950.520	0.560	-79626.531	43725.490
AH				
Female vs Male[#]	9830.509	0.347	-11017.746	30678.764
GDX vs Sham[##]	6991.586	0.496	-13545.796	27528.969
High fat diet vs Regular chow[###]	6079.521	0.560	-14807.017	26966.059
Fast vs Ab libitum[####]	599.040	0.954	-20287.498	21485.578
VMH				
Female vs Male[#]	8833.935	0.639	-28921.823	46589.693
GDX vs Sham[##]	6705.915	0.718	-30486.859	43898.688

High fat diet vs Regular chow[###]	-14866.204	0.432	-52691.292	22958.884
Fast vs Ab libitum[####]	-19095.426	0.314	-56920.514	18729.662
DMH				
Female vs Male[#]	-2075.195	0.135	-4826.375	675.985
GDX vs Sham[##]	267.179	0.843	-2442.978	2977.336
High fat diet vs Regular chow[###]	1587.169	0.252	-1169.064	4343.401
Fast vs Ab libitum[####]	-487.298	0.723	-3243.531	2268.934

Here we use male mice as reference. ## Here we use sham mice as reference. ### Here we use

regular chow as reference. ####Here we use fed mice as reference.

Figure Legends

Figure 1. Distribution of WFA-labelled PNNs in mouse hypothalamus. Representative fluorescent

microscopic images showing WFA-labelled PNNs (green) and DAPI counterstaining (purple) in a series of

coronal brain sections at the level of -0.82 mm (A), -1.34 mm (B), -1.58 mm (C) and -1.82 mm (D) relative

to the Bregma. Scale bars = 200 μm. AH, the anterior hypothalamus; ARH, the arcuate hypothalamic

nucleus; DMH, the dorsomedial hypothalamic nucleus; LH, the lateral hypothalamus; PeFAH, the

perifornical area of the anterior hypothalamus (PeFAH); PVH, the paraventricular hypothalamic nucleus;

TE, the terete hypothalamic nucleus; VMH, the ventromedial hypothalamic nucleus.

Figure 2. GDX reduces PNNs in the ARH. (A-D) Representative fluorescence microscopic images showing

WFA-labelled PNNs (green) in the ARH of female sham (A), female OVX (B), male sham (C) and male CAST

mice (D). Scale bars = 200 μm. ARH, the arcuate hypothalamic nucleus. (E) Quantification of the total

WFA fluorescence intensity in the ARH from sham vs. GDX mice. N = 24 or 25 mice per group. Data are

presented with mean ± SEM with individual data points. **, $P<0.01$ in unpaired two-tailed Student's t-

tests. (F) Quantification of the WFA fluorescence intensity at various rostral-caudal levels of the ARH

from sham vs. GDX mice. N = 24 or 25 mice per group. Data are presented with mean ± SEM. *, $P<0.05$

and **, $P<0.01$ in unpaired two-tailed Student's t-tests at each level. (G) Quantification of the total WFA

fluorescence intensity in the ARH from female or male mice with sham or GDX surgeries. N = 12 or 13

mice per group. Data are presented with mean ± SEM with individual data points. *, $P<0.05$ in two-way

ANOVA analysis followed by Holm-Sidak post hoc tests.

Figure 3. PNNs in the TE display a sexual dimorphism. (A-D) Representative fluorescence microscopic images showing WFA-labelled PNNs (green) in the TE of female sham (A), male sham (B), female OVX (C), and male CAST mice (D). Scale bars = 200 µm. TE, the terete hypothalamic nucleus. (E) Quantification of the total WFA fluorescence intensity in the TE from female vs. male mice. N = 24 or 25 mice per group. Data are presented with mean ± SEM with individual data points. **, $P<0.01$ in unpaired two-tailed Student's t-tests. (F) Quantification of the WFA fluorescence intensity at various rostral-caudal levels of the TE from female vs. male mice. N = 24 or 25 mice per group. Data are presented with mean ± SEM. *, $P<0.05$ and **, $P<0.01$ in unpaired two-tailed Student's t-tests at each level. (G) Quantification of the total WFA fluorescence intensity in the TE from female or male mice with sham or GDX surgeries. N = 12 or 13 mice per group. Data are presented with mean ± SEM with individual data points. **, $P<0.01$ in two way ANOVA analysis followed by Sidak post hoc tests.

Figure 4. HFD feeding increases TE PNNs in female mice. (A-D) Representative fluorescence microscopic images showing WFA-labelled PNNs (green) in the TE of female mice that were fed chow ad libitum (A), HFD ad libitum sham (B), chow after an overnight fasting (C), and HFD after an overnight fasting (D). Scale bars = 200 µm. TE, the terete hypothalamic nucleus. (E) Quantification of the total WFA fluorescence intensity in the TE from female mice with various dietary interventions. N = 6 mice per group. Data are presented with mean ± SEM with individual data points. *, $P<0.05$ in two-way ANOVA analysis followed by Sidak post hoc tests. (F) Quantification of the WFA fluorescence intensity at various rostral-caudal levels of the TE from female mice fed with chow or HFD ad libitum. N = 6 mice per group. Data are presented with mean ± SEM. *, $P<0.05$ and **, $P<0.01$ in unpaired two-tailed Student's t-tests at each level. (G) Quantification of the total WFA fluorescence intensity in the TE from male mice with

667   various dietary interventions. N = 6 or 7 mice per group. Data are presented with mean ± SEM with

668   individual data points. No significance was detected by in two way ANOVA analysis.

Figure 1

Figure 2

Figure 3

Figure 4