In search of an honest butterfly: sexually selected wing coloration and

reproductive traits from wild populations of the cabbage white butterfly

Anne E. Espeset[1] and Matthew L. Forister[1,2]

[1] Biology Department, Ecology, Evolution, and Conservation Biology Program, University of

Nevada, Reno, NV 89557 USA

[2] Corresponding Author: forister@gmail.com

Running Title: butterfly wings and honest signals

Abstract

Sexual selection is an important and well-studied topic and is central to many theories on mate selection and individual behavior. Relatively little is known about the impacts that human-induced rapid environmental change are having on secondary sexually selected characteristics. In particular, we lack a clear understanding of the effects of the introduction of once-limiting nutrients to wild populations. Honest signals function as an indicator of mate quality when there are differences in nutrient acquisition and allocation to secondary traits. We used the cabbage white butterfly, *Pieris rapae* (L.), to investigate differences in color and testes size as an index of reproductive potential. We collected individuals from four sites in California and Nevada to capture variation among areas with high and low cropland and potential nitrogen availability. Differences in mean testes size among sites raise the possibility that individuals from sites surrounded by high agricultural areas have smaller testes than individuals from less agriculturally-developed areas. Coloration variables and testes size were positively associated, consistent with the hypothesis that nitrogen-based coloration in the cabbage white is an honest mating signal. However, variation among sites in that relationship suggests complexities that need further exploration, including the possibility that the signal is not of equal value in all populations. Thus these results advance our understanding of complex relationships among human-induced environmental change and sexual selection in the wild.

Key words: *Pieris rapae*, butterfly, honest signal, land use, nitrogen

Some of the most dramatic and inspiring aspects of the animal world are sexually selected traits,

including mammal horns, bird feathers, and the colors of insect wings, which have all been

appreciated by natural scientists and human societies (Darwin 1896, Johnstone 1995). Thus, it is

not surprising that a range of theories have been developed to make sense of this diversity, with

one of the most influential being Zahavi's handicap principle of sexual selection (Zahavi 1975).

This is the idea that because secondary sexual signals are costly and not all individuals are able

to produce these expensive traits, their presence serves as an honest signal of individual quality

to potential mates. The handicap principle of sexual selection has been successfully investigated

in a number of natural systems. For example, in many bird species it has been observed that

longer tails (Moller and de Lope 1994) and testosterone-dependent ornaments (Folstad and

Karter 1992, Mougeot et al. 2007) are indicators of individual quality.

In addition to assuming variation among individuals in quality and the existence of

expensive traits, honest-signaling theory relies on the ability of the signal receiver to distinguish

high-quality signals as well the existence of a benefit for the signal receiver for mating with such

individuals. In some cases, the benefit may be directly realized through nuptial gifts or other

such provisioning during or after mating (Arnqvist and Nilsson 2000, Stålhandske 2001, Fedorka

and Mousseau 2002, Lewis et al. 2014). In other cases, evolutionary complexity arises because

the benefit may be through genes that will be passed on to offspring with associated

improvements to their fitness, as in the sexy sons hypothesis (Weatherhead and Robertson 1979).

For instance, individuals that can expend a higher amount of energy acquiring nutrients for a

signal show conspecific individuals (potential mates) that their offspring will also be able to

express similar genes and will be able to expend similar amounts of energy to obtain costly

nutrients (Scriber and Slansky 1981). The ability to acquire nutrients has impacts on individual

fitness (Boggs and Freeman 2005) and population dynamics (Batzli 1983, Pedersen and Greives

2008) and may be especially important under stressful conditions including in areas exposed to

high levels of human-induced change (Boggs and Freeman 2005, Boggs 2009, Boggs and

Niitepõld 2014).

In the era of human-induced rapid environmental change, examples are being reported in

which honest signals and associated "handicaps" have changed or might be affected in coming

years (Sih et al. 2011, Tuomainen and Candolin 2011). Of particular interest in this context is the

fact that human activity, through industry, agriculture and private land management, is having

enormous effects on nutritional landscapes (Snell-Rood et al. 2015). In disturbed areas, such

modification comes about through food waste (Robb et al. 2008), road salting (Snell-Rood et al.

2014), vehicle-associated aerial chemical deposition, and fertilizer application, among other

processes. Fertilizer deposition and pollution are flooding natural and disturbed areas with once-

limiting nutrients, such as phosphorous and nitrogen (Vitousek et al. 1997, Smil 2000, Galloway

et al. 2008, Yuan et al. 2018), and our understanding of the implications of such nutritional

changes on sexually selected characteristics is growing (Snell-Rood et al. 2015), especially in

terms of honest signaling and sexual selection (Goos et al. 2016, Espeset et al. 2019).

In this study, we ask if a putative honest signal is associated with a measure of mating

potential, and if that relationship (between advertisement and mating potential) differs among

populations. *Pieris rapae* (L.), the cabbage white butterfly, was chosen as our model system

because of its mate-choice strategy and previous information on sexually selected characteristics

of this species. For example, we know that *P. rapae* females select mates based on wing

coloration (Morehouse and Rutowski 2010b) and that the characteristic white and yellow wing

color of butterflies in the family Pieridae is from the pterin (or pteridine) pigment compound

class (Stavenga et al. 2004, Wijnen et al. 2007). Pterins are made up of heterocyclic compounds

packed with nitrogen. Therefore, when investigating wing coloration in terms of honest

signaling, all individuals who have high amounts of pterin pigments in their wings must have had

access to sufficient amounts of nitrogen that could be allocated to this secondary trait (a trait that

is not directly important for the survival and reproduction of an organism (Zahavi 1975)). We

can expect that the availability of sufficient nitrogen would mean that organisms could invest in

advertising (secondary sexual traits) while still investing in basal reproductive traits, such as

testes size, thus producing a correlation among individuals between secondary sexually-selected

traits and reproductive value or potential. Furthermore (and of key interest in the context of

landscapes flooded with anthropogenically-derived nutrients), this picture of honest signaling

and sexual selection may shift if all individuals in a population have access to excessive amounts

of a previously limiting nutrient (discussed further below).

In this study we address the following: (1) Does cabbage white wing coloration predict

testes size? This is a key element of honest signaling theory applied to the cabbage white system,

and addressing this question is our first motivation for the present study. Secondarily, we ask: (2)

Do wing coloration, sexual organs, and the relationship between coloration and reproductive

potential vary among populations located near high agricultural areas and areas with lower

agricultural activity? (3) Finally, we investigate variation in spermatophore count as an index of

repeated matings among study sites which potentially provides information on variation in

mating behavior.

In general (and in answer to question 1), we hypothesize that butterflies with larger testes

will have increased brightness, hue, and saturation, signifying the honesty of wing coloration for

reproductive quality. Additionally (in answer to question 2), we expected to see butterflies with

larger testes and more colorful wings to be from populations with higher atmospheric nitrogen

deposition (e.g. developed and highly populated areas) or land use associated with a higher direct

fertilizer input (e.g. croplands). We anticipated seeing less variation among individuals from

high-nitrogen areas and greater variation among individuals from natural or undisturbed areas

due to the more heterogeneous nature of nutrient availability in non-agricultural areas (see Fig. 1

for an illustration of one way in which these among-population differences might be realized).

Furthermore, if individuals from high-nitrogen areas indeed have uniformly larger testes and

brighter wings, then it is possible that the honest signaling relationship between testes and

coloration will be absent in these areas (Fig. 1B), and even (finally) it is possible that sexual

selection will thereby have been relaxed on wing coloration and individuals in high-nitrogen

areas could actually be less bright relative to natural areas, if selection on wing color has been

relaxed (the latter possibility is not illustrated in Fig. 1, see note in legend). We raise these

possibilities while acknowledging the complexity of the evolutionary processes involved in

organisms responding to altered environments, of which we are capturing a snapshot, at best,

with the number of populations sampled.

Materials and Methods

Specimens

Butterflies were collected at two sites in northern California and two in northern Nevada between

June and July 2018 (Figure 1) and transported in glassine envelopes to a lab at the University of

Nevada, Reno. Sites were selected to encompass variation in land use intensity, especially

regional agricultural activity. Once in the lab, bodies and wings were separated; wings were

stored in a dark drawer to avoid light degradation and bodies were stored in a -20°C freezer until

further processing.

Wing Measurements

Wing coloration was measured using a JAZ spectrophotometer. Visible spectra were measured

and processed as follows. Spectra were first trimmed to wavelengths of 300nm-700nm (the

visible light spectra) and smoothed using the PAVO package in R (with a smoothing span of 0.2

to maintain biological variation in reflectance measurements but reducing excess noise). Three

colorimetric variables were calculated following previous work by other researchers in this area

(Morehouse and Rutowski 2010a, Tigreros 2013, Espeset et al. 2019). In summary, midpoint

(R50, associated with brightness) was the average percent reflectance between 300-375 and 450-

550nm; LR50, associated with hue, was calculated as the associated wavelength to R50; and

BR50, associated with saturation, was calculated by finding the slope of the line tangent to the

point at R50. These measurements have been shown to be associated with pterin amount through

comparisons of pterin-extracted wings and non-pterin extracted wings, and through comparisons

of granule counts in scanning electron microscope images (Morehouse and Rutowski 2010b).

Wings were photographed and measured from the basal tip to apex tip, and wing length was used

as a proxy for body size.

Dissections

Abdomens were added to approximately 1 mL 10% PBS buffer (Fischer Scientific). Testes and

accessory glands were removed from males and spermatophores were removed from females

under a dissecting scope in 1X PBS buffer. Reproductive organs were again stored in 10% PBS

buffer and at -20°C until processed. Testes were weighed to the nearest microgram on a Mettler

Toledo microbalance. Spermatophore counts were taken at time of dissection.

Analyses

All analyses were conducted using R v.3.5.1. Linear models with population as a categorical

factor were used on testes weight to investigate differences among populations, with post-hoc

comparisons calculated using Tukey's honest significant different (HSD) test. A similar set of

analyses were used on spermatophore counts to examine population differences. Levene's tests

for homogeneity of variance were run as pairwise comparisons to determine differences in

variance among sites in color variables for male wings (with respect to hypotheses described

above and in Fig. 1).

Additional linear models were used to determine if colorimetric variables could predict

testes weight. Because colorimetric variables were highly correlated amongst themselves, we

ran three separate linear models for testes weight with each predictor colorimetric variable (R50,

LR50, and BR50); each model included population as a categorical predictor and wing length as

a covariate. We investigated the possibility that the relationship between color variables and

testes size varied among populations by testing for an interaction between population and color

variables. For all models, the distribution of residuals was examined for normality, and variance

inflation factors were evaluated to ensure independence of predictors. Slopes and associated

standard errors (from testes size predicted by color variables) were used to visualize differences

among populations in the strength of the putative honest signal.

Results

We caught and processed a total of 367 individuals from four different sites, two from northern

Nevada (on the edge of the city of Reno, and Washoe Lake) and two from north-central

California (the Davis Community Gardens, and Sutter National Wildlife Refuge), with numbers

of individuals as follows: 59 males and 28 females from Reno, 68 males and 30 females from

Washoe, 62 males and 27 females from Davis, and 54 males and 43 females from Sutter.

Individual level differences

Brighter (R50), longer wavelengths (hue; LR50), and more saturated (BR50) wings were

associated with larger testes sizes. In Fig. 2, we show relationships between testes weight and

color variables separately for each site as well as across all sites and individuals (the solid black

lines in Fig. 2), with results from the larger scale (across all individuals) as follows: R50, $F_{1,227}$ =

8.06, R^2 = 0.034, p = 0.005; LR50, $F_{1,227}$ = 15.03, R^2 = 0.062, p < 0.001; BR50, $F_{1,227}$ = 4.58, R^2

= 0.020, p = 0.033). Full models with site and wing length as additional variables (see Table 1)

explained considerably more variation in testes size (compared to color variables alone, as in Fig.

2), but the most powerful variable in those models was wing length. We did not detect an

interaction between color variables and collection site (R50, $F_{3,176}$ = 0.21, p = 0.89; LR50, $F_{3,176}$

= 0.81, p = 0.49; BR50, $F_{3,176}$ = 0.27, p = 0.85), even though (as in Fig. 2) the association

between color and testes size did appear to vary in magnitude among sites.

Population level differences

Testes size varied across sites: the California sites (with greater proximity to agriculture) had

overall smaller testes (and were similar to each other), while individuals from the Nevada sites

had larger testes, with individuals from Washoe being the largest (Fig. 3A). In general, the

Nevada sites are surrounded by less agriculture than the California locations, and the Washoe

site was the least agriculturally-developed of all (specimens were collected near a wetland and

terminal desert lake).

All populations had at least one individual with three spermatophores and all specimens

had at least one spermatophore found in dissections (except for one single individual from our

Reno site), with a median of one spermatophore dissected per population. Sites did not differ in

the average number of spermatophores dissected from females (Fig. 3B) suggesting that

populations at the different sites are similar in behavior with respect to the number of times

females successfully mate.

As an alternative visualization of relationships between testes and color variables, slopes

for each site from our above analyses of testes weight by R50, LR50, and BR50 variables (Fig.

2) are shown in Fig. 3C along with standard errors around the slope estimates. We found

considerable variation in slopes among populations, with variables that differ from the zero line

(i.e. standard errors not overlapping zero) suggesting honest signaling because a measure of wing

color was positively related to testes size, our measure of reproductive quality. This was evident

in the Reno and Davis populations (for BR50) in particular, and also for LR50.

We found variation among populations for all three color variables investigated, and that

was particularly true for R50 (see independent contrasts in Fig. 3D), with the brightest

individuals being from our Washoe site, Davis having the least bright individuals, and our other

two sites having intermediate brightness (Fig. 3D). We did not detect differences among sites in

hue or saturation in the simple contrasts used for visualization in Fig 3 (panels E and F), but it

should be noted that an overall effect of site on those color variables was detected in full models

(with other covariates) reported in Table 1. Using Levene's test for homogeneity of variance, we

did not detect differences among locations in the three wing color variables.

Discussion

Because mate choice, and reproduction in general, is important for population growth and

persistence, any changes to sexually selected signals are potentially important for our

understanding of population dynamics. One example is an increased frequency in the production

of interspecific hybrids when individuals are no longer able to distinguish conspecifics as a result

of increases in human disturbances (Van der Sluijs et al. 2011, Grabenstein and Taylor 2018).

Evidence that humans are also affecting such mating signals and systems through nutritional

changes to natural and modified landscapes is growing (Snell-Rood et al. 2015, Goos et al. 2016,

Espeset et al. 2019). In this study we looked at differences in the relationship between wing

coloration and testes size among populations in the cabbage white butterfly, a species with

extensive exposure to anthropogenic habitats. We collected specimens from two sites associated

with low cropland and two sites associated with high cropland to both quantify the extent to

which wing color is an honest signal in this system and to determine if there were differences in

the relationship between color and testes size in these different habitat types. We predicted that

individuals from areas with higher cropland surrounding collection sites would have brighter,

whiter, and more saturated wings with less variation among individuals as compared to

individuals from areas surrounded by low cropland.

Trait honesty within the cabbage white butterfly has been previously suggested

(Morehouse and Rutowski 2010b), although in that study male quality was not quantified. In

another study (Tigreros 2013), a relationship between spermatophore content and coloration was

not detected, however those authors did not measure color and reproductive tissue from the same

individual. Our results provide evidence that testes size is indeed positively associated with

coloration when measurements are taken from the same individuals. This could indicate that

coloration in cabbage white butterflies is, in fact, an honest indicator of reproductive investment,

although that relationship (between color and reproductive investment) is subtle (Fig. 2). We are

here making the assumption that testes size is a proxy for reproductive value in the context of

spermatophore quality (Gage 1994), an important "nuptial" gift in these butterflies in low-

nitrogen available areas (Tigreros et al. 2013).

When considering site-level relationships between color and testes, we did not detect an

overall interaction between site and color variables (predicting testes size), but we did find that

the honest-signaling relationship was notably stronger in some locations (see slopes in Fig. 3C,

and site-specific lines in Fig. 2). Reno had the strongest association of testes size with each color

variable, and Davis had a positive trend, while Washoe and Sutter sites showed much weaker or

nonexistent relationships between wing coloration and testes size (Fig. 2, Fig. 3C). With respect

to surrounding land-use, our Washoe site had the lowest proportion of cropland (0.125%) within

a 10km buffer around our collection location, followed by Reno (0.416%), then Davis (71.1%),

and finally Sutter (85.9%) with the highest proportion of cropland (National Land Cover

Database; www.mrlc.gov). Given the data currently in hand, it is not clear why our two

intermediate sites (neither most intensive crop area nor the area with the least cropland) showed

the strongest evidence of honest signaling, and an inconsistent effect across agricultural and non-

agricultural areas. We discuss possible interpretations below and acknowledge that this is a

worthwhile area for future study.

The difference in honest signaling among sites could suggest that individuals from the

very high agricultural land-use area (Sutter) have been exposed to relatively high amounts of

nitrogen for a longer period of time than any other site, relaxing the value of investing in an

expensive, sexually-selected secondary trait that no longer acts as an honest signal of mate

quality. Our Davis site may still hold some trait honesty but less than our Reno site, which has

been exposed to agricultural land and fertilizer deposition, but possibly for a shorter amount of

time. The fact that our site with the lowest assumed nitrogen availability (Washoe) did not show

trait honesty is perplexing, and likely highlights the need for further investigation and direct

measurement of levels of available nitrogen in varying natural landscapes. The Washoe site was

low density (in terms of adult butterflies), thus another possibility might be investment in testes

rather than wing color in an area were competition for mates is low.

In addition to the possibility that variable nitrogen availability (and variable lengths of

time that nitrogen input has been elevated at different sites) has reduced the value of male wing

color as an honest signal, it is also possible that nitrogen availability has had a direct effect on

the utility of nitrogen-based "nuptial gifts" (spermatophores) to females because they are able to

sequester a larger amount of nitrogen themselves. These nitrogen-flooded areas could also have

higher nitrogen-containing nectar sources (i.e. higher concentrations of amino-acids (Baker and

Baker 1977, Gardener and Gillman 2001)), with which individuals can replenish nutrient

depletions as adults and can increase reproductive success or fecundity (Jervis and Boggs 2005,

Mevi-Schütz and Erhardt 2005). More studies are needed to fully understand the dynamics of

increased nitrogen, including availability and sequestration at different life stages associated with

populations in different habitats (Bertazzini and Forlani 2016).

It would also be beneficial to examine mate-selection in areas exposed to different

amounts and histories of nitrogen availability. The relationship between colorimetric variables in

the visual spectra and female observation is still not completely understood in butterflies

(Stavenga and Arikawa 2006); however, we do know that females should be able to differentiate

individuals that vary in color within the visible range, which is what we have focused on

(Morehouse and Rutowski 2010b). We found positive relationships between testes size and

color, suggesting that it would indeed be beneficial for females to differentiate between

individuals of high and low quality based on color. A behavioral analysis of female mate

selection across populations in areas with differing nitrogen availability would help us

understand these signals better.

In our study, we focused on habitats that potentially differ in nutrient availability, but it is

important to remember that there are numerous other factors that could be causing the

differences in testes size we see in this study. Due to the labor-intensive nature of quantifying

both mate quality (testes size) and wing color, our investigations here were limited to four

locations. Thus we can conclude that the relationship between mate quality and the mating signal

varies among locations, but we cannot yet explain that variation, and expect that it involves

many factor not measured here, e.g. pesticide use, climate, elevation, maternal effects, and

others. The importance of additional, unmeasured factors is also supported by the observation

that individuals from agricultural areas tended to have smaller testes and less colorful wings (Fig.

3). The possibility of multiple factors driving variation in reproductive traits combined with the

possibility that different populations are at different points along an evolutionary trajectory

(possibly leading to the de-valuing of an honest signal) highlights the complexity and value of

studying sexually selected signals in the age of the Anthropocene.

Acknowledgments

This material is based upon work supported by the National Science Foundation Graduate

Research Fellowship under Grant No. 1447692 to AE. Thank you to Dr. Emilie Snell-Rood and

her lab for space and microscope use for reproductive dissections, thanks for Dr. Anne Leonard

for use of her JAZ spectrophotometer, and thanks to Dr. Art Shapiro for advice on when and

where to find the most butterflies.

References Cited

Arnqvist, G., and T. Nilsson. **2000**. The evolution of polyandry: multiple mating and female fitness in insects. Anim. Behav. 60: 145–164.

Baker, H. G., and I. Baker. **1977**. Intraspecific constancy of floral nectar amino acid complements. Bot. Gaz. 138: 183–191.

Batzli, G. O. 1983. Responses of arctic rodent populations to nutritional factors. Oikos. 396–406.

Bertazzini, M., and G. Forlani. **2016**. Intraspecific variability of floral nectar volume and composition in rapeseed (Brassica napus L. var. oleifera). Front. Plant Sci. 7: 288.

Boggs, C. L. 2009. Understanding insect life histories and senescence through a resource allocation lens. Funct. Ecol. 23: 27–37.

Boggs, C. L., and K. D. Freeman. **2005**. Larval food limitation in butterflies: effects on adult resource allocation and fitness. Oecologia. 144: 353–361.

Boggs, C. L., and K. Niitepõld. **2014**. Insights from stable isotopic tracers on reproductive allocation under stress.

Darwin, C. 1896. The descent of man and selection in relation to sex. D. Appleton.

Espeset, A., M. E. Kobiela, K. L. Sikkink, T. Pan, C. Roy, and E. C. Snell-Rood. **2019**. Anthropogenic increases in nutrients alter sexual selection dynamics: A case study in butterflies. Behav. Ecol. 30: 598–608.

Fedorka, K. M., and T. A. Mousseau. **2002**. Material and genetic benefits of female multiple mating and polyandry. Anim. Behav. 64: 361–367.

Folstad, I., and A. J. Karter. **1992**. Parasites, bright males, and the immunocompetence handicap. Am. Nat. 139: 603–622.

Gage, M. J. G. 1994. Associations between body size, mating pattern, testis size and sperm lengths across butterflies. Proc. R. Soc. London. Ser. B Biol. Sci. 258: 247–254.

Galloway, J. N., A. R. Townsend, J. W. Erisman, M. Bekunda, Z. Cai, J. R. Freney, L. A. Martinelli, S. P. Seitzinger, and M. A. Sutton. **2008**. Transformation of the nitrogen cycle: recent trends, questions, and potential solutions. Science. 320: 889–892.

Gardener, M. C., and M. P. Gillman. **2001**. Analyzing variability in nectar amino acids: composition is less variable than concentration. J. Chem. Ecol. 27: 2545–2558.

Goos, J. M., R. D. Cothran, and P. D. Jeyasingh. **2016**. Sex-specific nutrient use and preferential allocation of resources to a sexually selected trait in Hyalella amphipods. J. Exp. Biol. 219: 649–657.

Grabenstein, K. C., and S. A. Taylor. **2018**. Breaking barriers: causes, consequences, and experimental utility of human-mediated hybridization. Trends Ecol. Evol. 33: 198–212.

Jervis, M. A., and C. L. Boggs. **2005**. Linking nectar amino acids to fitness in female butterflies. Trends Ecol. Evol. 20: 585–587.

Johnstone, R. A. 1995. Honest advertisement of multiple qualities using multiple signals. J.
Theor. Biol. 177: 87–94.
Lewis, S. M., K. Vahed, J. M. Koene, L. Engqvist, L. F. Bussiere, J. C. Perry, D. Gwynne,
and G. U. C. Lehmann. 2014. Emerging issues in the evolution of animal nuptial gifts.
Biol. Lett. 10: 20140336.
Mevi-Schütz, J., and A. Erhardt. 2005. Amino acids in nectar enhance butterfly fecundity: a
long-awaited link. Am. Nat. 165: 411–419.
Moller, A. P., and F. de Lope. 1994. Differential costs of a secondary sexual character: an
experimental test of the handicap principle. Evolution (N. Y). 48: 1676–1683.
Morehouse, N. I., and R. L. Rutowski. 2010a. Developmental responses to variable diet
composition in a butterfly: the role of nitrogen, carbohydrates and genotype. Oikos. 119:
636–645.
Morehouse, N. I., and R. L. Rutowski. 2010b. In the eyes of the beholders: female choice and
avian predation risk associated with an exaggerated male butterfly color. Am. Nat. 176:
768–784.
Mougeot, F., L. PÉREZ-RODRÍGUEZ, J. Martinez-Padilla, F. Leckie, and S. M. Redpath.
2007. Parasites, testosterone and honest carotenoid-based signalling of health. Funct. Ecol.
21: 886–898.
Pedersen, A. B., and T. J. Greives. 2008. The interaction of parasites and resources cause
crashes in a wild mouse population. J. Anim. Ecol. 77: 370–377.
Robb, G. N., R. A. McDonald, D. E. Chamberlain, and S. Bearhop. 2008. Food for thought:
supplementary feeding as a driver of ecological change in avian populations. Front. Ecol.
Environ. 6: 476–484.
Scriber, J. M., and F. Slansky. 1981. The nutritional ecology of immature insects. Annu. Rev.
Entomol. 26: 183–211.
Sih, A., M. C. O. Ferrari, and D. J. Harris. 2011. Evolution and behavioural responses to
human-induced rapid environmental change. Evol. Appl. 4: 367–387.
Van der Sluijs, I., S. M. Gray, M. C. P. Amorim, I. Barber, U. Candolin, A. P. Hendry, R.
Krahe, M. E. Maan, A. C. Utne-Palm, and H.-J. Wagner. 2011. Communication in
troubled waters: responses of fish communication systems to changing environments. Evol.
Ecol. 25: 623–640.
Smil, V. 2000. Phosphorus in the environment: natural flows and human interferences. Annu.
Rev. energy Environ. 25: 53–88.
Snell-Rood, E. C., A. Espeset, C. J. Boser, W. A. White, and R. Smykalski. 2014.
Anthropogenic changes in sodium affect neural and muscle development in butterflies.
Proc. Natl. Acad. Sci. 111: 10221–10226.
Snell-Rood, E., R. Cothran, A. Espeset, P. Jeyasingh, S. Hobbie, and N. I. Morehouse. 2015.
Life-history evolution in the anthropocene: Effects of increasing nutrients on traits and
trade-offs. Evol. Appl. 8: 635–649.
Stålhandske, P. 2001. Nuptial gift in the spider Pisaura mirabilis maintained by sexual selection.
Behav. Ecol. 12: 691–697.
Stavenga, D. G., and K. Arikawa. 2006. Evolution of color and vision of butterflies. Arthropod
Struct. Dev. 35: 307–318.
Stavenga, D. G., S. Stowe, K. Siebke, J. Zeil, and K. Arikawa. 2004. Butterfly wing colours:
scale beads make white pierid wings brighter. Proc. R. Soc. London. Ser. B Biol. Sci. 271:
1577–1584.

Tigreros, N. 2013. Linking nutrition and sexual selection across life stages in a model butterfly
system. Funct. Ecol. 27: 145–154.
Tigreros, N., E. M. Sass, and S. M. Lewis. **2013**. Sex-specific response to nutrient limitation
and its effects on female mating success in a gift-giving butterfly. Evol. Ecol. 27: 1145–
1158.
Tuomainen, U., and U. Candolin. **2011**. Behavioural responses to human-induced
environmental change. Biol. Rev. 86: 640–657.
Vitousek, P. M., J. D. Aber, R. W. Howarth, G. E. Likens, P. A. Matson, D. W. Schindler,
W. H. Schlesinger, and D. G. Tilman. **1997**. Human alteration of the global nitrogen
cycle: sources and consequences. Ecol. Appl. 7: 737–750.
Weatherhead, P. J., and R. J. Robertson. **1979**. Offspring quality and the polygyny threshold:"
the sexy son hypothesis". Am. Nat. 113: 201–208.
Wijnen, B., H. L. Leertouwer, and D. G. Stavenga. **2007**. Colors and pterin pigmentation of
pierid butterfly wings. J. Insect Physiol. 53: 1206–1217.
Yuan, Z., S. Jiang, H. Sheng, X. Liu, H. Hua, X. Liu, and Y. Zhang. **2018**. Human
perturbation of the global phosphorus cycle: changes and consequences. Environ. Sci.
Technol. 52: 2438–2450.
Zahavi, A. 1975. Mate selection—a selection for a handicap. J. Theor. Biol. 53: 205–214.
425

426
427
428

Figure legends

430
431

Fig. 1. Expectations for the evolution of an honest signal (A and B) and a map (C) of collection
locales. Values graphed in panels (A) and (B) are artificial, made to illustrate the following ideas.
In (A), wing color by land use, we expect an increase in color in association with higher nitrogen
availability at agricultural sites (green points). We also hypothesize a smaller variance within
these populations because all individuals have access to excess nitrogen. In (B), slopes of the
relationship between color by testes weight are a measure of signal honesty (more positive slopes
indicate a relation between traits in which males with more reproductive potential are also more
intensely colored). In areas where there is excess nitrogen (agricultural areas) we would expect
the correlation of color by testes weight to be zero or closer to zero (because the signal has lost
value). In more nitrogen-limited areas, in contrast, we would expect to see a positive relationship
between color and testes weight corresponding to signal honesty. Points in (C) are actual
collection locations of *P. rapae* obtained across different proportions of cropland in Northern
California and Northern Nevada. Blue represents sites and areas with low proportion of cropland
(Washoe and Reno) and green signifies high levels of cropland (Davis and Sutter). Note that the
expectations illustrated in the first two panels are a potential snapshot of a complex process. A
further possibility (not illustrated here) is that loss of the honest signal in populations where all
individuals have access to excess nitrogen could ultimately relax selection on wing coloration
and lead to reduced brightness in those areas (i.e., a further step in the process than illustrated in
panel (A).

Fig. 2. Teste weights by colorimetric variables (i.e. R50: midpoint; LR50: wavelength; and
BR50: slope) for all male individuals from Davis, Sutter, Washoe, and Reno sites (see Fig. 1
map). Relationships across all individuals in each plot are represented by the solid black lines.
Relationships for subsets of individuals (by site) are shown as colored lines, with relationships
associated with $P < 0.05$ shown as solid lines.

Fig. 3. Within and among population variation in key variables studied, including reproductive
traits (A and B) and colorimetrics (D-F). In panels A-E and F, points are jittered for visualization
and lower case letters within plots are shown if post hoc comparisons indicated differences
among populations at $P < 0.05$. Panel C shows among-population differences in the slope of
testes weight by color variables (see legend for symbols) where more positive values suggest an
honest signal (a relationship between testes weight and wing color).

464
465
466

Table 1. Results from linear models predicting testes size with color variables, wing length, and site as a categorical factor (see Fig. 1 for locations studied).

	Testes by R50 Model		Testes by LR50 Model		Testes by B50 Model	
	F-value$_{(Df)}$	P value	F-value$_{(Df)}$	P value	F-value$_{(Df)}$	P value
Color Variable	$7.64_{(1)}$	P=0.006	$12.6_{(1)}$	P<0.001	$2.86_{(1)}$	P=0.09
Wing Length	$36.6_{(1)}$	P<0.001	$30.1_{(1)}$	P<0.001	$35.8_{(1)}$	P<0.001
Site	$8.81_{(3)}$	P<0.001	$10.7_{(3)}$	P<0.001	$9.87_{(3)}$	P<0.001
	$F_{(5,179)}=14.13$; $R^2=0.28$; p<0.001		$F_{(5,179)}=14.97$; $R^2=0.29$; p<0.001		$F_{(5,179)}=13.65$; $R^2=0.28$; p<0.001	

Fig. 1.

Fig. 2.

Fig. 3.