Integrative multi-omics profiling reveals cAMP-independent mechanisms regulating

hyphal morphogenesis in *Candida albicans*

Kyunghun Min[1], Thomas F. Jannace[1,5], Haoyu Si[1,6], Krishna R. Veeramah[2], John D. Haley[3,4],

and James B. Konopka[1]*

[1] Department of Microbiology and Immunology, Renaissance School of Medicine, Stony Brook

University (SUNY), Stony Brook, NY, USA

[2] Department of Ecology and Evolution, Stony Brook University (SUNY), Stony Brook, NY, USA

[3] Department of Pathology, Renaissance School of Medicine, Stony Brook University (SUNY),

Stony Brook, NY, USA

[4] Biological Mass Spectrometry Shared Resource, Renaissance School of Medicine, Stony

Brook University (SUNY), Stony Brook, NY, USA

[5] Present address: Department of Quality Assurance, Biogen, Morrisville, NC, USA

[6] Present address: School of Biological Sciences, University of Nebraska-Lincoln, Lincoln, NE,

USA

* Corresponding author: james.konopka@stonybrook.edu

Short Title: Regulation of *C. albicans* hyphal morphogenesis

ABSTRACT

Microbial pathogens grow in a wide range of different morphologies that provide distinct advantages for virulence. In the fungal pathogen *Candida albicans*, adenylyl cyclase (Cyr1) is thought to be a master regulator of the switch to invasive hyphal morphogenesis and biofilm formation. However, faster growing *cyr1*Δ/Δ pseudorevertant (PR) mutants were identified that form hyphae in the absence of cAMP. Isolation of additional PR mutants revealed that their improved growth was due to loss of one copy of *BCY1*, the negative regulatory subunit of protein kinase (A) from the left arm of chromosome 2. Furthermore, hyphal morphogenesis was improved in some of PR mutants by multigenic haploinsufficiency resulting from loss of large regions of the left arm of chromosome 2, including global transcriptional regulators. Interestingly, hyphal-associated genes were also induced in a manner that was independent of cAMP. This indicates that basal protein kinase A activity is an important prerequisite to induce hyphae, but activation of the cAMP pathway is not needed. Instead, phosphoproteomic analysis indicated that the Cdc28 cyclin-dependent kinase and the casein kinase 1 family member Yck2 play key roles in promoting polarized growth. In addition, integrating transcriptomic and proteomic data reveals hyphal stimuli induce increased production of key transcription factors that contribute to polarized morphogenesis.

AUTHOR SUMMARY

The human fungal pathogen *Candida albicans* switches between budding and

filamentous hyphal morphologies to gain advantages for virulence and survival in the host.

Although adenylyl cyclase has been thought to be a master regulator that controls this switch,

we identified *C. albicans* pseudorevertant mutants that grow better and form hyphae in the

absence of adenylyl cyclase and cAMP. The mutant cells were also able to induce hyphal-

associated genes in the absence of cAMP that are needed for virulence. Integrating information

from different omics approaches identified cAMP-independent mechanisms that promote hyphal

growth. This includes phosphoproteomic studies that revealed key roles for the Cdc28 cyclin-

dependent kinase and casein kinase 1 in promoting hyphal growth. In addition, integrating

transcriptomic and proteomic data revealed that post-transcriptional mechanisms regulate the

levels of a set of key transcription factors that are important for hyphal induction, suggesting a

novel form of translational regulation. These studies better define the pathways that stimulate *C.*

*albican*s to switch from budding to hyphal growth, which is important for invasion into tissues,

escape from the immune system, and biofilm formation.

INTRODUCTION

Fungal pathogens are capable of transitioning between different morphologies that provide distinct advantages for virulence and survival in the host [1, 2]. For example, the fungal pathogen *Candida albicans* switches from budding to hyphal growth, which promotes virulence as the long hyphal filaments mediate invasion into tissues, escape from immune cells, and biofilm formation [1-6]. A variety of stimuli promote hyphal morphogenesis, including serum, CO_2, alkaline pH, peptidoglycan breakdown products, N-acetylglucosamine (GlcNAc), and contact with a solid matrix [3, 7, 8]. In *C. albicans*, adenylyl cyclase (Cyr1) is thought to be a master regulator of hyphal growth [3, 5, 9-11]. Deletion of adenylyl cyclase (*cyr1Δ/Δ*) blocks hyphal formation, and addition of millimolar levels of exogenous cAMP induces it [12]. However, interpreting the role of Cyr1 and cAMP is complicated by the fact that the *cyr1Δ/Δ* mutants grow very poorly [13]. Furthermore, a recent study showed that faster growing *cyr1Δ/Δ* pseudorevertant (PR) mutants form hyphae in the absence of Cyr1 and cAMP [14]. Thus, although cAMP is capable of inducing hyphae, other pathways can contribute to stimulating the formation of filamentous hyphal cells [14].

In *C. albicans*, cAMP formed from ATP by Cyr1 acts by binding to the negative regulatory subunit (Bcy1) of protein kinase A (PKA), thereby releasing the PKA catalytic subunits (Tpk1 and Tpk2) to phosphorylate target proteins [11]. Most current models propose that the cAMP-PKA pathway induces the expression of a special set of genes that promote hyphal growth [2, 3, 11, 13, 15, 16]. This has been supported by the ability of many transcription factors (TFs) to regulate this switch. For example, PKA was reported to phosphorylate the Efg1 TF, which is required for hyphal induction [17, 18]. However, it has been surprising that none of the hyphal-induced genes has been shown to play a direct role in the morphogenesis machinery that determines cell shape [19, 20]. The common core set of genes stimulated by a group of different hyphal inducers in *C. albicans* do not account for the transition to hyphal growth [19]. In addition, our recent studies showed that a mutant that

cannot catabolize GlcNAc can be stimulated by GlcNAc to form hyphae without induction of

hyphal-specific genes [20]. Other studies have shown that some hyphal regulatory TFs can be

bypassed under special environmental conditions [21, 22], or by altered expression of the Cak1

protein kinase [23], the absence of a variant histone H3 [24], or mutation of the Ssn3 subunit of

the Mediator transcription complex [25]. This indicates that although TFs are important for

creating the proper physiological environment for cells to undergo hyphal growth, the target

genes are not directly involved in promoting filamentous morphogenesis.

Other studies have implicated several protein kinases in hyphal signaling that are not

directly regulated by cAMP [2, 3, 26]. For example, the cyclin-dependent kinase Cdc28

promotes hyphal growth in part by phosphorylating several proteins involved in polarized

morphogenesis [26-34]. Therefore, to better define the cAMP-dependent and independent

mechanisms of hyphal signaling, we isolated a set of new *cyr1Δ/Δ* PR mutants that displayed

variable levels of improved growth rates and ability to form hyphae. The wild type and PR

mutant cells were compared with four omics approaches: genomics combined with gene-

mapping by CRISPR-Cas9 to identify genetic changes in PR mutants, transcriptomics to better

define the role of cAMP in hyphal gene regulation, proteomics and phosphoproteomics to reveal

key kinases for hyphal induction. Furthermore, integrative transcriptomics and proteomics

profiling indicates that there are increases in the levels of key transcription factors, possibly due

to changes in their translation rate, that are induced by hyphal signaling. Altogether, these

results reveal how multiple processes in the gene expression pathway can modulate cellular

differentiation.

RESULTS

Large deletions in the left arm of chromosome 2 bypass the need for cAMP to improve

the growth rate and to stimulate hyphal morphogenesis.

We previously reported that slow growing *cyr1*Δ/Δ mutant strains give rise to faster

growing cells, which we termed pseudorevertants (PRs) [14]. Interestingly, these spontaneous

PRs could form hyphae even though they lack Cyr1 and cAMP [14]. To better define the

underlying mechanisms, we isolated additional PR strains. These PR mutants were isolated

from independent cultures which were started from different single colonies of the *cyr1*Δ/Δ strain.

We found that they differed in their growth rates and ability to be induced to form hyphae (Fig

1A, 1B and S1 Fig). Based on these phenotypic differences, the new PR mutants could be

grouped into 4 classes (Table 1). Surprisingly, the degree of improved growth rate of the PR

mutants did not correlate with their ability to form filamentous cells, indicating that the

mechanisms of suppression were complex. The faster growing Class 1 mutant doubled at a

nearly wild-type rate (1.5 h) but only 74% of the cells formed filamentous cells. In contrast, the

Class 2, Class 3, and Class 3+ PR mutants grew slower than the Class 1 mutants but were

more efficient at forming filamentous cells. For example, Class 2 mutants grew at a doubling

time of 1.8 h yet 93% of the cells were filamentous. Class 3 and 3+ produced filamentous cells

but they were shorter and thicker than the WT hyphae.

Whole genome sequencing of the diploid *C. albicans* strains identified distinct genetic

changes in the four classes of PR mutants. Interestingly, although they were different, all four

classes of mutants contained genetic changes that resulted in loss of expression of one copy of

BCY1, which encodes the negative regulatory subunit of cAMP-dependent PKA kinase. This

suggested that reduced expression of *BCY1* increased PKA activity resulting in improved

growth in the absence of cAMP. Similar results reported that a *bcy1*Δ mutant can rescue growth

of a *cyr1*Δ mutant in *S. cerevisiae* [35]. Class 1 mutants exhibited the simplest genetic change,

in that the improved growth correlated with a premature stop codon in *BCY1* on chromosome 2

(Fig 1C, S1 Table, and S1 Data). In contrast, the Class 2, 3, and 3+ PR mutants all lacked part

of the left arm of one copy of chromosome 2 that includes *BCY1*.

Western blot analysis showed that the Bcy1 protein levels were reduced by 50% in the

PR mutants, confirming that altered gene dosage leads to reduced production of Bcy1 (S2 Fig).

The reduction in Bcy1 did not impact the levels of Tpk2, one of the PKA catalytic subunits in the

PR mutants compared to the WT. Interestingly, *TPK2* was located in the middle of the

chromosome 2, which was duplicated in Class 3+ PRs. However, none of the other PKA

pathway components were located on chromosome 2. This supports the conclusion that the

haploinsufficiency of *BCY1* bypassed the need for cAMP for growth and hyphal induction in *C.*

albicans.

In the Class 2 and 3+ PRs, similar breakpoints were identified in the independently

isolated strains indicating that a specific DNA sequence induced the chromosomal deletion. For

example, PR12 and PR14 (class 2) lost the same 276-kb region of chromosome 2 although they

were isolated from independent cultures on different days. We hypothesized that a specific

DNA sequence induced the breakpoints at the same location. To better understand how the

large deletions of 270 kb and 590 kb were generated in the Class 2, 3, and 3+ PRs, we

inspected the genome sequences for matches to the *C. albicans* 23-bp telomere repeat

sequence (CACCAAGAAGTTAGACATCCGTA) [36] , which was found at the deletion end

points (S1 Table). Analysis of the breakpoint regions in chromosome 2 revealed 4 to 9-bp

matches to telomere seed sequences. However, the long repeat sequence, which was

previously reported to drive genome rearrangements, was not found near the deletion sites [37].

As described previously, this suggests that chromosome break healing by *de novo* telomere

addition contributes to the genome plasticity in *C. albicans* [38-40], which enables it to adapt to

adverse conditions or deleterious mutations such as *cyr1Δ/Δ*.

Gene mapping by CRISPR-Cas9 identified general transcription factors are involved in

hyphal regulation.

157   To confirm that the genetic changes in chromosome 2 improved growth and hyphal

158 induction in the PR mutants, transient expression of the CRISPR-Cas9 system [41, 42] was

159 utilized to recreate the loss of one copy of *BCY1* and the large deletions present in the Class 2,

160 3, and 3+ PR mutants. To facilitate large deletions, two single guide RNAs (sgRNAs) were

161 used to target Cas9 to cut immediately adjacent to the deletion end points detected in the

162 spontaneous PR mutants (Fig 2A). Analysis of these *cyr1Δ/Δ* mutants that were created to be

163 heterozygous *bcy1Δ*, 270kbΔ, and 590kbΔ in chromosome 2 showed that they recapitulated the

164 phenotypes of the corresponding spontaneous mutants PR13, PR12, and PR18, respectively

165 (Fig 2B-D and S1 Fig). This confirmed that haploinsufficiency of *BCY1* improved growth, and

166 that haploinsufficiency of the genes in the 270 kb region in the left arm of chromosome 2 was

167 sufficient to improve hyphal induction in the PR mutants.

168   Gene mapping strategies to identify the basis for the improved hyphal growth of the PR

169 mutants showed that this effect is due to multiple genetic changes (Fig 2E and S1C Fig). In the

170 first approach, smaller 90 kb sections of the 270 kb region were deleted from one copy of

171 chromosome 2 in a *bcy1Δ cyr1Δ/Δ* strain. However, these mutants were not better at producing

172 filamentous hyphal cells than the parental strain, suggesting that the improved hyphal

173 morphogenesis is multigenic. Heterozygous deletion of a larger segment of chromosome 2

174 from 90kb→260kb made the cells more filamentous, which partially narrowed the key region.

175 Smaller deletions designed to map the right end of the critical region of chromosome 2 showed

176 that a 90kb→250kb deletion mutant did not exhibit better hyphal growth. Haploinsufficiency of

177 the 10-kb region was therefore necessary to improve hyphal induction (although it was not

178 sufficient on its own).

179   Further mapping by deletion of the genes in this 10-kb region identified *SRB9* and *SPT5*

180 as contributing to in the improved hyphal phenotype (S3 Fig). *SRB9* encodes a subunit of the

181 RNA polymerase II Mediator complex and *SPT5* encodes a transcription elongation factor

182 complex subunit. Interestingly, the double heterozygous mutation of *SRB9* and *SPT5* in the

90kb→250kb deletion strain promoted improved hyphal growth, while the single heterozygous

mutations of each gene did not. According to previous analyses of haploinsufficiency genetic

interactions [43, 44], this suggest that the two genes function in the same pathway, consistent

with both *SRB9* and *SPT5* encoding subunits of the general transcription machinery.

Interestingly, mutation of a different subunit of the Mediator complex (*SSN3*) was identified in a

previous study because it restored the ability to form hyphae to a *cph1*Δ/Δ *efg1*Δ/Δ mutant strain

of *C. albicans* that lacks two key hyphal TFs [25]. Thus, the gene mapping results indicate that

global transcriptional regulators are important for creating a physiological state in the cell that is

more conducive to induction of the switch of hyphal morphogenesis [25, 45].

Adenylyl cyclase is not necessary for transcriptional induction of hyphal regulated genes

Based in part on previous studies of *cyr1*Δ/Δ mutant strains, adenylyl cyclase was

thought to be central to inducing the expression of a variety of genes, including virulence factors

such as adhesins and hyphal-specific genes that promote filamentous growth [2, 3, 13, 15, 16].

To examine this role of cAMP signaling, we performed RNA sequencing (RNA-seq) analysis of

the WT and PR mutants before and after stimulation with the hyphal inducer GlcNAc (Fig 3A

and S2 Data). Heatmap clustering and principal component analysis (PCA) of the normalized

RNA-seq dataset revealed that WT and PRs showed similar patterns in transcriptomes,

indicating that adenylyl cyclase is not required to induce the broad range of hyphal genes (Fig

3B and S4 Fig). In contrast, the *cyr1*Δ/Δ mutant clustered separately due to its expected defect

in regulating hyphal genes (Fig 3B). Data for a subset of hyphal genes, as well as control genes

are shown in Fig 3C. These data show that the transcriptomic profile of class 1 mutant PR13

(*bcy1**/*BCY1 cyr1*Δ/Δ) was similar to the WT in having a low basal level of hyphal gene

expression that was induced by hyphal stimulation. However, class 2 mutant PR12 (270kbΔ

*cyr1*Δ/Δ) was distinct in that it expressed a higher basal level of hyphal genes even in the

absence of the hyphal inducer GlcNAc (Fig 3C), which correlated with a higher basal level of

filamentous cell morphology (Fig 1B). The PR12 transcriptome was also distinct as the levels of

the genes in the 270-kb region were around 2-fold less than the WT level due to the

haploinsufficiency (see S2 Data). GlcNAc catabolic genes were highly expressed in all the

strains tested, including *cyr1*Δ/Δ (Fig 3C), which is consistent with previous studies showing that

their regulation is cAMP independent [46, 47]. Altogether, these data indicate that an

appropriate basal level of PKA activity is needed for cells to respond transcriptionally to a hyphal

inducer, rather than a requirement for stimulation of Cyr1 to produce cAMP.

Quantitative phosphoproteomics identified protein kinases required for normal hyphal

growth

To better understand how the switch to hyphal morphogenesis is regulated, we

compared protein production and phosphorylation during hyphal induction. WT and PR13

strains were treated with or without GlcNAc to form hyphae (Fig 4A), and then a label-free mass

spectrometry approach (LC-MS/MS) was used to detect 3,434 proteins and 17,580

phosphopeptides (S3 Data). To differentiate experimental variances from biological alterations,

we compared the replicate data of WT cells treated with GlcNAc from both the whole proteome

and the phosphoproteome. The null comparisons of the biological replicates showed high

correlations ($r > 0.9$) (Fig 4B). We first examined the phosphorylation status of Efg1, since

genetic studies implicated this TF as being a target of PKA and Cdc28 kinases on positions

T179 and T206 positions, respectively [17, 34]. However, our phosphoproteome data did not

detect phosphorylation at these sites in Efg1, consistent with the reports of Willger et al. [45]

and Cao et al. [48] (see S3 Data). Thus, three completely independent MS studies have not

detected phosphorylation of Efg1 which was implicated from genetic evidence. Interestingly,

during hyphal induction Efg1 was dephosphorylated in the C-terminal prion-like domain (PrLD),

which is a region of the Efg1 protein that is involved in the assembly of TF complexes by

formation of phase-separated condensates [49]. Altogether, these results indicate that models

for role of Efg1 regulation by phosphorylation during hyphal induction should be revised (see

Discussion).

We next used Gene Set Enrichment Analysis [50] to infer the activity of specific protein

kinases. The algorithm was based on the levels of phosphorylated peptides containing amino

acid sequence motifs characteristic of the substrate specificity of known protein kinases (Fig 4C).

Strikingly, in contrast to the current models [2, 3, 5], this analysis revealed PKA activity was not

significantly different before and after hyphal induction in the WT or PR13 strains (Fig 4D).

Furthermore, control studies showed that the levels of the PKA subunits Bcy1 and Tpk2

remained constant during hyphal induction (S2 Fig). Thus, there was no evidence to support

the stimulation of PKA in the PR mutants during hyphal induction. In contrast, Cdc28 kinase

activity was increased during hyphal induction in the WT strain (Fig 4D), consistent with

previous studies implicating this cyclin dependent kinase in hyphal regulation [27, 28, 34, 51].

To examine the role of Cdc28 further, we deleted the hyphal-induced cyclin *HGC1*. The

hgc1Δ/Δ mutation caused a stronger hyphal defect in the PR13 mutant compared to the WT

control cells (S5AB Fig), indicating that PR13 cells are more reliant on cyclin-dependent activity

for hyphal induction than are the WT cells. Despite of the abnormal morphology, the *hgc1Δ/Δ*

mutant produced a similar number of filamentous cells as the WT control strain (~95%).

However, the PR13 *hgc1Δ/Δ* mutant produced a lower level of filamentous cells (36%)

compared to the PR13 strain (61%).

Interestingly, many of the potential Cdc28 phosphosites also contained a second motif

indicating they are targets for Yck2, a member of the casein kinase I family (Fig 4E-G). Recent

studies in *C. albicans* have characterized the role of Yck2 in modulating morphogenesis,

virulence, fungal cell-wall stress responses, and resistance to clinical antifungal drugs [52-54].

Deletion of *YCK2* strongly affected hyphal growth in both the WT and PR13 strains (S5AB Fig).

The *yck2Δ/Δ* cells displayed an elongated cell morphology, but did not switch to a true hyphal

morphology in the presence of the hyphal inducers serum or GlcNAc. Interestingly, both

kinases are predicted to phosphorylate 11 sites in 8 proteins that are all thought to have a role

in polarized growth (S2 Table). We created phospho-site mutations of *BNI1* and *MOB2*

because these genes are required for normal hyphal growth [32, 55]. However, mutation of the

individual phosphorylation sites did not cause a detectable hyphal defect, suggesting that

mutations of additional target proteins may be required to cause a strong defect in hyphal

morphogenesis (S5C Fig).

Hyphal stimulation induces translation of TFs that regulate morphological switching

The transcriptomic and proteomic data sets were examined to determine how the

abundance of mRNAs and their corresponding proteins correlated during the switch to hyphal

morphogenesis (Fig 5A). There was a relatively weak correlation (Pearson correlation

coefficient, r = 0.39) between mRNA and protein abundances in *C. albicans* (Fig 5B). Typical

Pearson correlation coefficients (r) for mammalian cells are about 0.6 and for *S. cerevisiae* it is

about 0.7 [56]. This indicated that *C. albicans* has dynamic post-transcriptional regulatory

mechanisms. Since protein levels are a function of mRNA abundance, translation rate, and

protein turnover, we approximated the translation rate with the protein/mRNA level ratio (this

approximation neglects protein degradation, see Fig 5C). Integration of RNA-seq and label-free

LC-MS/MS data gave an opportunity to estimate the translation rate of 3,267 genes (S4 Data).

The translation rate spanned over five orders of magnitude across genes (Fig 5D). There was a

negative correlation between mRNA abundance and translation rate, while there was a positive

correlation between protein abundance and translation rate. Although it has been reported that

5' UTR length plays an important role in translation efficiency [57-59], there was no correlation

between 5' UTR length and translation rate in our analysis.

We next examined whether the translation rate changed for TFs that are known to

regulate hyphal morphogenesis. As a group, the predicted translation rate of the 11 hyphal

regulator TFs did not change significantly after hyphal induction (Fig 5E). However, there was a

huge variation in the translation rates of the hyphal regulator TFs before hyphal stimulation and

some of them were very low. Strikingly, translation rates of these TFs increased dramatically

during hyphal growth while mRNA levels did not change much compared to the translation rate

(Fig 5F and S6 Fig). The mean translation rates of ribosomal proteins and total proteins were

similar, indicating selective translation regulation of the TFs. The Ume6, Sfl1, Brg1, Tec1, and

Fkh2 proteins were not detected in the uninduced cells while their mRNA levels were similar to

the median of total mRNAs (S4 Data). Further work will be required to confirm whether these

effects are due to translation or another mechanism, such as increased protein stability [60, 61].

However, the results demonstrate that a post-transcriptional mechanism regulates key

transcription factors that are important for hyphal morphogenesis.

DISCUSSION

The ability of *C. albicans* to switch between budding and hyphal morphologies

contributes to virulence by promoting invasive growth, escape from immune cells, and biofilm

formation [1-6]. The prevailing model has been that the Cyr1 adenylyl cyclase acts as a master

regulator by integrating signals from different hyphal inducers [3, 5, 9-11]. Although Cyr1 clearly

plays an important role, the failure of *cyr1Δ/Δ* cells to form hyphae has often been interpreted to

indicate that cAMP plays an essential role. However, the phenotype of *cyr1Δ/Δ* cells is

complicated by the fact that they grow slowly and have an abnormal physiology [12, 13]. The

identification of faster growing *cyr1Δ/Δ* PR mutants provided strong proof that inducers including

serum, GlcNAc, and alkaline pH can stimulate hyphae in the total absence of Cyr1 and cAMP

[14]. In this study we demonstrated that the improved growth of the PR mutants is due to

mutation or deletion of one of the two copies of *BCY1*, which encodes the negative regulatory

subunit of PKA (Fig 1 and Fig 2). The lower level of Bcy1 in the PR mutants was confirmed by

Western blot analysis (S2 Fig) and RNA-seq data (Fig 3). These results are also interesting in

that genomic analysis revealed that the different classes of PR mutants showed distinct

mechanisms for mutation of *BCY1* or deletion of parts of the left arm of chromosome 2 (S1

Table), which underscore how the plasticity of the *C. albicans* genome enables it to adapt to

different kinds of stress from mutations or antifungal drugs [37, 38, 62, 63]. It was also

significant that the degree of improved growth of the different PR mutant classes did not

correlate with their ability to form hyphae. This indicated that additional genes affected by the

deletions in chromosome 2 also contributed to the ability of PR mutants to form hyphae.

Current models based on genetic evidence predict that PKA phosphorylates the Efg1 TF

to stimulate it to induce expression of hyphal genes [2, 3, 11, 13, 15, 16]. However, we failed to

detect phosphorylation of Efg1 in WT or PR13 cells, consistent with two previous

phosphoproteomic studies [45, 48]. In spite of this, the PR mutants were not only able to form

hyphae, they also induced the expression of the expected set of hyphal-associated genes that

includes many virulence factors (Fig 3C). A further limitation of the current models is that

several studies now question the role of hyphal-induced genes in promoting hyphal growth. For

example, the common core set of genes stimulated by a group of different hyphal inducers in *C.*

albicans do not account for the transition to hyphal growth [19]. In addition, under certain

conditions GlcNAc can stimulate hyphae without induction of hyphal-specific genes [20]. There

also appears to be a lack of TF specificity, as some hyphal regulatory TFs can be bypassed

under special environmental conditions [21], or by altered expression of the Cak1 protein kinase

[23] or the absence of a variant histone H3 [24]. In addition, hyphal growth was restored in a

cph1Δ/Δ efg1Δ/Δ mutant by a mutation in the *SSN3* subunit of the Mediator complex [25].

Consistent with this, we found that haploinsufficiency of two global transcriptional regulators,

SRB9 (a component of Mediator) and *SPT5*, contributed to the improved filamentous growth of

the Class 2 mutants (S3 Fig). Altogether, these data suggest a new model that TFs are

important for creating the proper physiological environment for cells to undergo hyphal growth,

rather than inducing specific target genes that are directly involved in promoting filamentous

morphogenesis (Fig. 6).

A perplexing aspect of hyphal induction is that it can be induced by such a broad range

of different environmental and nutritional conditions (e.g. 37˚C, alkaline pH, nitrogen levels,

GlcNAc, certain amino acids) [2, 3]. We suggest that there are many different pathways that

can lead to a physiological state that induces hyphal growth, which helps to explain why there

does not appear to be a common transcriptional response to different hyphal inducers.

However, TFs can play a role in inducing the appropriate state as mutation of *TUP1* or

overexpression of *UME6* is sufficient to induce hyphal growth [16, 64]. These TF mutants

display changes in a wide range of genes that can impact cell physiology. In addition, studies

presented in Fig. 5 indicate a role for positive feedback in that induction of hyphal growth can

lead to increased production of hyphal TFs (Fig. 6). This model also helps to explain other data

that metabolic pathway mutants can influence hyphal morphogenesis, such as mitochondrial

mutants [2, 65-67].

A further prediction of the model shown in Fig. 6 is that the new physiological state is

expected to lead to activation of protein kinases such as Cdc28 and Yck2. These kinases have

been shown previously to be important for inducing hyphal growth in WT cells, and our

phosphoproteomic data indicate that they are important in the absence of cAMP. In fact, it

appears that PR mutants are more reliant on Hgc1-Cdc28 than are WT cells (S5 Fig). An

interesting aspect of the phosphoproteomic data is that we identified a subset of proteins which

contain peptides that are phosphorylated on sites for both Cdc28 and Yck1. Although mutating

single sites in two of these proteins that are implicated in hyphal morphogenesis did not cause a

strong phenotype, it is possible that there are synergistic effects of Cdc28 and Yck2 via

regulation of a larger set of substrate proteins. Altogether, these results indicate that future

studies on hyphal signaling should focus on the roles that post-transcriptional and post-

translational regulation play in promoting the changes in the morphogenesis pathways that

mediate highly polarized hyphal growth.

 MATERIALS AND METHODS

 Strains, culture conditions, and hyphal analysis

The *C. albicans* strains used in this study are described in S3 Table. *C. albicans* strains

were streaked fresh from storage at -80°C onto YPD (1% yeast extract, 2% peptone, 2%

glucose) agar plates and then grown in rich YPD medium or in complete synthetic medium (SC)

made with yeast nitrogen base, select amino acids, and uridine. *C. albicans* transformants were

selected on YPD plus 200 µg/ml nourseothricin (NAT; Gold Biotechnology). NAT-sensitive

derivatives of cells carrying the SAT flipper were then obtained by growing cells on maltose agar

medium (1% yeast extract, 2% peptone, 2% maltose) for 2 days, and then several colonies

were spread on YPD plates containing 25 µg/ml of NAT and incubated for 2 days at 30°C [42].

NAT-sensitive cells formed smaller colonies than NAT-resistant parental strains.

Strain doubling times were determined from growth curves in YPD at 30°C. Fresh

overnight cultures were diluted to a final concentration of ~0.05 OD_{660} in 5 ml YPD in test tubes

and then incubated at 30°C on a tube roller. The OD_{660} was measured hourly for ~8 hours until

OD_{660} was >1.0. The reported doubling times are the averages of two biological replicates each

performed in duplicate.

To analyze hyphal formation in liquid media, cells were grown overnight at 30°C in YPD,

and then inoculated into SC + 50 mM glucose + 10% bovine calf serum, or SC + 50 mM GlcNAc

medium. Samples were then incubated at 37°C for the indicated time and then images were

captured using a Zeiss Axio Observer 7 inverted microscope equipped with a 40× objective

differential interference contrast (DIC) optics and a Zeiss AxioCam 702 digital camera. We

counted both hyphae and pseudohyphae as filaments to make the image analysis consistent. *C.*

albicans cells were grown and induced to make hyphae in the same conditions for transcriptome,

proteome, and phosphoproteome analyses. Galactose was used as a carbon source because

glucose represses expression of the GlcNAc transporter *NGT1*, which limits hyphal induction

[68]. *C. albicans* cells were grown at 37°C to log phase (OD_{660} of 0.2 ~ 0.4) in SC + 50 mM

galactose medium. For hyphal induction, pre-warmed 0.5 M GlcNAc solution was added into

the culture to make final concentration of 50 mM and the cultures were incubated for 2 h at 37°C.

Cells were harvested by centrifugation, washed three times with ice-cold ultrapure water, snap

frozen on dry ice, and stored at –80°C. Replicate experiments were conducted independently

on different days.

Strain construction

Deletion mutant strains were created using transient expression of CRISPR-Cas9 to

facilitate the genome engineering and limit transformation-induced genomic changes in *C.*

albicans [41, 69]. The methods were performed essentially as described previously [41, 70].

Briefly, *CaCAS9* expression and sgRNA expression cassettes were co-transformed into cells

along with a repair template. The *CaCAS9* gene was codon optimized for expression in *C.*

albicans [71]. The *CaCAS9* expression cassette was PCR amplified from the plasmid pV1093,

which was a kind gift from Dr. Valmik Vyas [71]. The sgRNA expression cassette was

constructed through single-joint PCR from the plasmid pV1093 [41]. We used 20-bp target

sequences of the sgRNA, as reported previously by Vyas et al. [72] to target Cas9 to make a

DNA double strand break at the target site (S4 Table). When deleting large regions of

chromosome 2 for gene mapping, two sgRNAs were used to cut each side of the target region

(see Supplemental Methods). Repair templates were synthesized using the plasmid pGR-NAT

as a template, which contains the *SAT1* flipper cassette [42]. The primers were designed to

include 60 to 80 bases of homology to the sequences upstream or downstream from the target

region. The oligonucleotides used in this study are listed in S5 Table. PCR was conducted with

Ex Taq (TaKaRa Bio, Inc.) in accordance with the CRISPR protocol (Supplemental Methods).

BNI1 and *MOB2* were mutated to prevent phosphorylation at the indicated sites using

the CRISPR-Cas9 method developed by Vyas et al. [71] The 20-bp target sequence of sgRNA

was cloned into pV1093 and correct clones were verified by DNA sequencing. The CRISPR

expression plasmids were linearized by digestion with *Kpn1* and *Sac1* before transformation for

efficient targeting to the *ENO1* locus. Repair templates were generated with two 60-base

oligonucleotides containing a 20-base overlap at their 3′ ends centered on the desired site to

mutate a phosphorylation site. The repair templates were constructed by PCR primer extension

to contain the desired phospho-site mutation as well as a unique restriction site to facilitate

identification of appropriate transformants.

PCR products and linearized plasmids for transformation were purified and concentrated

using a GeneJET PCR purification kit (Thermo Fisher Scientific, Inc.). Electroporation was used

to introduce the DNA into cells following a previously described method [73]. An

electrocompetent cell suspension (40 µl) was added to aliquoted DNA, placed in 0.1 cm-gap

electroporation cuvettes, and electroporated on a Bio-Rad Gene Pulser at 1.5 kV. One milliliter

of 0.5× YPD containing 1 M sorbitol was added immediately to the cuvette, and then the cell

mixture was incubated for 3 h at 30°C before plating onto YPD + 200 µg/ml NAT agar. Nat^r

transformants were selected, and PCR genotyping of the transformants verified the genome

editing. When deleting large regions of chromosome 2 for gene mapping, we confirmed that the

deletions are heterozygous. As described in a previous paper [41], two PCR detection

strategies were designed to detect the wild-type allele and the deletion allele. More than 10

independent transformants were tested for each deletion and around half of them had both WT

and deletion alleles (heterozygous deletion). We did not find any homozygous deletion mutants,

presumably due to the essential genes present in these regions of chromosome 2.

Genome analysis

A 1 µg aliquot of the genomic DNA was prepared using the Illumina TruSeq PCR-free

DNA HT sample preparation kit with 450 bp insert size. Intact genomic DNA was sheared using

the Covaris sonicator (adaptive focused acoustics), followed by end-repair and bead-based size

selection of fragmented molecules. The size selected fragments then had adenines added to

the 3' ends of the DNA, Illumina sequence adaptors ligated onto the fragments, followed by

PCR amplification and final library QC. A majority of the steps in this process were performed

on the Caliper SciClone NGSx workstation (PerkinElmer), a robotics system developed and

validated for automated library preparation. The library QC included a measurement of the

average size of library fragments using the Agilent BioAnalyzer, estimation of the total

concentration of DNA by PicoGreen (ThermoFisher), and a measurement of the yield and

efficiency of the adaptor ligation process with a quantitative PCR assay (Kapa) using primers

specific to the adaptor sequence. Sequencing was performed on the Illumina MiSeq instrument

at 2×150 bp read length.

Reads were first trimmed and quality filtered using skewer v0.1.127 [74] with the

following parameters (-q 20 -Q 20 -l 50 -n yes). Reads were then mapped to the *C. albicans*

SC5314 genome (Assembly 21) [75] using BWA mem v0.7.15 [76]. Duplicates were marked

and read groups added using the PICARD MarkDuplicates and AddOrReplaceReadGroups

programs. Indels were realigned using GATK 3.3 RealignerTargerCreator and IndelRealigner

[77]. A set of high-quality variants with variant scores of >100 was identified using GATK 3.3

HaplotypeCaller on all samples simultaneously. These were then set as known sites when

performing base quality recalibration using GATK 3.3 PrintReads, AnalyzeCovariates and

BaseRecalibrator to produce finished bam files. GATK 3.3 HaplotypeCaller was then used

again to call variants across samples to produce a final vcf file. snpEff 4.2 was used to annotate

variants using a custom built database based on the *C. albicans* SC5314 genome (Assembly 21)

[75]. Previously, we did not report the change of ploidy for PR2 [14]. In the present study we

had higher read coverage (>90$\times$), so we were able to detect changes of the ploidy and to

identify the junction regions.

Transcriptome analysis

RNA extraction, library preparations, and sequencing reactions were conducted at

GENEWIZ, LLC. (South Plainfield, NJ, USA). Total RNA was extracted from fresh frozen cell

pellets using Qiagen RNeasy Plus Universal mini kit following manufacturer's instruction

(Qiagen). Extracted RNA samples were quantified using Qubit 2.0 Fluorometer (Life

Technologies) and RNA integrity was checked using Agilent TapeStation 4200 (Agilent

Technologies). RNA sequencing libraries were prepared using the NEBNext Ultra II RNA

Library Prep Kit for Illumina following manufacturer's instructions (NEB). Briefly, mRNAs were

first enriched with Oligo(dT) beads. Enriched mRNAs were fragmented for 15 min at 94°C.

First strand and second strand cDNAs were subsequently synthesized. cDNA fragments were

end repaired and adenylated at 3' ends, and universal adapters were ligated to cDNA fragments,

followed by index addition and library enrichment by limited-cycle PCR. The sequencing

libraries were validated on the Agilent TapeStation (Agilent Technologies), and quantified by

using Qubit 2.0 Fluorometer (Invitrogen) as well as by quantitative PCR (KAPA Biosystems).

The sequencing libraries were pooled and clustered on 1 lane of a flowcell. After clustering, the

flowcell was loaded on the Illumina HiSeq instrument (4000 or equivalent) according to

manufacturer's instructions. The samples were sequenced using a 2×150bp Paired End (PE)

configuration. Image analysis and base calling were conducted by the HiSeq Control Software

(HCS). Raw sequence data (.bcl files) generated from Illumina HiSeq was converted into fastq

files and de-multiplexed using Illumina's bcl2fastq 2.17 software. One mismatch was allowed for

index sequence identification.

We performed quality profiling, adapter trimming, read filtering, and base correction for

raw data using an all-in-one FASTQ preprocessor, *fastp* [78]. The high-quality paired-end reads

were mapped to the *C. albicans* SC5314 genome (Assembly 22) [79] using HISAT2 [80].

StringTie [81] was used to assemble the read alignments obtained in the previous step and

estimate transcript abundances. Absolute mRNA abundance was expressed as fragments per

kilobase of transcript per million mapped reads (FPKM). Differential expression analyses were

conducted using DESeq2 [82] package from Bioconductor [83] on R.

Western blotting

The fresh frozen cell pellets were lysed using the same volume of 2✕ SDS lysis buffer (5%

SDS, 20% glycerol, 125 mM Tris-HCl, pH 6.8) supplemented with cOmplete EDTA-free

protease inhibitor cocktail (Roche), 10 mM activated sodium orthovanadate, 2.5 mM sodium

pyrophosphate, 1 mM β-glycerophosphate, 1% phosphatase inhibitor cocktail 2 (Millipore-

Sigma), 1% phosphatase inhibitor cocktail 3 (Millipore-Sigma). Zirconia beads were added, and

the cells were mechanically disrupted by 3 rounds of 1-min bead beating and 1-min cooling on

ice. The samples were boiled at 95°C for 5 min. After centrifugation at 14,000 rpm for 5 min at

4°C to remove cellular debris, the supernatant was transferred to a new tube and this step was

repeated. The protein concentration was measured using a BCA protein assay kit (Thermo

Fisher Scientific). 2-Mercaptoethanol and bromophenol blue were added to 5% and 0.002%

final concentration, respectively. The samples were boiled at 95°C for 5 min. Equal amounts of

proteins were resolved by SDS-PAGE using the Mini-PROTEAN Tetra Cell system (Bio-Rad)

and Tris/glycine/SDS running buffer. Subsequently, proteins were transferred to a nitrocellulose

membrane (Amersham) using a semi-dry transfer apparatus. The membranes were blocked

with 5% nonfat dry milk in TBS-T buffer (20 mM Tris-HCl, pH 7.6, 150 mM NaCl, 0.2% Tween-

20) for 30 min and probed with CaBcy1 rabbit polyclonal antibody (GenScript) or CaTpk2 rabbit

polyclonal antibody (GenScript), at 1:1,000 dilution in 5% nonfat dry milk in TBS-T buffer plus

0.5% sodium azide for 2 hours at room temperature. The membranes were then washed three

times with TBS-T and incubated with an IRDye 800CW goat anti-rabbit IgG secondary antibody

(LI-COR Biosciences) diluted at 1:10,000 in TBS-T. The membranes were washed three times

with TBS-T, stored in TBS buffer, and visualized by scanning with an Odyssey CLx infrared

imaging system (LI-COR Biosciences). For Coomassie stained gels, SDS-PAGE gels were

performed as described and then stained in Coomassie Brilliant Blue solution (0.1% Coomassie

R-250, 50% methanol, 40% water, 10% glacial acetic acid) overnight. Gels were incubated for

6 h in destaining solution (50% water, 40% methanol, 10% glacial acetic acid), rinsed with water,

and analyzed by scanning with an Odyssey CLx infrared imaging system (LI-COR Biosciences).

Resulting images were analyzed and quantified using ImageStudio software (LI-COR

Biosciences).

Preparation of samples for mass spectrometry

The frozen cell pellet samples were processed by Tymora Analytical Operations (West

Lafayette, IN). For lysis, 200 µL of lysis buffer (8M urea in 50mM Tris-Cl pH 7.5, supplemented

with phosphatase inhibitor cocktail 3 [Millipore-Sigma]) was added to each of the pellets and

pipetted up and down several time to lyse a portion of the pellet. The samples were incubated

for 10 min at 37°C, pulse-sonicated several times with a sonicator probe, and incubated again

for 10 min at 37°C. The lysed samples were then centrifuged at 16,000 g for 10 min to remove

debris and the supernatant portions collected. BCA assay was carried out to calculate the

protein concentration and all samples were normalized by protein amount to 200 µg each. Then

5 mM dithiothreitol was added and the proteins were incubated at 37°C for 15 min to reduce the

cysteine residues, and then alkylated by incubation in 15 mM iodoacetamide at room

temperature for 45 min in the dark. The samples were diluted 3-fold with 50 mM

triethylammonium bicarbonate and digested with Lys-C (Wako) at 1:100 (w/w) enzyme-to-

protein ratio for 3 h at 37°C. The samples were further diluted 3-fold with 50 mM

triethylammonium bicarbonate and trypsin was added to a final 1:50 (w/w) enzyme-to-protein

ratio for overnight digestion at 37°C. After digestion, the samples were acidified with

trifluoroacetic acid to a pH < 3 and desalted using Top-Tip C18 tips (Glygen) according to

manufacturer's instructions. A 1% portion of each sample was used for direct proteomics

analysis and the remainder of each sample was used for phosphopeptide enrichment. The

samples were dried completely in a vacuum centrifuge and stored at -80°C. The 99% portion of

each sample was used for phosphopeptides enrichment using PolyMAC Phosphopeptide

Enrichment Kit (Tymora Analytical) according to the manufacturer's instructions.

Liquid chromatography with tandem mass spectrometry (LC-MS/MS) analysis

The full phosphopeptide sample and 1 µg of the peptide sample each was dissolved in

10.5 µl of 0.05% trifluoroacetic acid with 3% (v/v) acetonitrile containing spiked-in indexed

Retention Time Standard artificially synthetized peptides (Biognosys). The spiked-in 11-

peptides standard mixture was used to account for any variation in retention times and to

normalize abundance levels among samples. Ten microliters of each sample were injected into

an Ultimate 3000 nano UHPLC system (Thermo Fisher Scientific). Peptides were captured on a

2-cm Acclaim PepMap trap column and separated on a heated 50-cm Acclaim PepMap column

(Thermo Fisher Scientific) containing C18 resin. The mobile phase buffer consisted of 0.1%

formic acid in ultrapure water (buffer A) with an eluting buffer of 0.1% formic acid in 80% (v/v)

acetonitrile (buffer B) run with a linear 90-min gradient of 6–30% buffer B at flow rate of 300

nL/min. The UHPLC was coupled online with a Q-Exactive HF-X mass spectrometer (Thermo

Fisher Scientific). The mass spectrometer was operated in the data-dependent mode, in which

a full-scan MS (from m/z 375 to 1,500 with the resolution of 60,000) was followed by MS/MS of

the 15 most intense ions (30,000 resolution; normalized collision energy - 28%; automatic gain

control target [AGC] - 2E4, maximum injection time - 200 ms; 60 sec exclusion).

LC-MS data processing

The raw files were searched directly against the *C. albicans* database updated in 2019

with no redundant entries, using Byonic (Protein Metrics) and Sequest search engines loaded

into Proteome Discoverer 2.3 software (Thermo Fisher Scientific). MS1 precursor mass

tolerance was set at 10 ppm, and MS2 tolerance was set at 20ppm. Search criteria included a

static carbamidomethylation of cysteines (+57.0214 Da), and variable modifications of

phosphorylation of S, T and Y residues (+79.996 Da), oxidation (+15.9949 Da) on methionine

residues and acetylation (+42.011 Da) at N terminus of proteins. Search was performed with

full trypsin/P digestion and allowed a maximum of two missed cleavages on the peptides

analyzed from the sequence database. The false-discovery rates of proteins and peptides were

set at 0.01. All protein and peptide identifications were grouped and any redundant entries were

removed. Only unique peptides and unique master proteins were reported.

Label-free quantitation analysis

All data were quantified using the label-free quantitation node of Precursor Ions

Quantifier through the Proteome Discoverer v2.3 (Thermo Fisher Scientific). For the

quantification of proteomic and phosphoproteomic data, the intensities of peptides were

extracted with initial precursor mass tolerance set at 10 ppm, minimum number of isotope peaks

as 2, maximum ΔRT of isotope pattern multiplets – 0.2 min, PSM confidence FDR of 0.01, with

hypothesis test of ANOVA, maximum RT shift of 5 min, pairwise ratio-based ratio calculation,

and 100 as the maximum allowed fold change. The abundance levels of all

peptides/phosphopeptides and proteins/phosphoproteins were normalized to the spiked-in

internal iRT standard. For calculations of fold-change between the groups of proteins, total

protein or phosphoprotein abundance values were added together and the ratios of these sums

were used to compare proteins within different samples. When the protein abundance could not

be determined, pseudo value (1) was given for logarithm.

Phosphoproteomic data analysis

Phosphopeptide abundance values were normalized by protein abundance values [84,

85]. Fifteen-amino acid sequences around the phosphorylation site was extracted from the

phosphoproteome using a custom Microsoft Excel spreadsheet. rmotifx [86] package on R was

used to find overrepresented patterns from the 15-aa sequence set. Potential substrates of 6

kinases (Cdc28, PKA, Yck2, MAPK, Cbk1, and Gin4) were predicted from the

phosphoproteome based on their consensus phosphorylation motifs. The following consensus

motifs were retrieved from the Scansite 4.0 database [87]; Cdc28 ([S/T]-P-x-K/R); PKA

(R/K-R/K-x-[S/T]); Yck2 ([S/T]-x-x-[S/T]); MAPK (P-x-[S/T]-P); Cbk1 (H-x-K/R-x-x-[S/T] or H-x-x-

K/R-x-[S/T]); Gin4 (R-S-x-[S/T]). We next utilized a gene set enrichment analysis algorithm,

GSEA [50, 88], to infer kinase activity based on the substrate phosphorylation levels from the

phosphoproteome.

Data availability

Requests for materials should be addressed to the corresponding author. The datasets

generated in this study can be found under NCBI BioProject PRJNA706677, including whole-

genome sequence data and RNA-seq data. Source data are provided with this paper.

ACKNOWLEDGEMENTS

This work was supported by Public Health Service grants from the National Institutes of Health

awarded to J.B.K. (R01GM116048 and R01AI047837). We thank the members of our

laboratories for their helpful comments on the manuscript. We are grateful to Jihye An for help

with microscopic image analysis and Kyunghye Min for assistance with illustrations.

AUTHOR CONTRIBUTIONS

K.M. and J.B.K conceived the study. K.M., T.F.J., H.S., K.R.V., J.D.H., and J.B.K. investigated

the study. K.M., K.R.V., and J.D.H. formally analyzed the study. K.M. and J.B.K. wrote the

original draft of the manuscript. K.M., T.F.J., H.S., K.R.V., J.D.H., and J.B.K reviewed and

edited the manuscript. J.B.K supervised the study.

DECLARATION OF INTERESTS

The authors declare no competing interests

FIGURE CAPTIONS

Fig 1. Genetic changes in chromosome 2 bypassed the need for cAMP to improve growth and hyphal induction in *cyr1*Δ/Δ PR mutants.

(A) Doubling times were measured in liquid YPD medium at 30°C. Shown is the mean ± SD (standard deviation) of 4 independent experiments. Statistical analysis was performed using one-way ANOVA with Dunnett's multiple comparisons test comparing the strains with the wild-type control (WT); ^{NS} p > 0.05, * p < 0.05, ** p < 0.01, *** p < 0.001. The doubling time of the strains shown was significantly shorter when compared to the *cyr1*Δ/Δ background (p < 0.0001).

(B) The strains indicated at the top were grown in the liquid medium indicated on the left, and then hyphal induction was assessed microscopically. Cells were grown in liquid medium containing 15% serum or 50 mM N-acetylglucosamine (GlcNAc) to induce hyphal growth. Cells were incubated at 37°C for 3 h and then photographed. Scale bar, 10 μm.

(C) Copy number variation analysis based upon read depth across the genome. Copy number estimates scaled to genome ploidy (Y-axis) and chromosome location (X-axis) were plotted using YMAP [89]. Numbers and symbols below chromosomes indicate chromosomal position (Mb), *BCY1* gene (green arrows), centromere locus (indentations in the chromosome cartoon), major repeat sequence position (black circles), and rDNA locus (blue circles, ChrR).

Fig 2. Gene mapping by CRISPR-Cas9 demonstrates that haploinsufficiency of genes in chromosome 2 improved hyphal induction in *cyr1*Δ/Δ PR mutants.

(A) A schematic diagram showing deleted sequences on chromosome 2.

(B) The strains indicated at the top were grown in the liquid medium indicated on the left, and then hyphal induction was assessed microscopically. Cells were grown in liquid medium containing 15% serum or 50 mM GlcNAc to induce hyphal growth at 37°C for 3 h and then photographed. Scale bar, 10 μm.

(C) Doubling times were measured in liquid YPD medium at 30°C. Shown is the mean ± SD of

6 independent experiments. The doubling time of the strains shown was significantly shorter

when compared to the *cyr1Δ/Δ* background ($p < 0.01$).

(D) Graph indicating the percent of filamentous cells after growth in GlcNAc medium described

in panel B. Shown is the mean ± SD of at least 3 independent experiments with at least 100

cells counted for each condition.

(E) Gene mapping by CRISPR-Cas9 identified a 10-kb region (yellow square) that is involved in

the filamentous phenotype. The left panel shows deleted sequences on chromosome 2. The

right panel shows the percent of filamentous cells in liquid GlcNAc medium at 37°C; green,

weak hyphal induction; yellow, intermediate hyphal induction; pink, strong hyphal induction.

(A and E), Note that deletions are heterozygous; the cells retain a wild-type version of

chromosome 2. Numbers and symbols indicate gene deletion (Δ), CRISPR cut site (scissors),

large genomic deletion (dotted line), *BCY1* gene (green circles), and chromosomal position (kb).

(C and D) Statistical analysis was performed using one-way ANOVA with Dunnett's multiple

comparisons test comparing the strains with the WT or parental strain; $^{NS} p > 0.05$, $^{*} p < 0.05$, $^{**} p$

< 0.01, $^{***} p < 0.001$.

Fig 3. Adenylyl cyclase is not necessary for transcriptional regulation of hyphal-induced

genes.

(A) Experimental scheme of transcriptomic analysis. Cells were grown at 37°C in liquid

galactose medium and then 50 mM GlcNAc was added for 2 h to induce hyphae.

(B) Cluster analysis of differentially expressed genes (DEGs).

(C) Summary of differential expression analysis. The numbers in colored boxes are $\log_2$

difference of transcript levels compared to the wild-type –GlcNAc condition. Grey numbers

indicate the difference is not significantly different (adjusted $p > 0.1$).

Fig 4. Quantitative phosphoproteomics detected activity of two key protein kinases in hyphal induction.

(A) Experimental scheme of parallel proteomic and phosphoproteomic analyses. Cells were grown at 37°C in liquid galactose medium and then 50 mM GlcNAc was added for 2 h to induce hyphae. For phosphorylation analysis, phosphopeptides were enriched by polymer-based metal-ion affinity capture (PolyMAC).

(B) Representative null comparisons of the biological replicates show high reproducibility. We compared the replicate data of WT+GlcNAc_1 and WT+GlcNAc_2 from both whole proteome and phosphoproteome.

(C) Representative results of kinase-substrate enrichment analysis. Gene set enrichment analysis (GSEA) algorithm [50] was used to identify significantly enriched or depleted groups of phosphorylated substrates. The potential substrates of both Cdc28 and Yck2 kinases were more phosphorylated (up-regulated) during the hyphal induction in the WT and PR13 cells.

(D) The color-coded diagram illustrates changes of the selected kinase activities indicated by the scale bar at the bottom (< 25% false discovery rate [FDR]). Kinase activity inference was based on the collective phosphorylation changes of their identified substrates.

(E) Venn diagrams of phosphopeptides upregulated during hyphal induction ($\log_2$ fold change > 0.5). Note that 605 phosphopeptides were upregulated in both WT and PR13 strains.

(F) Overrepresented motifs were extracted from the 605 phosphopeptides, using the motif-x algorithm [86] ($p < 10^{-6}$). 44 phosphopeptides had the Cdc28 consensus motif ([S/T]-P-x-K/R) indicating they were potential substrates.

(G) Venn diagrams of the upregulated Cdc28 substrates ([S/T]-P-x-K/R) or Yck2 substrates (S/T-x-x-[S/T]). 11 phosphopeptides were potential substrates of both Cdc28 and Yck2 kinases.

Fig 5. Integrated transcriptomic and proteomic analyses indicate that hyphal stimulation induces translation of hyphal regulator TFs.

(A) A basic gene expression model with key parameters (adapted from REF. [56, 90]). The

mRNA is transcribed with rate V_{sr} and degraded with a half-life represented by $t_{r1/2}$. The protein

is translated proportionally to the mRNA abundance with the rate constant K_{sp} and degraded

with a half-life of $t_{p1/2}$.

(B) An across-gene correlation analysis comparing estimates of absolute mRNA abundance

(expressed in fragments per kilobase of transcript per million mapped reads (FPKM)) to protein

abundance (expressed as label free quantitation (LFQ)) in exponentially growing WT cells. r,

Pearson correlation coefficient.

(C) Mathematical expression of mRNA (Equation 1) and protein abundances (Equation 2) as a

function of key gene expression parameters, including cell doubling time T_{cc}, as detailed in REF.

[90]. Rearranging Equation 2 yields the dependency of protein-mRNA abundance ratio on the

translation rate constant K_{sp}, cell doubling time T_{cc}, and protein half-life $t_{p1/2}$ (Equation 3). We

neglected protein degradation assuming that protein replacement is generally driven by dilution

due to cell division in exponentially growing *C. albicans* cells ($t_{p1/2} \gg T_{cc}$). Substituting the cell

doubling time (T_{cc} = 1.5 h in *C. albicans*) yields Equation 4. Therefore, we can approximate the

translation rate with the protein-to-mRNA level ratio.

(D) Across-gene correlation analyses comparing protein-to-mRNA ratio versus mRNA

abundance, protein abundance, and 5' UTR length (from left to right). r, Pearson correlation

coefficient.

(E) Protein-to-mRNA ratios of 11 selected hyphal regulator transcription factors (TFs) during

GlcNAc induction in WT. The protein-to-mRNA ratios of the 5 hyphal regulators were two

orders of magnitude lower than the median in –GlcNAc control, indicating their translation rate

was very low before hyphal induction. The p-value was calculated using two-sided unpaired t-

test (hyphal regulator TFs) and z-test (ribosomal proteins and total proteins).

(F) The relative change in protein-to-mRNA ratio for the selected 11 hyphal regulators is shown

in yellow, relative changes in mRNA expression are shown in blue. The protein-to-mRNA ratio

of *UME6*, *SFL1*, *BRG1*, *TEC1*, and *FKH2* increased dramatically (log2 fold change > 4) during

hyphal induction while mRNA levels did not.

Fig 6. Integrative multi-omics profiling revealed new pathways that induce *C. albicans*

hyphal formation in a cAMP-independent manner.

New insights into the mechanisms of hyphal morphogenesis were obtained by performing

integrative multi-omics profiling in *C. albicans*. Genetic changes in PR mutants were identified

by performing genome analysis, gene-mapping by CRISPR-Cas9, and transcriptome analysis.

These studies showed that gene dosage effects on *BCY1* (negative regulator of PKA) and

global transcription mechanisms promoted hyphal growth in the absence of cAMP. Integrating

transcriptomic and proteomic data revealed that the translation rate of some hyphal regulatory

TFs increased dramatically during hyphal induction while their mRNA levels did not, suggesting

a novel mechanism of translational regulation promotes hyphal growth. Parallel comparisons of

protein production and phosphorylation revealed that phosphorylation by Hgc1-Cdc28 kinase

and yeast casein kinase 1 (Yck2) increased during hyphal induction, even in the absence of

adenylyl cyclase, consistent with genetic studies indicating that these kinases play an important

role in hyphal growth. Altogether, these data suggest a new model that various hyphal

inducers and TFs are predicted to induce a new physiological state that promotes polarized

filamentous growth, in part due to changes in protein phosphorylation.

SUPPLEMENTAL INFORMATION TITLES AND LEGENDS

S1 Fig. Hyphal induction rate of the *C. albicans* strains.

(A) The plot shows the percent of filamentous cells in YPD medium at 30°C (-control).

(B) Graph indicating the percent of filamentous cells after growth in GlcNAc medium. Cells

were grown in liquid medium containing 50 mM GlcNAc to induce hyphal growth at 37°C for 2 h

and then filamentous cells were counted.

(C) The percent of filamentous cells in YPD medium at 30°C; green, weak filamentation; yellow,

intermediate filamentation; pink, strong filamentation. Deletions indicated on the left are on

chromosome 2 and they are heterozygous; the cells retain a wild-type version of chromosome 2.

(A, B, and C) Shown is the mean ± SD of at least 3 independent experiments with at least 100

cells counted for each condition. Statistical analysis was performed using one-way ANOVA with

Dunnett's multiple comparisons test comparing the strains with the WT or parental strain; [NS] $p >$

0.05, [**] $p < 0.01$, [***] $p < 0.001$.

S2 Fig. Expression levels of PKA subunits during hyphal induction.

(A) Western blot detection of the negative regulatory subunit (Bcy1) and the catalytic subunit

(Tpk2) of PKA. Cells were grown at 37°C in liquid galactose medium and then 50 mM GlcNAc

was added for 2 h to induce hyphae. The sizes of the protein standards (kDa) are indicated on

the right of each blot. Images shown are representative of three independent experiments.

(B-D) Relative levels of Bcy1 (B), Tpk2 (C), and Tpk2/Bcy1 ratio (D) compared to the WT 0-h

samples. Shown is the mean ± SD of 3 independent experiments. Expression levels were

normalized to total proteins on Coomassie-stained gels. Statistical analysis was performed

using one-way ANOVA with Dunnett's multiple comparisons test comparing the strains with the

WT; [NS] $p > 0.05$, [*] $p < 0.05$, [**] $p < 0.01$, [***] $p < 0.001$.

S3 Fig. Gene-mapping analysis of the 10-kb region of chromosome 2 identified a role for

***SRB9* and *SPT5* in hyphal induction in the PR mutants.**

The single, double, and triple heterozygous deletion mutants of *SRB9*, *SPT5*, and *SSL1* were

created in KM20 strain background (*Chr2L 90kb→250kbΔ bcy1Δ/BCY1 cyr1Δ/Δ*). The plot

shows the percent of filamentous cells in GlcNAc medium. Cells were grown in liquid medium

containing 50 mM GlcNAc to induce hyphal growth at 37°C for 2 h and then filamentous cells

were counted.; yellow, intermediate hyphal induction; pink, strong hyphal induction. Statistical

analysis was performed using one-way ANOVA with Dunnett's multiple comparisons test

comparing the strains with the parental strain; [NS] $p > 0.01$, [***] $p < 0.001$.

S4 Fig. Correlation analysis and principal component analysis (PCA) of the normalized

RNA-seq dataset.

(A) Representative null comparisons of the biological replicates show very high reproducibility (r

≈ 1.00) in the RNA-seq dataset. Each dot represents the transcript level of individual gene in

the scatter plots. We compared the biological replicates of WT–control and WT+GlcNAc. rLog,

regularized log transformation.

(B) PCA plot shows clusters of biological replicates based on their similarity in transcriptome.

S5 Fig. Deletion of Cdc28 cyclin (*HGC1*) and casein kinase 1 (*YCK2*) disrupt normal

hyphal growth.

(A) Deletion of Cdc28 cyclin (*HGC1*) and casein kinase 1 (*YCK2*) disrupt normal hyphal growth

in WT and PR13 backgrounds.

(B) The plot shows the percent of filamentous cells in YPD medium at 30°C (-control) and after

2-h growth in GlcNAc medium at 37°C (+GlcNAc). Shown is the mean ± SD of at least 3

independent experiments with at least 100 cells counted for each condition. Statistical analysis

was performed using one-way ANOVA with Dunnett's multiple comparisons test comparing the

strains with the WT or parental strain; [ns] $p > 0.05$, [****] $p < 0.0001$.

(C) Phospho-mutants of Mob2 and Bni1 did not show an obvious defect in hyphal growth.

(A and C) The strains indicated at the top were grown in the liquid medium indicated on the left,

and then hyphal induction was assessed microscopically. Cells were grown in liquid medium

containing 15% serum or 50 mM N-acetylglucosamine (GlcNAc) to induce hyphal growth. Cells

were incubated at 37°C for 2 h and then photographed. Scale bar, 10 μm.

S6 Fig. Protein-to-mRNA ratios of hyphal regulator TFs during GlcNAc induction in PR13.

The relative change in protein-to-mRNA ratio for the selected 11 hyphal regulators TFs is shown

in yellow, relative changes in mRNA expression are shown in blue. The protein-to-mRNA ratio

of *RIM101* and *RFG1* increased dramatically (log2 fold change > 4) during hyphal induction

while mRNA levels did not.

S1 Table. Genome analysis summary.

[a] 23-bp telomere repeat sequence, CACCAAGAAGTTAGACATCCGTA

S2 Table. Potential phosphorylation substrates of Cdc28 and Yck2 during hyphal

induction

[a] Descriptions and deletion phenotypes were obtained from *Candida* Genome Database.

S3 Table. *C. albicans* strains used in this study

S4 Table. Target sequences of the sgRNAs

S5 Table. The oligonucleotides used in this study

S1 Data. SNP analysis of the PR mutants

S2 Data. RNA-seq data

S3 Data. Quantitative phosphoproteomic data

S4 Data. Whole-proteomic data and the protein-mRNA abundance correlation

832 **Table 1. Phenotypic and genetic variations of *cyr1Δ/Δ* pseudorevertants (PRs)**

Strain	PR class	Doubling time (h)[a]	Filamentation (%)[b]	Short genotype[c]
WT		1.4 ± 0.07	98 ± 2.3	Prototrophic wild type strain
cyr1Δ/Δ		3.4 ± 0.36 [***]	0.0 [***]	*cyr1Δ/Δ* (parental strain)
PR13, PR19	Class 1	1.5 ± 0.19	74 ± 4.4 [***]	*bcy1*/BCY1 cyr1Δ/Δ*
PR12, PR14	Class 2	1.8 ± 0.09 [**]	93 ± 2.3	Monosomy of ~276 kb of Chr2L; *cyr1Δ/Δ*
PR18	Class 3	2.3 ± 0.06 [***]	93 ± 4.2	Monosomy of ~590 kb of Chr2L; *cyr1Δ/Δ*
PR16	Class 3+	2.4 ± 0.18 [***]	82 ± 16.4 [**]	Monosomy of ~557 kb of Chr2L; trisomy of ~1.3 Mb of Chr2; *cyr1Δ/Δ*

[a] Doubling times were measured in liquid YPD medium at 30°C. Shown is the mean ± SD of 4 independent experiments.

[b] The percent of filamentous cells was measured 3 h after growth in GlcNAc medium at 37°C. Shown is the mean ± SD of at least 3

independent experiments with at least 100 cells counted for each condition.

[c] An asterisk indicates a nonsense mutation. Chr2L, left arm of chromosome 2.

The p-value was calculated using one-way ANOVA with Dunnett's multiple comparisons test comparing the strains with the WT

control; [**]$p < 0.01$, [***]$p < 0.001$.

Fig 1. Genetic changes in chromosome 2 bypassed the need for cAMP to improve growth and hyphal induction in *cyr1Δ/Δ* PR mutants.

Fig 2. Gene mapping by CRISPR-Cas9 demonstrates that haploinsufficiency of genes in chromosome 2 improved hyphal induction in *cyr1*Δ/Δ PR mutants.

856 **Fig 3. Adenylyl cyclase is not necessary for transcriptional regulation of hyphal-induced**

857 **genes.**

Fig 4. Quantitative phosphoproteomics detected activity of two key protein kinases in

hyphal induction.

Fig 5. Integrated transcriptomic and proteomic analyses indicate that hyphal stimulation

induces translation of hyphal regulator TFs.

872 **Fig 6. Integrative multi-omics profiling revealed new pathways that induce *C. albicans***

873 **hyphal formation in a cAMP-independent manner.**

REFERENCES

1.   Min K, Neiman AM, Konopka JB. Fungal pathogens: Shape-shifting invaders. Trends Microbiol. 2020;28(11):922-33. Epub 2020/06/01. doi: 10.1016/j.tim.2020.05.001. PubMed PMID: 32474010; PubMed Central PMCID: PMCPMC7572602.

2.   Noble SM, Gianetti BA, Witchley JN. *Candida albicans* cell-type switching and functional plasticity in the mammalian host. Nat Rev Microbiol. 2017;15(2):96-108. Epub 2016/11/22. doi: 10.1038/nrmicro.2016.157. PubMed PMID: 27867199; PubMed Central PMCID: PMCPMC5957277.

3.   Sudbery PE. Growth of *Candida albicans* hyphae. Nat Rev Microbiol. 2011;9(10):737-48. Epub 2011/08/17. doi: 10.1038/nrmicro2636. PubMed PMID: 21844880.

4.   da Silva Dantas A, Lee KK, Raziunaite I, Schaefer K, Wagener J, Yadav B, et al. Cell biology of *Candida albicans*-host interactions. Curr Opin Microbiol. 2016;34:111-8. Epub 2016/11/05. doi: 10.1016/j.mib.2016.08.006. PubMed PMID: 27689902; PubMed Central PMCID: PMCPMC5660506.

5.   Lu Y, Su C, Liu H. *Candida albicans* hyphal initiation and elongation. Trends Microbiol. 2014;22(12):707-14. Epub 2014/09/30. doi: 10.1016/j.tim.2014.09.001. PubMed PMID: 25262420; PubMed Central PMCID: PMCPMC4256103.

6.   Naseem S, Douglas LM, Konopka JB. *Candida albicans rvs161Δ* and *rvs167Δ* endocytosis mutants are defective in invasion into the oral cavity. mBio. 2019;10(6):e02503-19. Epub 2019/11/14. doi: 10.1128/mBio.02503-19. PubMed PMID: 31719181; PubMed Central PMCID: PMCPMC6851284.

7.   Biswas S, Van Dijck P, Datta A. Environmental sensing and signal transduction pathways regulating morphopathogenic determinants of *Candida albicans*. Microbiol Mol Biol Rev. 2007;71(2):348-76. Epub 2007/06/08. doi: 10.1128/MMBR.00009-06. PubMed PMID: 17554048; PubMed Central PMCID: PMCPMC1899878.

8.   Min K, Naseem S, Konopka JB. N-Acetylglucosamine regulates morphogenesis and virulence pathways in fungi. J Fungi (Basel). 2019;6(1):8. Epub 2019/12/28. doi: 10.3390/jof6010008. PubMed PMID: 31878148; PubMed Central PMCID: PMCPMC7151181.

9.   Wang Y. Fungal adenylyl cyclase acts as a signal sensor and integrator and plays a central role in interaction with bacteria. PLoS Pathog. 2013;9(10):e1003612. Epub 2013/10/17. doi: 10.1371/journal.ppat.1003612. PubMed PMID: 24130478; PubMed Central PMCID: PMCPMC3795026.

10.   Niimi M. Dibutyryl cyclic AMP-enhanced germ tube formation in exponentially growing *Candida albicans* cells. Fungal Genet Biol. 1996;20(1):79-83. Epub 1996/03/01. doi: 10.1006/fgbi.1996.0013. PubMed PMID: 8634947.

11.   Huang G, Huang Q, Wei Y, Wang Y, Du H. Multiple roles and diverse regulation of the Ras/cAMP/protein kinase A pathway in *Candida albicans*. Mol Microbiol. 2019;111(1):6-16. Epub 2018/10/10. doi: 10.1111/mmi.14148. PubMed PMID: 30299574.

12.   Rocha CR, Schroppel K, Harcus D, Marcil A, Dignard D, Taylor BN, et al. Signaling through adenylyl cyclase is essential for hyphal growth and virulence in the pathogenic fungus *Candida albicans*. Mol Biol Cell. 2001;12(11):3631-43. Epub 2001/11/06. doi: 10.1091/mbc.12.11.3631. PubMed PMID: 11694594; PubMed Central PMCID: PMCPMC60281.

13.   Harcus D, Nantel A, Marcil A, Rigby T, Whiteway M. Transcription profiling of cyclic AMP signaling in *Candida albicans*. Mol Biol Cell. 2004;15(10):4490-9. Epub 2004/07/23. doi: 10.1091/mbc.e04-02-0144. PubMed PMID: 15269278; PubMed Central PMCID: PMCPMC519143.

14.   Parrino SM, Si H, Naseem S, Groudan K, Gardin J, Konopka JB. cAMP-independent signal pathways stimulate hyphal morphogenesis in *Candida albicans*. Mol Microbiol. 2017;103(5):764-79. Epub 2016/11/27. doi: 10.1111/mmi.13588. PubMed PMID: 27888610; PubMed Central PMCID: PMCPMC5323341.

15. Lu Y, Su C, Solis NV, Filler SG, Liu H. Synergistic regulation of hyphal elongation by
hypoxia, CO_2, and nutrient conditions controls the virulence of *Candida albicans*. Cell Host
Microbe. 2013;14(5):499-509. Epub 2013/11/19. doi: 10.1016/j.chom.2013.10.008. PubMed
PMID: 24237696; PubMed Central PMCID: PMCPMC4049569.
16. Carlisle PL, Banerjee M, Lazzell A, Monteagudo C, Lopez-Ribot JL, Kadosh D.
Expression levels of a filament-specific transcriptional regulator are sufficient to determine
Candida albicans morphology and virulence. Proc Natl Acad Sci U S A. 2009;106(2):599-604.
Epub 2009/01/01. doi: 10.1073/pnas.0804061106. PubMed PMID: 19116272; PubMed Central
PMCID: PMCPMC2626749.
17. Bockmuhl DP, Ernst JF. A potential phosphorylation site for an A-type kinase in the Efg1
regulator protein contributes to hyphal morphogenesis of *Candida albicans*. Genetics.
2001;157(4):1523-30. Epub 2001/04/06. PubMed PMID: 11290709; PubMed Central PMCID:
PMCPMC1461612.
18. Lassak T, Schneider E, Bussmann M, Kurtz D, Manak JR, Srikantha T, et al. Target
specificity of the *Candida albicans* Efg1 regulator. Mol Microbiol. 2011;82(3):602-18. Epub
2011/09/20. doi: 10.1111/j.1365-2958.2011.07837.x. PubMed PMID: 21923768.
19. Martin R, Albrecht-Eckardt D, Brunke S, Hube B, Hunniger K, Kurzai O. A core
filamentation response network in *Candida albicans* is restricted to eight genes. PLoS One.
2013;8(3):e58613. Epub 2013/03/22. doi: 10.1371/journal.pone.0058613. PubMed PMID:
23516516; PubMed Central PMCID: PMCPMC3597736.
20. Naseem S, Araya E, Konopka JB. Hyphal growth in *Candida albicans* does not require
induction of hyphal-specific gene expression. Mol Biol Cell. 2015;26(6):1174-87. Epub
2015/01/23. doi: 10.1091/mbc.E14-08-1312. PubMed PMID: 25609092; PubMed Central
PMCID: PMCPMC4357515.
21. Giusani AD, Vinces M, Kumamoto CA. Invasive filamentous growth of *Candida albicans*
is promoted by Czf1p-dependent relief of Efg1p-mediated repression. Genetics.
2002;160(4):1749-53. Epub 2002/04/26. PubMed PMID: 11973327; PubMed Central PMCID:
PMCPMC1462044.
22. Chen C, Zeng G, Wang Y. G1 and S phase arrest in *Candida albicans* induces
filamentous growth via distinct mechanisms. Mol Microbiol. 2018;110(2):191-203. Epub
2018/08/08. doi: 10.1111/mmi.14097. PubMed PMID: 30084240.
23. Woolford CA, Lagree K, Xu W, Aleynikov T, Adhikari H, Sanchez H, et al. Bypass of
Candida albicans filamentation/biofilm regulators through diminished expression of protein
kinase Cak1. PLoS Genet. 2016;12(12):e1006487. Epub 2016/12/10. doi:
10.1371/journal.pgen.1006487. PubMed PMID: 27935965; PubMed Central PMCID:
PMCPMC5147786.
24. Rai LS, Singha R, Sanchez H, Chakraborty T, Chand B, Bachellier-Bassi S, et al. The
Candida albicans biofilm gene circuit modulated at the chromatin level by a recent molecular
histone innovation. PLoS Biol. 2019;17(8):e3000422. Epub 2019/08/10. doi:
10.1371/journal.pbio.3000422. PubMed PMID: 31398188; PubMed Central PMCID:
PMCPMC6703697.
25. Wartenberg A, Linde J, Martin R, Schreiner M, Horn F, Jacobsen ID, et al.
Microevolution of *Candida albicans* in macrophages restores filamentation in a nonfilamentous
mutant. PLoS Genet. 2014;10(12):e1004824. Epub 2014/12/05. doi:
10.1371/journal.pgen.1004824. PubMed PMID: 25474009; PubMed Central PMCID:
PMCPMC4256171.
26. Wang Y. Hgc1-Cdc28-how much does a single protein kinase do in the regulation of
hyphal development in *Candida albicans*? J Microbiol. 2016;54(3):170-7. Epub 2016/02/28. doi:
10.1007/s12275-016-5550-9. PubMed PMID: 26920877.
27. Zheng XD, Lee RT, Wang YM, Lin QS, Wang Y. Phosphorylation of Rga2, a Cdc42 GAP,
by CDK/Hgc1 is crucial for *Candida albicans* hyphal growth. EMBO J. 2007;26(16):3760-9.

Epub 2007/08/04. doi: 10.1038/sj.emboj.7601814. PubMed PMID: 17673907; PubMed Central
PMCID: PMCPMC1952229.
28.   Bishop A, Lane R, Beniston R, Chapa-y-Lazo B, Smythe C, Sudbery P. Hyphal growth in
Candida albicans requires the phosphorylation of Sec2 by the Cdc28-Ccn1/Hgc1 kinase. EMBO
J. 2010;29(17):2930-42. Epub 2010/07/20. doi: 10.1038/emboj.2010.158. PubMed PMID:
20639857; PubMed Central PMCID: PMCPMC2944046.
29.   Caballero-Lima D, Sudbery PE. In *Candida albicans*, phosphorylation of Exo84 by Cdk1-
Hgc1 is necessary for efficient hyphal extension. Mol Biol Cell. 2014;25(7):1097-110. Epub
2014/02/07. doi: 10.1091/mbc.E13-11-0688. PubMed PMID: 24501427; PubMed Central
PMCID: PMCPMC3967973.
30.   Sinha I, Wang YM, Philp R, Li CR, Yap WH, Wang Y. Cyclin-dependent kinases control
septin phosphorylation in *Candida albicans* hyphal development. Dev Cell. 2007;13(3):421-32.
Epub 2007/09/04. doi: 10.1016/j.devcel.2007.06.011. PubMed PMID: 17765684.
31.   Zeng G, Wang YM, Wang Y. Cdc28-Cln3 phosphorylation of Sla1 regulates actin patch
dynamics in different modes of fungal growth. Mol Biol Cell. 2012;23(17):3485-97. Epub
2012/07/13. doi: 10.1091/mbc.E12-03-0231. PubMed PMID: 22787279; PubMed Central
PMCID: PMCPMC3431942.
32.   Gutierrez-Escribano P, Gonzalez-Novo A, Suarez MB, Li CR, Wang Y, de Aldana CR, et
al. CDK-dependent phosphorylation of Mob2 is essential for hyphal development in *Candida
albicans*. Mol Biol Cell. 2011;22(14):2458-69. Epub 2011/05/20. doi: 10.1091/mbc.E11-03-0205.
PubMed PMID: 21593210; PubMed Central PMCID: PMCPMC3135472.
33.   Gonzalez-Novo A, Correa-Bordes J, Labrador L, Sanchez M, Vazquez de Aldana CR,
Jimenez J. Sep7 is essential to modify septin ring dynamics and inhibit cell separation during
Candida albicans hyphal growth. Mol Biol Cell. 2008;19(4):1509-18. Epub 2008/02/01. doi:
10.1091/mbc.E07-09-0876. PubMed PMID: 18234840; PubMed Central PMCID:
PMCPMC2291409.
34.   Wang A, Raniga PP, Lane S, Lu Y, Liu H. Hyphal chain formation in *Candida albicans*:
Cdc28-Hgc1 phosphorylation of Efg1 represses cell separation genes. Mol Cell Biol.
2009;29(16):4406-16. Epub 2009/06/17. doi: 10.1128/MCB.01502-08. PubMed PMID:
19528234; PubMed Central PMCID: PMCPMC2725740.
35.   Matsumoto K, Uno I, Oshima Y, Ishikawa T. Isolation and characterization of yeast
mutants deficient in adenylate cyclase and cAMP-dependent protein kinase. Proc Natl Acad Sci
U S A. 1982;79(7):2355-9. Epub 1982/04/01. doi: 10.1073/pnas.79.7.2355. PubMed PMID:
6285379; PubMed Central PMCID: PMCPMC346192.
36.   McEachern MJ, Hicks JB. Unusually large telomeric repeats in the yeast *Candida
albicans*. Mol Cell Biol. 1993;13(1):551-60. Epub 1993/01/01. doi: 10.1128/mcb.13.1.551.
PubMed PMID: 8417351; PubMed Central PMCID: PMCPMC358934.
37.   Todd RT, Wikoff TD, Forche A, Selmecki A. Genome plasticity in *Candida albicans* is
driven by long repeat sequences. Elife. 2019;8:e45954. Epub 2019/06/08. doi:
10.7554/eLife.45954. PubMed PMID: 31172944; PubMed Central PMCID: PMCPMC6591007.
38.   Andaluz E, Bellido A, Gomez-Raja J, Selmecki A, Bouchonville K, Calderone R, et al.
Rad52 function prevents chromosome loss and truncation in *Candida albicans*. Mol Microbiol.
2011;79(6):1462-82. Epub 2011/01/29. doi: 10.1111/j.1365-2958.2011.07532.x. PubMed PMID:
21272099; PubMed Central PMCID: PMCPMC3564047.
39.   Pennaneach V, Putnam CD, Kolodner RD. Chromosome healing by de novo telomere
addition in *Saccharomyces cerevisiae*. Mol Microbiol. 2006;59(5):1357-68. Epub 2006/02/14.
doi: 10.1111/j.1365-2958.2006.05026.x. PubMed PMID: 16468981.
40.   Legrand M, Jaitly P, Feri A, d'Enfert C, Sanyal K. *Candida albicans*: an emerging yeast
model to study eukaryotic genome plasticity. Trends Genet. 2019;35(4):292-307. Epub
2019/03/04. doi: 10.1016/j.tig.2019.01.005. PubMed PMID: 30826131.

41.   Min K, Ichikawa Y, Woolford CA, Mitchell AP. *Candida albicans* gene deletion with a
transient CRISPR-Cas9 system. mSphere. 2016;1(3):e00130-16. Epub 2016/06/25. doi:
10.1128/mSphere.00130-16. PubMed PMID: 27340698; PubMed Central PMCID:
PMCPMC4911798.
42.   Reuss O, Vik A, Kolter R, Morschhauser J. The *SAT1* flipper, an optimized tool for gene
disruption in *Candida albicans*. Gene. 2004;341:119-27. Epub 2004/10/12. doi:
10.1016/j.gene.2004.06.021. PubMed PMID: 15474295.
43.   Mani R, St Onge RP, Hartman JLt, Giaever G, Roth FP. Defining genetic interaction.
Proc Natl Acad Sci U S A. 2008;105(9):3461-6. Epub 2008/02/29. doi:
10.1073/pnas.0712255105. PubMed PMID: 18305163; PubMed Central PMCID:
PMCPMC2265146.
44.   Glazier VE, Murante T, Murante D, Koselny K, Liu Y, Kim D, et al. Genetic analysis of
the *Candida albicans* biofilm transcription factor network using simple and complex
haploinsufficiency. PLoS Genet. 2017;13(8):e1006948. Epub 2017/08/10. doi:
10.1371/journal.pgen.1006948. PubMed PMID: 28793308; PubMed Central PMCID:
PMCPMC5565191.
45.   Willger SD, Liu Z, Olarte RA, Adamo ME, Stajich JE, Myers LC, et al. Analysis of the
Candida albicans Phosphoproteome. Eukaryot Cell. 2015;14(5):474-85. Epub 2015/03/10. doi:
10.1128/EC.00011-15. PubMed PMID: 25750214; PubMed Central PMCID: PMCPMC4421004.
46.   Su C, Lu Y, Liu H. N-acetylglucosamine sensing by a GCN5-related *N*-acetyltransferase
induces transcription via chromatin histone acetylation in fungi. Nat Commun. 2016;7:12916.
Epub 2016/10/04. doi: 10.1038/ncomms12916. PubMed PMID: 27694804; PubMed Central
PMCID: PMCPMC5063960.
47.   Naseem S, Gunasekera A, Araya E, Konopka JB. N-Acetylglucosamine (GlcNAc)
induction of hyphal morphogenesis and transcriptional responses in *Candida albicans* are not
dependent on its metabolism. J Biol Chem. 2011;286(33):28671-80. Epub 2011/06/28. doi:
10.1074/jbc.M111.249854. PubMed PMID: 21700702; PubMed Central PMCID:
PMCPMC3190674.
48.   Cao C, Wu M, Bing J, Tao L, Ding X, Liu X, et al. Global regulatory roles of the
cAMP/PKA pathway revealed by phenotypic, transcriptomic and phosphoproteomic analyses in
a null mutant of the PKA catalytic subunit in *Candida albicans*. Mol Microbiol. 2017;105(1):46-64.
Epub 2017/04/04. doi: 10.1111/mmi.13681. PubMed PMID: 28370450.
49.   Frazer C, Staples MI, Kim Y, Hirakawa M, Dowell MA, Johnson NV, et al. Epigenetic cell
fate in *Candida albicans* is controlled by transcription factor condensates acting at super-
enhancer-like elements. Nature Microbiol. 2020;5(11):1374-89. doi: 10.1038/s41564-020-0760-
7.
50.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene
set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression
profiles. Proc Natl Acad Sci U S A. 2005;102(43):15545-50. Epub 2005/10/04. doi:
10.1073/pnas.0506580102. PubMed PMID: 16199517; PubMed Central PMCID:
PMCPMC1239896.
51.   Greig JA, Sudbery IM, Richardson JP, Naglik JR, Wang Y, Sudbery PE. Cell cycle-
independent phospho-regulation of Fkh2 during hyphal growth regulates *Candida albicans*
pathogenesis. PLoS Pathog. 2015;11(1):e1004630. Epub 2015/01/27. doi:
10.1371/journal.ppat.1004630. PubMed PMID: 25617770; PubMed Central PMCID:
PMCPMC4305328.
52.   Blankenship JR, Fanning S, Hamaker JJ, Mitchell AP. An extensive circuitry for cell wall
regulation in *Candida albicans*. PLoS Pathog. 2010;6(2):e1000752. Epub 2010/02/09. doi:
10.1371/journal.ppat.1000752. PubMed PMID: 20140194; PubMed Central PMCID:
PMCPMC2816693.

53.   Jung SI, Rodriguez N, Irrizary J, Liboro K, Bogarin T, Macias M, et al. Yeast casein
kinase 2 governs morphology, biofilm formation, cell wall integrity, and host cell damage of
Candida albicans. PLoS One. 2017;12(11):e0187721. Epub 2017/11/07. doi:
10.1371/journal.pone.0187721. PubMed PMID: 29107946; PubMed Central PMCID:
PMCPMC5673188.
54.   Caplan T, Lorente-Macias A, Stogios PJ, Evdokimova E, Hyde S, Wellington MA, et al.
Overcoming fungal echinocandin resistance through inhibition of the non-essential stress kinase
Yck2. Cell Chem Biol. 2020;27(3):269-82. Epub 2020/01/12. doi:
10.1016/j.chembiol.2019.12.008. PubMed PMID: 31924499; PubMed Central PMCID:
PMCPMC7093242.
55.   Li CR, Wang YM, De Zheng X, Liang HY, Tang JC, Wang Y. The formin family protein
CaBni1p has a role in cell polarity control during both yeast and hyphal growth in *Candida*
albicans. J Cell Sci. 2005;118(Pt 12):2637-48. Epub 2005/05/26. doi: 10.1242/jcs.02393.
PubMed PMID: 15914538.
56.   Buccitelli C, Selbach M. mRNAs, proteins and the emerging principles of gene
expression control. Nat Rev Genet. 2020;21(10):630-44. Epub 2020/07/28. doi:
10.1038/s41576-020-0258-4. PubMed PMID: 32709985.
57.   Carlisle PL, Kadosh D. *Candida albicans* Ume6, a filament-specific transcriptional
regulator, directs hyphal growth via a pathway involving Hgc1 cyclin-related protein. Eukaryot
Cell. 2010;9(9):1320-8. Epub 2010/07/27. doi: 10.1128/EC.00046-10. PubMed PMID: 20656912;
PubMed Central PMCID: PMCPMC2937344.
58.   Khajuria RK, Munschauer M, Ulirsch JC, Fiorini C, Ludwig LS, McFarland SK, et al.
Ribosome levels selectively regulate translation and lineage commitment in human
hematopoiesis. Cell. 2018;173(1):90-103. Epub 2018/03/20. doi: 10.1016/j.cell.2018.02.036.
PubMed PMID: 29551269; PubMed Central PMCID: PMCPMC5866246.
59.   Desai PR, Lengeler K, Kapitan M, Janssen SM, Alepuz P, Jacobsen ID, et al. The 5'
untranslated region of the *EFG1* transcript promotes its translation to regulate hyphal
morphogenesis in *Candida albicans*. mSphere. 2018;3(4):e00280-18. Epub 2018/07/07. doi:
10.1128/mSphere.00280-18. PubMed PMID: 29976646; PubMed Central PMCID:
PMCPMC6034079.
60.   Hossain S, Lash E, Veri AO, Cowen LE. Functional connections between cell cycle and
proteostasis in the regulation of *Candida albicans* morphogenesis. Cell Reports.
2021;34(8):108781. doi: https://doi.org/10.1016/j.celrep.2021.108781.
61.   Lu Y, Su C, Ray S, Yuan Y, Liu H. CO2 signaling through the Ptc2-Ssn3 axis governs
sustained hyphal development of *Candida albicans* by reducing Ume6 phosphorylation and
degradation. mBio. 2019;10(1):e02320-18. Epub 2019/01/17. doi: 10.1128/mBio.02320-18.
PubMed PMID: 30647154; PubMed Central PMCID: PMCPMC6336421.
62.   Selmecki A, Forche A, Berman J. Aneuploidy and isochromosome formation in drug-
resistant *Candida albicans*. Science. 2006;313(5785):367-70. Epub 2006/07/22. doi:
10.1126/science.1128242. PubMed PMID: 16857942; PubMed Central PMCID:
PMCPMC1717021.
63.   Todd RT, Selmecki A. Expandable and reversible copy number amplification drives rapid
adaptation to antifungal drugs. Elife. 2020;9:e58349. Epub 2020/07/21. doi:
10.7554/eLife.58349. PubMed PMID: 32687060; PubMed Central PMCID: PMCPMC7371428.
64.   Braun BR, Johnson AD. Control of filament formation in *Candida albicans* by the
transcriptional repressor *TUP1*. Science. 1997;277(5322):105-9. Epub 1997/07/04. doi:
10.1126/science.277.5322.105. PubMed PMID: 9204892.
65.   Grahl N, Demers EG, Lindsay AK, Harty CE, Willger SD, Piispanen AE, et al.
Mitochondrial activity and Cyr1 are key regulators of Ras1 activation of *C. albicans* virulence
pathways. PLoS Pathog. 2015;11(8):e1005133. Epub 2015/09/01. doi:

10.1371/journal.ppat.1005133. PubMed PMID: 26317337; PubMed Central PMCID: PMCPMC4552728.

66.   Arkowitz RA, Bassilana M. Recent advances in understanding *Candida albicans* hyphal growth. F1000Res. 2019;8. Epub 2019/05/28. doi: 10.12688/f1000research.18546.1. PubMed PMID: 31131089; PubMed Central PMCID: PMCPMC6530606.

67.   Tao L, Zhang Y, Fan S, Nobile CJ, Guan G, Huang G. Integration of the tricarboxylic acid (TCA) cycle with cAMP signaling and Sfl2 pathways in the regulation of CO_2 sensing and hyphal development in *Candida albicans*. PLoS Genet. 2017;13(8):e1006949. Epub 2017/08/09. doi: 10.1371/journal.pgen.1006949. PubMed PMID: 28787458; PubMed Central PMCID: PMCPMC5567665.

68.   Alvarez FJ, Konopka JB. Identification of an N-acetylglucosamine transporter that mediates hyphal induction in *Candida albicans*. Mol Biol Cell. 2007;18(3):965-75. Epub 2006/12/29. doi: 10.1091/mbc.e06-10-0931. PubMed PMID: 17192409; PubMed Central PMCID: PMCPMC1805087.

69.   Marton T, Maufrais C, d'Enfert C, Legrand M. Use of CRISPR-Cas9 to target homologous recombination limits transformation-induced genomic changes in *Candida albicans*. mSphere. 2020;5(5):e00620-20. Epub 2020/09/04. doi: 10.1128/mSphere.00620-20. PubMed PMID: 32878930; PubMed Central PMCID: PMCPMC7471004.

70.   Min K, Biermann A, Hogan DA, Konopka JB. Genetic analysis of *NDT80* family transcription factors in *Candida albicans* using new CRISPR-Cas9 approaches. mSphere. 2018;3(6):e00545-18. Epub 2018/11/23. doi: 10.1128/mSphere.00545-18. PubMed PMID: 30463924; PubMed Central PMCID: PMCPMC6249646.

71.   Vyas VK, Barrasa MI, Fink GR. A *Candida albicans* CRISPR system permits genetic engineering of essential genes and gene families. Sci Adv. 2015;1(3):e1500248. Epub 2015/05/16. doi: 10.1126/sciadv.1500248. PubMed PMID: 25977940; PubMed Central PMCID: PMCPMC4428347.

72.   Vyas VK, Bushkin GG, Bernstein DA, Getz MA, Sewastianik M, Barrasa MI, et al. New CRISPR mutagenesis strategies reveal variation in repair mechanisms among fungi. mSphere. 2018;3(2):e00154-18. Epub 2018/04/27. doi: 10.1128/mSphere.00154-18. PubMed PMID: 29695624; PubMed Central PMCID: PMCPMC5917429.

73.   De Backer MD, Maes D, Vandoninck S, Logghe M, Contreras R, Luyten WH. Transformation of *Candida albicans* by electroporation. Yeast. 1999;15(15):1609-18. Epub 1999/11/26. doi: 10.1002/(sici)1097-0061(199911)15:15<1609::aid-yea485>3.3.co;2-p. PubMed PMID: 10572258.

74.   Jiang H, Lei R, Ding SW, Zhu S. Skewer: a fast and accurate adapter trimmer for next-generation sequencing paired-end reads. BMC Bioinformatics. 2014;15:182. Epub 2014/06/14. doi: 10.1186/1471-2105-15-182. PubMed PMID: 24925680; PubMed Central PMCID: PMCPMC4074385.

75.   Skrzypek MSB, J.; Binkley, G.; Miyasato, S.R.; Simison, M.; Sherlock, G. Candida Genome Database 2017 [cited 2017 2017]. Available from: http://www.candidagenome.org.

76.   Li H, Durbin R. Fast and accurate long-read alignment with Burrows-Wheeler transform. Bioinformatics. 2010;26(5):589-95. Epub 2010/01/19. doi: 10.1093/bioinformatics/btp698. PubMed PMID: 20080505; PubMed Central PMCID: PMCPMC2828108.

77.   DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, et al. A framework for variation discovery and genotyping using next-generation DNA sequencing data. Nat Genet. 2011;43(5):491-8. Epub 2011/04/12. doi: 10.1038/ng.806. PubMed PMID: 21478889; PubMed Central PMCID: PMCPMC3083463.

78.   Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics. 2018;34(17):i884-i90. Epub 2018/11/14. doi: 10.1093/bioinformatics/bty560. PubMed PMID: 30423086; PubMed Central PMCID: PMCPMC6129281.

79.   Skrzypek MS, Binkley J, Binkley G, Miyasato SR, Simison M, Sherlock G. The Candida
Genome Database (CGD): incorporation of Assembly 22, systematic identifiers and visualization
of high throughput sequencing data. Nucleic Acids Res. 2017;45(D1):D592-D6. Epub
2016/10/16. doi: 10.1093/nar/gkw924. PubMed PMID: 27738138; PubMed Central PMCID:
PMCPMC5210628.
80.   Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory
requirements. Nat Methods. 2015;12(4):357-60. Epub 2015/03/10. doi: 10.1038/nmeth.3317.
PubMed PMID: 25751142; PubMed Central PMCID: PMCPMC4655817.
81.   Pertea M, Kim D, Pertea GM, Leek JT, Salzberg SL. Transcript-level expression analysis
of RNA-seq experiments with HISAT, StringTie and Ballgown. Nat Protoc. 2016;11(9):1650-67.
Epub 2016/08/26. doi: 10.1038/nprot.2016.095. PubMed PMID: 27560171; PubMed Central
PMCID: PMCPMC5032908.
82.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550. Epub 2014/12/18. doi:
10.1186/s13059-014-0550-8. PubMed PMID: 25516281; PubMed Central PMCID:
PMCPMC4302049.
83.   Gentleman RC, Carey VJ, Bates DM, Bolstad B, Dettling M, Dudoit S, et al.
Bioconductor: open software development for computational biology and bioinformatics.
Genome Biol. 2004;5(10):R80. Epub 2004/10/06. doi: 10.1186/gb-2004-5-10-r80. PubMed
PMID: 15461798; PubMed Central PMCID: PMCPMC545600.
84.   Wu R, Dephoure N, Haas W, Huttlin EL, Zhai B, Sowa ME, et al. Correct interpretation of
comprehensive phosphorylation dynamics requires normalization by protein expression
changes. Mol Cell Proteomics. 2011;10(8):M111.009654. Epub 2011/05/10. doi:
10.1074/mcp.M111.009654. PubMed PMID: 21551504; PubMed Central PMCID:
PMCPMC3149096.
85.   Tan H, Yang K, Li Y, Shaw TI, Wang Y, Blanco DB, et al. Integrative proteomics and
phosphoproteomics profiling reveals dynamic signaling networks and bioenergetics pathways
underlying T cell activation. Immunity. 2017;46(3):488-503. Epub 2017/03/14. doi:
10.1016/j.immuni.2017.02.010. PubMed PMID: 28285833; PubMed Central PMCID:
PMCPMC5466820.
86.   Wagih O, Sugiyama N, Ishihama Y, Beltrao P. Uncovering phosphorylation-based
specificities through functional interaction networks. Mol Cell Proteomics. 2016;15(1):236-45.
Epub 2015/11/18. doi: 10.1074/mcp.M115.052357. PubMed PMID: 26572964; PubMed Central
PMCID: PMCPMC4762521.
87.   Obenauer JC, Cantley LC, Yaffe MB. Scansite 2.0: Proteome-wide prediction of cell
signaling interactions using short sequence motifs. Nucleic Acids Res. 2003;31(13):3635-41.
Epub 2003/06/26. doi: 10.1093/nar/gkg584. PubMed PMID: 12824383; PubMed Central PMCID:
PMCPMC168990.
88.   Mootha VK, Lindgren CM, Eriksson KF, Subramanian A, Sihag S, Lehar J, et al. PGC-
1alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated
in human diabetes. Nat Genet. 2003;34(3):267-73. Epub 2003/06/17. doi: 10.1038/ng1180.
PubMed PMID: 12808457.
89.   Abbey DA, Funt J, Lurie-Weinberger MN, Thompson DA, Regev A, Myers CL, et al.
YMAP: a pipeline for visualization of copy number variation and loss of heterozygosity in
eukaryotic pathogens. Genome Med. 2014;6(11):100. Epub 2014/12/17. doi: 10.1186/s13073-
014-0100-8. PubMed PMID: 25505934; PubMed Central PMCID: PMCPMC4263066.
90.   Baum K, Schuchhardt J, Wolf J, Busse D. Of gene expression and cell division time: A
mathematical framework for advanced differential gene expression and data analysis. Cell Syst.
2019;9(6):569-79 e7. Epub 2019/09/16. doi: 10.1016/j.cels.2019.07.009. PubMed PMID:
31521604.
1227