

1 **Allicin From Garlic Disrupts the Cytoskeleton in Plants**

2 Ulrike Noll¹, Miriam Schreiber^{1,2}, Monika Hermanns¹, Christopher A. Mertes¹, Alan J.

3 Slusarenko¹ and Martin C.H. Gruhlke^{1*}

4

5

6 ¹ Department of Plant Physiology, RWTH Aachen University, Worringer Weg 1, 52056 Aachen,

7 Germany

8 ² Present address: College of Life Sciences, University of Dundee, Division of Plant Science, Errol

9 Road, Invergowrie, Dundee, UK

10 *corresponding author

11

12

13

1 **Abstract**

2 Allicin is a defence substance produced by garlic cells upon injury. It is a thiosulfinate showing
3 redox-activity and a broad range of antimicrobial and biocidal activity. It is known that allicin
4 efficiently oxidizes thiol-groups and it has been described as a redox toxin. In order to learn
5 more about the effect of allicin on plants we used pure synthesized allicin, and investigated
6 cytoplasmic streaming in sterile filaments of *Tradescantia fluminensis*, organelle movement
7 using transgenic *Arabidopsis* with organelle-specific GFP-tags, and effects on actin and tubulin
8 in the cytoskeleton using GFP-tagged lines. Auxin distribution in roots was investigated using
9 PIN1:GFP, PIN3:GFP, DR5:GFP and DII-VENUS *Arabidopsis* reporter lines.

10 Allicin inhibited cytoplasmic streaming in *T. fluminensis* and organelle movement of
11 peroxisomes and the Golgi apparatus in a concentration-dependent manner, inhibited root
12 growth and destroyed the correct root tip distribution of auxin.

13 We speculate that the cytoskeleton can be a primary “receptor” for allicin’s oxidizing properties
14 and as a consequence cytoskeleton-dependent cellular processes are disrupted.

15

16

17

18

19

20

21 **Key Words:** Allicin; Redox; *Tradescantia*; *Arabidopsis*; Cytoskeleton; Auxin

22

Allicin and root development

1 Introduction

2 Actin and tubulin are major structural components of the cytoskeleton and their polymerization
3 and depolymerization are important, amongst other things, for cytoplasmic streaming and
4 intracellular trafficking (Allen & Allen, 1978). Soon after the discovery of actin it was shown that
5 oxidizing agents not only inhibited the polymerization of globular actin, but also depolymerized
6 actin filaments (Feuer et al. 1948). The importance of sulfhydryl groups for protoplasmic
7 streaming in plant cells was reported in 1964 by Abe (Abe 1964) who showed the effect of the
8 specific thiol-trapping reagent *p*-chloromercuribenzoate in bringing cytoplasmic streaming to a
9 standstill in internodal cells of *Nitella flexilis*, leaf cells of *Elodea densa*, root hair cells of
10 *Hydrocharis morsus ranae*, and stamen hair cells of *Tradescantia reflexa*. Elements of the
11 cytoskeleton are known to be highly sensitive to oxidation (Farah and Amberg 2007; Farah et al.
12 2011) and redox-mediated posttranslational modifications of the cytoskeleton affect
13 polymerization (Johansson and Lundberg 2007; Wang et al. 2001). The redox regulation of actin
14 and tubulin dynamics is now a well characterized phenomenon (Wilson et al. 1916; Landino et
15 al. 2004).

16 Allicin from garlic is the most prominent sulfur-containing bioactive compound in freshly
17 damaged garlic tissue (Cavallito, 1944; Block 1992). It oxidizes thiol-groups in a thiol-disulfide-
18 exchange-like manner leading to *S*-thioallylated adducts (Gruhlke et al. 2011; Gruhlke et al.
19 2010; Miron et al. 2010; Pinto et al. 2006; Winkler et al. 1992). Since allicin is a strongly
20 antimicrobial compound with a broad-spectrum of activity, it has the potential for application in
21 agricultural pest control (Auger et al. 2004; Freeman and Koder 1995; Portz et al. 2008;

Allicin and root development

1 Slusarenko et al. 2011; Slusarenko et al. 2008). Allicin's degradation products, primarily
2 polysulfanes, ajoenes or vinyldithiins, can also show biological activity and can be active as
3 antimicrobial components. The effect of these sulfur-containing components in allelopathic
4 interactions with other organisms can be of importance, as is observed in particular for wild
5 garlic (*Allium ursinum* L.), attributable to the exudation of sulfur-containing secondary
6 metabolites (Block 2010).

7 In animal cells it was shown that allicin, which acts as a "redox-toxin" (Gruhlke et al. 2010),
8 affects the integrity and function of the cytoskeleton (Prager-Khoutorsky et al. 2007). It was
9 reported that 0.5 μM allicin preferentially affected the tubulin cytoskeleton, while the actin
10 cytoskeleton was a minor target (*ibid.*). However, at a higher 25 μM concentration of allicin the
11 actin cytoskeleton was also reported to be disrupted (Sela et al. 2004). Furthermore, 16
12 cytoskeletal proteins were shown to be S-thioallylated by 100 μM allicin in human Jurkat cells
13 within 10 minutes of exposure (Gruhlke et al. 2019).

14 The actin cytoskeleton is of particular importance for the correct polar transport of auxin in
15 relation to PIN efflux carrier recycling (Vieten et al. 2007; Nick et al. 2009). Auxin also exerts an
16 influence on the regulation of actin turnover (Rahman et al. 2007). The high redox sensitivity of
17 actin is probably significant for the observation that auxin transport is related to the state of
18 the total cellular glutathione pool, the most important thiol-based redox buffer of the cell
19 (Koprivova et al. 2010). Thus, treatment of *Arabidopsis* seedlings with buthionine-sulphoximine
20 (BSO), a specific inhibitor of glutathione biosynthesis, led to a clear concentration-dependent
21 inhibition of root growth, and a loss of the auxin efflux carriers PIN1, PIN2 and PIN7 and,

Allicin and root development

1 consequently, a destruction of the correct auxin distribution in the root tip (*ibid.*). This
2 observation correlates with the finding that mutants in glutathione biosynthesis form
3 shortened roots, such as is seen in the weak allele mutants *pad2* and *cad2* of the Arabidopsis
4 *GSH1* gene which have reduced glutathione levels (Bashandy et al. 2010). Mutants of the strong
5 *rm1* (root-meristemless) allele of *GSH1*, are almost unable to produce glutathione, are no
6 longer able to maintain the root apical meristem, and consequently cannot form a root (Cheng
7 et al. 1995; Vernoux et al. 2000).

8 Due to the need to understand the phytotoxicity of allicin with regard to a possible use in
9 organic farming as well as the allelopathic effects of these substances, we are interested in
10 understanding the molecular causes of the effects of allicin on plants. It has already been
11 reported that allicin inhibits *Arabidopsis* primary and lateral root growth in a concentration-
12 dependent manner (Borlinghaus et al. 2014, Leontiev et al. 2018). In the work reported here we
13 analyzed the effect of allicin, applied as a solution of chemically synthesized pure allicin, on the
14 cytoskeleton and, in consequence, cytoskeleton-dependent cellular processes like cytoplasmic
15 streaming, organelle movement and auxin distribution which might lead to the root growth-
16 inhibiting effects of allicin.

17

18 **Materials and Methods**

19 Plant material and cultivation

20 *Tradescantia fluminensis* VELL. was cultivated in the greenhouse. For induction of flowering, the
21 plants were grown in nutrient-poor soil.

Allicin and root development

1 Mutants and different GFP-tagged Arabidopsis lines used in this study are listed in Table 1. The
2 plants were grown on solid Hoagland-medium (1 mM CaNO₃; 5,1 mM KNO₃; 0,5 mM MgSO₄; 0,5
3 mM MgCl₂; 0,13 mM (NH₄)H₂PO₄; 30 μM NH₃NO₃; 31 μM NaOH; 22 μM EDTA-Iron-Sodium Salt;
4 9,7 μM BH₃O₃; 22 μM FeSO₄; 2 μM MnSO₄; 0,31 μM ZnCl₂; 0,21 μM CuSO₄; 0,14 μM Na₂MoO₄;
5 86 nM Co(NO₃)₂, containing 8 g/L Plant Agar (Duchefa, Haarlem, The Netherlands). After
6 stratification for two days at 4°C, the seedlings were grown under long day conditions (16 h
7 dark, 8 h light; 22°C) for two days before treatment.

8

mutant	Phenotype	Reference
ABD2-GFP	GFP-fusion protein with actin binding protein ABD2 allows visualization of actin cytoskeleton	Voigt et al., 2005
Tubulin-GFP	GFP-fusion with beta-tubulin allows visualization of tubulin cytoskeleton	Ueda et al., 1999
GFP-PTS	Fusion of GFP with peroxisomal targeting sequence allows visualization of peroxisomes	Mano et al., 2002
<i>pad2</i>	weak GSH1 allele, homozygote has ~30 % GSH content of the wild type	Glazebrook and Ausubel 1994
<i>axr1-12</i>	No auxin-perception	Timpte et al., 1995
<i>aux1-2</i>	Lack of auxin-importer AUX1-2.	Timpte et al., 1995
<i>DR5::GFP</i>	GFP expression dependent on the auxin-responsive DR5-promoter.	Friml et al., 2003
<i>DII-VENUS</i>	Degradation of the fluorescent protein depending on the presence of auxin	Brunoud et al., 2012

9

10

11 Synthesis of allicin and quantification of allicin

Allicin and root development

1 Chemical synthesis of allicin as described previously (Gruhlke et al. 2010). Purity of chemically
2 synthesized allicin was >95% (*ibid.*).

3

4 Measurement of protoplasmic streaming

5 As a measure of the influence of allicin on protoplasmic flow, the time was recorded after which
6 cytoplasmic flow ceased on exposure to allicin. For this purpose, individual sterile flower
7 filaments were plucked off with tweezers and microscopied as a water preparation. To prevent
8 the specimen from being crushed, the cover glass was slightly raised by small amounts of
9 plasticine at the corners. Treatment with allicin or garlic juice was carried out by using a filter
10 paper to draw off the water under the coverslip and replacing it with allicin solutions of the
11 appropriate concentration. In the same way, the allicin solution was exchanged for distilled
12 water for the regeneration experiments. Microscopy was carried out at 400x magnification
13 using transmitted light microscopy. Separate staining was not necessary.

14

15 Confocal Laser Scan microscopy

16 For GFP-visualization, a Confocal Laser Scan Microscope (TCS SP1, Leica GmbH, Wetzlar,
17 Germany) was used with an excitation wavelength of 488 nm and emission filter of 500 – 550
18 nm. For microscopy, a 63x water immersion objective was used and processed with ImageJ
19 (<http://rsbweb.nih.gov/ij>).

20 For analysis of the cytoskeleton, two-day-old seedlings were used; the seedlings were placed on
21 a slide in water or in an allicin solution of the indicated concentration. The analysis was carried

Allicin and root development

1 out after 30 minutes. For the PIN and auxin localization study, seedlings were placed on
2 Hoagland medium containing a concentration of 2 mM. The seedlings were incubated for 24 h
3 under short day conditions (22°C, 16 h dark, 8 h light) before microscopic analysis.

4

5 Growth assays

6 Arabidopsis seeds were surface sterilised with ethanol and spread with a toothpick on solid
7 Hoagland medium. Plates were stored at 4°C for two days for stratification before incubation at
8 22°C for a further two days under long-day conditions (16 h light/8 h dark). After this time,
9 seedlings were carefully transferred with forceps to Hoagland medium plates containing allicin
10 at a final concentration of 0.5 mM and further incubated under these conditions. Root length
11 was measured at day 0 and after three days.

12

13 Statistics

14 Statistical analyses were carried out using the programme SigmaStat. Data were tested for
15 statistical significance using a multivariate data analysis (ANOVA) with a significance level
16 $p < 0.05$.

1 Results & Discussion

2 Protoplasmic streaming in *Tradescantia fluminensis* VELL.


3 As a model system for the influence of allicin on the cytoskeleton, we chose the sterile hyaline
4 filaments from the flower of *T. fluminensis* (shown in Figure 1 A, B & C). The time that elapsed
5 until cytoplasmic streaming ceased after exposure to allicin solutions of various
6 concentrations was considered as a measure of the relative activity of allicin on the
7 cytoskeleton. Particles carried in the cytoplasmic stream observed to follow locomotion (Fig. 1
8 D, E). Soon after the hyaline filaments of *T. fluminensis* were exposed the flow slowed down
9 and stopped entirely after a certain time, which was dependent on the allicin concentration. As
10 the concentration of allicin increased, the time required for the protoplasmic flow to come to a
11 standstill decreased (Fig. 1D). While a concentration of 0.5 mM allicin led to a cessation of
12 protoplasmic flow after approx. 13 minutes, a treatment with 4 mM allicin led to a complete
13 cessation of protoplasmic flow after approx. 2 minutes (Fig. 1D).
14 A further observation for 5, 10 and 30 minutes after treatment showed that in the presence of
15 allicin no new onset of protoplasmic disturbance was observed.

16

17 **Figure 1: (A) (B) & (C) Sterile filaments in the flower of *Tradescantia fluminensis* are made up**
18 **of hyaline cells and are well suited to observe protoplasmic streaming. The scale bar in (C) =**
19 **0.1 mm (D) Allicin leads to a concentration- and time-dependent decrease in protoplasmic**
20 **flow. The time that elapses at a certain concentration of allicin until protoplasmic flow stops**
21 **was measured, showing a correlation between the allicin concentration and the time that**
22 **elapses until flow stops. (E) Movement of structures in the protoplasmic stream, in this case a**
23 **fork in cytoplasmic strands, can be used to measure the protoplasmic streaming.**

24

1



2

3 The effect of allicin on the protoplasmic streaming was partially reversible

4 In the following experiments it was investigated whether the effect of allicin was reversible by
5 washing out the allicin. Filaments of *Tradescantia* were treated with allicin of a final
6 concentration of 0.25 mM. After cytoplasmic flow stopped the allicin solution was washed by
7 capillary suction with filter paper drawing in pure water at the coverslip edge to exchange the
8 bathing solution. After about 12 minutes, protoplasmic flow began again in isolated cells,

Allicin and root development


1 although at a slower rate than before allicin treatment (not shown). Thus, it can be concluded
2 that the inhibition of protoplasmic flow by allicin can be reversible after a certain regeneration
3 phase; at a concentration higher than 0.25 mM allicin, this reversal did not occur even after a
4 long observation period (>30 minutes).

5

6 Influence of allicin on the movement of peroxisomes in leaves

7 Since the movement of organelles, such as peroxisomes, also takes place within the framework
8 of protoplasmic flow, an Arabidopsis line was used whose peroxisomes were tagged with GFP
9 (Mano *et al.*, 2002). This makes their movement traceable. Leaves of this Arabidopsis line were
10 infiltrated by vacuum infiltration with either water as a control treatment or an allicin solution
11 of the indicated concentration and the movement of the organelles was observed under the
12 confocal laser scanning microscope (Fig. 2 A, B and C). While in the control it can clearly be seen
13 by observing the curved trail of peroxisome movements between the arrows that the
14 peroxisomes moved substantially over the 10 min observation period (Fig. 2 A), in the sample
15 infiltrated with 4 mM allicin, movement of the peroxisomes was not seen (Fig. 2 B).

16




17

1 **Figure 2: Movement of GFP-labelled peroxisomes in *Arabidopsis thaliana* treated with either**
2 **(A) water as a control or (B) 4 mM alliin. The images show a stack of 20 images each, taken**
3 **30 seconds apart. (C) For reference, a single image of the control treatment is shown**
4 **superimposed on the light microscopy image of the leaf cells.**

5
6

7 Allicin inhibits primary and lateral root growth and root hair development.



8

9 **Figure 3: Allicin inhibits primary root growth and root hair development. Seeds were**
10 **germinated for two days on filter paper discs which were then lifted and transferred to fresh**
11 **medium containing alliin (A) or cumene hydroperoxide at the stated concentrations. (B) Wild**

Allicin and root development

1 **type Col-0 was compared to the GSH-deficient *pad2* mutant. The scale bars = 1 mm. N=20,**
2 **experiment was repeated three times with similar results.**

3

4 The morphology of Arabidopsis wild type Col-0 seedlings germinated for two days then exposed
5 to 0.5 mM allicin was very reminiscent of the morphology of the *rm1* mutant (Cheng et al.
6 1995). The roots of wild type Col-0 ceased growing in the presence of allicin, while the shoot
7 axis appeared to be largely unaffected and developed normally (Fig. 3 A). Compared to the
8 untreated control, the roots of plants exposed to allicin are generally much shorter and showed
9 no lateral roots or root hairs. The prooxidant cumene hydroperoxide also led to a similarly
10 shortened root (Fig. 3 B). Root growth in the *pad 2* mutant was also similarly inhibited by allicin
11 and cumene hydroperoxide under the conditions tested (3B).


12

13 Effect of allicin on actin- and tubulin cytoskeleton in *Arabidopsis thaliana*

14 To understand in more detail how allicin affects root development and whether this is
15 associated with the putative effect on the cytoskeleton based on the observation in
16 *Tradescantia*, *Arabidopsis* lines were used whose actin or tubulin filaments have been visualised
17 with by beta-tubulin:GFP fusion (Ueda et al., 1999) and GFP:actin-binding protein expression
18 (Voigt et al., 2005).

19 Confocal microscopy of the actin cytoskeleton clearly shows typical filamentous structures in
20 the root of the untreated control (Fig.4 4 A). After exposure to 2 mM allicin for 24 h, the
21 filamentous structures were destroyed; instead, GFP aggregate structures were visible at the

1 margins of the cells. Allicin thus destroys the integrity of the actin cytoskeleton in roots of
2 Arabidopsis.
3 In addition to the actin cytoskeleton, the tubulin cytoskeleton was also examined in an
4 analogous manner. Here, too, the typical filament pattern was seen in the untreated control;
5 treatment with allicin resulted in a dissolution of these filament structures and a diffuse fine-
6 particulate distribution of GFP fluorescence, in contrast to the aggregate formation observed
7 for actin.



8

1 **Figure 4: Effect of 2 mM allicin on the actin and tubulin cytoskeleton after exposure for 24**
2 **hours. While controls exposed to water only show a typical filamentous structure of both**
3 **actin and tubulin, this is completely dissolved after 24 h exposure to allicin.**Effect of allicin on
4 auxin distribution PIN localization


5 The cytoskeleton fulfils a wide variety of tasks in the cell and is involved, among other things, in
6 transport processes in the cell. In order to investigate the influence of cytoskeletal integrity -
7 here primarily concerning actin - on important cellular transport processes, we chose the
8 localisation of PIN proteins, whose localisation is crucial for regulated auxin transport in tissues.
9 Again, GFP-tagged versions of the proteins, in this case PIN1 and PIN3, were used to track their
10 localisation *in vivo*. In addition, the expression of the constructs is under the respective native
11 promoter, so that the localisation of the expression can also be followed at tissue level.

12 In the untreated control, PIN1 shows expression mainly in the region of the central cylinder (Fig.
13 5 A); at the cellular level, the strongest fluorescence was observed on the basipetal side of the
14 cell. Exposure to allicin of a concentration of 2 mM led to a complete loss of this clear
15 localisation pattern and GFP fluorescence is weakly distributed throughout the cell (Fig. 5 B).

16 This corresponds exactly to the observations made by Koprivova during treatment with BSO,
17 suggesting that allicin is also active here via the glutathione pool (Koprivova et al., 2010).

18 In contrast to PIN1, the localisation of PIN3 is also found in the area of the cortex, but it is more
19 prominent in the area of the root apical meristem. Again, fluorescence is strongest at the
20 basipetal end of the cell (Fig. 5 C).

1 As observed with PIN1, complete disruption of PIN localisation has also occurred with PIN3 by
2 treatment with 2 mM allicin. The GFP signal, which indicates the localisation of the PIN1
3 protein, is diffusely present at both the cellular and tissue levels (Fig. 5 D). These results exactly
4 mirror the effects of the GSH1 inhibitor buthione sulfoximine (BSO) on PIN distribution (Kopriva
5 *et al.*, 2010). BSO leads to a reduction in cellular GSH levels and this mirrors the result of allicin
6 titrating out and oxidizing the GSH pool (Gruhlke *et al.*, 2010).



7

8 **Figure 5: Influence of allicin on the localisation of PIN1 and PIN3 proteins and on auxin**
9 **distribution in the root tip. (A) While the localisation of PIN1 in the control is restricted to the**
10 **central cylinder, a diffuse localisation of PIN1 is seen in (B) allicin-exposed plants. (C) The**
11 **localisation of PIN3 in the control is seen in the central cylinder and in the meristematic zone**
12 **of the root tip, but under allicin exposure (D) the localisation of PIN 3 protein is also diffuse.**
13 **This is also reflected in the localisation of auxin. In control plants, where GFP expression is**
14 **under the auxin-responsive DR5 promoter, a typical distribution of auxin is seen in the root**
15 **tip, meristematic and quiescent zone (E), whereas this clear localisation is abolished under**

Allicin and root development

1 **allicin exposure (F). The detection of auxin distribution in the root by means of DII-venus**
2 **indicates in a complementary way that no auxin is present where the fluorescent protein is**
3 **observed, because auxin leads to the degradation of the protein. Accordingly, in the control**
4 **(G), fluorescence is seen in the marginal areas of the root, but not in the meristematic and**
5 **quiescent zone, whereas under allicin exposure, fluorescence is observed everywhere in the**
6 **cells (H). All treated samples were exposed to 2 mM allicin for 24 h.**

7
8
9

10

11 Effect of allicin on auxin distribution

12 Based on the observation that allicin destroys the localisation of PIN proteins, and because PIN
13 proteins are essential for the directional transport of auxin, the final aim was to investigate
14 whether aberrant localisation of auxin occurs after allicin treatment. Two independent methods
15 based on fluorescent proteins were used to check the localisation of allicin. The DR5::GFP
16 reporter system is based on the fact that auxin induces the activity of the DR5 promoter and
17 thus drives GFP expression. GFP fluorescence thus indexes the presence of auxin.

18 Untreated Arabidopsis roots show a clear localisation of auxin around the quiescent zone of the
19 root apical meristem, in the central cylinder and the epidermal layers of the root (Fig. 5E).

20 Treatment with allicin resulted in the localisation of GFP fluorescence being restricted to the
21 root apex; auxin-dependent expression of GFP is no longer found in the quiescent and division
22 zone of the root apex (Fig. 5F). This shows that allicin causes a clear redistribution of auxin
23 localisation at the root tip.

24 As an alternative method to detect the localisation of auxin at the tissue level, the DII-Venus
25 construct was used, in which a "fast maturing" form of yellow-fluorescent protein (YFP) is


Allicin and root development

1 expressed in frame with an IAA protein under the control of a constitutive promoter. Because
2 IAA proteins are degraded in the presence of auxin, a lack of fluorescence of the VENUS protein
3 indicates a high concentration of auxin, whereas a strong fluorescence of the protein indicates
4 a low concentration of auxin. The results obtained by means of DII Venus corresponded to the
5 statements we also perceived with DR5:GFP. While the control showed that auxin was present
6 in the root meristem area, the distribution was diffuse after allicin exposure (Fig. 5G).

7

8 Effect of allicin on mutants impaired in auxin homeostasis and signaling.

9 Because we observed an effect on auxin distribution and PIN localization after allicin treatment
10 (Fig. 5), we investigated the effects of allicin on the auxin transport mutant *aux1-2* and the
11 auxin resistant mutant *axr1-12* in comparison to wt Col-0. Whereas the PIN proteins are
12 functional in auxin efflux at the basal cell pole, the AUX1 protein transports auxin into the cell
13 from the apoplast at the apical pole.



1

2 **Figure 6: (A) Influence of alliin on *aux1-2* and *axr1-12* mutants. (n=20) the experiments were**
3 **repeated three times with similar results.**

4

5 The Col-0 wild type shows that its exposure to 0.5 mM alliin leads to a significantly shortened

6 root. The *aux1-2* mutant, already has a significantly shortened root in the control compared to

7 the Col-0 but in contrast, root length was not further affected by the presence of alliin, *i.e.* this

8 mutant showed a degree of alliin resistance (Fig. 6 A) and this suggests that the root

9 phenotype we observe with alliin may be dependent upon auxin perhaps over-accumulating to

10 supraoptimal levels and inhibiting root growth. The *axr1-12* mutant, on the other hand, shows

Alliin and root development

1 no significant change in root length phenotype compared to the wild type in the control, but
2 treatment with allicin causes the roots to be shortened, but to a much lesser extent than in the
3 Col-0 control. This suggests that the influence of allicin on auxin transport is partly responsible
4 for the observation that roots grow less well under allicin exposure (Fig. 6 B). Possibly, another
5 auxin receptor can partially complement the phenotype of *axr1-12*. To test this hypothesis, Col-
6 0 and *axr1-12* mutants were exposed to 20 μ M cytochalasin, which leads to disruption of the
7 actin cytoskeleton. As a control, it was examined whether the solvent DMSO at the
8 concentration used (1% v/v) would lead to an effect, which was not the case. While the Col-0
9 wild type showed significantly shortened roots after treatment with cytochalasin, the *axr1-12*
10 mutant does not, suggesting that it is disruption of the actin cytoskeleton alters auxin
11 localisation. Accordingly, a mutant that is not an auxin receptor mutant shows no phenotype in
12 terms of root length upon destruction of the actin cytoskeleton. These results suggest that the
13 effect allicin has on root development correlates, at least in part, with its influence on auxin
14 distribution in the root dependent upon allicin's effects on the cytoskeleton.

15

16

17

1 **Acknowledgement**

2 Financial support from RWTH Aachen University is acknowledged.

3 **References**

4 Abe S (1964) The effect of p-chloromercuribenzoate on rotational protoplasmic streaming in
5 plant

6 cells. *Protoplasma* 58: 483-492.

7 Allen NS, Allen RD (1978) Cytoplasmic Streaming in Green Plants. *Ann. Rev. Biophys. Bioeng.*

8 7: 497-526.

9 Auger J, Arnault I, Diwo-Allain S, Ravier MI, Molia F, Pettiti M (2004) Insecticidal and
10 fungicidal

11 potential of *Allium* substances as biofumigants. *Agroindustria* 3: 5-8.

12 Bashandy T, Guillemot J, Vernoux T, Caparros-Ruiz D, Ljung K, Meyer Y, Reichheld J-P

13 (2010) Interplay between the NADP-Linked Thioredoxin and Glutathione Systems in

14 *Arabidopsis* Auxin Signaling. *Plant Cell* 22: 376-391.

15 Block E (1992) The Organosulfur Chemistry of the Genus *Allium* - Implications for the Organic

16 Chemistry of Sulfur. *Angew. Chem. Int. Edit.* 31: 1135-1178.

17 Block E (2010) *Garlic and Other Alliums - The Lore and the Science*. Royal Society of

18 Chemistry,

19 Cambridge.

20 Borlinghaus J, Albrecht F, Gruhlke M, Nwachukwu I, Slusarenko A (2014) Allicin: Chemistry

21 and Biological Properties. *Molecules* 19: 12591-12618.

22 Brunoud G, Wells DM, Oliva M, Larrieu A, Mirabet V, Burrow AH, Beeckman T, Kepinski S,

Allicin and root development

- 1 Traas J, Bennett MJ, Vernoux T. (2012) A novel sensor to map auxin response and
2 distribution at high spatio-temporal resolution. *Nature* 482(7383): 103-6, DOI:
3 10.1038/nature10791. PMID: 22246322.
- 4 Cavallito CJ, Bailey JH (1944) Allicin, the Antibacterial Principle of *Allium sativum* L.
5 Isolation,
6 Physical Properties and Antibacterial Action. *J. Am. Chem. Soc.* 66: 1950-1951.
- 7 Cheng JC, Seeley KA, Sung ZR (1995) RML1 and RML2, *Arabidopsis* Genes Required for Cell
8 Proliferation at the Root Tip. *Plant Physiol.* 107: 365-376.
- 9 Farah ME, Amberg DC (2007) Conserved actin cysteine residues are oxidative stress sensors that
10 can regulate cell death in yeast. *Mol. Biol. Cell.* 18: 1359-1365.
- 11 Farah ME, Sirotkin V, Haarer B, Kakhniashvili D, Amberg DC (2011) Diverse protective roles
12 of
13 the actin cytoskeleton during oxidative stress. *Cytoskeleton* 68: 340-354.
- 14 Feuer G, Molnar F, Pettko E, Straub FB (1948) Studies on the composition and polymerization
15 of
16 actin. *Hung. Acta Physiol.* 1:150–163.
- 17 Freeman F, Kodera Y (1995) Garlic Chemistry: Stability of S(2-Propenyl) 2-Propene-1-
18 sulfinothioate (Allicin) in Blood, Solvents, and Simulated Physiological Fluids *J. Agric.*
19 *Food Chem.* 43: 2332-2338.
- 20 Friml J, Vieten A, Sauer M *et al.* Efflux-dependent auxin gradients establish the apical–basal
21 axis
22 of *Arabidopsis*. *Nature* 426: 147–153, DOI: 10.1038/
23 Glazebrook J, Ausubel FM (1994) Isolation of phytoalexin-deficient mutants of *Arabidopsis*
 Allicin and root development

1 thaliana and characterization of their interactions with bacterial pathogens. Proc. Natl.
2 Acad. Sci. USA, 91: 8955-8959.

3 Gruhlke MCH, Antelmann H, Bernhardt J, Kloubert V, Rink L, Slusarenko AJ (2019) The
4 human
5 allicin-proteome: S-thioallylation of proteins by the garlic defence substance allicin and
6 its biological effects. Free Radic. Biol. Med. 131: 144-153.

7 Gruhlke MCH, Nwachukwu I, Arbach M, Anwar A, Noll U, Slusarenko AJ (2011) Allicin from
8 Garlic, Effective in Controlling Several Plant Diseases, is a Reactive Sulfur Species
9 (RSS) that Pushes Cells into Apoptosis. In: Dehne (ed.) Modern Fungicides and
10 Antifungal Compounds VI. Deutsche Phytomedizinische Gesellschaft, Braunschweig.

11 Gruhlke MCH, Portz D, Stitz M, Anwar A, Schneider T, Jacob C, Schlaich NL, Slusarenko AJ
12 (2010) Allicin disrupts the cell's electrochemical potential and induces apoptosis in yeast.
13 Free Rad. Biol. Med. 49: 1916-1924.

14 Johansson M, Lundberg M (2007) Glutathionylation of beta-actin via a cysteinyl sulfenic acid
15 intermediary. BCM Biochemistry 8: 26.

16 Koprivova A, Mugford S, Kopriva S (2010) Arabidopsis root growth dependence on glutathione
17 is linked to auxin transport. Plant Cell Rep. 29: 1157-1167.

18 Landino L.M, Robinson SH, Skreslet TE, Cabral DM (2004) Redox modulation of tau and
19 microtubule-associated protein-2 by the glutathione/glutaredoxin reductase system.
20 Biochem. Biophys. Res. Commun. 323(1): 112-7, DOI: 10.1016/j.bbrc.2004.08.065.

21 Leontiev R, Hohaus N, Jacob C, Gruhlke MCH, Slusarenko AJ (2018) A Comparison of the
22 Antibacterial and Antifungal Activities of Thiosulfinate Analogues of Allicin. Sci. Rep.
23 **8**: 6763, [DOI: 10.1038/s41598-018-25154-9](https://doi.org/10.1038/s41598-018-25154-9).

Allicin and root development

- 1 Mano S, Nakamori C, Hayashi M, Kato A, Kondo M, Nishimura, M (2002) Distribution and
2 Characterization of Peroxisomes in Arabidopsis by Visualization with GFP: Dynamic
3 Morphology and Actin-Dependent Movement. *Plant Cell Physiol.* 43: 331-341.
- 4 Miron T, Listowsky I, Wilchek M (2010) Reaction mechanisms of allicin and allyl-mixed
5 disulfides with proteins and small thiol molecules. *Eur. J. Med. Chem.* 45: 1912-1918.
- 6 Nick P, Han M-J, An G (2009) Auxin Stimulates Its Own Transport by Shaping Actin Filaments.
7 *Plant Physiol.* 151: 155-167, DOI: 10.1104/pp.109.140111.
- 8 Pinto JT, Krasnikov BF, Cooper AJL (2006) Redox-sensitive proteins are potential targets of
9 garlic-derived mercaptocysteine derivatives. *J. Nutr.* 136: 835S--841S.
- 10 Portz D, Koch E, Slusarenko AJ (2008) Effects of garlic (*Allium sativum* L.) juice containing
11 allicin on *Phytophthora infestans* (Mont.deBary) and on downy mildew of cucumber
12 caused by *Pseudoperonospora cubensis* (Berk. & M.A. Curtis) Rostovzev. *Eur. J. Plant*
13 *Pathol.* 122: 197–206.
- 14 Prager-Khoutorsky M, Goncharov I, Rabinkov A, Mirelman D, Geiger B, Bershadsky AD
15 (2007)
16 Allicin inhibits cell polarization, migration and division via its direct effect on
17 microtubules. *Cytoskeleton* 64: 321-337.
- 18 Rahman A, Bannigan A, Sulaman W, Pechter P, Blancaflor EB, Baskin TI (2007) Auxin, actin
19 and growth of the *Arabidopsis thaliana* primary root. *Plant J.* 50: 514-528.
- 20 Sela U, Ganor S, Hecht I, Brill A, Miron T, Rabinkov A, Wilchek M, Mirelman D, Lider O,
21 Hershkoviz R (2004) Allicin inhibits SDF-1 α -induced T cell interactions with fibronectin
22 and endothelial cells by down-regulating cytoskeleton rearrangement, Pyk-2
23 phosphorylation and VLA-4 expression. *Immunology* 111: 391–399.

- 1 Slusarenko AJ, Anwar A, Portz D, Noll U (2011) Garlic - A Call to Arms. *Outlooks on Pest*
2 *Management* 22: 83-86.
- 3 Slusarenko AJ, Patel A, Portz D (2008) Control of plant diseases by natural products: Allicin
4 from
5 garlic as a case study. *Eur. J. Plant Pathol.* 121: 313-322.
- 6 Timpte C, Lincoln C, Pickett FB, Turner J, Estelle M. (1995) The AXR1 and AUX1 genes of
7 *Arabidopsis* function in separate auxin-response pathways. *Plant J.* 8: 561-69.
- 8 Ueda K, Matsuyama T, Hashimoto T (1999) Visualization of microtubules in living cells of
9 transgenic *Arabidopsis thaliana*. *Protoplasma.* 206: 201-206.
- 10 Vernoux T, Wilson RC, Seeley KA, Reichheld JP, Muroy S, Brown S, Maughan SC, Cobbett
11 CS,
12 Montagu MV, Inzé D, May MJ, Sung ZR (2000) The ROOT
13 MERISTEMLESS1/CADMIUM SENSITIVE2 gene defines a glutathione-dependent
14 pathway involved in initiation and maintenance of cell division during postembryonic
15 root development. *Plant Cell* 12: 97—110.
- 16 Vieten A, Sauer M, Brewer PB, Friml J (2007) Molecular and cellular aspects of auxin-transport-
17 mediated development. *Trends Plant Sci.* 12: 160-168.
- 18 Voigt B, Timmers ACJ, Amaj JS, Müller J, Baluska F, Menzel D (2005) GFP-FABD2 fusion
19 construct allows in vivo visualization of the dynamic actin cytoskeleton in all cells of
20 *Arabidopsis* seedlings. *Eur. J. Cell Biol.* 84: 595–608
- 21 Wang J, Boja ES, Tan W, Tekle E, Fales HM, English S, Mieyal JJ, Chock PB (2001) Reversible
22 glutathionylation regulates actin polymerization in A431 cells. *J. Biol. Chem.* 276:
23 47763-47766.

1 Wilson C, Terman JR, González□Billault C, Ahmed G (2016), Actin filaments—A target for
2 redox

3 regulation. *Cytoskeleton* 73: 577-595, DOI: 10.1002/cm.21315.

4 Winkler G, Iberl B, Knobloch K (1992) Reactivity of Allicin and its Transformation Products
5 with

6 Sulphydryl Groups, Disulfide Groups, and Human Blood. *Planta Med.* 58: 665.

7

- 1 Supplemental Material
- 2 Videos showing the inhibitory effect of allicin on the movement of GFP-tagged peroxisomes in
- 3 Arabidopsis leafs.

