Systematic exploration of *Escherichia coli* phage-host interactions with the BASEL phage collection

Enea Maffei[1,#], Aisylu Shaidullina[1,#], Marco Burkolter[1], Valentin Druelle[1], Luc Willi[1], Fabienne Estermann[1], Sarah Michaelis[2], Hubert Hilbi[2], David S. Thaler[1,3], Alexander Harms[1]*

[1]Biozentrum, University of Basel, Basel, Switzerland

[2]Institute of Medical Microbiology, University of Zürich, Zürich, Switzerland

[3]Program for the Human Environment, Rockefeller University, New York City, USA

[#]These authors contributed equally.

*Correspondence should be addressed to Dr. Alexander Harms (alexander.harms@unibas.ch).

Short title: The BASEL phage collection links bacteriophage genomes and phenotypes

Abstract

Bacteriophages, the viruses infecting bacteria, hold great potential for the treatment of multidrug-resistant bacterial infections and other applications due to their unparalleled diversity and recent breakthroughs in their genetic engineering. However, fundamental knowledge of molecular mechanisms underlying phage-host interactions is mostly confined to a few traditional model systems and did not keep pace with the recent massive expansion of the field. The true potential of molecular biology encoded by these viruses has therefore remained largely untapped, and phages for therapy or other applications are often still selected empirically. We therefore sought to promote a systematic exploration of phage-host interactions by composing a well-assorted library of 66 newly isolated phages infecting the model organism *Escherichia coli* that we share with the community as the BASEL collection (BActeriophage SElection for your Laboratory). This collection is largely representative of natural *E. coli* phage diversity and was intensively characterized phenotypically and genomically alongside ten well-studied traditional model phages. We experimentally determined essential host receptors of all phages, quantified their sensitivity to eleven defense systems across different layers of bacterial immunity, and matched these results to the phages' host range across a panel of pathogenic enterobacterial strains. Our results reveal clear patterns in the distribution of phage phenotypes and genomic features that highlight systematic differences in the potency of different immunity systems and point towards the molecular basis of receptor specificity in several phage groups. Strong trade-offs were detected between fitness traits like broad host recognition and resistance to bacterial immunity that might drive the divergent adaptation of different phage groups to specific niches. We envision that the BASEL collection will inspire future work exploring the biology of bacteriophages and their hosts by facilitating the discovery of underlying molecular mechanisms as the basis for an effective translation into biotechnology or therapeutic applications.

Introduction

Bacteriophages, the viruses infecting bacteria, are the most abundant biological entities on earth with key positions in all ecosystems and carry large part of our planet's genetic diversity in their genomes [1-3]. Out of this diversity, a few phages infecting *Escherichia coli* became classical models of molecular biology with roles in many fundamental discoveries and are still major workhorses of research today [4]. The most prominent of these are the seven "T phages" T1 – T7 [5, reviewed in reference 6] and bacteriophage lambda [7]. Like the majority of known phages, these classical models are tailed phages or *Caudovirales* that use characteristic tail structures to bind host surface receptors and to inject their genomes from the virion head into the host cell. Three major virion morphotypes of *Caudovirales* are known, myoviruses with a contractile tail, siphoviruses with a long and flexible tail, and podoviruses with a very short, stubby tail [2] (Fig 1A). While the T phages are all so-called lytic phages and kill their host to replicate at each infection event, lambda is a temperate phage and can either kill the host to directly replicate or decide to integrate into the host's genome as a prophage for transient passive replication by vertical transmission in the so-called lysogen [2, 8]. These two alternative lifestyles as lytic or temperate phages have major implications for viral ecology and evolution: While lytic phages have primarily been selected to overcome host defenses and maximize virus replication, temperate phages characteristically encode genes that increase the lysogens' fitness, e.g., by providing additional bacterial immunity systems to fight other phages [8, 9].

The ubiquity of phage predation has driven the evolution of a vast arsenal of bacterial immunity systems targeting any step of phage infection [9, 10]. Inside host cells, phages encounter two lines of defense of which the first primarily comprises restriction-modification (RM) systems or CRISPR-Cas, the bacterial adaptive immunity, that directly attack viral genomes [11] (Fig 1B). A second line of defense is formed by diverse abortive infection (Abi) systems that protect the host population by triggering an altruistic suicide of infected cells when sensing viral infections [9-12] (Fig 1B). While RM systems and CRISPR-Cas are highly abundant and have been successfully adapted for biotechnology (honored with

61 Nobel Prizes in 1978 and 2020, respectively), the molecular mechanisms underlying the function of

62 collectively abundant, but each individually rare, Abi systems have remained elusive with few exceptions

63 [9-12].

Fig 1. The three morphotypes of *Caudovirales* and two lines of defense in bacterial immunity.
(A) The virions of tailed phages or *Caudovirales* can be assigned to three general morphotypes including myoviruses (contractile tail), siphoviruses (long and flexible, non-contractile tail), and podoviruses (short and stubby tail). **(B)** The life cycle of a typical lytic phage begins with reversible attachment to a so-called primary receptor on the bacterial cell surface (1), usually via lateral tail fibers at the virion. Subsequently, irreversible attachment to secondary or terminal receptors usually depends on structures at the end of the tail, e.g., short tail fibers for many myoviruses and central tail fibers or tail tip proteins for siphoviruses (2 ; see also (A)). After genome injection (3), the phage takes over the host cell, replicates, and releases the offspring by host cell lysis (4). Inside the host cell, the bacteriophage faces two lines of host defenses, first bacterial immunity systems that try to clear the infection by directly targeting the phage genome (5) and then abortive infection systems that kill the infected cell when a viral infection is sensed (6).

Research on phages has expanded at breathless pace over the last decade with a focus on

biotechnology and on clinical applications against bacterial infections ("phage therapy") [13, 14]. Besides

or instead of the few traditional model phages, many researchers now employ comparably poorly described,

newly isolated phages that are often only used for a few studies and available only in their laboratory. The

consequence of this development is a rapidly growing amount of very patchy data such as, e.g., the currently

more than 14'000 available unique phage genomes [15] for which largely no linked phenotypic data are

available. Despite the value of proof-of-principle studies and a rich genome database, this lack of

systematic, interlinked data in combination with the diversity of bacteriophages makes it very difficult to

gain a mechanistic understanding of phage biology or to uncover patterns in the data that would support the

discovery of broad biological principles beyond individual models.

As an example, phage isolates for treating a specific case of bacterial infection are necessarily

chosen largely empirically due to the lack of systematic data about relevant phage properties. Currently,

the selection of native phages and their engineering for therapeutic applications primarily focus on a lytic

lifestyle, a broad host range, and very occasionally on biofilm- or cell wall-degrading enzymes that are

comparably well understood genetically and mechanistically [13, 14, 16, 17]. However, the molecular

mechanisms and genetic basis underlying other desired features such as resistance to different bacterial

immunity systems and, in general, the distribution of all these features across different groups of phages

have remained understudied. Given the notable incidence of treatment failure in phage therapy [18-20], a

better understanding of the links between phage taxonomy, genome sequence, and phenotypic properties

seems timely to select more effective native phages for therapeutic applications and to expand the potential

of phage engineering.

In this work we therefore present the BASEL (Bacteriophage Selection for your Laboratory)

collection as a reference set of 66 newly isolated lytic bacteriophages that infect the laboratory strain *E. coli*

K-12 and make it accessible to the scientific community. We provide a systematic phenotypic and genomic

characterization of these phages alongside ten classical model phages regarding host receptors, sensitivity

and resistance to bacterial immunity, and host range across diverse enterobacteria. Our results highlight

clear phenotypic patterns between and within taxonomic groups of phages that reveal strong trade-offs

between important bacteriophage traits. These findings greatly expand our understanding of bacteriophage

ecology, evolution, and their interplay with bacterial immunity systems. We therefore anticipate that our

work will not only establish the BASEL collection as a reference point for future studies exploring

fundamental bacteriophage biology but also promote a rational application of phage therapy based on an

improved selection and engineering of bacteriophages.

Results

Composition of the BASEL collection

The first aim of our study was to generate a collection of new phage isolates infecting the

ubiquitously used *E. coli* K-12 laboratory strain that would provide representative insight into the diversity

of tailed, lytic phages by covering all major groups and containing a suitable selection of minor ones.

Similar collections exist, e.g., for *E. coli* in form of the ECOR collection of 72 *E. coli* strains that are

commonly used to study how this model organism varies in certain traits or can deal with / evolves under

certain conditions [21, 22]. Though nothing truly comparable is available for phages, it is notable that the

seven T phages had originally been chosen in the early 1940s with the explicit aim of providing a reference

point that would enable comparative, systematic research on bacteriophages [reviewed in reference 6].

However, while these and the other classical model phages such as lambda have been invaluable to uncover

many fundamental principles of molecular biology, their number and taxonomic range are too limited to

serve as a representative reference even for bacteriophages infecting *E. coli*.

As a first step, we generated a derivative of *E. coli* K-12 without any of the native barriers that

might limit or bias phage isolation unfavorably. This strain, *E. coli* K-12 MG1655 ΔRM, therefore lacks

the O-antigen glycan barrier (see below), all restriction systems, as well as the RexAB and PifA Abi systems

(see *Materials and Methods* as well as S1 Text). *E. coli* K-12 ΔRM was subsequently used to isolate

hundreds of phages from environmental samples such as river water or compost, but mostly from the inflow

of different sewage treatment facilities (Fig 2A and *Materials and Methods*).

Fig 2. Overview of the BASEL collection.
(A) Illustration of the workflow of bacteriophage isolation, characterization, and selection that resulted in the BASEL collection (details in *Materials and Methods*). **(B)** Taxonomic overview of the bacteriophages included in the BASEL collection and their unique Bas## identifiers. Newly isolated phages are colored according to their family while well-studied reference phages are shown in grey.

Previous bacteriophage isolation studies had already provided deep insight into the diversity of

tailed, lytic *E. coli* phages in samples ranging from sewage over diverse natural environments to infant guts

or blood and urine of patients in tertiary care [23-29]. Despite the wide diversity of known *E. coli* phages

[30], nearly all phage isolates reported in these studies belonged to five major groups and were either

myoviruses of 1) *Tevenvirinae* or 2) *Vequintavirinae* subfamilies and close relatives, 3) large siphoviruses

of the *Markadamsvirinae* subfamily within *Demerecviridae*, or small siphoviruses of 4) diverse

Drexlerviridae subfamilies or 5) the genera *Dhillonvirus*, *Nonagvirus*, and *Seuratvirus* of the *Siphoviridae*

family. Podoviruses of any kind were rarely reported, and if, then were mostly *Autographiviridae* isolated

using enrichment cultures that are known to greatly favor such fast-growing phages [24, 31]. This pattern

does not seem to be strongly biased by any given strain of *E.coli* as isolation host because a large, very

thorough study using diverse *E. coli* strains reported essentially the same composition of taxonomic groups

[23].

Whole-genome sequencing of around 120 different isolates from our isolation experiments largely

reproduced this pattern, which suggests that several intrinsic limitations of our approach did not strongly

affect the spectrum of phages that we sampled (S3 Text). After eliminating closely related isolates, the

BASEL collection was formed as a set of 66 new phage isolates (Fig 2; see also *Materials and Methods*

and S5 Table). We deliberately did not give full proportional weight to highly abundant groups like

Tevenvirinae so that the BASEL collection is not truly representative in a narrow quantitative sense. Instead,

we included as many representatives of rarely isolated groups (e.g., podoviruses) as possible to increase the

biological diversity of phages in our collection that could be an important asset for studying the genetics of

phage / host interactions or for unraveling genotype / phenotype relationships. Besides these 66 new isolates

(serially numbered as Bas01, Bas02, etc.; Fig 2 and S5 Table), we included a panel of ten classical model

phages in our genomic and phenotypic characterization and view them as an accessory part of the BASEL

collection. Beyond the T phages (without T1 that is a notorious laboratory contaminant [32]), we included

well-studied podovirus N4 and obligately lytic mutants of the three most commonly studied temperate

phages lambda, P1, and P2 [5, 7, 33, 34] (Fig 2; see also S5 Table).

Overview: Identification of phage surface receptors

The infection cycle of most tailed phages begins with host recognition by reversible adsorption to

a first "primary" receptor on host cells (often a sugar motif on surface glycans) followed by irreversible

binding to a terminal or "secondary" receptor directly on the cell surface which results in DNA injection

(Fig 1B) [35]. Importantly, the adsorption to many potential hosts is blocked by long O-antigen chains on

the LPS and other exopolysaccharides that effectively shield the cell surface unless they can be degraded

or specifically serve as the phage's primary receptor [35-38]. Tailed phages bind to primary and secondary

receptors using separate structures at the phage tail which display the dedicated receptor-binding

proteins (RBPs) in the form of tail fibers, tail spikes (smaller than fibers, often with enzymatic domains),

or central tail tips [comprehensively reviewed in reference 35]. Most commonly, surface glycans such as

the highly variable O-antigen chains or the enterobacterial common antigen (ECA) are bound as primary

receptors. Secondary receptors on Gram-negative hosts are near-exclusively porin-family outer

membrane proteins for siphoviruses and core LPS sugar structures for podoviruses, while myoviruses

were found to use either one of the two depending on phage subfamily or genus [35, 39, 40]. Notably, no

host receptor is known for the vast majority of phages that have been studied, but understanding the

genetic basis and molecular mechanisms underlying host recognition as the major determinant of phage

host range is a crucial prerequisite for host range engineering or a rational application of phage therapy

[41]. Since tail fibers and other host recognition modules are easily identified in phage genomes, we used

phage receptor specificity as a model to demonstrate the usefulness of systematic phenotypic data with

the BASEL collection as a key to unlock information hidden in the genome databases.

We therefore first experimentally determined the essential host receptor(s) of all phages in the

BASEL collection before analyzing the phages' genomes for the mechanisms underlying receptor

specificity. Briefly, the dependence on surface proteins was assessed by plating each phage on a set of more

than fifty single-gene mutants (S1 Table) or whole-genome sequencing of spontaneously resistant bacterial

clones (see *Materials and Methods*). The role of host surface glycans was quantified with *waaG* and *waaC*

mutants that display different truncations of the LPS core, a *wbbL(+)* strain with restored O16-type O-

antigen expression, and a *wecB* mutant that is specifically deficient in production of the ECA [42-44] (Fig

3A, S1 Table, and *Materials and Methods*).

Fig 3. Overview of *E. coli* surface glycan variants and the immunity systems used in this study.
(A) The surface glycans of different *E. coli* K-12 MG1655 variants are shown schematically (details in running text and *Materials and Methods*). Note that the *E. coli* K-12 MG1655 laboratory wildtype does not merely display the K12-type core LPS (classical rough LPS phenotype) but also the most proximal D-glucose of the O16-type O-antigen. **(B)** Key features of the six RM systems (each two of type I, type II, and type III) and the five Abi systems used for the phenotyping of this study are summarized schematically. Recognition sites of RM systems have either been determined experimentally or were predicted in REBASE (red nucleotides: methylation sites; dotted lines: cleavage sites) [47-49, 128]. The Abi systems have been characterized to very different extent but constitute the most well-understood representatives of these immunity systems of *E. coli* [10, 50].

Overview: Phenotyping of sensitivity / resistance to bacterial immunity systems

Some bacterial immunity systems are very common among different strains of a species (like

certain types of RM systems for *E. coli*), while others – especially the various Abi systems – have each a

very patchy distribution but are no less abundant if viewed together [11]. Each bacterial strain therefore

encodes a unique repertoire of a few very common and a larger number of rarer, strain-specific immunity

systems, but it is unknown how far these systems impact the isolation of phages or their efficacy in

therapeutic applications. A systematic view of the potency and target range of different immunity systems

and how the diverse groups of phages differ in sensitivity / resistance to these systems might enable us to

select or engineer phages with a higher and more reliable potency for phage therapy or biotechnology. As

an example, previous work showed that *Tevenvirinae* or *Seuratvirus* and *Nonagvirus* phages exhibit broad

resistance to RM systems due to the hypermodification of cytosines or guanosines in their genomes,

respectively, which could be an interesting target for phage engineering [45, 46]. However, it is unknown

whether this mechanism of RM resistance (or any other viral anti-immunity function) actually results in a

measurably broader phage host range.

We therefore systematically quantified the sensitivity of all phages of the BASEL collection against

a panel of eleven immunity systems and scored their infectivity on a range of pathogenic enterobacteria that

are commonly used as model systems (see *Materials and Methods* and Fig 3B). Shortly, we tested six RM

systems by including each two type I, type II, and type III systems that differ in the molecular mechanisms

of DNA modification and cleavage [10, 47-49] (Fig 3B). Besides these, we included the most well-studied

Abi systems of *E. coli*, RexAB of the lambda prophage, PifA of the F-plasmid, as well as the Old, Tin, and

Fun/Z systems of the P2 prophage (Fig 3B). Previous work suggested that RexAB and PifA sense certain

proteins of phages T4 and T7, respectively, to trigger host cell death by membrane depolarization [10].

Conversely, Old possibly senses the inhibition of RecBCD / ExoV during lambda infections and might kill

by tRNA degradation, Fun/Z abolishes infections by phage T5 via an unknown mechanism, and Tin poisons

the replicative ssDNA binding protein of *Tevenvirinae* [50]. Beyond *E. coli* K-12, we quantified the

infectivity of the BASEL collection on uropathogenic *E. coli* (UPEC) strains UTI89 and CFT073,

enteroaggregative *E. coli* (EAEC) strain 55989, alternative laboratory strain *E. coli* B REL606, and

Salmonella enterica subsp. *enterica* serovar Typhimurium strains 12023s and SL1344 (see *Materials and*

Methods and S1 Table).

Properties of the *Drexlerviridae* family

Phages of the *Drexlerviridae* family (previously also known as T1 superfamily [30]) are small

siphoviruses with genome sizes of ca. 43-52 kb (Figs 4A-D). The BASEL collection contains thirteen new

Drexlerviridae phages that are broadly spread out across the various subfamilies and genera of this family

(Fig 4B). Though it has apparently never been directly demonstrated, it seems likely that the *Drexlerviridae*

use their lateral tail fibers in the same way as their larger cousins of the *Markadamsvirinae* (see below) to

contact specific O-antigen glycans as primary receptors without depending on this interaction for host

recognition [51] (Fig 4A). Consistently, the locus encoding the lateral tail fibers is very diverse among

Drexlerviridae and the proteins forming these tail fibers are only highly related at the far N-terminus of the

distal subunits where these are likely attached to the tail (S1B Fig). Among the *Drexlerviridae* that we

isolated, only JakobBernoulli (Bas07) shows robust plaque formation on *E. coli* K-12 MG1655 with

restored O16-type O-antigen expression, suggesting that its lateral tail fibers can use this O-antigen as

primary receptor (Figs 4A and 4D).

Just like the much larger *Markadamsvirinae* siphoviruses (see below), *Drexlerviridae* phages use

a small set of outer membrane porins as their secondary / final receptors for irreversible adsorption and

DNA injection. To the best of our knowledge, previous work had only identified the receptors of T1 and

IME18 (FhuA), IME347 (YncD), IME253 (FepA), and of LL5 as well as TLS (TolC) while for an additional

phage, RTP, no protein receptor could be identified [40, 52-54]. Using available single-gene mutants, we

readily determined the terminal receptor of eleven out of our thirteen *Drexlerviridae* phages as FhuA, BtuB,

YncD, and TolC (Figs 4B and 4C) but failed for two others, AugustePiccard (Bas01) and JeanPiccard

(Bas02). However, whole-genome sequencing of spontaneously resistant *E. coli* mutants showed that

resistance was linked to mutations in the gene coding for LptD, the LPS export channel [36], strongly

suggesting that this protein was the terminal receptor of these phages (Fig 5 and *Materials and Methods*).

Fig 4. Overview of *Drexlerviridae* phages.
(A) Schematic illustration of host recognition by *Drexlerviridae*. **(B)** Maximum-Likelihood phylogeny of *Drexlerviridae* based on several core genes with bootstrap support of branches shown if > 70/100. Newly isolated phages of the BASEL collection are highlighted by red phage icons and the determined or proposed terminal receptor specificity is highlighted at the phage names using the color code highlighted in (C). The phylogeny was rooted based on a representative phylogeny including *Dhillonvirus* sequences as outgroup (S1A Fig). **(C)** On the left, the seven identified receptors of small siphoviruses are shown with a color code that is also used to annotate demonstrated or predicted receptor specificity in the phylogenies of Fig 4B and Figs 6A + 6C. On the right, we show representative *bona fide* RBP loci that seem to encode the receptor specificity of these small siphoviruses (with the same color code). Note that the loci linked to each receptor are very similar while the genetic arrangement differs considerably between loci linked to different terminal host receptors (see also S1C Fig). **(D)** The results of quantitative phenotyping experiments with *Drexlerviridae* phages regarding sensitivity to altered surface glycans and bacterial immunity systems are presented as efficiency of plating (EOP). Data points and error bars represent average and standard deviation of at least three independent experiments.

Similar to the central tail fibers of *Markadamsvirinae*, the RBPs of *Drexlerviridae* are thought to

be displayed at the distal end of a tail tip protein related to well-studied GpJ of bacteriophage lambda [51,

54, 55]. The details of this host recognition module had remained elusive, though it was suggested that (like

for T5 and unlike for lambda) dedicated RBPs are non-covalently attached to the J-like protein and might

be encoded directly downstream of the *gpJ* homologs together with cognate superinfection exclusion

proteins [54, 55]. By comparing the genomes of all *Drexlerviridae* phages with experimentally determined

surface protein receptors, we were able to match specific allelic variants of this *bona fide* RBP loci to each

known surface receptor of this phage family (Fig 4C). Notably, for the three known phages targeting TolC

including DanielBernoulli (Bas08), the RBP locus is absent and apparently functionally replaced by a C-

terminal extension of the GpJ-like tail tip protein that probably directly mediates receptor specificity like

GpJ of bacteriophage lambda (Fig 4C) [56, 57]. Interestingly, similar RBP loci with homologous alleles are

also found at the same genomic locus in small *Siphoviridae* of *Dhillonvirus*, *Nonagvirus*, and *Seuratvirus*

genera (see below) where they also match known receptor specificity without exception (Figs 4-6 and S1C).

Pending experimental validation, we therefore conclude that these distantly related groups of small

siphoviruses share a common, limited repertoire of RBPs that enables receptor specificity of most

representatives to be predicted *in silico*. As an example, it seems clear that phage RTP targets LptD as

terminal receptor (Figs 4B and S1D). The poor correlation of predicted receptor specificity with the

phylogenetic relationships of small siphoviruses (Figs 4B, 6B, and 6D) is indicative of frequent horizontal

transfer of these RBP loci. This highlights the modular nature of this host recognition system which might

enable the targeted engineering of receptor specificity to generate "designer phages" for different

applications as previously shown for siphoviruses infecting *Listeria* [14, 58]. LptD would be a particularly

attractive target for such engineering because it is strictly essential under all conditions, highly conserved,

and heavily constrained due to multiple interactions that are critical for its functionality [36].

Fig 5. LptD is a commonly targeted terminal receptor of small siphoviruses.
(A) Whole-genome sequencing of bacterial mutants exhibiting spontaneous resistance to seven small siphoviruses with no previously known receptor revealed different mutations or small deletions in the essential gene *lptD* that encodes the LptD LPS export channel. Top agar assays with two representative mutants in comparison to the ancestral *E. coli* K-12 BW25113 strain were performed with serial tenfold dilutions of twelve different phages (undiluted high-titer stocks at the bottom and increasingly diluted samples towards the top). Both mutants display complete resistance to the seven small siphoviruses of diverse genera within *Drexlerviridae* and *Siphoviridae* families that share the same *bona fide* RBP modules (S1C Fig) while no other phage of the BASEL collection was affected. In particular, we excluded indirect effects, e.g., via changes in the LPS composition in the *lptD* mutants, by confirming that five LPS-targeting phages of diverse families (see below) showed full infectivity on all strains. **(B)** The amino acid sequence alignment of wildtype LptD with the two mutants highlighted in (A) shows that resistance to LptD-targeting phages is linked to small deletions in or adjacent to regions encoding extracellular loops as defined in previous work [147], suggesting that they abolish the RBP-receptor interaction.

Once inside the host cell, our results show that *Drexlerviridae* phages are highly diverse in their

sensitivity or resistance to diverse bacterial immunity systems (Fig 4D). While few representatives like

FritzSarasin (Bas04) or PeterMerian (Bas05) are highly resistant, most *Drexlerviridae* are very sensitive to

any kind of RM systems sometimes including the type I machineries that are largely unable to target any

other phage that we tested (Fig 4D). Previous work suggested that *Drexlerviridae* might employ DNA

methyltransferases as a defense strategy against host restriction [24], and these phages indeed encode

variable sets of N6-adenine and C5-cytosine methyltransferases. However, we found no obvious pattern

that would link DNA methyltransferases or any other specific genomic features to restriction resistance /

sensitivity. Instead, sensitivity and resistance to bacterial immunity strongly correlate with *Drexlerviridae*

phylogeny: While *Hanrivervirus* and *Warwickvirus* genera of *Tempevirinae* are highly resistant and

Tunavirinae show comparably intermediate sensitivity, the other phages (in particular *Braunvirinae*, but

also *Tlsvirus* phage DanielBernoulli (Bas08)) are highly sensitive (Figs 4B and 4D).

Properties of *Siphoviridae* genera *Dhillonvirus*, *Nonagvirus*, and *Seuratvirus*

Phages of the *Dhillonvirus*, *Seuratvirus*, and *Nonagvirus* genera within the *Siphoviridae* family are

small siphoviruses that are superficially similar to *Drexlerviridae* and have genomes with characteristic

size ranges of 43-46 kb (*Dhillonvirus*), 56-61 kb (*Seuratvirus*), and 56-64 kb (*Nonagvirus*; see Fig 6A-E).

Our twelve isolates included in the BASEL collection are spread out broadly across the phylogenetic ranges

of these genera (Figs 6B and 6D). Similar to *Drexlerviridae* (and *Markadamsvirinae*, see below), we

suggest that they also recognize glycan motifs at the O-antigen as their primary receptor on different host

strains (Fig 6A). This notion is strongly supported by the remarkable variation exhibited by the different

genomes at the lateral tail fiber locus (S2A Fig), as observed previously [23, 24]. None of these phages can

infect *E. coli* K-12 MG1655 with restored O16-type O-antigen expression, but (like for *Drexlerviridae*)

some require an intact LPS core for infectivity (Figs 6C and 6F). Experimental identification of the terminal

receptor of all small *Siphoviridae* confirmed that receptor specificity seems to be encoded by the same

system of *bona fide* RBP loci downstream of *gpJ* as for the *Drexlerviridae* (Figs 4B, 6B, and 6D). Many of

these phages target LptD or FhuA as terminal receptors while others, unlike any *Drexlerviridae*, bind to

LamB (Figs 6C and 6F). Notably, three related *Nonagvirus* phages encode a distinct *bona fide* RBP module

that could not be matched to any known terminal receptor (S2B Fig), suggesting that these phages might

target a different protein.

Fig 6. Overview of *Siphoviridae* genera *Dhillonvirus*, *Nonagvirus*, and *Seuratvirus*.

(A) Schematic illustration of host recognition by small siphoviruses. **(B)** Maximum-Likelihood phylogeny of the *Dhillonvirus* genus based on a whole-genome alignment with bootstrap support of branches shown if > 70/100. Newly isolated phages of the BASEL collection are highlighted by red phage icons and the determined or proposed terminal receptor specificity is highlighted at the phage names using the color code highlighted in Fig 4C. The phylogeny was rooted between phage WFI and all others based on a representative phylogeny including *Drexlerviridae* sequences as outgroup (S1A Fig). **(C)** The results of quantitative phenotyping experiments with *Dhillonvirus* phages regarding sensitivity to altered surface glycans and bacterial immunity systems are presented as efficiency of plating (EOP). **(D)** Maximum-Likelihood phylogeny of the *Nonagvirus* and *Seuratvirus* genera based on a whole-genome alignment with bootstrap support of branches shown if > 70/100. Newly isolated phages of the BASEL collection are highlighted by red phage icons and the determined or proposed terminal receptor specificity is highlighted at the phage names using the color code highlighted in Fig 4C. The phylogeny was rooted between the two genera. **(E)** *Nonagvirus* and *Seuratvirus* phages share a core 7-deazaguanosine biosynthesis pathway involving FolE, QueD, QueE, and QueC which synthesizes dPreQ$_0$ that is inserted into their genomes by DpdA. In *Nonagvirus* phages, the fusion of QueC with a glutamate amidotransferase (Gat) domain to Gat-QueC results in the modification with dG$^+$ instead of dPreQ$_0$ [45]. **(F)** The results of quantitative phenotyping experiments with *Nonagvirus* and *Seuratvirus* phages regarding sensitivity to altered surface glycans and bacterial immunity systems are presented as efficiency of plating (EOP). In (C) and (F), data points and error bars represent average and standard deviation of at least three independent experiments.

In difference to *Dhillonvirus* phages that are relatives of *Drexlerviridae* [23] (see also S1A Fig),

Nonagvirus and *Seuratvirus* phages are closely related genera with genomes that are 10-15 kb larger than

those of the other small siphoviruses in the BASEL collection (S5 table) [59]. This difference in genome

size is largely due to genes encoding their signature feature, the 7-deazaguanine modification of

2′-deoxyguanosine (dG) in their genomes into 2′-deoxy-7-cyano-7-deazaguanosine (dpreQ$_0$) for

Seuratvirus phages and into 2′-deoxyarchaosine (dG$^+$) for *Nonagvirus* phages [45] (Fig 6E). These

modifications were shown to provide considerable (dG$^+$) or at least moderate (dpreQ$_0$) protection against

restriction of genomic DNA of *Seuratvirus* phage CAjan and *Nonagvirus* phage 9g *in vitro*, although

chemical analyses showed that only around one third (CAjan) or one fourth (9g) of the genomic dG content

is modified [45] (Fig 6E). Consistently, we found that both *Nonagvirus* and *Seuratvirus* phages are

remarkably resistant to type I and type II RM systems in our infection experiments, particularly if compared

to other small siphoviruses of *Dhillonvirus* or *Drexlerviridae* groups (Figs 4D, 6C, and 6F). Among

Nonagvirus and *Seuratvirus* phages, only ChristophMerian (Bas19, EcoKI) and VogelGryff (Bas25, EcoRI)

show some sensitivity to these RM systems. However, all phages of these genera are highly sensitive to

type III RM systems (Fig 6F). This observation does not necessarily indicate a lower sensitivity of type III

RM systems to guanosine modifications *per se* but is likely simply a consequence of the much larger

number of type III RM recognition sites in their genomes (S5 Table, see also Fig 3B).

Properties of *Demerecviridae: Markadamsvirinae*

Phages of the *Markadamsvirinae* subfamily of the *Demerecviridae* family (previously known as

T5-like phages [30]) are large siphoviruses with genomes of 101-116 kb length (Fig 7A-C). Our nine new

isolates in the BASEL collection (plus well-studied phage T5) are well representative of the two major

genera *Eseptimavirus* and *Tequintavirus* and their subclades (Fig 7B). These phages characteristically use

their lateral tail fibers to bind each one or a few types of O-antigen very specifically as their primary host

receptor, but this interaction is not essential for hosts like *E. coli* K-12 laboratory strains that don't express

an O-antigen barrier [51] (Fig 7A). As expected and observed previously [51, 60], the diversity of O-antigen

glycans is reflected in a high genetic diversity at the lateral tail fiber locus of the *Markadamsvirinae* of the

BASEL collection (S3A Fig). Only one of these phages, IrisVonRoten (Bas32), can infect *E. coli* K-12 with

restored expression of O16-type O-antigen, suggesting that it can recognize this glycan as its primary

receptor, but several others can infect different *E. coli* strains with smooth LPS or even *Salmonella* (see

below in Fig 12). Similar to what was proposed for the different small siphoviruses described above, the

terminal receptor specificity of T5 and relatives is determined by dedicated RBPs attached non-covalently

to the tip of a straight central tail fiber [51]. Compared to their smaller relatives, the repertoire of terminal

receptors among *Markadamsvirinae* seems limited – the vast majority target BtuB (shown for, e.g., EPS7

[61] and S132 [62]), and each a few others bind FhuA (T5 itself [51] and probably S131 [62]) or FepA (H8

[63] and probably S124 [62]). Sequence analyses of the RBP locus allowed us to predict the terminal

receptor of all remaining *Markadamsvirinae*, confirming that nearly all target BtuB (Fig 7B and S3B Fig).

309   It has long been known that phage T5 is largely resistant to RM systems, and this was thought to

be due to elusive DNA protection functions encoded in the early-injected genome region largely shared by

T5 and other *Markadamsvirinae* [64]. Surprisingly, we find that this resistance to restriction is a shared

feature only of the *Tequintavirus* genus, while their sister genus *Epseptimavirus* shows detectable yet

variable sensitivity in particular to the type II RM system EcoRV (Fig 7B). It seems likely that this

difference between the genera is due to the largely different number of EcoRV recognition sites in

Epseptimavirus genomes (70-90 sites, S5 Table) and *Tequintavirus* genomes (4-18 sites, S5 Table). This

observation suggests that efficient DNA ligation and RM site avoidance [65] play important roles in the

restriction resistance of T5 and relatives besides their putative DNA protection system. Remarkably, all

Markadamsvirinae are invariably sensitive to the Fun/Z immunity system of phage P2 which had already

been shown for T5 previously [50].

Fig 7. Overview of *Demerecviridae* subfamily *Markadamsvirinae*.
(A) Schematic illustration of host recognition by T5-like siphoviruses. **(B)** Maximum-Likelihood phylogeny of the *Markadamsvirinae* subfamily of *Demerecviridae* based on several core genes with bootstrap support of branches shown if > 70/100. Phages of the BASEL collection are highlighted by little phage icons and the determined or proposed terminal receptor specificity is highlighted at the phage names using the color code highlighted at the right side (same as for the small siphoviruses). The phylogeny was rooted between the *Epseptimavirus* and *Tequintavirus* genera. **(C)** The results of quantitative phenotyping experiments with *Markadamsvirinae* phages regarding sensitivity to altered surface glycans and bacterial immunity systems are presented as efficiency of plating (EOP). Data points and error bars represent average and standard deviation of at least three independent experiments.

Properties of *Myoviridae: Tevenvirinae*

Phages of the *Tevenvirinae* subfamily within the *Myoviridae* family are large myoviruses with characteristically prolate capsids and genomes of 160-172 kb size that infect a wide variety of Gram-negative hosts [30] (Figs 8A-E). The kinked lateral tail fibers of these phages contact primary receptors on the bacterial cell surface that are usually surface proteins like OmpC, Tsx, and FadL for prototypic *Tevenvirinae* phages T4, T6, and T2, respectively, but can also be sugar motifs in the LPS like in case of T4 when OmpC is not available [66] (Fig 8A). Robust interaction with the primary receptor unpins the short

tail fibers from the myovirus baseplate which enables their irreversible adsorption to terminal receptors in

the LPS core followed by contraction of the tail sheath. Consequently, the phage tail penetrates the cell

envelope and the viral genome is injected via a syringe-like mechanism [35] (Fig 8A).

Our thirteen newly isolated *Tevenvirinae* phages in the BASEL collection mostly belong to

different groups of the *Tequatrovirus* genus that also contains well-studied reference phages T2, T4, and

T6, but two isolates were assigned to the distantly related *Mosigvirus* genus (Fig 8B). As expected, the

infectivity of our thirteen isolates and of T2, T4, and T6 reference phages shows strong dependence on

each one of a small set of *E. coli* surface proteins that have previously been described as *Tevenvirinae*

primary receptors, i.e., Tsx, OmpC, OmpF, OmpA, and FadL [66] (Fig 8B). Interestingly, while for some

of these phages the absence of the primary receptor totally abolished infectivity, others still showed

detectable yet greatly reduced plaque formation (S4A Fig). It seems likely that these differences are caused

by the ability of some *Tevenvirinae* lateral tail fibers to contact several primary receptors such as, e.g.,

OmpC and the truncated *E. coli* B LPS core in case of T4 [66, 67].

Specificity for the secondary receptor depends on the short tail fibers that, in case of T4, target the

lipid A – Kdo region deep in the enterobacterial LPS core [68]. Given that this region is still present in the

waaC mutant, the most deep-rough mutant of *E. coli* K-12 that is viable (Fig 3A), it is unsurprising that T4

and some other *Tevenvirinae* did not seem to show a dependence on the LPS core in our experiments (Fig

8C). However, some *Tequatrovirus* and all tested *Mosigvirus* isolates required an intact inner core of the

host LPS for infectivity (Fig 8C). This phenotype is correlated with an alternative allele of the short tail

fiber gene that varies between *Tevenvirinae* phages irrespective of their phylogenetic position (Fig 8D). We

therefore suggest that those phages encoding the alternative allele express a short tail fiber that targets parts

of the LPS core above the lipid A – Kdo region (Figs 8D and 8E; see also Fig 3A).

Fig 8. Overview of the *Myoviridae* subfamily *Tevenvirinae*.

(A) Schematic illustration of host recognition by T4-like myoviruses. **(B)** Maximum-Likelihood phylogeny of the *Tevenvirinae* subfamily of *Myoviridae* based on a curated whole-genome alignment with bootstrap support of branches shown if > 70/100. The phylogeny was rooted between the *Tequatrovirus* and *Mosigvirus* genera. Phages of the BASEL collection are highlighted by little phage icons and experimentally determined primary receptor specificity is highlighted at the phage names using the color code highlighted at the top left. Primary receptor specificity of *Tevenvirinae* depends on RBPs expressed either as a C-terminal extension of the distal half fiber (T4 and other OmpC-targeting phages) or as separate small fiber tip adhesins [66], but sequence analyses of the latter remained ambiguous. We therefore only annotated experimentally determined primary receptors (see also S4A Fig) [53, 66]. **(C)** The results of quantitative phenotyping experiments with *Tevenvirinae* phages regarding sensitivity to altered surface glycans and bacterial immunity systems are presented as efficiency of plating (EOP). Data points and error bars represent average and standard deviation of at least three independent experiments. **(D)** The Maximum-Likelihood phylogeny *Tevenvirinae* short tail fiber proteins reveals two homologous, yet clearly distinct, clusters that correlate with the absence (variant #1, like T4) or presence (variant #2) of detectable LPS core dependence as shown in (C).

(E) The results of (D) indicate that variant #1, as shown for T4, binds the deep lipid A – Kdo region of the enterobacterial LPS core, while variant #2 binds a more distal part of the (probably inner) core.

Besides receptor specificity, the phenotypes of our diverse *Tevenvirinae* phages were highly

homogeneous. The hallmark of this *Myoviridae* subfamily is a high level of resistance to DNA-targeting

immunity like RM systems because of cytosine hypermodifications (hydroxymethyl-glucosylated for

Tequatrovirus and hydroxymethyl-arabinosylated for *Mosigvirus*) [46, 69]. Consistently, no or only very

weak sensitivity to any of the six RM systems was detected for any tested *Tevenvirinae* phage (Fig 8B).

The weak sensitivity to EcoRV observed for AugustSocin (Bas38) and phage T2 does, unlike for

Markadamsvirinae (see above), not correlate with a higher number of recognition sites and therefore

possibly depends on another feature of these phages such as, e.g., differences in DNA ligase activity (S5

Table). Conversely, all tested *Tevenvirinae* were sensitive to the Fun/Z and Tin Abi systems (Fig 8B). Tin

was previously shown to specifically target the DNA replication of T-even phages [50], and we found that

indeed all *Tevenvirinae* but no other tested phage are sensitive to this Abi system (Fig 8B). No sensitivity

was observed for RexAB, the iconic Abi system targeting *rIIA/B* mutants of phage T4 [10] (S4B),

suggesting that this Abi system is generally unable to affect wildtype *Tevenvirinae* phages (Fig 8C).

Properties of *Myoviridae: Vequintavirinae* and relatives

Phages of the *Vequintavirinae* subfamily within the *Myoviridae* family are myoviruses with

genomes of 131-140 kb size that characteristically encode three different sets of lateral tail fibers [23, 70]

(Figs 9A-E). Besides *Vequintavirinae* of the *Vequintavirus* genus, this feature is shared by two groups of

phages that are closely related to these *Vequintavirinae sensu stricto*: One group forms a cluster around

phage phAPEC8 (recently proposed as a new genus *Phapecoctavirus* within *Myoviridae* [23]), the other

group is their sister clade including phage phi92 [71] that is unclassified (Fig 9B). Despite not being

Vequintavirinae by current taxonomic classification, we are covering them together due to their

considerable similarities and propose to classify the phi92-like phages (including PaulScherrer (Bas60)) as

Nonagintaduovirus genus within the *Vequintavirinae*.

Fig 9. Overview of the *Myoviridae* subfamily *Vequintavirinae* and relatives.

(A) Schematic illustration of host recognition by *Vequintavirinae* and related myoviruses. **(B)** Maximum-Likelihood phylogeny of the *Vequintavirinae* subfamily of *Myoviridae* and relatives based on a curated whole-genome alignment with bootstrap support of branches shown if > 70/100. The phylogeny was rooted between the *Vequintavirus* genus and the two closely related, unclassified groups at the bottom. Newly isolated phages of the BASEL collection are highlighted by green phage icons. **(C)** The results of quantitative phenotyping experiments with *Vequintavirinae* and phage PaulScherrer regarding sensitivity to altered surface glycans and bacterial immunity systems are presented as efficiency of plating (EOP). Data points and error bars represent average and standard deviation of at least three independent experiments. **(D)** Amino acid sequence alignment of the lateral tail fiber Gp64 of phage N4 (*Enquatrovirus*, see below) and a lateral tail fiber conserved among *Vequintavirinae* and relatives (representatives shown). The proteins share a predicted poly-GlcNAc deacetylase domain as identified by Phyre2 [136]. **(E)** *Vequintavirinae sensu stricto* (represented by rV5 and Jeff Schatz) encode two paralogous short tail fiber proteins and a tail fiber chaperone that are homologous to the corresponding locus in phi92-like phages incl. PaulScherrer and, ultimately, to short tail fiber GpS and chaperone GpU of Mu(+) (which targets a different glucose in the K12-type LPS core GpS [95, 96]).

Three different, co-expressed lateral tail fibers have been directly visualized in the cryogenic

electron microscopy structure of phi92, and large orthologous loci are found in all *Vequintavirinae* and

relatives [23, 70, 71]. The considerable repertoire of glycan-hydrolyzing protein domains in these tail fiber

proteins has been compared to a "nanosized Swiss army knife" and is probably responsible for the

exceptionally broad host range of these phages and their ability to infect even diverse capsulated strains of

enterobacteria [23, 52, 70, 71]. However, it is far from clear which genes code for which components of

the different lateral tail fibers. While three lateral tail fiber genes exhibit considerable allelic variation

between phage genomes and might therefore encode the distal parts of the tail fibers with the receptor-

binding domains, the biggest part of the lateral tail fiber locus is highly conserved including diverse proteins

with sugar-binding or glycan-hydrolyzing domains (S5 Fig). This observation fits well with our finding that

the lysis host range of all *Vequintavirinae* available to us is remarkably homogeneous and suggests that, in

the absence of capsules and other recognized exopolysaccharides, these phages do not vary widely in their

host recognition (see below in *Host range across pathogenic enterobacteria and* E. coli *B*). Interestingly,

all *Vequintavirinae* in the BASEL collection are detectably inhibited when infecting the *wecB* knockout

(albeit to variable degree), strongly suggesting that the ECA is a shared primary receptor (Figs 9A and 9B).

This phenotype and the specific lysis host range of *Vequintavirinae* are shared with podoviruses of the

Enquatrovirus family (see below). Notably, one of the conserved lateral tail fiber proteins of the

Vequintavirinae is homologous to the lateral tail fiber protein of *Enquatrovirus* phages (Fig 9D), suggesting

that these two groups of phages might target the ECA in a similar way.

The terminal receptor of *Vequintavirinae* has been unraveled genetically for phage LL12, a close

relative of rV5 (Fig 9B), and seems to be at the first, heptose-linked glucose of the LPS outer core which is

shared by all *E. coli* core LPS types [52, 72] (Fig 9A). Consistently, we found that all *Vequintavirinae* in

the BASEL collection are completely unable to infect the *waaC* and *waaG* mutants with core LPS defects

that result in the absence of this sugar (Fig 9C, see also Fig 3A). This observation would be compatible

with the idea that all (tested) *Vequintavirinae* and also PaulScherrer of the phi92-like phages use the same

secondary receptor. Indeed, the two very similar short tail fiber paralogs of *Vequintavirinae sensu stricto*

do not show any considerable allelic variation between genomes (S5A Fig; unlike, e.g., the *Tevenvirinae*

short tail fibers presented in Figs 8D and 8E) and are also closely related to the single short tail fiber protein

of phi92-like phages or *Phapecoctavirus* (Fig 9E).

Unlike large myoviruses of the *Tevenvirinae* subfamily, the *Vequintavirinae* and relatives are

exceptionally susceptible to type II and type III RM systems, albeit with (minor) differences in the pattern

of sensitivity and resistance from phage to phage (Fig 9C). We therefore see no evidence for effective

mechanisms of these phages to overcome RM systems like, as proposed previously, covalent DNA

modifications which could be introduced by the notable repertoire of sugar-related enzymes found in their

genomes [24]. Similarly, the *Vequintavirinae* are invariably sensitive to the Fun/Z Abi system.

Properties of the *Autographiviridae* family and *Podoviridae: Enquatrovirus*

Phages of the *Autographiviridae* family and the genus *Enquatrovirus* in the *Podoviridae* family are

podoviruses that have been well studied in form of their representatives T3 and T7 (*Autographiviridae*

subfamily *Studiervirinae*) and N4 (*Enquatrovirus*). While they differ in their genome size (ca. 37-41 kb for

Studiervirinae, ca. 68-74 kb for *Enquatrovirus*) and each exhibit characteristic unique features, the overall

mode of infection of these podoviruses is the same: Lateral tail fibers of the virion contact bacterial surface

glycans at the cell surface for host recognition after which direct contact of the stubby tail with a terminal

receptor on the host triggers irreversible adsorption and DNA injection (Fig 10A).

This process has been studied in detail for the archetype of all *Autographiviridae*, phage T7, though

several open questions remain. The lateral tail fibers of this phage contact a receptor in rough LPS of *E.*

coli K-12 that is not fully understood, possibly because several alternative and overlapping sugar motifs in

the K-12 LPS core can be targeted [73-75] (Fig 10A). Conformational changes in the stubby tail tube

triggered by this receptor interaction then initiate the injection of the phage genome from the virion [75,

76]. Remarkably, at first several internal virion proteins are ejected and then fold into an extended tail that

spans the full bacterial cell envelope which, in a second step, enables DNA injection into the host cytosol

[77] (Fig 10A). Notably, our knowledge of this process does not allow the distinction between a "primary"

receptor for host recognition and a "secondary" terminal receptor for irreversible adsorption and DNA

injection. However, other *Autographiviridae* follow this classical scheme more closely and feature

enzymatic domains at their tail fibers or tail spikes that likely mediate attachment to surface glycans as

primary receptors followed by enzyme-guided movement towards the cell surface [38, 78].

The *Autographiviridae* are a large family of podoviruses hallmarked (and named) with reference

to their single-subunit T3/T7-type RNA polymerase that plays several key roles for the phage infection but

has also become a ubiquitous tool in biotechnology [79]. All *Autographiviridae* isolates of the BASEL

collection belong to several different genera within the very broad *Studiervirinae* subfamily that also

contains the classical T7 and T3 phages which we included as references (Fig 10B). Because phage T3

recognizes the peculiarly truncated R1-type LPS core of *E. coli* B (see also below) and cannot infect K-12

strains, we generated a T3(K12) chimera that encodes the lateral tail fiber gene of T7 similar as was reported

previously by others (see *Materials and Methods*) [80]. As expected, all tested *Autographiviridae* use core

LPS structures as host receptor and show impaired plaque formation on *waaC* and *waaG* mutants, but in

almost all cases some infectivity is retained even on the *waaC* mutant (Fig 10C). This suggests that, as

postulated for T7 [73-75], these phages are not strictly dependent on a single glycan motif but might

recognize a broader range target structures at the LPS core.

The overall pattern of restriction sensitivity and resistance is similar for all tested

Autographiviridae. While type I and type II RM systems are largely ineffective, type III RM systems show

remarkable potency across all phage isolates. Good part of the resistance to type II RM systems is likely

due to the near-complete absence of EcoRI and EcoRV recognition sites in the genomes of these phages as

described previously [65] (S5 Table) with the exception of ten EcoRV sites in T3(K12) which,

consequently, cause massive restriction (Fig 10C). The relatively few recognition sites for type I RM

systems do not result in considerable sensitivity for any phage, because at the *gp0.3* locus all of them either

encode an Ocr-type DNA mimic type I restriction inhibitor like T7 (*Teseptimavirus*) or an

S-adenosylmethionine (SAM) hydrolase that deprives these RM systems of their substrate like T3 (all other

genera) [9, 81, 82]. Notably, phages JacobBurckhardt (Bas63) and its close relative T7 are the only phages

tested in this work that show any sensitivity to the PifA Abi system encoded on the *E. coli* K-12 F-plasmid

(Fig 10C). Previous work showed that sensitivity of T7 to PifA immunity – as opposed to T3 which is

resistant – depends on the dGTPase Gp1.2 of the phage, though the major capsid protein Gp10 also seems

to play some role in sensitivity [83, 84]. Consequently, we find that phages T7 and JacobBurckhardt, but

not closely related *Teseptimavirus* JeanTingely (Bas62), encode a distinct variant of dGTPase Gp1.2 that

likely causes their sensitivity to PifA (Fig 10D).

Bacteriophage N4 is the archetype of *Enquatrovirus* phages that are hallmarked by using a large,

virion-encapsidated RNA polymerase for the transcription of their early genes [33], and our new

Enquatrovirus isolate AlfredRasser (Bas67) is a very close relative of N4 (Fig 10E). Based primarily on

genetic evidence, phage N4 is thought to initiate infections by contacting the host's ECA with its lateral tail

fibers [73, 85, 86] (Fig 10A). We indeed confirmed a remarkable dependence of N4 and AlfredRasser on

wecB (Fig 10C) and detected homology between the glycan deacetylase domain of the N4 lateral tail fiber

and a lateral tail fiber protein of *Vequintavirinae* which also seem to use the ECA as their primary receptor

(see above, Figs 9 and 10C). Similar to the O-antigen deacetylase tail fiber of its relative G7C (Fig 10E),

the enzymatic activity of the N4 tail fiber might provide a directional movement towards the cell surface

that is essential for the next steps of infection [38, 87] (Fig 10A). At the cell surface, the elusive outer

membrane porin NfrA was suggested to be the terminal receptor of phage N4 based on the findings that it

is required for N4 infection and interacts with the stubby tail of this phage [73, 85, 88]. To the best of our

knowledge, this would make phage N4 the first podovirus with an outer membrane protein as terminal

receptor [39, 40], though the absence of homologs to the tail extension proteins of *Autographiviridae* indeed

suggests a somewhat different mode of DNA injection (Fig 10A).

Fig 10. Overview of *Autographiviridae* phages and *Podoviridae* genus *Enquatrovirus*.
(A) Schematic illustration of host recognition by *Autographiviridae* and *Enquatrovirus* phages. **(B)** Maximum-Likelihood phylogeny of the *Studiervirinae* subfamily of *Autographiviridae* based on several core genes with bootstrap support of branches shown if > 70/100. The phylogeny was midpoint-rooted between the clade formed by *Teseptimavirus* and *Teetrevirus* and the other genera. Phages of the BASEL collection are highlighted by little phage icons. **(C)** The results of quantitative phenotyping experiments with *Autographiviridae* regarding sensitivity to altered surface glycans and bacterial immunity systems are presented as efficiency of plating (EOP). Data points and error bars represent average and standard deviation of at least three independent experiments. **(D)** Amino acid sequence alignment and Maximum-Likelihood phylogeny of Gp1.2 orthologs in all tested *Autographiviridae* phages. Phage JeanTinguely belongs to the *Teseptimavirus* genus but encodes an allele of *gp1.2* that is closely related to those of the *Berlinvirus* genus, possibly explaining its resistance to PifA (see (C)). **(E)** Maximum-Likelihood phylogeny of the *Enquatrovirus* genus and related groups of *Podoviridae* based on several core genes with bootstrap support of branches shown if > 70/100. The phylogeny was midpoint-rooted between the distantly related *Jwalphavirus* genus and the others. Phages of the BASEL collection are highlighted by little phage icons.

With regard to bacterial immunity, *Enquatrovirus* phages are highly resistant to any tested antiviral

defenses with exception to a slight sensitivity to the EcoP1_I type III RM system (Fig 10C). While their

resistance to tested type II RM systems is due to the absence of EcoRI or EcoRV recognition sites

(S5 Table), the genetic basis for their only slight sensitivity to type III RM systems is not known. We note

that *Enquatrovirus* phages encode an *rIIAB* locus homologous to the long-known yet poorly understood

rIIAB locus found in many large myoviruses which, in case of phage T4, provides resistance to the RexAB

Abi system of phage lambda [89] (S4B and S4C Figs). Both phage N4 and AlfredRasser are indifferent to

the presence of (O16-type) O-antigen or the tested truncations of the K-12 LPS core at the host cell surface

(Fig 10C), suggesting that LPS structures play no role for their infection process.

Properties of *Myoviridae*: *Ounavirinae* and classical temperate phages

The *Felixounavirus* genus in the *Ounavirinae* subfamily of *Myoviridae* comprises a group of

phages with genomes of 84-91 kb and characteristically straight lateral tail fibers of which *Salmonella*

phage Felix O1 has been most well studied [30, 90] (Figs 11A and 11B). Previous work aimed at the

isolation of *E. coli* phages differed greatly in the reported abundance of *Felixounavirus* phages, ranging

from no detection [23] over a moderate number of isolates in most studies [25, 91] to around one fourth of

all [24]. Across our phage sampling experiments we found a single *Felixounavirus* phage,

JohannRWettstein (Bas61), which is rather distantly related to Felix O1 within the eponymous genus (Fig

11B). Prototypic phage Felix O1 has been used for decades in *Salmonella* diagnostics because it lyses

almost every *Salmonella* strain but only very few other *Enterobacteriaceae* (reviewed in reference [90]).

The mechanism of its host recognition and the precise nature of its host receptor(s) have remained elusive,

but it is generally known to target bacterial LPS and not any kind of protein receptors [90]. While many

Ounavirinae seem to bind O-antigen glycans of smooth LPS, the isolation of multiple phages of this

subfamily on *E. coli* K-12 with rough LPS and on *E. coli* B with an even further truncated LPS indicate

that the functional expression of O-antigen is not generally required for their host recognition [24, 90, 92].

Phage JohannRWettstein is only slightly inhibited on *E. coli* K-12 with restored O16-type O-antigen,

suggesting that it can either use or bypass these glycans, and totally depends on an intact LPS core (Fig

11E). In addition, it is remarkably sensitive to several tested RM systems and shares sensitivity to the Fun/Z

Abi system with all other tested *Myoviridae* and the *Markadamsvirinae* (Fig 11E).

Fig 11. Overview of *Myoviridae: Ounavirinae* and classic temperate phages.
(A) Schematic illustration of host recognition by *Ounavirinae: Felixounavirus* phages. Note that the illustration shows short tail fibers simply in analogy to *Tevenvirinae* or *Vequintavirinae* (Figs 8A and 9A), but any role for such structures has not been explored for Felix O1 and relatives. **(B)** Maximum-Likelihood phylogeny of the *Ounavirinae* subfamily of *Myoiviridae* based on several core genes with bootstrap support of branches shown if > 70/100. The phylogeny was midpoint-rooted between *Kolesnikvirus* and the other genera. Our new isolate JohannRWettstein is highlighted by a green phage icon. **(C,D)** Schematic illustration of host recognition by classic temperate phages lambda, P1, and P2. Note the absence of lateral tail fibers due to a mutation in lambda PaPa laboratory strains [94]. **(E,F)** The results of quantitative phenotyping experiments with JohannRWettstein and classic temperate phages regarding sensitivity to altered surface glycans and bacterial immunity systems are presented as efficiency of plating (EOP). Data points and error bars represent average and standard deviation of at least three independent experiments.

Besides the lytic T phages, temperate phages lambda, P1, and P2 have been extensively studied

both as model systems for fundamental biology questions as well as regarding the intricacies of their

infection cycle [7, 34, 50]. Phage lambda was a prophage encoded in the original *E. coli* K-12 isolate and

forms siphovirus particles that display the GpJ tail tip to contact the LamB porin as the terminal receptor

for DNA injection [7, 93], while the lateral tail fibers (thought to contact OmpA as primary receptor) are

missing in most laboratory strains of this phage due to a mutation [7, 94] (Fig 11C). We reproduced the

dependence of our lambda*vir* variant on LamB and, as reported previously, found that an intact inner core

of the LPS was required for infectivity [73 and literature cited therein] (Fig 11F and S5 Table). Myoviruses

P1 and P2 were both shown to require a rough LPS phenotype to contact their receptors in the LPS core,

but the exact identity of these receptors has not been unraveled [40, 50, 73, 95]. While the molecular details

of the infection process after adsorption have not been well studied for P1, previous work showed that an

interaction of the P2 lateral tail fibers with the LPS core of K-12 strains triggers penetration of the outer

membrane by the tail tip [40, 50, 96] (Fig 11D). Consistently, our results confirm that mutations

compromising the integrity of the K-12 core LPS abolished bacterial sensitivity to P1*vir* and P2*vir* [50, 73,

95] (Fig 11F). The quantification of these phages' sensitivity to different immunity systems revealed that

they are remarkably sensitive to all tested RM systems (Fig 11F), a property only shared with a few

Drexlerviridae (Fig 4D). As expected from the considerably lower number of restriction sites, the

sensitivity was less pronounced for type I RM systems (Fig 11F and S5 Table). Phage lambda was

additionally specifically sensitive to the Old Abi system of the P2 prophage (Fig 11F), as shown previously

[50].

Host range across pathogenic enterobacteria and laboratory wildtype *E. coli* B

The host range of bacteriophages has repeatedly been highlighted as a critical feature for phage

therapy because broad infectivity can enable phages to be used against different strains of the same pathogen

without repeated sensitivity testing "analogous to the use of broad-spectrum antibiotics" [16]. Intuitively,

across strains of the same host species the infectivity of phages depends on their ability to successfully bind

the variable surface structures of host cells and to overpower or evade strain-specific bacterial immunity

[16, 97]. While simple qualitative tests assessing the lysis host range primarily inform about host

recognition alone, the more laborious detection of robust plaque formation as a sign of full infectivity is the

gold standard of host range determination [16, 97]. We therefore challenged a panel of commonly used

pathogenic enterobacterial strains with the BASEL collection and recorded the phages' lysis host range as

 well as plaque formation separately to gain insight into their host recognition and the ability to overcome

 immunity barriers inside host cells (Fig 12; see *Materials and Methods*).

Fig 12. Host range of phages in the BASEL collection.

(A) Surface glycans of the enterobacterial strains used in this work (see *Materials and Methods* for details on how the illustration of glycan chains was composed). **(B)** The ability of all phages in the BASEL collection to infect different enterobacteria was studied qualitatively (lysis host range; top) and, more stringently, based on the ability to form plaques (bottom). Top: The observartion of lysis zones with high-titer lysate (>10^9 pfu/ml) in at least three independent experiments is indicated by colored bars. Bottom: The infectivity of BASEL collection phages on diverse enterobacterial hosts was quantified as efficiency of plating (EOP). Data points and error bars represent average and standard deviation of at least three independent experiments. Since the data obtained with *Salmonella* Typhimurium 12023s and SL1344 were indistinguishable, we only show the results of one representative strain (*S.* Typhimurium 12023s).

As expected from previous work, *Vequintavirinae* phages showed an outstanding lysis host range

[23, 25, 70, 71] and invariably infected *E. coli* UTI89, *E. coli* 55989, and *E. coli* K-12 with restored O16-

type O-antigen (Figs 9C and 12B). The diversity of infected hosts suggests that *Vequintavirinae* can readily

bypass the O-antigen barrier, but the homogeneity of their lysis host ranges is clearly at odds with the

polyvalent nature of their virions that display several different lateral tail fibers and variable RBPs [70, 71]

(S5 Fig). Notably, we showed that *Enquatrovirus* phages share the exact same host range and an

insensitivity to the O-antigen barrier with *Vequintavirinae* (Figs 10C and 12B), possibly because these

phages target the highly conserved ECA as primary receptor using a homologous tail fiber (Fig 9D).

Recognition of the ECA molecules among smooth LPS (Fig 12A) might enable *Vequintavirinae* and

Enquatrovirus to bypass the O-antigen barrier and move to the cell surface along this glycan chain, possibly

using the deacetylase domain of their shared tail fiber as previously described for *Salmonella* phage P22

and the N4 relative G7C [38, 87] (Fig 10E). If true, the *Vequintavirinae* would be effectively monovalent

on the tested hosts and might primarily use their "nanosized Swiss army knife" of tail fibers to bind and

overcome capsules or other more specialized exopolysaccharides [70, 71]. Given the remarkable lysis host

range of *Vequintavirinae* and *Enquatrovirus*, it would be valuable to explore the molecular basis of their

host recognition in future studies in order to use this knowledge for phage host range engineering [14].

A decently broad lysis host range was also observed for *Tevenvirinae* phages (Fig 12B), in line

with previous work [23], though more scattered and far less remarkable than for the *Vequintavirinae*. For

both groups of phages it is astonishing how poorly their broad host recognition is reflected by actual plaque

formation (Fig 12B). Despite robust "lysis-from-without" in some cases [98], no reliable plaque formation

on any other host than *E. coli* K-12 was observed for *Vequintavirinae*, and the range of plaque formation

of *Tevenvirinae* was also severely contracted relative to the lysis host range (Fig 12B).

Besides *Vequintavirinae* and *Enquatrovirus* that can apparently bypass the O-antigen barrier, we

find that the lysis host range of around a third of the tested phages (24 / 61) included at least one host strain

with smooth LPS, while the O16-type O-antigen of *E. coli* K-12 blocked infections by all but five of these

phages scattered across the different taxonomic groups (5 / 61; Fig 12; see also Figs 4 and 6-11). These

results confirm an important role of the O-antigen as a formidable barrier to bacteriophage infection that

can only be overcome by specific recognition with tail fibers to breach it, e.g., via enzymatic activities [35].

Notably, less than half of the phages that could lyse any strain besides the *E. coli* K-12 ΔRM isolation host

showed robust plaque formation on that strain (Fig 12B), probably due to different layers of bacterial

immunity. In comparison to *Vequintavirinae* and *Tevenvirinae*, it is surprising how the lysis host range of

Markadamsvirinae is almost exactly reflected in their range of plaque formation (Fig 12B). One of these

phages, SuperGirl (Bas29) is the only phage in the BASEL collection that shows robust plaque formation

on *Salmonella* Typhimurium (Fig 12B).

The *E. coli* B lineage comprises a number of laboratory strains including REL606 that have, like

E. coli K-12, lost O-antigen expression and even part of its R1-type LPS core during domestication but are

significant as the original hosts of the T phages [5, 99]. In line with the barrier function of the O-antigen

(see above), the vast majority of phages in the BASEL collection can at least lyse this strain with the notable

exception of *Vequintavirinae* and *Enquatrovirus* (Fig 12B). We can only speculate about the molecular

basis of this observation, but maybe the *Vequintavirinae* are unable to use the truncated core LPS of this

strain for their infections (Figs 12A and 12B; compare Fig 9C). If inherited from earlier *E. coli* B strains, it

seems likely that this total resistance to *Vequintavirinae* is the reason why the T phages contain no

representative of this group despite their abundance. Similarly, the observation that phage T5 with its FhuA

receptor (Fig 7B) is a rather unusual member of the *Markadamsvirinae* can be explained by the loss of *btuB*

in older variants of *E. coli* B, but this mutation reverted in an ancestor of *E. coli* B REL606 [100].

Consequently, this modern *E. coli* B strain is sensitive to the various BtuB-targeting *Markadamsvirinae*

(Fig 12B). Similarly, *E. coli* B strains lack expression of *ompC* and the REL606 strain additionally lacks

functional *tsx* that both encode common primary receptors of the *Tevenvirinae* (Fig 8B) [66, 100]. Though

phage T4 itself can use the *E. coli* B LPS core as a primary receptor when OmpC is absent [67], the inability

of several other *Tevenvirinae* to infect *E. coli* B REL606 corresponds well to the dependency on OmpC and

Tsx as primary receptors (Figs 8B, 12B, and S4A). A number of additional phages of, e.g., the *Nonagvirus*

genus can lyse but not form plaques on *E. coli* B REL606 (Fig 12B). Based on previous work, we suggest

that this phenotype is caused by immunity systems encoded in the specific repertoire of cryptic prophages

of *E. coli* B strains that differ considerably between *E. coli* B and K-12 strains and are known to harbor

active immunity systems [100].

Discussion

Remarkable patterns of bacteriophage receptor specificity

Our results regarding the terminal receptor specificity of different groups of siphoviruses inherently

present the question why all these phages target less than ten of the more than 150 outer membrane proteins

of *E. coli* K-12 of which around two dozen are porins [39, 40, 101] (Figs 4C and 7B). Similarly,

Tevenvirinae use only five different outer membrane proteins as primary receptors (Fig 8B) [39, 40].

Previous work suggested that this bias might be linked to the abundance of the targeted proteins, both via

a preference for particularly numerous proteins (favoring phage adsorption) or very scarce ones (avoid

competition among phages) [102]. However, we feel that this stark bias in phage preference might be largely

driven by functional constraints. Notably, the large siphoviruses of *Markadamsvirinae* bind exactly a subset

of those terminal receptors targeted by the small siphoviruses despite using non-homologous RBPs, while

there is no overlap between these proteins and the primary receptors of the *Tevenvirinae* [39, 40] (Figs 4-8).

It seems therefore likely that the receptors targeted by siphoviruses need to have certain properties, e.g.,

favoring DNA injection, that greatly limit the repertoire of suitable candidates. We envision that future

studies might unravel the molecular mechanisms of how siphoviruses recognize and use outer membrane

proteins as terminal receptors on Gram-negative hosts similar to, e.g., how it has been studied for

myoviruses and podoviruses that directly puncture the outer membrane [68, 77].

Many open questions remain regarding the host recognition and receptor specificity of *E. coli*

phages that could be tackled with a combination of systematic phenotypic and genomic analyses. Why do

some small siphoviruses in *Drexlerviridae* and *Siphoviridae* but none of the larger ones of *Demerecviridae:*

Markadamsvirinae strongly depend on an intact LPS core, seemingly independent of primary receptor

usage or terminal receptor specificity (Figs 4D, 6C, 6F, and 7B)? Is it possible to perform analyses like we

did it for the RBPs of siphoviruses or *Tevenvirinae* for the much more variable lateral tail fibers (S1B, S2A,

and S3A Figs) that probably bind to the around 200 O-antigen types of *E. coli* [103]? And is there a

genetically encoded specificity for the inner membrane channels used by siphoviruses (like ManYZ by

phage lambda or PtsG for HK97 [7]) similar to the mix-and-match of RBPs and outer membrane proteins?

Addressing these questions in future studies would not only greatly expand our knowledge regarding the

molecular basis of bacteriophage ecology and evolution, but would also help making use of this knowledge

to optimize the application of phages in biotechnology and for therapeutic applications, e.g., for "phage

steering" [14, 104].

Bacteriophage sensitivity and resistance to host immunity

Our systematic analysis of the sensitivity and resistance profiles across the different taxonomic of

phages of the BASEL collection revealed several strong patterns that are informative about underlying

molecular mechanisms. While some groups of phages like *Tevenvirinae* or *Nonagvirus* and *Seuratvirus*

genera are highly homogeneous in their profiles of sensitivity or resistance (Figs 6F and 8C) – probably

largely driven by conserved DNA modifications – others are more heterogeneous but never showed merely

random differences. For example, the two genera of *Markadamsvirinae* differ systematically in their

resistance to EcoRV (Fig 7C), likely due to vastly different numbers of recognition sites (S5 Table). Specific

deviations of single phages from these patterns are sometimes readily explained like, e.g., the PifA

sensitivity of T7 and JacobBurckhardt (due to their *gp1.2* allele, Figs 10C and 10D) or the EcoRV sensitivity

of T3 (as the only podovirus failing in RM site avoidance regarding EcoRV; S5 Table and Fig 10C).

Notably, podoviruses generally show a remarkable lack of recognition sites of type I and type II RM systems

(S5 Table) which makes them phenotypically resistant [65], but this evolutionary strategy does not seem to

be effective for type III RM systems (Fig 11C), possibly due to the abundance of their (shorter) recognition

sequences (Fig 3B and S5 Table). Overall, we observed that phages with bigger genomes such as

Tevenvirinae, *Vequintavirinae*, and *Markadamsvirinae* are broadly targeted by Abi systems, while smaller

siphoviruses or podoviruses are either not or only sparingly targeted (Figs 4 and 6-11, see also S3 Text).

Temperate phages in general seem to be highly sensitive to any kind of host immunity (Fig 11F), possibly

because their evolution is less driven by selection to overcome host defenses but rather by optimizing the

lysogens' fitness, e.g., by providing additional bacterial immunity systems [8, 9].

Regarding the immunity systems themselves, we confirmed the intuitive expectation that the

potency of RM systems is linked to the number of recognition sites in each phage genome unless they are

masked by DNA modifications. Consequently, large phages with many recognition sites (like

Vequintavirinae) are exceptionally sensitive to RM systems (Fig 8C), while type I RM systems perform

very poorly against most phages because their long recognition sites are rare (Fig 3B and S5 Table; see also

Figs 4 and 6-11). For several groups of phages like *Markadamsvirinae* and *Tevenvirinae*, the EcoRV system

is the only RM system with significant impact (Figs 7C and 8C). We suggest that this is due to the blunt-

cutting activity of EcoRV that is much more difficult to seal for DNA repair than the sticky ends introduced

by the other tested RM systems (Fig 3B), though phage DNA ligases are known to differ in their efficiency

on different kinds of DNA breaks [105].

One surprising result of our phenotyping was the remarkably poor target range of the most well-

studied Abi systems of *E. coli* K-12, RexAB and PifA, that each affected only two or three phages (Figs 4

and 6-11). We do not think that this finding is an artifact of, e.g., the genetic constructs that we used,

because both systems protected its host very well against the phages that they were known to target [12]:

An *rIIAB* mutant of T4 was highly sensitive to RexAB (S4B Fig), and phage T7 was unable to infect a host

with PifA (Fig 10C). In stark difference to RexAB and PifA, the three Abi systems of phage P2 protected

each against a considerable number of phages, especially Fun/Z that inhibited all *Tevenvirinae*,

Vequintavirinae, and *Markadamsvirinae* plus the *Felixounavirus* JohannRWettstein (Figs 4 and 6-11).

Despite this remarkable potency, the molecular mechanism of Fun/Z immunity has remained unknown [50].

A common first step to unraveling the activities of an Abi system is the analysis of insensitive phage escape

mutants which can be particularly insightful if they were isolated from several very different phages [106].

It seems clear that the BASEL collection as a well-assorted set of very diverse phages could be an effective

tool for this purpose and also in general for a first phenotypic profiling of novel immunity systems that are

commonly tested against much smaller, less well-defined sets of phages [106, 107].

Though the technicalities of our immunity phenotyping are subject to a few minor caveats (see

Materials and Methods and S3 Text), our results demonstrate how the BASEL collection can be used to

explore the biology of bacterial immunity systems and the underlying molecular mechanisms. Several

important open questions remain: What is the evolutionarily or mechanistic reason for the stark differences

in Abi system target range? To which extent do the different layers of bacterial immunity limit

bacteriophage host range? Why are some groups of phages like the *Tevenvirinae* so strongly targeted by

seemingly unrelated Abi systems and other groups of phages apparently not at all? And how well would it

be possible, based on systematic phenotypic data of the BASEL collection or extensions thereof, to predict

the sensitivity of newly isolated phages to diverse immunity systems just based on their genome sequences?

Trade-offs between phage traits limit their effective host range

Our results show very clearly that seemingly advantageous phage traits such as the broad host

recognition of *Vequintavirinae* or the remarkable resistance of *Tevenvirinae* to RM systems do not confer

these phages a particularly large effective host range (Figs 8C and 12B). This observation can be explained

by strong trade-offs between different phage fitness traits as proposed already in earlier work [108]. Unlike

the hypothetical "Darwinian Demon" that would maximize all fitness traits simultaneously and dominate

its ecosystem alone [109], the limits imposed by these trade-offs instead drive the adaptation of different

phage groups towards specific niches that enable their long-term coexistence [108]. In marine

environments, the coexistence of a few specialized, highly successful phage groups was proposed to be

stable in space and time because extinction of individual phages would usually result in their replacement

by relatives from the same, highly adapted group [110]. This "royal family model" could also explain why

the same groups of *E. coli* phages sampled in our work have already been found again and again in previous

studies that sampled diverse other environments [23-29] (Fig 2).

With our data we can at least shed light on the molecular basis of the proximal causes of the

trade-offs between phage fitness traits that limit very broad effective host ranges in the selected example:

Vequintavirinae phages are highly sensitive to various RM systems (Fig 9C), while the *Tevenvirinae* appear

to be a major target of different Abi systems and have to evade a variety of those already only to infect

regular *E. coli* K-12 (Figs 8C, S4B, and S4C; see also S3 Text) [9, 10, 12]. The deeper evolutionary or

mechanistic basis of our observations is not clear – as an example, phages of *Nonagvirus* and *Seuratvirus*

genera with genomes that are far less than half the size of *Vequintavirinae* afford full pathways for

guanosine modifications that protect them reasonably well from RM systems (Figs 6E and 6F). It is not

intuitive why natural selection has not resulted in similarly effective mechanisms for the *Vequintavirinae*,

particularly because their promiscuity in host adsorption should bring them into very frequent contact with

the diverse RM systems that form a major pillar of *E. coli* immunity [11, 49]. A good illustration for this

paradox is phage PaulScherrer (Bas60), a phi92-like relative of the *Vequintavirinae* (Fig 9B), that is the

only phage in the BASEL collection lysing all tested host strains but without robust plaque formation on

any strain besides *E. coli* K-12 (Fig 12B).

The observed trade-offs between desired phage traits suggest that newly isolated phages might

greatly profit from improvements by experimental evolution or genetic engineering to help them escape

these constraints at least long enough for applications like phage therapy [14]. Such improvements seem to

be principally feasible without immediate loss of fitness because, e.g., phage FriedrichZschokke (Bas41)

of the *Tevenvirinae* displays only weak sensitivity to the Tin and Fun/Z Abi systems that are highly potent

against its relatives (Fig 8C). However, targeted improvements of phage properties directly depend on

fundamental knowledge of the molecular mechanisms underlying receptor specificity, the phages' anti-

immunity systems, and their host counterparts that are frequently unavailable [14, 41]. Future studies

targeting the fundamental biology of phage-host interactions could therefore be directly helpful for such

applications, and for *E. coli* phages the BASEL collection might be a suitable tool for such research.

A different way to rationalize the paradox highlighted above is that the observed trade-offs might

be less pronounced in natural environments, e.g., when bacteria are starved, stressed, in biofilms, or in some

other physiological state not well reflected by our laboratory experiments. Consistently, the insufficiency

of regular phage phenotyping using standard laboratory growth conditions has already been recognized

previously, but only limited data comparing, e.g., different host physiologies are available [16, 111]. Further

studies exploring how phage infectivity, host range, and sensitivity to immunity systems change under

different growth conditions would therefore be important to better understand the challenges for successful

phage therapy and, more generally, the ecology and evolution of phages in natural environments.

Concluding remarks

A recent landmark study systematically explored the genetic profile of host requirements for several

E. coli phages and uncovered multiple unexpected and exciting twists of phage biology such as the

dependence of phage N4 on high levels of the second messenger cyclic di-GMP [73]. Our current study

based on the BASEL collection represents a complementary approach to explore the biology of *E. coli*

phages not with elaborate genome-wide screens on the host side but rather by combining phenotypic and

genomic analyses of a well-assorted set of bacteriophages. With this strategy we achieved significant

advances in bacteriophage biology regarding the recognition of host receptors, the sensitivity or resistance

of phages to host immunity, and how these factors come together to determine bacteriophage host range.

Our work therefore establishes the BASEL collection as a powerful tool to explore new aspects of

bacteriophage biology by unraveling links between phage phenotypes and their genome sequences.

Furthermore, our extensive characterization of the most abundant lineages of *E. coli* phages also provides

a useful field guide for teaching and outreach activities analogous to the successful SEA-PHAGES initiative

[112] and the diverse student and high school projects that have enabled our study (see *Acknowledgments*

and S5 Table).

Materials and Methods

Preparation of culture media and solutions

Lysogeny Broth (LB) was prepared by dissolving 10 g/l tryptone, 5 g/l yeast extract, and 10 g/l

sodium chloride in Milli-Q H_2O and sterilized by autoclaving. LB agar plates were prepared by

supplementing LB medium with agar at 1.5% w/v before autoclaving. The M9RICH culture medium was

conceived as a variant of regular M9 medium [113] supplemented with trace elements and 10% v/v LB

medium prepared without NaCl to promote the growth of diverse enterobacterial strains. It was prepared

from sterilized components by mixing (for 50 ml) 33.75 ml Milli-Q H_2O, 10 ml 5x M9 salts solution, 5 ml

LB medium without NaCl, 500 µl 40% w/v D-glucose solution, 100 µl 1 M $MgSO_4$, and 5 µl 1 M $CaCl_2$

using sterile technique. Unless indicated otherwise, all components were sterilized by filtration (0.22 µm).

Phosphate-buffered saline (PBS) was prepared as a solution containing 8 g/l NaCl, 0.2 g/l KCl,

1.44 g/l $NA_2HPO_4 \cdot 2H_2O$, and 0.24 g/l KH_2PO_4 with the pH adjusted to 7.4 using 10 M NaOH and sterilized

by autoclaving. SM buffer was prepared as 0.1 M NaCl, 10 mM $MgSO_4$, and 0.05 M Tris (pH 7.5) using

sterile technique.

Bacterial handling and culturing

745   *Escherichia coli* and *Salmonella* Typhimurium strains were routinely cultured in LB medium at

746   37°C in glass culture tubes or Erlenmeyer flasks with agitation at 170rpm. For all phenotyping assays, the

747   bacteria were instead grown in M9RICH which supports robust growth of all strains to high cell densities.

748   LB agar plates were routinely used as solid medium. Selection for genetic modifications or plasmid

749   maintenance was performed with ampicillin at 50 µg/ml, kanamycin at 25 µg/ml, and zeocin at 50 µg/ml.

Bacteriophage handling and culturing

Bacteriophages were generally cultured using the double-agar overlay method [114] with a top agar prepared as LB agar with only 0.5% w/v agar supplemented with 20 mM MgSO$_4$ and 5 mM CaCl$_2$. Top agar plates were incubated at 37°C and plaques were counted as soon as they were identified by visual inspection. However, the plates were always incubated for at least 24 hours to record also slow-growing plaques. We routinely used the improvements of classical phage techniques published by Kauffman and Polz [115].

High-titer stocks of bacteriophages were generated using the plate overlay method. Briefly, top agar plates were set up to grow almost confluent plaques of a given phage and then covered with 12 ml of SM buffer. After careful agitation for 24 hours at 4°C, the suspension on each plate was pipetted off and centrifuged at 8'000g for 10 minutes. Supernatants were sterilized with few drops of chloroform and stored in the dark at 4°C. For archiving, bacteriophages were stored as virocells at -80°C [116]

Bacterial strains and strain construction

All bacterial strains used in this work are listed in S1 Table, all oligonucleotide primers in S2 Table, and all plasmids in S3 Table.

Escherichia coli K-12 MG1655 ΔRM

The standard laboratory strain *E. coli* K-12 MG1655 (CGSC #6300) was engineered into a more permissive host for bacteriophage isolation by knocking out the EcoKI type I restriction-modification system (encoded by *hsdRMS*) and the McrA, Mrr, and McrBC type IV restriction systems using lambda red recombineering [117] (see Text S1 and our previous work for more technical details [118]). The resulting strain lacks all known *E. coli* K-12 restriction systems and was therefore called *E. coli* K-12 ΔRM. To enable isolation of bacteriophages with tropism for the sex pilus of the F-plasmid conjugation system, we supplied *E. coli* K-12 MG1655 ΔRM with a variant of the F-plasmid in which the *pifA* immunity gene (which might otherwise have interfered with bacteriophage isolation) had been replaced with a zeocin

resistance cassette by one-step recombineering (see S1 Text for details). The strain was additionally transformed with plasmid pBR322_ΔP*tet* of reference [119] as an empty vector control for experiments with plasmid-encoded immunity systems (see below).

Escherichia coli K-12 MG1655 ΔRM mutants with altered surface glycans

To test the effect of altered surface glycans of bacteriophage infection, we generated derivatives of *E. coli* K-12 MG1655 ΔRM in which genes linked to specific glycans were knocked out. The *waaC* and *waaG* genes of the LPS core biosynthesis pathway were knocked out to generate mutants displaying a deep-rough or extremely deep-rough phenotype (Fig 3A). Expression of the O16-type O-antigen of *E. coli* K-12 was restored by precisely removing the IS5 element disrupting *wbbL* [44] (Fig 3A). These mutants were generated by two-step recombineering (see details in S1 Text and a list of all strains in S1 Table). A strain specifically lacking the enterobacterial common antigen (ECA) was obtained by flipping out the kanamycin resistance cassette of the *wecB* mutant of the KEIO collection using plasmid pCP20 [117, 120]. This strain is deficient in WecB, the enzyme synthesizing UDP-ManNAcA (UDP-N-acetyl-D-mannosaminuronic acid), which is specifically required for the synthesis of the ECA but no other known glycan of *E. coli* K-12 [43].

Escherichia coli K-12 BW25113 *btuB* and *tolC* knockout mutants

btuB and *tolC* knockout mutants isogenic with the KEIO collection were generated to use these strains lacking known bacteriophage receptors for qualitative top agar assays (see below). Details of the strain construction are provided in S1 Text.

Other enterobacterial strains used for bacteriophage phenotyping

E. coli B REL606 is a commonly used laboratory strain with a parallel history of domestication to *E. coli* K-12 MG1655 [99]. *E. coli* UTI89 (ST95, O18:K1:H7) and *E. coli* CFT073 (ST73, O6:K2:H1; we used the variant with restored *rpoS* described previously [121]) are commonly used as model strains for uropathogenic *E. coli* (UPEC) and belong to phylogroup B2 [122]. *E. coli* 55989 (phylogroup B1, ST678,

798   O104:H4) is commonly used as a model strain for enteroaggregative *E. coli* (EAEC) and closely related to

799   the Shiga toxin-producing *E. coli* which caused the 2011 outbreak in Germany [123, 124]. *Salmonella*

800   *enterica* subsp. *enterica* serovar Typhimurium strains 12023s (also known as ATCC 14028) and SL1344

801   are both commonly used in laboratory experiments but exhibit phylogenetic and biological differences

802   [125].

Plasmid construction

Plasmid vectors were generally cloned following the method of Gibson et al. ("Gibson Assembly")

[126] in which two or more linear fragments (usually PCR products) are ligated directionally guided by

short 25 bp overlaps. Initially, some plasmids were also constructed using classical restriction-based

molecular cloning. Briefly, a PCR-amplified insert and the vector backbone were each cut with appropriate

restriction enzymes (New England Biolabs). After dephosphorylation of the backbone (using FastAP

dephosphorylase; Thermo Scientific), insert and backbone were ligated using T4 DNA ligase (Thermo

Scientific). Local editing of plasmid sequences was performed by PCR with partially overlapping primers

as described by Liu and Naismith (127). *E. coli* strain EC100 *pir(+)* was used as host for all clonings. The

successful construction of every plasmid was confirmed by Sanger Sequencing. A list of all plasmids used

in this study is found in S3 Table and their construction is summarized in S4 Table. The sequences of all

oligonucleotide primers used in this study are listed in S2 Table.

For the series of plasmids encoding the eleven different bacterial immunity systems studied in this

work (see Fig 3B), we used the EcoRI and EcoRV constructs of Pleška, Qian et al. [119] as templates and,

consequently, the corresponding empty vector pBR322_ΔP*tet* as general cloning backbone and

experimental control. The different immunity systems were generally cloned together with their own

transcriptional promoter region. However, the *rexAB* genes are transcribed together with the cI repressor of

the lambda prophage [7]. We therefore cloned them directly downstream of the P*tet* promoter of the

pBR322 backbone and obtained a functional construct (validated in S4B Fig). The EcoKI, EcoRI, EcoRV,

and EcoP1_I RM systems have been studied intensively in previous work [47-49]. Besides the type IA RM

system EcoKI we also cloned EcoCFT_I that is nearly identical to the well-characterized type IB RM

system EcoAI but encoded in the genome of *E. coli* CFT073 that was available to us [47, 128]. Similarly,

the EcoCFT_II system was identified as a type III RM system in the *E. coli* CFT073 genome using REBASE

[49, 128]. Due to problems with toxicity, some immunity systems were cloned not into pBR322_ΔP*tet* but

rather into a similar plasmid carrying a low-copy SC101 origin of replication (pAH186SC101e [121], see

S3 and S4 Tables as well as the considerations in S3 Text). Since we failed to obtain any functional construct

for the ectopic expression of *pifA* (as evidenced by lack of immunity against T7 infection), we instead

replaced the F(*pifA::zeoR*) plasmid in *E. coli* K-12 MG1655 ΔRM with a wildtype F plasmid that had

merely been tagged with kanamycin resistance and encodes a functional *pifA* (pAH200e).

Bacteriophage isolation

Basic procedure

Bacteriophages were isolated from various different samples between July 2019 and November

2020 using *E. coli* K-12 MG1655 ΔRM as the host (see S5 Table for details) using a protocol similar to

common procedures in the field [16]. Phage isolation was generally performed without an enrichment step

to avoid biasing the isolation towards fast-growing phages (but see below).

For aqueous samples we directly used 50 ml, while samples with major solid components (like soil

or compost) were agitated overnight at 4°C in 50 ml of PBS to release viral particles. Subsequently, all

samples were centrifuged at 8'000 g for 15 minutes to spin down particles larger than viruses. The

supernatants were sterilized treated with 5% v/v chloroform which safely inactivates any bacteria as well

as enveloped viruses but will generally leave most *Caudovirales* intact [16]. Subsequently, viral particles

were precipitated by adding 1 ml of a 2 M $ZnCl_2$ solution per 50 ml of sample, mixing shortly by inversion,

and incubating the suspension at 37°C without agitation for 15 minutes [129]. After precipitation, the

samples were centrifuged again at 8'000 g for 15 minutes and the supernatant was discarded. The pellets

were carefully resuspended in each 500 µl of SM buffer by agitation at 4°C for 15 minutes. Subsequently,

the suspensions were cleared quickly using a tabletop spinner and mixed with 500 µl of bacterial overnight

culture (resuspended in fresh LB medium to induce resuscitation). After incubation at room temperature

for 15 minutes to promote phage adsorption, each mixture was added to 9 ml of pre-warmed top agar and

poured onto a pre-warmed square LB agar plate (ca. 12 cm x 12 cm). After solidification, the plates were

incubated at 37°C for up to 24 hours.

Isolation of bacteriophage clones

Bacteriophages were visible as plaques forming in the dense bacterial growth of the top agar. For

isolation of bacteriophage clones, they were picked from clearly separated plaques of diverse morphologies

with sterile toothpicks and propagated at least three times via single plaques on top agars of the isolation

host strain *E. coli* K-12 MG1655 ΔRM. To avoid isolating temperate phages or phages that are poorly

adapted to *E. coli* hosts, we only picked clear plaques (indicative of lytic phages) and discarded isolates

that showed poor plaque formation [16].

Isolation of *Autographiviridae* using enrichment cultures

The direct plating procedure outlined above never resulted in the isolation of phages belonging to

the *Autographiviridae* like iconic T phages T3 and T7. Given that these phages are known for fast

replication and high burst sizes [130], we therefore performed a series of enrichment culture isolation

experiments to obtain phages forming the characteristically large, fast-growing plaques of

Autographiviridae. For this purpose, we prepared M9 medium using a 5x M9 salts solution and chloroform-

sterilized sewage plant inflow instead of water (i.e., containing ca. 40 ml of sewage plant inflow per 50 ml

of medium) and supplemented it with 0.4% w/v D-glucose as carbon source. 50 ml cultures were set up by

inoculating these media with each 1 ml of an *E. coli* K-12 MG1655 ΔRM overnight culture and agitated

the cultures at 37°C for 24 hours. Subsequently, the cultures were centrifuged at 8'000 g for 15 minutes

and each 50 µl of supernatant was plated with the *E. coli* K-12 MG1655 ΔRM isolation strain in a top agar

on one square LB agar plate. After incubation at 37°C for three or four hours, the first *Autographiviridae*

plaques characteristically appeared (before most other plaques) and were picked and propagated as

described above. Using this procedure, we isolated four different new *Autographiviridae* isolates (see S5

Table and Fig 10B).

Composition of the BASEL collection

Bacteriophages were mostly isolated and characterized from randomly picked plaques in direct

selection experiments, but we later adjusted the procedure to specifically isolate *Autographiviridae* which

were the only major group of phages previously shown to infect *E. coli* K-12 missing from our collection

(see above). After every set of 10-20 phages that had been isolated, we performed whole-genome

sequencing and preliminary phylogenetic analyses to keep an overview of the growing collection (see

below). In total, more than 120 different bacteriophages were sequenced and analyzed of which we selected

66 tailed, lytic phage isolates to compose the BASEL collection (see Fig 2 and S5 Table for details). Phages

closely related to other isolates were deliberately excluded unless they displayed obvious phenotypic

differences such as, e.g., a different host receptor. In addition to the 66 newly isolated bacteriophages, ten

classical model phages were included for genomic and phenotypic characterization, and we view these

phages as an accessory part of the BASEL collection. These ten phages were six of the seven T phages

(excluding T1 because it is a notorious laboratory contaminant [32]), phage N4, and obligately lytic mutants

of the three well-studied temperate phages lambda, P1, and P2 [5, 7, 33, 34] (Fig 2; see also S5 Table). To

generate the T3(K12) chimera, the 3' end of *gp17* lateral tail fiber gene of phage T7 was cloned into low-

copy plasmid pUA139 with flanking regions exhibiting high sequence similarity to the phage T3 genome

(generating pUA139_T7(*gp17*), see S4 Table). Phage T3 was grown on *E. coli* B REL606 transformed with

this plasmid and then plated on *E. coli* K-12 MG1655 ΔRM to isolate recombinant clones. Successful

exchange of the parental T3 *gp17* allele with the variant of phage T7 was confirmed by Sanger Sequencing.

Qualitative top agar assays

The lysis host range of isolated bacteriophages on different enterobacterial hosts and their ability

to infect strains of a set of KEIO collection mutants lacking each one surface protein (or isogenic mutants

that were generated in this study, see S1 Table and S1 Text) were tested by qualitative top agar assays. For

this purpose, top agars were prepared for each bacterial strain on LB agar plates. For by overlaying them

with top agar supplemented with a suitable bacterial inoculum. For regular round Petri dishes (ca. 9.4 cm

diameter) we used 3 ml of top agar supplemented with 100 µl of bacterial overnight culture, while for larger

square Petri dishes (ca. 12 cm x 12 cm) we used 9 ml of top agar supplemented with 200 µl of overnight

culture. After solidification, each 2.5 µl of undiluted high-titer stocks of all tested bacteriophages ($>10^9$

pfu/ml) were spotted onto the top agar plates and dried into the top agar before incubation at 37°C for at

least 24 hours. If lysis zones on any enterobacterial host besides our *E. coli* K-12 ΔRM reference strain

were observed, we quantified phage infectivity in efficiency of plating assays (see below). Whenever a

phage failed to show lysis on a mutant strain lacking a well-known phage receptor, we interpreted this result

as indicating that the phage infecting depends on this factor as host receptor.

Efficiency of Plating assays

The infectivity of a given bacteriophage on a given host was quantified by determining the

efficiency of plating (EOP), i.e., by quantitatively comparing its plaque formation on a certain experimental

host to plaque formation on reference strain *E. coli* K-12 MG1655 ΔRM carrying F(*pifA::zeoR*) and

pBR322_ΔP*tet* [131]. Experimental host strains we identical to the reference strain with the difference that

they either carried plasmids encoding a certain bacterial immunity system (S3 Table) or had a chromosomal

modification changing surface glycan expression (Fig 3A and S1 Table).

For quantitative phenotyping, top agars were prepared for each bacterial strain on LB agar plates

by overlaying them with top agar (LB agar containing only 0.5% agar and additionally 20 mM $MgSO_4$ as

well as 5 mM $CaCl_2$; stored at 60°C) supplemented with a suitable bacterial inoculum. For regular round

Petri dishes (ca. 9.4 cm diameter) we used 3 ml of top agar supplemented with 100 µl of bacterial overnight

culture, while for larger square Petri dishes (ca. 12 cm x 12 cm) we used 9 ml of top agar supplemented

with 200 µl of overnight culture. While the top agars were solidifying, serial dilutions of bacteriophage

stocks (previously grown on *E. coli* K-12 MG1655 ΔRM to erase any EcoKI methylation) were prepared

in sterile phosphate-buffered saline (PBS). Subsequently, each 2.5 µl of all serial dilutions were spotted on

all top agar plates and dried into the top agar before incubation at 37°C for at least 24 hours. Plaque

formation was recorded repeatedly throughout this time (starting after 3 hours of incubation for fast-

growing phages). The EOP of a given phage on a certain host was determined by calculating the ratio of

plaques obtained on this host over the number of plaques obtained on the reference strain *E. coli* K-12

MG1655 ΔRM carrying F(*pifA::zeoR*) and pBR322_ΔP*tet* [131].

When no plaque formation could be unambiguously recorded by visual inspection, the EOP was

determined to be below detection limit even if the top agar showed lysis from without (i.e., lysis zones

caused by bacterial cell death without phage infection at an efficiency high enough to form plaques [98]).

However, for all non-K12 strains of *E. coli* as well as *Salmonella* Typhimurium we determined the lysis

host range (i.e., the range of hosts on which lysis zones were observed) besides the numerical determination

of EOP (see above). Occasionally, we found that certain phage / host pairs were on the edge between merely

strong lysis from without and very poor plaque formation. Whenever in doubt, we recorded the result

conservatively as an EOP below detection limit (e.g., for phage FriedrichMiescher on *E. coli* 55989,

Enquatrovirus phages and EmilHeitz on *E. coli* UTI89, and all *Vequintavirinae* on the host expressing the

Fun/Z Abi system).

Bacteriophage genome sequencing and assembly

Genomic DNA of bacteriophages was prepared from high-titer stocks using the Norgen Biotek

Phage DNA Isolation Kit according to the manufacturer's guidelines and sequenced at the Microbial

Genome Sequencing Center (MiGS) using the Illumina NextSeq 550 platform. Trimmed sequencing reads

were assembled using the Geneious Assembler implemented in Geneious Prime 2021.0.1 with a coverage

of typically 50-100x (S5 Table). Usually, circular contigs (indicating a complete assembly due to the fusion

of characteristically repeated sequence at the genome ends [132]) were easily obtained using the "Medium

Sensitivity / Fast" setting. Consistently incomplete assemblies or local ambiguities were solved by PCR

amplification using the high-fidelity polymerase Phusion (NEB) followed by Sanger Sequencing. For

annotation and further analyses, sequences were linearized with the 5' end set either to the first position of

the small terminase subunit gene or the first position of the operon containing the small terminase subunit

gene.

Bacteriophage genome annotation

A preliminary, automated annotation of the genes in all genomes was generated using MultiPhate

[133] and then manually refined. For this purpose, whole-genome alignments of all new isolates within a

given group of phages and well-studied and / or well-annotated references were generated using

progressiveMauve [134] implemented in Geneious 2021.0.1 and used to inform the annotation based on

identified orthologs. *Bona fide* protein-coding genes without clear functional annotation were translated

and analyzed using the blastp tool on the NCBI server (https://blast.ncbi.nlm.nih.gov/Blast.cgi), the

InterPro protein domain signature database [135], as well as the Phyre2 fold recognition server [136;

http://www.sbg.bio.ic.ac.uk/~phyre2/html/page.cgi?id=index]. tRNA genes were predicted using tRNA-

ScanSE in the [137; http://lowelab.ucsc.edu/tRNAscan-SE/] and spanins were annotated with help of the

SpaninDataBase tool [138]. While endolysin genes were easily recognized by homology to lysozyme-like

proteins and other peptidoglycan hydrolases, holins were more difficult to identify when no holins were

annotated in closely related bacteriophage genomes. In these cases, we analyzed all small proteins (<250

amino acids) for the presence of transmembrane helices – a prerequisite for the functionality of known

holins [139] – and studied their possible relationships to previously described holins using blastp and

InterPro. In most but not all cases, *bona fide* holins encoded close to endolysin and / or spanin genes could

be identified. The annotation of our genomes in a comparative genomics setup made it easily possible to

identify the boundaries of introns associated with putative homing endonucleases and to precisely identify

inteins (see also S5 Table). The annotated genome sequences of all 66 newly isolated phages as well as

T3(K12) have been submitted to the NCBI GenBank database under accession numbers listed in S5 Table.

Bacteriophage naming and taxonomy

Newly isolated bacteriophages were named according to rules and conventions in the field and

classified in line with the rules of the International Committee on the Taxonomy of Viruses (ICTV) [79,

140] (S5 Table). As a first step of taxonomic classification, phages were roughly sorted by family and genus

based on whole-genome blastn searches against the non-redundant nucleotide collection database

[https://blast.ncbi.nlm.nih.gov/Blast.cgi; see also reference 141]. For each such broad taxonomic group, we

selected a set of reference sequences from NCBI GenBank that would cover the diversity of this group and

that always included all members which had already previously been studied intensively. Subsequently, we

generated whole-genome alignments of these sets of sequences using progressiveMauve implemented in

Geneious Prime 2021.0.1 [134]. These alignments or merely regions encoding highly conserved marker

genes such as the terminase subunits, portal protein gene, major capsid protein gene, or other suitable loci

were then used to generate Maximum-Likelihood phylogenies to unravel the evolutionary relationships

between the different bacteriophage isolates and database references (see S2 Text for details). Care was

taken to avoid genome regions that are recognizably infested with homing endonucleases or that showed

obvious signs of having recently moved by horizontal gene transfer (e.g., an abrupt shift in the local

sequence identity to the different other genomes in the alignment). Clusters of phage isolates observed in

these phylogenies generally correlated very well with established taxonomy as inferred from NCBI

taxonomy (https://www.ncbi.nlm.nih.gov/taxonomy), the ICTV [https://talk.ictvonline.org/taxonomy/;

reference 142], and other reports in the literature [30]. To classify our phage isolates on the species level,

we generated pairwise whole-genome alignments using progressiveMauve implemented in Geneious Prime

2021.0.1 [134] with genomes of related phages as identified in our phylogenies. From these alignments, the

nucleotide sequence identity was determined as the query coverage multiplied by the identity of aligned

segments [24]. When the genomes were largely syntenic and showed >95% nucleotide sequence identity,

we classified our isolates as the same species as their close relative [141], at least if this phage has been

assigned to a species by the ICTV [142].

Most phages were named in honor of scientists and other historically relevant personalities with

links to the city of Basel, Switzerland, where the majority of phages had been isolated. Despite efforts to

name many phages after female personalities, a considerable gender imbalance remains. This imbalance is

a consequence of the biases in how science and history have been made and recorded before the 20[th] century.

Sequence alignments and phylogenetic analyses

The NCBI GenBank accession numbers of all previously published genomes used in this study are

listed in S6 Table. Sequence alignments of different sets of homologous genes were generated using

MAFFT v7.450 implemented in Geneious Prime 2021.0.1 [143]. Whenever required, poor or missing

annotations in bacteriophage genomes downloaded from NCBI GenBank were supplemented using the

ORF finder tool of Geneious Prime 2021.0.1 guided by orthologous sequence parts of related genomes.

Alignments were set up using default settings typically with the fast FFT-NS-2 algorithm and 200PAM/k=2

or BLOSUM62 scoring matrices for nucleotide and amino acid sequences, respectively. Subsequently,

alignments were curated manually to improve poorly aligned sequence stretches and to mask non-

homologous parts.

For phylogenetic analyses, sequence alignments of orthologous stretches from different genomes

(genes, proteins, or whole genome) were used to calculate Maximum-Likelihood phylogenies with PhyML

3.3.20180621 implemented in Geneious Prime 2021.0.1 [144]. Phylogenies were calculated with the

HYK85 substitution model for nucleotide sequences and with the LG substitution model for amino acid

sequences. For the inference of phylogenetic relationships between phage genomes, we sometimes used

curated whole-genome alignments, but the infestation with homing endonucleases and the associated gene

conversion made this approach impossible for all larger genomes. Instead, we typically used curated

sequence alignments of several conserved core genes (on nucleotide or amino acid level, depending on the

distance between the genomes) as the basis for Maximum-Likelihood phylogenies. The detailed procedures

for each phylogeny shown in this article are described in S2 Text.

LPS structures of enterobacterial strains

Escherichia K-12 MG1655 (serogroup A) codes for a K12-type LPS core and an O16-type

O-antigen, but functional expression of the O-antigen was lost during laboratory adaptation due to

inactivation of *wbbL* by an IS5 insertion so that only a single terminal GlcNAc is attached to the LPS core

[44, 72]. *E. coli* strains UTI89 (serogroup B2), CFT073 (serogroup B2), and 55989 (serogroup B1) express

O-antigens of the O18, O6, and O104 types, respectively [122, 123]. Their core LPS types were determined

to be R1 (UTI89 and CFT073) and R3 (55989) by BLAST searches with diagnostic marker genes similar

to the PCR-based approached described previously [72]. For the illustration in Fig 12A, the structures of

LPS cores and linked O-antigen polysaccharides were drawn as described in the literature with typical O-

antigen chain lengths of 15 - 25 repeat units [72, 103, 145]. For the O18-type of O-antigen polysaccharides,

different subtypes with slight differences in the repeat unit have been described [103], but to the best of our

knowledge it has remained elusive which subtype is expressed by *E. coli* UTI89. In Fig 12A we therefore

chose an O18A type as exemplary O-antigen polysaccharide for *E. coli* UTI89. Similar to the laboratory

adaptation of *E. coli* K-12 MG1655, strains of the *E. coli* B lineage like *E. coli* B REL606 lack O-antigen

expression due to an IS1 insertion in *wbbD* but have an additional truncation in their R1-type LPS core due

to a second IS1 insertion in *waaT* that leaves only two glucoses in the outer core (Fig 12A) [67, 99].

Salmonella enterica subsp. *enterica* serovar Typhimurium strains 12023s (also known as ATCC 14028)

and SL1344 share the common *S.* Typhimurium LPS core and *Salmonella* serogroup B / O4 O-antigen [42,

146] (Fig 12A).

Bacterial genome sequencing and analyses to identify host receptors

While top agar assays with *E. coli* mutants carrying defined deletions in genes coding for different

surface proteins or genes involved in the LPS core biosynthesis readily identified the host receptor of most

bacteriophages (see above), few small siphoviruses of the *Drexlerviridae* family and the *Dhillonvirus*,

Nonagvirus, and *Seuratvirus* genera of the *Siphoviridae* family could not be assigned to any surface protein

as secondary receptor. This was the case for phages AugustePiccard (Bas01) and JeanPiccard (Bas02) of

Drexlerviridae, TheodorHerzl (Bas14) and Oekolampad (Bas18) of *Dhillonvirus*, ChristophMerian (Bas19)

and FritzHoffmann (Bas20) of *Nonagvirus*, and VogelGryff (Bas25) of *Seuratvirus*. However, as

siphoviruses infecting Gram-negative bacteria it seemed highly likely that these phages would use an outer

membrane porin as their secondary receptor [40]. We therefore isolated resistant mutants of *E. coli* K-12

BW25113 by plating bacteria on LB agar plates which had been densely covered with high-titer lysates of

each of these bacteriophages. While verifying that the resistance of isolated bacterial clones was specific to

the phage on which they had been isolated, we found that many clones were fully resistant specifically but

without exception to all these small siphoviruses with yet no known host receptor, suggesting that they all

targeted the same receptor.

For multiple of these phage-resistant clones, genomic DNA was prepared using the GenElute

Bacterial Genomic DNA Kit (Sigma-Aldrich) according to the manufacturer's guidelines and sequenced at

the Microbial Genome Sequencing Center (MiGS) using the Illumina NextSeq 550 platform. Sequencing

reads were assembled to the *E. coli* K-12 MG1655 reference genome (NCBI GenBank accession

U000096.3) using Geneious Prime 2021.0.1 with a coverage of typically 100-200x to identify the mutations

underlying their phage resistance. These analyses uncovered a diversity of in-frame deletions in *lptD* for

the different mutant clones (Fig 5).

Quantification and statistical analysis

Quantitative data sets were analyzed by calculating mean and standard deviation of at least three

independent biological replicates for each experiment. Detailed information about replicates and statistical

analyses for each experiment is provided in the figure legends.

Acknowledgments

1063

The authors are grateful to Prof. Urs Jenal, Prof. Marek Basler, Prof. Martin Loessner, and Dr. Julie

Sollier for valuable input and critical reading of the manuscript. The authors are deeply indebted to Prof.

Urs Jenal and Prof. Christoph Dehio for sharing multiple different *E. coli* strains and plasmids (S1 and S3

Tables). Prof. Calin Guet is acknowledged for sharing plasmids (S3 Table). The authors thank high school

students participating in the Biozentrum Summer Science Academy 2019, Julia Harms, and Dr. Thomas

Schubert for assistance with the isolation of multiple bacteriophages described in this study (S5 Table).

Salmonella Typhimurium strains 12023s and SL1344 were obtained from Prof. Dirk Bumann and Prof.

Mederic Diard, respectively, while *E. coli* B REL606 was obtained from Dr. Jenna Gallie. Phage P2*vir* was

generously shared by Dr. Nicolas Wenner and Prof. Jay Hinton. The Coli Genetic Stock Center (CGSC)

and the Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ) are acknowledged for

sharing different strains of *E. coli* or different bacteriophages, respectively (S1 and S5 Tables). Dr. Christian

Beisel and Dr. Geoffrey Fucile are acknowledged for advice regarding DNA sequencing. The authors are

grateful to ARA Basel, ARA Canius (Lenzerheide), and ARA Limeco (Dietikon) for providing samples of

sewage plant inflow and to Universitätsspital Basel for hospital sewage.

References

1.   Mahmoudabadi G, Phillips R. A comprehensive and quantitative exploration of thousands of viral genomes. Elife. 2018;7. doi: 10.7554/eLife.31955. PubMed PMID: 29624169; PubMed Central PMCID: PMCPMC5908442.

2.   Dion MB, Oechslin F, Moineau S. Phage diversity, genomics and phylogeny. Nat Rev Microbiol. 2020;18(3):125-38. Epub 2020/02/06. doi: 10.1038/s41579-019-0311-5. PubMed PMID: 32015529.

3.   Cobian Güemes AG, Youle M, Cantu VA, Felts B, Nulton J, Rohwer F. Viruses as Winners in the Game of Life. Annu Rev Virol. 2016;3(1):197-214. Epub 2016/10/16. doi: 10.1146/annurev-virology-100114-054952. PubMed PMID: 27741409.

4.   Keen EC. A century of phage research: bacteriophages and the shaping of modern biology. Bioessays. 2015;37(1):6-9. doi: 10.1002/bies.201400152. PubMed PMID: 25521633; PubMed Central PMCID: PMCPMC4418462.

5.   Demerec M, Fano U. Bacteriophage-Resistant Mutants in *Escherichia Coli*. Genetics. 1945;30(2):119-36. Epub 1945/03/01. PubMed PMID: 17247150; PubMed Central PMCID: PMCPMC1209279.

6.   Abedon ST. The murky origin of Snow White and her T-even dwarfs. Genetics. 2000;155(2):481-6. Epub 2000/06/03. PubMed PMID: 10835374; PubMed Central PMCID: PMCPMC1461100.

7.   Casjens SR, Hendrix RW. Bacteriophage lambda: Early pioneer and still relevant. Virology. 2015;479-480:310-30. doi: 10.1016/j.virol.2015.02.010. PubMed PMID: 25742714; PubMed Central PMCID: PMCPMC4424060.

8.   Howard-Varona C, Hargreaves KR, Abedon ST, Sullivan MB. Lysogeny in nature: mechanisms, impact and ecology of temperate phages. ISME J. 2017;11(7):1511-20. doi: 10.1038/ismej.2017.16. PubMed PMID: 28291233; PubMed Central PMCID: PMCPMC5520141.

9.   Hampton HG, Watson BNJ, Fineran PC. The arms race between bacteria and their phage foes. Nature. 2020;577(7790):327-36. Epub 2020/01/17. doi: 10.1038/s41586-019-1894-8. PubMed PMID: 31942051.

10.   Dy RL, Richter C, Salmond GP, Fineran PC. Remarkable Mechanisms in Microbes to Resist Phage Infections. Annu Rev Virol. 2014;1(1):307-31. doi: 10.1146/annurev-virology-031413-085500. PubMed PMID: 26958724.

11.   Bernheim A, Sorek R. The pan-immune system of bacteria: antiviral defence as a community resource. Nat Rev Microbiol. 2020;18(2):113-9. Epub 2019/11/07. doi: 10.1038/s41579-019-0278-2. PubMed PMID: 31695182.

12.   Lopatina A, Tal N, Sorek R. Abortive Infection: Bacterial Suicide as an Antiviral Immune Strategy. Annu Rev Virol. 2020;7(1):371-84. Epub 2020/06/20. doi: 10.1146/annurev-virology-011620-040628. PubMed PMID: 32559405.

13.   Schmidt C. Phage therapy's latest makeover. Nat Biotechnol. 2019;37(6):581-6. Epub 2019/05/10. doi: 10.1038/s41587-019-0133-z. PubMed PMID: 31068679.

14.   Kilcher S, Loessner MJ. Engineering Bacteriophages as Versatile Biologics. Trends Microbiol. 2019;27(4):355-67. Epub 2018/10/17. doi: 10.1016/j.tim.2018.09.006. PubMed PMID: 30322741.

15.   Millard A. Phage Genomes - Dec2020 2020 [16.02.2021]. Available from: http://millardlab.org/bioinformatics/bacteriophage-genomes/phage-genomes-dec2020/.

16.   Hyman P. Phages for Phage Therapy: Isolation, Characterization, and Host Range Breadth. Pharmaceuticals (Basel). 2019;12(1). Epub 2019/03/14. doi: 10.3390/ph12010035. PubMed PMID: 30862020; PubMed Central PMCID: PMCPMC6469166.

17.   Gordillo Altamirano FL, Barr JJ. Phage Therapy in the Postantibiotic Era. Clin Microbiol Rev.
2019;32(2). Epub 2019/01/18. doi: 10.1128/CMR.00066-18. PubMed PMID: 30651225; PubMed Central
PMCID: PMCPMC6431132.
18.   Kortright KE, Chan BK, Koff JL, Turner PE. Phage Therapy: A Renewed Approach to Combat
Antibiotic-Resistant Bacteria. Cell Host Microbe. 2019;25(2):219-32. Epub 2019/02/15. doi:
10.1016/j.chom.2019.01.014. PubMed PMID: 30763536.
19.   Aslam S, Lampley E, Wooten D, Karris M, Benson C, Strathdee S, et al. Lessons Learned From the
First 10 Consecutive Cases of Intravenous Bacteriophage Therapy to Treat Multidrug-Resistant Bacterial
Infections at a Single Center in the United States. Open Forum Infect Dis. 2020;7(9):ofaa389. Epub
2020/10/03. doi: 10.1093/ofid/ofaa389. PubMed PMID: 33005701; PubMed Central PMCID:
PMCPMC7519779.
20.   Pires DP, Costa AR, Pinto G, Meneses L, Azeredo J. Current challenges and future opportunities of
phage   therapy.   FEMS   Microbiol   Rev.   2020;44(6):684-700.   Epub   2020/05/31.   doi:
10.1093/femsre/fuaa017. PubMed PMID: 32472938.
21.   Ochman H, Selander RK. Standard reference strains of *Escherichia coli* from natural populations.
J Bacteriol. 1984;157(2):690-3. PubMed PMID: 6363394; PubMed Central PMCID: PMCPMC215307.
22.   Tenaillon O, Skurnik D, Picard B, Denamur E. The population genetics of commensal *Escherichia
coli*. Nat Rev Microbiol. 2010;8(3):207-17. doi: 10.1038/nrmicro2298. PubMed PMID: 20157339.
23.   Korf IHE, Meier-Kolthoff JP, Adriaenssens EM, Kropinski AM, Nimtz M, Rohde M, et al. Still
Something to Discover: Novel Insights into *Escherichia coli* Phage Diversity and Taxonomy. Viruses.
2019;11(5).   doi: 10.3390/v11050454.   PubMed   PMID: 31109012;   PubMed   Central   PMCID:
PMCPMC6563267.
24.   Olsen NS, Forero-Junco L, Kot W, Hansen LH. Exploring the Remarkable Diversity of Culturable
Escherichia coli Phages in the Danish Wastewater Environment. Viruses. 2020;12(9). Epub 2020/09/10.
doi: 10.3390/v12090986. PubMed PMID: 32899836; PubMed Central PMCID: PMCPMC7552041.
25.   Mathieu A, Dion M, Deng L, Tremblay D, Moncaut E, Shah SA, et al. Virulent coliphages in 1-year-
old children fecal samples are fewer, but more infectious than temperate coliphages. Nat Commun.
2020;11(1):378. Epub 2020/01/19. doi: 10.1038/s41467-019-14042-z. PubMed PMID: 31953385; PubMed
Central PMCID: PMCPMC6969025.
26.   Sørensen PE, Van Den Broeck W, Kiil K, Jasinskyte D, Moodley A, Garmyn A, et al. New insights
into the biodiversity of coliphages in the intestine of poultry. Sci Rep. 2020;10(1):15220. Epub 2020/09/18.
doi: 10.1038/s41598-020-72177-2.   PubMed   PMID: 32939020;   PubMed   Central   PMCID:
PMCPMC7494930.
27.   Michniewski S, Redgwell T, Grigonyte A, Rihtman B, Aguilo-Ferretjans M, Christie-Oleza J, et al.
Riding the wave of genomics to investigate aquatic coliphage diversity and activity. Environ Microbiol.
2019;21(6):2112-28. Epub 2019/03/19. doi: 10.1111/1462-2920.14590. PubMed PMID: 30884081;
PubMed Central PMCID: PMCPMC6563131.
28.   Smith R, O'Hara M, Hobman JL, Millard AD. Draft Genome Sequences of 14 *Escherichia coli* Phages
Isolated   from   Cattle   Slurry.   Genome   Announc.   2015;3(6).   Epub   2016/01/02.   doi:
10.1128/genomeA.01364-15. PubMed PMID: 26722010; PubMed Central PMCID: PMCPMC4698387.
29.   Pacifico C, Hilbert M, Sofka D, Dinhopl N, Pap IJ, Aspock C, et al. Natural Occurrence of *Escherichia
coli*-Infecting Bacteriophages in Clinical Samples. Front Microbiol. 2019;10:2484. Epub 2019/11/19. doi:
10.3389/fmicb.2019.02484. PubMed PMID: 31736918; PubMed Central PMCID: PMCPMC6834657.
30.   Grose JH, Casjens SR. Understanding the enormous diversity of bacteriophages: the tailed phages
that infect the bacterial family *Enterobacteriaceae*. Virology. 2014;468-470:421-43. Epub 2014/09/23. doi:
10.1016/j.virol.2014.08.024. PubMed PMID: 25240328; PubMed Central PMCID: PMCPMC4301999.

31.   Olsen NS, Hendriksen NB, Hansen LH, Kot W. A New High-Throughput Screening Method for
Phages: Enabling Crude Isolation and Fast Identification of Diverse Phages with Therapeutic Potential.
PHAGE. 2020;1(3):137-48. doi: 10.1089/phage.2020.0016.
32.   Drexler H. Bacteriophage T1. In: Calendar R, editor. The Bacteriophages. Boston, MA: Springer US;
1988. p. 235-58.
33.   Wittmann J, Turner D, Millard AD, Mahadevan P, Kropinski AM, Adriaenssens EM. From Orphan
Phage to a Proposed New Family-the Diversity of N4-Like Viruses. Antibiotics (Basel). 2020;9(10). Epub
2020/10/04. doi: 10.3390/antibiotics9100663. PubMed PMID: 33008130; PubMed Central PMCID:
PMCPMC7650795.
34.   Bertani G. Lysogeny at mid-twentieth century: P1, P2, and other experimental systems. J Bacteriol.
2004;186(3):595-600. Epub 2004/01/20. doi: 10.1128/jb.186.3.595-600.2004. PubMed PMID: 14729683;
PubMed Central PMCID: PMCPMC321500.
35.   Nobrega FL, Vlot M, de Jonge PA, Dreesens LL, Beaumont HJE, Lavigne R, et al. Targeting
mechanisms of tailed bacteriophages. Nat Rev Microbiol. 2018;16(12):760-73. Epub 2018/08/15. doi:
10.1038/s41579-018-0070-8. PubMed PMID: 30104690.
36.   Henderson JC, Zimmerman SM, Crofts AA, Boll JM, Kuhns LG, Herrera CM, et al. The Power of
Asymmetry: Architecture and Assembly of the Gram-Negative Outer Membrane Lipid Bilayer. Annu Rev
Microbiol. 2016;70:255-78. Epub 2016/07/01. doi: 10.1146/annurev-micro-102215-095308. PubMed
PMID: 27359214.
37.   Letarov AV, Kulikov EE. Adsorption of Bacteriophages on Bacterial Cells. Biochemistry (Mosc).
2017;82(13):1632-58. Epub 2018/03/11. doi: 10.1134/S0006297917130053. PubMed PMID: 29523063.
38.   Broeker NK, Barbirz S. Not a barrier but a key: How bacteriophages exploit host's O-antigen as an
essential receptor to initiate infection. Mol Microbiol. 2017;105(3):353-7. Epub 2017/06/16. doi:
10.1111/mmi.13729. PubMed PMID: 28618013.
39.   Bertozzi Silva J, Storms Z, Sauvageau D. Host receptors for bacteriophage adsorption. FEMS
Microbiol Lett. 2016;363(4). Epub 2016/01/13. doi: 10.1093/femsle/fnw002. PubMed PMID: 26755501.
40.   Hantke K. Compilation of Escherichia coli K-12 outer membrane phage receptors - their function
and   some   historical   remarks.   FEMS   Microbiol   Lett.   2020;367(2).   Epub   2020/02/06.   doi:
10.1093/femsle/fnaa013. PubMed PMID: 32009155.
41.   Gordillo Altamirano FL, Barr JJ. Unlocking the next generation of phage therapy: the key is in the
receptors. Curr Opin Biotechnol. 2020;68:115-23. Epub 2020/11/18. doi: 10.1016/j.copbio.2020.10.002.
PubMed PMID: 33202354.
42.   Bertani B, Ruiz N. Function and Biogenesis of Lipopolysaccharides. EcoSal Plus. 2018;8(1). Epub
2018/08/02. doi: 10.1128/ecosalplus.ESP-0001-2018. PubMed PMID: 30066669; PubMed Central PMCID:
PMCPMC6091223.
43.   Rai AK, Mitchell AM. Enterobacterial Common Antigen: Synthesis and Function of an Enigmatic
Molecule. mBio. 2020;11(4). Epub 2020/08/14. doi: 10.1128/mBio.01914-20. PubMed PMID: 32788387;
PubMed Central PMCID: PMCPMC7439462.
44.   Liu D, Reeves PR. *Escherichia* coli K12 regains its O antigen. Microbiology. 1994;140 (Pt 1):49-57.
doi: 10.1099/13500872-140-1-49. PubMed PMID: 7512872.
45.   Hutinet G, Kot W, Cui L, Hillebrand R, Balamkundu S, Gnanakalai S, et al. 7-Deazaguanine
modifications protect phage DNA from host restriction systems. Nat Commun. 2019;10(1):5442. Epub
2019/12/01. doi: 10.1038/s41467-019-13384-y. PubMed PMID: 31784519; PubMed Central PMCID:
PMCPMC6884629.
46.   Weigele P, Raleigh EA. Biosynthesis and Function of Modified Bases in Bacteria and Their Viruses.
Chem Rev. 2016;116(20):12655-87. Epub 2016/06/21. doi: 10.1021/acs.chemrev.6b00114. PubMed
PMID: 27319741.

47.   Loenen WA, Dryden DT, Raleigh EA, Wilson GG. Type I restriction enzymes and their relatives. Nucleic Acids Res. 2014;42(1):20-44. Epub 2013/09/27. doi: 10.1093/nar/gkt847. PubMed PMID: 24068554; PubMed Central PMCID: PMCPMC3874165.

48.   Pingoud A, Wilson GG, Wende W. Type II restriction endonucleases--a historical perspective and more. Nucleic Acids Res. 2014;42(12):7489-527. Epub 2014/06/01. doi: 10.1093/nar/gku447. PubMed PMID: 24878924; PubMed Central PMCID: PMCPMC4081073.

49.   Rao DN, Dryden DT, Bheemanaik S. Type III restriction-modification enzymes: a historical perspective. Nucleic Acids Res. 2014;42(1):45-55. Epub 2013/07/19. doi: 10.1093/nar/gkt616. PubMed PMID: 23863841; PubMed Central PMCID: PMCPMC3874151.

50.   Christie GE, Calendar R. Bacteriophage P2. Bacteriophage. 2016;6(1):e1145782. Epub 2016/05/05. doi: 10.1080/21597081.2016.1145782. PubMed PMID: 27144088; PubMed Central PMCID: PMCPMC4836473.

51.   Goulet A, Spinelli S, Mahony J, Cambillau C. Conserved and Diverse Traits of Adhesion Devices from *Siphoviridae* Recognizing Proteinaceous or Saccharidic Receptors. Viruses. 2020;12(5). Epub 2020/05/10. doi: 10.3390/v12050512. PubMed PMID: 32384698; PubMed Central PMCID: PMCPMC7291167.

52.   Piya D, Lessor L, Koehler B, Stonecipher A, Cahill J, Gill JJ. Genome-wide screens reveal *Escherichia coli* genes required for growth of T1-like phage LL5 and V5-like phage LL12. Sci Rep. 2020;10(1):8058. Epub 2020/05/18. doi: 10.1038/s41598-020-64981-7. PubMed PMID: 32415154; PubMed Central PMCID: PMCPMC7229145.

53.   Li P, Lin H, Mi Z, Xing S, Tong Y, Wang J. Screening of Polyvalent Phage-Resistant *Escherichia coli* Strains Based on Phage Receptor Analysis. Front Microbiol. 2019;10:850. Epub 2019/05/21. doi: 10.3389/fmicb.2019.00850. PubMed PMID: 31105661; PubMed Central PMCID: PMCPMC6499177.

54.   Wietzorrek A, Schwarz H, Herrmann C, Braun V. The genome of the novel phage Rtp, with a rosette-like tail tip, is homologous to the genome of phage T1. J Bacteriol. 2006;188(4):1419-36. Epub 2006/02/03. doi: 10.1128/JB.188.4.1419-1436.2006. PubMed PMID: 16452425; PubMed Central PMCID: PMCPMC1367250.

55.   Hamdi S, Rousseau GM, Labrie SJ, Tremblay DM, Kourda RS, Ben Slama K, et al. Characterization of two polyvalent phages infecting *Enterobacteriaceae*. Sci Rep. 2017;7:40349. Epub 2017/01/17. doi: 10.1038/srep40349. PubMed PMID: 28091598; PubMed Central PMCID: PMCPMC5238451.

56.   Wang J, Hofnung M, Charbit A. The C-terminal portion of the tail fiber protein of bacteriophage lambda is responsible for binding to LamB, its receptor at the surface of *Escherichia coli* K-12. J Bacteriol. 2000;182(2):508-12. Epub 2000/01/12. doi: 10.1128/jb.182.2.508-512.2000. PubMed PMID: 10629200; PubMed Central PMCID: PMCPMC94303.

57.   Meyer JR, Dobias DT, Weitz JS, Barrick JE, Quick RT, Lenski RE. Repeatability and contingency in the evolution of a key innovation in phage lambda. Science. 2012;335(6067):428-32. Epub 2012/01/28. doi: 10.1126/science.1214449. PubMed PMID: 22282803; PubMed Central PMCID: PMCPMC3306806.

58.   Dunne M, Rupf B, Tala M, Qabrati X, Ernst P, Shen Y, et al. Reprogramming Bacteriophage Host Range through Structure-Guided Design of Chimeric Receptor Binding Proteins. Cell Rep. 2019;29(5):1336-50 e4. Epub 2019/10/31. doi: 10.1016/j.celrep.2019.09.062. PubMed PMID: 31665644.

59.   Sazinas P, Redgwell T, Rihtman B, Grigonyte A, Michniewski S, Scanlan DJ, et al. Comparative Genomics of Bacteriophage of the Genus *Seuratvirus*. Genome Biol Evol. 2018;10(1):72-6. Epub 2017/12/23. doi: 10.1093/gbe/evx275. PubMed PMID: 29272407; PubMed Central PMCID: PMCPMC5758909.

60.   Golomidova AK, Kulikov EE, Prokhorov NS, Guerrero-Ferreira Rcapital Es C, Knirel YA, Kostryukova ES, et al. Branched Lateral Tail Fiber Organization in T5-Like Bacteriophages DT57C and DT571/2 is

Revealed by Genetic and Functional Analysis. Viruses. 2016;8(1). Epub 2016/01/26. doi: 10.3390/v8010026. PubMed PMID: 26805872; PubMed Central PMCID: PMCPMC4728585.

61.   Hong J, Kim KP, Heu S, Lee SJ, Adhya S, Ryu S. Identification of host receptor and receptor-binding module of a newly sequenced T5-like phage EPS7. FEMS Microbiol Lett. 2008;289(2):202-9. Epub 2008/11/26. doi: 10.1111/j.1574-6968.2008.01397.x. PubMed PMID: 19025561.

62.   Gencay YE, Gambino M, Prussing TF, Brondsted L. The genera of bacteriophages and their receptors are the major determinants of host range. Environ Microbiol. 2019;21(6):2095-111. Epub 2019/03/20. doi: 10.1111/1462-2920.14597. PubMed PMID: 30888719.

63.   Rabsch W, Ma L, Wiley G, Najar FZ, Kaserer W, Schuerch DW, et al. FepA- and TonB-dependent bacteriophage H8: receptor binding and genomic sequence. J Bacteriol. 2007;189(15):5658-74. Epub 2007/05/29. doi: 10.1128/JB.00437-07. PubMed PMID: 17526714; PubMed Central PMCID: PMCPMC1951831.

64.   Davison J. Pre-early functions of bacteriophage T5 and its relatives. Bacteriophage. 2015;5(4):e1086500. Epub 2016/02/24. doi: 10.1080/21597081.2015.1086500. PubMed PMID: 26904381; PubMed Central PMCID: PMCPMC4743489.

65.   Rusinov IS, Ershova AS, Karyagina AS, Spirin SA, Alexeevski AV. Avoidance of recognition sites of restriction-modification systems is a widespread but not universal anti-restriction strategy of prokaryotic viruses. BMC Genomics. 2018;19(1):885. Epub 2018/12/12. doi: 10.1186/s12864-018-5324-3. PubMed PMID: 30526500; PubMed Central PMCID: PMCPMC6286503.

66.   Trojet SN, Caumont-Sarcos A, Perrody E, Comeau AM, Krisch HM. The gp38 adhesins of the T4 superfamily: a complex modular determinant of the phage's host specificity. Genome Biol Evol. 2011;3:674-86. Epub 2011/07/13. doi: 10.1093/gbe/evr059. PubMed PMID: 21746838; PubMed Central PMCID: PMCPMC3157838.

67.   Washizaki A, Yonesaki T, Otsuka Y. Characterization of the interactions between *Escherichia coli* receptors, LPS and OmpC, and bacteriophage T4 long tail fibers. Microbiologyopen. 2016;5(6):1003-15. Epub 2016/06/09. doi: 10.1002/mbo3.384. PubMed PMID: 27273222; PubMed Central PMCID: PMCPMC5221442.

68.   Hu B, Margolin W, Molineux IJ, Liu J. Structural remodeling of bacteriophage T4 and host membranes during infection initiation. Proc Natl Acad Sci U S A. 2015;112(35):E4919-28. Epub 2015/08/19. doi: 10.1073/pnas.1501064112. PubMed PMID: 26283379; PubMed Central PMCID: PMCPMC4568249.

69.   Thomas JA, Orwenyo J, Wang LX, Black LW. The Odd "RB" Phage-Identification of Arabinosylation as a New Epigenetic Modification of DNA in T4-Like Phage RB69. Viruses. 2018;10(6). Epub 2018/06/13. doi: 10.3390/v10060313. PubMed PMID: 29890699; PubMed Central PMCID: PMCPMC6024577.

70.   Kropinski AM, Waddell T, Meng J, Franklin K, Ackermann HW, Ahmed R, et al. The host-range, genomics and proteomics of *Escherichia coli* O157:H7 bacteriophage rV5. Virol J. 2013;10:76. Epub 2013/03/19. doi: 10.1186/1743-422X-10-76. PubMed PMID: 23497209; PubMed Central PMCID: PMCPMC3606486.

71.   Schwarzer D, Buettner FF, Browning C, Nazarov S, Rabsch W, Bethe A, et al. A multivalent adsorption apparatus explains the broad host range of phage phi92: a comprehensive genomic and structural analysis. J Virol. 2012;86(19):10384-98. Epub 2012/07/13. doi: 10.1128/JVI.00801-12. PubMed PMID: 22787233; PubMed Central PMCID: PMCPMC3457257.

72.   Amor K, Heinrichs DE, Frirdich E, Ziebell K, Johnson RP, Whitfield C. Distribution of core oligosaccharide types in lipopolysaccharides from *Escherichia coli*. Infect Immun. 2000;68(3):1116-24. Epub 2000/02/26. doi: 10.1128/iai.68.3.1116-1124.2000. PubMed PMID: 10678915; PubMed Central PMCID: PMCPMC97256.

73.   Mutalik VK, Adler BA, Rishi HS, Piya D, Zhong C, Koskella B, et al. High-throughput mapping of the
phage resistance landscape in *E. coli*. PLoS Biol. 2020;18(10):e3000877. Epub 2020/10/14. doi:
10.1371/journal.pbio.3000877. PubMed PMID: 33048924; PubMed Central PMCID: PMCPMC7553319
following competing interests: VKM, AMD, and APA consult for and hold equity in Felix Biotechnology,
Inc.
74.   Qimron U, Marintcheva B, Tabor S, Richardson CC. Genomewide screens for *Escherichia coli* genes
affecting growth of T7 bacteriophage. Proc Natl Acad Sci U S A. 2006;103(50):19039-44. Epub 2006/12/01.
doi: 10.1073/pnas.0609428103. PubMed PMID: 17135349; PubMed Central PMCID: PMCPMC1748173.
75.   Gonzalez-Garcia VA, Pulido-Cid M, Garcia-Doval C, Bocanegra R, van Raaij MJ, Martin-Benito J, et
al. Conformational changes leading to T7 DNA delivery upon interaction with the bacterial receptor. J Biol
Chem. 2015;290(16):10038-44. Epub 2015/02/24. doi: 10.1074/jbc.M114.614222. PubMed PMID:
25697363; PubMed Central PMCID: PMCPMC4400320.
76.   Cuervo A, Fabrega-Ferrer M, Machon C, Conesa JJ, Fernandez FJ, Perez-Luque R, et al. Structures
of T7 bacteriophage portal and tail suggest a viral DNA retention and ejection mechanism. Nat Commun.
2019;10(1):3746. Epub 2019/08/23. doi: 10.1038/s41467-019-11705-9. PubMed PMID: 31431626;
PubMed Central PMCID: PMCPMC6702177.
77.   Hu B, Margolin W, Molineux IJ, Liu J. The bacteriophage t7 virion undergoes extensive structural
remodeling during infection. Science. 2013;339(6119):576-9. Epub 2013/01/12. doi:
10.1126/science.1231887. PubMed PMID: 23306440; PubMed Central PMCID: PMCPMC3873743.
78.   Scholl D, Merril C. The genome of bacteriophage K1F, a T7-like phage that has acquired the ability
to replicate on K1 strains of *Escherichia coli*. J Bacteriol. 2005;187(24):8499-503. Epub 2005/12/03. doi:
10.1128/JB.187.24.8499-8503.2005.   PubMed   PMID: 16321955;   PubMed   Central   PMCID:
PMCPMC1317022.
79.   Adriaenssens EM, Sullivan MB, Knezevic P, van Zyl LJ, Sarkar BL, Dutilh BE, et al. Taxonomy of
prokaryotic viruses: 2018-2019 update from the ICTV Bacterial and Archaeal Viruses Subcommittee. Arch
Virol. 2020;165(5):1253-60. Epub 2020/03/13. doi: 10.1007/s00705-020-04577-8. PubMed PMID:
32162068.
80.   Ando H, Lemire S, Pires DP, Lu TK. Engineering Modular Viral Scaffolds for Targeted Bacterial
Population Editing. Cell Syst. 2015;1(3):187-96. Epub 2016/03/15. doi: 10.1016/j.cels.2015.08.013.
PubMed PMID: 26973885; PubMed Central PMCID: PMCPMC4785837.
81.   Studier FW, Movva NR. SAMase gene of bacteriophage T3 is responsible for overcoming host
restriction. J Virol. 1976;19(1):136-45. Epub 1976/07/01. doi: 10.1128/JVI.19.1.136-145.1976. PubMed
PMID: 781304; PubMed Central PMCID: PMCPMC354840.
82.   Bandyopadhyay PK, Studier FW, Hamilton DL, Yuan R. Inhibition of the type I restriction-
modification enzymes EcoB and EcoK by the gene 0.3 protein of bacteriophage T7. J Mol Biol.
1985;182(4):567-78. Epub 1985/04/20. doi: 10.1016/0022-2836(85)90242-6. PubMed PMID: 2989534.
83.   Schmitt CK, Molineux IJ. Expression of gene 1.2 and gene 10 of bacteriophage T7 is lethal to F
plasmid-containing *Escherichia coli*. J Bacteriol. 1991;173(4):1536-43. Epub 1991/02/01. doi:
10.1128/jb.173.4.1536-1543.1991. PubMed PMID: 1995595; PubMed Central PMCID: PMCPMC207293.
84.   Molineux IJ, Spence JL. Virus-plasmid interactions: mutants of bacteriophage T3 that abortively
infect plasmid F-containing (F+) strains of *Escherichia coli*. Proc Natl Acad Sci U S A. 1984;81(5):1465-9.
Epub 1984/03/01. doi: 10.1073/pnas.81.5.1465. PubMed PMID: 6324192; PubMed Central PMCID:
PMCPMC344857.
85.   Kiino DR, Rothman-Denes LB. Genetic analysis of bacteriophage N4 adsorption. J Bacteriol.
1989;171(9):4595-602. Epub 1989/09/01. doi: 10.1128/jb.171.9.4595-4602.1989. PubMed PMID:
2670887; PubMed Central PMCID: PMCPMC210256.

86.   Kiino DR, Licudine R, Wilt K, Yang DH, Rothman-Denes LB. A cytoplasmic protein, NfrC, is required
for bacteriophage N4 adsorption. J Bacteriol. 1993;175(21):7074-80. Epub 1993/11/01. doi:
10.1128/jb.175.21.7074-7080.1993. PubMed PMID: 8226648; PubMed Central PMCID: PMCPMC206835.
87.   Prokhorov NS, Riccio C, Zdorovenko EL, Shneider MM, Browning C, Knirel YA, et al. Function of
bacteriophage G7C esterase tailspike in host cell adsorption. Mol Microbiol. 2017;105(3):385-98. Epub
2017/05/18. doi: 10.1111/mmi.13710. PubMed PMID: 28513100.
88.   McPartland J, Rothman-Denes LB. The tail sheath of bacteriophage N4 interacts with the
Escherichia coli receptor. J Bacteriol. 2009;191(2):525-32. Epub 2008/11/18. doi: 10.1128/JB.01423-08.
PubMed PMID: 19011026; PubMed Central PMCID: PMCPMC2620810.
89.   Shinedling S, Parma D, Gold L. Wild-type bacteriophage T4 is restricted by the lambda rex genes.
J Virol. 1987;61(12):3790-4. Epub 1987/12/01. doi: 10.1128/JVI.61.12.3790-3794.1987. PubMed PMID:
2960831; PubMed Central PMCID: PMCPMC255994.
90.   Whichard JM, Weigt LA, Borris DJ, Li LL, Zhang Q, Kapur V, et al. Complete genomic sequence of
bacteriophage felix o1. Viruses. 2010;2(3):710-30. Epub 2010/03/01. doi: 10.3390/v2030710. PubMed
PMID: 21994654; PubMed Central PMCID: PMCPMC3185647.
91.   Kaczorowska J, Casey E, Neve H, Franz C, Noben JP, Lugli GA, et al. A Quest of Great Importance-
Developing a Broad Spectrum *Escherichia coli* Phage Collection. Viruses. 2019;11(10). Epub 2019/09/29.
doi: 10.3390/v11100899. PubMed PMID: 31561510; PubMed Central PMCID: PMCPMC6832132.
92.   Simoliunas E, Vilkaityte M, Kaliniene L, Zajanckauskaite A, Kaupinis A, Staniulis J, et al. Incomplete
LPS Core-Specific Felix01-Like Virus vB_EcoM_VpaE1. Viruses. 2015;7(12):6163-81. Epub 2015/12/04. doi:
10.3390/v7122932. PubMed PMID: 26633460; PubMed Central PMCID: PMCPMC4690856.
93.   Blattner FR, Plunkett G, 3rd, Bloch CA, Perna NT, Burland V, Riley M, et al. The complete genome
sequence of *Escherichia coli* K-12. Science. 1997;277(5331):1453-62. Epub 1997/09/05. doi:
10.1126/science.277.5331.1453. PubMed PMID: 9278503.
94.   Hendrix RW, Duda RL. Bacteriophage lambda PaPa: not the mother of all lambda phages. Science.
1992;258(5085):1145-8. doi: 10.1126/science.1439823. PubMed PMID: 1439823.
95.   Sandulache R, Prehm P, Kamp D. Cell wall receptor for bacteriophage Mu G(+). J Bacteriol.
1984;160(1):299-303. Epub 1984/10/01. doi: 10.1128/JB.160.1.299-303.1984. PubMed PMID: 6384194;
PubMed Central PMCID: PMCPMC214716.
96.   North OI, Sakai K, Yamashita E, Nakagawa A, Iwazaki T, Buttner CR, et al. Phage tail fibre assembly
proteins employ a modular structure to drive the correct folding of diverse fibres. Nat Microbiol.
2019;4(10):1645-53. Epub 2019/06/19. doi: 10.1038/s41564-019-0477-7. PubMed PMID: 31209305.
97.   Hyman P, Abedon ST. Bacteriophage host range and bacterial resistance. Adv Appl Microbiol.
2010;70:217-48. Epub 2010/04/03. doi: 10.1016/s0065-2164(10)70007-1. PubMed PMID: 20359459.
98.   Abedon ST. Lysis from without. Bacteriophage. 2011;1(1):46-9. doi: 10.4161/bact.1.1.13980.
PubMed PMID: 21687534; PubMed Central PMCID: PMCPMC3109453.
99.   Jeong H, Barbe V, Lee CH, Vallenet D, Yu DS, Choi SH, et al. Genome sequences of *Escherichia coli*
B strains REL606 and BL21(DE3). J Mol Biol. 2009;394(4):644-52. Epub 2009/09/30. doi:
10.1016/j.jmb.2009.09.052. PubMed PMID: 19786035.
100.   Studier FW, Daegelen P, Lenski RE, Maslov S, Kim JF. Understanding the differences between
genome sequences of *Escherichia coli* B strains REL606 and BL21(DE3) and comparison of the *E. coli* B and
K-12 genomes. J Mol Biol. 2009;394(4):653-80. doi: 10.1016/j.jmb.2009.09.021. PubMed PMID:
19765592.
101.   Sueki A, Stein F, Savitski MM, Selkrig J, Typas A. Systematic Localization of *Escherichia coli*
Membrane Proteins. mSystems. 2020;5(2). Epub 2020/03/05. doi: 10.1128/mSystems.00808-19. PubMed
PMID: 32127419; PubMed Central PMCID: PMCPMC7055658.

102. Kortright KE, Chan BK, Turner PE. High-throughput discovery of phage receptors using transposon
insertion sequencing of bacteria. Proc Natl Acad Sci U S A. 2020;117(31):18670-9. Epub 2020/07/18. doi:
10.1073/pnas.2001888117. PubMed PMID: 32675236; PubMed Central PMCID: PMCPMC7414163.
103. Liu B, Furevi A, Perepelov AV, Guo X, Cao H, Wang Q, et al. Structure and genetics of *Escherichia*
coli O antigens. FEMS Microbiol Rev. 2020;44(6):655-83. Epub 2019/11/30. doi: 10.1093/femsre/fuz028.
PubMed PMID: 31778182; PubMed Central PMCID: PMCPMC7685785.
104. Gurney J, Brown SP, Kaltz O, Hochberg ME. Steering Phages to Combat Bacterial Pathogens.
Trends Microbiol. 2020;28(2):85-94. Epub 2019/11/21. doi: 10.1016/j.tim.2019.10.007. PubMed PMID:
31744662; PubMed Central PMCID: PMCPMC6980653.
105. Bauer RJ, Zhelkovsky A, Bilotti K, Crowell LE, Evans TC, Jr., McReynolds LA, et al. Comparative
analysis of the end-joining activity of several DNA ligases. PLoS One. 2017;12(12):e0190062. Epub
2017/12/29. doi: 10.1371/journal.pone.0190062. PubMed PMID: 29284038; PubMed Central PMCID:
PMCPMC5746248.
106. Millman A, Bernheim A, Stokar-Avihail A, Fedorenko T, Voichek M, Leavitt A, et al. Bacterial
Retrons Function In Anti-Phage Defense. Cell. 2020;183(6):1551-61 e12. Epub 2020/11/07. doi:
10.1016/j.cell.2020.09.065. PubMed PMID: 33157039.
107. Doron S, Melamed S, Ofir G, Leavitt A, Lopatina A, Keren M, et al. Systematic discovery of
antiphage defense systems in the microbial pangenome. Science. 2018;359(6379). Epub 2018/01/27. doi:
10.1126/science.aar4120. PubMed PMID: 29371424; PubMed Central PMCID: PMCPMC6387622.
108. Keen EC. Tradeoffs in bacteriophage life histories. Bacteriophage. 2014;4(1):e28365. Epub
2014/03/13. doi: 10.4161/bact.28365. PubMed PMID: 24616839; PubMed Central PMCID:
PMCPMC3942329.
109. Law R. Optimal Life Histories Under Age-Specific Predation. The American Naturalist.
1979;114(3):399-417.
110. Breitbart M, Bonnain C, Malki K, Sawaya NA. Phage puppet masters of the marine microbial realm.
Nat Microbiol. 2018;3(7):754-66. Epub 2018/06/06. doi: 10.1038/s41564-018-0166-y. PubMed PMID:
29867096.
111. Mizuno CM, Luong T, Cederstrom R, Krupovic M, Debarbieux L, Roach DR. Isolation and
Characterization of Bacteriophages That Infect *Citrobacter rodentium*, a Model Pathogen for Intestinal
Diseases. Viruses. 2020;12(7). Epub 2020/07/12. doi: 10.3390/v12070737. PubMed PMID: 32650458;
PubMed Central PMCID: PMCPMC7412075.
112. Jordan TC, Burnett SH, Carson S, Caruso SM, Clase K, DeJong RJ, et al. A broadly implementable
research course in phage discovery and genomics for first-year undergraduate students. mBio.
2014;5(1):e01051-13. Epub 2014/02/06. doi: 10.1128/mBio.01051-13. PubMed PMID: 24496795;
PubMed Central PMCID: PMCPMC3950523.
113. M9 minimal medium (standard). Cold Spring Harbor Protocols. 2010;2010(8):pdb.rec12295. doi:
10.1101/pdb.rec12295.
114. Kropinski AM, Mazzocco A, Waddell TE, Lingohr E, Johnson RP. Enumeration of bacteriophages by
double agar overlay plaque assay. Methods Mol Biol. 2009;501:69-76. doi: 10.1007/978-1-60327-164-6_7.
PubMed PMID: 19066811.
115. Kauffman KM, Polz MF. Streamlining standard bacteriophage methods for higher throughput.
MethodsX. 2018;5:159-72. Epub 2019/01/10. doi: 10.1016/j.mex.2018.01.007. PubMed PMID: 30622914;
PubMed Central PMCID: PMCPMC6318102.
116. Golec P, Dabrowski K, Hejnowicz MS, Gozdek A, Los JM, Wegrzyn G, et al. A reliable method for
storage of tailed phages. J Microbiol Methods. 2011;84(3):486-9. Epub 2011/01/25. doi:
10.1016/j.mimet.2011.01.007. PubMed PMID: 21256885.

117. Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in *Escherichia coli* K-12
using PCR products. Proc Natl Acad Sci USA. 2000;97(12):6640-5. doi: 10.1073/pnas.120163297. PubMed
PMID: 10829079; PubMed Central PMCID: PMC18686.

118. Harms A, Fino C, Sørensen MA, Semsey S, Gerdes K. Prophages and Growth Dynamics Confound
Experimental Results with Antibiotic-Tolerant Persister Cells. MBio. 2017;8(6). doi: 10.1128/mBio.01964-
17. PubMed PMID: 29233898; PubMed Central PMCID: PMCPMC5727415.

119. Pleska M, Qian L, Okura R, Bergmiller T, Wakamoto Y, Kussell E, et al. Bacterial Autoimmunity Due
to a Restriction-Modification System. Curr Biol. 2016;26(3):404-9. Epub 2016/01/26. doi:
10.1016/j.cub.2015.12.041. PubMed PMID: 26804559.

120. Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al. Construction of *Escherichia coli* K-
12 in-frame, single-gene knockout mutants: the Keio collection. Mol Syst Biol. 2006;2:2006 0008. doi:
10.1038/msb4100050. PubMed PMID: 16738554; PubMed Central PMCID: PMC1681482.

121. Fino C, Vestergaard M, Ingmer H, Pierrel F, Gerdes K, Harms A. PasT of *Escherichia coli* sustains
antibiotic tolerance and aerobic respiration as a bacterial homolog of mitochondrial Coq10.
Microbiologyopen. 2020;9(8):e1064. Epub 2020/06/20. doi: 10.1002/mbo3.1064. PubMed PMID:
32558363; PubMed Central PMCID: PMCPMC7424257.

122. Wiles TJ, Kulesus RR, Mulvey MA. Origins and virulence mechanisms of uropathogenic *Escherichia
coli*. Exp Mol Pathol. 2008;85(1):11-9. Epub 2008/05/17. doi: 10.1016/j.yexmp.2008.03.007. PubMed
PMID: 18482721; PubMed Central PMCID: PMCPMC2595135.

123. Grad YH, Godfrey P, Cerquiera GC, Mariani-Kurkdjian P, Gouali M, Bingen E, et al. Comparative
genomics of recent Shiga toxin-producing *Escherichia coli* O104:H4: short-term evolution of an emerging
pathogen. mBio. 2013;4(1):e00452-12. Epub 2013/01/24. doi: 10.1128/mBio.00452-12. PubMed PMID:
23341549; PubMed Central PMCID: PMCPMC3551546.

124. Bernier C, Gounon P, Le Bouguenec C. Identification of an aggregative adhesion fimbria (AAF) type
III-encoding operon in enteroaggregative *Escherichia coli* as a sensitive probe for detecting the AAF-
encoding operon family. Infect Immun. 2002;70(8):4302-11. Epub 2002/07/16. doi:
10.1128/iai.70.8.4302-4311.2002. PubMed PMID: 12117939; PubMed Central PMCID: PMCPMC128174.

125. Branchu P, Bawn M, Kingsley RA. Genome Variation and Molecular Epidemiology of *Salmonella
enterica* Serovar Typhimurium Pathovariants. Infect Immun. 2018;86(8). Epub 2018/05/23. doi:
10.1128/IAI.00079-18. PubMed PMID: 29784861; PubMed Central PMCID: PMCPMC6056856.

126. Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, 3rd, Smith HO. Enzymatic assembly of
DNA molecules up to several hundred kilobases. Nat Methods. 2009;6(5):343-5. Epub 2009/04/14. doi:
10.1038/nmeth.1318. PubMed PMID: 19363495.

127. Liu H, Naismith JH. An efficient one-step site-directed deletion, insertion, single and multiple-site
plasmid mutagenesis protocol. BMC Biotechnol. 2008;8:91. doi: 10.1186/1472-6750-8-91. PubMed PMID:
19055817; PubMed Central PMCID: PMC2629768.

128. Roberts RJ, Vincze T, Posfai J, Macelis D. REBASE--a database for DNA restriction and modification:
enzymes, genes and genomes. Nucleic Acids Res. 2015;43(Database issue):D298-9. Epub 2014/11/08. doi:
10.1093/nar/gku1046. PubMed PMID: 25378308; PubMed Central PMCID: PMCPMC4383893.

129. Czajkowski R, Ozymko Z, Lojkowska E. Application of zinc chloride precipitation method for rapid
isolation and concentration of infectious *Pectobacterium* spp. and *Dickeya* spp. lytic bacteriophages from
surface water and plant and soil extracts. Folia Microbiol (Praha). 2016;61(1):29-33. Epub 2015/06/24.
doi: 10.1007/s12223-015-0411-1. PubMed PMID: 26099750; PubMed Central PMCID: PMCPMC4691450.

130. You L, Suthers PF, Yin J. Effects of *Escherichia coli* physiology on growth of phage T7 *in vivo* and *in
silico*. J Bacteriol. 2002;184(7):1888-94. Epub 2002/03/13. doi: 10.1128/jb.184.7.1888-1894.2002.
PubMed PMID: 11889095; PubMed Central PMCID: PMCPMC134924.

131.   Abedon ST, Katsaounis TI. Basic Phage Mathematics. Methods Mol Biol. 2018;1681:3-30. Epub
2017/11/15. doi: 10.1007/978-1-4939-7343-9_1. PubMed PMID: 29134583.
132.   Garneau JR, Depardieu F, Fortier LC, Bikard D, Monot M. PhageTerm: a tool for fast and accurate
determination of phage termini and packaging mechanism using next-generation sequencing data. Sci
Rep. 2017;7(1):8292. Epub 2017/08/16. doi: 10.1038/s41598-017-07910-5. PubMed PMID: 28811656;
PubMed Central PMCID: PMCPMC5557969.
133.   Ecale Zhou CL, Malfatti S, Kimbrel J, Philipson C, McNair K, Hamilton T, et al. multiPhATE:
bioinformatics pipeline for functional annotation of phage isolates. Bioinformatics. 2019;35(21):4402-4.
Epub 2019/05/16. doi: 10.1093/bioinformatics/btz258. PubMed PMID: 31086982; PubMed Central
PMCID: PMCPMC6821344.
134.   Darling AE, Mau B, Perna NT. progressiveMauve: multiple genome alignment with gene gain, loss
and rearrangement. PLoS One. 2010;5(6):e11147. Epub 2010/07/02. doi: 10.1371/journal.pone.0011147.
PubMed PMID: 20593022; PubMed Central PMCID: PMCPMC2892488.
135.   Mitchell AL, Attwood TK, Babbitt PC, Blum M, Bork P, Bridge A, et al. InterPro in 2019: improving
coverage, classification and access to protein sequence annotations. Nucleic Acids Res.
2019;47(D1):D351-D60. Epub 2018/11/07. doi: 10.1093/nar/gky1100. PubMed PMID: 30398656; PubMed
Central PMCID: PMCPMC6323941.
136.   Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ. The Phyre2 web portal for protein
modeling, prediction and analysis. Nat Protoc. 2015;10(6):845-58. doi: 10.1038/nprot.2015.053. PubMed
PMID: 25950237; PubMed Central PMCID: PMCPMC5298202.
137.   Chan PP, Lowe TM. tRNAscan-SE: Searching for tRNA Genes in Genomic Sequences. Methods Mol
Biol. 2019;1962:1-14. Epub 2019/04/26. doi: 10.1007/978-1-4939-9173-0_1. PubMed PMID: 31020551;
PubMed Central PMCID: PMCPMC6768409.
138.   Kongari R, Rajaure M, Cahill J, Rasche E, Mijalis E, Berry J, et al. Phage spanins: diversity,
topological dynamics and gene convergence. BMC Bioinformatics. 2018;19(1):326. Epub 2018/09/17. doi:
10.1186/s12859-018-2342-8. PubMed PMID: 30219026; PubMed Central PMCID: PMCPMC6139136.
139.   Cahill J, Young R. Phage Lysis: Multiple Genes for Multiple Barriers. Adv Virus Res. 2019;103:33-
70. Epub 2019/01/13. doi: 10.1016/bs.aivir.2018.09.003. PubMed PMID: 30635077; PubMed Central
PMCID: PMCPMC6733033.
140.   Adriaenssens E, Brister JR. How to Name and Classify Your Phage: An Informal Guide. Viruses.
2017;9(4). doi: 10.3390/v9040070. PubMed PMID: 28368359; PubMed Central PMCID: PMCPMC5408676.
141.   Tolstoy I, Kropinski AM, Brister JR. Bacteriophage Taxonomy: An Evolving Discipline. Methods Mol
Biol. 2018;1693:57-71. Epub 2017/11/10. doi: 10.1007/978-1-4939-7395-8_6. PubMed PMID: 29119432.
142.   Lefkowitz EJ, Dempsey DM, Hendrickson RC, Orton RJ, Siddell SG, Smith DB. Virus taxonomy: the
database of the International Committee on Taxonomy of Viruses (ICTV). Nucleic Acids Res.
2018;46(D1):D708-D17. Epub 2017/10/19. doi: 10.1093/nar/gkx932. PubMed PMID: 29040670; PubMed
Central PMCID: PMCPMC5753373.
143.   Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in
performance and usability. Mol Biol Evol. 2013;30(4):772-80. Epub 2013/01/19. doi:
10.1093/molbev/mst010. PubMed PMID: 23329690; PubMed Central PMCID: PMCPMC3603318.
144.   Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms and
methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. Syst Biol.
2010;59(3):307-21. Epub 2010/06/09. doi: 10.1093/sysbio/syq010. PubMed PMID: 20525638.
145.   Heinrichs DE, Yethon JA, Whitfield C. Molecular basis for structural diversity in the core regions of
the lipopolysaccharides of *Escherichia coli* and *Salmonella enterica*. Mol Microbiol. 1998;30(2):221-32.
Epub 1998/10/29. doi: 10.1046/j.1365-2958.1998.01063.x. PubMed PMID: 9791168.

146.   Micoli F, Ravenscroft N, Cescutti P, Stefanetti G, Londero S, Rondini S, et al. Structural analysis of
O-polysaccharide chains extracted from different *Salmonella* Typhimurium strains. Carbohydr Res.
2014;385:1-8. Epub 2014/01/05. doi: 10.1016/j.carres.2013.12.003. PubMed PMID: 24384528.
147.   Storek KM, Chan J, Vij R, Chiang N, Lin Z, Bevers J, 3rd, et al. Massive antibody discovery used to
probe structure-function relationships of the essential outer membrane protein LptD. Elife. 2019;8. Epub
2019/06/27.   doi: 10.7554/eLife.46258.   PubMed   PMID: 31237236;   PubMed   Central   PMCID:
PMCPMC6592684.
148.   Eriksson JM, Haggard-Ljungquist E. The multifunctional bacteriophage P2 cox protein requires
oligomerization   for   biological   activity.   J Bacteriol.   2000;182(23):6714-23.   Epub   2000/11/14.   doi:
10.1128/jb.182.23.6714-6723.2000.   PubMed   PMID: 11073917;   PubMed   Central   PMCID:
PMCPMC111415.
149.   Silhavy   TJ.   Gene   fusions.   J   Bacteriol.   2000;182(21):5935-8.   Epub   2000/10/13.   doi:
10.1128/jb.182.21.5935-5938.2000. PubMed PMID: 11029410; PubMed Central PMCID: PMCPMC94724.
150.   Crick FH, Barnett L, Brenner S, Watts-Tobin RJ. General nature of the genetic code for proteins.
Nature. 1961;192:1227-32. Epub 1961/12/30. doi: 10.1038/1921227a0. PubMed PMID: 13882203.
151.   Benzer S, Champe SP. A change from nonsense to sense in the genetic code. Proc Natl Acad Sci U
S   A.   1962;48:1114-21.   Epub   1962/07/15.   doi: 10.1073/pnas.48.7.1114.   PubMed   PMID: 13867417;
PubMed Central PMCID: PMCPMC220916.
1555

Figures

Fig 1. The three morphotypes of *Caudovirales* and two lines of defense in bacterial immunity.

(A) The virions of tailed phages or *Caudovirales* can be assigned to three general morphotypes including myoviruses (contractile tail), siphoviruses (long and flexible, non-contractile tail), and podoviruses (short and stubby tail). **(B)** The life cycle of a typical lytic phage begins with reversible attachment to a so-called primary receptor on the bacterial cell surface (1), usually via lateral tail fibers at the virion. Subsequently, irreversible attachment to secondary or terminal receptors usually depends on structures at the end of the tail, e.g., short tail fibers for many myoviruses and central tail fibers or tail tip proteins for siphoviruses (2 ; see also (A)). After genome injection (3), the phage takes over the host cell, replicates, and releases the offspring by host cell lysis (4). Inside the host cell, the bacteriophage faces two lines of host defenses, first bacterial immunity systems that try to clear the infection by directly targeting the phage genome (5) and then abortive infection systems that kill the infected cell when a viral infection is sensed (6).

Fig 2. Overview of the BASEL collection.

(A) Illustration of the workflow of bacteriophage isolation, characterization, and selection that resulted in the BASEL collection (details in *Materials and Methods*). **(B)** Taxonomic overview of the bacteriophages included in the BASEL collection and their unique Bas## identifiers. Newly isolated phages are colored according to their family while well-studied reference phages are shown in grey.

Fig 3. Overview of *E. coli* surface glycan variants and the immunity systems used in this study.

(A) The surface glycans of different *E. coli* K-12 MG1655 variants are shown schematically (details in running text and *Materials and Methods*). Note that the *E. coli* K-12 MG1655 laboratory wildtype does not merely display the K12-type core LPS (classical rough LPS phenotype) but also the most proximal

D-glucose of the O16-type O-antigen. **(B)** Key features of the six RM systems (each two of type I, type II,

and type III) and the five Abi systems used for the phenotyping of this study are summarized schematically.

Recognition sites of RM systems have either been determined experimentally or were predicted in REBASE

(red nucleotides: methylation sites; dotted lines: cleavage sites) [47-49, 128]. The Abi systems have been

characterized to very different extent but constitute the most well-understood representatives of these

immunity systems of *E. coli* [10, 50].

Fig 4. Overview of *Drexlerviridae* phages.

(A) Schematic illustration of host recognition by *Drexlerviridae*. **(B)** Maximum-Likelihood

phylogeny of *Drexlerviridae* based on several core genes with bootstrap support of branches shown if >

70/100. Newly isolated phages of the BASEL collection are highlighted by red phage icons and the

determined or proposed terminal receptor specificity is highlighted at the phage names using the color code

highlighted in (C). The phylogeny was rooted based on a representative phylogeny including *Dhillonvirus*

sequences as outgroup (S1A Fig). **(C)** On the left, the seven identified receptors of small siphoviruses are

shown with a color code that is also used to annotate demonstrated or predicted receptor specificity in the

phylogenies of Fig 4B and Figs 6A + 6C). On the right, we show representative *bona fide* RBP loci that

seem to encode the receptor specificity of these small siphoviruses (with the same color code). Note that

the loci linked to each receptor are very similar while the genetic arrangement differs considerably between

loci linked to different terminal host receptors (see also S1C Fig). **(D)** The results of quantitative

phenotyping experiments with *Drexlerviridae* phages regarding sensitivity to altered surface glycans and

bacterial immunity systems are presented as efficiency of plating (EOP). Data points and error bars

represent average and standard deviation of at least three independent experiments.

Fig 5. LptD is a commonly targeted terminal receptor of small siphoviruses.

1602 **(A)** Whole-genome sequencing of bacterial mutants exhibiting spontaneous resistance to seven

1603 small siphoviruses with no previously known receptor revealed different mutations or small deletions in the

1604 essential gene *lptD* that encodes the LptD LPS export channel. Top agar assays with two representative

mutants in comparison to the ancestral *E. coli* K-12 BW25113 strain were performed with serial tenfold dilutions of twelve different phages (undiluted high-titer stocks at the bottom and increasingly diluted samples towards the top). Both mutants display complete resistance to the seven small siphoviruses of diverse genera within *Drexlerviridae* and *Siphoviridae* families that share the same *bona fide* RBP modules (S1C Fig) while no other phage of the BASEL collection was affected. In particular, we excluded indirect effects, e.g., via changes in the LPS composition in the *lptD* mutants, by confirming that five LPS-targeting phages of diverse families (see below) showed full infectivity on all strains. **(B)** The amino acid sequence alignment of wildtype LptD with the two mutants highlighted in (A) shows that resistance to LptD-targeting phages is linked to small deletions in or adjacent to regions encoding extracellular loops as defined in previous work [147], suggesting that they abolish the RBP-receptor interaction.

Fig 6. Overview of *Siphovirida*e genera *Dhillonvirus*, *Nonagvirus*, and *Seuratvirus*.

      (A) Schematic illustration of host recognition by small siphoviruses. **(B)** Maximum-Likelihood phylogeny of the *Dhillonvirus* genus based on a whole-genome alignment with bootstrap support of branches shown if > 70/100. Newly isolated phages of the BASEL collection are highlighted by red phage icons and the determined or proposed terminal receptor specificity is highlighted at the phage names using the color code highlighted in Fig 4C. The phylogeny was rooted between phage WFI and all others based on a representative phylogeny including *Drexlerviridae* sequences as outgroup (S1A Fig). **(C)** The results of quantitative phenotyping experiments with *Dhillonvirus* phages regarding sensitivity to altered surface glycans and bacterial immunity systems are presented as efficiency of plating (EOP). **(D)** Maximum-Likelihood phylogeny of the *Nonagvirus* and *Seuratvirus* genera based on a whole-genome alignment with bootstrap support of branches shown if > 70/100. Newly isolated phages of the BASEL collection are highlighted by red phage icons and the determined or proposed terminal receptor specificity is highlighted at the phage names using the color code highlighted in Fig 4C. The phylogeny was rooted between the two genera. **(E)** *Nonagvirus* and *Seuratvirus* phages share a core 7-deazaguanosine biosynthesis pathway involving FolE, QueD, QueE, and QueC which synthesizes $dPreQ_0$ that is inserted into their genomes by

DpdA. In *Nonagvirus* phages, the fusion of QueC with a glutamate amidotransferase (Gat) domain to Gat-

QueC results in the modification with dG^+ instead of $dPreQ_0$ [45]. **(F)** The results of quantitative

phenotyping experiments with *Nonagvirus* and *Seuratvirus* phages regarding sensitivity to altered surface

glycans and bacterial immunity systems are presented as efficiency of plating (EOP). In (C) and (F), data

points and error bars represent average and standard deviation of at least three independent experiments.

Fig 7. Overview of *Demerecviridae* subfamily *Markadamsvirinae*.

(A) Schematic illustration of host recognition by T5-like siphoviruses. **(B)** Maximum-Likelihood

phylogeny of the *Markadamsvirinae* subfamily of *Demerecviridae* based on several core genes with

bootstrap support of branches shown if > 70/100. Phages of the BASEL collection are highlighted by little

phage icons and the determined or proposed terminal receptor specificity is highlighted at the phage names

using the color code highlighted at the right side (same as for the small siphoviruses). The phylogeny was

rooted between the *Epseptimavirus* and *Tequintavirus* genera. **(C)** The results of quantitative phenotyping

experiments with *Markadamsvirinae* phages regarding sensitivity to altered surface glycans and bacterial

immunity systems are presented as efficiency of plating (EOP). Data points and error bars represent average

and standard deviation of at least three independent experiments.

Fig 8. Overview of the *Myoviridae* subfamily *Tevenvirinae*.

(A) Schematic illustration of host recognition by T4-like myoviruses. **(B)** Maximum-Likelihood

phylogeny of the *Tevenvirinae* subfamily of *Myoviridae* based on a curated whole-genome alignment with

bootstrap support of branches shown if > 70/100. The phylogeny was rooted between the *Tequatrovirus*

and *Mosigvirus* genera. Phages of the BASEL collection are highlighted by little phage icons and

experimentally determined primary receptor specificity is highlighted at the phage names using the color

code highlighted at the top left. Primary receptor specificity of *Tevenvirinae* depends on RBPs expressed

either as a C-terminal extension of the distal half fiber (T4 and other OmpC-targeting phages) or as separate

small fiber tip adhesins [66], but sequence analyses of the latter remained ambiguous. We therefore only

annotated experimentally determined primary receptors (see also S4A Fig) [53, 66]. **(C)** The results of

quantitative phenotyping experiments with *Tevenvirinae* phages regarding sensitivity to altered surface glycans and bacterial immunity systems are presented as efficiency of plating (EOP). Data points and error bars represent average and standard deviation of at least three independent experiments. **(D)** The Maximum-Likelihood phylogeny *Tevenvirinae* short tail fiber proteins reveals two homologous, yet clearly distinct, clusters that correlate with the absence (variant #1, like T4) or presence (variant #2) of detectable LPS core dependence as shown in (C). **(E)** The results of (D) indicate that variant #1, as shown for T4, binds the deep lipid A – Kdo region of the enterobacterial LPS core, while variant #2 binds a more distal part of the (probably inner) core.

Fig 9. Overview of the *Myoviridae* subfamily *Vequintavirinae* and relatives.

(A) Schematic illustration of host recognition by *Vequintavirinae* and related myoviruses. **(B)** Maximum-Likelihood phylogeny of the *Vequintavirinae* subfamily of *Myoviridae* and relatives based on a curated whole-genome alignment with bootstrap support of branches shown if > 70/100. The phylogeny was rooted between the *Vequintavirus* genus and the two closely related, unclassified groups at the bottom. Newly isolated phages of the BASEL collection are highlighted by green phage icons. **(C)** The results of quantitative phenotyping experiments with *Vequintavirinae* and phage PaulScherrer regarding sensitivity to altered surface glycans and bacterial immunity systems are presented as efficiency of plating (EOP). Data points and error bars represent average and standard deviation of at least three independent experiments. **(D)** Amino acid sequence alignment of the lateral tail fiber Gp64 of phage N4 (*Enquatrovirus*, see below) and a lateral tail fiber conserved among *Vequintavirinae* and relatives (representatives shown). The proteins share a predicted poly-GlcNAc deacetylase domain as identified by Phyre2 [136]. **(E)** *Vequintavirinae sensu stricto* (represented by rV5 and Jeff Schatz) encode two paralogous short tail fiber proteins and a tail fiber chaperone that are homologous to the corresponding locus in phi92-like phages incl. PaulScherrer and, ultimately, to short tail fiber GpS and chaperone GpU of Mu(+) (which targets a different glucose in the K12-type LPS core GpS [95, 96]).

Fig 10. Overview of *Autographiviridae* phages and *Podoviridae* genus *Enquatrovirus*.

(A) Schematic illustration of host recognition by *Autographiviridae* and *Enquatrovirus* phages. **(B)** Maximum-Likelihood phylogeny of the *Studiervirinae* subfamily of *Autographiviridae* based on several core genes with bootstrap support of branches shown if > 70/100. The phylogeny was midpoint-rooted between the clade formed by *Teseptimavirus* and *Teetrevirus* and the other genera. Phages of the BASEL collection are highlighted by little phage icons. **(C)** The results of quantitative phenotyping experiments with *Autographiviridae* regarding sensitivity to altered surface glycans and bacterial immunity systems are presented as efficiency of plating (EOP). Data points and error bars represent average and standard deviation of at least three independent experiments. **(D)** Amino acid sequence alignment and Maximum-Likelihood phylogeny of Gp1.2 orthologs in all tested *Autographiviridae* phages. Phage JeanTinguely belongs to the *Teseptimavirus* genus but encodes an allele of *gp1.2* that is closely related to those of the *Berlinvirus* genus, possibly explaining its resistance to PifA (see (C)). **(E)** Maximum-Likelihood phylogeny of the *Enquatrovirus* genus and related groups of *Podoviridae* based on several core genes with bootstrap support of branches shown if > 70/100. The phylogeny was midpoint-rooted between the distantly related *Jwalphavirus* genus and the others. Phages of the BASEL collection are highlighted by little phage icons.

Fig 11. Overview of *Myoviridae: Ounavirinae* and classic temperate phages.

(A) Schematic illustration of host recognition by *Ounavirinae: Felixounavirus* phages. Note that the illustration shows short tail fibers simply in analogy to *Tevenvirinae* or *Vequintavirinae* (Figs 8A and 9A), but any role for such structures has not been explored for Felix O1 and relatives. **(B)** Maximum-Likelihood phylogeny of the *Ounavirinae* subfamily of *Myoiviridae* based on several core genes with bootstrap support of branches shown if > 70/100. The phylogeny was midpoint-rooted between *Kolesnikvirus* and the other genera. Our new isolate JohannRWettstein is highlighted by a green phage icon. **(C,D)** Schematic illustration of host recognition by classic temperate phages lambda, P1, and P2. Note the absence of lateral tail fibers due to a mutation in lambda PaPa laboratory strains [94]. **(E,F)** The results of quantitative phenotyping experiments with JohannRWettstein and classic temperate phages regarding

sensitivity to altered surface glycans and bacterial immunity systems are presented as efficiency of plating

(EOP). Data points and error bars represent average and standard deviation of at least three independent

experiments.

Fig 12. Host range of phages in the BASEL collection.

(A) Surface glycans of the enterobacterial strains used in this work (see *Materials and Methods* for

details on how the illustration of glycan chains was composed). **(B)** The ability of all phages in the BASEL

collection to infect different enterobacteria was studied qualitatively (lysis host range; top) and, more

stringently, based on the ability to form plaques (bottom). Top: The observartion of lysis zones with high-

titer lysate ($>10^9$ pfu/ml) in at least three independent experiments is indicated by colored bars. Bottom:

The infectivity of BASEL collection phages on diverse enterobacterial hosts was quantified as efficiency

of plating (EOP). Data points and error bars represent average and standard deviation of at least three

independent experiments. Since the data obtained with *Salmonella* Typhimurium 12023s and SL1344 were

indistinguishable, we only show the results of one representative strain (*S.* Typhimurium 12023s).

Data reporting

All data generated or analyzed during this study are included in this published article.

Accession numbers

The annotated genome sequences of all 66 newly isolated phages as well as T3(K12) have been

submitted to the NCBI GenBank database under accession numbers listed in S5 Table.

Supporting Information Captions

S1 Table. List of all bacterial strains used in this study.

The abbreviations in the selection column indicate the drug and its concentration that were used. Amp = ampicillin, Cam = chloramphenicol, Kan = kanamycin, Zeo = zeocin; 25 / 50 / 100 refer to 25 µg/ml, 50 µg/ml, and 100 = 100 µg/ml, respectively. The following mutants of the KEIO collection were used for qualitative top agar assays but are not included in the strain list because no phage showed any growth phenotype on them: *ompW::kanR*, *phoE::kanR*, *flgG::kanR*, *fepA::kanR*, *hofQ::kanR*, *cirA::kanR*, *fhuE::kanR*, *fiu::kanR*, *ompN::kanR*, *pgaA::kanR*, *chiP::kanR*, *ompL::kanR*, *yddB::kanR*, *fecA::kanR*, *uidC::kanR*, *nanC::kanR*, *yfaZ::kanR*, *bglH::kanR*, *bcsC::kanR*, *cusC::kanR*, *gfcE::kanR*, *mdtP::kanR*, *ompG::kanR*, *ompX::kanR*, *yfeN::kanR*, *csgF::kanR*, *wza::kanR*, *flu::kanR*, *nmpC::kanR*, *eaeH::kanR*, *ydiY::kanR*, *yiaT::kanR*, *yaiO::kanR*, *mdtQ::kanR*, *pgaB::kanR*, *mipA::kanR*, *pldA::kanR*, *yzcX::kanR*, *ydeT::kanR*, *blc::kanR*, *gspD::kanR*, *yjgL::kanR*

S2 Table. List of all oligonucleotide primers used in this study.

S3 Table. List of all plasmids used in this study.

The abbreviations in the selection column indicate the drug and its concentration that were used. Amp = ampicillin, Cam = chloramphenicol, Kan = kanamycin, Zeo = zeocin; 25 / 50 / 100 refer to 25 µg/ml, 50 µg/ml, and 100 = 100 µg/ml, respectively.

S4 Table. Construction of all plasmids generated in this study.

S5 Table. List of all phages used in this study.

The table lists all details regarding the taxonomic classification, isolation / source, host receptors, and genomic features of all phages used in this study. As *Caudovirales*, all these phages are classified as *Viruses* / *Duplodnaviria* / *Heunggongvirae* / *Uroviricota* / *Caudoviricetes* / *Caudovirales*, so only the classification on the level of family and below as defined by the ICTV [https://talk.ictvonline.org/taxonomy/; reference 142] is listed. For the reference phages, genome sizes and estimation of RM system recognition sites are based on the reference genomes in NCBI GenBank as indicated. Note that for phage T5 the reference genome includes the 10'219bp terminal repeat twice, unlike

all other *Markadamsvirinae* genomes that we listed. The P2*vir* phage had been isolated as a spontaneous mutant of a P2 *cox3* lysogen (supposedly unable to excise [148]) and was analyzed by whole-genome sequencing (as described in *Materials and Methods*), revealing few single-nucleotide differences to the reference genome, the expected *cox3* mutation, and a one-basepair insertion in the lysogeny repressor gene *C* that results in a premature stop codon. Recognition sites of RM systems were identified using Geneious Prime 2021.0.1. The numbers include the two orientations of palindromic recognition sequences separately and do not account for the fact that the counts are not fully comparable, e.g., because type III RM systems require two unmodified recognition sites in head-to-head orientation for cleavage [49]. Given the high number of type III RM recognition sites in all genomes because of their short length (5 nt in case of both EcoCFT_II and EcoP1_I, see Fig 3B), this limitation is largely theoretical.

S6 Table. List of all phage genomes used in the *in silico* analyses.

S1 Text. Construction of bacterial mutant strains.

S2 Text. Generation of the Maximum-Likelihood phylogenies shown in this article.

S3 Text. Technical considerations regarding the composition of the BASEL collection and the phenotyping of bacterial defense systems.

S1 Figure. Supplemental data for Figs 4 and 5.

(A) Maximum-Likelihood phylogeny of *Drexlerviridae* and the *Dhillonvirus* genus of *Siphoviridae* based on several core genes with bootstrap support of branches shown if > 70/100. It is clearly apparent that *Drexlerviridae* are split into two major clades, one formed by *Braunvirinae* and *Rogunavirinae* and another one formed by *Tempevirinae*, *Tunavirinae*, plus a few other groups. Given that the phylogenies strongly agree on all major branches, the root of the *Drexlerviridae* phylogeny shown in Fig 4B was placed between these two major clades. **(B)** The locus encoding lateral tail fibers was analyzed in a sequence alignment of the thirteen *Drexlerviridae* phage genomes of the BASEL collection (see *Materials and Methods*). It is clearly visible that the upstream and downstream regions (encoding genes involved in

recombination as well as primase / helicase proteins for genome replication) are highly conserved and fully

syntenic, with exception of small insertions in a few sequences. Conversely, only the most 5' end of the

largest lateral tail fiber protein gene is very similar among all analyzed genomes (green circle), while the

rest shows neither synteny nor clear homology across all genomes. **(C)** The *bona fide* RBP loci downstream

of the *gpJ* homolog are shown for all small siphoviruses (*Drexlerviridae* and *Siphoviridae* of *Dhillonvirus*,

Nonagvirus, and *Seuratvirus* genera) where we had experimentally determined the terminal receptor

(together with selected representatives where previous work had determined the receptor specificity). **(D)**

The *bona fide* RBP locus of *E. coli* phage RTP was aligned to the homologous locus of phage

AugustePiccard (Bas01) as described in *Materials and Methods*. For the region comprising *rtp44* and *rtp45*

of phage RTP, the pairwise identity of the two nucleotide sequences is ca. 93%.

S2 Figure. Supplemental data for Fig 6.

(A) The locus encoding lateral tail fibers was analyzed in sequence alignments of the five

Dhillonvirus phage genomes of the BASEL collection (top) and the seven *Nonagvirus + Seuratvirus* phage

genomes of the BASEL collection (bottom) as described in *Materials and Methods*. In both cases two clear

dips in overall sequence similarity are obvious, once at the *bona fide* RBP locus and then at the lateral tail

fiber locus downstream of the far 5' end of its first gene. **(B)** Schematic comparison of representative *bona*

fide RBP loci as shown in S1C Fig to the corresponding allele of *E. coli* phages JenK1, HdK1, and HdsG1

that does not match clearly match any of them.

S3 Figure. Supplemental data for Fig 7.

(A) The locus encoding lateral tail fibers was analyzed in a sequence alignment of the

Demerecviridae: Markadamsvirinae phage genomes of the BASEL collection as described in *Materials*

and Methods. Sequence identity is high upstream and downstream of the lateral tail fiber locus (with

exception of presence / absence of a few putative homing endonucleases) but drops considerably at the

lateral tail fiber genes. Note that, as described previously, the lateral tail fibers can either be composed of a

single large polypeptide or by two (or more) separate proteins [51, 60]. The same diversity in architecture

of lateral tail fibers can also be seen at the corresponding loci of small siphoviruses (S1B and S2A Figs).

(B) The illustration shows a phylogeny of the RBPs of all *Markadamsvirinae* phages shown in Fig 7B.

Briefly, the RBP genes of all genomes (invariably encoded directly upstream of the terminase genes) were

translated, aligned, and then used to generate a phylogeny as described in *Materials and Methods*. Three

clear clusters emerge, one including all phages known to bind BtuB (left), one including all phages known

to bind FepA (top right), and one including all phages known to bind FhuA (bottom right). Based on similar

analyses by others [62], we conclude that the position of RBPs in these clusters is predictive of terminal

receptor specificity of the phages encoding them.

S4 Figure. Supplemental data for Fig 8.

(A) Top agar assays with different surface protein mutants of the KEIO collection in comparison

to the ancestral *E. coli* K-12 BW25113 strain were performed with serial tenfold dilutions of all

Tevenvirinae phages used in this study (undiluted high-titer stocks at the bottom and increasingly diluted

samples towards the top). The phages show strongly or totally abolished growth on each one of the mutants

which identifies the primary receptor of each phage (also indicated by the color code highlighted on the

right). **(B)** Top agar assays of reference strain *E. coli* K-12 ΔRM carrying empty vector pBR322_ΔP*tet* or

pAH213_rexAB were performed with serial tenfold dilutions of phage T4 wildtype, a T4 variant encoding

an apparently hypomorphic *rIIAB* fusion (see (C)), and phage T5 (as control). The *rIIAB* mutant of phage

T4 is unable to form plaques on the *rexAB*-expressing host and shows only "lysis from without" [98], while

the T4 wildtype and phage T5 are not affected. **(D)** A T4 phage mutant was erroneously obtained from a

culture collection instead of the wildtype and encoded a peculiar *rII* allele with fusion of the *rIIA* and *rIIB*

open reading frames (shown as orange arrows; genome sequence determined by whole-genome

sequencing). Since such a mutant seems unlikely to arise spontaneously during shipping, we find it likely

that this phage strain is related to the *rIIAB* fusion mutants employed for discovery of the triplet nature of

the genetic code that were once commonly used [concisely reviewed in reference 149, 150]. Notably,

position and size of the deletion fusing *rIIA* and *rIIB* are indistinguishable from the sketch drawn by Benzer

and Champe for the *rIIAB* fusion mutant *r1589* which was used in the aforementioned work [151]. Unlike

T4 wildtype, the *rIIAB* fusion mutant is susceptible to *rexAB* when ectopically expressed from a plasmid

vector (see (C)) and therefore validates functionality of the *rexAB* construct.

S5 Figure. Supplemental data for Fig 9.

The illustration shows a sequence alignment of the locus encoding lateral tail fiber genes in phage

rV5 and new isolates DrSchubert, AlexBoehm, and JeffSchatz that broadly cover the phylogenetic range

of this genus (Fig 9B). It extends from the tail tape measure protein of phage rV5 (*gp49*) to the last large

lateral tail fiber gene (*gp27*) [70, 71]. Note that most genes are highly conserved including the lateral tail

fiber component with sugar deacetylase domain (compare Fig 9D; around position 8'000 in this alignment)

or the short tail fiber locus (compare Fig 9E). Only the three largest lateral tail fiber genes show considerable

allelic variation that is very strong for two of them (positions 10'000-14'000 and 16'000-19'000) and

moderate for another one (positions 29'000 to 34'000).