Differential Roles of Positive and Negative Valence Systems in the Development of Post-Traumatic Stress Psychopathology

Ziv Ben-Zion[1,2], Ofir Shany[1,3], Roee Admon[4,5], Nimrod Jackob Keynan[1,6], Netanell Avisdris[1,7], Shira Reznik Balter[1], Arieh Y. Shalev[8], Israel Liberzon[9] & Talma Hendler[1,2,3,10]

[1]Sagol Brain Institute Tel-Aviv, Wohl Institute for Advanced Imaging, Tel Aviv Sourasky Medical Center, Tel-Aviv, Israel

[2]Sagol School of Neuroscience, Tel-Aviv University, Tel-Aviv, Israel

[3]School of Psychological Sciences, Faculty of Social Sciences, Tel-Aviv University, Tel-Aviv, Israel

[4]School of Psychological Sciences, University of Haifa, Haifa, Israel

[5]The Integrated Brain and Behavior Research Center (IBBRC), University of Haifa, Haifa, Israel

[6]Department of Psychiatry and Behavioral Sciences, Stanford University School of Medicine, Stanford, CA, USA

[7]School of Computer Science and Engineering, the Hebrew University of Jerusalem, Jerusalem, Israel

[8]Department of Psychiatry, NYU Langone Medical Center, New York, NY, USA

[9]Department of Psychiatry, Texas A&M Health Science Center, TX, USA

[10]Sackler Faculty of Medicine, Tel-Aviv University, Tel-Aviv, Israel

Corresponding Author:

Prof. Talma Hendler, M.D. Ph.D.

Professor of Psychiatry and Neuroscience, Sagol School of Neuroscience, Sackler Faculty of Medicine, and School of Psychological Sciences, Tel Aviv University, P.O. Box 39040, Tel Aviv, Israel; Director, Sagol Brain Institute, Wohl Institute for Advanced Imaging, Tel Aviv Sourasky Medical Center, Weizmann 6, Tel Aviv, 64239, Israel;

Phone: +972 (0)3 697 3549. Email: talma@tlvmc.gov.il , thendler@post.tau.ac.il

Abstract

Negative and positive valence systems (NVS and PVS) pertain to processing of aversive and rewarding stimuli, respectively. Post-Traumatic Stress Disorder (PTSD) has been typically associated with abnormalities of the NVS, mostly related to heightened threat processing, yet more recent work also suggests deficits in PVS functionality in PTSD,

mainly in the form of reduced reward functioning. The current study examined the hypothesis that individuals' ability to recover from a potentially traumatic event relies on promoting reward-seeking behaviors (i.e., PVS) alongside diminished threat assessment (i.e., NVS), during the first year following trauma, a critical period for PTSD development or recovery. To do so, we longitudinally tracked behavioral and neural responses among 171 adult civilians with early post-traumatic stress symptoms at 1-, 6- and 14-months following trauma exposure (TP1, TP2, and TP3, respectively). At each time-point, participants played a naturalistic game encompassing dynamic provocation of risk-taking, punishments and rewards in an fMRI setting. Results showed that greater amygdala activation and functional connectivity with the lateral orbitofrontal cortex (lOFC) in response to punishments (i.e., hyperactive NVS) at TP1 were associated with more severe post-traumatic stress symptoms at both TP1 and TP3 (but not at TP2), and specifically with more hyperarousal and intrusion symptoms. On the other hand, decreased ventral striatum (VS) activity and functional connectivity with the ventromedial prefrontal cortex (vmPFC) in response to rewards (i.e., hypoactive PVS) at TP1 were associated with more severe post-traumatic stress symptoms at TP3 (but not at TP1 or TP2), and specifically with more avoidance symptoms. Explainable machine learning revealed the primacy of the PVS over the NVS at TP1 in predicting PTSD symptom development from TP1 to TP3. Behaviorally, fewer risky choices in the

task were associated with more severe symptoms at TP1, but not at TP2 or TP3. Finally, an integrative exploratory analysis revealed that reduction in risky choices in the task (from TP1 to TP2) moderated the relation between NVS hyperactivity at TP1 and symptom severity at TP3. Altogether, our results support the idea that trauma exposure might alter both NVS and PVS processing. While NVS presents early heightened saliency processing in the immediate aftermath of trauma, early PVS only affects the long-term outcome of traumatic stress. These insights inform possible mechanism-driven therapeutic strategies for PTSD, addressing not only negative but also positive valence processing.

Introduction

The concept of negative and positive valence systems (NVS and PVS, respectively) originated in psychology over a century ago, yet more recently was incorporated into the field of clinical neuroscience[1] and the Research Domain Criteria (RDoC)[2,3]. The NVS is responsible for responses to aversive situations or contexts, evoking negative feelings such as fear, anxiety, and loss, whereas the PVS is in charge of responses to positive motivational situations or contexts such as reward-seeking,

consummatory behavior, and reward learning. Valence estimation could be challenging in real-life as stimuli often evoke mixed or even conflicting emotions and consequence behaviors. Traumatic stress might hinder accurate valence estimations[4-6], as it increases vigilance and drains cognitive resources[7,8]. While this may serve survival in the immediate aftermath of stress, transition into reward-driven behavior over time, despite the presence of a heightened threat, is necessary for promoting stress resilience[9-13]. Indeed, chronic stress psychopathologies such as Post-Traumatic Stress Disorder (PTSD) are often characterized by a tendency to sacrifice potential rewards to avoid aversive encounters [14-17]. Here, we examine the idea that individuals' ability to recover from a potentially traumatic event relies on differential processing of the PVS and NVS by assessing neurobehavioral manifestations of these systems separately and together, in the early aftermath of trauma.

Substantial evidence links PTSD to abnormal NVS, consistently showing increased sensitivity to various negative stimuli (e.g. symptom provocation, fearful faces)[18,19] among PTSD patients, indicating elevated NVS, a phenomenon that may relate to the clinical symptoms of hyperarousal and intrusion (i.e., re-experiencing)[20-23]. Furthermore, studies using decision-making tasks demonstrated an association between PTSD and increased behavioral aversion to risk[24,25] and ambiguous losses[26], possibly reflecting the generalized (not only trauma-specific) oversensitivity of the NVS, often seen in PTSD patients. At the

neural level, the role of the NVS in PTSD has been repeatedly documented as abnormally heightened amygdala activation and hyperactive salience network (e.g., anterior insula, dorsal anterior cingulate cortex) in response to negative stimuli[20,27-29]. Furthermore, aberrant functional connectivity of the amygdala with the prefrontal cortex (PFC) in response to negative stimuli was observed in PTSD, specifically with the orbitofrontal cortex (OFC) in humans and the para-limbic cortex in animals, suggesting disrupted emotion regulatory capacity[30-33].

More recent work suggests that PTSD might also involve abnormalities in the PVS, as indicated by deficient reward anticipation, decreased approach (reward-seeking) behavior, and diminished hedonic responses to rewarding outcomes[34,35]. Reward processing is known to involve the mesocorticolimbic pathway, represented by dopamine projections from the ventral tegmental area (VTA) to the ventral striatum (VS), including the nucleus accumbens (NAcc), and further to ventromedial/orbital frontal brain structures[36,37]. While decreased VS activation to positive stimuli was demonstrated initially in depressed individuals mostly in relation to anhedonia[38,39], it was also reported in PTSD patients in response to monetary gains[40,41] and happy faces[42]. Recent studies further pointed to aberrant functional connectivity between the VS and the ventromedial PFC (vmPFC) in PTSD, suggesting a compromised reward circuitry in this disorder[43,44].

Although both systems seem to have a role in PTSD, their relative contribution to the development of post-traumatic psychopathology remains largely unknown, due to several substantial clinical and methodological challenges. First, only a small portion (around 20%) of those with early stress symptoms go on to develop chronic PTSD[45,46]. Second, even within the group of PTSD patients, clinical phenotypes are heterogeneous[47,48], and different symptom manifestations (e.g. hyperarousal vs. avoidance) might be related to different neurobehavioral processes (e.g. NVS vs. PVS). Third, the typical cross-sectional designs used in PTSD research cannot infer the immediate response to trauma nor on any potential dynamics that may occur during the first year post-trauma, a critical period that determines who will develop PTSD and who will recover[49,50]. While there is an increase in longitudinal studies of PTSD in recent years, most of them focused on either NVS or PVS and did not combine clinical, behavioral, and neural measurements altogether in trying to unravel the unique role of each system in the same person over time.

To overcome these challenges, we conducted a large-scale longitudinal fMRI study of recent trauma survivors (see study protocol[51]). A sample of n=171 adult civilians were screened for early PTSD symptoms, suggestive of chronic PTSD risk[52,53], within 10–14 days following their release from a general hospital's emergency room (ER). Participants were longitudinally tracked at 1-, 6- and 14-months

following trauma exposure (TP1, TP2, and TP3, respectively) as they underwent fMRI scanning while playing a naturalistic interactive gambling choice game encompassing safe or risky choices resulting in outcomes of reward (win) or punishment (loss). To depict the neural processing of the two valence systems, the NVS was represented by the amygdala's response to punishments vs. rewards, while the PVS was represented by VS response to rewards vs. punishments. This paradigm, termed here 'Safe or Risky Domino Choice' (SRDC), was previously shown to elicit VS response to rewarding outcomes and amygdala response to punishment outcomes[25,54-57]. More so, in order to win, individuals' behavior in the SRDC game must include both safe and risky choices (determined by the type of domino chip chosen and its subsequent exposure or not by the opponent), enabling a behavioral marker of the balance between the PVS and NVS activation.

Our first aim was to find neurobehavioral indicators of NVS and PVS abnormalities associated with post-traumatic symptom severity shortly after exposure (TP1). Based on previous findings[24-26,58], we hypothesized that more severe post-traumatic symptoms would be associated with greater response of the amygdala to punishment vs. reward, decreased response of the VS to reward vs. punishment, and altered functional connectivity of the VS and the amygdala with the PFC. In light of our previous finding with the SRDC game in soldiers[25], we expected that individuals with more severe symptoms at TP1 would take fewer risky-

choices in the game, reflecting combined abnormality of hyperactive NVS and hypoactive PVS. Our second aim was to reveal the relationship between early NVS and PVS abnormalities and post-traumatic stress symptom development (i.e., beyond initial severity) within the first year following trauma exposure (TP2 and TP3). We hypothesized that greater response (activity and connectivity) of the amygdala to punishment vs. reward, decreased response (activity and connectivity) of the VS to reward vs. punishment, and decreased risky choice behavior at TP1, would be predictive of more severe post-traumatic symptoms at TP2 and TP3. Specifically, we expected that individual differences in NVS activation (i.e., amygdala response to punishment vs. reward) would be associated with symptom clusters of hyperarousal and intrusion, whereas differences in PVS activation (i.e., VS response to reward vs. punishment) would be associated with avoidance symptoms.

Methods

Participants. The study group included adult survivors of potentially traumatic events who were admitted to Tel-Aviv Sourasky Medical Center's Emergency Room (ER). The most common trauma type among participants was motor vehicle accidents (n=137, 80%), while other traumatic events included assaults, terror attacks, drowning, mass casualty incidents, animal attacks, robbery, and electrocution. Exclusion

criteria included head trauma or coma exceeding 30 minutes, incompatibility for MRI scanning, history of substance abuse, current or past psychotic disorder, or chronic PTSD diagnosis pre-admission to ER. Only trauma survivors without a known medical condition that interfered with their ability to give informed consent or to cooperate with screening and/or treatment were included. A total of 171 participants completed clinical and neuroimaging assessments within one-month following their traumatic incident (TP1). Of these, 39 individuals were excluded due to missing (n=16) or partial (n=5) functional scan while performing the SRDC game; poor quality of the functional scan (e.g., movements, artifacts, etc.) (n=6); missing or poor structural scan (n=5); missing or partial behavioral data of the paradigm (n=5); not understanding the instructions properly (n=1); and missing clinical data (n=1), for a final sample size of 132 individuals at TP1. Of these 132 participants, 115 and 112 participants completed clinical interviews at TP2 and TP3 (respectively). Participants' demographic and clinical characteristics across the three time-points (TP1, TP2, and TP3) are presented in Table 1. Of note, 6 individuals were excluded from the neural analysis at TP2 and 22 at TP3, due to missing/partial/poor quality of the functional scan, missing / poor quality structural scan, or missing/partial behavioral data of the SRDC game, for a final sample size of 109 and 90 individuals with valid neural data at TP2 and TP3.

Procedure. A member of the research team identified potential trauma-exposed individuals via the ER computerized medical records. Within 10-14 days of trauma exposure, approximately n=4,000 potential participants were contacted by telephone for initial screening. Acute PTSD symptoms (indicative of the risk for PTSD development[52]) were assessed using a modified dichotomous version of the PTSD Checklist (PCL) questionnaire[59]. Out of 4,000 potential participants, only 600 met PTSD symptom criteria (except for the 1-month duration) based on the phone-screening interview. Participants who did not meet any of the exclusion criteria were invited to participate in both comprehensive face-to-face clinical assessment and an fMRI scan, within one-month post-trauma (TP1). We preferentially enrolled survivors who met PTSD diagnosis in the face-to-face clinical interview, but also enrolled some individuals with sub-threshold PTSD (n=35). Two identical follow-up meetings, including both clinical and neural assessments, were conducted at 6- and 14-months following trauma (TP2 and TP3, respectively). For more details, see Ben Zion et al. (2019).[51]

Clinical Assessments. PTSD diagnosis and severity at each time-point were determined by a comprehensive clinical interview conducted by trained and certified clinical interviewers. A continuous measure of total symptom severity was obtained by summing individual items' scores of the Clinician-Administered PTSD Scale for DSM-5 (CAPS-5)[60], the current

gold-standard for PTSD diagnosis. Total scores were further computed for each of the DSM-5 symptom clusters: intrusion (cluster B), avoidance (cluster C), negative alterations in cognition and mood (cluster D), and hyperarousal (cluster E).

fMRI Paradigm. During scanning, participants played a 2-player gambling game for 14 minutes (termed herby Safe or Risk Domino Choice game; SRDC), in which they were required to take risky choices to win (see Figure 1). The effectiveness of the SRDC to detect individuals' sensitivity to risk, punishment and reward was previously validated in both healthy and clinical populations[25,54-57,61]. Participants were told that their opponent is the experimenter who decides whether to expose their choice or not, thus their choices can increase their chances of winning. In fact, however, the computer randomly generated the opponent's responses in a predetermined pattern to allow a balanced design (exposing the player's choices 50% of the time). The SRDC paradigm involved four intervals: decision-making (deciding which chip to choose), decision-execution (taking risky or safe chips), anticipation of an outcome (waiting passively for the opponent's decision to expose or not the player's choice), and response to an outcome (receiving a reward or punishment, defined by the chosen chip and opponent's decision). In this study, our focus was on the decision-making interval for behavioral indexing (i.e., individual tendency to take risky vs safe choices) and on

the neural responses to the outcome (rewards vs. punishments). At the beginning of each game, 12 random domino chips were assigned to the participant, while one domino "master" chip (constant throughout the game) appeared at the top left corner of the board. In each round of the game, players had to choose one chip and place it face down adjacent to the master chip. They then had to wait for the opponent's response (i.e., anticipation), to see whether the opponent challenges this choice by uncovering the chosen chip or not (i.e., outcome). Players were able to win this competitive game if they were able to successfully dispose of all 12 chips within 4 minutes. Each assigned chip either matched the master chip (had at least one of the master chip's numbers) or was non-matching. Since the master chip was constant throughout the game, it was only possible to win by choosing both matching and non-matching chips. In the game context, matching chips were considered ''safe'' moves since they were associated with rewards, while non-matching chips were considered ''risky'' moves since they were associated with punishments. Accordingly, based on players' choices, there were two possible anticipation periods, "risky anticipation" following a non-matching choice or "safe anticipation" following a matching choice. Based on players' choices and opponent's response, there were four possible consequences (outcomes) per game round: (1) <u>Show of a non-matching chip (i.e., main punishment):</u> the choice of a non-matching chip was exposed and the player was punished by receiving back the selected chip

plus 2 additional chips from the deck; (2) No-show of a non-matching chip (i.e., relative reward): the choice of a non-matching chip remained unexposed and only the selected chip was disposed of, so the player was relatively rewarded as he got away with a non-matching choice; (3) Show of a matching chip (i.e., main rewards): the choice of a matching chip was exposed and the player was rewarded by the disposal of the selected chip and one additional random chip from the game board; (4) No-show of a matching chip (i.e., relative punishment): the choice of a matching chip was not exposed and only the selected match chip was disposed of, so the player was relatively punished as he could have disposed of a non-matching chip instead. For more details, see supplementary methods.

Behavioral Analysis of the Paradigm. To characterize individuals' behavioral choices during the game, a *risky choice index* was defined as the ratio between the number of risky choices (e.g., choosing a non-matching chip) and the total number of choices throughout the entire game (e.g., choosing either a matching or non-matching chip), multiplied by 100 (to obtain percentage). Rounds in which participants had no actual choice between safe and risky choices were excluded (i.e., when there were only matching or only non-matching chips on the screen). Previous work using the SRDC game in healthy individuals showed an average risky choice index of 50% across individuals[54]. We thus assume here that a balanced PVS/NVS processing is needed to achieve an index of 50%, while

an index greater than 50% represents a bias towards more risky behavior, and less than 50% represents a bias towards safer behavior (i.e. risk aversion and avoidance tendencies).

$$risky\ choice\ index\ (\%) = \frac{\#\ non\ matching\ chips}{\#\ non\ matching\ chips + \#\ matching\ chips} * 100$$

fMRI Data Acquisition. Whole-brain functional and anatomical images were acquired using a 3.0 Tesla Siemens MRI system (MAGNETOM Prisma, Germany) with a 20-channel head coil at Tel-Aviv Sourasky Medical Center. Functional images were acquired in an interleaved order (anterior to posterior), using a T2*-weighted gradient-echo planar imaging pulse sequence (TR/TE=2000/28ms, flip angle= 90°, voxel size 2.2×2.2×2.2mm, FOV=220×220mm, slice thickness=3mm, 36 slices per volume). A T1-weighted three-dimensional anatomical image was also collected, using a magnetization prepared rapid gradient echo (MPRAGE) sequence (TR/TE=2400/2.29ms, flip angle = 8°, voxel size 0.7 × 0.7 × 0.7 mm, FOV = 224 × 224 mm), enabling optimal localization of the functional effects.

fMRI Data Analysis. Preprocessing was conducted using FMRIPREP version 1.5.8[62], a Nipype based tool[63]. The preprocessing procedures included spatial realignment of the echo planar imaging volumes, co-registration with the structural image, segmentation, normalization, and spatial smoothing (for full details, see supplementary methods). The

preprocessed imaging data was further analyzed using Statistical Parametric Mapping 12 (SPM12). The size of the effect for each condition for each participant was computed by a general linear model (GLM), which included the different conditions of the game: "choose", "ready", "go", "picked match", "picked non-match", "show match", "show non-match", "no show match" and "no show non-match". Individual statistical parametric maps were calculated for the a-priori defined contrasts of interest of "rewards vs. punishments" (receiving both rewarding outcomes > receiving both punishing outcomes) and "punishments vs. rewards" (receiving both punishing outcomes > receiving both rewarding outcomes). This was done for all participants at each time-point separately (n=132 at TP1, n=109 at TP2, and n=90 at TP3).

ROI analysis. Based on previous findings using the same paradigm[25,54-57,61], two main regions of interest (ROIs) were defined using the Human Brainnetome atlas (developed by Fan and colleagues[64]) and California Institute of Technology 168 (CIT-168) atlas (developed by Pauli and colleagues[65]). In the condition of rewards vs. punishments, the focus was on the left and right ventral striatum (VS), composed of the ventral caudate from the Human Brainnetome atlas (regions 219-220) and the nucleus accumbens from the CIT-168 atlas. In the opposite condition of punishments vs. rewards, the focus was on the left and right amygdala, composed of the medial (regions 211-212) and lateral amygdala (regions 213-214) from the Human Brainnetome atlas. MarsBaR region of interest

toolbox for SPM[66] was used to extract participants' contrast activations (average beta weight) from each ROI separately for each time point. Analyses were performed separately for each hemisphere (left and right amygdala, left and right VS). For more details regarding whole-brain and ROI analyses, see supplementary methods.

Functional Connectivity Analysis. Examination of how the brain region interacts in a task-dependent manner was performed using generalized psychophysiological interaction (gPPI) as implemented in CONN toolbox[67,68] (for full details, see supplementary methods). ROI to ROI analysis was performed using the two main a-priori ROIs, amygdala and VS, as seed regions and two prefrontal a-priori ROIs, ventromedial prefrontal cortex (vmPFC) and lateral orbitofrontal cortex (lOFC), as target regions. These regions were chosen based on previous literature reporting medial- medial-lateral segregation in the human prefrontal cortex, indicating that rewards are processed in the medial part of the PFC, while punishments are processed in the OFC[69-73]. The two target regions were anatomically defined by the Human Brainnetome atlas[64]: The vmPFC was composed of regions 41-42, 47-48, whereas the lOFC was composed of regions 43-44, 45-46, 51-52. For more details, see supplementary methods.

Statistical Analysis. IBM SPSS Statistics for Windows[74] and R[75] software were both used for the statistical procedures. Participants with extreme

scores of ±3 standard deviations from the mean in the relevant variables were defined as outliers and excluded from the analysis. For all statistical tests, α=0.05 was used with either one-sided a-priori hypotheses or two-sided non-directional hypotheses (according to the a-priori hypotheses outlined at the end of the introduction section). Benjamini-Hochberg False Discovery Rate (FDR) correction (q<0.05)[76] was calculated to control for multiple comparisons.

Predictor Importance Ranking. To examine the contribution of early neural activations (at TP1) and rank their importance for the prediction of PTSD symptom severity at the study's endpoint (TP3), we used SHAP (Shapley Additive exPlanation)[77], a state-of-the-art methodology in the field of explainable machine learning. SHAP estimates "Shapely" values, which provide a surrogate for the individual additive contribution of each feature to the prediction. In other words, SHAP's rank order informs which feature values influence the prediction the most while accounting for the influence of all other feature values and controlling for the order in which features are added to the model[77].

Results

Neurobehavioral indicators of PVS and NVS for PTSD severity shortly after trauma

To find the neural indicators of the PVS and NVS associated with PTSD symptom severity shortly after exposure (TP1), partial correlations were computed between neural indices (mean activations of the VS and amygdala to rewards vs. punishments, as well as their connectivity patterns), and PTSD symptom severity (i.e., CAPS-5 total scores), while controlling for participants' age, gender, and trauma type (i.e., covariates). Results revealed a significant positive correlation between the amygdala's responses to punishments (vs. rewards) and symptom severity at TP1 *(n=128; left amygdala: r=0.155, p=0.043, p_{FDR}=0.043; right amygdala: r=0.162, p=0.035, p_{FDR}=0.043)*, indicating that individuals with more severe symptoms showed greater amygdala activation (Fig. 2a). In contrast to our expectation, VS activation in the contrast of rewards vs. punishments was not significantly associated with symptom severity at TP1 *(n=131; left VS: r=0.022, p=0.401; right VS: r=0.048, p=0.297)* (Fig. 2b). When examining functional connectivity patterns with predetermined PFC regions, we found that stronger amygdala-lOFC connectivity in the contrast of punishments vs. rewards was associated with more severe symptoms *(n=124; right amygdala – left lOFC: r=0.254, p=0.005, q_{-FDR}<0.05)*, indicating that greater connectivity was associated with greater symptom severity at TP1 (Fig. 3c). VS functional connectivity with the PFC in the contrast of rewards vs.

punishments was not related to total symptom severity (n=122, q_{FDR}>0.05). For whole-brain results, please refer to supplementary results, supplementary table 1, and supplementary figure 1.

To assess the relation between risky choice index, a behavioral measure reflecting balance/imbalance between the PVS and NVS, and PTSD symptom severity, we used both independent t-test between individuals with and without PTSD, and a partial correlation between the behavior and symptoms while controlling for participants' age, gender, and trauma type. At the group level, a two-sample independent t-test revealed a significant difference in risky choice index between 'PTSD' (n=97, dark gray) and 'non-PTSD' (n=35, light gray) groups *(t=2.833, p=0.006)* (Fig. 3a). While individuals who did not qualify for PTSD diagnosis showed an average choice of 50/50 between risky and safe choices *(M=49.69%, SE=2.00%)*, those with PTSD at TP1 displayed a significantly decreased risky choice index *(M=42.73%, SE=1.40%)* with a bias towards safer behavior. Notably, similar differences were observed between the groups when examining risky choice index following rewards *(t=3.466, p=0.001)* and following punishments *(t=2.828, p=0.005)*, separately. Specifically, while non-PTSD individuals (n=35) showed balanced behavior both after receiving rewards *(M=50.39%, SE=2.53%)* and punishments *(M=52.09%, SD=2.77%)*, those with PTSD at TP1 (n=97) took significantly fewer risks after receiving rewards *(M=39.32%, SE=1.96%)* and punishments *(M=43.04%, SD=1.64%)*. At the individual

level, a negative correlation was found between risky choice index and CAPS-5 total scores at TP1 *(n=132, r=-0.185, p=0.018)*, indicating that individuals with more severe post-traumatic stress symptoms made less risky choices in the SDRC game (Fig. 3b). This behavioral tendency towards safe behavior was particularly associated with more severe avoidance (CAPS-5 cluster C) *(r=-0.244, p=0.003, p_{FDR}=0.012)* and intrusive symptoms (CAPS-5 cluster B) *(r=-0.212, p=0.016, p_{FDR}=0.032)*, but not with other symptom clusters (Fig. 3b).

Neural abnormalities of PVS and NVS predict first-year PTSD development

To assess the relationship between early neural PVS and NVS abnormalities (TP1) and PTSD symptom development within the first year following trauma exposure (TP2 and TP3), partial correlations were computed between neural indices at TP1 and PTSD symptom severity at TP2 and TP3, while controlling for participants' age, gender, trauma type, and initial symptom severity (CAPS-5 total scores at TP1). In line with our hypothesis, both hyperactive NVS (i.e., amygdala's activation to punishment) and hypoactive PVS (i.e., VS activation to reward) at TP1 were significantly predictive of more severe PTSD symptoms at TP3. Specifically, greater left amygdala activation at TP1 was associated with higher CAPS-5 total scores at TP3 *(n=108; r=0.197, p=0.022)* (see Fig. 4a); and decreased right VS activation at TP1 was associated with higher CAPS-5 total scores at TP3 *(n=111, r=-0.235, p=0.007)* (see Fig. 4b).

Individuals with hyperactive NVS or hypoactive PVS early after trauma (TP1) were thus prone to develop more severe symptoms a year later (TP3).

Testing specific symptom clusters, greater amygdala activation to punishments at TP1 was associated with more severe hyperarousal *(r=0.176, p=0.037, p_{FDR}=0.074)* and intrusion *(r=0.217, p=0.027, p_{FDR}=0.074)* at TP3 (Fig. 4a), whereas decreased VS activation to rewards at TP1 was linked to more avoidance at TP3 *(r=-0.285, p=0.001, p_{FDR}=0.004)* (Fig. 4b). Of note, neither amygdala activation to punishments nor the VS activation to rewards at TP1 were significantly associated with CAPS-5 total scores at TP2 *(n=114; left amygdala: r=-0.021, p=0.413; right amygdala: r=-0.146, p=0.320; left VS: r=0.065, p=0.249; right VS: r=0.006, p=0.475)*, nor with any specific CAPS-5 symptom clusters.

Examining the power of functional connectivity patterns of the PVS and NVS at TP1 for predicting PTSD at TP3 revealed such a relation only for the PVS. Specifically, decreased VS-vmPFC connectivity during rewards vs. punishments at TP1 was associated with more severe post-traumatic stress symptoms at TP3 *(n=108, right VS – right vmPFC: r=-0.292, p=0.003, q_{-FDR}<0.05)*, indicating that individuals with less VS-vmPFC connectivity at TP1 developed more severe PTSD symptoms at TP3 (see Fig. 4c). Amygdala functional connectivity with the lOFC in the contrast of

punishments vs. rewards at TP1 was not related to PTSD symptom severity at TP3 (n=110, $q_{-FDR}>0.05$).

To assess the relation between risky choice index and PTSD symptom severity, we used both independent t-test between individuals with and without PTSD, and a partial correlation between behavior and symptoms while controlling for participants' age, gender, trauma type, and initial symptom severity. At the group level, no significant difference was found between PTSD and non-PTSD groups at TP2 *(t=0.760, p=0.449)* or TP3 *(t=1.120, p=0.265)*. Similarly, at the behavioral level, no significant correlations were found between risky choice index at TP1 and PTSD symptoms at TP2 *(n=115, r=-0.039, p=0.341)* or TP3 *(n=112, r=-0.073, p=0.226)*.

Finally, we tested the relative contribution of amygdala and VS functionality (activation and connectivity) at TP1 for PTSD symptom severity per CAPS-5 scores at TP3, using linear regression with the baseline neurobehavioral indices of PVS and NVS from TP1, which significantly predicted post-traumatic stress symptoms at TP3: left amygdala activation to punishments vs. rewards (Fig. 4a), right VS activation to rewards vs. punishments (Fig. 4b), and right VS-right vmPFC functional connectivity during rewards vs. punishments (Fig. 4c). As expected, all three variables together at TP1 accounted for a significant amount of variance of CAPS-5 total scores at TP3 *(n=105, R²=0.200, $F_{3,101}=8.398$, p<0.001)*. To identify the relative importance of these three

neural predictors, we calculated importance values using the SHAP[77] analytic approach (see Predictor Importance Ranking in Methods). In terms of absolute feature importance, PVS abnormalities showed greater importance compared to NVS abnormalities. VS-vmPFC connectivity during rewards vs. punishments at TP1 was the best predictor of subsequent CAPS scores at TP3, followed by the right VS average response to rewards vs. punishments at TP1, and lastly the left amygdala's response to punishments vs. rewards at TP1 (Fig. 4d, right panel). Notably, when comparing the variances of VS activity and connectivity to amygdala activity, it seems that while the importance/contribution of the VS differed greatly between individuals (SHAP values ranging from −6 to +6), the amygdala had a small contribution in most participants (SHAP values between −2 to +2), and a large contribution to only a minority (Fig. 4d, left panel).

Integrative modeling of brain, behavior and PTSD development

In light of the null finding of the relationship between risky choice index at TP1 and PTSD symptoms at TP2 or TP3, we further explored the possibility that behavioral change in risky choice in the SRDC game (TP2–TP1 or TP3–TP1) moderates the relation between early NVS or PVS activity (at TP1) and post-traumatic stress symptom severity at the study's endpoint (TP3). Moderation effects were tested with PROCESS macro for SPSS[78,79] using hierarchical multiple regression analysis with centered

variables and a centered interaction term. Specifically, 8 moderation models were tested, including 4 different neural activations (left and right amygdala and VS at TP1) and 2 different behavioral measures (change in risky choice index from TP1 to TP2, and from TP1 to TP3).

Of these 8 moderation models tested, only one was statistically significant after correcting for multiple comparisons *(q-FDR<0.05)* (see Fig. 5a). This regression model consisted of left amygdala activation to punishments vs. rewards at TP1, change in risky choice index from TP1 to TP2, and the interaction between them, together with four covariates (age, gender, trauma type, and initial symptom severity), accounted for a significant amount of variance in symptom severity at TP3 *(n=90; R²=0.400, F$_{7,82}$=7.805, p<0.001).* A significant interaction (moderation) effect was detected between amygdala activation at TP1 and risk-taking behavior (change from TP1 to TP2) in predicting CAPS-5 total scores at TP3 *(b=-1.074, t$_{82}$=-3.311, p=0.001, p$_{FDR}$=0.011)* (see supplementary table 2). This significant interaction was further probed by testing the conditional effects of the left amygdala activation at different values of the behavioral change in risky choice index (Mean-1SD, Mean, Mean+1SD) (see Fig. 5b). For individuals who demonstrated a shift towards less risky choices over time (negative change in TP2-TP1 risky choice index, blue line), the amygdala's activation to punishments at TP1 was associated with more symptoms at TP3 *(t=3.754, p=0.003).* However, for those with no major change (green line) or with a shift

towards more risky behavior from TP1 to TP (red line), the relation between amygdala activation at TP1 and CAPS-5 total scores at TP3 was not significant *(p>0.05)* (Fig. 5b).

Discussion

The longitudinal design of this fMRI study, along with the use of a naturalistic gambling task in a large cohort of recent trauma survivors, enabled a comprehensive multi-parametric assessment of the relationships between the two valence systems' functionality and the development of PTSD. The NVS was represented by the amygdala's response to punishments vs. rewards, whereas the PVS was represented by the VS response to rewards vs. punishments. Our results demonstrate the differential contribution of the valence systems in the early manifestation of post-traumatic psychopathology (TP1), as well as in predicting the development of symptom severity (beyond initial symptoms) over the first year following traumatic exposure (TP3). At TP1, we found a leading contribution of the NVS neural processing to post-traumatic stress symptom severity (Fig. 2). While heightened processing in NVS (shown as hyperactive amygdala in response to punishment and its connectivity with the lOFC) was associated with more severe PTSD, diminished processing of PVS (as indicated by VS activity to reward and its connectivity with the vmPFC) was not associated with early symptoms.

Behaviorally, individuals with more severe symptoms at TP1 showed less risky choices (Fig.3), possibly reflecting an imbalance between NVS and PVS. Neural activation of the two valence systems at TP1 showed an opposite association with the development of post-traumatic stress symptoms at TP3 (beyond initial symptom severity) (Fig. 4). Hyperactive NVS and hypoactive PVS at TP1 were thus both associated with more severe symptoms at TP3. In a combined regression model aiming to predict symptom severity at TP3, early PVS neural abnormalities at TP1 were more important to the prediction model compared to early NVS abnormalities (Fig. 4d). Although we did not find risky-choice behavior at TP1 to be predictive of PTSD symptoms at TP3, an exploratory moderation analysis revealed that individuals with both early NVS (and not PVS) abnormality (TP1) and a reduction in risky choice behavior (from TP1 to TP2) were prone to develop more severe PTSD symptoms at TP3 (Fig. 5).

Contribution of NVS and PVS to early PTSD manifestations

The large cohort of symptomatic individuals in this study (n=132) provided the opportunity to examine the variability in neurobehavioral measures of PVS and NVS in relation to post-traumatic stress symptom severity as early as one-month after trauma (TP1). Consistent with the vast literature on increased amygdala response to negative stimuli in PTSD[20,27–29,80,81], hyperactive amygdala to punishments during the SRDC

game was associated with increased general post-traumatic stress symptom severity. Furthermore, increased amygdala-OFC functional connectivity while receiving punishments was also linked to greater PTSD severity. The OFC has a known role in the modulation of the amygdala[82,83] during volitional suppression of negative emotion[84] and in the presence of threatening stimuli[85], and its connectivity with the amygdala is involved in processing negative outcomes that signal a need for a behavioral change[72,73]. Indeed, disturbed fronto-amygdalar connectivity was previously observed in patients with PTSD[86,87] and other anxiety disorders[88,89], but this might not be specific as it was also found in depression[90,91] and bipolar disorders[92]. Although the current study cannot confidently disentangle causes from consequences of traumatic stress, the causal role of the NVS was implicated in previous prospective studies, showing for example that hyperactive amygdala in soldiers prior to exposure to stressful military experience was associated with more PTSD symptoms post-exposure[25,93].

Decreased risk-taking during the game (less than 50%) was also associated with more severe PTSD symptoms at TP1. This safer than risky choice tendency was apparent after both reward and punishment outcomes, suggesting an outcome independent behavior at an early stage. This supports the specific association with avoidance symptoms. It is interesting to note that trauma-exposed individuals who did not meet PTSD diagnostic criteria (n=35) at TP1 demonstrated risk taking-behavior

of around 50% throughout the whole game (half safe and half risky choices), similar to previous studies in healthy individuals using the same SRDC game[54-57,61]. This adaptive flexible behavior was optimal for wining the SRDC paradigm, as the computer randomly generated the opponent's responses, exposing the player's choices 50% of the time (see fMRI Paradigm under Methods). In contrast, individuals diagnosed with PTSD at TP1 (n=97) showed significantly less risk-taking across the game and thus reduced their chance of winning in the end. We suspect that this behavioral abnormality might reflect an imbalance between the valence systems, combining increased threat from negative outcomes (heightened NVS) and reluctance to take risks in order to achieve rewards which might not elicit a great enough hedonic response (diminished PVS). Such imbalance is in line with previous work in chronic PTSD patients reporting increased behavioral aversion to risky monetary gains and ambiguous monetary losses[24,26]. This further corresponds to the idea of Stein & Paulus (2009)[15] that trauma exposure might alter the homeostatic balance in motivation behavior towards less approach and more avoidance, which in turn might lead to the development of chronic PTSD. To elucidate the exact mechanism that underlies this behavioral abnormality, future studies should compare more formal behavioral indices of reinforcement learning and reward-seeking and valuation.

Early NVS and PVS neural abnormalities predict PTSD first year development

Using a longitudinal design, we were able to demonstrate that neural responsivity of NVS and PVS components shortly after trauma predicted post-traumatic stress symptom severity at more than one year following the traumatic event. Specifically, increased amygdala response to punishment and decreased VS response to reward at TP1 were associated with more severe symptoms at TP3 (beyond initial symptom severity) (Figure 2). These opposite effects of the NVS and PVS allude to a similar finding by Admon et al. (2013)[25] in a-priori healthy soldiers, in which increased PTSD-related symptoms after-exposure to stressful military experience corresponded to increased amygdala response to risk, both pre- and post-exposure, and to decreased VS response to reward, only post-exposure. Likewise, the amygdala's response to punishment at TP1 here was related to PTSD symptoms both at TP1 and TP3, whereas VS response to reward at TP1 was not associated with symptoms at the same time-point, but was predictive of subsequent symptoms at TP3. Both studies, in line with a suggested casual model of PTSD development[93], suggest that while heightened NVS neural activity represents a predisposing risk factor for this disorder, diminished PVS neural activity may contribute to long-term consequence following the trauma. Furthermore, decreased VS-vmPFC connectivity at TP1 was associated with more severe symptoms at TP3. The VS and vmPFC are two prominent

nodes of the reward circuit, involved in value computations and decision-making processes[94,95]. Human neuroimaging studies have repeatedly demonstrated coincident activation and functional connectivity between these regions[96,97], with animal studies further showing that the vmPFC modulates VS activity[98-100]. Intriguingly, Pujara et al. (2016)[101] reported that vmPFC damage was associated with decreased VS volume and diminished response to reward, further supporting a causal role of the vmPFC for reward processing via the VS. Of particular interest is the relationship found between decreased PVS activation at TP1 to avoidance symptoms at TP3 (beyond total severity). Recent animal studies suggested that dopaminergic neurons in the VS regulate approach-avoidance behavior under goal-conflict situations[102,103]. Comparably in humans, VS activity measured by fMRI discriminated between personality tendencies for approach or avoidance under naturalistic high goal-conflict situations[104].

Looking at the brain's reward system in the context of resilience, PVS activation was shown to mitigate subsequent stress responses to a wide range of stressors in animals and humans[12,13,105,106]. It has been suggested that recruitment of positivity accelerated recovery shortly after stress, by assisting the return to homeostasis[12,13,105-108]. This implies that PVS processing may interrupt the development of stress-related psychopathology, even in the presence of heightened NVS[109]. One mechanism for such an effect has been recently suggested by a

longitudinal study showing that trauma exposure impacted prospective relationships between markers of the reward circuitry function and affective symptom trajectories[110]. Specifically, trauma exposure moderated the prospective relationships between VS and amygdala activations to reward prediction error and hypo/mania severity trajectory[110]. Yet, the association of this interaction to PTSD severity following trauma is unclear.

Altogether it seems that post-traumatic psychopathology might involve neural deficits in the NVS, PVS, or both, which might underlie abnormalities in different key processes, reflected in different clinical phenotypes of this disorder. Utilizing explainable machine learning, we found that VS activity and connectivity in response to rewards vs. punishments early after trauma were more important to the prediction of PTSD development a year later, compared to the amygdala activity and connectivity in response to punishments vs. rewards (Fig. 4f). While PTSD research to date has mostly focused on the NVS (e.g., fear, threat), our findings suggest the important role of the PVS in the risk of developing the disorder in the first year after trauma. Of note, the neural model of specific brain responses to reward and punishment is a simplification of the human positive and negative valence systems, involving multiple brain areas and networks and the interactions between them[1,111,112]. Future studies should examine the neural response using a network perspective or a data-driven whole-brain approach.

Beyond the association of PVS and NVS with overall PTSD symptom severity (i.e., CAPS-5 total scores), our results suggest a relationship between neurobehavioral abnormalities of these systems and specific clusters of PTSD symptoms, alluding to specific underlying mental processes. While NVS hyperactivation was associated mostly with symptom clusters of hyperarousal and intrusion, both at TP1 and TP3 (see Fig. 2 and 4), the PVS hypoactivation was particularly associated with avoidance symptoms (Fig. 4). Hyperarousal and intrusion are well-known as hallmarks of magnified threat processing in PTSD[7,113,114], while avoidance marks deficient reward processing[104,115,116], respectively standing for abnormally intensified NVS and weakened PVS. This finding suggests that the activation of NVS/PVS as early as one month after trauma could point to specific PTSD abnormalities a year later, thus guiding precise and personalized early intervention.

Interestingly, amygdala and VS activations at TP1 did not significantly predict PTSD symptom severity at TP2. This null-result might be explained by the dynamic clinical manifestations during the first year following trauma exposure, displaying an overall progressive reduction in PTSD symptom severity, but large inter-individual variability[117-120]. An intermediary point of 6-months post-trauma might be too early to capture the tangible chronic PTSD subtype, whereas 14-months may portray a more stable representation of the chronic disorder, as it was shown to predict over 90% of the expected recovery from PTSD[121,122]. A

similar trend of null-results at TP2 was observed in a previous analysis of this data set examining neuroanatomical risk factors for PTSD[123].

The possible cognitive mechanism underlying imbalanced NVS/PVS in PTSD

The decrease in risky behavior was associated with both intrusion and avoidance symptoms (corresponding to NVS and PVS processing, respectively) (see Fig. 3), supporting the idea that risky choice index reflects a balance between the two systems. To further explore the relationship between risky choice bias and the neural mechanism of the valence systems, we employed an integrative moderation model. Results showed that a decrease in risky behavior over time (from TP1 to TP2) moderated the relationship between early hyperactive NVS (at TP1) and subsequent post-traumatic stress symptom development (at TP3) (Fig. 5 a&b). It remains to be determined what drives such a behavioral bias following traumatic stress exposure. Speculatively, a possible mechanism involves cognitive flexibility, the ability to change and adapt one's behavior in response to changing environments and stimuli[124,125]. Such flexibility may facilitate goal-directed behavior (e.g., approach or avoidance) that promotes the survival and wellbeing of the organism even in the face of danger[126-128]. Previous work has highlighted the importance of early cognitive flexibility compared to other neurocognitive domains,

as it was shown to be the most significant cognitive predictor of PTSD symptoms at 14-months post-trauma[129,130]. Furthermore, early neurocognitive intervention improved cognitive flexibility which in turn reduced PTSD symptoms, suggesting its role as a modifiable target preceding and underlying the development of post-traumatic psychopathology[130,131]. Hence, enhancement of cognitive flexibility may enable more risky approach behavior toward prospective rewards, possibly by recruitment of the positive system. This idea corresponds to theoretical accounts regarding the importance of PVS in stress recovery by broadening attention and building cognitive and social resources[132,133]. Future research may investigate cognitive flexibility as a potential mechanism of balance between neural PVS and NVS concerning the development of stress-related psychopathology. A common paradigm that captures both reward sensitivity and cognitive flexibility is the probabilistic reversal learning (PRL) paradigm[134-136]. Recent animal work using the PRL paradigm observed that while reward increased flexibility and learning, the presence of aversive punishing stimulus decreased learning performance, increased perseveration, and impaired error detection[137]. As stress is known to impair general learning mechanisms, Harms et al. (2017)[138] showed that stress-exposed adolescents performing the PRL paradigm learned associations of rewards and punishments more slowly (compared to typically developing peers), and showed profound deficits in reversing learned stimulus-response

associations. Finally, cognitive flexibility training improved reversal learning in the PRL paradigm and extinction retention memory (a hallmark cognitive effect of trauma)[139].

Conclusion

This study provides a mechanistic neurobehavioral investigation of PVS and NVS in a large cohort of recent trauma survivors that also takes into consideration dynamics in post-traumatic stress symptoms during the first year following trauma. Results include specific behavioral and neural patterns (activation and connectivity) at one-month after trauma, associated with symptom severity at the same time and predictive of post-trauma psychopathology development over a year later. This work also revealed connections between these neurobehavioral indices and specific symptom clusters, suggesting different NVS/PVS processes underlying different clinical phenotypes of PTSD. As the neurobehavioral mechanisms of the human response to positive and negative outcomes are intrinsically linked, novel therapeutic strategies for PTSD should benefit from addressing symptoms on both fronts. While most interventions and treatments for PTSD focus on reducing the hyperactive negative valence system, strategies aiming to increase the positive valence system may further promote stress resilience and recovery[109].

Acknowledgments: This work was supported by award number R01-MH-103287 from the National Institute of Mental Health (NIMH) given to AS (PI), IL and TH (co-Investigators, subcontractors), and had undergone critical review by the NIMH Adult Psychopathology and Disorders of Aging study section. This work was also supported by funding to the Human Brain Project from the European Union Seventh Framework Program (FP7/2007-2013) under grant agreement no. 604102 (HBP). Sagol School of Neuroscience at Tel-Aviv University, Sagol Brain Institute at Tel-Aviv Sourasky Medical Center, and the Human Brain Project (HBP) from the European Union Seventh Framework Program supported the authors' fellowships. The authors would like to thank the research team at Tel-Aviv Sourasky Medical Center – including Nili Green, Mor Halevi, Sheli Luvton, Yael Shavit, Olga Nevenchannaya, Iris Rashap, Efrat Routledge and Ophir Leshets – for their major contribution in carrying out this research, including subjects' recruitment and screening, and performing clinical and neural assessments. We also want to extend our gratitude to all the participants of this study, who completed all the assessments at three different time-points after experiencing a traumatic event.

Ethics: The study was approved by the ethics committee in the local Medical Center (Reference number 0207/14). All participants gave written informed consent in accordance with the Declaration of Helsinki, and received financial remuneration at the end of each time-point (TP1, TP2 and TP3). The study was registered under ClinicalTrials.gov database

(registration ID#: NCT03756545; https://clinicaltrials.gov/ct2/show/NCT03756545).

Conflict of Interest: The authors declare that they have no financial disclosures and no conflict of interests.

References

1. Tye KM. Neural Circuit Motifs in Valence Processing. *Neuron*. 2018;100(2):436-452. doi:10.1016/j.neuron.2018.10.001

2. Insel T, Cuthbert B, Garvey M, et al. Research Domain Criteria (RDoC): Toward a new classification framework for research on mental disorders. *Am J Psychiatry*. 2010;167(7):748-751. doi:10.1176/appi.ajp.2010.09091379

3. Cuthbert BN, Insel TR. Toward the future of psychiatric diagnosis: The seven pillars of RDoC. *BMC Med*. 2013;11(1):126. doi:10.1186/1741-7015-11-126

4. Reynolds SM, Berridge KC. Emotional environments retune the valence of appetitive versus fearful functions in nucleus accumbens. *Nat Neurosci*. 2008;11(4):423-425. doi:10.1038/nn2061

5. Shabel SJ, Wang C, Monk B, Aronson S, Malinow R. Stress transforms lateral habenula reward responses into punishment signals. *Proc Natl Acad Sci U S A*. 2019;116(25):12488-12493. doi:10.1073/pnas.1903334116

6. Symmonds M, Bossaerts P, Dolan RJ. A behavioral and neural evaluation of prospective decision-making under risk. *J Neurosci*. 2010;30(43):14380-14389. doi:10.1523/JNEUROSCI.1459-10.2010

7. Levy I, Schiller D. Neural Computations of Threat. *Trends Cogn Sci*. 2021;25(2):151-171. doi:10.1016/j.tics.2020.11.007

8. Shields GS, Sazma MA, Yonelinas AP. The effects of acute stress on core executive functions: A meta-analysis and comparison with cortisol. *Neurosci Biobehav Rev*. 2016;68:651-668. doi:10.1016/j.neubiorev.2016.06.038

9. Dreisbach G, Goschke T. How Positive Affect Modulates Cognitive Control: Reduced Perseveration at the Cost of Increased Distractibility. *J Exp Psychol Learn Mem Cogn*. 2004;30(2):343-353. doi:10.1037/0278-7393.30.2.343

10. Goschke T. Voluntary action and cognitive control from a cognitive neuroscience perspective. In: *Voluntary Action: Brains, Minds, and Sociality.* ; 2003:49-85.

11. Hommel B. Between Persistence and Flexibility. *Adv Motiv Sci*.

2015:33–67. doi:10.1016/bs.adms.2015.04.003

12. Speer ME, Mauricio R. Reminiscing about positive memories buffers acute stress responses. *Nat Hum Behav*. 2017;1(5). doi:10.1038/s41562-017-0093

13. Tugade MM, Fredrickson BL. Resilient Individuals Use Positive Emotions to Bounce Back From Negative Emotional Experiences. *J Pers Soc Psychol*. 2004;86(2):320–333. doi:10.1037/0022-3514.86.2.320

14. McNaughton N, Corr PJ. The neuropsychology of fear and anxiety: A foundation for Reinforcement Sensitivity Theory. In: *The Reinforcement Sensitivity Theory of Personality*. ; 2008:44–94. doi:10.1017/CBO9780511819384.003

15. Stein MB, Paulus MP. Imbalance of Approach and Avoidance: The Yin and Yang of Anxiety Disorders. *Biol Psychiatry*. 2009;66(12):1072–1074. doi:10.1016/j.biopsych.2009.09.023

16. Weaver SS, Kroska EB, Ross MC, et al. Sacrificing reward to avoid threat: Characterizing PTSD in the context of a trauma-related approach-avoidance conflict task. *J Abnorm Psychol*. 2020;129(5):457–468. doi:10.1037/abn0000528

17. Aupperle RL, Paulus MP. Neural systems underlying approach and avoidance in anxiety disorders. *Dialogues Clin Neurosci*. 2010;12(4):517–531. doi:10.31887/dcns.2010.12.4/raupperle

18. Hayes JP, Hayes SM, Mikedis AM, et al. Quantitative meta-analysis of neural activity in posttraumatic stress disorder. *Biol Mood Anxiety Disord*. 2012;2(1):9. doi:10.1186/2045-5380-2-9

19. Hayes JP, VanElzakker MB, Shin LM. Emotion and Cognition Interactions in PTSD: A Review of Neurocognitive and Neuroimaging Studies. *Front Integr Neurosci*. 2012;6:89. doi:10.3389/fnint.2012.00089

20. Ronzoni G, del Arco A, Mora F, Segovia G. Enhanced noradrenergic activity in the amygdala contributes to hyperarousal in an animal model of PTSD. *Psychoneuroendocrinology*. 2016;70:1–9. doi:10.1016/j.psyneuen.2016.04.018

21. Bruce SE, Buchholz KR, Brown WJ, Yan L, Durbin A, Sheline YI. Altered emotional interference processing in the amygdala and

insula in women with Post-Traumatic Stress Disorder. *NeuroImage Clin.* 2013;2(1):43-49. doi:10.1016/j.nicl.2012.11.003

22. Hayes JP, LaBar KS, McCarthy G, et al. Reduced hippocampal and amygdala activity predicts memory distortions for trauma reminders in combat-related PTSD. *J Psychiatr Res.* 2011;45(5):660-669. doi:10.1016/j.jpsychires.2010.10.007

23. Forster GL, Simons RM, Baugh LA. Revisiting the Role of the Amygdala in Posttraumatic Stress Disorder. In: *The Amygdala - Where Emotions Shape Perception, Learning and Memories.* ; 2017. doi:10.5772/67585

24. Jia R, Ruderman L, Gordon C, et al. A Shift from Value- to Saliency-Neural Encoding of Subjective Value in Combat Veterans with PTSD during Decision Making under Uncertainty. *bioRxiv.* 2020:2020.04.14.041467. doi:10.1101/2020.04.14.041467

25. Admon R, Lubin G, Rosenblatt JD, et al. Imbalanced neural responsivity to risk and reward indicates stress vulnerability in humans. *Cereb Cortex.* 2013;23(1):28-35. doi:10.1093/cercor/bhr369

26. Ruderman L, Ehrlich DB, Roy A, Pietrzak RH, Harpaz-Rotem I, Levy I. Posttraumatic Stress Symptoms and Aversion To Ambiguous Losses in Combat Veterans. *Depress Anxiety.* 2016;33(7):606-613. doi:10.1002/da.22494

27. Zuj D V., Palmer MA, Lommen MJJ, Felmingham KL. The centrality of fear extinction in linking risk factors to PTSD: A narrative review. *Neurosci Biobehav Rev.* 2016;69:15-35. doi:10.1016/j.neubiorev.2016.07.014

28. LeDoux JE. Emotion circuits in the brain. *Annu Rev Neurosci.* 2000;23:155-184. doi:10.1146/annurev.neuro.23.1.155

29. Graham BM, Milad MR. The study of fear extinction: Implications for anxiety disorders. *Am J Psychiatry.* 2011;168(12):1255-1265. doi:10.1176/appi.ajp.2011.11040557

30. Rabellino D, Densmore M, Frewen PA, Théberge J, McKinnon MC, Lanius RA. Aberrant functional connectivity of the amygdala complexes in PTSD during conscious and subconscious processing of trauma-related stimuli. *PLoS One.* 2016.

doi:10.1371/journal.pone.0163097

31. Gilboa A, Shalev AY, Laor L, et al. Functional connectivity of the prefrontal cortex and the amygdala in posttraumatic stress disorder. *Biol Psychiatry*. 2004. doi:10.1016/j.biopsych.2003.08.004

32. Stevens JS, Jovanovic T, Fani N, et al. Disrupted amygdala-prefrontal functional connectivity in civilian women with posttraumatic stress disorder. *J Psychiatr Res*. 2013. doi:10.1016/j.jpsychires.2013.05.031

33. Ben-Zion Z, Zeevi Y, Keynan NJ, et al. Multi-domain potential biomarkers for post-traumatic stress disorder (PTSD) severity in recent trauma survivors. *Transl Psychiatry*. 2020. doi:10.1038/s41398-020-00898-z

34. Nawijn L, van Zuiden M, Frijling JL, Koch SBJ, Veltman DJ, Olff M. Reward functioning in PTSD: A systematic review exploring the mechanisms underlying anhedonia. *Neurosci Biobehav Rev*. 2015;51:189-204. doi:10.1016/j.neubiorev.2015.01.019

35. Seidemann R, Duek O, Jia R, Levy I, Harpaz-Rotem I. The Reward System and Post-Traumatic Stress Disorder: Does Trauma Affect the Way We Interact With Positive Stimuli? *Chronic Stress*. 2021;5:247054702199600. doi:10.1177/2470547021996006

36. Ikemoto S. Brain reward circuitry beyond the mesolimbic dopamine system: A neurobiological theory. *Neurosci Biobehav Rev*. 2010;35(2):129-150. doi:10.1016/j.neubiorev.2010.02.001

37. Nikolova YS, Bogdan R, Brigidi BD, Hariri AR. Ventral striatum reactivity to reward and recent life stress interact to predict positive affect. *Biol Psychiatry*. 2012;72(2):157-163. doi:10.1016/j.biopsych.2012.03.014

38. Pizzagalli DA, Holmes AJ, Dillon DG, et al. Reduced caudate and nucleus accumbens response to rewards in unmedicated individuals with major depressive disorder. *Am J Psychiatry*. 2009;166(6):702-710. doi:10.1176/appi.ajp.2008.08081201

39. Stoy M, Schlagenhauf F, Sterzer P, et al. Hyporeactivity of ventral striatum towards incentive stimuli in unmedicated depressed patients normalizes after treatment with escitalopram. *J Psychopharmacol*. 2012;26(5):677-688.

doi:10.1177/0269881111416686

40. Elman I, Lowen S, Frederick BB, Chi W, Becerra L, Pitman RK. Functional Neuroimaging of Reward Circuitry Responsivity to Monetary Gains and Losses in Posttraumatic Stress Disorder. *Biol Psychiatry*. 2009;66(12):1083–1090. doi:10.1016/j.biopsych.2009.06.006

41. Sailer U, Robinson S, Fischmeister FPS, et al. Altered reward processing in the nucleus accumbens and mesial prefrontal cortex of patients with posttraumatic stress disorder. *Neuropsychologia*. 2008;46(11):2836–2844. doi:10.1016/j.neuropsychologia.2008.05.022

42. Felmingham KL, Falconer EM, Williams L, et al. Reduced amygdala and ventral striatal activity to happy faces in PTSD is associated with emotional numbing. *PLoS One*. 2014;9(9). doi:10.1371/journal.pone.0103653

43. Mehta ND, Stevens JS, Li Z, et al. Inflammation, reward circuitry and symptoms of anhedonia and PTSD in trauma-exposed women. *Soc Cogn Affect Neurosci*. 2020. doi:10.1093/scan/nsz100

44. Zhu X, Helpman L, Papini S, et al. Altered resting state functional connectivity of fear and reward circuitry in comorbid PTSD and major depression. *Depress Anxiety*. 2017. doi:10.1002/da.22594

45. American Psychiatric Association, Association AP. *Diagnostic And Statistical Manual of Mental Disorders : DSM-5.* American Psychiatric Association; 2013. https://ci.nii.ac.jp/naid/10010095609/.

46. Rab SL, Admon R. Parsing inter- and intra-individual variability in key nervous system mechanisms of stress responsivity and across functional domains. *Neurosci Biobehav Rev*. 2020. doi:10.1016/j.neubiorev.2020.09.007

47. Galatzer-Levy IR, Bryant RA. 636,120 Ways to Have Posttraumatic Stress Disorder. *Perspect Psychol Sci*. 2013;8(6):651–662. doi:10.1177/1745691613504115

48. Young G, Lareau C, Pierre B. One Quintillion Ways to Have PTSD Comorbidity: Recommendations for the Disordered DSM-5. *Psychol Inj Law*. 2014;7(1):61–74. doi:10.1007/s12207-014-9186-y

49. Pitman RK, Rasmusson AM, Koenen KC, et al. Biological studies of post-traumatic stress disorder. *Nat Rev Neurosci.* 2012;13(11):769-787. doi:10.1038/nrn3339

50. Zoladz P, Diamond D. Current status on behavioral and biological markers of PTSD. *Neurosci Biobehav Rev.* 2013.

51. Ben-Zion Z, Fine NB, Keynan NJ, et al. Neurobehavioral Moderators of Post-Traumatic Stress Disorder Trajectories: Prospective fMRI Study of Recent Trauma Survivors. *Eur J Psychotraumatol.* 2019;10(1). doi:10.1080/20008198.2019.1683941

52. Shalev AY, Gevonden M, Ratanatharathorn A, et al. Estimating the risk of PTSD in recent trauma survivors: results of the International Consortium to Predict PTSD (ICPP). *World Psychiatry.* 2019;18(1):77-87. doi:10.1002/wps.20608

53. Shalev AY, Ankri Y, Israeli-Shalev Y, Peleg T, Adessky R, Freedman S. Prevention of posttraumatic stress disorder by early treatment: Results from the Jerusalem trauma outreach and prevention study. *Arch Gen Psychiatry.* 2012;69(2):166-176. doi:10.1001/archgenpsychiatry.2011.127

54. Kahn I, Yeshurun Y, Rotshtein P, Fried I, Ben-Bashat D, Hendler T. The role of the amygdala in signaling prospective outcome of choice. *Neuron.* 2002;33(6):983-994.

55. Assaf M, Kahn I, Pearlson GD, et al. Brain activity dissociates mentalization from motivation during an interpersonal competitive game. *Brain Imaging Behav.* 2009;3(1):24-37. doi:10.1007/s11682-008-9047-y

56. Gonen T, Admon R, Podlipsky I, Hendler T. From animal model to human brain networking: Dynamic causal modeling of motivational systems. *J Neurosci.* 2012;32(21):7218-7224. doi:10.1523/JNEUROSCI.6188-11.2012

57. Thaler A, Gonen T, Mirelman A, et al. Altered reward-related neural responses in non-manifesting carriers of the Parkinson disease related LRRK2 mutation. *Brain Imaging Behav.* 2019;13(4):1009-1020. doi:10.1007/s11682-018-9920-2

58. Admon R, Lubin G, Stern O, et al. Human vulnerability to stress depends on amygdala's predisposition and hippocampal plasticity.

Proc Natl Acad Sci. 2009;106(33):14120-14125. doi:10.1073/pnas.0903183106

59. Blanchard EB, Jones-Alexander J, Buckley TC, Forneris CA. Psychometric properties of the PTSD checklist (PCL). *Behav Res Ther*. 1996;34(8):669-673. doi:10.1016/0005-7967(96)00033-2

60. Weathers FW, Bovin MJ, Lee DJ, et al. The Clinician-Administered PTSD Scale for DSM-5 (CAPS-5): Development and initial psychometric evaluation in military veterans. *Psychol Assess*. 2018;30(3):383-395. doi:10.1037/pas0000486

61. Hyatt CJ, Assaf M, Muska CE, et al. Reward-related dorsal striatal activity differences between former and current cocaine dependent individuals during an interactive competitive game. *PLoS One*. 2012;7(5):1-15. doi:10.1371/journal.pone.0034917

62. Esteban O, Markiewicz CJ, Blair RW, et al. fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nat Methods*. 2019;16(1):111-116. doi:10.1038/s41592-018-0235-4

63. Gorgolewski K, Burns CD, Madison C, et al. Nipype: A flexible, lightweight and extensible neuroimaging data processing framework in Python. *Front Neuroinform*. 2011;5. doi:10.3389/fninf.2011.00013

64. Fan L, Li H, Zhuo J, et al. The Human Brainnetome Atlas: A New Brain Atlas Based on Connectional Architecture. *Cereb Cortex*. 2016;26(8):3508-3526. doi:10.1093/cercor/bhw157

65. Pauli WM, Nili AN, Michael Tyszka J. Data Descriptor: A high-resolution probabilistic in vivo atlas of human subcortical brain nuclei. *Sci Data*. 2018;5. doi:10.1038/sdata.2018.63

66. Brett M, Anton JL, Valabregue R, Poline JB. Region of interest analysis using an SPM toolbox. *Neuroimage*. 2002.

67. McLaren DG, Ries ML, Xu G, Johnson SC. A generalized form of context-dependent psychophysiological interactions (gPPI): A comparison to standard approaches. *Neuroimage*. 2012. doi:10.1016/j.neuroimage.2012.03.068

68. Whitfield-Gabrieli S, Nieto-Castanon A. Conn: A Functional Connectivity Toolbox for Correlated and Anticorrelated Brain Networks. *Brain Connect*. 2012;2(3):125-141.

doi:10.1089/brain.2012.0073

69. Cheng W, Rolls ET, Qiu J, et al. Medial reward and lateral non-reward orbitofrontal cortex circuits change in opposite directions in depression. *Brain*. 2016;139(12):3296–3309. doi:10.1093/brain/aww255

70. Kim SH, Yoon HS, Kim H, Hamann S. Individual differences in sensitivity to reward and punishment and neural activity during reward and avoidance learning. *Soc Cogn Affect Neurosci*. 2014;10(9):1219–1227. doi:10.1093/scan/nsv007

71. Yacubian J, Gläscher J, Schroeder K, Sommer T, Braus DF, Büchel C. Dissociable systems for gain- and loss-related value predictions and errors of prediction in the human brain. *J Neurosci*. 2006;26(37):9530–9537. doi:10.1523/JNEUROSCI.2915-06.2006

72. Elliott R, Dolan RJ, Frith CD. Dissociable functions in the medial and lateral orbitofrontal cortex: Evidence from human neuroimaging studies. *Cereb Cortex*. 2000;10(3):308–317. doi:10.1093/cercor/10.3.308

73. O'Doherty J, Kringelbach ML, Rolls ET, Hornak J, Andrews C. Abstract reward and punishment representations in the human orbitofrontal cortex. *Nat Neurosci*. 2001;4(1):95–102. doi:10.1038/82959

74. IBM Corp. IBM SPSS Statistics for Windows, Version 26.0. *2019*. 2019.

75. R Core Team (2020). R: A language and environment for statistical computing. *R A Lang Environ Stat Comput R Found Stat Comput Vienna, Austria*. 2020.

76. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J R Stat Soc Ser B*. 1995;57(1):289–300. doi:10.1111/j.2517-6161.1995.tb02031.x

77. Lundberg SM, Lee SI. A unified approach to interpreting model predictions. In: *Advances in Neural Information Processing Systems*. Vol 2017-Decem. ; 2017:4766–4775.

78. Preacher KJ, Hayes AF. Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behav Res Methods*. 2008;40(3):879–891.

doi:10.3758/BRM.40.3.879

79. Hayes AF. *Introduction to Mediation, Moderation and Conditional Process Analysis*. Vol 53.; 2013. doi:10.5539/ass.v11n9p207

80. Liberzon I, Sripada CS. The functional neuroanatomy of PTSD: a critical review. *Prog Brain Res*. 2007;167:151–169. doi:10.1016/S0079-6123(07)67011-3

81. Shin LM, Liberzon I. The Neurocircuitry of Fear, Stress and Anxiety Disorders. *Neuropsychopharmacology*. 2010;35(1):169–191. doi:10.1038/npp.2009.83

82. Amiel Rosenkranz J, Grace AA. Cellular mechanisms of infralimbic and prelimbic prefrontal cortical inhibition and dopaminergic modulation of basolateral amygdala neurons in vivo. *J Neurosci*. 2002;22(1):324–337. doi:10.1523/jneurosci.22-01-00324.2002

83. Quirk GJ, Likhtik E, Pelletier JG, Paré D. Stimulation of medial prefrontal cortex decreases the responsiveness of central amygdala output neurons. *J Neurosci*. 2003;23(25):8800–8807. doi:10.1523/jneurosci.23-25-08800.2003

84. Phan KL, Fitzgerald DA, Nathan PJ, Moore GJ, Uhde TW, Tancer ME. Neural substrates for voluntary suppression of negative affect: A functional magnetic resonance imaging study. *Biol Psychiatry*. 2005;57(3):210–219. doi:10.1016/j.biopsych.2004.10.030

85. Garcia R, Vouimba RM, Baudry M, Thompson RF. The amygdala modulates prefrontal cortex activity relative to conditioned fear. *Nature*. 1999;402(6759):294–296. doi:10.1038/46286

86. Shin LM, Wright CI, Cannistraro PA, et al. A functional magnetic resonance imaging study of amygdala and medial prefrontal cortex responses to overtly presented fearful faces in posttraumatic stress disorder. *Arch Gen Psychiatry*. 2005;62(3):273–281. doi:10.1001/archpsyc.62.3.273

87. Williams LM, Kemp AH, Felmingham K, et al. Trauma modulates amygdala and medial prefrontal responses to consciously attended fear. *Neuroimage*. 2006;29(2):347–357. doi:10.1016/j.neuroimage.2005.03.047

88. Hahn A, Stein P, Windischberger C, et al. Reduced resting-state functional connectivity between amygdala and orbitofrontal cortex

in social anxiety disorder. *Neuroimage*. 2011;56(3):881–889. doi:10.1016/j.neuroimage.2011.02.064

89. Monk CS, Telzer EH, Mogg K, et al. Amygdala and ventrolateral prefrontal cortex activation to masked angry faces in children and adolescents with generalized anxiety disorder. *Arch Gen Psychiatry*. 2008;65(5):568–576. doi:10.1001/archpsyc.65.5.568

90. Frodl T, Bokde ALW, Scheuerecker J, et al. Functional Connectivity Bias of the Orbitofrontal Cortex in Drug-Free Patients with Major Depression. *Biol Psychiatry*. 2010;67(2):161–167. doi:10.1016/j.biopsych.2009.08.022

91. Drevets WC, Price JL, Furey ML. Brain structural and functional abnormalities in mood disorders: Implications for neurocircuitry models of depression. *Brain Struct Funct*. 2008;213(1–2):93–118. doi:10.1007/s00429-008-0189-x

92. Versace A, Thompson WK, Zhou D, et al. Abnormal Left and Right Amygdala-Orbitofrontal Cortical Functional Connectivity to Emotional Faces: State Versus Trait Vulnerability Markers of Depression in Bipolar Disorder. *Biol Psychiatry*. 2010;67(5):422–431. doi:10.1016/j.biopsych.2009.11.025

93. Admon R, Milad MR, Hendler T. A causal model of post-traumatic stress disorder: disentangling predisposed from acquired neural abnormalities. *Trends Cogn Sci*. 2013;17(7):337–347. doi:10.1016/J.TICS.2013.05.005

94. Pauli WM, O'Reilly RC, Yarkoni T, Wager TD. Regional specialization within the human striatum for diverse psychological functions. *Proc Natl Acad Sci U S A*. 2016;113(7):1907–1912. doi:10.1073/pnas.1507610113

95. Clithero JA, Rangel A. Informatic parcellation of the network involved in the computation of subjective value. *Soc Cogn Affect Neurosci*. 2013;9(9):1289–1302. doi:10.1093/scan/nst106

96. Cauda F, Cavanna AE, D'agata F, Sacco K, Duca S, Geminiani GC. Functional connectivity and coactivation of the nucleus accumbens: A combined functional connectivity and structure-based meta-analysis. *J Cogn Neurosci*. 2011;23(10):2864–2877. doi:10.1162/jocn.2011.21624

97. Choi EY, Thomas Yeo BT, Buckner RL. The organization of the human striatum estimated by intrinsic functional connectivity. *J Neurophysiol.* 2012;108(8):2242-2263. doi:10.1152/jn.00270.2012

98. Voorn P, Vanderschuren LJMJ, Groenewegen HJ, Robbins TW, Pennartz CMA. Putting a spin on the dorsal-ventral divide of the striatum. *Trends Neurosci.* 2004;27(8):468-474. doi:10.1016/j.tins.2004.06.006

99. Gabbott PLA, Warner TA, Jays PRL, Salway P, Busby SJ. Prefrontal cortex in the rat: Projections to subcortical autonomic, motor, and limbic centers. *J Comp Neurol.* 2005;492(2):145-177. doi:10.1002/cne.20738

100. Ghazizadeh A, Ambroggi F, Odean N, Fields HL. Prefrontal cortex mediates extinction of responding by two distinct neural mechanisms in accumbens shell. *J Neurosci.* 2012;32(2):726-737. doi:10.1523/JNEUROSCI.3891-11.2012

101. Pujara MS, Philippi CL, Motzkin JC, Baskaya MK, Koenigs M. Ventromedial prefrontal cortex damage is associated with decreased ventral striatum volume and response to reward. *J Neurosci.* 2016;36(18):5047-5054. doi:10.1523/JNEUROSCI.4236-15.2016

102. Nguyen D, Fugariu V, Erb S, Ito R. Dissociable roles of the nucleus accumbens D1 and D2 receptors in regulating cue-elicited approach-avoidance conflict decision-making. *Psychopharmacology (Berl).* 2018;235(8):2233-2244. doi:10.1007/s00213-018-4919-3

103. Nguyen D, Alushaj E, Erb S, Ito R. Dissociative effects of dorsomedial striatum D1 and D2 receptor antagonism in the regulation of anxiety and learned approach-avoidance conflict decision-making. *Neuropharmacology.* 2019;146:222-230. doi:10.1016/j.neuropharm.2018.11.040

104. Gonen T, Soreq E, Eldar E, Ben-Simon E, Raz G, Hendler T. Human mesostriatal response tracks motivational tendencies under naturalistic goal conflict. *Soc Cogn Affect Neurosci.* 2016;11(6):961-972. doi:10.1093/scan/nsw014

105. Fredrickson BL, Levenson RW. Positive Emotions Speed Recovery from the Cardiovascular Sequelae of Negative Emotions. *Cogn Emot.*

1998. doi:10.1080/026999398379718

106. Fredrickson BL, Mancuso RA, Branigan C, Tugade MM. The undoing effect of positive emotions. *Motiv Emot*. 2000. doi:10.1023/A:1010796329158

107. Hughes AA, Kendall PC. Effect of a Positive Emotional State on Interpretation Bias for Threat in Children With Anxiety Disorders. *Emotion*. 2008. doi:10.1037/1528-3542.8.3.414

108. Falkenstern M, Schiffrin HH, Nelson SK, Ford L, Keyser C. Mood over matter: Can happiness be your undoing? *J Posit Psychol*. 2009. doi:10.1080/17439760902992415

109. Dutcher JM, Creswell JD. The role of brain reward pathways in stress resilience and health. *Neurosci Biobehav Rev*. 2018;95:559-567. doi:10.1016/j.neubiorev.2018.10.014

110. Eckstrand KL, Forbes EE, Bertocci MA, et al. Trauma impacts prospective relationships between reward-related ventral striatal and amygdala activation and 1-year future hypo/mania trajectories. *Biol Psychiatry*. November 2020. doi:10.1016/j.biopsych.2020.11.017

111. Berridge KC. Affective valence in the brain: modules or modes? *Nat Rev Neurosci*. 2019;20(4):225-234. doi:10.1038/s41583-019-0122-8

112. Baxter MG, Murray EA. The amygdala and reward. *Nat Rev Neurosci*. 2002;3(7):563-573. doi:10.1038/nrn875

113. Rabellino D, Densmore M, Frewen PA, Théberge J, Lanius RA. The innate alarm circuit in post-traumatic stress disorder: Conscious and subconscious processing of fear- and trauma-related cues. *Psychiatry Res - Neuroimaging*. 2016;248:142-150. doi:10.1016/j.pscychresns.2015.12.005

114. Lanius RA, Rabellino D, Boyd JE, Harricharan S, Frewen PA, McKinnon MC. The innate alarm system in PTSD: conscious and subconscious processing of threat. *Curr Opin Psychol*. 2017;14:109-115. doi:10.1016/j.copsyc.2016.11.006

115. Romanczuk-Seiferth N, Koehler S, Dreesen C, Wüstenberg T, Heinz A. Pathological gambling and alcohol dependence: Neural disturbances in reward and loss avoidance processing. *Addict Biol*.

2015;20(3):557–569. doi:10.1111/adb.12144

116. Simon JJ, Walther S, Fiebach CJ, et al. Neural reward processing is modulated by approach- and avoidance-related personality traits. *Neuroimage*. 2010;49(2):1868–1874. doi:10.1016/j.neuroimage.2009.09.016

117. Hepp U, Moergeli H, Buchi S, et al. Post-traumatic stress disorder in serious accidental injury: 3-Year follow-up study. *Br J Psychiatry*. 2008;192(5):376–383. doi:10.1192/bjp.bp.106.030569

118. Yehuda R, McFarlane AC, Shalev AY. Predicting the development of posttraumatic stress disorder from the acute response to a traumatic event. *Biol Psychiatry*. 1998;44(12):1305–1313. doi:10.1016/S0006-3223(98)00276-5

119. Koren D, Arnon I, Klein E. Long term course of chronic posttraumatic stress disorder in traffic accident victims: A three-year prospective follow-up study. *Behav Res Ther*. 2001;39(12):1449–1458. doi:10.1016/S0005-7967(01)00025-0

120. Perkonigg A, Pfister H, Stein MB, et al. Longitudinal course of posttraumatic stress disorder and posttraumatic stress disorder symptoms in a community sample of adolescents and young adults. *Am J Psychiatry*. 2005;162(7):1320–1327. doi:10.1176/appi.ajp.162.7.1320

121. Shalev AY, Freedman S. PTSD Following Terrorist Attacks: A Prospective Evaluation. *Am J Psychiatry*. 2005;162(6):1188–1191. doi:10.1176/appi.ajp.162.6.1188

122. Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson CB. Posttraumatic Stress Disorder in the National Comorbidity Survey. *Arch Gen Psychiatry*. 1995;52(12):1048–1060. doi:10.1001/archpsyc.1995.03950240066012

123. Ben-Zion Z, Artzi M, Niry D, et al. Neuroanatomical Risk Factors for Post Traumatic Stress Disorder (PTSD) in Recent Trauma Survivors. *Biol Psychiatry Cogn Neurosci Neuroimaging*. 2019:721134. doi:10.1101/721134

124. Cañas JJ, Quesada JF, Antolí A, Fajardo I. Cognitive flexibility and adaptability to environmental changes in dynamic complex problem-solving tasks. *Ergonomics*. 2003;46(5):482–501.

doi:10.1080/0014013031000061640

125. Cañ JJ, Antoli A, Fajardo I, Salmerón L. Cognitive inflexibility and the development and use of strategies for solving complex dynamic problems: Effects of different types of training. *Theor Issues Ergon Sci*. 2005;6(1):95–108. doi:10.1080/14639220512331311599

126. Berridge KC, Kringelbach ML. Neuroscience of affect: Brain mechanisms of pleasure and displeasure. *Curr Opin Neurobiol*. 2013;23(3):294–303. doi:10.1016/j.conb.2013.01.017

127. Berridge KC, Kringelbach ML. Pleasure Systems in the Brain. *Neuron*. 2015;86(3):646–664. doi:10.1016/j.neuron.2015.02.018

128. Lang PJ, Bradley MM. Emotion and the motivational brain. *Biol Psychol*. 2010;84(3):437–450. doi:10.1016/j.biopsycho.2009.10.007

129. Ben-Zion Z, Zeevi Y, Keynan NJ, et al. Potential Neurocognitive Biomarkers for Post Traumatic Stress Disorder (PTSD) severity in recent trauma survivors. *bioRxiv*. 2019. doi:10.1101/721068

130. Ben-Zion Z, Fine NB, Keynan NJ, et al. Cognitive Flexibility Predicts PTSD Symptoms: Observational and Interventional Studies. *Front Psychiatry*. 2018;9:477. doi:10.3389/fpsyt.2018.00477

131. Joseph J, Moring J, Bira L. Cognitive Flexibility as a Key Factor in the Conceptualization and Treatment of PTSD. *Curr Psychiatry Rev*. 2015. doi:10.2174/1573400511666150629104921

132. Fredrickson BL. What good are positive emotions? *Rev Gen Psychol*. 1998. doi:10.1037/1089-2680.2.3.300

133. Fredrickson BL, Tugade MM, Waugh CE, Larkin GR. What good are positive emotions in crisis? *J Pers Soc Psychol*. 2003.

134. Cools R, Clark L, Owen AM, Robbins TW. Defining the Neural Mechanisms of Probabilistic Reversal Learning Using Event-Related Functional Magnetic Resonance Imaging. *J Neurosci*. 2002;22(11):4563–4567. doi:10.1523/jneurosci.22-11-04563.2002

135. Clark L, Robbins TW. Decision-making deficits in drug addiction. *Trends Cogn Sci*. 2002;6(9):361–363. doi:10.1016/S1364-6613(02)01960-5

136. Lesage E, Aronson SE, Sutherland MT, Ross TJ, Salmeron BJ, Stein

EA. Neural signatures of cognitive flexibility and reward sensitivity following nicotinic receptor stimulation in dependent smokers a randomized trial. *JAMA Psychiatry*. 2017;74(6):632–640. doi:10.1001/jamapsychiatry.2017.0400

137. Jager A, Dam SA, Van Der Mierden S, et al. Modulation of cognitive flexibility by reward and punishment in BALB/cJ and BALB/cByJ mice. *Behav Brain Res*. 2020;378:112294. doi:10.1016/j.bbr.2019.112294

138. Harms MB, Shannon Bowen KE, Hanson JL, Pollak SD. Instrumental learning and cognitive flexibility processes are impaired in children exposed to early life stress. *Dev Sci*. 2018;21(4):e12596. doi:10.1111/desc.12596

139. Chaby LE, Karavidha K, Lisieski MJ, Perrine SA, Liberzon I. Cognitive flexibility training improves extinction retention memory and enhances cortical dopamine with and without traumatic stress exposure. *Front Behav Neurosci*. 2019;13:24. doi:10.3389/fnbeh.2019.00024

Tables

Table 1. **Participants' demographic and clinical characteristics.** The table summarizes characteristics of participants included in the final analyses across the three time-points. Means and standard deviations of participants' age, gender (Female: Male), and PTSD severity (CAPS-5 total scores), at 1-, 6- and 14-months post-trauma (TP1, TP2 and TP3). Additionally, the percentage of motor-vehicle accidents of individuals diagnosed with PTSD (%MVA's, %PTSD) are reported for each time-point separately.

Measure	TP1 (n=132)		TP2 (n=115)		TP3 (n=112)	
	M	*SD*	*M*	*SD*	*M*	*SD*
Age	33.52	11.01	33.73	11.08	33.56	11.27
Gender (F:M)	63:69	–	55:60	–	56:56	–
CAPS-5 Total	24.91	11.68	14.97	10.89	10.69	10.10
% MVA's	89% (n=117)		88% (n=101)		88% (n=99)	
% PTSD	74% (n=97)		35% (n=40)		24% (n=27)	

Figures Titles and Legends

Figure 1. Safe or Risky Domino Choice (SRDC) Paradigm. Each round of the game is composed of four intervals. First, the participant choose which chip to play next (i.e., decision-making), either a matching choice (e.g., a chip with least one of the master chip's numbers) or a non-matching choice. Next, he moves the cursor to the chosen chip and places it facing down adjacent to the master chip (i.e., decision-execution). The participant then waits for the opponent's response (i.e., anticipation of an outcome) and sees whether the opponent challenges

this choice by uncovering the chosen chip or not (i.e., response to outcome). Participant's choices and opponent's responses are interactively determined by the flow of the game round after round, creating a natural progression of a game situation that lasts 4 min or until the player wins the game by disposing of all his chips. Each player played consecutively for 14 min (approximately 3–4 game rounds).

Figure 2. Neural Indicators of Positive and Negative Valence Systems for Post-Traumatic Stress Symptom Severity Shortly after Trauma Exposure. a. Partial regression scatter plots depicting the relation between CAPS-5 total scores at TP1 (y-axis) and mean beta values for the left and right amygdala activation in response to punishments vs. rewards (x-axis). Values on axes are unstandardized residuals. The anatomical amygdala ROI which was used for this analysis is presented on a coronal view of the brain (in red). Each dot represents one subject. **b.** Partial regression scatter plots depicting the relation between CAPS-5 total scores at TP1 (y-axis) and mean beta values for the left and right ventral striatum activation in response to rewards vs. punishments (a-axis). Values on axes are unstandardized residuals. The anatomical ventral striatum ROI which was used for this analysis is presented on a coronal view of the brain (in green). Each dot represents one subject. **c.** Partial regression scatter plots depicting the relation between CAPS-5 total scores at TP1 (y-axis) and mean beta values for the right amygdala – left lateral OFC

connectivity in response to punishments vs. rewards at TP1 (x-axis). Values on both axes are unstandardized residuals. The anatomical ROIs which were used for this analysis, right amygdala (red) and left lateral OFC (violet), are presented on an axial view of the brain. Each asterisk represents one subject.

Figure 3. Behavioral Indicators of Positive and Negative Valence Systems for Post-Traumatic Stress Symptom Severity Shortly after Trauma Exposure. a. Boxplots with individual data points representing average risky choice index (%) for individuals which met PTSD diagnosis at TP1 and those who did not ('PTSD', n=97, dark gray; 'No PTSD', n=35, light gray). **b.** Partial regression plot depicting the relation between individuals' risky choice indexes at TP1 (%, x-axis) and their total CAPS-5 scores (y-axis) at TP1, while controlling for effects of age, gender and trauma type (covariates). Values on both axes are unstandardized residuals. On the left, a bar plot presenting the correlations between risky choice index and all four PTSD symptom clusters at TP1 according to CAPS-5: intrusion (B), avoidance (C), negative alterations in cognition and mood (D), and hyperarousal symptoms (E). Pearson correlation coefficients (r) are presented above each bar. *p-FDR<0.05.

Figure 4. Neural abnormalities of PVS and NVS predict first-year PTSD development. a. Partial regression scatter plot depicting the relation

between CAPS-5 total scores at TP3 (y-axis) and mean beta values for the left amygdala activation in response to punishments vs. rewards at TP1 (x-axis), while controlling for effects of age, gender and trauma type (covariates). Values on both axes are unstandardized residuals. On the left, a bar plot presenting the correlations between left amygdala activation and all four PTSD symptom clusters at TP3 according to CAPS-5: intrusion (B), avoidance (C), negative alterations in cognition and mood (D), and hyperarousal symptoms (E). Pearson correlation coefficients (r) are presented above each bar. *p-FDR<0.05. **b.** Partial regression scatter plot depicting the relation between CAPS-5 total scores at TP3 (y-axis) and mean beta values for the right ventral striatum activation in response to rewards vs. punishments at TP1 (x-axis), while controlling for covariates (see in a). On the left, a bar plot presenting the correlations between right ventral striatum's activation and all four PTSD symptom clusters at TP1 according to CAPS-5. Pearson correlation coefficients (r) are presented above each bar. *p-FDR<0.05. **c.** Partial regression scatter plot depicting the relation between CAPS-5 total scores at TP3 (y-axis) and mean beta values for the right ventral striatum – right vmPFC connectivity in response to rewards vs. punishments at TP1 (x-axis), while controlling for covariates (see in a). The corresponding predefined anatomical ROIs, right VS (green) and right vmPFC (yellow), are presented next to the plot. **d. Explainable machine learning.** On the right, Absolute feature importance as calculated by SHAP, pointing to the importance of

neural features at TP1 in predicting CAPS-5 total scores at TP3. The larger the SHAP value, the more important the feature is to discriminate between individuals with different symptom severity (CAPS-5 total scores). On the left, SHAP importance summary dot plot displaying features that influenced the linear regression model predictions of PTSD symptom severity (CAPS-5 total scores) at TP3. Features are first sorted by their global impact (y-axis). For every individual from the n=105 included in our sample, a dot represents the attribution value for each feature from low (blue) to high (red).

Figure 5. Integrative modeling of brain, behavior and PTSD development. **a.** Integrative moderation model of neural NVS, behavior and clinical symptoms: behavioral change in risky choice index (from TP1 to TP2) significantly moderated the relation between valence specific left amygdala activation at TP1 and PTSD symptoms at TP3, beyond age, gender, trauma type and initial symptom severity. **b.** Conditional effects of TP1 left amygdala's activation to punishments vs. rewards on TP3 CAPS-5 total scores at different values of n=90 individuals' change in risky choice index (TP2-TP1) (blue = more risky behavior, Mean-1SD; green= no change in risky behavior, Mean; red = less risky behavior, Mean+1SD). All variables were centered prior to the analysis. Change in risky choice index is presented as a categorical variable with 3 levels for

illustration purposes, even though it was used as continuous variable in the analyses. *p >.05

a t=2.833, p=0.006

b $$Risky\ Choice\ Index\ (\%) = \frac{\#\ non\ matching\ chips}{\#\ non\ matching\ chips\ +\ \#\ matching\ chips} * 100$$

r=-0.185, p=0.018, n=132

a

Behavior (TP2-TP1):
Risky Choice Index

Neural NVS (TP1):
**Left Amygdala
Punishment>Reward**

Clinical Severity (TP3):
Total CAPS-5 Scores

b

Change in Risky Choice Index (TP2-TP1)

More Risky   No Change   Less Risky

Total CAPS-5 Scores at TP3

Left Amygdala Activation at TP1
Punishments > Rewards