Title: DNA methylation in *Ensifer* species during free-living growth and during nitrogen-fixing

symbiosis with *Medicago* spp.

Running title: DNA methylation in the genus *Ensifer*

Authors: George C. diCenzo[a,b,*], Lisa Cangioli[b], Quentin Nicoud[c], Janis H.T. Cheng[a], Matthew J.

Blow[d], Nicole Shapiro[d], Tanja Woyke[d], Emanuele G Biondi[c], Benoît Alunni[c], Alessio Mengoni[b,1],

and Peter Mergaert[c,1]

[1] These authors contributed equally to this work.

Affiliations: [a] Department of Biology, Queen's University, Kingston ON, K7L 3N6, Canada

[b]   Department of Biology, University of Florence, Florence, 50019, Italy

[c]   Université Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell, Gif-sur-

Yvette, 91198, France

[d]   U.S. Department of Energy Joint Genome Institute, Berkeley CA, 94720, USA

* Corresponding author: George C. diCenzo

19   Queen's University, Department of Biology

20   Biosciences Complex, Room 2433

21   116 Barrie Street

22   Kingston, ON, K7P0S7, Canada

23   george.dicenzo@queensu.ca

24   +1 (613) 533-6000 x78529

26   **Keywords:** Rhizobia, Symbiotic nitrogen fixation, DNA methylation, Cell cycle regulation, CcrM

ABSTRACT

Methylation of specific DNA sequences is ubiquitous in bacteria and has known roles in immunity and regulation of cellular processes, such as the cell cycle. Using single-molecule real-time sequencing, six genome-wide methylated motifs were identified across four *Ensifer* strains, five of which were strain-specific. Only the GANTC motif, recognized by the cell cycle-regulated CcrM methyltransferase, was methylated in all strains. In actively dividing cells, methylation of GANTC motifs increased progressively from the *ori* to *ter* regions in each replicon, in agreement with a cell cycle-dependent regulation of CcrM. In contrast, there was near full genome-wide GANTC methylation in the early stage of symbiotic differentiation. This was followed by a moderate decrease in the overall extent methylation and a progressive decrease in chromosomal GANTC methylation from the *ori* to *ter* regions in later stages of differentiation. We interpret these observations as evidence of dysregulated and constitutive CcrM activity during terminal differentiation, and we hypothesize that it is a driving factor for endoreduplication of terminally differentiated bacteroids.

INTRODUCTION

Methylation of genomic DNA is a pervasive phenomenon found in eukaryotes (Greenberg and Bourc'his, 2019; Tang et al., 2012; Zhang et al., 2018), archaea (Blow et al., 2016), and bacteria (Blow et al., 2016; Sánchez-Romero et al., 2015). The biological roles of DNA methylation are most extensively studied in mammals, where it contributes to normal development and disease via its impact on gene expression (Gopalakrishnan et al., 2008). In bacteria, DNA methylation is best known for its role in restriction-modification (R-M) systems that are thought to provide defence against phage infection and limit horizontal gene transfer through the degradation of invading non-methylated DNA (Vasu and Nagaraja, 2013). Several methyltransferases (MTases) of R-M systems have also been implicated in phase variation in pathogens through modulating gene expression (Atack et al., 2018). A recent study of over 200 bacterial and archaeal species identified orphan MTases not belonging to R-M systems in nearly half of the genomes (Blow et al., 2016). To date, biological functions have been attributed to very few orphan MTases, namely, the Dam MTase of the γ-Proteobacteria and the CcrM MTase of the α-Proteobacteria (Adhikari and Curtis, 2016). The Dam MTase of *Escherichia coli* is notable for its role in regulation of DNA replication (Campbell and Kleckner, 1990; Kang et al., 1999) and DNA repair (Lahue et al., 1989) by modulating the activity of other DNA-binding proteins. The CcrM MTase was first identified in *Caulobacter crescentus* (Zweiger et al., 1994), with homologs since identified in diverse α-Proteobacteria (Brilli et al., 2010; Wright et al., 1997). CcrM activity was shown to be cell cycle regulated in *C. crescentus* and *Agrobacterium tumefaciens* (Kahng and Shapiro, 2001; Zweiger et al., 1994), leading to methylation of its cognate DNA motif (the pentanucleotide GANTC) specifically during a short period at the end of DNA replication. This leads to a switching of GANTC sites between fully methylated (methylated on both strands) and hemi-methylated

(methylated only on the template strand) as a result of DNA replication (Kozdon et al., 2013),

which serves to modulate gene expression in a cell cycle-dependent fashion (Fioravanti et al.,

2013; Gonzalez et al., 2014; Gonzalez and Collier, 2013). Over- and under-expression of *ccrM*

results in defects in DNA replication and cell division (Gonzalez and Collier, 2013; Kahng and

Shapiro, 2001; Wright et al., 1997), while its complete loss is lethal under some conditions.

The rhizobia are a polyphyletic group of α-Proteobacteria and β-Proteobacteria that can

both live free in the soil and enter into an endosymbiotic interaction with legumes (Wang and

Young, 2019). This interaction begins following an exchange of signals between the free-living

partners (Oldroyd, 2013), and it culminates in the formation of a new organ known as a root nodule

within which the cytoplasm of plant cells contain thousands of N_2-fixing bacteria called bacteroids.

Bacteroid formation results in the differential expression of more than a thousand genes (Barnett

et al., 2004; Roux et al., 2014) and global changes in cellular metabolism (diCenzo et al., 2020).

In legumes of the Inverted-Repeat Lacking Clade (IRLC) and the Dalbergioid clade of the family

Papilionoideae, bacteroid development involves an additional process of terminal differentiation

(Czernic et al., 2015; Mergaert et al., 2006); in other legume clades, bacteroid differentiation is

less pronounced and is reversible. Terminal bacteroid development, in contrast to reversible

bacteroid formation, involves cell enlargement (bacteroids are 5- to 10-fold longer than their free-

living counterparts) and genome endoreduplication (resulting in up to 24 copies of the genome per

cell), indicative of a cell cycle transition occurring during differentiation (Mergaert et al., 2006).

Indeed, the correct expression of cell cycle regulators in *Ensifer* (syn. *Sinorhizobium*) *meliloti*, a

symbiont of *Medicago* species of the IRLC, is essential for the formation of functional bacteroids

(Kobayashi et al., 2009; Pini et al., 2013), while over-expression of CcrM or disruption of the

master cell cycle regulator CtrA can give rise to bacteroid-like morphology in free-living cells

(Pini et al., 2015; Wright et al., 1997). Additionally, mutants in the *E. meliloti* cell cycle regulators

divJ, *cbrA*, and *cpdR1*, encoding three negative regulators of CtrA, form non-functional nodules

in which bacteroids do not differentiate properly (Gibson et al., 2006; Kobayashi et al., 2009, p. 1;

Pini et al., 2013), and genes encoding several cell cycle regulators (including CcrM) are strongly

downregulated in bacteroids (Roux et al., 2014). The differentiation and cell cycle switch of

bacteroids is controlled by the legume host through the production of a large family of peptides,

known as Nodule-specific Cysteine-Rich (NCR) peptides (Farkas et al., 2014; Penterman et al.,

2014; Van de Velde et al., 2010).

Multiple studies have provided evidence that changes in the methylation status of the DNA

of legume nodule cells contributes to symbiotic development (Nagymihaly et al., 2017; Pecrix et

al., 2018; Satgé et al., 2016). Conversely, it remains unknown if methylation of rhizobium DNA

contributes to the regulation of N_2-fixation or bacteroid development. We are aware of only one

study (Davis-Richardson et al., 2016) comparing DNA methylation of a rhizobium

(*Bradyrhizobium diazoefficiens* USDA110) between free-living and symbiotic states (soybean

nodules), and comparing these changes with differential expression data. Intriguingly, the authors

identified a DNA motif that was methylated specifically in bacteroids (Davis-Richardson et al.,

2016). However, no clear evidence was presented that methylation of this (or any other) motif is

involved in transcriptional regulation, and the number of genes both differentially expressed and

differentially methylated in bacteroids did not appear to be different than expected by chance.

While these data may suggest that DNA methylation does not play a major role in regulating N_2-

fixation by rhizobia, they do not address the role of DNA methylation in terminal bacteroid

differentiation, as *B. diazoefficiens* undergoes reversible differentiation in soybean nodules

(Barrière et al., 2017).

Here, we use Pacific Biosciences Single-Molecule Real-Time (SMRT) sequencing to

detect genome-wide patterns of DNA methylation in four strains belonging to the genus *Ensifer*.

Our results indicate that DNA methylation is poorly conserved across the genus and suggests that

DNA methylation is not a major mechanism of regulating gene expression in these organisms aside

from cell cycle control. However, analysis of bacteroid samples led us to hypothesize that

constitutive activation of the CcrM MTase may be a contributing factor driving terminal

differentiation.

118   **RESULTS**

The methylomes of the genus *Ensifer*

Our experimental design, as summarized in the Materials and Methods and **Figure S1**, was

developed to support an investigation into multiple potential roles of DNA methylation in plant-

associated bacteria from the genus *Ensifer* through the use of SMRT sequencing. This was

accomplished by: i) including DNA samples isolated from phylogenetically diverse wild-type

strains, ii) examining DNA methylation in a single strain across multiple conditions (exponential

phase growth versus stationary phase; growth with sucrose versus growth with succinate), iii)

investigating the impact of a large-scale genome reduction on DNA methylation patterns, and iv)

isolating DNA from bacteroids purified from legume nodules.

We began with base modification analyses of four wild-type strains from three species,

including three nodule-forming strains (*E. meliloti* Rm2011, *E. meliloti* FSM-MA, *E. fredii*

NGR234) and one plant-associated, non-symbiotic strain (*E. adhaerens* OV14). To ensure

consistency, all strains were grown to mid-exponential phase in a common minimal medium with

succinate as the carbon source. A total of six methylated motifs were identified, of which five were

m6A modifications and one was a m4C modification (**Table 1**). Five of the six motifs were

methylated specifically in one strain. Only the GANTC motif, recognized by the highly conserved

cell cycle-regulated CcrM methyltransferase (Wright et al., 1997; Zweiger et al., 1994), was

methylated in all four strains. To further examine the conservation, or lack thereof, of DNA

modification across the genus *Ensifer*, we examined the distribution of methyltransferases in the

model species *E. meliloti*. Based on gene annotations, we identified 24 genes encoding putative

MTases in a previous pangenome analysis of 20 *E. meliloti* strains (**Table S1**) (diCenzo et al.,

2019). Of these 24 genes, only one (*ccrM*) was found in all 20 strains, while four were found in

two strains and 19 were found in a single strain. These results suggest that DNA methylation is

unlikely to play a biologically significant role in the genus *Ensifer* aside from cell cycle control

via CcrM-mediated methylation, and phage defence.

In support of most DNA methylation not having a regulatory function in the genus *Ensifer*,

none of the motifs methylated in *E. meliloti* Rm2011 were enriched in the promoter regions of

genes previously shown to be differentially expressed when grown with glucose vs. succinate

(diCenzo et al., 2017). Similarly, except for the GANTC motif as discussed below, no global effect

of carbon source (sucrose [glycolytic] vs. succinate [gluconeogenic]) was observed on the DNA

methylation pattern of *E. meliloti* Rm2011 (**Figure S2**). Similarly no global differences in DNA

methylation were detected between *E. meliloti* Rm2011 and RmP3496, a Rm2011 derivative

lacking the pSymA and pSymB replicons that together account for 45% of the genome content of

E. meliloti (diCenzo et al., 2014) (**Figure S3**).

Table 1. Methylated motifs identified in this study.

Motif[*]	Type[†]	Count[¥]	Frequency (motifs/kb)
E. meliloti 2011			
G**A**NTC CTN**A**G	m6A	11,169	1.67
RCG**C**CTC YGCGGAG	m4C	3,943	0.59
CGC**A**(N5)GTG GCGT(N5)C**A**C	m6A	1,085	0.16
E. meliloti FSM-MA			
G**A**NTC CTN**A**G	m6A	11,215	1.67
TCG**A**(N8)TCGA AGCT(N8)**A**GCT	m6A	2,612	0.39
E. fredii NGR234			
G**A**NTC CTN**A**G	m6A	11,111	1.61
CAG**A**(N7)GTTG GTCT(N7)C**A**AC	m6A	188	0.03
E. adhaerens OV14			
G**A**NTC CTN**A**G	m6A	8,475	1.10
WNCCG**A**TG WNGGCTAC	m6A	4,596	0.60

155 * The methylated nucleotides are indicated in boldface font.

156 † Indicates whether the modification is a N[6]-methyladenoside (m6A) or N[4]-methylcytosine (m4C).

157 ¥ The total times the motif appears in the genome, regardless of methylation status.

Cell cycle regulation by the CcrM methyltransferase

A progressive increase in the extent of methylation (herein defined as the estimated fraction of

reads mapping to a motif that were methylated) of GANTC sites was observed from the *ori* to *ter*

regions of the chromosomes of all four strains during mid-exponential growth (**Figures 1, S4-S6**).

There was a local drop in GANTC methylation around the 1.5 Mb mark in the *E. meliloti* FSM-

MA chromosome (**Figure S4**); however, this was seen in only two of three replicates and

corresponded to a region of high sequencing depth (**Figure S7**), suggesting the result is a

sequencing artefact. In contrast to exponential phase cultures, GANTC sites displayed near full

methylation (averaging ~ 95%) across the genome during early stationary phase, while all other

motifs displayed near full methylation (averaging 95-99%) across the genome regardless of growth

state (**Figures 1, S4-S6**). The observed pattern of GANTC methylation indicates a progressive

switch from full to hemi-methylated states as DNA replication proceeds (model provided as

Figure S8), confirming that the CcrM methyltransferase of the family *Rhizobiaceae* is cell cycle

regulated as demonstrated in *C. crescentus* (Kozdon et al., 2013; Mohapatra et al., 2014; Zweiger

et al., 1994). Interestingly, the genome-wide pattern of GANTC methylation displayed a smaller

variation in the extent of methylation from the *ori* to *ter* regions in *E. meliloti* Rm2011 when grown

with sucrose compared to succinate as the carbon source (**Figure S2**). While this observation could

suggest metabolic regulation of CcrM activity, we instead hypothesize, as displayed in **Figure**

S8A, that it is due to DNA replication being initiated later in the cell cycle when *E. meliloti* is

provided sucrose, as recent observations showed that central carbon metabolism influences the

rate of DNA polymerase processivity and timing of DNA replication initiation in *Bacillus subtilis*

(Nouri et al., 2018).

Notably, the GANTC methylation pattern differed across replicons within each genome.

For example, in *E. meliloti* Rm2011 the extent of GANTC methylation ranged from 0.80 to 0.98

for pSymB and 0.80 to 0.96 for pSymA, while for the chromosome the range was from 0.71 to

0.99 (**Figure 1**). This result suggests that replication of each replicon is asynchronous, with

replication of the secondary replicons being initiated later in the cell cycle than that of the

chromosome (model provided as **Figure S8**).

Figure 1. Genome-wide DNA methylation of *E. meliloti* Rm2011. (**A-F**) The extent of methylation is shown, using a 10 kb sliding window, of GANTC sites (black) and CGCA(N$_5$)GTG sites (grey) across the chromosome (**A,D**), pSymB (**B,E**), and pSymA (**C,F**) replicons of exponential phase (**A-C**) or early stationary phase (**D-F**) *E. meliloti* Rm2011. Averages from three biological replicates are shown. The red (GANTC) and blue (CGCA(N$_5$)GTG) lines are polynomial regression lines calculated in R using the "rlm" method and the formula "y~poly(x,2)". (**G-I**) Cumulative GC skews, shown using a 10 kb sliding window, across the *E. meliloti* Rm2011 chromosome (**G**), pSymB (**H**), and pSymA (**I**) replicons.

A previous study identified 462 cell cycle-regulated genes in *E. meliloti* through RNA-sequencing of synchronized cell populations (De Nisco et al., 2014), which were classified into six groups based on the timing of their expression. We identified 111 cell cycle-regulated genes, belonging to 78 transcripts, that contained at least one GANTC in the predicted promoter regions (defined as the 125 bp upstream of the transcript; **Dataset S1**), and distribution of these 111 genes across the six cell cycle gene expression groups was unbiased (De Nisco et al., 2014). As these 111 genes are both cell cycle regulated and contain a GANTC site, they represent an initial candidate CcrM regulon in *E. meliloti* although further work is required to validate the CcrM regulon.

We found it striking that the *E. adhaerens* OV14 genomes had 2,636 to 2,740 fewer

GANTC sites than the other three strains, despite having the largest genome size. Normalized by

genome length, there are 1.10 GANTC sites per kb in the *E. adhaerens* OV14 genome (**Table 1**),

which is similar to the 1.12 GANTC sites per kb in *C. crescentus* NA1000. In contrast, the three

legume symbionts contained more than 1.60 GANTC sites per kb across their genomes (**Table 1**).

This result prompted us to examine the frequency and distribution of GANTC sites across 157

Ensifer genomes. As defined previously (Fagorzi et al., 2020), the genus *Ensifer* can be broadly

sub-divided into two monophyletic clades; the "symbiotic" clade (113 strains) in which nearly all

strains are legume symbionts, and the "non-symbiotic" clade (44 strains) in which nearly all strains

are non-symbionts (**Figure 2A**). Consistent with previous results (Gonzalez et al., 2014), GANTC

sites occurred less frequently in all genomes (0.90 to 1.83 GANTC sites per kb) than expected in

a random sequence of nucleotides (~ 3.5 GANTC sites per kb). Moreover, GANTC sites were ~

2-fold more common in intergenic regions than in coding regions (**Figure 2B**). Strikingly, there

was a strong and statistically significant difference (p-value $< 1 \times 10^{-10}$; two-sample t-test) in the

frequency of GANTC sites across the genomes of strains belonging to the symbiotic clade

compared to strains of the non-symbiotic clades (**Figure 2B**), with an overall average of 1.70 and

1.06 GANTC sites per kb in the symbiotic and non-symbiotic clades, respectively. The difference

in the frequency of GANTC sites between the two clades could not be explained by differences in

the GC content of these organisms, as both clades had an average GC content of 61.9% (**Figure**

S9), suggesting that the difference reflects underlying differences in the evolution, and possibly

the biology, of these two clades.

Figure 2. GANTC frequency in the genus *Ensifer*. (**A**) An unrooted maximum likelihood phylogeny of 17 representative *Ensifer* strains. The phylogeny represents the bootstrap best tree following 100 bootstrap replicates, prepared on the basis of the concatenated nucleotide alignments of 1566 core genes. Values represent the bootstrap support. N_2-fixing legume symbionts were identified by the presence of the symbiotic genes *nodABC* and *nifHDK*. They are indicated in blue, while red denotes non-symbiotic strains. The four wild-type strains used in this study are indicated with arrows. (**B**) Box plots summarizing the frequency of GANTC sites (presented as GANTC sites per kb) in 157 *Ensifer* strains is shown. The monophyletic "symbiotic" and "non-symbiotic" clades as defined previously (Fagorzi et al., 2020), are represented by 113 and 44 genomes respectively. The densities of GANTC sites across the entire genome (pink), within coding regions (blue), and within intergenic regions (yellow) are shown. Statistically different values ($p < 0.05$) are denoted by uppercase letters as determined by a one-way ANOVA followed by a Tukey's HSD post hoc test.

Contributions of DNA methylation to bacteroid differentiation

The only previously published study to examine the role of rhizobium DNA methylation during symbiosis using SMRT sequencing did so in a symbiosis where the bacteria do not undergo terminal differentiation (Barrière et al., 2017; Davis-Richardson et al., 2016). To evaluate whether DNA methylation potentially contributes to regulation of terminal differentiation, we determined

the DNA methylation patterns of *E. meliloti* Rm2011 and *E. meliloti* FSM-MA bacteroids purified

from whole *Medicago sativa* nodules. *E. meliloti* FSM-MA bacteroids were additionally purified

from *Medicago truncatula* nodules to determine whether the host plant influences bacteroid DNA

methylation. *E. meliloti* Rm2011 bacteroids were not isolated from *M. truncatula* nodules as,

unlike FSM-MA, Rm2011 forms a poor symbiosis with *M. truncatula* and produces moderately

differentiated bacteroids in this host (Kazmierczak et al., 2017; Moreau et al., 2008).

Moreover, we exploited the spatially distinct developmental zones that are present in

indeterminate nodules (Vasse et al., 1990), like those formed by *M. sativa* and *M. truncatula*. At

the tip of these nodules, a bacteria-free meristem is present, responsible for the continuous growth

of the nodule. Immediately below is the infection and differentiation zone II where nodule cells

become infected and bacteria differentiate into the large, polyploid bacteroids. The tip and zone II

of nodules appears white. Adjacent to the white zone II is the easily recognizable pinkish zone III

(due to the presence of the oxygen-carrying leghemoglobin) where mature bacteroids fix nitrogen.

This nodule tissue organization provided an opportunity to examine how DNA methylation

patterns may differ between differentiating and differentiated bacteroids. To this end, *E. meliloti*

Rm2011 and FSM-MA bacteroids were isolated from nodules hand-sectioned at the white-pink

border; bacteroids isolated from the white sections represent the infecting and differentiating

bacteroids (zone II) while those isolated from the pink sections represent the mature, hence

terminally differentiated, N_2-fixing bacteroids (zone III).

Fluorescence microscopy and flow cytometry confirmed that nodule sectioning resulted in

the isolation of distinct bacteroid populations (**Figures S10-S13**). Nearly all of the bacteroids

isolated from whole-nodule samples and zone III samples were enlarged and polyploid, and most

were positive for propidium iodide (PI) staining as expected for terminally differentiated

bacteroids (Mergaert et al., 2006). In contrast, bacteroids of the zone II samples contained a mix

of cell types differing in their size, ploidy level, and PI staining. These data confirmed that the

whole-nodule samples and zone III samples consisted predominately of mature N_2-fixing

bacteroids, whereas the zone II samples contained a mix of cells at various stages of bacteroid

differentiation.

With the exception of the GANTC sites (described below), no global difference was

observed in the methylation patterns of bacteroids versus free-living cells (**Figures 3, 4, S14-S17**).

Although a lower percentage of each motif was detected as methylated in the bacteroid samples

compared to free-living samples, this was correlated with lower sequencing depth (**Table S2**) and

thus unlikely to be biologically meaningful. Unlike *B. diazoefficiens* USDA110 (Davis-Richardson

et al., 2016), we did not identify motifs that were methylated specifically in the *E. meliloti*

bacteroids. In addition, we found little evidence for any of the methylated motifs being enriched

in the promoter regions of *E. meliloti* Rm2011 genes up-regulated or down-regulated in bacteroids

relative to free-living cells, as identified in published transcriptomic data for *E. meliloti* Rm1021

(Barnett et al., 2004), a near-isogenic relative of strain Rm2011 also derived from the nodule

isolate SU47. These data suggest that most DNA methylation is unlikely to be a significant factor

in directly regulating gene expression in *E. meliloti* bacteroids.

Figure 3. Chromosome-wide DNA methylation of *E. meliloti* Rm2011 bacteroids. The extent of methylation of (**A-E**) GANTC, (**F-J**) CGCA(N$_5$)GTG, and (**K-O**) RCGCCTC motifs across the *E. meliloti* Rm2011 chromosome is shown using a 10 kb sliding window. Averages from three biological replicates are shown for free-living and whole nodule samples; data represents one replicate for the zone II and zone III nodule sections. (**A,F,K**) Free-living cells harvested in mid-exponential phase. (**B,G,L**) Free-living cells harvested in early stationary phase. (**C,H,M**) Bacteroids isolated from *M. sativa* zone II nodule sections. (**D,I,N**) Bacteroids isolated from *M. sativa* zone III nodule sections. (**E,J,O**) Bacteroids isolated from *M. sativa* whole nodule samples. The red lines are polynomial regression lines calculated in R using the "rlm" method and the formula "y~poly(x,2)". Data for pSymB and pSymA are shown in Figures S14 and S15.

Figure 4. Chromosome-wide DNA methylation of *E. meliloti* FSM-MA bacteroids. The extent of methylation of (**A-F**) GANTC and (**G-L**) TCGA(N_8)TCGA motifs across the *E. meliloti* FSM-MA chromosome is shown using a 10 kb sliding window. Averages from three biological replicates are shown for free-living and whole nodule samples; data represents one replicate for the zone II and zone III nodule sections. (**A,G**) Free-living cells harvested in mid-exponential phase. (**B,H**) Free-living cells harvested in early stationary phase. (**C,I**) Bacteroids isolated from *M. sativa* zone II nodule sections. (**D,J**) Bacteroids isolated from *M. sativa* zone III nodule sections. (**E,K**) Bacteroids isolated from *M. sativa* whole nodule samples. (**F,L**) Bacteroids isolated from *M. truncatula* whole nodule samples. The red lines are polynomial regression lines calculated in R using the "rlm" method and the formula "y~poly(x,2)". Data for pSymB and pSymA are shown in Figures S16 and S17.

CcrM methyltransferase activity is dysregulated during terminal differentiation

Bacteroid development involves cell enlargement and genome endoreduplication, indicative of a cell cycle transition occurring during differentiation (Mergaert et al., 2006). Indeed, expression of *ccrM* and other genes encoding cell cycle regulators vary across stages of bacteroid development and are strongly downregulated in mature nitrogen-fixing *E. meliloti* bacteroids (Roux et al., 2014). We were therefore interested in examining whether GANTC methylation by the CcrM MTase was

disrupted in bacteroids. Our data revealed a surprising genome-wide pattern of GANTC

methylation in *E. meliloti* Rm2011 and FSM-MA bacteroids, which differed from free-living cells

in either the exponential or stationary phases of growth (**Figures 3, 4, S14-S17**). The majority of

GANTC sites had moderate to high levels of methylation in zone II, zone III, or whole-nodule

samples, averaging 0.71 to 0.95 across each replicon (**Figures 3, 4, S14-S17, and Tables S3, S4**).

Most distinctive, a progressive decrease in the extent of chromosomal methylation of the GANTC

sites was observed from the *ori* to *ter* in nearly all bacteroid samples, revealing a characteristic

smiling pattern, which differs from the patterns seen in exponential (frowning pattern, i.e., a

progressive increase from *ori* to *ter*) and stationary (consistent methylation) phase cells. The

exception was the *E. meliloti* FSM-MA zone II bacteroid sample, which displayed a consistently

high level of GANTC methylation across the genome (**Figure 4C**). This pattern, which is different

from those in exponential phase cells as well as mature bacteroids, could correspond to the

methylation status of an early stage of bacteroid differentiation. We did not observe the same

pattern in the *E. meliloti* Rm2011 zone II samples. As noted earlier (**Figures S10-S13**), the zone

II samples contain cells at various stages of differentiation. Given that terminal differentiation is

associated with an up to 24-fold increase in DNA content, small increases in the proportion of cells

at late stages of differentiation could mask the DNA methylation pattern of the cells at early stages

of differentiation. Thus, we hypothesize that the Rm2011 zone II sample captures a later stage of

differentiation than that captured by the FSM-MA zone II sample. Supporting this hypothesis, the

distribution of DNA content per cell in the flow cytometry data was flatter for *E. meliloti* Rm2011

zone II bacteroids compared to *E. meliloti* FSM-MA zone II bacteroids (**Figure S18**), which

suggests that the former sample represents a broader range of differentiation stages than the latter

sample. In contrast to GANTC, the extent of methylation of the second m6A modified motif in

each genome was consistently high, irrespective of condition or replicon (**Figures 3, 4, S14-S17,**

and Tables S3, S4). Similarly, sequencing depth was consistent across the length of each replicon

(**Figure S7**). These observations indicate that the changes in GANTC methylation patterns are

biologically meaningful and not simply a sequencing artefact.

To further explore changes in CcrM activity during bacteroid differentiation, we took

advantage of a collection of *M. truncatula* mutant plant lines (*dnf1*, *dnf2*, *dnf4*, *dnf5*, *dnf7*) whose

nodules contain bacteria blocked at various stages of differentiation (Bourcy et al., 2013;

Domonkos et al., 2013; Horváth et al., 2015; Kim et al., 2015; Lang and Long, 2015; Starker et

al., 2006; Wang et al., 2010). Microscopy and flow cytometry data was consistent with past work

(Lang and Long, 2015) showing that bacteroids were blocked at the earliest to latest stages of

differentiation in mutant plant lines in the order *dnf1* → *dnf5* → *dnf2* → *dnf7* → *dnf4* (**Figure 5**).

Nodule bacteria of *M. truncatula dnf1* mutant plants were small with one or two haploid genome

copies per cell (i.e., ploidy level = 1 or 2) (**Figures 5A, 5G**), suggesting that the cell population

was dominated by actively dividing cells that had not yet begun differentiation. Indeed, the

GANTC methylation pattern of these cells (**Figures 5M, S19, S20**) resembled the frowning pattern

of exponentially growing free-living cells (**Figure 4A**). Although the nodule bacteria of *M.*

truncatula dnf5 mutant plants were also small and undifferentiated into bacteroids, the majority of

cells had a ploidy level of one (**Figures 5B, 5H**), suggesting these cells had ceased replication but

had not yet begun the process of endoreduplication. GANTC methylation was consistently high

across the chromosome of bacteria purified from *dnf5* nodules (**Figure 5N**), similar to stationary

phase free-living cells (**Figure 4B**) and indicating that terminal differentiation is preceded by full

GANTC methylation.

Figure 5. Bacteroid morphology and chromosomal GANTC methylation in *E. meliloti* bacteroids purified from *M. truncatula dnf* mutant nodules. Data is shown for *E. meliloti* FSM-MA bacteroids purified from (**A,G,M,S**) *M. truncatula dnf1* mutant nodules, (**B,H,N,T**) *M. truncatula dnf5* mutant nodules, (**C,I,O,U**) *M. truncatula dnf2* mutant nodules, (**D,J,P,V**) *M. truncatula dnf7* mutant nodules, (**E,K,Q,W**) *M. truncatula dnf4* mutant nodules, and (**F,L,R,X**) *M. truncatula* A17 wild-type nodules. (**A-F**) Micrographs of *E. meliloti* FSM-MA bacteroids

stained with the DNA binding dye DAPI. The scale bar represents 30 μm. (**G-L**) Pseudo-coloured

scatterplots displaying the cell morphology (X-axis) and DNA content (Y-axis) of *E. meliloti*

FSM-MA bacteroids, as determined based on flow cytometry analysis of DAPI stained cells. The

red dashed ellipses indicate the position of undifferentiated bacteria as in culture (not shown) or in

the *dnf1* mutant nodules (lower left ellipse) or fully mature bacteroids as in the A17 wild-type

nodules (top right ellipse). (**M-X**) The extent of methylation of (**M-R**) GANTC or (**S-X**)

TCGA(N_8)TCGA motifs across the *E. meliloti* FSM-MA chromosome, shown using a 10 kb

sliding window. The red lines are polynomial regression lines calculated in R using the "rlm"

method and the formula "y~poly(x,2)". Data for pSymB and pSymA are shown in Figures S19

and S20.

The nodule bacteria purified from *M. truncatula dnf2* mutant nodules were a mix of

undifferentiated and partially differentiated bacteroids, which were polyploid to an extent similar

to bacteroids purified from wild-type A17 nodules (**Figure 5I** compared to **Figure 5L**); however,

their cell size was much smaller (**Figure 5C** compared to **Figure 5F**). This was similar to

differentiating bacteroids purified from *M. truncatula* and *M. sativa* zone II nodule sections, many

of which had high ploidy without a corresponding increase in cell size (**Figures S11, S13**). The

GANTC methylation pattern of bacteroids from *dnf2* nodules (**Figure 5O**) was also similar to that

of zone II nodule sections. There was a consistently high extent of GANTC methylation across the

chromosome averaging 0.870, which was less than that of bacteroids purified from *dnf5* nodules

(0.956) but higher than that of bacteroids purified from wild-type A17 nodules (0.804) (**Table S5**),

and without the smiling pattern. The nodule bacteria purified from *M. truncatula dnf7* and *dnf4*

nodules also contained a mix of undifferentiated bacteria and fully differentiated bacteroids

(**Figures 5D, 5E, 5J, 5K**), with the number of undifferentiated bacteria greater in *dnf7* nodules

compared to *dnf4* nodules. The GANTC methylation pattern of bacteroids purified from *dnf7* and

dnf4 nodules was similar to that of bacteroids purified from A17 nodules (**Figures 5P-Q**). Overall,

we interpret the data from bacteroids purified from section nodules and *M. truncatula dnf* mutant

nodules as suggesting that CcrM is dysregulated during terminal bacteroid differentiation and that

CcrM is constitutively active during endoreduplication.

Chromosome, pSymB, and pSymA sequencing depth are unequal in *E. meliloti* bacteroids

We noticed that in each bacteroid sample, the average extent of GANTC methylation for the

chromosomes of the two strains were lower (by 0.04 to 0.13) than that of pSymA or pSymB, and

unlike the chromosome, the extent of GANTC methylation was relatively constant across pSymA

and pSymB (**Figures S14-S17** compared to **Figures 3 and 4**). These results suggest that, unlike in

free-living cells, replication of the three replicons is not well coordinated during terminal

differentiation. In agreement with this hypothesis, the mean sequencing depth across pSymA and

pSymB was on average ~ 33% lower than that of the chromosome in all replicates of the *E. meliloti*

whole-nodule bacteroid samples (**Table 2**). Similarly, the mean sequencing depth across pSymA

and pSymB was on average ~ 23% lower than that of the chromosome for the polyploid bacterial

cell populations purified from *M. truncatula dnf2*, *dnf7*, and *dnf4* mutant nodules, but not for the

haploid/diploid bacterial cell populations purified from *M. truncatula dnf1* and *dnf5* mutant

nodules (**Table 2**). Assuming sequencing depth is correlated with copy number, this observation

suggests that *E. meliloti* bacteroids carry approximately two copies each of pSymA and pSymB

per three copies of the chromosome.

Table 2. Relative sequencing depth of each *E. meliloti* replicon.

Strain	Condition	Relative mean sequencing depth *		
		Chromosome	pSymA	pSymB
Rm2011	Mid-exponential	1.00 ± 0.00	1.00 ± 0.03	0.98 ± 0.03
Rm2011	Stationary phase	1.00 ± 0.00	1.06 ± 0.02	1.02 ± 0.03
Rm2011	*M. sativa* bacteroids	1.00 ± 0.00	0.64 ± 0.02	0.62 ± 0.02
FSM-MA	Mid-exponential	1.00 ± 0.00	1.02 ± 0.03	0.87 ± 0.05
FSM-MA	Stationary phase	1.00 ± 0.00	1.10 ± 0.00	1.01 ± 0.03
FSM-MA	*M. sativa* bacteroids	1.00 ± 0.00	0.79 ± 0.04	0.71 ± 0.02
FSM-MA	*M. truncatula* bacteroids	1.00 ± 0.00	0.66 ± 0.02	0.59 ± 0.01
FSM-MA	*dnf1* bacterial cells	1.00	0.98	0.90
FSM-MA	*dnf5* bacterial dells	1.00	0.96	0.94
FSM-MA	*dnf2* bacterial dells	1.00	0.81	0.73
FSM-MA	*dnf7* bacterial dells	1.00	0.82	0.71
FSM-MA	*dnf4* bacterial dells	1.00	0.80	0.73
FSM-MA	A17 bacteroids	1.00	0.76	0.67

* Sequencing depth is presented relative to the sequencing depth of the chromosome in the same sample. Values are the means of triplicate samples ± standard deviation, except for the third section of the table for which numbers are based on a single replicate.

DISCUSSION

In this study, we examined the genome-wide DNA methylation patterns in the free-living cells of four *Ensifer* strains, and in bacteroids of two *E. meliloti* strains, and detected a total of six methylated motifs. We were able to predict cognate MTases for most of the motifs based on genome annotations, the exception being the WNCCGATG motif of *E. adhaerens* OV14. The CGCA(N$_5$)GTG motif of *E. meliloti* Rm2011 is presumably methylated by Smc02296 (HsdM), a predicted m6A MTase belonging to the HsdRSM type I R-M system that is known to be functional and reduce transformation efficiency (Brumwell et al., 2019; Ferri et al., 2010). The RCGCCTC motif of *E. meliloti* Rm2011 is possibly methylated by Smc03763, a predicted cysteine-specific MTase located upstream of the gene *vsr* that putatively encodes a very short patch repair protein. Neither of these proteins are found in the other three strains examined here. The motifs

TCGA(N_8)TCGA of *E. meliloti* FSM-MA and CAGA(N_7)GTTG of *E. fredii* NGR234 are likely

methylated by SMB554_16155 and NGR_c01340, respectively, which are 88% identical at the

amino acid level. Homologs of these two proteins are not found in the other two strains.

Except for GANTC, each methylated motif was detected as methylated only in a single

strain. Moreover, MTases, apart from CcrM, are not conserved among *E. meliloti* strains. The lack

of conservation suggests that most DNA methylation does not have a major regulatory role in the

genus *Ensifer*, aside from its role in cell cycle regulation. Supporting this conclusion, no motif was

enriched in the promoter regions of symbiosis, carbon source, or cell cycle-regulated genes, and

we did not detect any motifs that were methylated specifically in bacteroids. However, we cannot

rule out that one or more methylated motifs may influence specific gene expression during free-

living growth, differentiation, or N_2-fixation through extended motifs or proximity to other

promoter elements, similar to the interplay between CcrM and GcrA during cell cycle regulation

in *C. crescentus* (Fioravanti et al., 2013; Haakonsen et al., 2015).

As previously published studies have provided evidence for a cell cycle transition

occurring during terminal bacteroid differentiation (Mergaert et al., 2006), we were particularly

interested in CcrM, a cell cycle-regulated MTase that is broadly conserved in the α-Proteobacteria,

and its cognate DNA motif, GANTC. By identifying GANTC sites in the promoter regions of a

previously determined set of 462 cell cycle-regulated genes (De Nisco et al., 2014), we defined a

candidate CcrM regulon in *E. meliloti* consisting of 111 genes. However, further studies are

required to better delineate the CcrM regulon in *E. meliloti* as the presence of a GANTC site is not

diagnostic of CcrM regulation; GANTC sites were found in the promoter regions of 904 transcripts

that did not display cell cycle regulation, and the promoter regions of cell cycle regulated genes

were not enriched in GANTC sites relative to the whole *E. meliloti* genome. Studies in *C.*

crescentus suggest that the impact of the fully or hemi-methylated status of GANTC sites on gene

expression is mediated, at least in part, through modulating the activity of the transcriptional

regulator GcrA (Fioravanti et al., 2013; Haakonsen et al., 2015). However, not all promoter sites

containing a GANTC motif are regulated by GcrA in *C. crescentus*, with the relationship

dependent on an extended YGAKTCK motif and the precise position of this motif relative to other

promoter elements (Haakonsen et al., 2015). Likely, CcrM-mediated gene regulation in the genus

Ensifer is similarly dependent upon additional sequence elements beyond the GANTC motif.

Consistent with past observations (Gonzalez et al., 2014), GANTC sites were under-

represented in the genomes of 157 *Ensifer* strains, particularly within coding regions. More

surprising, however, was the strong difference in the frequency of GANTC sites between the

previously defined (Fagorzi et al., 2020) symbiotic and non-symbiotic clades in the genus *Ensifer*,

with the frequency of GANTC sites being ~ 60% higher in the symbiotic clade. As methylation of

the GANTC motif by CcrM is known to influence gene expression in *C. crescentus* (Gonzalez et

al., 2014), our observations suggest that CcrM has a greater impact on modulating gene expression

in the symbiotic clade compared to the non-symbiotic clade. Although further work is required to

understand the biological significance of the greater frequency of GANTC sites in the symbiotic

clade, it is tempting to speculate it is associated with legume symbiosis.

Our data is consistent with CcrM activity differing during terminal bacteroid differentiation

compared to free-living cells. The overall moderate to high rates of GANTC methylation in all *E.*

meliloti bacteroid samples, coupled with the lack of a chromosome-wide pattern in the *E. meliloti*

FSM-MA zone II sample, leads us to hypothesize that CcrM remains constitutively active

throughout most of terminal differentiation. This hypothesis is supported by the results for nodule

bacteria purified from *M. truncatula dnf* mutant nodules, which showed that differentiation is

preceded by full GANTC methylation and that GANTC methylation remains high (but moderately

lower) during endoreduplication followed by another moderate drop in GANTC methylation in

late stages of differentiation. Considering that over-expression of CcrM can give rise to bacteroid-

like morphology in free-living cells (Wright et al., 1997), we hypothesize that constitutive CcrM

MTase activity is one (of potentially multiple) factor(s) driving polyploidization of bacteroids

(**Figure S8**). However, further studies monitoring CcrM abundance and artificially manipulating

ccrM expression throughout bacteroid differentiation are required to conclusively determine if

CcrM is constitutively active during terminal differentiation and the importance of this activity to

the promotion of endoreduplication.

CcrM activity is confined to a short window in the cell cycle since the *ccrM* gene is

expressed in the late phase of genome replication (De Nisco et al., 2014) and the CcrM protein is

degraded by the Lon protease prior to cell division (Wright et al., 1996). Thus, constitutive CcrM

MTase activity in differentiating bacteroids could be obtained through an aberrant expression of

the gene or alternatively a lack of proteolytic degradation of the CcrM protein. In agreement with

the latter possibility, Lon protease was identified as a target of the NCR247 peptide (Farkas et al.,

2014). It is tempting to speculate that NCR peptides like NCR247 inhibit Lon protease activity

post-translationally, thereby stabilizing CcrM and triggering bacteroid differentiation. However,

the CcrM MTase does not appear to remain active in fully differentiated bacteroids, with the lower

GANTC methylation near the chromosomal *ter* regions suggesting that loss of CcrM MTase

activity occurs slightly prior to completion of genome endoreduplication (model provided as

Figure S8). These hypotheses are consistent with *M. truncatula – E. meliloti* nodule zone-specific

RNA-sequencing data (Roux et al., 2014), which showed that *ccrM* expression in the root distal

portion of zone II is ~ 2-fold higher than in the root proximal portion of zone II, and ~ 10-fold

higher than in zone III. The ~ 10-fold difference in *ccrM* expression across nodule zones suggests

to us that the level of *ccrM* expression during early stages of bacteroid differentiation is

biologically significant, a prerequisite for the constitutive CcrM activity that we hypothesize.

Our analyses also provide insight into the genome replication dynamics of *E. meliloti*

during free-living growth and terminal bacteroid differentiation. Notably, flow cytometry data of

E. meliloti bacteroids purified from zone II nodule sections and *M. truncatula dnf2* nodules suggest

that endoreduplication and cell enlargement largely occur subsequently, not concurrently. Genome

replication might be a much faster process than cell growth or alternatively, endoreduplication

might be required to drive cell enlargement. Moreover, our data are consistent with a loss of

coordination of replication of the three replicons during terminal bacteroid differentiation, leading

to unequal copy numbers with two copies of pSymA and pSymB per three copies of the

chromosome in bacteroids. This relative change in replicon copy number occurs concomitantly

with differentiation and polyploidization, as supported by the relative abundance of the replicons

differing in nodule bacterial cells that have experienced endoreduplication (i.e., cells retrieved

from *M. truncatula dnf2*, *dnf7*, and *dnf4* mutant nodules) but not in bacterial cells that had not yet

undergone endoreduplication (i.e., cells retrieved from *M. truncatula dnf1* and *dnf5* mutant

nodules). This differs from free-living cells, where copy number of the three replicons was

approximately equal based on average sequencing depth. In contrast to our results, a previous

comparison of the relative abundance of the three replicons in free-living *E. meliloti* Rm1021

versus bacteroids, no changes were detected using comparative genome hybridization with

microarrays (Mergaert et al., 2006). We believe that the difference between our present data and

the previous analysis is due to the subtlety of the differences and the lower sensitivity of the

microarray hybridization method compared to high throughput sequencing.

542 We also observed that during free-living exponential growth, the extent of GANTC

543 methylation at the *ori* of pSymA and pSymB is higher than at the *ori* of the chromosome, while

544 the *ter* of the pSymA and pSymB has a slightly lower extent of GANTC methylation than the *ter*

545 of the chromosome. As GANTC methylation occurs at a fixed stage of the cell cycle corresponding

546 to the end of chromosome replication, our observations indicate that pSymA and pSymB

547 replication is initiated later in the cell cycle than initiation of chromosome replication, while their

548 replication terminates slightly before completion of chromosomal replication and the activation of

549 CcrM (model provided as **Figure S8**). These results provide additional support for the notion of

550 spatiotemporal regulation of DNA replication and partitioning in the multipartite *E. meliloti*

551 genome as proposed previously (De Nisco et al., 2014; Frage et al., 2016). Similarly, replication

552 of chromosome II of *Vibrio cholerae* is delayed relative to chromosome I, leading to the replication

553 of these two replicons terminating at approximately the same time (Rasmussen et al., 2007). Thus,

554 co-ordinating the timing or replication termination may be a general feature of multipartite

555 genomes.

557         **MATERIALS AND METHODS**

558 **Experimental design**

559 The overall experimental design is summarized in **Figure S1**. Genomic DNA was isolated from

560 four wild-type *Ensifer* strains to explore how DNA methylation varies across this genus; to allow

561 direct comparison, the four strains were grown to mid-exponential phase in minimal medium with

562 succinate as a carbon source. To investigate how DNA methylation patterns differ between

563 actively dividing and non-dividing cells, genomic DNA was isolated from *E. meliloti* Rm2011

564 growth to either mid-exponential phase or stationary phase. Genomic DNA was isolated from *E.*

meliloti Rm2011 grown to mid-exponential phase with either a glycolytic (sucrose) or

gluconeogenic (succinate) carbon source to investigate whether DNA methylation may play a role

in regulating carbon metabolism. Furthermore, a *E. meliloti* Rm2011 derivative lacking the pSymA

and pSymB replicons (named RmP3496) was studied to gain insight into whether these replicons

contribute to DNA methylation patterns in *E. meliloti* Rm2011; these strains were grown with

sucrose (instead of succinate) as RmP3496 lacks the succinate transporter.

In addition to the free-living samples, *E. meliloti* bacteroid samples purified from legume

nodules were collected to investigate changes in DNA methylation during bacteroid differentiation

and nitrogen fixation. To do so, *E. meliloti* Rm2011 and *E. meliloti* FSM-MA bacteroids were

isolated from *M. sativa* whole nodules. *E. meliloti* FSM-MA bacteroids were additionally purified

from *M. truncatula* whole nodules to examine the impact of host plant on bacteroid DNA

methylation patterns. *E. meliloti* Rm2011 bacteroids were only isolated from *M. sativa* nodules as,

unlike FSM-MA, Rm2011 forms a poor symbiosis with *M. truncatula* (Kazmierczak et al., 2017;

Moreau et al., 2008). Moreover, *E. meliloti* Rm2011 and *E. meliloti* FSM-MA bacteroids were

isolated from *M. sativa* nodule sections (sectioned at the white – pink border to separate the root

distal infection and differentiation zone II [white] from the root proximal nitrogen-fixing zone III

[pink]) to facilitate an analysis of how DNA methylation patters differ between differentiating

bacteroids and fully differentiated and nitrogen-fixing bacteroids. This was followed by isolation

of *E. meliloti* FSM-MA bacteroids from whole nodules of *M. truncatula* mutant lines (*dnf1*, *dnf2*,

dnf4, *dnf5*, *dnf7*) to investigate DNA methylation patterns in nodule bacteria blocked as various

stages of differentiation.

Bacterial strains and growth conditions

Bacterial strains used in this work are listed in **Table S6**. All strains were routinely grown on TY

with 2 µM $CoCl_2$ as it was required for *E. meliloti* RmP3496 (diCenzo et al., 2014). The MM9

minimal medium (diCenzo et al., 2014) consisted of the following: 40 mM MOPS, 20 mM KOH,

19.2 mM NH_4Cl, 85.6 mM NaCl, 2 mM KH_2PO_4, 1 mM $MgSO_4$, 0.25 mM $CaCl_2$, 1 µg ml^{-1} biotin,

42 nM $CoCl_2$, 38 µM $FeCl_3$, 10 µM thiamine-HCl, and either 10 mM sucrose (MM9-sucrose) or

20 mM disodium succinate (MM9-succinate). Prior to inoculation of plants with *E. meliloti*, the

strains were grown in YEB medium (Krall et al., 2002).

DNA isolation from free-living cells

Overnight cultures of all strains grown in MM9-succinate or MM9-sucrose media were diluted

into 10 mL of the same medium to a starting OD_{600nm} of 0.025 (0.05 for RmP3496) and incubated

overnight at 30˚C with shaking (130 rpm). The next day, cultures were diluted into 40 mL of the

same medium in 100 mL flasks to the OD_{600nm} values as listed in **Table S7**. To obtain mid-

exponential phase samples, cultures were harvested after 15.5-16 hours of growth at OD_{600nm}

values between 0.37 and 0.69 (**Table S7**). To obtain stationary phase samples, cultures were

harvested after 24 hours of growth at OD_{600nm} values of ~ 1.4. In all cases, cultures were streaked

on TY plates to check for contamination and then centrifuged (8,200 *g*, 10 minutes, 4°C); the full

40 mL was centrifuged for mid-exponential phase cultures, whereas only 15 mL was centrifuged

for stationary phase cultures. Most of the supernatant was discarded, and the pellet resuspended in

the remaining volume, transferred to a 2 mL tube, centrifuged again (16,200 *g*, room temperature,

one minute), and the supernatant discarded. Three biological replicates, each starting from a

separate overnight culture, were performed. DNA was isolated using phenol:chloroform

extractions and ammonium acetate precipitations as described elsewhere (Cowie et al., 2006), and

the DNA pellets (following RNase A treatment) were resuspended in 200 µL of 10 mM Tris-HCl,

pH 8.5.

DNA isolation from bacteroids

M. sativa cv. Gabès and *M. truncatula* cv. A17 were used wild-type plants for all experiments. *M. truncatula dnf1*, *dnf2*, *dnf4*, *dnf5*, and *dnf7* mutants (Starker et al., 2006), derived from the A17 wild type, were used for collection of bacteroids blocked at various stages of differentiation. Seeds were scarified, surface sterilized, and germinated on Kalys agar as described previously (Kazmierczak et al., 2017). Fifty mL of overnight cultures of *E. meliloti* Rm2011 or FSM-MA, grown in YEB, were centrifuged (4,000 *g*, 20 minutes, room temperature) and resuspended in ~ 1,200 mL of sterile, distilled water to obtain a cell suspension at an OD_{600nm} of ~ 0.1. Germinated seedlings were immersed for one hour in the appropriate rhizobial cell suspension, and then planted in a perlite:sand (2:1) mixture. Plants were grown in a greenhouse for five to six weeks, with occasional watering with a 1 g L^{-1} nutrient solution (PlantProd solution [N-P-K, 0-15-40; Fertil]).

For whole nodule samples of wild type plants, pink nodules were collected from 53-60 plants per replicate 34-35 days post-inoculation; in the case of *dnf* mutants (and a matched wild-type A17 sample), nodules were collected from ~ 105 plants per genotype 23-24 days post-inoculation. Nodules were collected from the roots and stored in tubes in liquid nitrogen until collection was complete, at which point they were stored at -80°C until use. For sectioned nodule samples, pink nodules were collected from 103 *M. sativa* plants for each of the microsymbionts 35 to 40 days post-inoculation. Nodules were manually sectioned at the white-pink border. Nodule sections were stored in tubes over dry ice or liquid nitrogen until collection was complete, at which point they were stored at -80°C until use. Average plant shoot dry weights for all samples are listed in **Table S8**. Bacteroids were isolated from the nodule samples using Percoll gradient centrifugation as described elsewhere (Mergaert et al., 2006). The recovered bacteroids were resuspended in 50-100 µL of Bacteroid Extraction Buffer (BEB; 125 mM KCl, 50 mM Na-

succinate, 50 mM TES, pH 7.0), and either used immediately for microscopy, flow cytometry, and

DNA isolation, or stored at -80°C until use.

Nucleic acids were initially purified from most bacteroid samples using Epicentre

MasterPure™ Complete DNA and RNA Purification Kit, following the protocol for DNA isolation

from cell samples; the exceptions were bacteroids purified from *dnf* mutant nodules (and the

matched wild-type A17 sample), for which nucleic acids were isolated by using phenol:chloroform

extractions followed by ammonium acetate DNA precipitations as described elsewhere (Cowie et

al., 2006). For sectioned nodule samples, pure DNA was isolated by using the manufacturer's

protocol for the complete removal of RNA. For whole nodule samples, the isolated DNA was

further purified by treating the nucleic acids samples with RNase A, after which pure DNA was

isolated by using phenol:chloroform extractions followed by ammonium acetate DNA

precipitations or alternatively using the MasterPure™ DNA clean-up protocol for the DNA from

dnf mutant nodules and the matched wild-type A17 sample. In all cases, the final DNA pellets

were resuspended in 200 µL of 10 mM Tris-HCl, pH 8.5. Three biological replicates were

performed for bacteroids isolated from most whole nodules, whereas only one replicate was

performed for bacteroids isolated from sectioned nodules or *dnf* mutant nodules (and the matched

wild-type A17 sample) due to low quantities of starting materials.

DNA sequencing, modification detection, and motif analysis

DNA sequencing was performed at the U.S. Department of Energy Joint Genome Institute (JGI)

or in-house at the University of Florence (the stationary phase samples and *dnf* mutant nodules and

the matched wild-type A17 sample) using Pacific Biosciences (PacBio) sequencing technology

(Eid et al., 2009). Genomic DNA was sheared to 3 kb using a Covaris LS220 (Covaris Inc.,

Woburn, MA, USA) or 15 kb (for stationary phase samples and bacteroids isolated from *dnf* mutant

nodules and the matched wild-type A17 sample) using g-TUBEs (Covaris Inc., Woburn, MA,

USA). Sheared DNA was treated with exonuclease to remove single-stranded ends and DNA

damage repair mix followed by end repair and ligation of barcoded blunt adapters using SMRTbell

Template Prep Kit 2.0 (PacBio, Menlo Park, CA, USA). Libraries were purified with AMPure PB

beads (Beckman Coulter, Brea, CA, USA) and three or eight libraries with different barcodes were

pooled at equimolar ratios and purified with AMPure PB beads. For most samples, SMRTbell

template libraries were prepared using a Sequel Binding Kit 3.0 (PacBio, Menlo Park, CA, USA),

and sequenced on a Sequel instrument using a v3 or v4 sequencing primer, 1M v3 SMRT cells,

and Version 3.0 sequencing chemistry with 1x360 or 1x600 sequencing movie run times. The

exceptions were the *E. meliloti* Rm2011 zone II and *E. meliloti* FSM-MA zone III bacteroid

samples. For these samples, SMRTbell template libraries were prepared using a Sequel II Binding

Kit 2.0 (PacBio, Menlo Park, CA, USA), and then sequenced on a Sequel II instrument using the

tbd-sample dependent sequencing primer, 8M v1 SMRT cells, and Version 2.0 sequencing

chemistry with 1x900 sequencing movie run times.

DNA modification detection and motif analysis were performed using the PacBio SMRT

Link software (PacBio, Menlo Park, CA, USA). Briefly, raw reads were filtered using SFilter to

remove short reads and reads derived from sequencing adapters. Filtered reads were aligned

against the appropriate reference genome (**Table S2**) using BLASR (Chaisson and Tesler, 2012)

and modified sites were then identified through kinetic analysis of the aligned DNA sequence data

(Flusberg et al., 2010); the number of mapped bases per sample is provided in **Table S2**. Modified

sites were then grouped into motifs using MotifFinder. These motifs represent the recognition

sequences of MTase genes active in the genome (Clark et al., 2012). Downstream analyses were

performed using custom Perl and R scripts.

Flow cytometry

Flow cytometry was performed as described previously (Mergaert et al., 2006). Freshly prepared

bacteroid samples were diluted in 200 µL of BEB, heat-treated for 10 minutes in a 70°C water

bath, and then stained with the DNA-binding dye diamidino-2-phenylindole (DAPI). Cell size and

ploidy level of the bacteroid samples were determined using flow cytometry with a Beckman

Coulter CytoFLEX S instrument. Measurements consisted of 50,000 cells. Data analysis was

performed using the CytExpert 2.2.0.97 software.

Fluorescence microscopy

One µL of each freshly prepared bacteroid sample was mixed with 1 µL of 50 µg mL^{-1} DAPI or

with both 1 µL of 50 µg mL^{-1} DAPI and 1 µL of 100 µg mL^{-1} propidium iodide (PI), which are

both DNA binding dyes. Samples were visualized at 100x magnification under oil immersion using

a Nikon Eclipse 80*i* fluorescence microscope with the NIS-Elements BR 4.00.01 software and a

Digital Sight DS-U3 camera.

Phylogenetic analysis

The nucleotide fasta files of representative *Ensifer* species were downloaded from the National

Centre for Biotechnology Information (NCBI) Genome database. A core gene phylogeny was

constructed using a previously prepared pipeline (diCenzo et al., 2018) reliant on the use of Roary

3.11.3 (Page et al., 2015), Prokka 1.12-beta (Seemann, 2014), PRANK 140110 (Löytynoja, 2014),

trimAl (Capella-Gutiérrez et al., 2009), and RAxML 8.2.9 (Stamatakis, 2014). The phylogeny was

visualized with the iTol webserver (Letunic and Bork, 2016). Identification of nodulation

(*nodABC*) and nitrogenase genes (*nifHDK*) was performed with a published pipeline (diCenzo et

al., 2018) reliant on the use of HMMER 3.1b2 (Eddy, 2009), and the Pfam-A 31.0 (Finn et al.,

2016) and TIGERFAM 15.0 (Haft et al., 2013) databases.

DATA AVAILABILITY

Most sequencing data is available through the JGI Genome Portal (genome.jgi.doe.gov/portal/)

under Proposal 503835, as well as through NCBI (see **Table S2** for BioSample accessions). The

data for stationary phase cultures and bacteroids isolated from *dnf* mutant nodules are available

only through the NCBI (BioProject accessions PRJNA706182 and PRJNA705832; see **Table S2**

for BioSample accessions). All custom scripts to perform the analyses described in this study are

available through GitHub (github.com/diCenzo-Lab/003_2021_Ensifer_DNA_methylation), as

are the flow cytometry FCS files.

ACKNOWLEDGMENTS

We are grateful to E. Mullins (Teagasc, MTA2018233) for *E. adhaerens* OV14 and M. Bourge

(I2BC) for help with the flow cytometry experiments.

CONFLICT OF INTERESTS

The authors declare that they have no conflict of interest.

FUNDING

Institute (JGI), through the CSP New Investigator program (proposal: 503835). The work

conducted by the U.S. Department of Energy Joint Genome Institute, a DOE Office of Science

User Facility, is supported under Contract No. DE-AC02-05CH11231. This work was partially

supported by the "MICRO4Legumes" grant (Italian Ministry of Agriculture), by the grant

"Dipartimento di Eccellenza 2018–2022" from the Italian Ministry of Education, University and

Research (MIUR), and by a Discovery Grant from Natural Sciences and Engineering Research

Council of Canada. The present work benefited from the Imagerie-Gif core facility supported by

the Agence Nationale de la Recherche (ANR-11-EQPX-0029/Morphoscope, ANR-10-INBS-

04/FranceBioImaging, ANR-11-IDEX-0003-02/Saclay Plant Sciences). G.C.D was supported by

a NSERC postdoctoral fellowship and a European Molecular Biology Organization (EMBO) short

term fellowship for part of this work. L.C. was supported by the MICRO4Legumes grant (Italian

Ministry of Agriculture). Q.N. was supported by a PhD fellowship from the Paris-Saclay

University. J.H.T.C. was supported by a NSERC Undergraduate Summer Research Award. P.M.

and B.A. were supported by Saclay Plant Sciences (SPS) and grant ANR-17-CE20-0011 from the

Agence Nationale de la Recherche.

REFERENCES

Adhikari S, Curtis PD. 2016. DNA methyltransferases and epigenetic regulation in bacteria. *FEMS*

Microbiol Rev **40**:575–591. doi:10.1093/femsre/fuw023

Atack JM, Tan A, Bakaletz LO, Jennings MP, Seib KL. 2018. Phasevarions of bacterial pathogens:

methylomics sheds new light on old enemies. *Trends Microbiol* **26**:715–726.

doi:10.1016/j.tim.2018.01.008

Barnett MJ, Toman CJ, Fisher RF, Long SR. 2004. A dual-genome Symbiosis Chip for coordinate

751 study of signal exchange and development in a prokaryote–host interaction. *Proc Natl Acad*

752 *Sci USA* **101**:16636–16641.

753 Barrière Q, Guefrachi I, Gully D, Lamouche F, Pierre O, Fardoux J, Chaintreuil C, Alunni B,

754 Timchenko T, Giraud E, Mergaert P. 2017. Integrated roles of BclA and DD-

755 carboxypeptidase 1 in *Bradyrhizobium* differentiation within NCR-producing and NCR-

756 lacking root nodules. *Sci Rep* **7**:9063. doi:10.1038/s41598-017-08830-0

757 Blow MJ, Clark TA, Daum CG, Deutschbauer AM, Fomenkov A, Fries R, Froula J, Kang DD,

758 Malmstrom RR, Morgan RD, Pósfai J, Singh K, Visel A, Wetmore K, Zhao Z, Rubin EM,

759 Korlach J, Pennacchio LA, Roberts RJ. 2016. The epigenomic landscape of prokaryotes.

760 *PLOS Genet* **12**:e1005854. doi:10.1371/journal.pgen.1005854

761 Bourcy M, Brocard L, Pislariu CI, Cosson V, Mergaert P, Tadege M, Mysore KS, Udvardi MK,

762 Gourion B, Ratet P. 2013. *Medicago truncatula* DNF2 is a PI-PLC-XD-containing protein

763 required for bacteroid persistence and prevention of nodule early senescence and defense-

764 like reactions. *New Phytol* **197**:1250–1261. doi:10.1111/nph.12091

765 Brilli M, Fondi M, Fani R, Mengoni A, Ferri L, Bazzicalupo M, Biondi EG. 2010. The diversity

766 and evolution of cell cycle regulation in alpha-proteobacteria: a comparative genomic

767 analysis. *BMC Syst Biol* **4**:52. doi:10.1186/1752-0509-4-52

768 Brumwell SL, MacLeod MR, Huang T, Cochrane RR, Meaney RS, Zamani M, Matysiakiewicz

769 O, Dan KN, Janakirama P, Edgell DR, Charles TC, Finan TM, Karas BJ. 2019. Designer

770 *Sinorhizobium meliloti* strains and multi-functional vectors enable direct inter-kingdom

771 DNA transfer. *PLOS One* **14**:e0206781. doi:10.1371/journal.pone.0206781

772 Campbell JL, Kleckner N. 1990. *E. coli oriC* and the *dnaA* gene promoter are sequestered from

773 *dam* methyltransferase following the passage of the chromosomal replication fork. *Cell*

62:967–979. doi:10.1016/0092-8674(90)90271-F

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated alignment

trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**:1972–1973.

doi:10.1093/bioinformatics/btp348

Chaisson MJ, Tesler G. 2012. Mapping single molecule sequencing reads using basic local

alignment with successive refinement (BLASR): application and theory. *BMC Bioinform*

13:238. doi:10.1186/1471-2105-13-238

Clark TA, Murray IA, Morgan RD, Kislyuk AO, Spittle KE, Boitano M, Fomenkov A, Roberts

RJ, Korlach J. 2012. Characterization of DNA methyltransferase specificities using single-

molecule, real-time DNA sequencing. *Nucleic Acids Res* **40**:e29–e29.

doi:10.1093/nar/gkr1146

Cowie A, Cheng J, Sibley CD, Fong Y, Zaheer R, Patten CL, Morton RM, Golding GB, Finan

TM. 2006. An integrated approach to functional genomics: construction of a novel reporter

gene fusion library for *Sinorhizobium meliloti*. *Appl Environ Microbiol* **72**:7156–7167.

doi:10.1128/AEM.01397-06

Czernic P, Gully D, Cartieaux F, Moulin L, Guefrachi I, Patrel D, Pierre O, Fardoux J, Chaintreuil

C, Nguyen P, Gressent F, Da Silva C, Poulain J, Wincker P, Rofidal V, Hem S, Barrière

Q, Arrighi J-F, Mergaert P, Giraud E. 2015. Convergent evolution of endosymbiont

differentiation in Dalbergioid and inverted repeat-lacking clade legumes mediated by

nodule-specific cysteine-rich peptides. *Plant Physiol* **169**:1254–1265.

doi:10.1104/pp.15.00584

Davis-Richardson AG, Russell JT, Dias R, McKinlay AJ, Canepa R, Fagen JR, Rusoff KT, Drew

JC, Kolaczkowski B, Emerich DW, Triplett EW. 2016. Integrating DNA methylation and

gene expression data in the development of the soybean-*Bradyrhizobium* N_2-fixing

symbiosis. *Front Microbiol* **7**:518. doi:10.3389/fmicb.2016.00518

De Nisco NJ, Abo RP, Wu CM, Penterman J, Walker GC. 2014. Global analysis of cell cycle gene

expression of the legume symbiont *Sinorhizobium meliloti*. *Proc Natl Acad Sci USA*

111:3217–3224. doi:10.1073/pnas.1400421111

diCenzo GC, Debiec K, Krzysztoforski J, Uhrynowski W, Mengoni A, Fagorzi C, Gorecki A,

Dziewit L, Bajda T, Rzepa G, Drewniak L. 2018. Genomic and biotechnological

characterization of the heavy-metal resistant, arsenic-oxidizing bacterium *Ensifer* sp. M14.

Genes **9**:379. doi:10.3390/genes9080379

diCenzo GC, MacLean AM, Milunovic B, Golding GB, Finan TM. 2014. Examination of

prokaryotic multipartite genome evolution through experimental genome reduction. *PLOS*

Genet **10**:e1004742. doi:10.1371/journal.pgen.1004742

diCenzo GC, Muhammed Z, Østerås M, O'Brien SAP, Finan TM. 2017. A key regulator of the

glycolytic and gluconeogenic central metabolic pathways in *Sinorhizobium meliloti*.

Genetics **207**:961–974. doi:10.1534/genetics.117.300212

diCenzo GC, Tesi M, Pfau T, Mengoni A, Fondi M. 2020. Genome-scale metabolic reconstruction

of the symbiosis between a leguminous plant and a nitrogen-fixing bacterium. *Nat Commun*

11:2574. doi:10.1101/765271

diCenzo GC, Zamani M, Checcucci A, Fondi M, Griffitts J, Finan TM, Mengoni A. 2019. Multi-

disciplinary approaches for studying rhizobium - legume symbioses. *Can J Microbiol*

65:1–33. doi:10.1139/cjm-2018-0377

Domonkos A, Horvath B, Marsh JF, Halasz G, Ayaydin F, Oldroyd GED, Kalo P. 2013. The

identification of novel loci required for appropriate nodule development in *Medicago*

truncatula. BMC Plant Biol **13**:157. doi:10.1186/1471-2229-13-157

Eddy SR. 2009. A new generation of homology search tools based on probabilistic inference.

Genome Inform **23**:205–211. doi:10.1142/9781848165632_0019

Eid J, Fehr A, Gray J, Luong K, Lyle J, Otto G, Peluso P, Rank D, Baybayan P, Bettman B, Bibillo

A, Bjornson K, Chaudhuri B, Christians F, Cicero R, Clark S, Dalal R, deWinter A, Dixon

J, Foquet M, Gaertner A, Hardenbol P, Heiner C, Hester K, Holden D, Kearns G, Kong X,

Kuse R, Lacroix Y, Lin S, Lundquist P, Ma C, Marks P, Maxham M, Murphy D, Park I,

Pham T, Phillips M, Roy J, Sebra R, Shen G, Sorenson J, Tomaney A, Travers K, Trulson

M, Vieceli J, Wegener J, Wu D, Yang A, Zaccarin D, Zhao P, Zhong F, Korlach J, Turner

S. 2009. Real-time DNA sequencing from single polymerase molecules. *Science* **323**:133–

138. doi:10.1126/science.1162986

Fagorzi C, Ilie A, Decorosi F, Cangioli L, Viti C, Mengoni A, diCenzo GC. 2020. Symbiotic and

nonsymbiotic members of the genus *Ensifer* (syn. *Sinorhizobium*) are separated into two

clades based on comparative genomics and high-throughput phenotyping. *Genome Biol*

Evol **12**:2521–2534. doi:10.1093/gbe/evaa221

Farkas A, Maróti G, Dürgő H, Györgypál Z, Lima RM, Medzihradszky KF, Kereszt A, Mergaert

P, Kondorosi E. 2014. *Medicago truncatula* symbiotic peptide NCR247 contributes to

bacteroid differentiation through multiple mechanisms. *Proc Natl Acad Sci USA*

111:5183–5188. doi:10.1073/pnas.1404169111

Ferri L, Gori A, Biondi EG, Mengoni A, Bazzicalupo M. 2010. Plasmid electroporation of

Sinorhizobium strains: The role of the restriction gene *hsdR* in type strain Rm1021. *Plasmid*

63:128–135. doi:10.1016/j.plasmid.2010.01.001

Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, Potter SC, Punta M, Qureshi

M, Sangrador-Vegas A, Salazar GA, Tate J, Bateman A. 2016. The Pfam protein families

database: towards a more sustainable future. *Nucleic Acids Res* **44**:D279–D285.

doi:10.1093/nar/gkv1344

Fioravanti A, Fumeaux C, Mohapatra SS, Bompard C, Brilli M, Frandi A, Castric V, Villeret V,

Viollier PH, Biondi EG. 2013. DNA binding of the cell cycle transcriptional regulator

GcrA depends on N6-adenosine methylation in *Caulobacter crescentus* and other

Alphaproteobacteria. *PLOS Genet* **9**:e1003541. doi:10.1371/journal.pgen.1003541

Flusberg BA, Webster DR, Lee JH, Travers KJ, Olivares EC, Clark TA, Korlach J, Turner SW.

2010. Direct detection of DNA methylation during single-molecule, real-time sequencing.

Nat Methods **7**:461–465. doi:10.1038/nmeth.1459

Frage B, Döhlemann J, Robledo M, Lucena D, Sobetzko P, Graumann PL, Becker A. 2016.

Spatiotemporal choreography of chromosome and megaplasmids in the *Sinorhizobium*

meliloti cell cycle. *Mol Microbiol* **100**:808–823. doi:10.1111/mmi.13351

Gibson KE, Campbell GR, Lloret J, Walker GC. 2006. CbrA is a stationary-phase regulator of cell

surface physiology and legume symbiosis in *Sinorhizobium meliloti*. *J Bacteriol* **188**:4508–

4521. doi:10.1128/JB.01923-05

Gonzalez D, Collier J. 2013. DNA methylation by CcrM activates the transcription of two genes

required for the division of *Caulobacter crescentus*: regulation of cell division by DNA

methylation. *Mol Microbiol* **88**:203–218. doi:10.1111/mmi.12180

Gonzalez D, Kozdon JB, McAdams HH, Shapiro L, Collier J. 2014. The functions of DNA

methylation by CcrM in *Caulobacter crescentus*: a global approach. *Nucleic Acids Res*

42:3720–3735. doi:10.1093/nar/gkt1352

Gopalakrishnan S, Van Emburgh BO, Robertson KD. 2008. DNA methylation in development and

human disease. *Mutat Res* **647**:30–38. doi:10.1016/j.mrfmmm.2008.08.006

Greenberg MVC, Bourc'his D. 2019. The diverse roles of DNA methylation in mammalian

development and disease. *Nat Rev Mol Cell Biol* **20**:590–607. doi:10.1038/s41580-019-

0159-6

Haakonsen DL, Yuan AH, Laub MT. 2015. The bacterial cell cycle regulator GcrA is a σ [70]

cofactor that drives gene expression from a subset of methylated promoters. *Genes Dev*

29:2272–2286. doi:10.1101/gad.270660.115

Haft DH, Selengut JD, Richter RA, Harkins D, Basu MK, Beck E. 2013. TIGRFAMs and genome

properties in 2013. *Nucleic Acids Res* **41**:D387–D395. doi:10.1093/nar/gks1234

Horváth B, Domonkos Á, Kereszt A, Sz\Hucs A, Ábrahám E, Ayaydin F, Bóka K, Chen Y, Chen

R, Murray JD, Udvardi MK, Kondorosi E, Kaló P. 2015. Loss of the nodule-specific

cysteine rich peptide, NCR169, abolishes symbiotic nitrogen fixation in the *Medicago*

truncatula dnf7 mutant. *Proceedings of the National Academy of Sciences* **112**:15232–

15237. doi:10.1073/pnas.1500777112

Kahng LS, Shapiro L. 2001. The CcrM DNA methyltransferase of *Agrobacterium tumefaciens* Is

essential, and Its activity is cell cycle regulated. *J Bacteriol* **183**:3065–3075.

doi:10.1128/JB.183.10.3065-3075.2001

Kang S, Lee H, Han JS, Hwang DS. 1999. Interaction of SeqA and Dam methylase on the

hemimethylated origin of *Escherichia coli* chromosomal DNA replication. *J Biol Chem*

274:11463–11468. doi:10.1074/jbc.274.17.11463

Kazmierczak T, Nagymihaly M, Lamouche F, Barrière Q, Guefrachi I, Alunni B, Ouadghiri M,

Ibijbijen J, Kondorosi E, Mergaert P, Gruber V. 2017. Specific host-responsive

associations between *Medicago truncatula* accessions and *Sinorhizobium strains*. *Mol*

Plant Microbe Interact **30**:399–409. doi:10.1094/MPMI-01-17-0009-R

Kim M, Chen Y, Xi J, Waters C, Chen R, Wang D. 2015. An antimicrobial peptide essential for

bacterial survival in the nitrogen-fixing symbiosis. *Proceedings of the National Academy*

of Sciences **112**:15238–15243. doi:10.1073/pnas.1500123112

Kobayashi H, De Nisco NJ, Chien P, Simmons LA, Walker GC. 2009. *Sinorhizobium meliloti*

CpdR1 is critical for co-ordinating cell cycle progression and the symbiotic chronic

infection. *Mol Microbiol* **73**:586–600. doi:10.1111/j.1365-2958.2009.06794.x

Kozdon JB, Melfi MD, Luong K, Clark TA, Boitano M, Wang S, Zhou B, Gonzalez D, Collier J,

Turner SW, Korlach J, Shapiro L, McAdams HH. 2013. Global methylation state at base-

pair resolution of the *Caulobacter* genome throughout the cell cycle. *Proc Natl Acad Sci*

USA **110**:E4658–E4667. doi:10.1073/pnas.1319315110

Krall L, Wiedemann U, Unsin G, Weiss S, Domke N, Baron C. 2002. Detergent extraction

identifies different VirB protein subassemblies of the type IV secretion machinery in the

membranes of *Agrobacterium tumefaciens*. *Proc Natl Acad Sci USA* **99**:11405–11410.

doi:10.1073/pnas.172390699

Lahue R, Au K, Modrich P. 1989. DNA mismatch correction in a defined system. *Science*

245:160–164. doi:10.1126/science.2665076

Lang C, Long SR. 2015. Transcriptomic analysis of *Sinorhizobium meliloti* and *Medicago*

truncatula symbiosis using nitrogen nixation–deficient nodules. *Mol Plant Microbe*

Interact **28**:856–868. doi:10.1094/MPMI-12-14-0407-R

Letunic I, Bork P. 2016. Interactive tree of life (iTOL) v3: an online tool for the display and

annotation of phylogenetic and other trees. *Nucleic Acids Res* **44**:W242–W245.

doi:10.1093/nar/gkw290

Löytynoja A. 2014. Phylogeny-aware alignment with PRANK. In: Russell D, editor. Methods in

913   Molecular Biology (Methods and Protocols). Totowa, NJ: Humana Press. pp. 155–170.

914   Mergaert P, Uchiumi T, Alunni B, Evanno G, Cheron A, Catrice O, Mausset A-E, Barloy-Hubler

915   F, Galibert F, Kondorosi A, Kondorosi E. 2006. Eukaryotic control on bacterial cell cycle

916   and differentiation in the *Rhizobium*-legume symbiosis. *Proc Natl Acad Sci USA*

917   **103**:5230–5235. doi:10.1073/pnas.0600912103

918   Mohapatra SS, Fioravanti A, Biondi EG. 2014. DNA methylation in *Caulobacter* and other

919   Alphaproteobacteria during cell cycle progression. *Trends Microbiol* **22**:528–535.

920   doi:10.1016/j.tim.2014.05.003

921   Moreau D, Voisin A-S, Salon C, Munier-Jolain N. 2008. The model symbiotic association between

922   *Medicago truncatula* cv. Jemalong and *Rhizobium meliloti* strain 2011 leads to N-stressed

923   plants when symbiotic N2 fixation is the main N source for plant growth. *J Exp Bot*

924   **59**:3509–3522. doi:10.1093/jxb/ern203

925   Nagymihaly M, Veluchamy A, Györgypál Z, Ariel F, Jégu T, Benhamed M, Sz\Hucs A, Kereszt

926   A, Mergaert P, Kondorosi E. 2017. Ploidy-dependent changes in the epigenome of

927   symbiotic cells correlate with specific patterns of gene expression. *Proc Natl Acad Sci USA*

928   **114**:4543–4548. doi:10.1073/pnas.1704211114

929   Nouri H, Monnier A-F, Fossum-Raunehaug S, Maciąg-Dorszyńska M, Cabin-Flaman A, Képès F,

930   Węgrzyn G, Szalewska-Pałasz A, Norris V, Skarstad K, Janniere L. 2018. Multiple links

931   connect central carbon metabolism to DNA replication initiation and elongation in *Bacillus*

932   *subtilis*. *DNA Res* **25**:641–653. doi:10.1093/dnares/dsy031

933   Oldroyd GED. 2013. Speak, friend, and enter: signalling systems that promote beneficial

934   symbiotic associations in plants. *Nat Rev Microbiol* **11**:252–263. doi:10.1038/nrmicro2990

Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MTG, Fookes M, Falush D, Keane

JA, Parkhill J. 2015. Roary: rapid large-scale prokaryote pan genome analysis.

Bioinformatics **31**:3691–3693. doi:10.1093/bioinformatics/btv421

Pecrix Y, Staton SE, Sallet E, Lelandais-Brière C, Moreau S, Carrère S, Blein T, Jardinaud M-F,

Latrasse D, Zouine M, Zahm M, Kreplak J, Mayjonade B, Satgé C, Perez M, Cauet S,

Marande W, Chantry-Darmon C, Lopez-Roques C, Bouchez O, Bérard A, Debellé F,

Muños S, Bendahmane A, Bergès H, Niebel A, Buitink J, Frugier F, Benhamed M, Crespi

M, Gouzy J, Gamas P. 2018. Whole-genome landscape of *Medicago truncatula* symbiotic

genes. *Nat Plants* **4**:1017–1025. doi:10.1038/s41477-018-0286-7

Penterman J, Abo RP, De Nisco NJ, Arnold MFF, Longhi R, Zanda M, Walker GC. 2014. Host

plant peptides elicit a transcriptional response to control the *Sinorhizobium meliloti* cell

cycle during symbiosis. *Proc Natl Acad Sci USA* **111**:3561–3566.

doi:10.1073/pnas.1400450111

Pini F, De Nisco NJ, Ferri L, Penterman J, Fioravanti A, Brilli M, Mengoni A, Bazzicalupo M,

Viollier PH, Walker GC, Biondi EG. 2015. Cell cycle control by the master regulator CtrA

in *Sinorhizobium meliloti*. *PLOS Genet* **11**:e1005232.

Pini F, Frage B, Ferri L, De Nisco NJ, Mohapatra SS, Taddei L, Fioravanti A, Dewitte F, Galardini

M, Brilli M, Villeret V, Bazzicalupo M, Mengoni A, Walker GC, Becker A, Biondi EG.

2013. The DivJ, CbrA and PleC system controls DivK phosphorylation and symbiosis in

Sinorhizobium meliloti. *Mol Microbiol* **90**:54–71. doi:10.1111/mmi.12347

Rasmussen T, Jensen RB, Skovgaard O. 2007. The two chromosomes of *Vibrio cholerae* are

initiated at different time points in the cell cycle. *EMBO J* **26**:3124–3131.

doi:10.1038/sj.emboj.7601747

Roux B, Rodde N, Jardinaud M-F, Timmers T, Sauviac L, Cottret L, Carrère S, Sallet E, Courcelle

E, Moreau S, Debellé F, Capela D, de Carvalho-Niebel F, Gouzy J, Bruand C, Gamas P.

2014. An integrated analysis of plant and bacterial gene expression in symbiotic root

nodules using laser-capture microdissection coupled to RNA sequencing. *Plant J* **77**:817–

837. doi:10.1111/tpj.12442

Sánchez-Romero MA, Cota I, Casadesús J. 2015. DNA methylation in bacteria: from the methyl

group to the methylome. *Curr Opin Microbiol* **25**:9–16. doi:10.1016/j.mib.2015.03.004

Satgé C, Moreau S, Sallet E, Lefort G, Auriac M-C, Remblière C, Cottret L, Gallardo K, Noirot

C, Jardinaud M-F, Gamas P. 2016. Reprogramming of DNA methylation is critical for

nodule development in *Medicago truncatula*. *Nat Plants* **2**:16166.

doi:10.1038/nplants.2016.166

Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**:2068–2069.

doi:10.1093/bioinformatics/btu153

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* **30**:1312–1313. doi:10.1093/bioinformatics/btu033

Starker CG, Parra-Colmenares AL, Smith L, Mitra RM, Long SR. 2006. Nitrogen fixation mutants

of *Medicago truncatula* fail to support plant and bacterial symbiotic gene expression. *Plant*

Physiol **140**:671–680. doi:10.1104/pp.105.072132

Tang Y, Gao X-D, Wang Y, Yuan B-F, Feng Y-Q. 2012. Widespread existence of cytosine

methylation in yeast DNA measured by gas chromatography/mass spectrometry. *Anal*

Chem **84**:7249–7255. doi:10.1021/ac301727c

Van de Velde W, Zehirov G, Szatmari A, Debreczeny M, Ishihara H, Kevei Z, Farkas A, Mikulass

K, Nagy A, Tiricz H, Satiat-Jeunemaître B, Alunni B, Bourge M, Kucho K, Abe M, Kereszt

A, Maróti G, Uchiumi T, Kondorosi E, Mergaert P. 2010. Plant peptides govern terminal

differentiation of bacteria in symbiosis. *Science* **327**:1122–1126.

doi:10.1126/science.1184057

Vasse J, De Billy F, Camut S, Truchet G. 1990. Correlation between ultrastructural differentiation

of bacteroids and nitrogen fixation in alfalfa nodules. *J Bacteriol* **172**:4295–4306.

doi:10.1128/jb.172.8.4295-4306.1990

Vasu K, Nagaraja V. 2013. Diverse functions of restriction-modification systems in addition to

cellular defense. *Microbiol Mol Biol Rev* **77**:53–72. doi:10.1128/MMBR.00044-12

Wang D, Griffitts J, Starker C, Fedorova E, Limpens E, Ivanov S, Bisseling T, Long S. 2010. A

nodule-specific protein secretory pathway required for nitrogen-nixing symbiosis. *Science*

327:1126–1129. doi:10.1126/science.1184096

Wang ET, Young JPW. 2019. History of rhizobial taxonomy In: Wang ET, Tian CF, Chen WF,

Young JPW, Chen WX, editors. Ecology and Evolution of Rhizobia: Principles and

Applications. Singapore: Springer. pp. 23–39. doi:10.1007/978-981-32-9555-1_2

Wright R, Stephens C, Shapiro L. 1997. The CcrM DNA methyltransferase is widespread in the

alpha subdivision of proteobacteria, and its essential functions are conserved in *Rhizobium*

meliloti and *Caulobacter crescentus*. *J Bacteriol* **179**:5869–5877.

Wright R, Stephens C, Zweiger G, Shapiro L, Alley MR. 1996. *Caulobacter* Lon protease has a

critical role in cell-cycle control of DNA methylation. *Genes Dev* **10**:1532–1542.

doi:10.1101/gad.10.12.1532

Zhang H, Lang Z, Zhu J-K. 2018. Dynamics and function of DNA methylation in plants. *Nat Rev*

Mol Cell Biol **19**:489–506. doi:10.1038/s41580-018-0016-z

Zweiger G, Marczynski G, Shapiro L. 1994. A *Caulobacter* DNA methyltransferase that functions

only in the predivisional cell. *J Mol Biol* **235**:472–485. doi:10.1006/jmbi.1994.1007

1005

                                      TABLES AND FIGURES

1007   **Table 1.** Methylated motifs identified in this study.

Motif*	Type†	Count¥	Frequency (motifs/kb)
E. meliloti 2011			
G**A**NTC CTN**A**G	m6A	11,169	1.67
RCG**C**CTC YGCGGAG	m4C	3,943	0.59
CGC**A**(N5)GTG GCGT(N5)C**A**C	m6A	1,085	0.16
E. meliloti FSM-MA			
G**A**NTC CTN**A**G	m6A	11,215	1.67
TCG**A**(N8)TCGA AGCT(N8)**A**GCT	m6A	2,612	0.39
E. fredii NGR234			
G**A**NTC CTN**A**G	m6A	11,111	1.61
CAG**A**(N7)GTTG GTCT(N7)C**A**AC	m6A	188	0.03
E. adhaerens OV14			
G**A**NTC CTN**A**G	m6A	8,475	1.10
WNCCG**A**TG WNGGCTAC	m6A	4,596	0.60

1008   * The methylated nucleotides are indicated in boldface font.

1009   † Indicates whether the modification is a N^6-methyladenoside (m6A) or N^4-methylcytosine (m4C).

1010   ¥ The total times the motif appears in the genome, regardless of methylation status.

1011 **Table 2.** Relative sequencing depth of each *E. meliloti* replicon.

Strain	Condition	Relative mean sequencing depth *		
		Chromosome	pSymA	pSymB
Rm2011	Mid-exponential	1.00 ± 0.00	1.00 ± 0.03	0.98 ± 0.03
Rm2011	Stationary phase	1.00 ± 0.00	1.06 ± 0.02	1.02 ± 0.03
Rm2011	*M. sativa* bacteroids	1.00 ± 0.00	0.64 ± 0.02	0.62 ± 0.02
FSM-MA	Mid-exponential	1.00 ± 0.00	1.02 ± 0.03	0.87 ± 0.05
FSM-MA	Stationary phase	1.00 ± 0.00	1.10 ± 0.00	1.01 ± 0.03
FSM-MA	*M. sativa* bacteroids	1.00 ± 0.00	0.79 ± 0.04	0.71 ± 0.02
FSM-MA	*M. truncatula* bacteroids	1.00 ± 0.00	0.66 ± 0.02	0.59 ± 0.01
FSM-MA	*dnf1* bacterial cells	1.00	0.98	0.90
FSM-MA	*dnf5* bacterial dells	1.00	0.96	0.94
FSM-MA	*dnf2* bacterial dells	1.00	0.81	0.73
FSM-MA	*dnf7* bacterial dells	1.00	0.82	0.71
FSM-MA	*dnf4* bacterial dells	1.00	0.80	0.73
FSM-MA	A17 bacteroids	1.00	0.76	0.67

1012 * Sequencing depth is presented relative to the sequencing depth of the chromosome in the same

1013 sample. Values are the means of triplicate samples ± standard deviation, except for the third section

1014 of the table for which numbers are based on a single replicate.

Figure 1. Genome-wide DNA methylation of *E. meliloti* Rm2011. (**A-F**) The extent of methylation is shown, using a 10 kb sliding window, of GANTC sites (black) and CGCA(N$_5$)GTG sites (grey) across the chromosome (**A,D**), pSymB (**B,E**), and pSymA (**C,F**) replicons of exponential phase (**A-C**) or early stationary phase (**D-F**) *E. meliloti* Rm2011. Averages from three biological replicates are shown. The red (GANTC) and blue (CGCA(N$_5$)GTG) lines are polynomial regression lines calculated in R using the "rlm" method and the formula "y~poly(x,2)". (**G-I**) Cumulative GC skews, shown using a 10 kb sliding window, across the *E. meliloti* Rm2011 chromosome (**G**), pSymB (**H**), and pSymA (**I**) replicons.

1024

Figure 2. GANTC frequency in the genus *Ensifer*. (**A**) An unrooted maximum likelihood phylogeny of 17 representative *Ensifer* strains. The phylogeny represents the bootstrap best tree following 100 bootstrap replicates, prepared on the basis of the concatenated nucleotide alignments of 1566 core genes. Values represent the bootstrap support. N_2-fixing legume symbionts were identified by the presence of the symbiotic genes *nodABC* and *nifHDK*. They are indicated in blue, while red denotes non-symbiotic strains. The four wild-type strains used in this study are indicated with arrows. (**B**) Box plots summarizing the frequency of GANTC sites (presented as GANTC sites per kb) in 157 *Ensifer* strains is shown. The monophyletic "symbiotic" and "non-symbiotic" clades as defined previously (Fagorzi et al., 2020), are represented by 113 and 44 genomes respectively. The densities of GANTC sites across the entire genome (pink), within coding regions (blue), and within intergenic regions (yellow) are shown. Statistically different values ($p < 0.05$) are denoted by uppercase letters as determined by a one-way ANOVA followed by a Tukey's HSD post hoc test.

1038

Figure 3. Chromosome-wide DNA methylation of *E. meliloti* Rm2011 bacteroids. The extent

of methylation of (**A-E**) GANTC, (**F-J**) CGCA(N$_5$)GTG, and (**K-O**) RCGCCTC motifs across the

E. meliloti Rm2011 chromosome is shown using a 10 kb sliding window. Averages from three

biological replicates are shown for free-living and whole nodule samples; data represents one

replicate for the zone II and zone III nodule sections. (**A,F,K**) Free-living cells harvested in mid-

exponential phase. (**B,G,L**) Free-living cells harvested in early stationary phase. (**C,H,M**)

Bacteroids isolated from *M. sativa* zone II nodule sections. (**D,I,N**) Bacteroids isolated from *M.*

sativa zone III nodule sections. (**E,J,O**) Bacteroids isolated from *M. sativa* whole nodule samples.

The red lines are polynomial regression lines calculated in R using the "rlm" method and the

formula "y~poly(x,2)". Data for pSymB and pSymA are shown in Figures S14 and S15.

1049

Figure 4. Chromosome-wide DNA methylation of *E. meliloti* FSM-MA bacteroids. The extent of methylation of (**A-F**) GANTC and (**G-L**) TCGA(N$_8$)TCGA motifs across the *E. meliloti* FSM-MA chromosome is shown using a 10 kb sliding window. Averages from three biological replicates are shown for free-living and whole nodule samples; data represents one replicate for the zone II and zone III nodule sections. (**A,G**) Free-living cells harvested in mid-exponential phase. (**B,H**) Free-living cells harvested in early stationary phase. (**C,I**) Bacteroids isolated from *M. sativa* zone II nodule sections. (**D,J**) Bacteroids isolated from *M. sativa* zone III nodule sections. (**E,K**) Bacteroids isolated from *M. sativa* whole nodule samples. (**F,L**) Bacteroids isolated from *M. truncatula* whole nodule samples. The red lines are polynomial regression lines calculated in R using the "rlm" method and the formula "y~poly(x,2)". Data for pSymB and pSymA are shown in Figures S16 and S17.

1060

Figure 5. Bacteroid morphology and chromosomal GANTC methylation in *E. meliloti* bacteroids purified from *M. truncatula dnf* mutant nodules. Data is shown for *E. meliloti* FSM-MA bacteroids purified from (**A,G,M,S**) *M. truncatula dnf1* mutant nodules, (**B,H,N,T**) *M. truncatula dnf5* mutant nodules, (**C,I,O,U**) *M. truncatula dnf2* mutant nodules, (**D,J,P,V**) *M. truncatula dnf7* mutant nodules, (**E,K,Q,W**) *M. truncatula dnf4* mutant nodules, and (**F,L,R,X**)

M. truncatula A17 wild-type nodules. (**A-F**) Micrographs of *E. meliloti* FSM-MA bacteroids

stained with the DNA binding dye DAPI. The scale bar represents 30 μm. (**G-L**) Pseudo-coloured

scatterplots displaying the cell morphology (X-axis) and DNA content (Y-axis) of *E. meliloti*

FSM-MA bacteroids, as determined based on flow cytometry analysis of DAPI stained cells. The

red dashed ellipses indicate the position of undifferentiated bacteria as in culture (not shown) or in

the *dnf1* mutant nodules (lower left ellipse) or fully mature bacteroids as in the A17 wild-type

nodules (top right ellipse). (**M-X**) The extent of methylation of (**M-R**) GANTC or (**S-X**)

TCGA(N$_8$)TCGA motifs across the *E. meliloti* FSM-MA chromosome, shown using a 10 kb

sliding window. The red lines are polynomial regression lines calculated in R using the "rlm"

method and the formula "y~poly(x,2)". Data for pSymB and pSymA are shown in Figures S19

and S20.