Complete genome sequence of *Xylella taiwanensis* and comparative analysis

of virulence gene content with *Xylella fastidiosa*

Running Head: Genome analysis of *Xylella taiwanensis*

Ling-Wei Weng[1#], Yu-Chen Lin[2#], Chiou-Chu Su[3], Ching-Ting Huang[2], Shu-Ting

Cho[2], Ai-Ping Chen[2], Shu-Jen Chou[2], Chi-Wei Tsai[1*], Chih-Horng Kuo[2*]

[1] Department of Entomology, National Taiwan University, Taipei, Taiwan

[2] Institute of Plant and Microbial Biology, Academia Sinica, Taipei, Taiwan

[3] Division of Pesticide Application, Taiwan Agricultural Chemicals and Toxic

Substances Research Institute, Taichung, Taiwan

[#] Equal contribution

* Authors for Correspondence:

Chi-Wei Tsai; chiwei@ntu.edu.tw

Chih-Horng Kuo; chk@gate.sinica.edu.tw

Keywords: *Xylella*, Xanthomonadaceae, plant pathogens, pear leaf scorch,

genome, virulence

Abstract

The bacterial genus *Xylella* contains plant pathogens that are major threats to

agriculture in America and Europe. Although extensive research was conducted to

characterize different subspecies of *Xylella fastidiosa* (*Xf*), comparative analysis at

above-species levels were lacking due to the unavailability of appropriate data sets.

Recently, a bacterium that causes pear leaf scorch (PLS) in Taiwan was described

as the second *Xylella* species (i.e., *Xylella taiwanensis*; *Xt*). In this work, we report

the complete genome sequence of *Xt* type strain PLS229^T. The genome-scale

phylogeny provided strong support that *Xf* subspecies *pauca* (*Xfp*) is the basal

lineage of this species and *Xylella* was derived from the paraphyletic genus

Xanthomonas. Quantification of genomic divergence indicated that different *Xf*

subspecies share ~87-95% of their chromosomal segments, while the two *Xylella*

species share only ~66-70%. Analysis of overall gene content suggested that *Xt* is

most similar to *Xf* subspecies *sandyi* (*Xfs*). Based on the existing knowledge of *Xf*

virulence genes, the homolog distribution among 28 *Xylella* representatives was

examined. Among the 11 functional categories, those involved in secretion and

metabolism are the most conserved ones with no copy number variation. In contrast,

several genes related to adhesins, hydrolytic enzymes, and toxin-antitoxin systems

are highly variable in their copy numbers. Those virulence genes with high levels of

conservation or variation may be promising candidates for future studies. In

summary, the new genome sequence and analysis reported in this work contributed

to the study of several important pathogens in the family Xanthomonadaceae.

Contribution to the Field

Xylella fastidiosa is a plant-pathogenic bacterium with multiple subspecies that are major threats to agriculture in America and Europe. Although extensive research has been conducted, comparative analysis of this species with other bacteria is lacking due to the unavailability of known close relatives. In this work, we report the complete genome sequence of *Xylella taiwanensis*, a newly described species within the same genus. This new data set and our focused analysis helped to better understand the evolutionary relationships among different *Xylella* lineages and their genomic diversity. Moreover, detailed examination of their virulence genes identified those that are either highly conserved or variable, providing promising candidates for future studies to further investigate the molecular mechanisms of *Xylella* virulence.

Introduction

The gammaproteobacterium *Xylella fastidiosa* (*Xf*) is an insect-vectored plant

pathogen that resides in plant xylem and is fastidious (Wells et al., 1987). To date,

more than 563 plant species in 82 families have been reported as hosts for *Xf*

(European Food Safety Authority, 2018). *Xf* could be classified into at least five

subspecies; some of the notable examples include *Xf* subspecies *fastidiosa* (*Xff*) that

causes Pierce's disease (PD) of grapevine, *Xf* subspecies *pauca* (*Xfp*) that causes

citrus variegated chlorosis (CVC) and olive quick decline syndrome (OQDS), and *Xf*

subspecies *sandyi* (*Xfs*) that causes oleander leaf scorch (OLS). Because of their

economic and ecological impacts, substantial resources have been devoted to

related research. Notably, several large-scale studies were conducted to investigate

the genomic diversity and evolution of *Xf* (Denancé et al., 2019; Potnis et al., 2019;

Vanhove et al., 2019). Based on a comparison of 72 strains, the five *Xf* subspecies

harbor high levels of genetic diversity (Vanhove et al., 2019). With an average gene

content of ~2,150 per strain, the core genome (i.e., genes shared by >95% of the

strains) contains only ~900 genes, while the pangenome contains ~10,000 genes.

Moreover, although certain patterns of sequence divergence were found among

those subspecies (Denancé et al., 2019), extensive recombination occurred at the

levels of within- and between-subspecies (Potnis et al., 2019).

In contrast to the extensive genomic research at within-species level,

comparative studies of *Xf* at higher taxonomic levels are lacking. Under the current

taxonomy, *Xylella* belongs to the family Xanthomonadaceae and is most closely

related to *Xanthomonas* (Rodriguez-R et al., 2012; Anderson et al., 2013). However,

the genomic divergence between *Xylella* and *Xanthomonas* is very high in terms of

chromosomal organization, gene content, and sequence variation. Thus, extracting

biological insights from such comparisons is difficult. At within-genus level, *Xf* was

largely considered as the only species within this genus since it was formally

described in 1987 (Wells et al., 1987), which made between-species comparison

infeasible. Intriguingly, a *Xylella* lineage that causes pear leaf scorch (PLS) in

Taiwan was found to exhibit a slightly lower level of 16S rRNA gene sequence

identity at 97.8-98.6% when compared to different subspecies of *Xf* (Su et al., 2012).

In 2016, this PLS *Xylella* was formally reclassified as a novel species *Xylella*

taiwanensis (*Xt*) based on a polyphasic approach (Su et al., 2016). Although a draft

genome sequence of *Xt* was published earlier (Su et al., 2014), that draft assembly

was produced with only ~20-fold coverage of Roche/454 GS-FLX reads and is highly

fragmented (i.e., 85 contigs; N50 = 121 kb). Moreover, no comparative analysis of

gene content between *Xt* and *Xf* has been conducted.

To fill this gap, we determined the complete genome sequence of the type

strain of *Xt* (i.e., PLS229^T) for comparative analysis with its relatives. In addition to

providing a genome-level overview of their diversity and evolution, we utilized the

existing knowledge of *Xff* virulence genes and conducted detailed comparisons of

virulence gene content among different *Xylella* lineages.

Materials and Methods

The strain was acquired from the Bioresource Collection and Research

Centre (BCRC) in Taiwan (accession 80915). The procedures for genome

sequencing and comparative analysis were based on those described in our

previous studies (Lo et al., 2013; Lo et al., 2018; Cho et al., 2020). All bioinformatics

tools were used with the default settings unless stated otherwise.

Briefly, the strain was cultivated on PD2 medium as described (Su et al.,

2016) for DNA extraction using Wizard Genomic DNA Purification Kit (A1120;

Promega, USA). For Illumina sequencing, a paired-end library with a target insert

size of 550-bp was prepared using KAPA LTP Library Preparation Kit (KK8232;

Roche, Switzerland) without amplification, then sequenced using MiSeq Reagent

Nano Kit v2 (MS-103-1003; Illumina, USA) to obtain ~50X coverage. For Oxford

Nanopore Technologies (ONT) sequencing, the library was prepared using ONT

Ligation Kit (SQK-LSK109) and sequenced using MinION (FLO-MIN106; R9.4

chemistry and MinKNOW Core v3.6.0) to obtain ~228X coverage; Guppy v3.4.5 was

used for basecalling. The raw reads were combined for *de novo* assembly by using

Unicycler v0.4.8-beta (Wick et al., 2017). For validation, the Illumina and ONT raw

reads were mapped to the assembly using BWA v0.7.12 (Li and Durbin, 2009) and

Minimap2 v2.15 (Li, 2018), respectively. The results were programmatically checked

using SAMtools v1.2 (Li et al., 2009) and manually inspected using IGV v2.3.57

(Robinson et al., 2011). The finalized assembly was submitted to the National Center

for Biotechnology Information (NCBI) and annotated using their Prokaryotic Genome

Annotation Pipeline (PGAP) (Tatusova et al., 2016).

A total of 40 genomes, including 27 from *Xf*, were used for comparative

analysis (**Table 1**). Our taxon sampling mainly focused on the strains that could

represent the known *Xylella* diversity (Vanhove et al., 2019). Two other

Xanthomonadaceae genera were also included. For the closely-related

Xanthomonas, 10 species were selected to represent the key lineages (Parkinson et

al., 2009; Rodriguez-R et al., 2012). For the distantly-related *Pseudoxanthomonas*,

only two species were sampled.

Chromosomal level comparisons of nucleotide sequences were conducted

using fastANI v1.1 (Jain et al., 2018). Homologous gene clusters were identified

based on protein sequences using BLASTP v2.10.0+ (Camacho et al., 2009) and

OrthoMCL v1.3 (Li et al., 2003). For gene content comparisons, the homolog

clustering result was converted into a matrix of genomes by homolog clusters with

the value in each cell corresponding to the copy number. This matrix was converted

into a Jaccard distance matrix among genomes using VEGAN package v2.5-6 in R,

then processed using the principal coordinates analysis function in APE v5.4

(Paradis and Schliep, 2019) and visualized using ggplot2 v3.3.2 (Wickham, 2016).

For phylogenetic analysis, homologous sequences were aligned using MUSCLE

v3.8.31 (Edgar, 2004). The maximum likelihood inference was performed using

PhyML v.3.3.20180214 (Guindon and Gascuel, 2003); the proportion of invariable

sites and the gamma distribution parameter were estimated from the data set and

the number of substitute rate categories was set to four. The PROTDIST program of

PHYLIP v3.697 (Felsenstein, 1989) was used to calculate sequence similarities.

Results and Discussion

<u>Genome Characteristics</u>

Strain Xt PLS229^T has one 2,824,877-bp circular chromosome with 53.3%

G+C content; no plasmid was found. The annotation contains two complete sets of

16S-23S-5S rRNA genes, 49 tRNA genes, and 2,176 intact coding sequences

(CDSs). This genome size is near the upper range of those Xf representatives

(median: 2.54 Mb; range: 2.39-2.88 Mb) and much smaller compared to

$Xanthomonas$ spp. (median: 5.09 Mb; range: 3.76-5.35 Mb) (**Table 1**). Among all 40

representative Xanthomonadaceae genomes, the genome sizes and the numbers of

intact CDSs have a correlation coefficient of 0.989 ($p < 2.2e^{-16}$). Compared to those

Xf representatives with similar genome sizes (i.e., ~2.73-2.88 Mb), such as those

five *Xfp* strains from Brazil or the *Xfs* strain from the USA, the *Xt* PLS229^T genome

has fewer intact CDSs (i.e., 2,273-2,456 vs. 2,173) and fewer pseudogenes (i.e.,

153-242 vs. 132). It is unclear if these observations were caused by annotation

artifacts or have biological meaning.

<u>Molecular Phylogeny and Genome Divergence</u>

A total of 779 single-copy protein-coding genes were found to be shared by

the 40 Xanthomonadaceae genomes compared (**Table 1**). Based on the

concatenated alignment of the protein sequences derived from these genes, a robust

maximum likelihood phylogeny was inferred (**Figure 1**). The availability of this *Xt*

genome sequence provided a more appropriate outgroup to root the *Xf* phylogeny

and further supported that *Xfp* is the basal lineage (Denancé et al., 2019; Potnis et

al., 2019; Vanhove et al., 2019).

The genus *Xanthomonas* was known to be paraphyletic but the relationships

of its two major clades (i.e., represented by *Xanthomonas albilineans* and

Xanthomonas campestris, respectively) with *Xylella* were controversial (Pieretti et al.,

2009; Rodriguez-R et al., 2012). With our genome-scale phylogeny, it is clear that

Xylella is more closely related to *X. campestris* (**Figure 1**) and has experienced

genome reduction since their divergence (**Table 1**).

When the genetic divergence was measured by overall nucleotide sequence

conservation using fastANI (Jain et al., 2018), comparisons within each of the five *Xf*

subspecies found that 88.8-99.8% of the chromosomal segments are shared and

those segments have 98.5-100% average nucleotide identity (ANI) (**Figure 2**). For

between-subspecies comparisons, 86.6-94.8% of the chromosomal segments are

shared and those segments have 96.3-98.8% ANI. When those *Xf* subspecies were

compared to *Xt*, only 66.4-70.3% of the chromosomal segments are shared and

those segments have 82.9-83.4% ANI. These results are consistent with previous

findings (Su et al., 2016; Denancé et al., 2019) and provide further support to the

current taxonomy based on the 95% ANI threshold recommended for delineating

bacterial species (Jain et al., 2018).

Because the ANI approach provides low resolutions when the nucleotide

sequence identity drops to ~80% (Jain et al., 2018) and may not be appropriate for

cross-genus comparisons, we also evaluated divergence based on the protein

sequences of those 779 Xanthomonadaceae core genes. The two *Xylella* species

have ~88.8-89.1% protein sequence similarity, which is lower than the values

observed in the comparisons among those eight *X. campestris* clade representatives

(median: 93.8%; range: 92.6-97.2%), comparable to the *X. albilineans-X. hyacinthi*

comparison (88.6%), and higher than the *P. mexicana-P. spadix* comparison

(75.4%).

In addition to analysis of sequence divergence based on those 779 core

genes (**Figure 1**) and the entire chromosomes (**Figure 2**), the divergence in gene

content was also examined. The gene content comparisons were based on principal

coordinates analysis that examines copy number variation among all homologous

gene clusters in the entire pangenome and does not consider sequence divergence

within each homologous gene cluster. When all 40 Xanthomonadaceae genomes

were compared together based on their 11,455 homologous gene clusters, the

grouping patterns (**Figure 3A**) are consistent with the phylogenetic clades inferred

based on sequence divergence of the 779 core genes (**Figure 1A**). All 27 *Xf*

genomes form a tight cluster (**Figure 3A**) despite their differences in the number of

intact CDSs (range:1,911-2,456; av.± std. dev.: 2,156 ± 144) (**Table 1**). In contrast,

although the *Xt* genome has 2,176 intact CDSs, which is close to the average

observed among those 27 *Xf* representatives, it does not fall into the *Xf* cluster

(**Figure 3A**). This result indicates that the gene content divergence between these

two *Xylella* species is much higher than the divergence among *Xf* subspecies. For

the within-*Xylella* comparison based on 5,395 homologous gene clusters, the

grouping patterns are consistent with the taxonomic assignments and *Xt* is most

similar to *Xfs* (**Figure 3B**). It is interesting that *Xt* and *Xfs* are similar in having a

narrow host range (i.e., *Xt* is restricted to pear and *Xfs* is mostly known for oleander

infections), while other *Xf* subspecies can infect a wide range of hosts (Baldi and

Porta, 2017; European Food Safety Authority, 2018; Rapicavoli et al., 2018).

However, it is also important to note that the host range information may be limited

by sampling and experimental efforts. As more research results become available,

this information may be updated. For example, *Xfs*-related strains have been

reported to infect coffee (Jacques et al., 2016) and whether *Xt* can infect a wider

range of plants remains to be investigated.

Virulence Genes and Pathogenicity Factors

225   Based on the current knowledge of putative virulence genes and

pathogenicity factors identified in *Xf*, type IV pili (T4P) are important for twitching

motility and movements within infected plants (Meng et al., 2005; Li et al., 2007;

Burdman et al., 2011; Cursino et al., 2011; Hao et al., 2017). The *Xylella* T4P genes

are organized into four major gene clusters with 25 homologs and highly conserved

among the 28 representative genomes examined (**Figure 4**). Among these four T4P

gene clusters, cluster II that corresponds to the *pil-chp* operon (Cursino et al., 2011;

Hao et al., 2017) is the most conserved cluster with only two putative gene losses

(i.e., *chpB* in *Xfp* Salento-1 and *chpC* in *Xt*). A notable gene absence is PD1925 in

cluster IV, which encodes a hypothetical protein and is absent in all *Xfp* strains and

Xt. The only gene duplication observed involves a tandem duplication of the cluster

IV *pilA2* homolog in *Xt*. Intriguingly, *Xt* PLS229[T] is the only strain that has two copies

of *pilA2* and no *pilA1*. These two type IV pilin paralogs were shown to have different

functions in *Xff,* with *pilA1* affecting pilus number and location while *pilA2* is required

for twitching (Kandel et al., 2018). It is unclear if this *pilA2* duplication in *Xt* PLS229[T]

can complement its lack of *pilA1*.

In addition to the T4P genes, many other *Xylella* pathogenicity factors have

been identified and these additional virulence genes may be classified into 10 major

functional categories (Lee et al., 2014; Merfa et al., 2016; Chen et al., 2017;

Rapicavoli et al., 2018; Ge et al., 2021). Among these categories, secretion systems

and metabolism are the most conserved ones with no variation in gene copy number

across all *Xylella* representatives (**Figure 5**). Additionally, those genes involved in

regulatory systems are also highly conserved. In contrast, several genes related to

adhesins, hydrolytic enzymes, and toxin-antitoxin systems are highly variable in copy

numbers.

For more detailed examination, these putative virulence genes were classified

into 38 homologous gene clusters and six are absent in the *Xt* genome (**Figure 5**).

These include the genes that encode a putative adhesin (PD0986, hemagglutinin-

like protein), two hydrolytic enzymes (PD0956, serine protease; PD1485,

polygalacturonase), one pair of toxin-antitoxin (PD0370, motility quorum sensing

regulator MqsR ribonuclease; PD0371, MqsA antitoxin), and another separate toxin

(PD1100, endoribonuclease). Notably, the *mqsR-mqsA* toxin-antitoxin system genes

(Lee et al., 2014; Merfa et al., 2016) are differentially distributed among those *Xf*

subspecies. Homologs of these two genes are entirely conserved in all *Xff* and *Xf*

subspecies *morus* strains, present in five out of the nine *Xfp* strains, and completely

absent in *Xfs* and *Xf* subspecies *multiplex*.

Based on previous studies that characterized mutant phenotypes, PD0956

(Gouran et al., 2016) and PD1100 (Burbank and Stenger, 2017) are both

antivirulence factors and the loss of either one resulted in hypervirulence of *Xff* in

grapevines. Similarly, PD0370 is another antivirulence factor that reduces the

virulence of *Xfp* against citrus when overexpressed (Merfa et al., 2016). In contrast,

both PD0986 and PD1485 are critical for *Xff* virulence in grapevines. For PD0986,

this gene is absent in a *Xf* biocontrol strain EB92-1 that can infect and persist in

grapevines but causes only very slight symptoms. When PD0986 is cloned into

EB92-1, the transformant induces significantly increased symptoms that are

characteristic of PD (Zhang et al., 2015). For PD1485, the knockout mutant was

avirulent due to the loss of ability to systemically colonize grapevines (Roper et al.,

2007).

Two gene families appeared to have experienced copy number expansion in

the *Xt* genome. The first family includes homologs of PD1792 and PD2118, which

encode hemagglutinins. These adhesins are antivirulence factors that restrict *in*

planta movement by promoting self-aggregation; transposon-insertion mutants of *Xff*

PD1792 and PD2118 both exhibit hypervirulence in grapevines (Guilhabert and

Kirkpatrick, 2005). Among the representative *Xf* and *Xff* genomes, the median copy

numbers of this family are 3 and 8, respectively. In comparison, *Xt* has 12 copies. It

remains to be investigated if the copy number variation is linked to protein

expression level and virulence. The second family includes a Zot-like toxin

(PD0928). Similar to PD0986 (hemagglutinin-like protein), the biocontrol strain EB92-

1 lacks the homolog of PD0928 and the transformant that expresses this gene is

virulent (Zhang et al., 2015).

<u>Conclusions</u>

In conclusion, this work reported the complete genome sequence of an

important plant-pathogenic bacterium that is endemic to Taiwan. In addition to

providing the genomic resource that contributes to the study of this pathogen, this

species is the only known sister of *Xf*, which has extensive genetic variations and

devastating effects on agriculture worldwide. The availability of this new *Xt* genome

sequence provides critical genomic information of a key lineage that may improve

the study of *Xylella* evolution and the inference of *Xf* ancestral states. At above-

genus level, our genome-scale phylogenetic inference resolved the relationships

between *Xylella* and *Xanthomonas*, which are some of the key plant pathogens in

the family Xanthomonadaceae.

For gene content analysis, our comparison of the putative virulence genes

and pathogenicity factors among representative *Xylella* strains identified the genes

that exhibit high levels of conservation or diversity (**Figures 4 and 5**). These genes

are promising candidates for future functional studies to investigate the molecular

mechanisms of *Xylella* virulence. Previous characterizations of single-gene mutants,

particularly those conducted in *Xff*, have provided a strong foundation (Burdman et

al., 2011; Rapicavoli et al., 2018). However, it is important to note that the current

knowledge of *Xylella* virulence genes is mostly derived from to those strains that are

relatively easy to culture and transform, such that only limited diversity has been

investigated in molecular genetics studies. Moreover, infection experiments for the

investigation of gene functions were limited to a small number of plant species. For

further improvements, experimental studies that examine more diverse *Xylella*

lineages and plant hosts, as well as the combined effects of multiple virulence genes

will be critical.

Data Availability

314   The complete genome sequence of *Xylella taiwanensis* PLS229[T] has been

315   deposited in GenBank/ENA/DDBJ under the accession CP053627. The raw reads

316   have been deposited at the NCBI Sequence Read Archive under the accession

317   numbers SRR11805344 and SRR11805345.

Funding

320   The funding was provided by the Ministry of Science and Technology of

321   Taiwan (MOST 106-2923-B-002-005-MY3) to CWT and Academia Sinica to CHK.

322   The funders had no role in study design, data collection and interpretation, or the

323   decision to submit the work for publication.

Acknowledgments

326   The Illumina and Oxford Nanopore sequencing library preparation service was

327   provided by the Genomic Technology Core (Institute of Plant and Microbial Biology,

328   Academia Sinica). The Illumina MiSeq sequencing service was provided by the

329   Genomics Core (Institute of Molecular Biology, Academia Sinica).

Author Contribution

Conceptualization: CWT, CHK

Funding acquisition: CWT, CHK

Investigation: LWW, YCL, CTH

Methodology: CCS, STC, APC, SJC, CHK

Project administration: CWT, CHK

Resources: CCS, CWT, CHK

Supervision: CWT, CHK

Validation: LWW, YCL, CTH

Visualization: LWW, YCL, CTH

Writing – original draft: LWW, CHK

Writing – review & editing: LWW, YCL, CCS, CTH, STC, APC, SJC, CWT, CHK

Conflict of Interest

The authors declare no conflict of interest.

References

Anderson, I., Teshima, H., Nolan, M., Lapidus, A., Tice, H., Rio, T. G. D., et al.

(2013). Genome sequence of *Frateuria aurantia* type strain (Kondô 67^T), a

xanthomonade isolated from *Lilium auratium* Lindl. *Stand Genomic Sci* 9, 83.

doi:10.4056/sigs.4338002.

Baldi, P., and La Porta, N. (2017). *Xylella fastidiosa*: host range and advance in

molecular identification techniques. *Front Plant Sci* 8, 944.

doi:10.3389/fpls.2017.00944.

Burbank, L. P., and Stenger, D. C. (2017). The DinJ/RelE toxin-antitoxin system

suppresses bacterial proliferation and virulence of *Xylella fastidiosa* in

grapevine. *Phytopathology* 107, 388–394. doi:10.1094/PHYTO-10-16-0374-R.

Burdman, S., Bahar, O., Parker, J. K., and De La Fuente, L. (2011). Involvement of

type IV pili in pathogenicity of plant pathogenic bacteria. Genes 2, 706–735.

doi:10.3390/genes2040706.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., et

al. (2009). BLAST+: architecture and applications. *BMC Bioinformatics* 10,

421. doi:10.1186/1471-2105-10-421.

Chen, H., Kandel, P. P., Cruz, L. F., Cobine, P. A., and De La Fuente, L. (2017). The

major outer membrane protein MopB Is required for twitching movement and

affects biofilm formation and virulence in two *Xylella fastidiosa* strains. Mol

Plant Microbe Interact 30, 896–905. doi:10.1094/MPMI-07-17-0161-R.

Cho, S.-T., Zwolińska, A., Huang, W., Wouters, R. H. M., Mugford, S. T., Hogenhout,

S. A., et al. (2020). Complete genome sequence of "*Candidatus* Phytoplasma

asteris" RP166, a plant pathogen associated with rapeseed phyllody disease

in Poland. *Microbiol Resour Announc* 9. doi:10.1128/MRA.00760-20.

Cursino, L., Galvani, C. D., Athinuwat, D., Zaini, P. A., Li, Y., De La Fuente, L., et al.

(2011). Identification of an operon, Pil-Chp, that controls twitching motility and

virulence in *Xylella fastidiosa*. Mol Plant Microbe Interact 24, 1198–1206.

doi:10.1094/MPMI-10-10-0252.

Denancé, N., Briand, M., Gaborieau, R., Gaillard, S., and Jacques, M.-A. (2019).

Identification of genetic relationships and subspecies signatures in *Xylella*

fastidiosa. *BMC Genomics* 20, 239. doi:10.1186/s12864-019-5565-9.

Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and

high throughput. *Nucl Acids Res* 32, 1792–1797. doi:10.1093/nar/gkh340.

European Food Safety Authority (2018). Update of the *Xylella* spp. host plant

database. *EFSA Journal* 16, e05408. doi:10.2903/j.efsa.2018.5408.

Felsenstein, J. (1989). PHYLIP - phylogeny inference package (version 3.2).

Cladistics 5, 164–166.

Ge, Q., Cobine, P., and De La Fuente, L. (2021). The influence of copper

homeostasis genes *copA* and *copB* on *Xylella fastidiosa* virulence is affected

by sap copper concentration. Phytopathology. doi:10.1094/PHYTO-12-20-

0531-R.

Guindon, S., and Gascuel, O. (2003). A simple, fast, and accurate algorithm to

estimate large phylogenies by maximum likelihood. *Syst Biol* 52, 696–704.

doi:10.1080/10635150390235520.

Gouran, H., Gillespie, H., Nascimento, R., Chakraborty, S., Zaini, P. A., Jacobson,

A., et al. (2016). The secreted protease PrtA controls cell growth, biofilm

formation and pathogenicity in *Xylella fastidiosa*. *Sci Rep* 6, 31098.

doi:10.1038/srep31098.

Guilhabert, M. R., and Kirkpatrick, B. C. (2005). Identification of *Xylella fastidiosa*

antivirulence genes: hemagglutinin adhesins contribute to *X. fastidiosa* biofilm

maturation and colonization and attenuate virulence. *Mol Plant Microbe*

Interact 18, 856–868. doi:10.1094/MPMI-18-0856.

Hao, L., Athinuwat, D., Johnson, K., Cursino, L., Burr, T. J., and Mowery, P. (2017).

Xylella fastidiosa pil-chp operon is involved in regulating key structural genes

of both type I and IV pili. Vitis 56, 55–62. doi:10.5073/vitis.2017.56.55-62.

Jacques, M.-A., Denancé, N., Legendre, B., Morel, E., Briand, M., Mississipi, S., et

al. (2016). New coffee plant-infecting *Xylella fastidiosa* variants derived via

homologous recombination. Appl Environ Microbiol 82, 1556–1568.

doi:10.1128/AEM.03299-15.

Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T., and Aluru, S.

(2018). High throughput ANI analysis of 90K prokaryotic genomes reveals

clear species boundaries. *Nat Commun* 9, 5114. doi:10.1038/s41467-018-

07641-9.

Kandel, P. P., Chen, H., and De La Fuente, L. (2018). A short protocol for gene

knockout and complementation in *Xylella fastidiosa* shows that one of the type

IV pilin paralogs (PD1926) is needed for twitching while another (PD1924)

affects pilus number and location. Appl Environ Microbiol 84, e01167-18.

doi:10.1128/AEM.01167-18.

Lee, M. W., Tan, C. C., Rogers, E. E., and Stenger, D. C. (2014). Toxin-antitoxin

systems *mqsR/ygiT* and *dinJ/relE* of *Xylella fastidiosa*. Physiol Mol Plant

Pathol 87, 59–68. doi:10.1016/j.pmpp.2014.07.001.

Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*

34, 3094–3100. doi:10.1093/bioinformatics/bty191.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows–

Wheeler transform. *Bioinformatics* 25, 1754–1760.

doi:10.1093/bioinformatics/btp324.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., et al. (2009).

The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25,

2078–2079. doi:10.1093/bioinformatics/btp352.

Li, Yaxin., Hao, Guixia., Galvani, C. D., Meng, Y., De La Fuente, L., Hoch, H. C., et

al. (2007). Type I and type IV pili of *Xylella fastidiosa* affect twitching motility,

biofilm formation and cell–cell aggregation. Microbiology 153, 719–726.

doi:10.1099/mic.0.2006/002311-0.

Li, L., Stoeckert, C. J., and Roos, D. S. (2003). OrthoMCL: identification of ortholog

groups for eukaryotic genomes. *Genome Res* 13, 2178–2189.

doi:10.1101/gr.1224503.

Lo, W.-S., Chen, L.-L., Chung, W.-C., Gasparich, G. E., and Kuo, C.-H. (2013).

Comparative genome analysis of *Spiroplasma melliferum* IPMB4A, a

honeybee-associated bacterium. *BMC Genomics* 14, 22. doi:10.1186/1471-

2164-14-22.

Lo, W.-S., Gasparich, G. E., and Kuo, C.-H. (2018). Convergent evolution among

ruminant-pathogenic *Mycoplasma* involved extensive gene content changes.

Genome Biol Evol 10, 2130–2139. doi:10.1093/gbe/evy172.

Meng, Y., Li, Y., Galvani, C. D., Hao, G., Turner, J. N., Burr, T. J., et al. (2005).

Upstream migration of *Xylella fastidiosa* via pilus-driven twitching motility. J

Bacteriol 187, 5560–5567. doi:10.1128/JB.187.16.5560-5567.2005.

Merfa, M. V., Niza, B., Takita, M. A., and De Souza, A. A. (2016). The MqsRA toxin-

antitoxin system from *Xylella fastidiosa* plays a key role in bacterial fitness,

pathogenicity, and persister cell formation. Front Microbiol 7, 904.

doi:10.3389/fmicb.2016.00904.

Paradis, E., and Schliep, K. (2019). ape 5.0: an environment for modern

phylogenetics and evolutionary analyses in R. *Bioinformatics* 35, 526–528.

doi:10.1093/bioinformatics/bty633.

Parkinson, N., Cowie, C., Heeney, J., and Stead, D. (2009). Phylogenetic structure

of *Xanthomonas* determined by comparison of *gyrB* sequences. *Int J Syst*

Evol Microbiol 59, 264–274. doi:10.1099/ijs.0.65825-0.

Pieretti, I., Royer, M., Barbe, V., Carrere, S., Koebnik, R., Cociancich, S., et al.

(2009). The complete genome sequence of *Xanthomonas albilineans*

provides new insights into the reductive genome evolution of the xylem-limited

Xanthomonadaceae. *BMC Genomics* 10, 616. doi:10.1186/1471-2164-10-

616.

Potnis, N., Kandel, P. P., Merfa, M. V., Retchless, A. C., Parker, J. K., Stenger, D.

C., et al. (2019). Patterns of inter- and intrasubspecific homologous

recombination inform eco-evolutionary dynamics of *Xylella fastidiosa*. *ISME J*

13, 2319–2333. doi:10.1038/s41396-019-0423-y.

Rapicavoli, J., Ingel, B., Blanco-Ulate, B., Cantu, D., and Roper, C. (2018). *Xylella*

fastidiosa: an examination of a re-emerging plant pathogen. *Mol Plant Pathol*

19, 786–800. doi:https://doi.org/10.1111/mpp.12585.

Robinson, J. T., Thorvaldsdottir, H., Winckler, W., Guttman, M., Lander, E. S., Getz,

G., et al. (2011). Integrative genomics viewer. *Nat Biotech* 29, 24–26.

doi:10.1038/nbt.1754.

Rodriguez-R, L. M., Grajales, A., Arrieta-Ortiz, M. L., Salazar, C., Restrepo, S., and

Bernal, A. (2012). Genomes-based phylogeny of the genus *Xanthomonas*.

BMC Microbiol 12, 43. doi:10.1186/1471-2180-12-43.

Roper, M. C., Greve, L. C., Warren, J. G., Labavitch, J. M., and Kirkpatrick, B. C.

(2007). *Xylella fastidiosa* requires polygalacturonase for colonization and

pathogenicity in *Vitis vinifera* grapevines. *Mol Plant Microbe Interact* 20, 411–

419. doi:10.1094/MPMI-20-4-0411.

Su, C.-C., Chang, C.-J., Yang, W.-J., Hsu, S.-T., Tzeng, K.-C., Jan, F.-J., et al.

(2012). Specific characters of 16S rRNA gene and 16S–23S rRNA internal

transcribed spacer sequences of *Xylella fastidiosa* pear leaf scorch strains.

Eur J Plant Pathol 132, 203–216. doi:10.1007/s10658-011-9863-6.

Su, C.-C., Deng, W.-L., Jan, F.-J., Chang, C.-J., Huang, H., and Chen, J. (2014).

Draft genome sequence of *Xylella fastidiosa* pear leaf scorch strain in Taiwan.

Genome Announc. 2, e00166-14. doi:10.1128/genomeA.00166-14.

Su, C.-C., Deng, W.-L., Jan, F.-J., Chang, C.-J., Huang, H., Shih, H.-T., et al. (2016).

Xylella taiwanensis sp. nov., causing pear leaf scorch disease. *Int J Syst Evol*

Microbiol 66, 4766–4771. doi:10.1099/ijsem.0.001426.

Tatusova, T., DiCuccio, M., Badretdin, A., Chetvernin, V., Nawrocki, E. P.,

Zaslavsky, L., et al. (2016). NCBI prokaryotic genome annotation pipeline.

Nucleic Acids Res 44, 6614–6624. doi:10.1093/nar/gkw569.

Vanhove, M., Retchless, A. C., Sicard, A., Rieux, A., Coletta-Filho, H. D., De La

Fuente, L., et al. (2019). Genomic diversity and recombination among *Xylella*

fastidiosa subspecies. *Appl Environ Microbiol* 85, e02972-18.

doi:10.1128/AEM.02972-18.

Wells, J. M., Raju, B. C., Hung, H.-Y., Weisburg, W. G., Mandelco-Paul, L., and

Brenner, D. J. (1987). *Xylella fastidiosa* gen. nov., sp. nov: gram-negative,

xylem-limited, fastidious plant bacteria related to *Xanthomonas* spp. *Int J Syst*

Bacteriol 37, 136–143. doi:10.1099/00207713-37-2-136.

Wick, R. R., Judd, L. M., Gorrie, C. L., and Holt, K. E. (2017). Unicycler: Resolving

bacterial genome assemblies from short and long sequencing reads. *PLOS*

Comput Biol 13, e1005595. doi:10.1371/journal.pcbi.1005595.

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. New York, NY:

Springer-Verlag New York doi:10.1007/978-0-387-98141-3.

Zhang, S., Chakrabarty, P. K., Fleites, L. A., Rayside, P. A., Hopkins, D. L., and

Gabriel, D. W. (2015). Three new Pierce's Disease pathogenicity effectors

identified using *Xylella fastidiosa* biocontrol strain EB92-1. *PLOS ONE* 10,

e0133796. doi:10.1371/journal.pone.0133796.

 Figure Legends

Figure 1. Molecular phylogeny of *Xylella* and related species in the family

Xanthomonadaceae. The maximum likelihood phylogeny was based on 779 shared

single-copy genes, the concatenated alignment contains 252,319 aligned amino acid

sites. The genus *Pseudoxanthomonas* was included as the outgroup. (A) A

phylogram for illustrating the relationships among all 40 genomes analyzed. Each of

the *X. fastidiosa* subspecies was collapsed into a triangle for simplified visualization.

All internal nodes illustrated in this phylogram received >95% bootstrap support

based on 1,000 replicates. (B) A cladogram for illustrating the relationships among

those 28 *Xylella* genomes analyzed. Internal nodes with bootstrap values lower than

100% were labeled.

Figure 2. Genome similarity among the representative *Xylella* strains. The pairwise

comparisons were classified into three categories: (1) within the same *X. fastidiosa*

subspecies, (2) between different *X. fastidiosa* subspecies, and (3) between *X.*

fastidiosa and *X. taiwanensis*.

Figure 3. Principal coordinates analysis of gene content dissimilarity. The %

variance explained by each coordinate was provided in parentheses. (A) Based on

the 11,455 homologous gene clusters found among all 40 Xanthomonadaceae

genomes analyzed in this work. (B) Based on the 5,395 homologous gene clusters

found among the 28 *Xylella* genomes analyzed in this work.

Figure 4. Distribution of the type IV pili genes found among representative *Xylella*

genomes. *Xanthomonas oryzae* was included as the outgroup. The genes were

identified by the PD numbers based on the annotation of Temecula1 genome; those

with adjacent PD numbers are located in neighboring regions on the chromosome. A

total of four gene clusters (labelled as I-IV) located in syntenic regions were found to

be conserved. Gene names were provided when available; '*hypo*' indicates those

annotated as encoding hypothetical proteins. Patterns of gene presence and

absence were illustrated in the format of a heatmap. For gene absence, those with

identifiable pseudogenes were labelled accordingly. One case of tandem gene

duplication was observed for the PD_1926 (*pilA2*) homolog in the *Xylella taiwanensis*

genome.

Figure 5. Distribution of putative virulence genes and pathogenicity factors among

representative *Xylella* genomes. *Xanthomonas oryzae* was included as the outgroup.

The homologous gene clusters were identified by the PD numbers based on the

annotation of Temecula1 genome; gene names were provided when available. Gene

copy numbers were illustrated in the format of a heatmap; values higher than two

were labeled with the exact numbers. Two adhesin genes (i.e., PD1792 and

PD2118) were assigned to the same homologous gene cluster and were combined

for copy number calculation.

Table

Table 1. List of the genome sequences analyzed.

Species	Strain	Location	Accession	Assembly	Genome Size (bp)	CDS (Intact)	CDS (Pseudo)	CDS (All)
Xylella taiwanensis	PLS229	Taiwan: Houli, Taichung	GCA_013177435.1	Complete	2,824,877	2,176	132	2,308
Xylella fastidiosa subsp. *fastidiosa*	ATCC35879	USA: Florida	GCA_011801475.1	Complete	2,607,257	2,189	133	2,322
Xylella fastidiosa subsp. *fastidiosa*	Bakersfield-1	USA: Bakersfield, California	GCA_009664125.2	Complete	2,575,627	2,198	70	2,268
Xylella fastidiosa subsp. *fastidiosa*	GB514	USA: Texas	GCA_000148405.1	Complete	2,517,383	1,998	197	2,195
Xylella fastidiosa subsp. *fastidiosa*	GV230	Taiwan: Waipu, Taichung	GCA_014249995.1	Complete	2,514,993	2,092	61	2,153
Xylella fastidiosa subsp. *fastidiosa*	M23	USA: California	GCA_000019765.1	Complete	2,573,987	2,235	118	2,353
Xylella fastidiosa subsp. *fastidiosa*	Temecula1	USA: California	GCA_000007245.1	Complete	2,521,148	2,107	64	2,171
Xylella fastidiosa subsp. *morus*	MUL0034	USA: California	GCA_000698825.1	Complete	2,666,577	2,266	151	2,417
Xylella fastidiosa subsp. *morus*	Mul-MD	USA: Maryland	GCA_000567985.1	101 contigs	2,520,555	2,121	155	2,276
Xylella fastidiosa subsp. *multiplex*	AlmaEM3	USA: Georgia	GCA_006369915.1	30 contigs	2,479,954	2,049	141	2,190
Xylella fastidiosa subsp. *multiplex*	CFBP8418	France: Corse, Alata	GCA_001971465.1	271 contigs	2,513,969	2,199	298	2,497
Xylella fastidiosa subsp. *multiplex*	Dixon	USA: California	GCA_000166835.1	32 contigs	2,622,328	2,272	123	2,395
Xylella fastidiosa subsp. *multiplex*	Griffin-1	USA: Georgia	GCA_000466025.1	84 contigs	2,387,314	1,911	303	2,214
Xylella fastidiosa subsp. *multiplex*	M12	USA: California	GCA_000019325.1	Complete	2,475,130	2,041	116	2,157
Xylella fastidiosa subsp. *multiplex*	sycamore Sy-VA	USA: Virginia	GCA_000732705.1	128 contigs	2,475,880	2,068	178	2,246
Xylella fastidiosa subsp. *multiplex*	TOS14	Italy: Tuscany	GCA_007713995.1	80 contigs	2,445,518	2,013	159	2,172
Xylella fastidiosa subsp. *multiplex*	TOS4	Italy: Tuscany	GCA_007713905.1	77 contigs	2,445,114	2,017	158	2,175
Xylella fastidiosa subsp. *multiplex*	TOS5	Italy: Tuscany	GCA_007713945.1	72 contigs	2,443,867	2,017	152	2,169
Xylella fastidiosa subsp. *pauca*	3124	Brazil: Matao, Sao Paulo	GCA_001456195.1	Complete	2,748,594	2,273	179	2,452
Xylella fastidiosa subsp. *pauca*	9a5c	Brazil: Macaubal, Sao Paulo	GCA_000006725.1	Complete	2,731,750	2,333	153	2,486
Xylella fastidiosa subsp. *pauca*	De Donno	Italy: Apulia	GCA_002117875.1	Complete	2,543,738	2,092	152	2,244
Xylella fastidiosa subsp. *pauca*	Fb7	Argentina: Corrientes	GCA_001456335.3	Complete	2,699,320	2,178	286	2,464
Xylella fastidiosa subsp. *pauca*	Hib4	Brazil: Jarinu, Sao Paulo	GCA_001456315.1	Complete	2,877,548	2,456	185	2,641
Xylella fastidiosa subsp. *pauca*	J1a12	Brazil: Jales, Sao Paulo	GCA_001456235.1	Complete	2,867,237	2,421	242	2,663
Xylella fastidiosa subsp. *pauca*	Salento-1	Italy: Taviano, Lecce, Apulia	GCA_002954185.1	Complete	2,543,366	1,989	260	2,249
Xylella fastidiosa subsp. *pauca*	Salento-2	Italy: Ugento, Lecce, Apulia	GCA_002954205.1	Complete	2,543,566	2,033	212	2,245
Xylella fastidiosa subsp. *pauca*	U24D	Brazil: Ubarana, Sao Paulo	GCA_001456275.1	Complete	2,732,490	2,274	178	2,452
Xylella fastidiosa subsp. *sandyi*	Ann-1	USA: California	GCA_000698805.1	Complete	2,780,908	2,379	179	2,558
Xanthomonas albilineans	Xa-FJ1	China: Fujian	GCA_009931595.1	Complete	3,756,117	2,968	114	3,082
Xanthomonas arboricola	17	China: Jiangsu	GCA_000972745.1	Complete	5,254,865	4,330	162	4,492
Xanthomonas axonopodis	NCPPB 796	Mauritius	GCA_013177355.1	Complete	4,886,779	3,514	683	4,197
Xanthomonas campestris	MAFF106181	Japan: Aomori	GCA_013388375.1	Complete	4,942,039	4,041	75	4,116
Xanthomonas citri	GD3	China: Guangdong	GCA_000961335.1	Complete	5,223,748	4,219	141	4,360
Xanthomonas cucurbitae	ATCC 23378	USA: New York	GCA_009883735.1	Complete	4,615,492	3,702	234	3,936
Xanthomonas hortorum	B07-007	Canada: Monteregie, Quebec	GCA_002285515.1	Complete	5,250,904	4,241	218	4,459
Xanthomonas hyacinthi	CFBP 1156	Netherlands	GCA_009769165.1	Complete	4,963,026	4,025	282	4,307
Xanthomonas oryzae	PXO99A	Philippines	GCA_000019585.2	Complete	5,238,555	3,952	736	4,688
Xanthomonas vesicatoria	LMG911	New Zealand	GCA_001908725.1	Complete	5,349,905	4,340	159	4,499
Pseudoxanthomonas mexicana	GTZY	China: Beijing	GCA_014211895.1	Complete	3,936,186	3,575	37	3,612
Pseudoxanthomonas spadix	BD-a59	South Korea	GCA_000233915.4	Complete	3,452,554	3,048	85	3,133

A

Coordinate 2 (6.16%)

Coordinate 1 (59.1%)

X. citri
X. hortorum
X. vesicatoria
X. campestris
X. arboricola
X. cucurbitae
X. axonopodis
X. oryzae
X. albilineans
X. hyacinthi
P. spadix
P. mexicana

B

Coordinate 2 (18.2%)

Coordinate 1 (25.9%)

🔴 *Xylella taiwanensis*

🔵 *X. fastidiosa* subsp. *pauca*

🟠 *X. fastidiosa* subsp. *sandyi*

🟢 *X. fastidiosa* subsp. *fastidiosa*

🟡 *X. fastidiosa* subsp. *morus*

🟣 *X. fastidiosa* subsp. *multiplex*

⚪ *Xanthomonas*

⚫ *Pseudoxanthomonas*