1 **RBPMS2 is a conserved regulator of alternative splicing that promotes myofibrillar**
2 **organization and optimal calcium handling in cardiomyocytes**
3
4
5
6

7 Alexander A. Akerberg (PhD)[1,2,3], Michael Trembley (PhD)[1,3], Vincent Butty (MD, PhD)[4], Asya
8 Schwertner (PhD)[2,3], Long Zhao (PhD)[2,3], Manu Beerens (PhD)[2,3,5], Xujie Liu (PhD)[1,3],
9 Mohammed Mahamdeh (PhD)[2,3], Shiaulou Yuan (PhD)[2,3], Laurie Boyer (PhD)[4,6], Calum
10 MacRae (MD, PhD)[3,5], Christopher Nguyen (PhD)[2,3], William T. Pu (MD)[1,3,7], Caroline E. Burns
11 (PhD)[1,2,3,7*], C. Geoffrey Burns (PhD)[1,2,3*]
12
13
14
15
[1] Division of Basic and Translational Cardiovascular Research, Department of Cardiology, Boston
Children's Hospital, Boston, MA, USA
[2] Cardiovascular Research Center, Massachusetts General Hospital, Charlestown, MA, USA
[3] Harvard Medical School, Boston, MA, USA
[4] Department of Biology, Massachusetts Institute of Technology, Cambridge, MA, USA
[5] Division of Cardiovascular Medicine, Brigham and Women's Hospital, Boston, MA, USA
[6] Department of Biological Engineering, Massachusetts Institute of Technology, Cambridge, MA,
USA
[7] Harvard Stem Cell Institute, Cambridge, MA, USA
25
26
27
28
29 * Correspondence: Caroline and C. Geoffrey Burns: caroline.burns@childrens.harvard.edu,
30 geoff.burns@childrens.harvard.edu. 300 Longwood Ave., Boston MA 02115

31
32
33 Current Word Count: 10,455
34 AHA Subject Terms: Contractile function, myocardial biology, Genetics, Gene expression and
35 regulation, Heart failure

ABSTRACT

<u>Rationale:</u> The identification of novel cardiomyocyte-intrinsic factors that support heart function will expand the number of candidate genes and therapeutic options for heart failure, a leading cause of death worldwide.

<u>Objective:</u> To identify and characterize conserved regulators of cardiomyocyte function.

<u>Methods and Results:</u> We report that the RNA-binding protein RBPMS2 is required for myofibril organization and the regulation of intracellular calcium dynamics in both zebrafish embryos and human induced pluripotent stem cell-derived cardiomyocytes (hiPSC-CMs). A differential expression screen in zebrafish uncovered enrichment of *rbpms2* paralogs, *rbpms2a* and *rbpms2b,* in the myocardium. Double knock-out (*rbpms2*-null) embryos suffer from compromised ventricular filling during the relaxation phase of the cardiac cycle, which significantly reduces cardiac output. Whole transcriptome sequencing and validation studies revealed differential alternative splicing of several genes linked to cardiomyopathies in humans, including *myosin binding protein C3* (*mybpc3*) and *phospholamban* (*pln*), consistent with a role in causing the observed ventricular deficiencies. Further, *RBPMS*2-null hiPSC-CMs exhibit myofibril and calcium handling defects that are highly analogous to those observed in the *rbpms2*-null zebrafish ventricle.

<u>Conclusions:</u> Taken together, our data identify *RBPMS2* as a conserved and essential regulator of alternative splicing that is required for myofibrillar organization and optimal calcium handling from zebrafish to humans.

INTRODUCTION

Heart failure remains a common, costly, and growing public health burden [1]. While lifestyle choices that cause hypertension and myocardial infarction are substantial contributors to heart failure, inherited and *de novo* genetic mutations that cause primary diseases of the heart muscle, including hypertrophic cardiomyopathy (HCM) and dilated cardiomyopathy (DCM) are also important instigators of disease [2,3]. Causal mutations for HCM and DCM commonly reside within two classes of genes, those encoding components of the sarcomere, the basic contractile unit of cardiac muscle, and those encoding factors that regulate calcium handling, another critical determinant of cardiac contractility [2,4–6]. While the pathogenesis of mutations affecting sarcomeres and calcium handling is well-established in HCM and DCM, a significant percentage of cases remain idiopathic, because clinically recognized cardiomyopathy susceptibility genes remain mutation free in targeted sequencing panels. Therefore, ongoing efforts to identify novel molecular activities required for optimal myocardial contractility will highlight novel candidate genes for primary cardiomyopathies.

Alternative mRNA splicing has emerged in recent years as an important regulator of myocardial contractility in humans. Disruptions in the RNA-recognition motif (RRM)-containing protein RBM20 were recently discovered to cause dilated cardiomyopathy (DCM) by regulating alternative splicing of TITIN (TTN), the largest protein component of the sarcomere [7–9]. TTN transcripts within the heart can be separated into three distinct classes, each of which codes for a unique combination of functional domains [10]. Loss of Rbm20 in animal models results in aberrant splicing of TTN, leading to changes in ventricular morphology and diminished contractility in a manner consistent with observations in DCM patients. Clinical studies have further confirmed that genetic variation in the Rbm20 locus is associated with a higher risk of cardiomyopathy [11,12]. Alternatively, spliced transcripts of calcium/calmodulin-dependent kinase II delta (CAMK2D) were also shown to contribute to increased risk of arrhythmogenic cardiomyopathy in the absence of Rbm20 [7], demonstrating roles for Rbm20 in the regulation of

82 myocardial calcium handling and action potential propagation. Similar to Rbm20, another RRM-

83 containing protein, termed Rbm24, was shown to regulate the splicing of a subset of genes

84 encoding components of the sarcomere, including TTN, within postnatal mice[13]. Initial screens

85 for Rbm24 mutations in human cardiomyopathy patients suggest a potential link to disease,

86 however, these mutations were found to be relatively rare in the disease population [14]. Together

87 these data represent the first etiological links between splicing factors and cardiomyopathy,

88 thereby underscoring the importance for further research to identify the full array of splicing

89 factors that regulate cardiac form and function.

90   The importance of alternative splicing in the heart is further evidenced by the abundance

91 of isoform variation within cardiac-specific transcripts. Furthermore, many genes with

92 established roles in cardiomyopathy are known to produce multiple distinct isoforms within the

93 heart. For example, the cardiac troponin t (*TNNT2*) gene is alternatively spliced in a regulated

94 pattern during cardiac development and aberrant splicing can lead to cardiomyopathy in animal

95 models [15–17]. Other cardiomyopathy genes such as tropomyosin (*TPM1*) and myosin binding

96 protein C3 (*MYBPC3*) are predicted to be alternatively spliced in a similar fashion, however the

97 importance of these splicing events and the identity of the factors that govern them remain

98 unknown.

99   Here, we identify the RRM-containing protein Rbpms2 (RNA binding protein with multiple

100 splicing 2) as a crucial regulator of alternative splicing within the developing myocardium. We

101 show that Rbpms2a and Rbpms2b redundantly promote ventricular contractility within the

102 developing zebrafish heart. Further, we show that these Rbpms2 proteins regulate the splicing

103 of numerous genes that are associated with cardiac sarcomere assembly and ion transport,

104 including transcripts for known disease genes *mybpc3* and *phospholamban* (*pln)* [18,19]. We then

105 demonstrate that *rbpms2*-null animals exhibit sarcomere disorganization and calcium handling

106 deficits that are consistent with the misregulation of Rbpms2 targets.

METHODS

Data availability

All data and materials will be made publicly available upon publishing. RNA-sequencing

datasets summarized in Tables S1-5 were deposited into the NCBI GEO repository under

accession numbers GSE167416 and GSE167414.

Zebrafish husbandry and strains

Zebrafish were bred and maintained following animal protocols approved by the Institutional

Animal Care and Use Committees at Massachusetts General Hospital and Boston Children's

Hospital. All protocols and procedures followed the guidelines and recommendations outlined by

the Guide for the Care and Use of Laboratory Animals. The following zebrafish strains were

used in this study: *TgBAC(-36nkx2.5:ZsYellow)*[fb7] [20], *Tg(myl7:GFP)*[fb1] [formerly

Tg(cmlc2:GFP)[fb1]] [21], *Tg(myl7:NLS-eGFP)*[fb18] [formerly *Tg(cmlc2:nucGFP)*[fb18]] [22],

Tg(myl7:actn3b-EGFP)[sd10] [formerly *Tg(cmlc2:actinin3-EGFP)*] [23], *rbpms2a*[chb3] (this study), and

rbpms2b[chb4] (this study).

Generation and genotyping of *rbpms2a*[ch3] and rbpms2*b*[ch4] mutant alleles

Customized transcription activator-like effector nucleases (TALENs) [24–26] were used to induce

double-stranded breaks in regions of the *rbpms2a* and *rbpms2b* loci that encode the RNA-

recognition motifs. TALENs were designed to bind the sequences 5'-

TGCCTGTTGACATCAAGC-3' (+) and 5'-TTGAAGGGTCTGAACAGC-3' (-) in exon 2 of

rbpms2a and 5'- TGCCAACAGATATCAAAC-3' (+) and 5'-TTAAATGGTCGAAATAGC-3' (-) in

exon 2 of *rbpms2b*. TALEN constructs were constructed as described [26]. Messenger RNAs

encoding each TALEN pair were produced in vitro and co-injected into one-cell stage zebrafish

embryos as described [25] Germline transmission of TALEN-induced mutations was detected

using fluorescent PCR and DNA-fragment analysis as described [27]. The nucleotides deleted in

the mutant alleles *rbpms2a[chb3]* and *rbpms2b[chb4]* are shown (Fig. S4A,B). The primers utilized to

distinguish wild-type *rbpms2a* and *rbpms2b* from the mutant alleles *rbpms2a[chb3]* and

rbpms2b[chb4] by fluorescent PCR and DNA-fragment analysis are shown (Table S5). The wild-

type alleles produce amplicons of 97 base pairs (bp)(*rbpms2a*) and 76bp (*rbpms2b*), whereas

the mutant alleles produce amplicons of 83bp (*rbpms2a[chb3]*) and 74bp (*rbpms2b[chb4]*).

hiPSC maintenance and genome editing

All hiPSC lines were maintained at 37°C, 5% CO_2 in Essential 8 medium (Cat. # A1517001,

Thermo Fisher Scientific) and passaged every 3 – 4 days using Versene (Cat. # 15040066,

Thermo Fisher Scientific). Culture dishes were pre-coated with 0.5% GelTrex matrix solution

(Cat. # A1413302, Thermo Fisher Scientific). CRISPR/Cas9 genome editing was performed to

create isogenic lines. Briefly, guide RNA for *RBPMS2* (sequence 5'-

GTCTTGCAGTGAGCTTGATC-3') was synthesized using the T7 MegaShortScript transcription

kit (Thermo Fisher Scientific). We previously described methods [28] to generate a doxycycline -

inducible Cas9 line in the WTC-11 iPSC background (Coriell Institute). Targeted integration of

Cas9 into the AAVS1 locus was performed using Addgene plasmids #73500 and #48139. This

WTC-11 AAVS1[Cas9/+] hiPSC line was treated with doxycycline (2μg/ml) for 16 hours prior to

transfection to induce expression of an NLS-SpCas9 fusion protein. After induction,

approximately 0.5×10^6 cells were transfected with 10μg guide RNA using an Amaxa

nucleofector (Lonza Technologies). Cells were seeded at a low density to pick single clones.

Positive clones were confirmed by Sanger and amplicon sequencing. Stem cell marker

expression was validated in isolated cell lines by flow cytometry using anti-SSEA4-FITC (Cat. #

130-098-371, Miltenyi Biotec).

RESULTS

Differential expression analysis of FACS-purified zebrafish anterior lateral plate mesodermal

cells identifies cardiomyocyte-restricted expression of rbpms2a and rbpms2b

Lineage-restricted genes often perform specialized and essential functions in the

development, maintenance, or function of the populations they mark. To generate a candidate

list of zebrafish genes with essential activities in cardiopharyngeal progenitors or embryonic

cardiomyocytes, we sought to identify mRNAs with restricted expression to the subset of

anterior lateral plate mesodermal (ALPM) cells marked by the *nkx2.5:ZsYellow* transgene. This

population contains cardiopharyngeal progenitors for multiple cell types including second heart

field-derived ventricular cardiomyocytes, three outflow tract (OFT) lineages, a subset of

pharyngeal muscles, and endothelium of the pharyngeal arch arteries [20,29–32]. It also contains

early-differentiating ventricular cardiomyocytes derived from the first heart field [29,31,32].

To isolate ALPM cells labeled with the transgene, *Tg(nkx2.5:ZsYellow)* embryos were

dissociated to single cells at a developmental stage when ZsYellow fluorescence is restricted to

the ALPM [14-16 somites stage (ss); Fig. 1A]. Dissociated cells were separated by FACS into

ZsYellow-positive and ZsYellow-negative subpopulations before being subjected to RNA-

sequencing and differential expression analysis (Fig 1A; Fig S1A-C). From this, we identified

1848 protein-coding RNAs significantly enriched in the ZsYellow-positive population [fold

change (FC)>1.5; adjusted p-value<0.05; Fig. 1A; Fig. S1D; Table S1]. Among them were

several previously characterized lateral plate mesodermal markers that encode transcription

factors essential for vertebrate cardiopharyngeal development including *nkx2.5*, *hand2*,

gata4/5/6, *mef2ca/b*, *tbx1/5a/20*, *isl1*, and *fgf8a/10a/10b* (Fig. S1D; Table S1). Due to the

presence of early-differentiating cardiomyocytes in the ZsYellow-positive population, we also

identified numerous genes encoding sarcomere components [*cmlc1*, *myl7* (*cmlc2*), *myh7*

(*vmhc*), *myh6* (*amhc*), *acta1b*, *actc1a*, *actn2b*, *tnni1b*, *tnnt2a, tnnc1a, tpm4a, ttn.2*], calcium

handling proteins [*slc8a1a* (*ncx1*), *ryr2b, atp2a2a (serca2)*] and other factors expressed

preferentially in cardiomyocytes (*cx36.7*, *nppa/al/b*; Fig. S1D; Table S1). Not surprisingly, gene

ontology (GO) term enrichment analysis uncovered overrepresentation of biological process

terms "heart development", "angiogenesis", "heart contraction", "actin cytoskeletal organization",

and "sarcomere organization" in the ALPM-enriched gene set (Fig. S1E, see Table S1 for

complete GO term analysis). The successful identification of ALPM markers with well-

documented and indispensable roles in cardiopharyngeal progenitors or cardiomyocytes

suggests that novel molecular activities of equal importance are likely to be among the

uncharacterized ALPM-restricted genes.

Transcripts encoding <u>R</u>NA <u>b</u>inding <u>p</u>rotein with <u>m</u>ultiple <u>s</u>plicing (variants) 2a (Rbpms2a)

and Rbpms2b were enriched 14- and 5-fold, respectively, in the ZsYellow-positive population

(Fig. S1D; Table S1), suggesting that these RNA-recognition motif (RRM)-containing proteins

localize to the ALPM. Classified as ohnologs, *rbpms2a* and *rbpms2b* encode proteins that are

92% identical and perform redundant functions in oocyte differentiation [33]. Across the full-length

amino acid sequences, Rbpms2a and Rbpms2b share greater than 81% identity with human

RBPMS2 (Fig. S2A,B). The shared identity exceeds 92% in the RRMs (Fig. S2A,B). Such a high

degree of sequence conservation suggests that the molecular function of RBPMS2 proteins and

the properties of their binding site sequences are also likely to be conserved across evolution.

The founding member of the *RBPMS2* gene family, originally termed *hermes*, was

discovered in a differential expression screen designed to identify heart-specific transcripts in

Xenopus [34,35]. In situ hybridization revealed that *RBPMS2* localizes to the lateral plate

mesoderm and embryonic myocardium in both amphibian and avian species [34,36]. Hearts of

embryonic zebrafish were also recently reported to express *rbpms2a* and *rbpms2b* [33]. Despite

the high degree of amino-acid sequence conservation and consistent myocardial-restricted

expression pattern across vertebrates, *RBPMS2* has yet to be implicated in any aspect of

myocardial biology through genetic loss-of-function studies.

To validate the predicted ALPM localization of *rbpms2a* and *rbpms2b,* we performed in

situ hybridization for both genes at the same developmental stage used for profiling

nkx2.5:ZsYellow-positive ALPM cells (16 ss; Fig. S1). As anticipated, both transcripts localized

bilaterally to the ALPM (Fig. 1B,E) in a pattern mirroring early-differentiating cardiomyocytes in

at this stage [37]. Later-stage analysis revealed *rbpms2a* and *rbpms2b* expression in the heart

tube at 24-28 hours post fertilization (hpf; Fig. 1C,F; Fig. S3A,C) and in the atrium and ventricle

of the two-chambered heart at 48 hpf (Fig. 1D,G; Fig. S3B,D). Consistent with previous reports

[33,38], extra-cardiac expression was observed in the pronephric duct and retina (Fig. S3A-D)

To investigate the lineage specificity and subcellular localization of Rbpms2a and

Rbpms2b in the zebrafish heart, we immunostained wild-type embryos with a monoclonal

antibody raised against human RBPMS2, which cross reacts with zebrafish Rbpms2 (see

below). Consistent with the distribution of *rbpms2a/b* transcripts, Rbpms2 protein was detected

in the heart tube (Fig. 1H,I; Fig. S3E,F), two chambered heart (Fig. 1J,K; Fig. S3G), pronephric

duct (Fig. S3E-G) and retina (Fig. S3F,G). Double-immunostaining for Rbpms2 and the

myocardial marker cardiac Troponin T (CT3) revealed that Rbpms2 expression localizes to the

myocardium at all stages analyzed (Fig. 1H',I',J',K'). Weak expression of Rbpms2 was also

evident by 72 hpf in the OFT (i.e. bulbous arteriosus; Fig. 1K,K'). Examination of single optical

sections through the ventricular wall, which showed nuclear Rbpms2 signals surrounded by

Troponin T-positive cytoplasm, confirmed myocardial localization of Rbpms2, (Fig. 1L-L''').

Adjacent endocardial cells were devoid of Rbpms2 signals, which highlights the myocardial-

specific nature of Rbpms2 expression (Fig. 1L-L'''). The subcellular distribution of Rbpms2

appeared variable, ranging from a nucleocytoplasmic distribution to exclusively nuclear (Fig.

1H,I,J,K), but the significance of this observation is unknown. Lastly, we documented Rbpms2

expression in the nuclei and cytoplasm of cardiomyocytes in hearts from adult zebrafish (Fig.

1M,M'), suggesting that Rbpms2 functions in the myocardium continuously throughout the

lifespan of zebrafish rather than transiently during embryogenesis.

Generation of Rbpms2-null zebrafish

To discover if *rbpms2a* and *rbpms2b* are required for myocardial development or

function, we created mutant alleles of both genes using transcription activator-like effector

nucleases (TALENs) [24]. TALEN pairs were designed to target DNA sequences encoding N-

terminal regions of the RRM domains (Fig. S4A,B). We isolated the mutant alleles *rbpms2a^{chb3}*

and *rbpms2b^{chb4}*, which carry frame-shifting deletions and pre-mature stop codons that truncate

the proteins by >80% (Fig. 2A,B; Fig. S4A,B). The truncations also remove 82% of the RRMs,

suggesting that the RNA-binding activities of the mutant proteins are abolished or significantly

compromised. As a result, both alleles are likely to be null. Animals homozygous for either

rbpms2a^{chb3} or *rbpms2b^{chb4}* are grossly indistinguishable from siblings during all life stages,

which is consistent with a previous study that independently generated mutant alleles of

rbpms2a and *rbpms2b* [33].

To determine if *rbpms2a* and *rbpms2b* perform redundant functions in the heart, we

generated and monitored double homozygous embryos [i.e. *rbpms2a^{chb3/chb3}; rbpms2b^{chb4/chb4}*

animals, referred to hereafter as *rbpms2*-null, double-mutant, or double-knockout (DKO)

embryos or animals] for cardiac defects. DKO animals are grossly indistinguishable from control

siblings for the first ~30 hours of life. By ~33 hpf, they develop a characteristic indentation of the

yolk, which is indicative of mild pericardial edema (Fig. S4C,D). At this stage, double-mutants

are delayed in initiating blood flow even though peristalsis of the heart tube is readily evident. By

72 hpf, pericardial edema becomes progressively more severe (Fig. 2C,D), suggesting that

rbpms2-null animals suffer from a cardiac malformation or ongoing functional deficit.

Western blotting and whole mount immunostaining revealed that the zebrafish epitopes

recognized by the anti-human RBPMS2 antibody are completely absent in double mutants (Fig.

2E-G), demonstrating that the antibody cross reacts with zebrafish Rbpms2. It also suggests

that the mutant mRNAs or proteins are unstable, further suggesting that *rbpms2a^{chb3}* and

rbpms2b^{chb4} are bona fide null alleles.

To assess double-mutant viability, we monitored the survival of control-sibling and DKO

animals from embryonic stages to adulthood. Over 50% of *rbpms2*-null animals died by two

weeks of age (Fig. 2H). Eighty five percent of animals perished by 60 days post fertilization

(dpf), and the remaining small percentage of animals did not survive beyond 6 months post

fertilization. At all ages, death was generally accompanied by progressively severe pericardial

edema, suggesting a cardiac origin. Together, these data reveal that *rbpms2a* and *rbpms2b*

ohnologues perform redundant functions that are essential to zebrafish survival.

rbpms2a and rbpms2b are required for chamber expansion during ventricular filling

To evaluate DKO embryos for structural heart disease, we examined hearts in 48 hpf

control-sibling and *rbpms2*-null animals carrying the *myl7:GFP* transgene, which labels the

entire myocardium with GFP. Although the two-chambered heart in double mutants displayed

normal left-right patterning, the ventricle was displaced anteriorly relative to the atrium (Fig.

2I,J), likely reflecting stretching at the cardiac poles caused by pericardial edema. Atrial size

was not obviously altered in *rbpms2*-null animals, but mutant ventricles appeared smaller (Fig.

1I,J). To determine if this might reflect a diminished ventricular cardiomyocyte number, we

quantified cardiomyocytes in control-sibling and *rbpms2*-null animals carrying the *myl7:nucGFP*

transgene, which labels cardiomyocyte nuclei with GFP (Fig. 2K,L). From this, we learned that

ventricular, as well as atrial, cardiomyocyte numbers were unaltered in double mutants (Fig. 2K-

M; Fig. S5A), demonstrating that any deviations in chamber size are not attributable to altered

myocardial cell content. It also suggests that *rbpms2a* and *rbpms2b* are dispensable for

myocardial differentiation from the first and second heart fields, which is the principal method by

which cardiomyocytes are produced before 48 hpf [20,39–41]. These data demonstrate that early

cardiac morphogenesis is largely unperturbed in double mutant animals, with the only apparent

anatomical abnormality being a reduction in ventricular size.

To characterize potential functional defects in *rbpms2*-null hearts, we measured ejection

fraction and cardiac output in 48 hpf animals using our previously described Cardiac Functional

Imaging Network [CFIN; [42]]. To that end, we captured Z-stack dynamic images of GFP+ hearts

over several cardiac cycles in control-sibling and DKO *myl7:GFP* transgenic embryos using light

sheet fluorescence microscopy. CFIN extracted chamber volumes and identified the maximum

and minimum ventricular volumes throughout the cardiac cycle, referred to as end-diastolic

volume (EDV; maximum volume during cardiac relaxation) and end-systolic volume (ESV;

minimum volume during cardiac contraction), respectively. Whereas the ESV was unchanged in

DKO animals, the EDV was significantly reduced (Fig. 2N-R), demonstrating that the ventricle

fails to expand during the relaxation phase of the cardiac cycle. As a result, three indices of

ventricular function, all of which are calculated from EDV and ESV, were significantly reduced.

These include stroke volume (SV), the volume of blood expelled by the ventricle per beat

(SV=EDV-ESV), ejection fraction, the percentage of blood expelled by the ventricle per beat

(EF=SV/EDV), and cardiac output, the total volume of blood expelled by the ventricle per minute

(SV*heart rate) (Fig. 2R). Heart rate was unchanged in the mutant (Fig. 2R). CFIN analysis also

revealed an increase in atrial volume (Fig. S5B), which was not grossly obvious after fixation

(Fig. 2I,J) and is a common secondary consequence of impaired ventricular function [43–45]. These

data demonstrate that *rbpms2a* and *rbpms2b* are required for ventricular function in the

zebrafish embryo. Further, this requirement likely extends into adulthood because the longest-

living double-mutant animals exhibited several signs of heart failure such as stunted growth,

pericardial edema, and cardiac pathologies including compact, fibrotic ventricles and massively

enlarged atria (Fig. S6A-E).

Identification of differential alternative splicing events in rbpms2-null zebrafish

To identify transcriptomic abnormalities in the hearts of *rbpms2*-null embryos, we

performed bulk RNA-sequencing and differential expression analysis on control-sibling and

rbpms2-null animals at 33 hpf, the earliest developmental stage when DKO animals are

distinguishable by morphology (Fig. S4C,D). We reasoned that early-stage analysis would

maximize the chances of highlighting primary molecular defects without interference by

secondary responses to cardiac dysfunction. Importantly, we confirmed that the primitive

ventricle in 33 hpf *rbpms2*-null animals exhibits a functional deficit by documenting reduced

myocardial wall movement at the arterial pole of the heart tube (Fig. S7A). We also confirmed

that total cardiomyocyte numbers are unaltered at 33 hpf in double mutants (Fig. S7B), reducing

the chances that any transcriptomic abnormalities would be attributable to altered cellular

composition. Lastly, we profiled whole embryos, instead of purified hearts, because effective

protocols for isolating hearts or cardiomyocytes at this early stage do not currently exist.

Nonetheless, because *rbpms2a* and *rbpms2b* are expressed with exquisite specificity in the

heart, retina and pronephric duct (Fig. S3F), we anticipated that the transcriptional changes

occurring in the myocardium would be a discernable subset of those occurring in the whole

animal.

While RNA-sequencing and gene-level differential expression analysis highlighted only 2

downregulated transcripts in *rbpms2*-null animals (|FC|>1.5; p-adj<0.05; Table S2; Figure S8A),

isoform-level differential expression analysis yielded 78 downregulated and 74 upregulated

transcripts (|FC|>1.5; p-adj<0.05; Table S3; Fig. S8B). GO term analysis did not identify

overrepresentation of any terms in either gene set [False Discovery Fate (FDR)<0.05].

However, among the downregulated transcripts, there were three well-characterized myocardial

genes including *phospholamban* (*pln*; FC=-43.00; p-adj=1.53E-09), *atrial myosin heavy chain 6*

(*myh6*; FC=-2.17; p-adj=1.52E-11), and *natriuretic peptide precursor a* (*nppa*; FC=-1.57; p-

adj=0.019; Fig. S8B; Table S2). We confirmed downregulation of *nppa* by reverse-transcription

quantitative PCR (RT-qPCR) at 33hpf (Fig. S7C). However, this reduction was short lived

because two subsequent developmental stages showed increased *nppa* expression (Fig. S7C),

which is characteristic behavior for *nppa* in response to cardiac stress [46]. RT-qPCR analysis of

myh6 revealed a downward trending but statistically insignificant change in double mutants (Fig.

S7D). Ultimately, downregulation of either gene is unlikely to explain the ventricular phenotype

in *rbpms2*-null mutants because *nppa*-null zebrafish do not exhibit any gross cardiac defects [47]

and the primary defect in *myh6*-deficient animals is a silent atrium [48].

Of the two *pln* genes in the zebrafish genome (GRCZ11), *si:ch211-270g19.5*

(ENSDARG00000097256) and *pln* (ENSDARG00000069404), the latter is alternatively spliced

according to publicly available cDNA sequencing data. Interestingly, only one of the

alternatively-spliced *pln* isoforms (ENSDART00000141159; *pln-203*) was found to be

downregulated in *rbpms2*-null animals (FC=-43.00; p-adj1.53E-09; Fig. S8B; Table S2). The

other *pln* transcript (ENSDART00000133911**;** *pln-202***)** was unchanged (FC=1.37; p-adj>0.99;

Table S2), which highlighted the possibility that Rbpms2 functions in the zebrafish myocardium

to regulate alternative splicing, a molecular function previously ascribed to the closely related

RBPMS protein [49], but not to RBPMS2 itself [35].

To investigate this possibility directly, we performed replicate multivariate analysis of

transcript splicing (rMATS) [50] on the same RNA-sequencing dataset and identified 150

differential alternative splicing events (ASEs) in *rbpms2*-null animals (0 uncalled replicates;

FDR<0.01; |Incleveldifference|>0.1; Fig. 3A,B; Table S3). The most common ASE was skipped

exon (SE), representing over 40% of the total ASEs, followed by alternative 3' splice site (A3SS,

23%), alternative 5' splice site (A5SS, 21%), mutually exclusive exons (MXE, 10%), and

retained intron (RI, 6%) (Fig 3A,C). Gene ontology (GO) term enrichment analysis did not

identify overrepresentation of any terms among the differentially spliced genes. However, a

limited number of myocardial genes were found to be differentially spliced in *rbpms2*-null

animals including *myosin binding protein C3* (*mybpc3*) and *myomesin 2a* (*myom2a*), both of

which encode sarcomere components [51–54], and *solute carrier family 8 member 1a* [*slc8a1a*;

also termed the *sodium calcium exchanger 1* (*ncx1*)] which encodes a plasma membrane

transporter responsible for extruding calcium from cardiomyocytes during cardiac relaxation

[55,56].

In the case of *mybpc3*, exons 2 and exons 10 were identified as cassette exons whose

inclusion levels (Inclevels) in mature *mybpc3* transcripts were found to be significantly lower in

rbpms2-null animals (exon2 Inclevel=0.031; exon 10 Inclevel=0.057) when compared to those in

control siblings (exon2 Inclevel=0.557; exon10 Inclevel=0.296; exon 2 Incleveldifference=0.526,

FDR=4.77E-40; exon 10 Incleveldifference=0.24, FDR=2.38E-21; Fig. 3B; Fig.4A; Table S4).

The same was true for exon 16 of *myom2a* (control Inclevel=1, DKO Inclevel=0,

Incleveldifference=1, FDR 2.74E-20; Fig. 3B; Fig.4A; Table S4) and exon 5 of *slc8a1a* (control

Inclevel=1, DKO Inclevel=0.37, Incleveldifference=0.629, FDR=4.92E-14; Fig. 3B; Fig.4A; Table

S4). In all cases, exon inclusion was higher in in control siblings, which means that these exon

sequences were significantly reduced or absent in *rbpms2*-null animals.

Despite supporting evidence from differential isoform expression analysis (Fig. S8B;

Table S2), rMATS did not identify alternative splicing of *pln*, likely because it involves a terminal

exon (Fig. 4A), which rMATS is not designed to evaluate (Fig. 3A). Therefore, we employed the

mixture of isoforms (MISO) algorithm [57] to evaluate alternative splicing of *pln* in *rbpms2*-null

animals and found strong statistical evidence that the exon 3-containing *pln*-203 isoform

(ENSDART00000141159) is uniquely reduced in mutant animals. Specifically, the Bayes factor

was largely above the accepted cutoff of 5 for differences in ENSDART00000141159 versus

ENSDART00000133911 usage in WT and double-mutant animals across all five replicate pairs

(Table S5), which is strongly suggestive of differential alternative splicing. This conclusion was

also supported by genotype-specific differences in read densities across the locus (Fig. S9).

Collectively, these data demonstrate that *rbpms2*-null embryos display abnormal splicing of

genes that are essential to cardiac function.

Validation of differential alternative splicing events in rbpms2-null zebrafish

Next, we used splicing-sensitive RT-qPCR to validate differential alternative splicing of

mybpc3, *myom2a*, *slc8a1*, and *pln* in *rbpms2*-null animals at 48 hpf (Fig. 4A). To that end, we

designed two primer pairs for each transcript. One was designed to measure relative "total

transcript" levels between sibling control and *rbpms2*-null animals, independent of the

alternatively spliced exon. The other was designed to quantify relative "isoform-specific" levels

by having one of the primer-binding sites contained within the exon of reduced inclusion

according to rMATS or MISO (Fig. 4A). For all four genes, total transcript levels were not

significantly altered in *rbpms*2-null animals (Fig. 4B). However, the specific exon-containing

isoforms were reduced by at least 50% in all cases, and by greater than 95% in three cases,

indicative of almost-complete exon exclusion (Fig. 4C). Taken together, these data provide

strong evidence that Rbpms2 activity promotes the inclusion of specific exons into mature

cardiac transcripts. Importantly, we verified that *mybpc3* and *pln* are also abnormally spliced

earlier in development (33 hpf), when signs of cardiac distress first become visible in double

mutants, reducing the likelihood that abnormal splicing is a secondary consequence of cardiac

dysfunction (Fig. S7E,F).

To confirm that the DKO splicing abnormalities localize to the myocardium, we

performed splicing-sensitive in situ hybridization analysis for two genes, *mybpc3* and *pln*, with

rbpms2-dependent exons large enough to generate effective exon-specific anti-sense

riboprobes (i.e. exon 2 of *mybpc3* and exon 3 of *pln*; Fig. 4A). In a strategy analogous to the

splicing-sensitive qPCR, we designed a pair of riboprobes for each gene. One probe detected

the total transcript pools of *mybpc3* or *pln*, independent of the alternatively spliced exons

("Total-transcript" in situ hybridization). The other probe detected the alternatively spliced exon

sequences exclusively. In control-sibling animals at 33 hpf, total *mybpc3* transcripts were

localized to both cardiac and skeletal muscle (Fig. 4D,D'), and total *pln* transcripts were

detected exclusively in the heart (Fig. 4H,H'). These expression patterns were unaffected by the

loss of *rbpms2* (Fig. 4E,E',I,I'), which is consistent with the total transcript qPCR analysis (Fig.

4B). Isoform-specific in situ hybridization analysis revealed expression of the *rbpms2*-dependent

mybpc3 and *pln* exons exclusively in the hearts of control-sibling animals (Fig. 4F,F',J,J').

Consistent with the isoform-specific qPCR analysis (Fig. 4C), double-mutant animals were

completely devoid of these exon sequences (Fig. 4G,G',K,K'). Taken together, these data

implicate *rbpms2* as a positive regulator of exon inclusion in the zebrafish embryonic

myocardium.

For those alternative splicing events validated by qPCR or in situ hybridization (*mybpc3*,

myom2a, *pln*, and *slc8a1a*), exon exclusion in mutant hearts maintains the open reading frame,

indicating that the hearts of mutant animals express proteins lacking amino acid sequences

encoded by the alternatively spliced exons (Fig. 4E,E',G,G'I,I',K,K'). In the case of *mybpc3,*

exon 2 encodes a domain termed C0, which is contained exclusively within the cardiac isoform

of Myosin Binding Protein C in higher vertebrates (MYBPC3) but is absent from the skeletal

muscle isoform [MYBPC2; [58]]. Similarly, *mybpc3*'s exon 10 encodes cardiac-specific amino

acids that become phosphorylated by protein kinase A (PKA) in response to β-adrenergic

signaling [59,60]. For Myom2a, the short stretch of amino acids that disappears in DKO animals is

of unknown function. For *pln*, excluding exon 3 shortens the protein by 5 amino acids on the C-

terminus, which has been shown to increase SERCA inhibitory activity *in vitro* [6]. Lastly, the

slc8a1a exon with reduced inclusion in DKO animals is one of six conserved exons that are

alternatively spliced in higher vertebrates to control the Ca2+ sensitivity of the transporter [61,62].

Zebrafish rbpms2-null ventricular cardiomyocytes exhibit myofibrillar disarray and calcium

handling defects

Because Mybpc3 and Myom2a are sarcomere components, we evaluated 48 hpf

control-sibling and double-mutant hearts for myofibrillar defects after immunostaining with

antibodies that recognize thin filaments, thick filaments, or Z-disks. For all epitopes, prominent

striations were apparent in control-sibling animals reflecting the tandem arrays of sarcomeres

composing myofibrils, which themselves are transversely aligned and bundled near the cell

cortices (Fig. 5A,A',C-C",E). By contrast, despite the readily-evident detection of epitopes in

DKO hearts, striations were significantly reduced or absent because the myofibrils were in a

state of disarray (Fig. 5B,B',D-D",F). We confirmed that myofibril disarray was also present in

ventricular cardiomyocytes of double mutant animals at an earlier developmental stage of 33 hpf

(Fig. S7G-H'). Lastly, this defect is specific to ventricular cardiomyocytes because myofibril

organization in atrial cardiomyocytes of control and DKO embryos was grossly indistinguishable

(Fig. S7I-J'). Taken together, these data demonstrate that Rbpms2 activity is required for

establishing or maintaining the three-dimensional architecture of myofibrils in ventricular

cardiomyocytes of the zebrafish embryo.

Lastly, because *pln* and *slc8a1a* regulate calcium handing in cardiomyocytes [56,63–65], we

evaluated *rbpms2*-null hearts for abnormalities in the amplitude or duration of cytosolic calcium

transients, which stimulate cardiomyocyte contraction. To that end, control-sibling and *rbpms2*-

null hearts at 48 hpf were explanted and loaded with the dual-wavelength calcium-sensitive dye

Fura-2. High-speed imaging was performed to obtain a ratiometric indicator of fluctuating

calcium in cardiomyocytes over several cardiac cycles [66]. From this, we measured the duration

and amplitude of the calcium transients in the atrium and ventricle of control-sibling and *rbpms2*-

null hearts. Whereas these values were not significantly different in the atrium of double mutant

hearts (Fig. 6A,B,D-F,H), both the duration and amplitude of the calcium transients in double

mutant ventricles were reduced in the mutant ventricle (Fig. 6A-C; E-G), demonstrating that

Rbpms2 is indispensable for the regulation of calcium handling in ventricular cardiomyocytes of

the zebrafish embryo.

Human RBPMS2-null cardiomyocytes exhibit myofibrillar disarray and calcium handling defects

To determine if the cellular function of *RBPMS2* is conserved in human cardiomyocytes,

we targeted *RBPMS2* in human induced pluripotent stem cells (hiPSCs) using CRISPR/Cas9-

mediated genome editing and analyzed phenotypes that arose following differentiation into

hiPSC-derived cardiomyocytes (hiPSC-CMs). We targeted the DNA sequence encoding

RBPMS2's RNA recognition motif and isolated a clone with a biallelic 2 base-pair deletion (Fig.

S10A; *ΔRBPMS*), which truncates both the protein and the RRM by ~70% due to a premature

stop codon (Fig. 7A; Fig. S10B). Quantitative RT-PCR analysis revealed a significant 82%

reduction in *RBPMS2* transcripts in *ΔRBPMS* hiPSCs (Fig. 7B) and hiPSC-CMs (Fig. 7C),

suggesting that the mutant transcripts are susceptible to non-sense mediated decay. This

observation, in combination with the large size of the C-terminal truncation (Fig. 7A) strongly

suggest that *ΔRBPMS2* cells are null for *RBPMS2*.

To determine if *RBPMS2*-null human cardiomyocytes exhibit cellular phenotypes similar

to those observed in ventricular cardiomyocytes from *rbpms2*-null zebrafish (Fig. 5A-F), we

analyzed myofibril structure in control-parental and *ΔRBPMS2* hiPSC-CMs by double

immunostaining with antibodies that detect thin filaments or Z-disks. The hallmark striations

created by thin filaments, which were readily apparent in control cells (Fig. 7D,E,G), were

almost completely absent in *ΔRBPMS2* hiPSC-CMs (Fig. 7H,I,K). Moreover, whereas the Z-

disks in control cardiomyocytes were prominent, well defined and arrayed in parallel (Fig.

7D,F,G), those in *ΔRBPMS2* hiPSC-CMs were smaller, punctate, and disorganized (Fig.

7H,J,K). Taken together, these data uncover a conserved role for RBPMS2 in establishing or

maintaining myofibril organization in human cardiomyocytes.

Lastly, we evaluated control-parental and *ΔRBPMS2* hiPSC-CMs for defects in calcium

handling. hiPSC-CMs cells were loaded with the calcium-sensitive dye Fluo-4, paced at 1Hz

and imaged by line scanning confocal microscopy to capture intracellular Ca2+ dynamics (Fig.

7L,M). Similar to ventricular cardiomyocytes from *rbpms2*-null zebrafish, *ΔRBPMS2* hiPSC-CMs

exhibited significant decreases in Ca2+-transient amplitude (Fig. 7N) and transient duration

(Fig. 7O,P). These phenotypes were accompanied by decreases in time-to-peak (Fig. 7Q) and

maximum upstroke velocity (Fig. 7R). Lastly, without pacing, *ΔRBPMS2* hiPSC-CMs contracted

at a lower spontaneous beat frequency (Fig. 7S). These data demonstrate that human

cardiomyocytes rely on RBPMS2 function for optimal calcium handling. Taken together, the

overlap in sarcomere and calcium handling phenotypes observed between *Rbpms2*-null

zebrafish and *ΔRBPMS2* hiPSC-CMs suggest that the cellular function(s) of this myocardial

RNA-binding protein is conserved in human cardiomyocytes.

DISCUSSION

Our study identifies Rbpms2 proteins as critical RNA splicing factors that regulate tissue-

specific isoform expression within the ventricular myocardium. Specifically, we show that

zebrafish Rbpms2a and Rbpms2b are redundantly required for ventricular form and function in

embryonic zebrafish. We then show that Rbpms2 proteins function to promote the generation of

distinct isoforms within the heart, including a previously undocumented cardiac-specific isoform

of *mybpc3*. Further, we found that *rbpms2*-null hearts have disorganized sarcomeres and

disrupted calcium handling, which is phenotypically consistent with the broad classes of

alternatively spliced transcripts identified by our RNA-seq. Lastly, we discovered that RBPMS2-

deficient human iPSC-CMs exhibit defects in sarcomere structure and calcium handling that are

strikingly similar to what we observed in zebrafish ventricles.

Our characterization of Rbpms2 provides further evidence that RNA splicing within the

heart may be a much more pervasive and physiologically important mode of gene regulation

than previously appreciated. While our results reveal a clear connection between Rbpms2 and

RNA splicing, these proteins may function in additional capacities within tissues outside the

heart. Previous reports in zebrafish have shown that Rbpms2 proteins contribute to the

establishment of oocyte polarity and development by influencing the localization of RNAs such

as *bucky ball* (*buc*) to the Balbiani body[33,67,68]. Further research in xenopus and chicken have

described similar roles for Rbpms2 in RNA stability and localization within intestinal smooth

muscle, kidney cells, and retinal ganglion cells[38,69,70]. Interestingly, the localization of Rbpms2 to

stress granules has been observed in several of these models[69,71], indicating that this function

may be conserved in multiple tissues. It will be important to determine if RBPMS2 functions in

this capacity within the heart, or whether this is a tissue-specific function that is restricted to

other RBPMS2-positive organs such as the eye and nephron.

Even though we found that Rbpms2 regulates the splicing of genes whose orthologs are

known to cause cardiomyopathy, it is unclear whether mutations in RBPMS2 segregate with

human disease. Our data therefore warrants a careful re-evaluation of existing GWAS datasets

and provides new incentives to examine the RBPMS2 locus in CHD patients. However, given

the severity of cardiac phenotypes observed in our loss-of-function experiments, it remains

possible that embryonic lethality may confound GWAS results. Additionally, the upstream

regulators of Rbpms2 remain unknown. Further studies will be necessary to identify these

upstream factors in order to determine if they contribute to disease states through a more subtle

modulation of Rbpms2 function. Given that Rbpms2 is itself alternatively spliced in vertebrates,

it is also possible that its function may be regulated by yet another upstream splicing factor.

Collectively these studies will evaluate RBPMS2 function within the broader context of human

cardiomyopathy and heart failure.

ACKNOWLEDGEMENTS

Massachusetts General Hospital Computational and Integrative Biology DNA Core (Director:

Amy Avery).

FUNDING

AAA was supported by an American Heart Association Postdoctoral Fellowship

(20POST35110027). AAA and AS were supported by a National Institutes of Health (NIH)

training grant awarded to Massachusetts General Hospital (MGH) (T32HL007208; PI: A.

Rosenzweig). The Burns Laboratory was supported by NIH grants 7R01HL139806 (formerly

1R01HL139806) and 7R35HL135831 (formerly 1R35H135831) to CGB and CEB, respectively,

by awards from the Executive Committee on Research at MGH to CGB, and by Department of

Defense Peer-Reviewed Medical Research Program (PRMRP) grant PR190628 to CEB. CEB

was a d'Arbelhoff MGH Research Scholar. CGB was a Hassenfeld Cardiovascular Scholar at

MGH. The Burns Laboratory was supported, in part, by funds from the Boston Children's

Hospital Department of Cardiology.

MT was supported by an NIH training grant awarded to Boston Children's Hospital

(T32HL007572; PI: W. Pu). The bioinformatics analysis was supported in part by the Koch

Institute Support (core) Grant P30-CA14051 from the National Cancer Institute.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES

1. Virani SS, Alonso A, Aparicio HJ, Benjamin EJ, Bittencourt MS, Callaway CW, Carson AP,
Chamberlain AM, Cheng S, Delling FN, Elkind MSV, Evenson KR, Ferguson JF, Gupta DK, Khan SS, et
al. Heart Disease and Stroke Statistics—2021 Update: A Report From the American Heart Association.
Circulation. 2021:CIR0000000000000950.

2. Skrzynia C, Berg J, Willis M, Jensen B. Genetics and Heart Failure: A Concise Guide for the Clinician.
Current Cardiology Reviews. 2014;11(1):10–17.

3. Garfinkel AC, Seidman JG, Seidman CE. Genetic Pathogenesis of Hypertrophic and Dilated
Cardiomyopathy. *Heart Failure Clinics*. 2018;14(2):139–146.

4. Towbin JA, Bowles NE. Genetic abnormalities responsible for dilated cardiomyopathy. *Current*
Cardiology Reports. 2000;2(5):475–480.

5. Towbin JA. Molecular genetics of hypertrophic cardiomyopathy. *Current Cardiology Reports*.
2000;2(2):134–140.

6. Gorski PA, Ceholski DK, Hajjar RJ. Altered Myocardial Calcium Cycling and Energetics in Heart
Failure—A Rational Approach for Disease Treatment. *Cell Metabolism*. 2015;21(2):183–194.

7. Hoogenhof MMG van den, Beqqali A, Amin AS, Made I van der, Aufiero S, Khan MAF, Schumacher
CA, Jansweijer JA, Spaendonck-Zwarts KY van, Remme CA, Backs J, Verkerk AO, Baartscheer A, Pinto
YM, Creemers EE. RBM20 Mutations Induce an Arrhythmogenic Dilated Cardiomyopathy Related to
Disturbed Calcium Handling. *Circulation*. 2018;138(13):1330–1342.

8. Guo W, Schafer S, Greaser ML, Radke MH, Liss M, Govindarajan T, Maatz H, Schulz H, Li S, Parrish
AM, Dauksaite V, Vakeel P, Klaassen S, Gerull B, Thierfelder L, et al. RBM20, a gene for hereditary
cardiomyopathy, regulates titin splicing. *Nature Medicine*. 2012;18(5):766–773.

9. Maatz H, Jens M, Liss M, Schafer S, Heinig M, Kirchner M, Adami E, Rintisch C, Dauksaite V, Radke
MH, Selbach M, Barton PJR, Cook SA, Rajewsky N, Gotthardt M, et al. RNA-binding protein RBM20
represses splicing to orchestrate cardiac pre-mRNA processing. *Journal of Clinical Investigation*.
2014;124(8):3419–3430.

10. LeWinter MM, Granzier HL. Titin Is a Major Human Disease Gene. *Circulation*. 2013;127(8):938–
944.

11. Haas J, Frese KS, Peil B, Kloos W, Keller A, Nietsch R, Feng Z, Müller S, Kayvanpour E, Vogel B,
Sedaghat-Hamedani F, Lim W-K, Zhao X, Fradkin D, Köhler D, et al. Atlas of the clinical genetics of
human dilated cardiomyopathy. *European Heart Journal*. 2015;36(18):1123–1135.

12. Refaat MM, Lubitz SA, Makino S, Islam Z, Frangiskakis JM, Mehdi H, Gutmann R, Zhang ML,
Bloom HL, MacRae CA, Dudley SC, Shalaby AA, Weiss R, McNamara DM, London B, et al. Genetic
variation in the alternative splicing regulator RBM20 is associated with dilated cardiomyopathy. *Heart*
Rhythm. 2012;9(3):390–396.

13. Liu J, Kong X, Zhang M, Yang X, Xu X. RNA binding protein 24 deletion disrupts global alternative
splicing and causes dilated cardiomyopathy. *Protein & Cell*. 2019;10(6):405–416.

14. Gaertner A, Brodehl A, Milting H. Screening for mutations in human cardiomyopathy- is RBM24 a
new but rare disease gene? *Protein & Cell*. 2019;10(6):393–394.

15. Jin JP, Lin JJ. Isolation and Characterization of cDNA Clones Encoding Embryonic and Adult
Isoforms of Rat Cardiac Troponin T. *Journal of Biological Chemistry*. 1989;264(24):14471–14477.

16. Farza H, Townsend PJ, Carrier L, Barton PJ, Mesnard L, Bährend E, Forissier J-F, Fiszman M,
Yacoub MH, Schwartz K. Genomic Organisation, Alternative Splicing and Polymorphisms of the Human
Cardiac Troponin T Gene. *Journal of Molecular and Cellular Cardiology*. 1998;30(6):1247–1253.

17. Biesiadecki BJ, Elder BD, Yu Z-B, Jin J-P. Cardiac Troponin T Variants Produced by Aberrant
Splicing of Multiple Exons in Animals with High Instances of Dilated Cardiomyopathy*. *Journal of
Biological Chemistry*. 2002;277(52):50275–50285.

18. Niimura H, Bachinski LL, Sangwatanaroj S, Watkins H, Chudley AE, McKenna W, Kristinsson A,
Roberts R, Sole M, Maron BJ, Seidman JG, Seidman CE, Thierfelder L, Jarcho JA, Anastasakis A, et al.
Mutations in the Gene for Cardiac Myosin-Binding Protein C and Late-Onset Familial Hypertrophic
Cardiomyopathy. *The New England Journal of Medicine*. 1998;338(18):1248–1257.

19. MacLennan DH, Kranias EG. Phospholamban: a crucial regulator of cardiac contractility. *Nature
Reviews Molecular Cell Biology*. 2003;4(7):566–577.

20. Zhou Y, Cashman TJ, Nevis KR, Obregon P, Carney SA, Liu Y, Gu A, Mosimann C, Sondalle S,
Peterson RE, Heideman W, Burns CE, Burns CG. Latent TGF-β binding protein 3 identifies a second
heart field in zebrafish. *Nature*. 2011;474(7353):645–648.

21. Burns CG, Milan DJ, Grande EJ, Rottbauer W, MacRae CA, Fishman MC. High-throughput assay for
small molecules that modulate zebrafish embryonic heart rate. *Nature Chemical Biology*. 2005;1(5):263–
264.

22. González-Rosa JM, Sharpe M, Field D, Soonpaa MH, Field LJ, Burns CE, Burns CG. Myocardial
Polyploidization Creates a Barrier to Heart Regeneration in Zebrafish. *Developmental Cell*.
2018;44(4):433-446.e7.

23. Lin Y-F, Swinburne I, Yelon D. Multiple influences of blood flow on cardiomyocyte hypertrophy in
the embryonic zebrafish heart. *Developmental Biology*. 2012;362(2):242–253.

24. Sander JD, Cade L, Khayter C, Reyon D, Peterson RT, Joung JK, Yeh J-RJ. Targeted gene disruption
in somatic zebrafish cells using engineered TALENs. *Nature Biotechnology*. 2011;29(8):697–698.

25. Ma AC, Lee HB, Clark KJ, Ekker SC. High efficiency In Vivo genome engineering with a simplified
15-RVD GoldyTALEN design. Müller F, ed. *PloS one*. 2013;8(5):e65259.

26. Sanjana NE, Cong L, Zhou Y, Cunniff MM, Feng G, Zhang F. A transcription activator-like effector
toolbox for genome engineering. *Nature Protocols*. 2012;7(1):171–192.

27. Foley JE, Yeh J-RJ, Maeder ML, Reyon D, Sander JD, Peterson RT, Joung JK. Rapid Mutation of
Endogenous Zebrafish Genes Using Zinc Finger Nucleases Made by Oligomerized Pool ENgineering
(OPEN). *PLoS ONE*. 2009;4(2):e4348.

28. Wang G, Yang L, Grishin D, Rios X, Ye LY, Hu Y, Li K, Zhang D, Church GM, Pu WT. Efficient,
footprint-free human iPSC genome editing by consolidation of Cas9/CRISPR and piggyBac technologies.
Nature Protocols. 2017;12(1):88–103.

29. Guner-Ataman B, Paffett-Lugassy N, Adams MS, Nevis KR, Jahangiri L, Obregon P, Kikuchi K,
Poss KD, Burns CE, Burns CG. Zebrafish second heart field development relies on progenitor
specification in anterior lateral plate mesoderm and nkx2.5 function. *Development (Cambridge, England)*.
2013;140(6):1353–1363.

30. Paffett-Lugassy N, Singh R, Nevis KR, Guner-Ataman B, O'Loughlin E, Jahangiri L, Harvey RP,
Burns CG, Burns CE. Heart field origin of great vessel precursors relies on nkx2.5-mediated
vasculogenesis. *Nature Cell Biology*. 2013;15(11):1362–1369.

31. Schoenebeck JJ, Keegan BR, Yelon D. Vessel and blood specification override cardiac potential in
anterior mesoderm. *Developmental cell*. 2007;13(2):254–267.

32. Serbedzija GN, Chen JN, Fishman MC. Regulation in the heart field of zebrafish. *Development
(Cambridge, England)*. 1998;125(6):1095–1101.

33. Kaufman OH, Lee K, Martin M, Rothhämel S, Marlow FL. rbpms2 functions in Balbiani body
architecture and ovary fate. *PLoS genetics*. 2018;14(7):e1007489.

34. Gerber WV, Yatskievych TA, Antin PB, Correia KM, Conlon RA, Krieg PA. The RNA-binding
protein gene, hermes, is expressed at high levels in the developing heart. *Mechanisms of development*.
1999;80(1):77–86.

35. Akerberg AA, Burns CE, Burns CG. Exploring the Activities of RBPMS Proteins in Myocardial
Biology. *Pediatric cardiology*. 2019;40(7):1410–1418.

36. Wilmore HP, McClive PJ, Smith CA, Sinclair AH. Expression profile of the RNA-binding protein
gene hermes during chicken embryonic development. *Developmental dynamics : an official publication of
the American Association of Anatomists*. 2005;233(3):1045–1051.

37. Yelon D, Horne SA, Stainier DY. Restricted expression of cardiac myosin genes reveals regulated
aspects of heart tube assembly in zebrafish. *Developmental biology*. 1999;214(1):23–37.

38. Hörnberg H, Horck FW, Maurus D, Zwart M, Svoboda H, Harris WA, Holt CE. RNA-binding protein
Hermes/RBPMS inversely affects synapse density and axon arbor formation in retinal ganglion cells in
vivo. *The Journal of neuroscience : the official journal of the Society for Neuroscience*.
2013;33(25):10384–10395.

39. Pater E de, Clijsters L, Marques SR, Lin Y-F, Garavito-Aguilar ZV, Yelon D, Bakkers J. Distinct
phases of cardiomyocyte differentiation regulate growth of the zebrafish heart. *Development (Cambridge,
England)*. 2009;136(10):1633–1641.

40. Lazic S, Scott IC. Mef2cb regulates late myocardial cell addition from a second heart field-like
population of progenitors in zebrafish. *Developmental biology*. 2011;354(1):123–133.

41. Hami D, Grimes AC, Tsai H-J, Kirby ML. Zebrafish cardiac development requires a conserved
secondary heart field. *Development (Cambridge, England)*. 2011;138(11):2389–2398.

42. Akerberg AA, Burns CE, Burns CG, Nguyen C. Deep learning enables automated volumetric
assessments of cardiac function in zebrafish. *Disease models & mechanisms*. 2019;12(10):dmm040188.

43. Cuspidi C, Negri F, Sala C, Valerio C, Mancia G. Association of left atrial enlargement with left
ventricular hypertrophy and diastolic dysfunction: A tissue Doppler study in echocardiographic practice.
Blood Pressure. 2011;21(1):24–30.

44. Cuspidi C, Rescaldani M, Sala C. Prevalence of Echocardiographic Left-Atrial Enlargement in
Hypertension: A Systematic Review of Recent Clinical Studies. *American Journal of Hypertension*.
2013;26(4):456–464.

45. Huttner IG, Wang LW, Santiago CF, Horvat C, Johnson R, Cheng D, Frieling-Salewsky M von,
Hillcoat K, Bemand TJ, Trivedi G, Braet F, Hesselson D, Alford K, Hayward CS, Seidman JG, et al. A-
Band Titin Truncation in Zebrafish Causes Dilated Cardiomyopathy and Hemodynamic Stress
Intolerance. *Circulation: Genomic and Precision Medicine*. 2018;11(8):e002135.

46. Horsthuis T, Houweling AC, Habets PEMH, Lange FJ de, Azzouzi H el, Clout DEW, Moorman
AFM, Christoffels VM. Distinct regulation of developmental and heart disease-induced atrial natriuretic
factor expression by two separate distal sequences. *Circulation research*. 2008;102(7):849–859.

47. Grassini DR, Lagendijk AK, Angelis JED, Silva JD, Jeanes A, Zettler N, Bower NI, Hogan BM,
Smith KA. Nppa and Nppb act redundantly during zebrafish cardiac development to confine AVC marker
expression and reduce cardiac jelly volume. *Development (Cambridge, England)*.
2018;145(12):dev160739.

48. Berdougo E, Coleman H, Lee DH, Stainier DYR, Yelon D. Mutation of weak atrium/atrial myosin
heavy chain disrupts atrial function and influences ventricular morphogenesis in zebrafish. *Development
(Cambridge, England)*. 2003;130(24):6121–6129.

49. Nakagaki-Silva EE, Gooding C, Llorian M, Jacob AG, Richards F, Buckroyd A, Sinha S, Smith CW.
Identification of RBPMS as a mammalian smooth muscle master splicing regulator via proximity of its
gene with super-enhancers. *eLife*. 2019;8.

50. Shen S, Park JW, Lu Z, Lin L, Henry MD, Wu YN, Zhou Q, Xing Y. rMATS: robust and flexible
detection of differential alternative splicing from replicate RNA-Seq data. *Proceedings of the National
Academy of Sciences of the United States of America*. 2014;111(51):E5593-601.

51. Agarkova I, Perriard J-C. The M-band: an elastic web that crosslinks thick filaments in the center of
the sarcomere. *Trends in Cell Biology*. 2005;15(9):477–485.

52. Gautel M, Djinović-Carugo K. The sarcomeric cytoskeleton: from molecules to motion. *Journal of
Experimental Biology*. 2016;219(2):135–145.

53. Previs MJ, Previs SB, Gulick J, Robbins J, Warshaw DM. Molecular Mechanics of Cardiac Myosin-
Binding Protein C in Native Thick Filaments. *Science*. 2012;337(6099):1215–1218.

54. Previs MJ, Prosser BL, Mun JY, Previs SB, Gulick J, Lee K, Robbins J, Craig R, Lederer WJ,
Warshaw DM. Myosin-binding protein C corrects an intrinsic inhomogeneity in cardiac excitation-
contraction coupling. *Science Advances*. 2015;1(1):e1400205.

55. Ebert AM, Hume GL, Warren KS, Cook NP, Burns CG, Mohideen MA, Siegal G, Yelon D, Fishman
MC, Garrity DM. Calcium extrusion is critical for cardiac morphogenesis and rhythm in embryonic

zebrafish hearts. *Proceedings of the National Academy of Sciences of the United States of America*.
2005;102(49):17705–17710.

56. Langenbacher AD, Dong Y, Shu X, Choi J, Nicoll DA, Goldhaber JI, Philipson KD, Chen J-N.
Mutation in sodium–calcium exchanger 1 (NCX1) causes cardiac fibrillation in zebrafish. *Proceedings of*
the National Academy of Sciences of the United States of America. 2005;102(49):17699–17704.

57. Katz Y, Wang ET, Airoldi EM, Burge CB. Analysis and design of RNA sequencing experiments for
identifying isoform regulation. *Nature Methods*. 2010;7(12):1009–1015.

58. McNamara JW, Sadayappan S. Skeletal myosin binding protein-C: An increasingly important
regulator of striated muscle physiology. *Archives of Biochemistry and Biophysics*. 2018;660:121–128.

59. Gautel M, Zuffardi O, Freiburg A, Labeit S. Phosphorylation switches specific for the cardiac isoform
of myosin binding protein-C: a modulator of cardiac contraction? *The EMBO Journal*. 1995;14(9):1952–
1960.

60. Mohamed AS, Dignam JD, Schlender KK. Cardiac Myosin-Binding Protein C (MyBP-C):
Identification of Protein Kinase A and Protein Kinase C Phosphorylation Sites. *Archives of Biochemistry*
and Biophysics. 1998;358(2):313–319.

61. Kofuji P, Lederer WJ, Schulze DH. Mutually exclusive and cassette exons underlie alternatively
spliced isoforms of the Na/Ca exchanger. *Journal of Biological Chemistry*. 1994;269(7):5145–5149.

62. Quednau BD, Nicoll DA, Philipson KD. Tissue specificity and alternative splicing of the Na+/Ca2+
exchanger isoforms NCX1, NCX2, and NCX3 in rat. *American Journal of Physiology-Cell Physiology*.
1997;272(4):C1250–C1261.

63. Simmerman HKB, Jones LR. Phospholamban: Protein Structure, Mechanism of Action, and Role in
Cardiac Function. *Physiological Reviews*. 1998;78(4):921–947.

64. Toyoshima C, Asahi M, Sugita Y, Khanna R, Tsuda T, MacLennan DH. Modeling of the inhibitory
interaction of phospholamban with the Ca2+ ATPase. *Proceedings of the National Academy of Sciences*.
2003;100(2):467–472.

65. Koushik SV, Wang J, Rogers R, Moskophidis D, Lambert NA, Creazzo TL, Conway SJ. Targeted
inactivation of the sodium-calcium exchanger (Ncx1) results in the lack of a heartbeat and abnormal
myofibrillar organization. *The FASEB Journal*. 2001;15(7):1209–1211.

66. Russell JT. Imaging calcium signals in vivo: a powerful tool in physiology and pharmacology. *British*
Journal of Pharmacology. 2011;163(8):1605–1625.

67. Kosaka K, Kawakami K, Sakamoto H, Inoue K. Spatiotemporal localization of germ plasm RNAs
during zebrafish oogenesis. *Mechanisms of Development*. 2007;124(4):279–289.

68. Heim AE, Hartung O, Rothhämel S, Ferreira E, Jenny A, Marlow FL. Oocyte polarity requires a
Bucky ball-dependent feedback amplification loop. *Development*. 2014;141(4):842–854.

69. Farazi TA, Leonhardt CS, Mukherjee N, Mihailovic A, Li S, Max KEA, Meyer C, Yamaji M, Cekan
P, Jacobs NC, Gerstberger S, Bognanni C, Larsson E, Ohler U, Tuschl T. Identification of the RNA
recognition element of the RBPMS family of RNA-binding proteins and their transcriptome-wide mRNA
targets. *RNA (New York, N.Y.)*. 2014;20(7):1090–1102.

70. Notarnicola C, Rouleau C, Guen LL, Virsolvy A, Richard S, Faure S, Barbara PDS. The RNA-
Binding Protein RBPMS2 Regulates Development of Gastrointestinal Smooth Muscle. *Gastroenterology*.
2012;143(3):687-697.e9.

71. Furukawa MT, Sakamoto H, Inoue K. Interaction and colocalization of HERMES/RBPMS with
NonO, PSF, and G3BP1 in neuronal cytoplasmic RNP granules in mouse retinal line cells. *Genes to
Cells*. 2015;20(4):257–266.

72. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics. 2010;26(6):841–842.

73. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-aware
quantification of transcript expression. *Nature Methods*. 2017;14(4):417–419.

74. Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq: transcript-level estimates
improve gene-level inferences. *F1000Research*. 2016;4:1521.

75. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR.
STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*. 2013;29(1):15–21.

76. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or without a
reference genome. *BMC Bioinformatics*. 2011;12(1):323.

77. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data
with DESeq2. *Genome Biology*. 2014;15(12):550.

78. Katz Y, Wang ET, Silterra J, Schwartz S, Wong B, Thorvaldsdóttir H, Robinson JT, Mesirov JP,
Airoldi EM, Burge CB. Quantitative visualization of alternative exon expression from RNA-seq data.
Bioinformatics. 2015;31(14):2400–2402.

79. Thisse C, Thisse B. High-resolution in situ hybridization to whole-mount zebrafish embryos. *Nature
protocols*. 2008;3(1):59–69.

80. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. *Nature
methods*. 2012;9(7):671–675.

81. Jahangiri L, Sharpe M, Novikov N, González-Rosa JM, Borikova A, Nevis K, Paffett-Lugassy N,
Zhao L, Adams M, Guner-Ataman B, Burns CE, Burns CG. The AP-1 transcription factor component
Fosl2 potentiates the rate of myocardial differentiation from the zebrafish second heart field.
Development. 2016;143(1):113–122.

82. Poss KD, Wilson LG, Keating MT. Heart Regeneration in Zebrafish. *Science*. 2002;298(5601):2188–
2190.

83. Akerberg AA, Henner A, Stewart S, Stankunas K. Histone demethylases Kdm6ba and Kdm6bb
redundantly promote cardiomyocyte proliferation during zebrafish heart ventricle maturation.
Developmental Biology. 2017;426(1):84–96.

84. Power RM, Huisken J. A guide to light-sheet fluorescence microscopy for multiscale imaging. *Nature*
Methods. 2017;14(4):360–373.

85. Huisken J, Stainier DYR. Even fluorescence excitation by multidirectional selective plane
illumination microscopy (mSPIM). *Optics Letters*. 2007;32(17):2608.

86. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR
and the 2(-Delta Delta C(T)) Method. *Methods (San Diego, Calif.)*. 2001;25(4):402–408.

87. Panáková D, Werdich AA, MacRae CA. Wnt11 patterns a myocardial electrical gradient through
regulation of the L-type Ca(2+) channel. *Nature*. 2010;466(7308):874–878.

    FIGURES AND LEGENDS

Figure 1 – Restricted expression of RNA-binding proteins Rbpms2a and Rbpms2b to the

zebrafish myocardium. (A) Workflow used to identify 1848 protein-coding RNAs enriched in

ZsYellow-positive (ZsY+) cardiopharyngeal progenitors or cardiomyocytes in the anterior lateral

plate mesoderm of 14-16 somite stage (ss) *Tg(nkx2.5:ZsYellow)* embryos. (B-G) Brightfield

images of 16 ss (B,E), 28 hours post fertilization (hpf; C,F), and 48 hpf (D,G) embryos

processed for in situ hybridization with riboprobes that detect *rbpms2a* (B-D) or *rbpms2b* (E-G).

Little to no variation in expression pattern was observed between animals in each group

(n>10/group). (H-K') Confocal projections of hearts in 24 hpf (H,H'), 33 hpf (I,I'), 48 hpf (J,J'),

and 72 hpf (K,K') zebrafish embryos co-immunostained with antibodies that detect Rbpms2

(magenta) or cardiac Troponin T (CT3 antibody; green). Single (H,I,J,K) and merged double

channel (H',I',J',K') images are shown. Little to no variation in expression pattern was observed

between animals in each group (n>10/group). (L-L''') Single optical section through the

ventricular wall of a 48 hpf embryo co-immunostained with antibodies that detect Rbpms2

(magenta) or cardiac Troponin T (CT3 antibody; green) and counterstained with DAPI (blue).

Single (L, L'), merged double (L''), and merged triple (L''') channel images are shown. Yellow

arrowheads highlight Rbpms2-positive cardiomyocyte nuclei surrounded by Troponin T-positive

cytoplasm. White arrowheads highlight Rbpms2-negative nuclei in adjacent endocardial cells

lining the ventricular chamber. 4/4 ventricles exhibited this Rbpms2 expression pattern. (M,M')

Confocal projections of a histological section through the ventricle of an adult zebrafish heart co-

immunostained with antibodies that detect Rbpms2 (magenta) or Myosin Heavy Chain (MHC;

MF20 antibody; green). Single channel (M) and merged double channel (M') images are shown.

18/18 sections, 6 from each of three hearts, exhibited this Rbpms2 expression pattern.

Abbreviations: FACS, fluorescence activated cell sorting, DE, differential expression; V,

ventricle; A, atrium; OFT, outflow tract. Scale bars=25μm.

Figure 2 – *rbpms2a* and *rbpms2b* are required for ventricular function. (A,B) Schematic
diagrams of Rbpms2a and Rbpms2b (top) and the predicted protein products of the *rbpms2a^chb3*
and *rbpms2b^chb4* alleles (bottom). The asterisks mark the locations of premature stop codons (*)
within the RNA-recognition motifs (RRMs) caused by frame-shifting deletions. The white box in
(A) shows the location of divergent amino acids prior to the stop codon. (C-D) Brightfield images
of control-sibling (C; CTRL) and *rbpms2*-null (*rbpms2a^chb3/chb3*; *rbpms2b^chb4/chb4*; D) embryos at
72 hours post fertilization (hpf). Arrowhead highlights pericardial edema in the mutant. Little to
no variation in phenotype was observed between animals in each group (n>10/group). Scale
bars=250μm. (E) Western blot showing Rbpms2 levels in 48 hpf wild-type (WT), CTRL, and
rbpms2-null embryos (top) relative to the loading control alpha-tubulin (bottom). (F-G) Confocal
projections of hearts in 48 hpf CTRL (F) and *rbpms2*-null (G) embryos immunostained with an
antibody that detects Rbpms2 (magenta) and counterstained with DAPI (blue). The dotted lines
delineate the myocardium as ascertained from the Rbpms2 signal (F) or *Tg(myl7:eGFP)*
reporter signal present in these animals (F,G; not shown). Little to no variation in expression
pattern was observed between animals in each group (n>10/group). (H) Kaplan Meier plot

showing the survival curves of CTRL (n=33) and *rbpms2*-null (n=33) animals. Statistical

significance was determined using a log-rank test. (I-L) Confocal projections of hearts in 48 hpf

CTRL (I,K) or *rbpms2*-null (J,L) embryos carrying the *myl7:GFP* (I,J) or *myl7:nucGFP* (K,L)

transgenes immunostained with an anti-GFP antibody. For (I,J), little to no variation in

phenotype was observed between animals in each group (n>10/group). (M) Dot plot showing

ventricular cardiomyocyte (CM) number in CTRL (n=6) and *rbpms2*-null (n=6) hearts at 48 hpf.

Error bars show one standard deviation. Statistical significance was determined by an unpaired,

two-tailed Student's t-test assuming equal variances. ns, not significant. (N-Q) Representative

single-plane light sheet fluorescence microscopy images of hearts in 48 hpf CTRL (N,O) and

rbpms2-null (P,Q) embryos carrying the *myl7:GFP* transgene. Images of hearts during

ventricular diastole (N,P) or systole (O,Q) are shown with Cardiac Functional Imaging Network

(CFIN) overlays of the atrial (blue) and ventricular (cyan) lumens. (R) Dot plots showing end-

diastolic volume (EDVs) and end-systolic volume (ESVs), stroke volume, ejection fraction, heart

rate, and cardiac output of ventricles in 48 hpf CTRL (n=6) and *rbpms2*-null (n=6) animals

measured with CFIN. Error bars show one standard deviation. Statistical significance was

determined by an unpaired, two-tailed Student's t-test assuming equal variances. ns, not

significant. Abbreviations: aa, amino acid; V, ventricle; A, atrium. Scale bars=25μm.

Figure 3 – Identification of differential alternative splicing events in *rbpms2*-null embryos.

(A) Schematic diagrams of the categories of alternative splicing events (ASEs) investigated by the RNA-seq Multivariate Analysis of Transcript Splicing (rMATS) algorithm including skipped exon (SE), alternative 3' splice site (A3SS), alternative 5' splice site (A5SS), mutually exclusive exon (MXE), and retained intron (RI). (B) Volcano plot showing the inclusion (Inc) level differences (control-sibling Inclevel - *rbpms2*-null Inclevel) and False Discovery Rates (FDR) for differential ASEs between 33 hours post fertilization (hpf) control-sibling and *rbpms2*-null animals (0 uncalled replicates; FDR<0.01; |Incleveldifference|>0.1;). The ASEs shown in bold were validated with splicing-sensitive qPCR and in situ hybridization. (C) Pie chart showing the proportions of differential ASEs in each category in *rbpms2*-null animals.

Figure 4 – Validation of differential alternative splicing events in *rbpms2*-null animals. (A) Schematic diagrams of alternative splicing events (ASEs) identified in control-sibling (CTRL) or *rbpms2*-null animals at 33 hours post fertilization (hpf) by the RNA-seq Multivariate Analysis of Transcript Splicing (rMATS) or mixture of isoforms (MISO) algorithms. The legend indicates that the isoforms shown on top (black) are significantly reduced or absent in *rbpms2*-null animals and replaced by the isoforms shown on the bottom (gray). (B) Bar graph showing the relative expression levels of the four genes shown in (A), independent of the alternatively spliced exons, between 48 hours post fertilization control-sibling and DKO animals. (C) Bar graph showing the relative expression levels of the indicated, exon-containing isoforms [(shown in black in (A)] of the same four genes between control-sibling and *rbpms2*-null animals at 48 hpf. For (B,C), error bars show one standard deviation. Statistical significance was determined by an unpaired, two-

tailed Student's t-test assuming equal variances. ns, not significant. (D-K') Brightfield images of

33 hpf CTRL (D,D',F,F',H,H'J,J') and *rbpms2*-null (E,E',G,G',I,I',K,K') embryos processed for in

situ hybridization with riboprobes that detect total pools (D-E'; H-I') or specific exon-containing

mRNA isoforms (F-G'; J-K') of *mybpc3* (D-G') or *pln* (H-K'). The cardiac regions in

(D,E,F,G,H,I,J,K) are shown at higher zoom in (D',E',F',G',H',I',J',K'). Red arrowheads in (D,E)

highlight expression of *mybpc3* in skeletal muscle, which is shown from a dorsal view in the

insets. Black arrowheads in (D-K) indicate the linear heart tube. Little to no variation in

expression pattern was observed between animals in each group (n>10/group). Scale

bars=100μm.

Figure 5 – Myofibrillar disarray in *rbpms2*-null ventricular cardiomyocytes. (A-B')

Confocal projections of ventricles in 48 hours post fertilization (hpf) control-sibling (CTRL) or

rbpms2-null embryos immunostained with an anti-TroponinT antibody (CT3) to visualize thin

filaments. Regions demarcated in (A) and (B) are enlarged in (A') and (B'). (C-D'') Confocal

projections of ventricles in 48 hpf CTRL and *rbpms2*-null *Tg(α-actn3b-EGFP)* animals co-

immunostained with antibodies that detect Tropomyosin (CH1; green) to visualize thin filaments

or GFP (magenta) to visualize Z-disks. (E-F) Confocal projections of ventricles in 48 hpf CTRL

and *rbpms2*-null embryos immunostained with an antibody that detects myosin heavy chain

(MF20) to visualize thick filaments. Little to no variation in protein distributions were observed

between animals in each group (n>10 animals/group). Scale bars=10μm.

Figure 6 – *rbpms2*-null ventricular cardiomyocytes exhibit reductions in Ca2+-transient

duration and amplitude. Color maps (A,B,E,F) and dot plots (C,D,G,H; n=10/group) of Ca2+-

transient duration (A-D) and Ca2+ release amplitude (E-F) in the ventricle (A-C; E-G) and atrium

(A,B,D,E,F,H) of 48 hours post fertilization control-sibling (CTRL; A,C,D,E,G,H) and *rbpms2*-null

(B-D; F-H) hearts. Color code depicts localized Ca2+-transient duration in milliseconds (ms;

A,B) or the ratiometric fluorescence indicator of Ca2+-release amplitude. Each dot represents

center-chamber measurements from one heart. Error bars show one standard deviation.

Statistical significance was determined by an unpaired, two-tailed Student's t-test assuming

equal variances. ns, not significant.

Figure 7 – Human *RBPMS2*-null cardiomyocytes exhibit myofibrillar disarray and Ca2+

handling defects similar to those in zebrafish *rbpms2*-null cardiomyocytes. (A,B)

Schematic diagrams of human RBPMS2 (top) and the predicted protein product of the

ΔRBPMS2 null allele created with CRISPR/Cas9-mediated genome editing (bottom). The

asterisk shows the location of a premature stop codon within the RNA-recognition motif (RRM)

caused by a frame-shifting 2 base pair deletion. The white box shows the location of divergent

amino acids prior to the stop codon. (B,C) Bar graphs showing the relative expression levels of

RBPMS2 in parental control (CTRL) and *ΔRBPMS2* human induced pluripotent stem cells

(hiPSCs; B) and cardiomyocytes (hiPSC-CMs; C) after 15 days of directed differentiation. Error

bars show one standard deviation. Statistical significance was determined by an unpaired, two-

tailed Student's t-test assuming equal variances. (D-K) Representative confocal projections of

CTRL (D-G) and *ΔRBPMS2* (H-K) hiPSC-CMs immunostained with antibodies that detect

TroponinT (CT3; green) or Alpha Actinin (anti-Sarcomeric Alpha Actinin; magenta) to visualize

thin filaments or Z-disks, respectively, and counterstained with DAPI (blue). Single (D-F; H-J)

and merged triple (G,K) channel images are shown. Little to no variation in staining pattern was

observed between cells in each experimental group (n>50 cells/group from at least three wells

from two separate differentiations were examined). (L,M) Representative traces of fluorescence

intensity over baseline (F/F$_o$; top) derived from line scans (bottom) of CTRL (L) and *ΔRBPMS2*

(M) hiPSC-CMs loaded with fluo-4 and paced at 1Hz. (N-S) Violin plots showing Ca2+-transient

amplitude (N), duration at 80% (O) and 50% (P) repolarization, time to peak amplitude (Q),

upstroke velocity (R), and un-paced spontaneous beating rate (S). Statistical significance was

determined by an unpaired, two-tailed Student's t-test assuming equal variances. n=80

cells/group, 40 each from two separate differentiations. Scale bars=25μm.