High Transcriptional Activity and Diverse Functional Repertoires of Hundreds of Giant Viruses in a Coastal Marine System

Anh D Ha, Mohammad Moniruzzaman, Frank O Aylward*

Department of Biological Sciences, Virginia Tech

926 West Campus Drive. Blacksburg, VA, 24061

*Correspondence: faylward@vt.edu

Running Title: Transcriptional Activity of Marine Giant Viruses

Keywords: Giant Viruses, NCLDV, *Nucleocytoviricota*, Viral Diversity, Kinesin, Myosin

Abstract

Viruses belonging to the *Nucleocytoviricota* phylum are globally distributed and include members with notably large genomes with complex functional repertoires. Recent studies have shown that these viruses are particularly diverse and abundant in marine systems, but the magnitude of actively replicating *Nucleocytoviricota* present in ocean habitats remains unclear. In this study, we compiled a curated database of 2,431 *Nucleocytoviricota* genomes and used it to examine the gene expression of these viruses in a 2.5-day metatranscriptomic time-series from surface waters of the California Current. We identified 145 viral genomes with high levels of gene expression, including 90 *Mimiviridae* and 53 *Phycodnaviridae*. In addition to recovering high expression of core genes involved in information processing that are commonly expressed during viral infection, we also identified transcripts of diverse viral metabolic genes from

pathways such as glycolysis, the TCA cycle, and the pentose phosphate pathway, suggesting that

virus-mediated reprogramming of central carbon metabolism is common in oceanic surface

waters. Surprisingly, we also identified viral transcripts with homology to actin, myosin, and

kinesin domains, suggesting that viruses may use them to manipulate host cytoskeletal dynamics

during infection. We performed phylogenetic analysis on the virus-encoded myosin and kinesin

proteins, which demonstrated that most belong to deep-branching viral clades, but that others

appear to have been acquired from eukaryotes more recently. Our results highlight a remarkable

diversity of active *Nucleocytoviricota* in a coastal marine system and underscore the complex

functional repertoires expressed by these viruses during infection.

Importance

The discovery of giant viruses has transformed our understanding of viral complexity. Although

viruses have traditionally been viewed as filterable infectious agents that lack metabolism, giant

viruses can reach sizes rivalling cellular lineages and possess genomes encoding central

metabolic processes. Recent studies have shown that giant viruses are widespread in aquatic

systems, but the activity of these viruses and the extent to which they reprogram host physiology

in situ remains unclear. Here we show that numerous giant viruses consistently express central

metabolic enzymes in a coastal marine system, including components of glycolysis, the TCA

cycle, and other pathways involved in nutrient homeostasis. Moreover, we found expression of

several viral-encoded actin, myosin, and kinesin genes, indicating viral manipulation of host

cytoskeleton during infection. Our study reveals a high activity of giant viruses in a coastal

marine system and indicates they are a diverse and underappreciated component of microbial

diversity in the ocean.

Introduction

Large dsDNA viruses of the family *Nucleocytoviricota*, commonly referred to as "giant viruses",

are a diverse group of double-stranded DNA viruses with virion sizes reaching 1.5 µm,

comparable to the sizes of many cellular lineages (1–4). These viruses are known to infect a

broad range of eukaryotic hosts; members of the *Poxviridae* and *Iridoviridae* families infect

numerous metazoans, the *Mimiviridae* and *Phycodnaviridae* are known to infect a wide range of

algae and other protists, and the *Asfarviridae* infect a mixture of metazoan and protist hosts (5–

7). Members of the *Nucleocytoviricota* typically harbor exceptionally large genomes that are

often > 300 kbp in length, and in some cases as large as 2.5 Mbp (8). Numerous studies have

noted the unusually complex genomic repertoires of viruses in this group that include many

genes typically found only in cellular lineages, such as those involved in the TCA cycle (9, 10),

glycolysis (10), amino acid metabolism (11), light sensing (12, 13), sphingolipid biosynthesis

(14, 15), eukaryotic cytoskeleton (16), and fermentation (16). These complex metabolic

repertoires are thought to play a role in the manipulation of host physiology during infection, in

effect transforming healthy cells into reprogrammed "virocells" that more efficiently produce

viral progeny (17, 18).

The genomic complexity of *Nucleocytoviricota* together with reports that they are abundant in

global marine environments has raised questions regarding the extent to which they manipulate

the physiology of their hosts and thereby influence global biogeochemical cycles (19). Cultivated

representatives of these viruses are known to infect a broad array of ecologically-important

eukaryotic algae, including dinoflagellates, haptophytes, brown algae, and chlorophytes, among

others (20–23), suggesting they could play important roles in shaping the activity and

composition of the microbial eukaryotic community. Even so, our understanding of the

distribution and activity of *Nucleocytoviricota* in marine systems has lagged behind that of other

viral groups owing to their large size, which precludes their presence in the small size fractions

typically analyzed in viral diversity studies (24). Some early studies were prescient in suggesting

that these viruses, and in particular those within the families *Mimiviridae* and *Phycodnaviridae*,

are a diverse and underappreciated component of viral diversity in the ocean (25–28). More

recent work has confirmed this view and revealed that a diverse range of *Nucleocytoviricota*

inhabit the global ocean and likely contribute to key processes such as algal bloom termination

and carbon export (10, 29–34). Some metatranscriptomic studies have also begun to note the

presence of *Nucleocytoviricota* transcripts in marine samples, confirming their activity (35–37).

Although these studies have vastly expanded our knowledge of the diversity and host range of

Nucleocytoviricota in the ocean, the activity of these viruses and the extent to which they use

cellular metabolic genes they encode during infection remains unclear.

In this study, we constructed a database of 2,436 annotated *Nucleocytoviricota* genomes for the

purpose of evaluating the gene expression landscape of these viruses. We then leveraged a

previously published metatranscriptome time-series from surface waters of the California

Current (37) to assess the daily activity of *Nucleocytoviricota* in this environment. We show that

hundreds of these viruses, primarily of the *Mimiviridae* and *Phycodnaviridae* families, were

consistently active during this period and frequently expressed genes involved in central carbon

metabolism, light harvesting, oxidative stress reduction, and lipid metabolism. Unexpectedly, we

also found expression of several viral-encoded cytoskeleton genes, including those that encode

the motor proteins myosin and kinesin, and we performed a phylogenetic analysis demonstrating

that *Nucleocytoviricota* commonly encode deep-branching enzymes in these protein families.

Our findings highlight the surprisingly high activity of *Nucleocytoviricota* in marine systems and

suggest they are an underappreciated component of viral diversity in the ocean.

Results and Discussion

We examined a metatranscriptomic dataset of 16 timepoints sampled over a 60-hour period that

were obtained from microbial communities previously reported for the California current system

(37). This dataset is ideal for examining the activity of marine *Nucleocytoviricota* because it

targeted the >5 µm size fraction, which is enriched in many of the eukaryotic plankton that these

viruses are known to infect. Altogether, we identified 145 *Nucleocytoviricota* genomes with

metatranscriptomic reads mapping over the sampling period, including 90 *Mimiviridae*, 53

Phycodnaviridae, 1 *Iridoviridae*, and 1 *Asfarviridae* (Fig 1). Expression level of the 145

genomes with mapping are summarized in Supplemental Data S1.

Of the 145 genomes detected, 11 are complete genomes of cultivated viruses, while 134 were

generated using cultivation-independent methods, emphasizing the importance of metagenome-

assembled viral genomes for expanding the number of available reference genomes. Almost half

(66) of the 145 *Nucleocytoviricota* genomes have genome or assembly sizes > 300 kbp (Fig. S1),

indicating that a large fraction of the active viruses have large genomes with potentially complex

functional repertoires. The abundance of each *Nucleocytoviricota* genome in the transcriptomes

was positively correlated to the number of genes that could be recovered for that virus (Fig. 1a),

which is expected given that only highly-expressed viral genes will be recovered for low-

abundance viruses in the dataset. Over 80% of the predicted genes in five genomes were

recovered throughout the sampling period (one *Mimiviridae* and four *Phycodnaviridae*),

indicating that the recovery of most transcripts is possible for some abundant viruses (Fig 1).

We constructed a multi-locus phylogenetic tree of the 145 viruses present in the transcriptomes

together with 1,458 references so that we could examine the phylogenetic distribution of the

active *Nucleocytoviricota* (Fig 2). The topology of the resulting tree is consistent with a previous

tree we constructed using similar methods (10). Given the large diversity of *Mimiviridae* and

Phycodnaviridae in the dataset, we grouped viruses in these families into sub-clades that we

have previously established (10). Consistent with our previous findings, we also refer to two

groups of *Phycodnaviridae* as early- vs late-branching because they form distinct monophyletic

groups in the tree. Of the 145 viruses that were abundant in the metatranscriptomes, *Mimiviridae*

Clade 1 (MC_01) and Late *Phycodnaviridae* Clade 1 (LP_01) were by far the most well-

represented (55 and 47 genomes, respectively, Fig 1B); interestingly, both of these clades also

include cultivated representatives that are known to infect marine protists, which suggests

possible hosts for the viruses in the transcriptomes; MC_01 contains haptophyte viruses that

infect the genera *Phaeocystis* and *Chrysochromulina* (38–40), while LP_01 contains

Prasinoviruses that are known to infect *Ostreococcus, Bathycoccus*, and *Micromonas* (41).

Transcripts mapping to isolate *Ostreococcus* viruses of LP_01 was previously reported in the

original analysis of these metatranscriptomes (37); in addition to recovering these genomes, we

also identified transcripts mapping to an additional 39 metagenome-derived *Nucleocytoviricota*

genomes in clade LP_01, highlighting the diversity of viruses within this clade that were active

in the same community during the sampling period. A wide range of other Mimivirus clades

were also present in addition to MC_01 (Fig 1c): MC_16, which is the canonical Mimivirus

clade and includes the original *Acanthamoeba polyphaga* Mimivirus and its close relatives,

MC_12 containing the recently-cultivated Tetraselmis virus (TetV) (16), MC_08, which includes

the recently-reported Choanoflagellate virus (42), MC_09, which includes *Aureococcus*

anophagefferens virus (AaV) (23), and Mimivirus clades 5, 7, and 13 containing no cultivated

representatives. One Iridovirus (GVMAG-M-3300009467-15) and one Asfarvirus (GVMAG-M-

3300027833-19) were also abundant in the transcriptomes, suggesting they may also infect

microbial eukaryotes in the community. The Asfarvirus GVMAG-M-3300027833-19 was also

recently identified in waters off the coast of South Africa (7), indicating that it may be broadly

distributed in the ocean. The presence of numerous divergent clades of *Nucleocytoviricota* in the

metatranscriptomes supports the view that marine systems harbor an immense diversity of these

viruses that are actively infecting a wide variety of protists hosts.

We examined summed whole-genome transcriptional activity of the 145 *Nucleocytoviricota* to

identify potential diel cycles of viral activity. We found 6 viruses that had significantly higher

expression during nighttime periods (Fig 3A, Mann-Whitney U test, corrected p-value < 0.1).

These transcriptional patterns were not strictly diel as their peak of expression was found at

different times during the night, but they nonetheless suggest that viral activity may be higher at

night for some of these viruses. We also analyzed whole-genome transcriptional activity using a

weighted network-based approach that groups viruses with similar temporal transcriptional

profiles into modules. This analysis revealed two primary modules, one of which contained 60

viruses and had a module-wide transcriptional profile with notable peaks at nighttime periods

(Module 1, Fig 3B). Previous work found diel cycling in marine bacteriophage that typically

peaked near dusk, with overall viral transcripts also increasing at night (43). Moreover, a culture-

based study of *Ostreococcus tauri* virus 5 found that viral transcripts increased markedly at night

(44). Two of the *Nucleocytoviricota* with higher nighttime expression in our analysis belong to

clade LP_01 together with cultivated Prasinoviruses that infect *Ostreococcus* and other

prasinophytes, suggesting that higher nighttime activity may be a common trait in this viral

clade. We found no significant diel cycles in the *Nucleocytoviricota* whole genome

transcriptional profiles when we tested for strict diel periodicity (RAIN, p-value < 0.1, see

Methods), suggesting that, although viral activity appears higher at night for some viruses,

pronounced and significant diel cycling of viral activity was not detectable. This does not

necessarily suggest that diel cycles are uncommon in marine *Nucleocytoviricota*; previous

studies have found that the detection of diel cycles in metatranscriptomic data can require longer

time-series (>20 time-points) and may only be feasible for the most abundant community

members for which gene expression is high (45). More work is therefore needed to examine the

potential role of diel cycling in these viruses in more detail.

Throughout the sampling period, we detected expression of numerous viral genes involved in

diverse metabolic processes (Fig 4, Fig S2, Supplemental Data S2). As expected, we found high

expression of viral core genes, including the major capsid protein, family B DNA Polymerase,

A32 packaging enzyme, VLTF3 transcriptional factor, and superfamily II helicase (Fig. 4); this

is consistent with both culture-based and culture-independent studies of *Nucleocytoviricota* that

have found most of these genes, and in particular the capsid proteins, to be highly expressed (44,

46–49). Chaperones from the Hsp70 and Hsp90 families were also highly expressed in numerous

Nucleocytoviricota, consistent with some previous transcriptome studies (47, 49); these likely

play a role in protein folding in virus factories, particularly for capsids (3, 49). We also detected

transcripts involved in mitigating cellular stress, such as glutathione-S-transferase (GST) and

superoxide dismutase (SOD). SOD is involved in the detoxification of reactive oxygen species

produced by hosts as a defense mechanism during infection, and it has been postulated that viral-

encoded SODs may allow some viruses to infect a broader array of hosts (50). GST may be

involved in the detoxification of electrophilic metabolites, and the host version of this gene was

also found to be overexpressed in several time-points of an infection experiment of *Aureococcus*

anophagefferens virus and its host (51).

We also detected numerous transporters predicted to target nitrogen, sulfur, and phosphorus-

containing nutrients, consistent with the view that *Nucleocytoviricota* manipulate the nutrient

environment of their hosts during infection. Among these, we identified expression of several

viral ammonium transporters; previous work with an ammonium transporter encoded by

Ostreococcus tauri virus 6 has shown that this gene is expressed during infection and

manipulates nutrient uptake by the host cell (52). Viral-encoded phosphate transporters are also

common in many marine *Nucleocytoviricota* (10, 53), and expression of these genes further

emphasizes their likely role in viral-mediated nutrient transport during infection. Other genes

involved in nutrient homeostasis were also well-represented in the metatranscriptomes, including

ferritin, glutaminase, glutamine synthetase, and Pho regulon components (Fig. 4), consistent with

recent findings that these genes are also common in these viruses (10).

Recent work has noted that many *Nucleocytoviricota* encode enzymes involved in central

metabolic pathways, and it has been postulated that these have a role in the reprogramming of

healthy cells into virocells during infection (10). We detected expression of many of these

central metabolic enzymes throughout the sampling period; genes involved in glycolysis, the

TCA cycle, the pentose phosphate pathway, and beta oxidation were all present in the

metatranscriptomes, and most were consistently expressed across timepoints, especially in

Mimiviridae clades (Fig 4, Fig S2). A recent study of TCA cycle enzymes encoded by a

Pandoravirus isolate has suggested that these enzymes have activity consistent with

bioinformatic predictions, and that some are packaged into virions and would therefore influence

host physiology soon after infection (54). Together with our results, this suggests that viral

manipulation of central metabolic processes is a widespread mechanism employed by many

Nucleocytoviricota, and that this is a common occurrence in viral infections in the ocean. This

raises the question of what proportion of the protist community is infected by *Nucleocytoviricota*

at any given time, and thereby has an altered metabolic state. Previous studies examining

infected hosts have found that 27-37% of cells can be infected by *Nucleocytovicota* during an

algal bloom (55, 56); given the possibility that multiple viruses may infect the same host

population, the total number of infected cells may be substantially higher. This leads to the

surprising possibility that the majority of protist populations in some marine systems may consist

of mostly "virocells" in which their metabolism is substantially altered by viruses that infect

them, at least some periods of high viral activity.

Surprisingly, we also found several transcripts matching to predicted cytoskeletal proteins,

including actin, myosin, and kinesin. One recent study identified several actin homologs in

Nucleocytoviricota ("viractins") and suggested that they played a role in the emergence of this

protein in extant eukaryotic lineages (57), but the identification of myosin and kinesin-like

proteins has not previously been reported. We found three expressed viral genes with predicted

myosin motor domains and one with kinesin domains, and subsequent analysis of all

Nucleocytoviricota genomes in our reference database recovered an additional 109 myosin and

200 kinesin homologs. To examine if these genes were recently acquired from cellular hosts, we

performed a phylogenetic analysis of the viral myosin and kinesin proteins with references; our

results indicate that the viral proteins typically form separate branches that are distinct from their

eukaryotic homologs, except in a few cases where it appears some *Nucleocytoviricota* have

acquired cellular copies more recently (Fig 5). This is consistent with recent studies that have

revealed a dynamic gene exchange between *Nucleocytoviricota* and their hosts that in some cases

dates back to the early diversification of eukaryotes (13, 58–60). The expression of viral genes

with homology to actin, myosin, and kinesin domains suggests that these viruses may use these

proteins to manipulate host cytoskeletal dynamics during infection; it is possible that these

proteins play a role in the establishment and maintenance of perinuclear viral factories, which are

responsible for viral morphogenesis, or for the subcellular trafficking of viral particles. In

Aureococcus anophagefferens cytoskeletal genes are downregulated during infection by the

Mimivirus AaV (51); if this is common across other host-virus systems it would suggests that

Nucleocytoviricota-encoded myosin, kinesin, and actin proteins may help to maintain the

functioning of the cytoskeleton during later periods of infection.

To potentially link viruses to their hosts, we performed a network-based analysis based on

correlations of whole-genome transcription of both viruses and eukaryotic plankton (Fig 6). This

was motivated by several studies that have noted that the transcriptional activity of many viruses

is often correlated to that of their host *in situ* (37, 46, 61). The resulting network links

Nucleocytoviricota to diverse potential hosts, including Dinophyta, Ciliophora, Stramenopiles,

and Pelagophyta. Many members of the Late *Phycodnaviridae* clustered in the same part of the

network as *Ostreococcus*, consistent with the prediction that most members of clade LP_01 are

Prasinoviruses. Several *Mimiviridae* in clade MC_01 were correlated with members of the

diatom genera *Leptocylindrus*, *Chaeotoceros*, and *Eucampia*, consistent with a recent co-

occurrence analysis from Tara Oceans data which suggested that some *Nucleocytoviricota* are

potentially associated with diatoms (62); this association remains speculative pending more

detailed analysis, however. In general, these results support the view that *Nucleocytoviricota*

likely infect diverse hosts in the environment, but specific host predictions should be treated with

caution pending validation with experimental approaches.

Conclusion

In this study, we identified 145 *Nucleocytoviricota* with high transcriptional activity over a 60-

hour period in surface waters of a coastal marine system. We analyzed only genomes for which >

10% of the predicted genes could be recovered at some point during the time-series; this

provides high confidence that the detected viruses were indeed present, but it also implies that

the actual number of active viruses is far higher than 145 because many viral transcripts remain

below the level of detection. This therefore further highlights the remarkable diversity of active

Nucleocytoviricota in this coastal system and implies more generally that these viruses are an

important component of global marine environments. In addition, we found expression of diverse

viral-encoded functional genes from central metabolic pathways as well as genes with similarity

to eukaryotic cytoskeletal components, emphasizing the complex mechanisms employed by giant

viruses to manipulate their hosts in the environment. One implication of these findings is that

these viruses also have a potentially important role in shaping marine food webs; the relative

impact of top-down vs bottom up controls in marine systems has long been debated (63), and the

ability of *Nucleocytoviricota* to infect both heterotrophic protists (i.e., grazers) and phototrophic

plankton (i.e. primary producers) suggests they can influence the relative impact of these forces.

Future work identifying the hosts of these viruses and their relative contribution to eukaryotic

plankton mortality rates will therefore be important avenues of future work. Overall, our findings

suggest that *Nucleocytoviricota* are an underappreciated component of marine systems that exert

an important influence on community dynamics and biogeochemical cycling.

Methods

Compilation of a *Nucleocytoviricota* Genome Database for Read Mapping

We compiled a set of *Nucleocytoviricota* genomes for transcript mapping that included

metagenome-assembled genomes (MAGs) as well as genomes of cultured isolates. For this we

first downloaded 2,074 genomes from Schulz et al. (60) and 501 genomes from Moniruzzaman

et al (10), two recent studies which generated numerous *Nucleocytoviricota* MAGs. We also

included all *Nucleocytoviricota* genomes available in NCBI RefSeq as of June 1st, 2020. Lastly,

we also included several *Nucleocytoviricota* genomes from select publications that were not yet

available in NCBI, such as the cPacV, ChoanoV, and AbALV viruses that have recently been

described (42, 47, 64).

After compiling this set, we dereplicated the genomes, since the presence of highly similar or

identical genomes can obfuscate the results of read mapping. For dereplication we compared all

Nucleocytoviricota genomes against each other using MASH v. 2.0 (65) ("mash dist" parameters

-k 16 and -s 300), and clustered genomes together using a single-linkage clustering, with all

genomes with a MASH distance of ≤ 0.05 linked together. The MASH distance of 0.05 was

chosen since it has been found that it corresponds to an average nucleotide identity (ANI) of 95%

(65), which has furthermore been suggested to be a useful threshold for distinguishing viral

species-like clusters (66). From each cluster, we chose the genome with the highest N50 contig

length as the representative. Prior to read mapping we also decontaminated the genomes through

analysis with ViralRecall v.2.o (67) (-c parameter), with all contigs with negative scores

removed. The final database used for read mapping contained 2,431 viral genomes.

Our read mapping strategy employed a translated homology search (DNA searched against

protein), and so we predicted proteins from all genomes using Prodigal 2.6.3 (68) (run

independently on each genome using default parameters), and then masked the protein database

using tantan v. 22 (69) (default parameters) to prevent spurious read mapping to low complexity

sequences. We then formatted the database for read mapping using the lastdb utility in LAST v.

1060 (70).

Metatranscriptome Dataset

We examined the metatranscriptomic dataset of 16 timepoints obtained from >5 µm size fraction

microbial communities previously reported for the California current system (37). In this study

community RNA samples were collected every four hours over a 60-hour time frame, starting 6

AM September 16 to 6 PM September 18. We downloaded and trimmed reads from each of the

16 metatranscriptome libraries with Trim Galore v. 0.6.4 with parameters "--length 50 -q 5 --

stringency 1", and after removing singleton reads we then mapped paired reads onto the

Nucleocytoviricota database using LASTAL v. 1060 (70) (parameters -m 10 -u 2 -Q 1 -F 15).

We only considered hits with percent identity > 95% and bit score > 50. Transcript mapping

workflows similar to this have been successfully employed in other studies (43, 45). For relative

abundance comparisons read mapping counts were converted to Transcripts per Kilobase per

Million (TPM) (71). To avoid the spurious detection of viral genomes in the metatranscriptomes

we only considered genomes for which >= 10% of the proteins in that genome had hits; this

cutoff was guided by recent work which has suggested that reads mapping to > 10% of a genome

are unlikely to be due to spurious mappings, and therefore are indicative of the presence of that

virus in a given sample (72). After performing this filtering, we arrived at a set of 145

Nucleocytoviricota reference genomes that were used in subsequent analysis.

Protein Annotations

We annotated proteins in our *Nucleocytoviricota* database by comparing them to the EggNOG

4.5 (73) and Pfam v. 31 (74) Hidden Markov Models (HMMs) using HMMER3 3.3 (parameter

"-E 1e-3" for the NOG search and "--cut_nc" for the Pfam search). In both cases only best hits

were retained. These annotations are available in Supplemental Data S2. Annotations were

manually reviewed to examine functions of interest, including energy metabolism, carbohydrate

metabolism, cytoskeleton, membrane transport, light harvesting, and oxidative stress regulation.

Hallmark *Nucleocytoviricota* proteins were identified using HMMs previously described (10).

Nucleocytoviricota **Phylogeny and Clade Delineation**

To provide phylogenetic context for the 145 genomes we identified in the metatranscriptomes,

and to delineate broader clades, we constructed a large mutli-locus phylogenetic tree of

Nucleocytoviricota using the marker genes DNA polymerase B (PolB), A32 packaging enzyme

(A32), Major Capsid Protein (MCP), Superfamily II helicase (SFII), and the VLTF3

transcription factor (VLTF3). Previous analyses have found that these marker genes are

widespread in *Nucleocytoviricota* and useful for phylogenetic analysis (10, 75, 76). For this tree

we used all 145 genomes present in the metatranscriptomes as well as select reference

Nucleocytoviricota sampled from across all of the main families (*Asfarviridae, Poxviridae,*

Pithoviridae, Marseilleviridae, Iridoviridae, Early and Late-branching *Phycodnaviridae*, and

Mimiviridae). We only included reference genomes if they contained four of five of the marker

genes we used, to ensure accurate phylogenetic estimation. The concatenated phylogeny was

generated with the ncldv_markersearch.py script (github.com/faylward/ncldv_markersearch), the

alignment was trimmed with TrimAl v. 1.4.rev22 (77) (parameter -gt 0.1), and the tree was

inferred using IQ-TREE (78). To infer the best model, we used the ModelFinder utility in IQ-

TREE (79), which selected the VT+F+I+G4 model. Support values were inferred using 1000

ultrafast bootstraps (80). We previously delineated family-level and subfamily-level clades for

the *Nucleocytoviricota* (10), and we used this nomenclature to do the same for this tree.

Virus-Host Transcriptional Network

To identify potential marine host-virus relationships, we computed a Pearson correlation matrix

between the total abundance of all genomes with gene expression across the 16 timepoints

(function cor() in R). The total abundance of genomes was calculated as the summed abundance

of all genes of the genome expressed. The abundance of individual genes was calculated by

normalizing the number of raw reads by the total number of raw reads per million in each

sample. Only correlations coefficients between genome pairs exceeding 0.8 were reported.

Tests for Diel Cycling and Day/Night Overexpression

To explore the transcriptional activities and possible diel expression pattern of the

Nucleocytoviricota community, we examined the TPM-normalized transcriptional profiles of

individual *Nucleocytoviricota* genes using RAIN (81). We did the same for each of the 145

Nucleocytoviricota genomes present in the transcriptomes by summing TPM expression values

across each genome. P-values for these tests were corrected using the Benjamini and Hochberg

method (82) as implemented by the p.adjust command in R.

In addition to strict tests for diel periodicity, we also sought to examine potential overexpression

of *Nucleocytoviricota* in daytime vs nighttime periods. For this we performed a Mann-Whitney

U-test on the summed gene expression of each virus, using timepoints of 0600, 1000, and 1400

as daytime and timepoints of 1800, 2200, and 0200 as nighttime (wilcox.test in R). For these

tests, we also corrected the p-values using the same method as the RAIN tests.

To find clusters (modules) of genomes with highly correlated temporal expression, we performed

the Weighted Gene Co-expression Network Analysis method (83) on the TPM-normalized

transcriptional profiles of the *Nucleocytoviricota* genome-wide transcriptional profiles. We used

a soft thresholding power of 14, parameters TOMType ="signed", reassignThreshold = 0,

minModuleSize=10 and mergeCutHeight = 0.25. We then used the moduleEigengenes utility in

the WGCNA package to summarize the genome module's average expression profile (module

eigengene).

Phylogeny of Myosin and Kinesin Proteins

To examine the evolutionary history of the *Nucleocytoviricota* proteins with myosin and kinesin

domains we identified all occurrences of these proteins in our viral genome database, including

those proteins that were not expressed in the metatranscriptomes. For this we only considered

proteins with matches to the PF00063 (myosin) and PF00225 (kinesin) (only scores greater than

the noise cutoff considered). For references we downloaded all the kinesin (COG5059) and

myosin (COG5022) homologs from the EggNOG database v4.5. Many of the species harbor

multiple of these proteins, and for the purpose of broad phylogenetic analysis, we randomly

selected one copy of these genes per species. Diagnostic phylogenetic trees were constructed

using FastTree v. 2 (84) implemented in ete3 toolkit (85) for the purpose of removing long

branches and redundant or small sequences (< 300 amino acids long). To construct the final

trees, we used ClustalOmega to align the sequences and trimAl (parameter -gt 0.1) to trim the

alignments. We used IQ-TREE to build maximum likelihood phylogenies with the model

LG+I+G4 and assessed the node support with 1000 ultrafast bootstrap replicates.

Data Availability

The datasets analyzed in this study are already publicly available and were accessed as described

in the Methods section.

Figure legends

Figure 1. (A) General statistics of the 145 *Nucleocytoviricota* genomes. Each dot represents a

viral genome, the x-axis shows the average abundance of the genome across 16 timepoints (log-

transformed TPM), the y-axis shows the percentage of genes recovered from the transcriptomes,

and the dot size correlates to the total number of genes recovered. (B) The taxa distribution of the

145 genomes. The x-axis represents the total number of viruses in each taxon, and the y-axis

shows the *Nucleocytoviricota* family (left) and clade (right). In the clade plot (right), clades with

less than two members are collapsed down into "Other" category. Abbreviations: MC:

Mimiviridae, LP: *Late Phycodnaviridae*.

Figure 2. Phylogeny of *Nucleocytoviricota* with mean read mapping (TPM, log10-transformed)

on the outermost bar plot. Family classification of *Nucleocytoviricota* is denoted on the outer

ring and clade number is shown on the inner ring.

Figure 3. Expression level of *Nucleocytoviricota* during the 16 timepoints studied (A) Whole-

genome expression profiles of six genomes with significant difference in expression level

between daytime and nighttime (Mann-Whitney U test p< 0.1). The x-axis represents the 16

timepoints and the y-axis represents the total expression (TPM). *Nucleocytoviricota* clade

classifications are shown in brackets. Abbreviations: MC: *Mimiviridae*, LP: *Late*

Phycodnaviridae. (B) Eigengene of the genome co-expression modules. The x-axis represents

the 16 timepoints and the y-axis represents the eigengene expression values.

Figure 4. Metabolic genes expressed in the transcriptomes. The x-axis shows different viral

clades and the y-axis denotes the functional annotation. The sizes of the bubbles represent the

total abundance of the gene in TPM (log-transformed) and the colors show the functional

category. Abbreviations: a) Clade assignments. AF: *Asfarviridae*, EP: *Early Phycodnaviridae,*

IR: *Iridoviridae,* MC: *Mimiviridae*, LP: *Late Phycodnaviridae*. b) Gene function. PFK:

Phosphofructokinase, PGM: Phosphoglycerate mutase, GAPDH: Glyceraldehyde 3-P

dehydrogenase, PEPCK: Phosphoenolpyruvate carboxykinase, FBP1: Fructose-1-6-

bisphosphatase, SDH: Succinate dehydrogenase, 6PGD: 6-phosphogluconate dehydrogenase,

LHCB: Chlorophyll a/b binding protein, ACAD: Acyl-CoA dehydrogenase, HAT: Histone

acetyltransferase, GST: Glutathione S-transferase.

Figure 5. Phylogeny of *Nucleocytoviricota* myosin (left) and kinesin (right) proteins together

with references available in the EggNOG database. The myosin and kinesin expressed are

colored red.

Figure 6. Host-virus expression correlation network. Each node represents a eukaryotic genome

(circle) or viral genome (square). The line between two genomes denotes a correlation

coefficient larger than 0.8. The size of the node represents the total abundance in TPM (log-

transformed) of the genome and the colors show the genome's taxonomy.

Supplemental Documents

Supplemental Figure S1. Histogram of genome size distribution of the 145 *Nucleocytoviricota*

viruses present in the metatranscriptomes.

Supplemental Figure S2. *Nucleocytoviricota* metabolic gene expression in each of the 16

timepoints. The y-axis shows the functional annotation; the x-axis denotes the timepoints with

the numbers after the underscores indicating the hours. The sizes of the bubbles represent the

total abundance in TPM (log-transformed) of the gene across all 16 timepoints and the colors

show the functional categories. Abbreviations: PFK: Phosphofructokinase, PGM:

Phosphoglycerate mutase, GAPDH: Glyceraldehyde 3-P dehydrogenase, PEPCK:

Phosphoenolpyruvate carboxykinase, FBP1: Fructose-1-6-bisphosphatase, SDH: Succinate

dehydrogenase, 6PGD: 6-phosphogluconate dehydrogenase, LHCB: Chlorophyll a/b binding

protein, ACAD: Acyl-CoA dehydrogenase, HAT: Histone acetyltransferase, GST: Glutathione

S-transferase.

Supplemental Data S1. Summary of expression level of the 145 viruses with mapping in the

metatranscriptomes. Column name abbreviations: *num_genes_expressed*: the number of genes

in the genome that were expressed in at least one transcriptome; *sum_TPM, mean_TPM, median*

TPM: the total, average and median abundances (in TPM) of genomes across 16 timepoints,

respectively; *num_timepoints_expressed*: number of timepoints at which the gene was expressed;

proteins: the total number of protein encoded in the viral genome; *percent_proteins*: the

percentage of proteins in the genome that had hits in at least one metatranscriptome;

NCLDV_markers: the presence or absence of the four marker genes A32, major capsid protein

(mcp), SFII, and PolB in the genome; *mwt_day_night_pval:* p-values of Mann-Whitney U test

for significantly different expression during daytime versus nighttime periods; *mwt_diff:* the

difference between the total expression during daytime versus nighttime periods; *overexpressed*:

the time period when the genome is significantly higher expressed according to Mann-Whitney

U test; *Cor_Euk*: names of eukaryotic species that have significant correlation in expression (i.e.

correlations coefficients exceeding 0.8) with the virus ; *Correlation_strength:* correlations

coefficient values corresponding to the Eukaryotes listed in the *Cor_Euk* column.

Supplemental Data S2. Information on *Nucleocytoviricota* genes expressed in the

transcriptomes: Total expression across all 16 timepoints (*Sum_TPM*), number of timepoints at

which the gene was expressed (*num_timepoints_expressed*), gene annotations and similarity bit

score with reference to the databases EggNOG 4.5 and Pfam v. 31, and marker gene annotations

(*NCLDV_marker*).

Acknowledgments

We acknowledge the use of the Virginia Tech Advanced Research Computing Center for

bioinformatic analyses performed in this study. We are thankful to the members of Aylward Lab

for their helpful suggestions.

References

1.   Fischer MG. 2016. Giant viruses come of age. Curr Opin Microbiol 31:50–57.

2.   Koonin EV. 2005. Virology: Gulliver among the Lilliputians. Current Biology.

3. Koonin EV, Yutin N. 2010. Origin and evolution of eukaryotic large nucleo-cytoplasmic

DNA viruses. Intervirology 53:284–292.

4. Koonin EV, Dolja VV, Krupovic M, Varsani A, Wolf YI, Yutin N, Zerbini FM, Kuhn JH.

2020. Global Organization and Proposed Megataxonomy of the Virus World. Microbiol

Mol Biol Rev 84.

5. Koonin EV, Yutin N. 2012. Nucleo-cytoplasmic Large DNA Viruses (NCLDV) of

Eukaryotes. eLS.

6. 2019. Evolution of the Large Nucleocytoplasmic DNA Viruses of Eukaryotes and

Convergent Origins of Viral Gigantism, p. 167–202. *In* Advances in Virus Research.

Academic Press.

7. Karki S, Moniruzzaman M, Aylward FO. 2021. Comparative Genomics and Environmental

Distribution of Large dsDNA viruses in the family Asfarviridae. Cold Spring Harbor

Laboratory.

8. Abergel C, Legendre M, Claverie J-M. 2015. The rapidly expanding universe of giant

viruses: Mimivirus, Pandoravirus, Pithovirus and Mollivirus. FEMS Microbiol Rev 39:779–

796.

9. Blanc-Mathieu R, Dahle H, Hofgaard A, Brandt D, Ban H, Kalinowski J, Ogata H, Sandaa

R-A. 2021. A persistent giant algal virus, with a unique morphology, encodes an

unprecedented number of genes involved in energy metabolism. J Virol

https://doi.org/10.1128/JVI.02446-20.

10.   Moniruzzaman M, Martinez-Gutierrez CA, Weinheimer AR, Aylward FO. 2020. Dynamic

524   genome evolution and complex virocell metabolism of globally-distributed giant viruses.

525   Nat Commun 11:1710.

526   11.   Moreau H, Piganeau G, Desdevises Y, Cooke R, Derelle E, Grimsley N. 2010. Marine

527   prasinovirus genomes show low evolutionary divergence and acquisition of protein

528   metabolism genes by horizontal gene transfer. J Virol 84:12555–12563.

529   12.   Yutin N, Koonin EV. 2012. Proteorhodopsin genes in giant viruses. Biol Direct 7:34.

530   13.   Rozenberg A, Oppermann J, Wietek J, Fernandez Lahore RG, Sandaa R-A, Bratbak G,

531   Hegemann P, Béjà O. 2020. Lateral Gene Transfer of Anion-Conducting

532   Channelrhodopsins between Green Algae and Giant Viruses. Curr Biol 30:4910–4920.e5.

533   14.   Wilson WH, Schroeder DC, Allen MJ, Holden MTG, Parkhill J, Barrell BG, Churcher C,

534   Hamlin N, Mungall K, Norbertczak H, Quail MA, Price C, Rabbinowitsch E, Walker D,

535   Craigon M, Roy D, Ghazal P. 2005. Complete genome sequence and lytic phase

536   transcription profile of a Coccolithovirus. Science 309:1090–1092.

537   15.   Monier A, Pagarete A, de Vargas C, Allen MJ, Read B, Claverie J-M, Ogata H. 2009.

538   Horizontal gene transfer of an entire metabolic pathway between a eukaryotic alga and its

539   DNA virus. Genome Res 19:1441–1449.

540   16.   Schvarcz CR, Steward GF. 2018. A giant virus infecting green algae encodes key

541   fermentation genes. Virology 518:423–433.

17.   Forterre P. 2013. The virocell concept and environmental microbiology. ISME J 7:233–236.

18.   Forterre P. 2011. Manipulation of cellular syntheses and the nature of viruses: The virocell

concept. C R Chim 14:392–399.

19.   Rosenwasser S, Ziv C, van Creveld SG, Vardi A. 2016. Virocell Metabolism: Metabolic

Innovations During Host-Virus Interactions in the Ocean. Trends Microbiol 24:821–832.

20.   Ogata H, Toyoda K, Tomaru Y, Nakayama N, Shirai Y, Claverie J-M, Nagasaki K. 2009.

Remarkable sequence similarity between the dinoflagellate-infecting marine girus and the

terrestrial pathogen African swine fever virus. Virol J 6:178.

21.   Sandaa RA, Heldal M, Castberg T, Thyrhaug R, Bratbak G. 2001. Isolation and

characterization of two viruses with large genome size infecting Chrysochromulina ericina

(Prymnesiophyceae) and Pyramimonas orientalis (Prasinophyceae). Virology 290:272–280.

22.   Brussaard CPD, Short SM, Frederickson CM, Suttle CA. 2004. Isolation and phylogenetic

analysis of novel viruses infecting the phytoplankton Phaeocystis globosa

(Prymnesiophyceae). Appl Environ Microbiol 70:3700–3705.

23.   Moniruzzaman M, LeCleir GR, Brown CM, Gobler CJ, Bidle KD, Wilson WH, Wilhelm

SW. 2014. Genome of brown tide virus (AaV), the little giant of the Megaviridae, elucidates

NCLDV genome expansion and host-virus coevolution. Virology 466-467:60–70.

24.   Claverie J-M, Abergel C. 2018. : An Expanding Family of Highly Diverse Large dsDNA

Viruses Infecting a Wide Phylogenetic Range of Aquatic Eukaryotes. Viruses 10.

25.   Ghedin E, Claverie J-M. 2005. Mimivirus relatives in the Sargasso sea. Virol J 2:62.

26.   Monier A, Larsen JB, Sandaa R-A, Bratbak G, Claverie J-M, Ogata H. 2008. Marine

mimivirus relatives are probably large algal viruses. Virol J 5:12.

27.   Williamson SJ, Rusch DB, Yooseph S, Halpern AL, Heidelberg KB, Glass JI, Andrews-

Pfannkoch C, Fadrosh D, Miller CS, Sutton G, Frazier M, Venter JC. 2008. The Sorcerer II

Global Ocean Sampling Expedition: metagenomic characterization of viruses within aquatic

microbial samples. PLoS One 3:e1456.

28.   Chen F, Suttle CA, Short SM. 1996. Genetic diversity in marine algal virus communities as

revealed by sequence analysis of DNA polymerase genes. Appl Environ Microbiol

62:2869–2874.

29.   Endo H, Blanc-Mathieu R, Li Y, Salazar G, Henry N, Labadie K, de Vargas C, Sullivan

MB, Bowler C, Wincker P, Karp-Boss L, Sunagawa S, Ogata H. 2020. Biogeography of

marine giant viruses reveals their interplay with eukaryotes and ecological functions. Nat

Ecol Evol 4:1639–1649.

30.   Kaneko H, Blanc-Mathieu R, Endo H, Chaffron S, Delmont TO, Gaia M, Henry N,

Hernández-Velázquez R, Nguyen CH, Mamitsuka H, Forterre P, Jaillon O, de Vargas C,

Sullivan MB, Suttle CA, Guidi L, Ogata H. 2021. Eukaryotic virus composition can predict

the efficiency of carbon export in the global ocean. iScience 24:102002.

31.   Mihara T, Koyano H, Hingamp P, Grimsley N, Goto S, Ogata H. 2018. Taxon Richness of

"Megaviridae" Exceeds those of Bacteria and Archaea in the Ocean. Microbes and

Environments.

32.   Li Y, Hingamp P, Watai H, Endo H, Yoshida T, Ogata H. 2018. Degenerate PCR Primers to

Reveal the Diversity of Giant Viruses in Coastal Waters. Viruses 10.

33. Laber CP, Hunter JE, Carvalho F, Collins JR, Hunter EJ, Schieler BM, Boss E, More K,

Frada M, Thamatrakoln K, Brown CM, Haramaty L, Ossolinski J, Fredricks H, Nissimov JI,

Vandzura R, Sheyn U, Lehahn Y, Chant RJ, Martins AM, Coolen MJL, Vardi A, DiTullio

GR, Van Mooy BAS, Bidle KD. 2018. Coccolithovirus facilitation of carbon export in the

North Atlantic. Nat Microbiol 3:537–547.

34. Sheyn U, Rosenwasser S, Lehahn Y, Barak-Gavish N, Rotkopf R, Bidle KD, Koren I,

Schatz D, Vardi A. 2018. Expression profiling of host and virus during a coccolithophore

bloom provides insights into the role of viral infection in promoting carbon export. ISME J

12:704–713.

35. Zeigler Allen L, McCrow JP, Ininbergs K, Dupont CL, Badger JH, Hoffman JM, Ekman M,

Allen AE, Bergman B, Venter JC. 2017. The Baltic Sea Virome: Diversity and

Transcriptional Activity of DNA and RNA Viruses. mSystems 2.

36. Carradec Q, Pelletier E, Da Silva C, Alberti A, Seeleuthner Y, Blanc-Mathieu R, Lima-

Mendez G, Rocha F, Tirichine L, Labadie K, Kirilovsky A, Bertrand A, Engelen S, Madoui

M-A, Méheust R, Poulain J, Romac S, Richter DJ, Yoshikawa G, Dimier C, Kandels-Lewis

S, Picheral M, Searson S, Tara Oceans Coordinators, Jaillon O, Aury J-M, Karsenti E,

Sullivan MB, Sunagawa S, Bork P, Not F, Hingamp P, Raes J, Guidi L, Ogata H, de Vargas

C, Iudicone D, Bowler C, Wincker P. 2018. A global ocean atlas of eukaryotic genes. Nat

Commun 9:373.

37. Kolody BC, McCrow JP, Allen LZ, Aylward FO, Fontanez KM, Moustafa A,

Moniruzzaman M, Chavez FP, Scholin CA, Allen EE, Worden AZ, Delong EF, Allen AE.

2019. Diel transcriptional response of a California Current plankton microbiome to light,

low iron, and enduring viral infection. ISME J 13:2817–2833.

38.   Santini S, Jeudy S, Bartoli J, Poirot O, Lescot M, Abergel C, Barbe V, Wommack KE,

608   Noordeloos AAM, Brussaard CPD, Claverie J-M. 2013. Genome of Phaeocystis globosa

609   virus PgV-16T highlights the common ancestry of the largest known DNA viruses infecting

610   eukaryotes. Proc Natl Acad Sci U S A 110:10800–10805.

611   39.   Stough JMA, Yutin N, Chaban YV, Moniruzzaman M, Gann ER, Pound HL, Steffen MM,

612   Black JN, Koonin EV, Wilhelm SW, Short SM. 2019. Genome and Environmental Activity

613   of a Chrysochromulina parva Virus and Its Virophages. Frontiers in Microbiology.

614   40.   Gallot-Lavallée L, Blanc G, Claverie J-M. 2017. Comparative Genomics of

615   Chrysochromulina Ericina Virus and Other Microalga-Infecting Large DNA Viruses

616   Highlights Their Intricate Evolutionary Relationship with the Established Mimiviridae

617   Family. J Virol 91.

618   41.   Weynberg KD, Allen MJ, Wilson WH. 2017. Marine Prasinoviruses and Their Tiny

619   Plankton Hosts: A Review. Viruses 9.

620   42.   Needham DM, Yoshizawa S, Hosaka T, Poirier C, Choi CJ, Hehenberger E, Irwin NAT,

621   Wilken S, Yung C-M, Bachy C, Kurihara R, Nakajima Y, Kojima K, Kimura-Someya T,

622   Leonard G, Malmstrom RR, Mende DR, Olson DK, Sudo Y, Sudek S, Richards TA,

623   DeLong EF, Keeling PJ, Santoro AE, Shirouzu M, Iwasaki W, Worden AZ. 2019. A

624   distinct lineage of giant viruses brings a rhodopsin photosystem to unicellular marine

625   predators. Proceedings of the National Academy of Sciences.

43.   Aylward FO, Boeuf D, Mende DR, Wood-Charlson EM, Vislova A, Eppley JM, Romano

AE, DeLong EF. 2017. Diel cycling and long-term persistence of viruses in the ocean's

euphotic zone. Proc Natl Acad Sci U S A 114:11446–11451.

44. Derelle E, Yau S, Moreau H, Grimsley NH. 2018. Prasinovirus Attack of Ostreococcus Is

Furtive by Day but Savage by Night. J Virol 92.

45. Aylward FO, Eppley JM, Smith JM, Chavez FP, Scholin CA, DeLong EF. 2015. Microbial

community transcriptional networks are conserved in three domains at ocean basin scales.

Proc Natl Acad Sci U S A 112:5443–5448.

46. Moniruzzaman M, Wurch LL, Alexander H, Dyhrman ST, Gobler CJ, Wilhelm SW. 2017.

Virus-host relationships of marine single-celled eukaryotes resolved from

metatranscriptomics. Nat Commun 8:16054.

47. Needham DM, Poirier C, Hehenberger E, Jiménez V, Swalwell JE, Santoro AE, Worden

AZ. 2019. Targeted metagenomic recovery of four divergent viruses reveals shared and

distinctive characteristics of giant viruses of marine eukaryotes. Philos Trans R Soc Lond B

Biol Sci 374:20190086.

48. Rodrigues RAL, Louazani AC, Picorelli A, Oliveira GP, Lobo FP, Colson P, La Scola B,

Abrahão JS. 2020. Analysis of a Marseillevirus Transcriptome Reveals Temporal Gene

Expression Profile and Host Transcriptional Shift. Front Microbiol 11:651.

49. Legendre M, Audic S, Poirot O, Hingamp P, Seltzer V, Byrne D, Lartigue A, Lescot M,

Bernadac A, Poulain J, Abergel C, Claverie J-M. 2010. mRNA deep sequencing reveals 75

new genes and a complex transcriptional landscape in Mimivirus. Genome Res 20:664–674.

50. Lartigue A, Burlat B, Coutard B, Chaspoul F, Claverie J-M, Abergel C. 2015. The

megavirus chilensis Cu,Zn-superoxide dismutase: the first viral structure of a typical

cellular copper chaperone-independent hyperstable dimeric enzyme. J Virol 89:824–832.

51. Moniruzzaman M, Gann ER, Wilhelm SW. 2018. Infection by a Giant Virus (AaV) Induces

Widespread Physiological Reprogramming in CCMP1984 A Harmful Bloom Algae. Front

Microbiol 9:752.

52. Monier A, Chambouvet A, Milner DS, Attah V, Terrado R, Lovejoy C, Moreau H, Santoro

AE, Derelle É, Richards TA. 2017. Host-derived viral transporter protein for nitrogen

uptake in infected marine phytoplankton. Proc Natl Acad Sci U S A 114:E7489–E7498.

53. Monier A, Welsh RM, Gentemann C, Weinstock G, Sodergren E, Armbrust EV, Eisen JA,

Worden AZ. 2012. Phosphate transporters in marine phytoplankton and their viruses: cross-

domain commonalities in viral-host gene exchanges. Environ Microbiol 14:162–176.

54. Aherfi S, Belhaouari DB, Pinault L, Baudoin J-P, Decloquement P, Abrahao J, Colson P,

Levasseur A, Lamb DC, Chabriere E, Raoult D, La Scola B. 2020. Tricarboxylic acid cycle

and proton gradient in Pandoravirus massiliensis: Is it still a virus? Cold Spring Harbor

Laboratory.

55. Vincent F, Sheyn U, Porat Z, Vardi A. 2020. Visualizing active viral infection reveals

diverse cell fates in synchronized algal bloom demise. Cold Spring Harbor Laboratory.

56. Gastrich MD, Leigh-Bell JA, Gobler, CH, Anderson, OR, Wilhelm, SW, Bryan M. 2004.

Viruses as potential regulators of regional brown tide blooms caused by the alga,

Aureococcus anophagefferens 27:112–119.

57. Da Cunha V, Gaia M, Ogata H, Jaillon O, Delmont TO, Forterre P. 2020. Giant viruses

encode novel types of actins possibly related to the origin of eukaryotic actin: the viractins.

Cold Spring Harbor Laboratory.

58. Moniruzzaman M, Weinheimer AR, Martinez-Gutierrez CA, Aylward FO. 2020.

Widespread endogenization of giant viruses shapes genomes of green algae. Nature

588:141–145.

59. Guglielmini J, Woo AC, Krupovic M, Forterre P, Gaia M. 2019. Diversification of giant

and large eukaryotic dsDNA viruses predated the origin of modern eukaryotes. Proc Natl

Acad Sci U S A 116:19585–19592.

60. Schulz F, Roux S, Paez-Espino D, Jungbluth S, Walsh DA, Denef VJ, McMahon KD,

Konstantinidis KT, Eloe-Fadrosh EA, Kyrpides NC, Woyke T. 2020. Giant virus diversity

and host interactions through global metagenomics. Nature.

61. Stough JMA, Kolton M, Kostka JE, Weston DJ, Pelletier DA, Wilhelm SW. 2018. Diversity

of Active Viral Infections within the Sphagnum Microbiome. Appl Environ Microbiol 84.

62. Meng L, Endo H, Blanc-Mathieu R, Chaffron S, Hernández-Velázquez R, Kaneko H, Ogata

H. 2020. Quantitative assessment of NCLDV–host interactions predicted by co-occurrence

analyses. Cold Spring Harbor Laboratory.

63. Lynam CP, Llope M, Möllmann C, Helaouët P, Bayliss-Brown GA, Stenseth NC. 2017.

Interaction between top-down and bottom-up control in marine food webs. Proc Natl Acad

Sci U S A 114:1952–1957.

64.   Matsuyama T, Takano T, Nishiki I, Fujiwara A, Kiryu I, Inada M, Sakai T, Terashima S,

Matsuura Y, Isowa K, Nakayasu C. 2020. A novel Asfarvirus-like virus identified as a

potential cause of mass mortality of abalone. Sci Rep 10:4620.

65.   Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, Phillippy AM.

2016. Mash: fast genome and metagenome distance estimation using MinHash. Genome

Biol 17:132.

66.   Bobay L-M, Ochman H. 2018. Biological species in the viral world. Proceedings of the

National Academy of Sciences.

67.   Aylward FO, Moniruzzaman M. 2021. ViralRecall-A Flexible Command-Line Tool for the

Detection of Giant Virus Signatures in 'Omic Data. Viruses 13.

68.   Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal:

prokaryotic gene recognition and translation initiation site identification. BMC

Bioinformatics 11:119.

69.   Frith MC. 2011. A new repeat-masking method enables specific detection of homologous

sequences. Nucleic Acids Res 39:e23.

70.   Kiełbasa SM, Wan R, Sato K, Horton P, Frith MC. 2011. Adaptive seeds tame genomic

sequence comparison. Genome Res 21:487–493.

71.   Mortazavi A, Williams BA, McCue K, Schaeffer L, Wold B. 2008. Mapping and

quantifying mammalian transcriptomes by RNA-Seq. Nat Methods 5:621–628.

72.   Tithi SS, Aylward FO, Jensen RV, Zhang L. 2018. FastViromeExplorer: a pipeline for virus

and phage identification and abundance profiling in metagenomics data. PeerJ 6:e4227.

73.   Huerta-Cepas J, Szklarczyk D, Forslund K, Cook H, Heller D, Walter MC, Rattei T, Mende

710   DR, Sunagawa S, Kuhn M, Jensen LJ, von Mering C, Bork P. 2016. eggNOG 4.5: a

711   hierarchical orthology framework with improved functional annotations for eukaryotic,

712   prokaryotic and viral sequences. Nucleic Acids Res 44:D286–93.

713   74.   El-Gebali S, Mistry J, Bateman A, Eddy SR, Luciani A, Potter SC, Qureshi M, Richardson

714   LJ, Salazar GA, Smart A, Sonnhammer ELL, Hirsh L, Paladin L, Piovesan D, Tosatto SCE,

715   Finn RD. 2019. The Pfam protein families database in 2019. Nucleic Acids Res 47:D427–

716   D432.

717   75.   Koonin EV, Yutin N. 2018. Multiple evolutionary origins of giant viruses. F1000Res 7.

718   76.   Yutin N, Wolf YI, Raoult D, Koonin EV. 2009. Eukaryotic large nucleo-cytoplasmic DNA

719   viruses: clusters of orthologous genes and reconstruction of viral genome evolution. Virol J

720   6:223.

721   77.   Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated

722   alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25:1972–1973.

723   78.   Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective

724   stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol

725   32:268–274.

726   79.   Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017.

727   ModelFinder: fast model selection for accurate phylogenetic estimates. Nat Methods

14:587–589.

80. Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving

the Ultrafast Bootstrap Approximation. Mol Biol Evol 35:518–522.

81. Thaben PF, Westermark PO. 2014. Detecting rhythms in time series with RAIN. J Biol

Rhythms 29:391–400.

82. Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B

(Methodological).

83. Langfelder P, Horvath S. 2008. WGCNA: an R package for weighted correlation network

analysis. BMC Bioinformatics 9:1–13.

84. Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-likelihood

trees for large alignments. PLoS One 5:e9490.

85. Huerta-Cepas J, Dopazo J, Gabaldón T. 2010. ETE: a python Environment for Tree

Exploration. BMC Bioinformatics 11:24.

Figure 1. (A) General statistics of the 145 *Nucleocytoviricota* genomes. Each dot represents a viral genome, the x-axis shows the average abundance of the genome across 16 timepoints (log-transformed TPM), the y-axis shows the percentage of genes recovered from the transcriptomes, and the dot size correlates to the total number of genes recovered. (B) The taxa distribution of the 145 genomes. The x-axis represents the total number of viruses in each taxon, and the y-axis shows the *Nucleocytoviricota* family (left) and clade (right). In the clade plot (right), clades with

less than two members are collapsed down into "Other" category. Abbreviations: MC: *Mimiviridae*, LP: *Late Phycodnaviridae*.

Figure 2. Phylogeny of *Nucleocytoviricota* with mean read mapping (TPM, log10-transformed) on the outermost bar plot. Family classification of *Nucleocytoviricota* is denoted on the outer ring and clade number is shown on the inner ring.

Figure 3. Expression level of *Nucleocytoviricota* during the 16 timepoints studied (A) Whole-genome expression profiles of six genomes with significant difference in expression level between daytime and nighttime (Mann-Whitney U test p< 0.1). The x-axis represents the 16 timepoints and the y-axis represents the total expression (TPM). *Nucleocytoviricota* clade classifications are shown in brackets. Abbreviations: MC: *Mimiviridae*, LP: *Late Phycodnaviridae*. (B) Eigengene of the genome co-expression modules. The x-axis represents the 16 timepoints and the y-axis represents the eigengene expression values.

Figure 4. Metabolic genes expressed in the transcriptomes. The x-axis shows different viral

clades and the y-axis denotes the functional annotation. The sizes of the bubbles represent the

total abundance of the gene in TPM (log-transformed) and the colors show the functional category. Abbreviations: a) Clade assignments. AF: *Asfarviridae*, EP: *Early Phycodnaviridae,* IR: *Iridoviridae,* MC: *Mimiviridae*, LP: *Late Phycodnaviridae*. b) Gene function. PFK: Phosphofructokinase, PGM: Phosphoglycerate mutase, GAPDH: Glyceraldehyde 3-P dehydrogenase, PEPCK: Phosphoenolpyruvate carboxykinase, FBP1: Fructose-1-6-bisphosphatase, SDH: Succinate dehydrogenase, 6PGD: 6-phosphogluconate dehydrogenase, LHCB: Chlorophyll a/b binding protein, ACAD: Acyl-CoA dehydrogenase, HAT: Histone acetyltransferase, GST: Glutathione S-transferase.

Figure 5. Phylogeny of *Nucleocytoviricota* myosin (left) and kinesin (right) proteins together with references available in the EggNOG database. The myosin and kinesin expressed are circled and colored green.

Figure 6. Host-virus expression correlation network. Each node represents a eukaryotic genome (circle) or viral genome (square). The line between two genomes denotes a correlation coefficient larger than 0.8. The size of the node represents the total abundance in TPM (log-transformed) of the genome and the colors show the genome's taxonomy.