Nanoscopic dopamine transporter distribution and conformation are inversely regulated by excitatory drive and D₂-autoreceptor activity

Matthew D. Lycas[1], Aske L. Ejdrup[1], Andreas T. Sørensen[1], Nicolai O. Haahr[1], Søren H. Jørgensen[1], Daryl A. Guthrie [2], Jonatan F. Støier[1], Christian Werner[3], Amy Hauck Newman[2], Markus Sauer[3], Freja Herborg[1] and Ulrik Gether[1*]

[1]Molecular Neuropharmacology and Genetics Laboratory, Department of Neuroscience, Faculty of Health and Medical Sciences, Maersk Tower 7.5, University of Copenhagen, DK-2200 Copenhagen, Denmark

[2]Medicinal Chemistry Section, Molecular Targets and Medications Discovery Branch, National Institute on Drug Abuse-Intramural Research Program, National Institutes of Health, Baltimore, Maryland 21224

[3]Department of Biotechnology and Biophysics, Biocenter, Julius-Maximilians-Universität Würzburg, Am Hubland, 97074 Würzburg, Germany

*Correspondence: gether@sund.ku.dk

Abstract

The nanoscopic organization and regulation of individual molecular components in presynaptic

varicosities of neurons releasing modulatory volume neurotransmitters like dopamine (DA) remain largely

elusive. Here we show by application of several single-molecule sensitive super-resolution microscopy

techniques to cultured neurons and mouse striatal slices, that the dopamine transporter (DAT), a key

protein in varicosities of dopaminergic neurons, exists in the membrane in dynamic equilibrium between

an inward-facing nanodomain-localized and outward-facing unclustered configuration. The balance

between these configurations is inversely regulated by excitatory drive and by DA D2-autoreceptor

activation in a manner dependent on Ca^{2+}-influx via N-type voltage-gated Ca^{2+}-channels. The DAT

nanodomains contain tens of transporters molecules and overlap with nanodomains of PIP2

(phosphatidylinositol-4,5-bisphosphate) but show little overlap with D2-autoreceptor, syntaxin-1 and

clathrin nanodomains. Summarized, the data reveal a mechanism for rapid alterations in nanoscopic DAT

distribution and show a striking link of this to the conformational state of the transporter.

Introduction

In its 15 years of utilization, super-resolution microscopy has improved our insights into the molecular

organization and dynamic regulation of pre- and postsynaptic structures in the brain. Indeed, it has

become clear that regulatory processes of fundamental importance for spatial and temporal control of

synaptic signaling occur at a nanoscale level. In excitatory synapses, for example, it is now believed that

AMPA-type and NMDA-type ionotropic glutamate receptors segregate into distinct nanodomains in the

postsynaptic density and that this co-organization plays a critical role in synaptic physiology[1,2]. Such

studies highlight the importance of observing past the diffraction limit of visible light while still leveraging

labeling strategies and live imaging compatibility that light microscopy provides[1-3].

Much less is known about the nanoscale architecture of the presynaptic compartment in neurons

releasing modulatory transmitters like dopamine (DA). DA neurons project from the midbrain to the basal

ganglia, the limbic system and the prefrontal cortex to mediate the role of DA in motor functions, reward

mechanisms and learning processes[4,5]. The DA neurons have remarkable axonal arbors with numerous

release sites (i.e. varicosities) that enable single DA neurons to affect the function of multiple target

neurons[6,7]. DA neurotransmission, however, differs from classical fast synaptic transmission by operating

perhaps primarily via "volume transmission", that is, DA is predominantly released from non-synaptic

release sites to act on target cells often located microns away[6,8]. It remains nonetheless elusive how the

different molecular components in the dopaminergic release sites need to be organized at a nanoscale

level and individually regulated to enable proper neurotransmitter release and reuptake.

A key regulator of DA neurotransmission, the dopamine transporter (DAT), is highly expressed in

the varicosities of DA neurons and removes released DA from the extracellular space. DAT has received

attention as the primary target for therapeutics, such as methylphenidate, and drugs of abuse such as

methamphetamine and cocaine[9,10]. Not surprisingly, DAT is subject to tight regulation through several

mechanisms involving post-translational modifications, protein-protein interactions and protein-lipid

interactions[9,11,12]. Moreover, we recently showed by application of super-resolution imaging that DAT

regulation might involve mechanisms that cannot be revealed by more classical methods[13]. Our data

showed that DAT localizes to discrete nanodomains in the plasma membrane of cultured DA neurons and

that activation of NMDA-receptors reduced DAT nanodomain localization[13]. Although this indicated that

nanoscale dynamics in dopamine varicosities could be critical for controlling DA neurotransmission, the

mechanisms and functional implications remain unknown.

Here we use several super-resolution imaging techniques, including Photoactivated Localization

Microscopy (PALM)[14], direct Stochastic Optical Reconstruction Microscopy (dSTORM)[15,16], 10 x Expansion

Microscopy (ExM)[17-19] and single-particle tracking dSTORM (spt-dSTORM)[20] together with new fluorescent

cocaine-analogues[21] and new data analysis tools, to investigate how DAT and associated molecular

components at DA varicosities are dynamically regulated. In summary, our experimental efforts

demonstrate that excitatory drive and D2-autoreceptor activity inversely regulate in a Ca^{2+} dependent

manner the conformational state of DAT and its localization in DA varicosities to PIP2 enriched

nanodomains separate from syntaxin-1 (STX-1), D2-autoreceptor or clathrin nanodomains. In this way,

our study uncovers a hitherto unknown level of regulation of an essential membrane protein in a

modulatory transmitter terminal.

Results

DAT shows nanodomain distribution in cultured neurons and striatal slices. There are a variety of

strategies to resolve biological structures past the diffraction limit of light, but each method is associated

with strengths, compromises, and potential artefacts. Our previous dSTORM work suggested that DAT

distributes into nanodomains at DA varicosities of cultured midbrain DA neurons[13]. To substantiate this

finding, we decided to assess DAT distribution across a variety of super resolution methods and to

ultimately investigate whether DAT is distributed into nanodomains in intact tissue. First, we assessed the

distribution of DAT regardless of potential antibody-induced clustering artifacts by transfecting cultured

DA neurons with DAT having the photoconvertible fluorescent protein mEOS2[22] fused to the N-terminus.

Images of tyrosine hydroxylase (TH) positive varicosities revealed a clustered distribution of the mEOS2-

DAT signal surrounding the cytosolic TH signal (Fig. 1A-D), similar to that seen by imaging the endogenous

DAT by dSTORM[13]. We next employed astigmatic 3D-dSTORM to show that the DAT nanodomains were

localized to the periphery of the varicosities, substantiating that DAT is found mainly in the plasma

membrane (Fig. 1E-H). To assess the distribution of DAT in intact tissue, we analyzed coronal mouse brain

slices (Fig. 1I-L). Widefield imaging exposed the DAT positive varicosities and by dSTORM we could

visualize DAT in single varicosities, showing a nanodomain distribution like that seen in cultured neurons

(Fig. 1I-L). To exclude that the observed distribution of DAT was not caused by artifacts introduced by the

single molecule localization microscopy approach (e.g., "overcounting" [23]), we turned to ExM where

nanoscale spatial resolution is obtained by expanding the tissue sample using a polymer system[19]. We

employed a 10x expansion protocol[18] to mouse striatal slices before DAT imaging using a spinning disk

confocal microscope (Fig. 1M-P). In the stained expanded sample, we captured the intermingled axonal

network of DA neurons while at the same time observing clustering of DAT into nanodomains in

varicosities (Fig. 1M-P). Summarized, the distribution of DAT into nanodomains in the plasma membrane

of DA neurons was substantiated by several high-resolution imaging techniques.

Having reaffirmed the clustered organization of DAT with different techniques and in intact brain

tissue, we decided to focus on the dSTORM technology in our subsequent studies because of its

quantitative nature and single-molecule sensitivity. To derive quantitative measures for the clustered

distribution of DAT , we employed the DBSCAN algorithm (density based spatial clustering of applications

with noise)[13,24] and extracted two key parameters from the analysis: the size of the nanoclusters and the

density of localizations within the clusters (Fig. 1Q-S). The size was quantified as the fraction of clusters

with a diameter >75 nm (Fig. 1R). This size was chosen as threshold to exclude any clusters that might be

the result of amplified labeling of single proteins and thus might arise from the antibody labeling

methodology. By assessing the density of localizations as well, we could determine changes in the number

of molecules in a cluster without a change in cluster size. Dense clusters were defined as clusters with >80

localizations within a radius of 50 nm (dense clusters in red) (Fig. 1S). We also employed Voronoi

tessellation to compare localization density distributions in a parameter free fashion [25]. In this use of

Voronoi tessellation, clusters themselves were not identified but localization density changes could be

detected with greater sensitivity (Fig. 1T).

We also wanted to estimate how many DAT molecules are present in an average DAT nanodomain

defined by the DBSCAN analysis. We used a protocol modified from[26,27] and labeled cultured DA neurons

with sequential dilutions of the primary DAT antibody. This enabled us to define that a minimum cluster

is made from ~three localizations and that an average large cluster contains ~33 copies of the antibody

corresponding to 50-100 DAT molecules as a rough estimate (Supplementary Fig. 1). The same data series

lent itself perfectly for cluster analysis by varied label density cluster verification, enabling us to

distinguish between random clustering arising from multiple observations of single fluorophores and true

clustering[28]. In agreement with our data in DAT expressing Cath.a-differentiated (CAD) cells[13], our analysis

supported that the clustering behavior of DAT in cultured neurons reflect true nanocluster formation

(Supplementary Fig. 2).

Nanodomain localized DAT is preferentially in an inward-facing conformation. To assess putative

functional implications of the nanoclustered distribution of DAT, we investigated whether this could

involve a conformational bias with nanoclustered DAT having a different conformation than unclustered

DAT. Cocaine and cocaine-like inhibitors are known to bind with higher affinity to DAT in the outward-

facing conformation than in the inward-facing conformation[29,30]. We envisioned accordingly (and in line

with[31]) that an indirect measure of the conformational state of DAT inside and outside the nanodomains

could be obtained by parallel imaging of binding to DAT of a fluorescently tagged cocaine-analogue and

of DAT itself; that is, as a result of its conformational bias the cocaine-analogue would be expected to

bind more to outward-facing than to inward-facing transporters at a subsaturating concentration. We

labeled cultured DA neurons and mEOS2-DAT expressing CAD cells with DG3-63 (10 nM), an analogue of

cocaine tagged with JF_{646} suited for dSTORM (Fig. 2A-D). The DG3-63 signal generally appeared somewhat

less clustered as compared to that observed by our other approaches (see Fig. 1). This might indicate a

preferential binding of DG3-63 to unclustered DAT, which was supported by dual-color images of DG3-63

labelled DAT and mEOS-DAT, showing a more dispersed DG3-63 signal as compared to a more clustered

mEOS2-DAT signal (Fig. 2E-G). Importantly, the DG3-63 signal was specific as it was essentially eliminated

by excess of unlabeled cocaine (Fig. 2H). To quantify whether DG3-63 preferentially bound to unclustered

DAT, we used a modification of the principles outlined by the colocalization extension of the Voronoi

tessellation algorithm [25] where associated DG3-63 was identified by including DG3-63 localizations within

25 nm from an mEOS2-DAT localization (Fig. 2I-K). The total mEOS2-DAT Voronoi tessellated area

distribution showed a preference for smaller tessellated areas consistent with a clustered distribution

(Fig. 2L). In contrast, the mEOS2-DAT-associated DG3-63 localizations resided on larger Voronoi

tessellated areas; hence, DG3-63 binding was not enriched in the DAT nanodomains containing the

highest density of DAT molecules (Fig. 2L). This suggests that nanoclustered DAT are less prone to bind

DG3-63, consistent with a more inward-facing transporter conformation or a less accessible binding site.

Interestingly, by analyzing the distribution of mEOS2-DAT after treatment with an excess, and thereby

saturating concentration, of cocaine, we observed a dispersal of the signal. This supports that forcing the

transporter into an outward-facing conformation may release the transporter from the nanodomains

(Supplementary Fig. 3).

We further rationalized that if the nanoscale DAT distribution reflects its molecular conformation,

then a redistribution would be expected for a DAT variant with a conformational bias. Accordingly, we

compared in transfected CAD cells clustering of WT DAT with that of a disease-associated DAT mutation

(D421N) known to be shifted towards an inward-facing conformation[32] (Fig. 2M-R). In agreement with the

idea that inward-facing DAT will display preferential localization to nanodomains, dSTORM after

immunolabeling revealed for D421N compared to WT both a higher fraction of large clusters (diameter

>75 nm) and a higher fraction of localizations in dense clusters (Fig. 2N, O, Q, R) without a change in total

number of detected localizations (Fig. 2P). Summarized, the data support that DAT preferentially assumes

an inward-facing conformation in DAT nanodomains while the conformation outside the nanodomains

conceivably is more outward-facing.

NMDA receptor activation declusters DAT in striatal slices. We next assessed whether DAT nanodomain

distribution was sensitive to NMDA stimulation in intact tissue (Fig. 2I-P) as we previously observed in

cultured neurons[13]. Notably, NMDA receptors are present in DA terminals where they can stimulate DA

release[33]. Acute coronal striatal slices were treated for 5 min with vehicle, NMDA or NMDA plus the NMDA

receptor antagonist (2R)-amino-5-phosphonovaleric acid (AP5) before DAT immunolabeling and dSTORM

(Fig. 3A). Because each dSTORM image contained a vast number of DAT positive varicosities, we

developed a new collection algorithm for automated and unbiased isolation of these varicosities, allowing

subsequent quantification of DAT clustering (Fig. 3B, C). Following NMDA treatment, both the fraction of

clusters >75 nm and the faction of localizations in dense clusters decreased; an effect that was muted by

the NMDA antagonist AP5 (Fig. 3D, E). Note that the comparisons were normalized to the control sample

from the individual imaging session to correct for unavoidable variability in sample preparations and

between imaging sessions. Importantly, the data were further supported by Voronoi tessellation showing

that NMDA, but not NMDA + AP5, reduced the fraction of smaller tessellated areas and thus clustering

(Fig. 3F). We conclude that local activation of NMDA receptors in intact tissue promotes dispersal of DAT

nanodomains.

DAT nanodomain localization is regulated by neuronal activity. We speculated whether persistent

changes in membrane potential and thereby in neuronal excitability would affect DAT nanodomain

distribution. To test this, we expressed the bacterial sodium channel (mNaChBac) or the inward-rectifier

potassium channel (Kir2.1)[34,35] in cultured DA neurons to make them more or less excitable, respectively

[34]. In agreement with selective expression in DA neurons, we confirmed that transduced cells were also

DAT positive (Supplementary Fig. 4). Electrophysiological recordings on single neurons showed that the

pacemaker firing activity of DA neurons[36] was enhanced in mNaChBac-expressing neurons and reduced

in Kir2.1 expressing neurons (Fig S4). Analysis by dSTORM of DAT distribution in mNaChBac expressing

neurons showed decreased clustering, while for Kir2.1 expressing neurons we saw a trend towards an

increased fraction of large clusters and a significant increase in localizations within large dense clusters

(Fig. 4A-D). Thus, DAT nanodomain localization is in cultured DA neurons regulated by persistent changes

in excitability or firing activity with increased activity leading to dispersal of DAT from nanodomains and

decreased activity leading to increased nanodomains localization.

DAT nanodomain localization is regulated by Ca^{2+}. Because of the effect of neuronal excitability, we

decided to investigate putative involvement of Ca^{2+} influx via voltage-gated Ca^{2+} channels (VGCC) in

NMDA-induced DAT declustering. Note that both P/Q type and N-type VGCCs should be present in DA

terminals [37]. DA neurons were stimulated with NMDA for 5 min with or without the N-type VGCC blocker

ω-conotoxin or the P/Q type VGCC blocker ω-agatoxin. While we observed no effect of ω-agatoxin, the

dispersing effect of NMDA on DAT clustering was essentially abolished by ω-conotoxin. Similarly, Voronoi

tessellation analysis showed that ω-conotoxin, but not ω-agatoxin, blocked the NMDA-induced shift away

from small, tessellated areas and thus blocked the dispersal of DAT nanodomains (Fig. 4E-J, Movies S1-2).

We next asked whether Ca^{2+} would be important *per se* by preincubating the neurons with the cell

permeable Ca^{2+}-chelator BAPTA-AM. Indeed, BAPTA-AM blunted the response to NMDA (Fig. 4K-L). Of

note, the pharmacological data fitted well with available single-cell mRNA sequencing data on DA neurons

supporting highest transcription of N-type VGCC in DA neurons (Fig. 4M). Taken together, NMDA-induced

dispersal of DAT nanodomains appears to occur via Ca^{2+}-dependent mechanism involving the activity of

N-type VGCCs.

DAT nanodomain localization is regulated by the D2 autoreceptor. D2-autoreceptors in DA neurons act

as negative feedback regulators decreasing neuronal excitability and firing activity by promoting opening

of inhibitory G-protein-activated inwardly rectifying potassium channels[38]. We surmised that D2-

autoreceptor activity might regulate localization of DAT to nanodomains and tested this by determining

the effect of the D2 receptor antagonist haloperidol and the D2 receptor agonist quinpirole. Interestingly,

incubation with 10 nM haloperidol for 5 min resulted in dispersal of DAT nanodomains with a reduced

fraction of clusters >75 m, fewer localizations in dense clusters and a reduced fraction of smaller Voronoi

tessellated areas (Fig. 5A-F, Movies S1, S3). This effect is likely the result of haloperidol blocking the effect

on the D2-autoreceptors of DA released in the culture as a consequence of DA neuronal pacemaker

activity as previously demonstrated for this culture of DA neurons[36]. Also, co-incubation with an excess of

quinpirole blocked the effect of haloperidol (Fig. 5A-F). Moreover, the effect of haloperidol was

accompanied by an increase in cytoplasmic free Ca^{2+} consistent with D2-autoreceptor activity suppressing

neuronal firing (Supplementary Fig. 5). This prompted us to test if the effect of haloperidol, like the effect

of NMDA, was dependent on Ca^{2+}- influx via N-type VGCCs. This appeared to be the case as ω-conotoxin,

but not ω-agatoxin, eliminated the effect of haloperidol (Fig. 5A-F). Summarized, our data support that

D2-autoreceptors are preoccupied with DA in the neuronal culture and that the activity of these receptors

can regulate DAT nanodomain localization via a Ca^{2+}-dependent mechanism.

Single particle tracking supports nanodomain dispersal in response to neuronal activity. To obtain a

better understanding of the dynamics underlying DAT nanodomain localization, we employed SPT-

dSTORM [20] to assess if dispersal of DAT from nanodomains is accompanied by increased lateral mobility

of DAT in the membrane. To allow visualization in live cells, we labeled DAT in DA neurons with a Janelia

Fluor®549-conjugated cocaine analogue, DG3-80, suited for live dSTORM[21] (Fig. 5G, H). Individual DAT

tracks were captured with lengths up to 1.6 sec (100 frames, 16 msec frame rate) with the majority being

10-20 frames (Fig. 5I, J). Separating tracks into those reflecting confined movements and those reflecting

freely moving particles showed that ~90% of the tracks described freely moving particles (Fig. 5K, L). Note

that this distribution (with most of the labeled molecules being freely moving) likely reflects preferential

binding of DG3-80 to unclustered DAT in the outward-facing conformation (Fig. 2). The mean jump

distance (i.e. the distance an individual molecule/particle has traveled between each image) was ~110

nm for all the recorded tracks and ~50 nm for static tracks only (Fig. 5M, N). Interestingly, a jump distance

of ~110 nm is so fast that a transporter molecule might move 7-14 μm during a transport cycle estimated

to last 1-2 sec[39]. Among the freely moving particles, there were even some that moved very fast with

mean jump distance of 200-350 nm (Fig. 5M-O). Both NMDA receptor activation and blockade of the D2-

autoreceptors with either haloperidol or the neutral antagonist UH-232 increased the population of DAT

molecules with mean jump distances >200 nm, indicative of a possible link between dispersal of DAT

nanodomains and increased lateral mobility. Note that the similar effect of UH-232 and haloperidol

strongly suggest that their effect reflects blockade of D2-autoreceptors activated by DA, which is released

because of DA neuronal pacemaker activity, and not inhibition of constitutive receptor activity[40].

DAT nanodomains overlap with PIP2 nanodomains. A key question is how the nanoscale distribution of

DAT relates to other molecular components of DA terminals. We first turned to the plasma membrane

phospholipid, PIP2, which can bind DAT and in doing so modulate DAT activity[41]. Dual-color dSTORM on

PIP2 and DAT in DA neurons showed as expected[42], a distribution of the PIP2 signal into nanodomains

(Fig. 6). Strikingly, these nanodomains showed a remarkable overlap with DAT nanodomains with the

Voronoi tessellation distributions revealing a large fraction of PIP2 localizations associated with dense

DAT localizations (Fig. 6A, B, F). As also seen for DAT domains, treatment with both NMDA and haloperidol

reduced PIP2 localizations in large dense clusters while haloperidol also reduced cluster size. Importantly,

all these changes were blocked by AP5 or quinpirole, respectively (Fig. 6B-D). The NMDA and haloperidol-

induced changes were reflected in the Voronoi tessellation distributions as well by showing dispersal of

the PIP2 nanodomains and reduced PIP2 association to nanoclustered DAT (Fig. 6E, F).

The D2R is also shows nanodomain distribution. We next analyzed the D2-autoreceptors that are found

both in the somatodendritic compartment of DA neurons and in DA terminals. Due to known challenges

with DA receptor antibodies, we first verified that our antibody was specific for the receptor in transfected

HEK293 cells (Supplementary Fig. 5). Subsequent dual-color dSTORM experiments on cultured DA

neurons revealed a clustered distribution of the D2-autoreceptors with nanodomains of a similar size as

those found for DAT (Fig. 7A, B). Furthermore, like our findings for DAT and PIP2, NMDA and haloperidol

promoted dispersal of the receptor domains (Fig. 7B-E). Relatively few of the nanodomains, however,

overlapped with the DAT domains as apparent from the Voronoi tessellation distributions (Fig. 7A, B, F),

and as for D2-autoreceptor association to DAT, we only observed marginal changes upon NMDA and

haloperidol treatment (Fig. 7F).

The findings prompted us to further assess the specificity of the pharmacological manipulations by

imaging additional key membrane components in DA terminals including clathrin[43] and the single

transmembrane SNARE protein syntaxin1 (STX-1). Imaging of clathrin revealed a nanoclustered

distribution, possibly corresponding to clathrin-coated pits[44], that showed little co-localization with DAT

and, importantly, essentially no change in distribution in response to NMDA (Supplementary Fig. 6). We

also recapitulated nanodomain distribution of STX-1 that is known to exist in such domains in presynaptic

terminals [45] as well as the protein has been proposed to bind DAT[46]. However, the STX-1 domains showed

little overlap with DAT and responded differently to NMDA with a decreased fraction of large clusters and

an increased fraction of localizations in dense clusters (Supplementary Fig. 6). Note, however, that NMDA

slightly increased localization of STX1 to unclustered DAT (i.e. the larger tessellated areas) according to

the Voronoi tesselated distributions (FIgure S6). In summary, several plasma membrane components of

DA terminals show a nanoclustered distribution that, however, appears to be differentially regulated.

Lycas et al.

Discussion

DA neurons have remarkable axonal arbors with numerous presynaptic terminals or varicosities that serve

as the principal architectural units responsible for release and reuptake of DA. Still, we have a surprisingly

poor understanding of the molecular organization of these varicosities, and what regulatory processes

that take place within these compartments less than ~1 μm in size. Indeed, most of these varicosities are

non-synaptic release sites and it remains unclear whether this results in less stringent requirement of the

spatial organization of the molecular machinery as compared to e.g. glutamatergic synapses[1-3]. Here, we

utilize several super-resolution techniques to provide detailed, quantitative insight into how a key

membrane protein, DAT, is subject to nanoscale regulation within these varicosities. This regulation

involves Ca^{2+}-dependent movements of DAT in and out of PIP2-enriched nanodomains in the plasma

membrane, which are governed by NMDA receptor and D2-autoreceptor activity and likely involves shift

of DAT between outward and inward-facing conformations.

To firmly substantiate that DAT is distributed into nanodomains in the plasma membrane of DA

neurons, we applied a range of different super-resolution microscopy techniques to both cultured

neurons and striatal slices. A varied label density cluster analysis[28] supported that our dSTORM procedure

reports true nanodomains, which was further confirmed by ExM, excluding potential artifacts from

multiple countings of the same fluorophore. A rough estimate of the number of DAT molecules suggested

the presence of 50-100 DAT molecules within each domain, which is in the same range estimated for

AMPA receptor domains in the postsynapse (~25 molecules)[47]. Importantly, by using the novel fluorescent

cocaine analogue DG3-63 and the disease-associated DAT mutant D421N, we were able to obtain

evidence that DAT displays a conformational bias depending on whether it is present in a nanodomain or

not. Based on these data, it is appealing to envision that a distribution of DAT between a nanoclustered,

inward-facing population of transporter molecules and an unclustered, "free" outward-facing population

of DAT molecules represents a means by which DAT activity can be regulated on a very fast time scale. A

possible question is whether this dynamic localization of DAT to nanodomains involves changes in DAT

oligomerization; however, we find this unlikely considering recent data supporting the existence in the

plasma membrane of DA dimers that are highly stable even over the course of minutes[48].

Of major importance, we validated in striatal slices the dispersing effect of NMDA on DAT

nanodomains by employing a novel Voronoi tessellation-based collection algorithm. In addition to

support that our findings in cultured DA neurons can be translated to intact tissue, the algorithm offers

possibilities for future detailed, quantitative analyses of protein distribution in neuronal terminals in

intact brain tissue. Another important observation was that both acute excitatory input, i.e. activation of

NMDA receptors, and sustained alterations of neuronal excitability, as evidenced by expression of either

mNaChBac or Kir2.1[34,35] in DA neurons, affect the nanoscale distribution of DAT in presynaptic varicosities.

We also observed a marked role of presynaptic D2R activity, further suggesting that the nanoscopic

distribution of a membrane protein, such as DAT, is continuously adapted by several mechanisms to the

immediate activity level of the neurons in which it is expressed. It may be considered that the effects of

the pharmacological manipulations on DAT clustering are quantitatively modest. However, the effects are

highly significant, as well as it should be considered that the numerous varicosities included in each

quantification may be very heterogenous and possibly differentially sensitive to treatments. Such

heterogeneity is directly supported by the finding that only a subset of DA terminals appeared active at

any given time[49]. Future efforts should further clarify this interesting issue.

Our data revealed a striking co-localization of DAT and PIP2 nanodomains, as well as a dispersing

effect of NMDA and haloperidol treatment on both types of domains. We also found that the effect of

haloperidol and NMDA on DAT nanodomains not only required opening of VGCCs but also was contingent

on the presence Ca^{2+}. Interestingly, an increase in cytosolic Ca^{2+} was previously shown to release

membrane protein sequestered PIP2, which might indicate a possible PIP2-dependent Ca^{2+}-regulation of

DAT nanodomain localization[50]. Indeed, the anionic lipid is known to form nanodomains that have been

suggested to be important for membrane protein nanoclustering, including STX-1[42]. Also, PIP2 is known

to regulate many cellular processes including ion channel and transporter function[51,52]. For DAT, it was

suggested that PIP2 binds to the DAT N-terminus and induces a structural change that encourages

phosphorylation and a reverse transport mode of DAT critical for amphetamine-induced DA efflux[41,51].

This is interesting in relation to our data that support an inward-facing configuration of nanodomain

localized DAT and thereby presumably a more efflux prone transporter[53]. Of note, our previous study

suggested a dependency on cholesterol for DAT nanodomain distribution[13]. Furthermore, previous data

have indicated that cholesterol binds to DAT and regulates its function by promoting an outward-facing

conformation[54]. A conceivable explanation is that cholesterol serves to compartmentalize the membrane

to enable a nanoclustered co-distribution of DAT and PIP2[55], and accordingly that cholesterol rather

interacts with unclustered, outward-facing DAT. It is even possible that rapid movements of DAT in and

out of nanodomains parallels the transport cycle shifting the transporter back and forth from an outward-

facing, substrate binding, conformation and unclustered distribution to an inward-facing, substrate

releasing (or efflux prone), conformation and clustered distribution. Indeed, our single particle tracking

data (Spt-STORM) show that the mean jump distance for a given particle is $r\sim110$ nm in a 16 ms time fram

(Fig. 5M), suggesting that a single DAT molecule moves 7-14 μm during a single transport cycle of 1-2

sec[39]. Thus, the movements in the membrane are fast enough for DAT molecules to move in and out of

nanodomains during a single transport cycle. Fast movements of DAT have also been supported with

classical single particle tracking techniques[21] although use of large quantum dots for labeling suggested

somewhat slower diffusion speeds[56].

Previous data have indicated that stimulation of D2-autoreceptors of both the D2 and D3 subtype

enhance DAT activity[38,57-59]. Here we provide evidence that in the DA neuronal cultures the D2-

autoreceptor is preoccupied with DA and that this promotes DAT nanoclustering, consistent with such an

enhancement; that is, if D2-autoreceptor activation causes an increase in DAT activity, this would

expectably lead to a shift of the DAT conformational equilibrium towards a more inward-facing

conformation and, hence, a likely shift towards a higher degree of nanodomain localization. However, our

data may challenge that the regulation of DAT by D2-autoreceptors involves a direct interaction as

previously suggested[57,58]. Although the D2-autoreceptor, like DAT, according to our dSTORM analyses, are

distributed into nanodomains, we observed only modest co-localization between the two proteins in the

DA terminals. The nanodomains responded, nonetheless, like DAT to NMDA and haloperidol, suggesting

that excitatory input promotes nanodomain dispersal, while D2 autoreceptor activation promotes

nanoclustering. We speculate that this increased clustering upon activation reflects assembly of the

receptor with intracellular signaling complexes and/or dynamics relating to desensitization and/or

internalization.

DAT nanodomains showed rather little co-localization with nanodomains of STX-1 and clathrin that

importantly also responded differently to NMDA. These differential effects support that the changes we

see for DAT, the D2-autoreceptor and PIP2 most likely reflect true molecular redistributions and unlikely

are the result of non-specific changes. The decrease in STX-1 nanocluster size and increased localization

within dense clusters in response to NMDA is consistent with other studies indicating that breakup of STX-

1 nanodomains, correlating with vesicle release, takes place, while at the same time more STX-1 is being

brought to the sites of release[45,60]. Of interest, the Voronoi tesselated distributions showed that NMDA

slightly increased localization of STX1 to unclustered DAT (i.e. the larger tessellated areas). This suggests

that the previously described direct interaction between DAT and STX-1, which have been proposed to

be important for amphetamine-induced efflux[46], may take place between the unclustered fractions of the

two proteins.

Summarized, by application of single-molecule sensitive super-resolution imaging we show

compelling evidence that the molecular machinery supporting volume transmission mediated by a

neuromodulator like DA is subject to spatial and temporal nanoscale regulation with putative major

impact on synaptic function. The conclusion is substantiated by detailed studies of DAT that according to

our results is subject to remarkable nanoscale regulatory processes governed by excitatory input,

membrane potential, Ca^{2+} and activity of the presynaptic D2 receptor. Taken together, the present results

not only constitute a new framework for further deciphering the nanoscale architecture of presynaptic

release sites and the underlying functional implications for the DA system but also add to the growing

acknowledgment of nanoscopic regulation as a still poorly explored area of fundamental importance for

neuronal signaling.

Methods

Molecular biology, virus generation and virus transduction. The cDNA encoding human DAT (synthetic

gene kindly provided by Dr. Jonathan Javitch, Columbia University, NY, USA[61] was inserted into pmEOS2-

C1 (RRID: Addgene#54510) to generate pmEOS2-hDAT C1 encoding hDAT with mEOS2 fused to the N-

terminus. Plasmids encoding WT hDAT and D421N hDAT (pRC/CMV hDAT and pRC/CMV hDAT D421N)

were generated as described[32]). The cDNA encoding the short transcriptional variant of human DA D2

receptor with an N-terminal FLAG tag (SF-D2R-S) was inserted into mammalian expression vector pcDNA

3.1(+) (Invitrogen)[62]. The Cre-dependent construct, pAAV-CAG-Flex-NES-jRGECO1a-WPREpA, was

obtained from Addgene (RRID: Addgene #100854). The plasmids, pAAV-hSyn-DIO-mNaChBac-T2A-

mKate2-WPREpA and pAAV-hSyn-DIO-Kir2.1-T2A-mKate2-WPREpA, were SLIC cloned[63] from pAAV-EF1α-

F-FLEX-mNaChBac-T2A-tdTomato (RRID: Addgene #60658) and pAAV-EF1α-F-FLEX-Kir2.1-T2A-tdTomato

(RRID: Addgene #60661), respectively, into a Cre-dependent AAV backbone plasmid with a mKate2

fluorophore instead of tdTomato. This was done to make them sufficiently short so that packaging into

an AAV capsid would be feasible (ITR to ITR length<4700 kbp). The pAAV-pTH-iCre-WPREpA construct was

created and cloned in-house and contains a truncated 488bp rat TH promoter encoding Cre recombinase.

In-house generated AAVs were produced using a FuGene6 (Promega) mediated triple plasmid co-

transfection method in HEK293t cells. Three days after transfection, cells were harvested and virus

purified using an adapted Iodixanol gradient purification protocol [64]. Genomic AAV titer was determined

by a PicoGreen-based method as described elsewhere [65] AAV2-retro-pTH-iCre mediated expression of

Cre-dependent jRGECO1 was verified by fluorescence microscopy and no differences were observed

across the different titers used. For transduction, the appropriate virus was applied to primary culture DA

neurons at an approximate concentration of 2*10E9 vg/mL at 1 to 3 days after preparation of the

neuronal cultures.

Primary cultures of DA neurons with cortical glia cells. Cultured glial cells were obtained from cortex and

cultured DA neurons were obtained from the midbrain of P1-P3 Wistar rats (Charles River, Germany).

Gender was mixed as the cultures were generated from a mixed sex populations of rat pups. For

preparation of cultured glial cells, brains were removed from euthanized P1-P3 pups and placed in

phosphate-buffered saline (PBS) on ice. The cortex of each brain was isolated and cut into millimeter sized

sections and collected in cold dissection medium (Hanks' Balanced Salt Solution with added Sodium

Pyruvate (1 mM), Penicillin-Streptomycin (Sigma P0781, 200 U/L), 10 mM HEPES, and 0.54% glucose)).

After collecting all, the dissection medium was replaced with glia cell medium (DMEM with HEPES "1965",

FBS (10%), Penicillin-Streptomycin (60 U/L)). The tissue was mechanically titrated through pipetting up

and down using a flame-polished Pasteur pipette followed by straining through a 70 μm cell trainer and

centrifugation for 10 min at 200-300 x g. The pellet was resuspended in 10 mL glia medium and strained

again before a second centrifugation for 10 min at 200-300 g and removal of the supernatant. The pellet

was resuspended and seeded into large T175 flasks at approximately 1.5 brains per flask with added glial

medium to a total of 20 mL. Glia cells were grown in an incubator at 37∘ C with 5% CO_2 until they were

70% confluent. The cells were either frozen in heat-inactivated serum containing 10% DMSO for later use

or applied to ethanol-cleaned and poly-D-lysine-coated coverslips and incubated for 1 week prior to DA

neuron addition. On the day before DA neuron addition, the glial cells were changed to DA neuron

medium (Neurobasal A with heat-inactivated FBS (1/100), GlutaMAX (1/100) B-27 (1/50), ascorbic acid

(200 μM), Penicillin-Streptomycin (60 U/L), kynurenic acid (0.5 mM)). Midbrain DA neurons were

generated using a protocol modified from[36]. Brains from rat pups (P1-P3) were isolated and placed in PBS

on ice. Midbrain sections were retrieved from each brain as the brains were cut while in dissection buffer.

The midbrain sections were transferred to papain solution (cysteine (1 mM), DNase (0.1 mg/mL), papain

(20 units/mL), NaCl (116 mM), KCl (5.4 mM), $CaCl_2$ (1.9 mM), NaHCO3 (26 mM), $NaH_2PO_4H_2O$ (2 mM),

$MgSO_4$ (1 mM), EDTA (0.5 mM), Glucose (25 mM), kynurenic acid (0.5 mM) in water) for 30 min at 37∘C.

The solution was replaced with prewarmed DA medium and samples were mechanically titrated with

flame polished Pasteur pipettes and the resulting suspension centrifuged at 200-300 x g for 10 min. The

pellet was resuspended in DA neuron medium and applied to each culture. Glia-derived neurotrophic

factor (GDNF) (100ng/mL of culture solution volume) was applied immediately after seeding of the

neurons. Cultures were incubated for 2 to 3 weeks prior to use. All procedures were performed according

to institutional guidelines at the Faculty of Health Sciences, University of Copenhagen.

Transfection of DA neurons. Primary cultures of DA neurons were transfected using magnetotransfection

at ~14 DIV[66]. NeuroMag reagent (OZ biosciences) (8 µL) was mixed with plasmid (4 µg) and DA neuron

medium (200 µL). The mixture was incubated for 20 min at RT before it was added to cultures grown in

6-well plates. The culture was positioned on top of a magnetic plate for 15 min after which the culture

before it was moved back to the cell incubator and used for immunostaining 24-48 hours later.

Immunostaining of primary cultures of DA neurons for dSTORM and PALM. The cell samples were fixed

in paraformaldehyde (3%) and washed three times in glycine (20 mM) and NH_4Cl (50 mM) in PBS.

Subsequently, the cells were washed in blocking buffer (5% Donkey serum, 1% BSA in PBS), and incubated

in blocking permeabilization buffer (blocking buffer with saponin (0.2%)). Primary antibody was applied

in blocking buffer for 60 min, followed by 3–5 min incubation in blocking buffer. Secondary antibody was

applied in blocking buffer for 45 min, and the sample was incubated 2x for 5 min in blocking buffer.

Samples were washed in PBS twice and post-fixated in paraformaldehyde (3%) for 15 min. Samples were

washed twice in glycine (20 mM) and NH_4Cl (50 mM) in PBS and stored in PBS at 4°C until imaging.

Mouse brain slices. Mouse brain slices were obtained from male C57BL/6 mice 12-16 weeks old. The

slices were prepared as described below. All procedures were performed according to institutional

guidelines at the Faculty of Health Sciences, University of Copenhagen.

Preparation and immunostaining of mouse brain slices for dSTORM. Mice were anesthetized with

isoflurane and perfused first with PBS followed by 4% paraformaldehyde (PFA) injected directly into the

heart. Brains were then placed in 4% PFA for 24 hours, followed by sucrose (30%) for 24 to 48 hours.

Subsequently, brains were snap-frozen on powdered dry ice and stored at -80°C. The brains were sliced

using a cryostat (Leica CM3050 S) to obtain 10 μm striatal coronal sections and each slice was immediately

fused to a 3-aminopropyltriethoxysilane coated coverslip and let dry at room temperature before being

stored in antifreeze (50% glycerol 50% PBS) at -20 ∘C. For immunostaining, brain sections were first rinsed

3x in PBS and subsequently washed 2x with 15 min incubation in glycine (20 mM) and NH_4Cl (50 mM) in

PBS. This was followed by 30 min wash in 10 mM trisodium citrate, pH 6.0, preheated to 80∘ C, 4x wash

in PBS and then blocking/permeabilization for 30 min in PBS with 5% donkey serum, 1% BSA and 0.3 %

Triton X-100. Primary antibody was applied overnight in blocking/permeabilization buffer followed by

washing in 5/5/90 min increments and application of secondary antibody overnight in

blocking/permeabilization buffer. Sections were again washed in 5/5/90 minute increments in

blocking/permeabilization buffer and then with 5/5 min increments with PBS. The sections were post-

fixated in paraformaldehyde (3%) and washed twice for 15 min in glycine (20 mM) and NH_4Cl (50 mM) in

PBS. Samples were lastly washed twice in PBS and stored in PBS at 4 ∘C until imaging.

Antibodies and labeling. Secondary antibodies (donkey anti-rat, Jackson 712-005-153, RRID:AB_2340631;

donkey anti-mouse, Jackson 715-005-151, RRID:AB_2340759); or donkey anti-rabbit, Jackson 711-005-

152, RRID:AB_2340585) were labeled with either NHS-ester conjugated Alexa647 or CF™568 (Biotium,

Freemont, CA) and then isolated through Zeba™ Spin Desalting Columns, 40K MWCO (ThermoFisher

87767). This was completed by first washing the column 3x with 300 µL of 100 mM NaHCO$_3$ in PBS.

Following the washes, 50 µg of antibody in 100 µL of PBS was added and spun at 1500 x g for 2 mins. The

flow-through was collected and combined with a 5-fold molar excess of NHS ester fluorophore. This

solution was incubated in the dark at RT, shaking, for 2.5 hours. A new spin column was then washed 3

times with NaN--3 (0.02%) in PBS. The antibody sample with the dye was then added to the column and

centrifuged for 2 min at 1500 x g. The protein concentration and label concentration of the final product

were measured on a NanoDrop 2000 Spectrophotometer (ThermoFisher). Labeled antibodies had

between 1 and 2 fluorophores attached.

We used the following antibodies and fluorophores for our imaging analysis:

Antibody Target	Product Number	Experiment	Dilution	Secondary antibody	Fluorophore
DAT	Millipore MAB369 RRID:AB_2190413	Primary culture dSTORM	1/200	5 µg/mL Donkey anti-Rat	Alexa647
DAT	Millipore MAB369 RRID:AB_2190413	IHC dSTORM	1/200	10 µg/mL Donkey anti-Rat	Alexa647
DAT	Millipore MAB369 RRID:AB_2190413	Expansion (ExM)	1/200	5 µg/mL Donkey anti-Rat	Atto 643N
TH	Millipore MAB318 RRID:AB_2201528	PALM/dSTORM	1/500	5 µg/mL Donkey anti-Mouse	Alexa647
STX-1	Synaptic Systems 110 011 RRID:AB_887844	Primary culture dSTORM	1/500	5 µg/mL Donkey anti-Mouse	CF568
Clathrin light chain	Synaptic Systems 113 011 RRID:AB_887706	Primary culture dSTORM	1/500	5 µg/mL Donkey anti-Mouse	CF568
PIP2	Abcam ab11039 RRID:AB_442848	Primary culture dSTORM	1/500	5 µg/mL Donkey anti-mouse	CF568
D2R	Synaptic Systems 376 203 RRID:AB_2636918	Primary culture dSTORM	1/500	5 µg/mL Donkey anti-rabbit	CF568
mKate2	Evrogen AB233 RRID:AB_2571743	Primary culture dSTORM	1/500	5 µg/mL Donkey anti-rabbit	CF568

477
478

Expansion Microscopy (ExM). For ExM, perfused brains were acquired by injecting first PBS and then 4%

PFA into the heart of anesthetized mice. Brains were postfixated in 4% PFA for 24 hours and immersed in

sucrose solution (30%) for 24 to 48 hours. Coronal brain slices (40 µm) were obtained with a cryostat. The

brain slices were stored, free floating, in antifreeze (50% glycerol 50% PBS) at -20◦C until use. Brain slices

were washed with PBS three times, and then permeabilized in blocking buffer (5% donkey serum, 1%

bovine serum albumin, 0.1% Triton X-100, in PBS). Brain slices were incubated overnight with primary

antibody in blocking buffer (5% donkey serum, 1% bovine serum albumin, 0.1% Triton X-100, in PBS, pH

7.4). Slices were washed 4x in blocking buffer with 30 min increments and incubated overnight with

secondary antibody in blocking buffer at 4◦C. Slices were again washed 4x in blocking buffer with 30 min

increments. Anchoring treatment was performed with Acryloyl-X-SE (0.1 mg/mL) in PBS at room

temperature overnight[18]. Gel solution was made by mixing N,N-dimethylacrylamide (1.335 g) and sodium

acrylate (0.32g) in 2.85 g of water. Oxygen was purged with nitrogen gas for 40 min. 2.7 mL of the gelling

solution were mixed with 0.3 mL 0.036 g/mL solution of potassium persulfate ($K_2S_2O_8$) solution and purged

of oxygen with nitrogen gas for 15 min on ice. Brain slices were transferred to the extra gelling solution

and placed flat on poly-D-Lysine coated glass slides. Excess gelling solution was removed, and glass

coverslips were placed on the edges of each slice to limit gel dispersion. All work from this point was

done in a cold room (4◦C). Aliquots of 500 µL of polymerization solution were combined with 2 µL of

TEMED was added and the solution vortexed before applying 80 µL to brain slices. A layer of parafilm

covered coverslip was placed overtop of the gel to fully surround the gel. The sample was kept in the cold

room for 15 min before being brought back out to room temperature. The container with the gels was

filled with nitrogen and wet paper to keep humidity and made airtight. Following gelation, samples were

carefully cut out from the chamber in a sim-card shape to ensure orientation. Gels were measured in size

and placed in digestion buffer overnight at room temperature (50mM TRIS, 800 mM guanidine HCl, 2 mM

CaCl2, 0.5% Triton X-100, proteinase K 8U/mL, pH 8.0). The gels were placed into individual dishes and

dialyzed with water many times over a day until they no longer expanded. Imaging was performed the

next day on a spinning disk confocal microscope (FEI CorrSight utilizing a C-Apochromat 63x/1.20 W Corr

M27 objective). Samples were places dry on poly-D-Lysine coated chambers and water was added to the

top of the gel just enough so the gel was hydrated but not so it would move around.

dSTORM. For dSTORM we used a buffer containing β-mercaptoehtanol and an enzymatic oxygen

scavenger system (10% (w/V) glucose, 1% (V/V) beta-mercaptoethanol, 2 mM cyclooctatetraene, 50 mM

Tris-HCl (pH 8), 10 mM NaCl, 34 µg mL^{-1} catalase, 28 µg mL^{-1} glucose oxidase). The imaging was performed

with an ECLIPSE Ti-E epifluoresence/TIRF microscope (NIKON, Japan) equipped with 405 nm, 488 nm 561,

and 647 nm lasers (Coherent, California, USA). All lasers are individually shuttered and collected in a single

fiber to the sample through a 1.49 NA, 100x, apochromat TIRF oil objective (NIKON). Single color imaging

was done with a dichroic mirror (z405/488/561/647 rpc) and the emitted light was filtered by a 710/80

nm bandpass filter. For dual-color dSTORM, we used a dichroic mirror with the range 350–412, 485–490,

558–564, and 637–660 nm (97,335 QUAD C-NSTORM C156921). The excitation light was filtered at the

wavelengths: 401 ± 24 nm, 488 ± 15 nm, 561 ±15 nm, 647 ± 24 nm. The emitted light was filtered at the

wavelengths: 425-475, 505-545, 578-625, and 664-787 nm, and secondly by an extra filter to decrease

noise (561 nm Longpass, Edge Basic, F76-561, AHF). A motorized piezo stage controlled by a near-infrared

light-adjusted perfect focus system (NIKON) is applied to the system to reduce any sample drift over time

in the z-direction. Single-color dSTORM images were constructed from 30,000 frames taken at a 16 ms

frame rate. Dual-color dSTORM images were constructed from 20,000 frames for each color taken at a 16

ms frame rate with each color alternating by frame. Photons were collected with an iXon3 897 EM-CCD

camera camera (Andor, United Kingdom). Laser powers used were 2.3 kW cm −2 for 647 nm, 1.0 kW cm−2

for 488 nm and for 561 nm. The 405 nm laser was used to incrementally increase blinking behavior at

power <0.1 kW cm−2. For 3D-dSTORM, a cylindrical lens was placed before the camera to impart

astigmatism. To verify the multicolor alignment of our microscope, dSTORM images of 100 nm Tetraspek

beads were taken. Localizations were fit to widefield illuminated Tetraspec beads and processed as our

dSTORM data is processed (Supplementary Fig. 7).

PALM. For PALM imaging of mEos-DAT, we obtained 5,000 consecutive frames with a frame rate of 33 Hz

to construct one image. The 488 nm excitation laser was held constant, 0.4 kW cm^{-2}, during the capture

of the image while the 405 nm activation laser was gradually increased to <0.1 kW cm^{-2}.

Localization fitting of dSTORM data. 2D localizations were fit with ThunderSTORM using local maximum

detection with a threshold of 1.5 *std(Wave.F1) and estimator of an integrated gaussian PSF. Sigma was

1.6, fit radius was 3 pixels, and weighted least squares was the method to fit the estimated gaussian to

the PSF[67]. 3D localization fitting was performed with fit3Dcspline[44]. 3D calibrations were made on 100 nm

TetraSpeck beads attached to a coverslip by first placing 50 µL of MgCl2 (1M) on the coverslip and then

adding 500 L of water with 0.9 µL of TetraSpeck beads to the coverslip. Drift for all data was corrected

with redundant cross correlation through Matlab[68]. Localizations were filtered for uncertainty being less

than 25 nm and then merged for localizations that were detected within 15 nm and 3 frames.

Data processing pipeline – dSTORM. To perform necessary localization fitting on the 1349 dSTORM

images included in this study, an in-house data processing pipeline was developed to streamline

operations. Localizations were fit to the raw data with the aforementioned settings in ThunderSTORM

through a self-written ImageJ macro. Although ThunderSTORM can perform the remaining processing

steps, a custom python script (utilizing Matlab functions) was written to perform cross correlation,

merging and filtering so that redundant cross correlation could be employed. The images were then

blinded by labeling them with a random integer and saving the cypher elsewhere. Through a separate

custom Matlab script, varicosities were selected from each image though a freehand selection.

Varicosities were identified by the local swelling of their structure compared to the rest of the extension.

We next used DBSCAN (density based spatial clustering of applications with noise)[13,24] to identify clusters

by size or for determining localizations in dense clusters through the use of a python script employing the

DBSCAN method from the sklearn python library. Voronoi tessellation [25] was employed through Matlab.

The Voronoi based density association algorithm was also written in house and using Matlab.

Varicosity identification in dSTORM DA neuron culture data. Automatic varicosity identification could

not discern between extension and varicosity, so these were selected through a freehand ROI taken of

blinded files. Localization data for all images in a given dataset were blinded by being randomly assigned

a number that was stored on a cypher. ROIs were then gathered without the knowledge of the image

identity, with this information restored in the analysis of the ROIs.

Varicosity identification in brain slice dSTORM data. As this was a computationally intensive process,

each image was divided into 100 quadrants in a 10x10 grid. Within each quadrant, the localizations were

sub-sampled down to 1000 random localizations, and the Voronoi tessalation was obtained for that

section. Tesselated areas were filtered for having an area less than 15000 nm^2. Tesselated shapes that

were touching one another were merged together. Shapes were now filtered for possessing an area

greater than 300000 nm^2. These ROIS were buffered by 50 nm along each edge to smooth the ROI.

Clustering analysis of dSTORM data. Cluster size was identified by performing DBSCAN on each varicosity

for a radius of 15 nm and 5 localizations being the cluster qualifier. The convex hull was gathered for each

cluster to find its area, and the fraction of clusters with a diameter >75 nm was used for the cluster size

analysis. Cluster size distributions were not shown as clusters reaching the detection limit of this method

are not accurate and cause a false perception on the cluster size distribution. Localizations in dense

clusters were found by applying DBSCAN to each varicosity with a radius of 50 nm and 80 localizations

being the qualifier. Voronoi tessellation was applied to each varicosity and areas were filtered for sizes

between 5 nm2 and 1000 nm2. Probability density functions were created for these distributions.

Density Association. Density association was found by performing Voronoi tessellation on target A, and

finding all points in target B that were 25 nm from any point in target A. Each of these target B points that

possessed A association were mapped to the Voronoi tessellated area from target A they resided in. The

probability density functions for target A and the areas from target A that had an associated target B were

created. This procedure was applied to compare the dSTORM signal of DG3_63 to mEOS2-DAT, STX1 to

DAT, D2R to DAT, PIP2 to DAT, and clathrin to DAT.

Quantification of DAT molecules in nanodomains. Primary DA neurons were labeled and imaged for

dSTORM with the DAT MAB369 antibody at dilutions 1:200, 1:400, 1:800, 1:1600, 1:3200. Varicosities

were manually selected from blinded data files. Clusters of localizations potentially arising from single

localizations were identified in the varicosities found in the 1:3200 dilution data set through DBSCAN with

the parameters [ε = 30 nm, nps = 2]. A histogram was made comparing the number of localizations found

per cluster from this cluster analysis, and 3 localizations was determined to be the estimate for the

number of localizations arising from a single monoclonal antibody in this image set. Clusters were then

found in the remaining dataset through the same method defined to identify cluster size. A histogram

was made to show the number of localizations present within the clusters found with a diameter >75 nm.

The peak of this histogram was ~100 localizations, suggesting there are roughly 33 primary antibodies

labeling DAT within a given cluster/nanodomain of this size.

Varied label density cluster verification. Varicosities used for the counting experiment were subject to

the varied label density cluster verification test[28]. Sig for gaussians was set to 15 nm and the cut gauss at

x times sig was set to 2. The threshold selected was 2.

Labeling and imaging primary cultures with DG3-63. Live primary cultures were labeled with DG3-63 (20

nM) [21] for 10 min in the dark at room temperature and washed 3 times with aCSF (NaCl (120 mM), KCl (5

mM), CaCl$_2$ (2 mM), MgCl$_2$ (2 mM), 1 mM NaH$_2$PO$_4$, HEPES (25 mM), glucose (30 mM), pH 7.4). Samples

were fixed with paraformaldehyde (3%) for 15 min followed by 3 washes with glycine (20 mM) and NH$_4$Cl

(50 mM) in PBS. Samples were imaged in normal dSTORM buffer.

Heterologous cell culture. Cath.a-differentiated (CAD) cells[69] (ATCC CRL-11179 or Sigma-Aldrich

08100805) originated from mouse (B6/D2 F1 hybrid) catecholaminergic neuronal tumor. The cells were

maintained in a 1:1 mixture of DMEM 1965 and Ham's F-12 medium (Invitrogen), both supplemented

with 10% FBS and 0.01 mg/mL gentamicin. HEK293 cells (human embryonic kidney cells, ATCC CRL-1573)

were grown as described[32].

Transfection of CAD cells and labeling with DG3-63. CAD cells were transfected with pmEOS2-hDAT

encoding mEOS-DAT or with pRC/CMV-hDAT or pRC/CMV-hDAT D421N encoding WT DAT and the DAT

mutant D421N, respectively[32]. The CAD cells (~1 million cells) were seeded the day before transfection in

a 25 cm^2 flask and washed on the day of transfection before addition of 5 mL of medium. For the

transfection, 1 µg of DNA was added to 100 µL of optiMEM as well as 3 µL Lipofectamine (Invitrogen) to

a separate 100 µL of optiMEM. After 5 min, the two solutions were mixed and incubated for 30 min before

the total mixture was added to the cells. The next day the cells were harvested and seeded on poly-D-

Lysine coated coverslips (100,000 cells per coverslip). For DG3-63 labeling experiments, the media was

removed the next day from the mEOS-DAT expressing cells and 1 mL of aCSF was added (NaCl (120 mM),

KCl (5 mM), CaCl$_2$ (2 mM), MgCl$_2$ (2 mM), HEPES (25 mM), glucose (30 mM), pH 7.4) before incubation for

10 min with or without 100 μM unlabeled cocaine. Following this, the cells were labeled with 10 nM DG3-

63 [21] for 10 min in aCSF. Samples were washed 3 times with aCSF and then fixed for 15 min in

paraformaldehyde (3%) followed by 2x wash with glycine (20 mM) and NH$_4$Cl (50 mM) in PBS. Samples

were imaged in dSTORM buffer as described above. For dSTORM of hDAT WT and D421N, cells the cover

slips samples were stained in the same fashion as primary DA cultures.

Pharmacological treatment of acute brain slices. Mice were euthanized with isoflurane and the live

brains were removed. Coronal sections, 300 μm thick, were cut with a vibratome in cold aCSF (NaCl (119

mM), KCl (2.5 mM), NaH$_2$PO$_4$ (24 mM), Glucose (12.5 mM), CaCl$_2$ (2 mM), MgCl$_2$ (2 mM)) that was being

aerated with carbogen gas. Brain slices were transferred to aCSF and maintained for 1 hour at room

temperature under constant carbogenation. Slices were then transferred to either aCSF or aCSF with AP5

(100 μM), both at 37∘ C for 10 min. Slices from normal aCSF were moved to aCSF with or without NMDA

(20 μM). The NMDA containing aCSF had reduced magnesium (0.3 mM MgCl2) and added glycine (100

μM). The samples from AP5 were transferred to aCSF with AP5 (100 μM) and NMDA (20μM), reduced

magnesium (0.3 mM MgCl$_2$), and added glycine (100 μM). These incubated for 5 min, and then the slices

were transferred to paraformaldehyde (3%) overnight at 4∘ C. Slices were transferred to a 30% sucrose

solution for 2 days a 4∘C and then placed into plastic molds and covered with Tissue-Tek O.C.T. and stored

at -80∘C. Samples were sliced at 10 μm on the cryostat and stained for dSTORM.

Pharmacological treatment of primary cultures. In general, treatments were performed by replacing the

media at room temperature with aCSF (NaCl (120 mM), KCl (5 mM), CaCl$_2$ (2 mM), MgCl$_2$ (2 mM), 1 mM

NaH$_2$PO$_4$, HEPES (25 mM), glucose (30 mM), pH 7.4) followed by stimulation with indicated compounds.

For the NMDA or haloperidol experiments with testing of VGCC inhibitors, cells were treated for 10 min

with either control aCSF, AP5 (100 μM) or quinpirole (50 μM), ω-conotoxin (1 μM), ω-agatoxin (1 μM).

Samples were then treated for 5 min with these same concentrations as cotreatment with either NMDA

(20 μM) or haloperidol (10 nM). NMDA treatments were done in aCSF with added glycine (100 μM). All

experiments using NMDA were done with aCSF with reduced magnesium (0.3 mM MgCl$_2$) for all

conditions. For the BAPTA-AM experiment, the pretreatment with BAPTA-AM (25 μM) was for 30 min

prior to the 5-min NMDA (20 μM) treatment.

SPT-dSTORM. Live primary cultures were labeled with DG3-80 (20 nM) for 10 min and then washed 3

times with aCSF (NaCl (120 mM), KCl (5 mM), CaCl$_2$ (2 mM), MgCl$_2$ (2 mM – 0.3 mM for NMDA related

experiments), 1 mM NaH$_2$PO$_4$, HEPES (25 mM), glucose (30 mM), pH 7.4). A suitable location to image

was found while the sample was kept in aCSF, and then SPTdSTORM imaging was done in a modified

dSTORM buffer (560 μg/mL glucose oxidase, 34 μg/mL catalase, 1.4 μL/mL β-Mercaptoethanol in aCSF).

Pharmacological treatments were done 5 min before imaging with either 20 μM NMDA with 100 μM

glycine, 200 nM haloperidol, or 1 μM UH-232. AP5 (100 μM) was applied for 10 min prior to cotreatment

with NMDA. Imaging was performed at room temperature on the NIKON TiE Eclipse TIRF microscope

described under dSTORM. Light with a wavelength of 561 nm at 0.3 kW cm^{-2} was utilized to promote

blinking of the fluorophores. Images were collected with 16 ms frame rate. Localizations were fit using

ThunderStorm as described above. Single molecule trajectories were found using the program swift [70].

Parameter used beyond the default were: "expdisplacement" : 300,"expnoiserate":

10,"maxblinkingduration" : 2,"pblink" : 0.5,"preappear" : 0.5,"pswitch" : 0.001,"precision":

30,"pruningbase" : 2,"pruningrate" : 0.2,"randomseed" : 42,"wdiffusion" : 2,"wdirected" : 1,"wfield" :

0,"wimmobile" : 1.

Electrophysiology recordings of DA neurons. DA neuronal cultures were transduced after 3 DIV (days in

vitro) with pAAV-pTH-iCre-WPREpA plus either pAAV-hSyn-DIO-mNaChBac-T2A-mKate2-WPREpA or

pAAV-hSyn-DIO-Kir2.1-T2A-mKate2-WPREpA. Electrophysiological recording by the whole-cell patch

clamp technique were done 2-3 weeks after transduction. The DA neurons expressing either mNaChBac

or Kir2.1 were identified based on mKate2 fluorescence using an upright microscope (Olympus BX51WI).

For the recordings, the cultures were submerged in circulating, heated, and oxygenated aCSF (in mM;

NaCl 125, KCl 2.5, $NaHCO_3$ 26, $CaCl_2$ 2, $MgCl_2$ 1, NaH_2PO_4 1.25, Glucose 25; 2 ml/min, 35°C). The aCSF was

supplemented with NBQX (20 μM). The patch glass electrode was filled with (in mM) K-gluconate 122,

Na_2-ATP 5, $MgCl_2$ 2.5, $CaCl_2$ 0.0003, Mg-Gluconate 5.6, K-Hepes 5, H-Hepes 5, and EGTA 1, and had tip

resistance of 4-6 MΩ. After break-in, the cells were held at 0 pA in current clamp mode to measure the

pacemaker activity. Recordings were acquired with a Multiclamp 700B amplifier and 1440A Digitizer.

Calcium Imaging of DA neurons. Primary DA neurons were prepared as described above but instead of

culturing the neurons on glass coverslips, they were cultured on MatTek 35 mm 14 mm glass microwell

(no 1.5) glass bottom dishes (MatTek). To enable measurement of Ca^{2+}-fluctuations specifically in DA

neurons, the neurons were transduced with an AAV encoding double-floxed jRGECO1a (AAV-CAG-Flex-

NES-jRGECO1a-WPREpA) together with an AAV encoding Cre recombinase under control of a truncated

TH promoter (AAV2-retro-pTH-iCre-WPREpA). The cell medium was changed to aCSF (NaCl (120 mM), KCl

(5 mM), CaCl2 (2 mM), $MgCl_2$ (2 mM), 1 mM NaH_2PO_4, HEPES (25 mM), glucose (30 mM), pH 7.4)

immediately prior to imaging. Imaging took place at room temperature on the NIKON TiE Eclipse TIRF

microscope described above. Once a suitable neuron was located, the sample was imaged with 561 nm

light (10 W cm^{-2}). Videos were collected with a 16 ms frame rate. Haloperidol was added mid-video so

that the sample had a total concentration of 10 nM Haloperidol. Data was collected from three

independent experiments. Two varicosities were identified per image.

RNA-seq mining and data analysis. All RNA sequencing data was accessed via the Gene Expression

Omnibus (GEO) under the following accession numbers: GSE108020, GSE115070, GSE76381, GSE116470,

GSE116138. Raw counts were converted to RPKM and quantile normalized using the preprocessCore

(3.11) package in R.

Statistical Analysis. Statistical details can be found in the figure legends and were computed with

GraphPad Prism 8. When employing one-way ANOVA, Dunnett's post hoc test was used for multiple

comparisons. The dSTORM comparisons were normalized to the control sample from the individual

imaging session. This was done to control for unavoidable variability in sample preparations and between

imaging sessions.

Experiment	Number of sample preparations	Number of dSTORM Images	Number of varicosities (neuron experiments) or cells (CAD cell experiment)
IHC dSTORM NMDA	3	102	4028
CAD Cell D421N	3	55	80
Membrane Potential	3	108	471
NMDA and VGCC	3	188	841
Haloperidol and VGCC	3	180	840
BAPTA-AM	3	95	208
STX1 NMDA	4	162	678
Clathrin NMDA	3	90	304
PIP2 Haloperidol	3	84	309
PIP2 NMDA	3	88	384
D2 NMDA	3	101	344

D2 Haloperidol	3	90	524

707 **Data Availability.** Custom written codes used for the data analysis can be found at:

708 https://github.com/GetherLab/Super-Resolution-Data-Analysis. . The datasets supporting the current

709 study have not been deposited in a public repository because of the file size but are available from the

710 corresponding author on request.

Lycas et al.

REFERENCES

Groc, L. & Choquet, D. Linking glutamate receptor movements and synapse function. *Science* **368** (2020).

Choquet, D. & Hosy, E. AMPA receptor nanoscale dynamic organization and synaptic plasticities. *Curr Opin Neurobiol* **63**, 137-145 (2020).

Liu, C. & Kaeser, P. S. Mechanisms and regulation of dopamine release. *Curr Opin Neurobiol* **57**, 46-53 (2019).

Bjorklund, A. & Dunnett, S. B. Dopamine neuron systems in the brain: an update. *Trends Neurosci* **30**, 194-202 (2007).

Tritsch, N. X. & Sabatini, B. L. Dopaminergic modulation of synaptic transmission in cortex and striatum. *Neuron* **76**, 33-50 (2012).

Descarries, L., Watkins, K. C., Garcia, S., Bosler, O. & Doucet, G. Dual character, asynaptic and synaptic, of the dopamine innervation in adult rat neostriatum: a quantitative autoradiographic and immunocytochemical analysis. *J Comp Neurol* **375**, 167-186 (1996).

Giguere, N. *et al.* Increased vulnerability of nigral dopamine neurons after expansion of their axonal arborization size through D2 dopamine receptor conditional knockout. *PLoS Genet* **15**, e1008352 (2019).

Borroto-Escuela, D. O. *et al.* Brain Dopamine Transmission in Health and Parkinson's Disease: Modulation of Synaptic Transmission and Plasticity Through Volume Transmission and Dopamine Heteroreceptors. *Front Synaptic Neurosci* **10**, 20 (2018).

Kristensen, A. S. *et al.* SLC6 neurotransmitter transporters: structure, function, and regulation. *Pharmacol Rev* **63**, 585-640 (2011).

Torres, G. E. & Amara, S. G. Glutamate and monoamine transporters: new visions of form and function. *Curr Opin Neurobiol* **17**, 304-312 (2007).

Bermingham, D. P. & Blakely, R. D. Kinase-dependent Regulation of Monoamine Neurotransmitter Transporters. *Pharmacol Rev* **68**, 888-953 (2016).

Torres, G. E. The dopamine transporter proteome. *J Neurochem* **97 Suppl 1**, 3-10 (2006).

Rahbek-Clemmensen, T. *et al.* Super-resolution microscopy reveals functional organization of dopamine transporters into cholesterol and neuronal activity-dependent nanodomains. *Nat Commun* **8**, 740 (2017).

Betzig, E. *et al.* Imaging intracellular fluorescent proteins at nanometer resolution. *Science* **313**, 1642-1645 (2006).

Heilemann, M. *et al.* Subdiffraction-resolution fluorescence imaging with conventional fluorescent probes. *Angew Chem Int Ed Engl* **47**, 6172-6176 (2008).

16   Huang, B., Bates, M. & Zhuang, X. Super-resolution fluorescence microscopy. *Annu Rev Biochem*
78, 993-1016 (2009).

17   Tillberg, P. W. *et al.* Protein-retention expansion microscopy of cells and tissues labeled using
standard fluorescent proteins and antibodies. *Nat Biotechnol* **34**, 987-992 (2016).

18   Truckenbrodt, S., Sommer, C., Rizzoli, S. O. & Danzl, J. G. A practical guide to optimization in X10
expansion microscopy. *Nat Protoc* **14**, 832-863 (2019).

19   Chen, F., Tillberg, P. W. & Boyden, E. S. Optical imaging. Expansion microscopy. *Science* **347**,
543-548 (2015).

20   Manley, S. *et al.* High-density mapping of single-molecule trajectories with photoactivated
localization microscopy. *Nat Methods* **5**, 155-157 (2008).

21   Guthrie, D. A. *et al.* Novel Fluorescent Ligands Enable Single-Molecule Localization Microscopy
of the Dopamine Transporter. *ACS Chem Neurosci* **11**, 3288-3300 (2020).

22   Baker, S. M., Buckheit, R. W., 3rd & Falk, M. M. Green-to-red photoconvertible fluorescent
proteins: tracking cell and protein dynamics on standard wide-field mercury arc-based
microscopes. *BMC Cell Biol* **11**, 15 (2010).

23   Shivanandan, A., Deschout, H., Scarselli, M. & Radenovic, A. Challenges in quantitative single
molecule localization microscopy. *FEBS Lett* **588**, 3595-3602 (2014).

24   Ester, M., Kriegel, H.-P., Sander, J. & Xu, X. in *Proc. 2nd International Conference on Knowledge*
Discovery and Data Mining   226–231 (AAAI Press, 1996).

25   Levet, F. *et al.* A tessellation-based colocalization analysis approach for single-molecule
localization microscopy. *Nat Commun* **10**, 2379 (2019).

26   Ehmann, N. *et al.* Quantitative super-resolution imaging of Bruchpilot distinguishes active zone
states. *Nat Commun* **5**, 4650 (2014).

27   Siddig, S. *et al.* Super-resolution imaging reveals the nanoscale organization of metabotropic
glutamate receptors at presynaptic active zones. *Science Advances* **6**, eaay7193 (2020).

28   Baumgart, F. *et al.* Varying label density allows artifact-free analysis of membrane-protein
nanoclusters. *Nat Methods* **13**, 661-664 (2016).

29   Newman, A. H. *et al.* Translating the atypical dopamine uptake inhibitor hypothesis toward
therapeutics for treatment of psychostimulant use disorders. *Neuropsychopharmacology* **44**,
1435-1444 (2019).

30   Beuming, T. *et al.* The binding sites for cocaine and dopamine in the dopamine transporter
overlap. *Nat Neurosci* **11**, 780-789 (2008).

31   Lebowitz, J. J. & Khoshbouei, H. Heterogeneity of dopamine release sites in health and
degeneration. *Neurobiol Dis* **134**, 104633 (2020).

32 Herborg, F., Andreassen, T. F., Berlin, F., Loland, C. J. & Gether, U. Neuropsychiatric disease-
associated genetic variants of the dopamine transporter display heterogeneous molecular
phenotypes. *J Biol Chem* **293**, 7250-7262 (2018).

33 Salamone, A. *et al.* Prolonged nicotine exposure down-regulates presynaptic NMDA receptors in
dopaminergic terminals of the rat nucleus accumbens. *Neuropharmacology* **79**, 488-497 (2014).

34 Lin, C.-W. *et al.* Genetically increased cell-intrinsic excitability enhances neuronal integration
into adult brain circuits. *Neuron* **65**, 32-39 (2010).

35 Xue, M., Atallah, B. V. & Scanziani, M. Equalizing excitation-inhibition ratios across visual cortical
neurons. *Nature* **511**, 596-600 (2014).

36 Rayport, S. *et al.* Identified postnatal mesolimbic dopamine neurons in culture: morphology and
electrophysiology. *J Neurosci* **12**, 4264-4280 (1992).

37 Turner, T. J., Adams, M. E. & Dunlap, K. Multiple Ca2+ channel types coexist to regulate
synaptosomal neurotransmitter release. *Proceedings of the National Academy of Sciences of the
United States of America* **90**, 9518-9522 (1993).

38 Beaulieu, J. M. & Gainetdinov, R. R. The physiology, signaling, and pharmacology of dopamine
receptors. *Pharmacol Rev* **63**, 182-217 (2011).

39 Prasad, B. M. & Amara, S. G. The dopamine transporter in mesencephalic cultures is refractory
to physiological changes in membrane voltage. *J Neurosci* **21**, 7561-7567 (2001).

40 Strange, P. G. Antipsychotic drug action: antagonism, inverse agonism or partial agonism. *Trends
Pharmacol Sci* **29**, 314-321 (2008).

41 Belovich, A. N. *et al.* A network of phosphatidylinositol (4,5)-bisphosphate (PIP2) binding sites
on the dopamine transporter regulates amphetamine behavior in Drosophila Melanogaster.
Molecular Psychiatry (2019).

42 van den Bogaart, G. *et al.* Membrane protein sequestering by ionic protein-lipid interactions.
Nature **479**, 552-555 (2011).

43 Maxfield, F. R. & McGraw, T. E. Endocytic recycling. *Nature Reviews Molecular Cell Biology* **5**,
121-132 (2004).

44 Li, Y. M. *et al.* Real-time 3D single-molecule localization using experimental point spread
functions. *Nat. Methods* **15**, 367-+ (2018).

45 Padmanabhan, P. *et al.* Need for speed: Super-resolving the dynamic nanoclustering of syntaxin-
1 at exocytic fusion sites. *Neuropharmacology* **169** (2020).

46 Binda, F. *et al.* Syntaxin 1A interaction with the dopamine transporter promotes amphetamine-
induced dopamine efflux. *Mol Pharmacol* **74**, 1101-1108 (2008).

47   Goncalves, J. *et al.* Nanoscale co-organization and coactivation of AMPAR, NMDAR, and mGluR
at excitatory synapses. *Proc Natl Acad Sci U S A* **117**, 14503-14511 (2020).

48   Das, A. K. *et al.* Dopamine transporter forms stable dimers in the live cell plasma membrane in a
phosphatidylinositol 4,5-bisphosphate-independent manner. *J Biol Chem* **294**, 5632-5642
(2019).

49   Pereira, D. B. *et al.* Fluorescent false neurotransmitter reveals functionally silent dopamine
vesicle clusters in the striatum. *Nat Neurosci* **19**, 578-586 (2016).

50   McLaughlin, S. & Murray, D. Plasma membrane phosphoinositide organization by protein
electrostatics. *Nature* **438** (2005).

51   Hamilton, P. J. *et al.* PIP2 regulates psychostimulant behaviors through its interaction with a
membrane protein. *Nat Chem Biol* **10**, 582-589 (2014).

52   Suh, B. C. & Hille, B. PIP2 is a necessary cofactor for ion channel function: how and why? *Annu
Rev Biophys* **37**, 175-195 (2008).

53   Robertson, S. D., Matthies, H. J. & Galli, A. A closer look at amphetamine-induced reverse
transport and trafficking of the dopamine and norepinephrine transporters. *Mol Neurobiol* **39**,
73-80 (2009).

54   Hong, W. C. & Amara, S. G. Membrane cholesterol modulates the outward facing conformation
of the dopamine transporter and alters cocaine binding. *J Biol Chem* **285**, 32616-32626 (2010).

55   Carquin, M., D'Auria, L., Pollet, H., Bongarzone, E. R. & Tyteca, D. Recent progress on lipid lateral
heterogeneity in plasma membranes: From rafts to submicrometric domains. *Progress in Lipid
Research* **62**, 1-24 (2016).

56   Thal, L. B. *et al.* Single Quantum Dot Imaging Reveals PKCbeta-Dependent Alterations in
Membrane Diffusion and Clustering of an Attention-Deficit Hyperactivity
Disorder/Autism/Bipolar Disorder-Associated Dopamine Transporter Variant. *ACS Chem
Neurosci* **10**, 460-471 (2019).

57   Lee, F. J. *et al.* Dopamine transporter cell surface localization facilitated by a direct interaction
with the dopamine D2 receptor. *Embo J* **26**, 2127-2136 (2007).

58   Bolan, E. A. *et al.* D2 receptors regulate dopamine transporter function via an extracellular
signal-regulated kinases 1 and 2-dependent and phosphoinositide 3 kinase-independent
mechanism. *Mol Pharmacol* **71**, 1222-1232 (2007).

59   Zapata, A. *et al.* Regulation of dopamine transporter function and cell surface expression by D3
dopamine receptors. *J Biol Chem* **282**, 35842-35854 (2007).

60   Maidorn, M., Olichon, A., Rizzoli, S. O. & Opazo, F. Nanobodies reveal an extra-synaptic
population of SNAP-25 and Syntaxin 1A in hippocampal neurons. *Mabs* **11**, 305-321 (2019).

61   Loland, C. J., Granas, C., Javitch, J. A. & Gether, U. Identification of intracellular residues in the
dopamine transporter critical for regulation of transporter conformation and cocaine binding. *J
Biol Chem* **279**, 3228-3238. (2004).

62   Klewe, I. V. *et al.* Recruitment of beta-arrestin2 to the dopamine D2 receptor: insights into anti-
psychotic and anti-parkinsonian drug receptor signaling. *Neuropharmacology* **54**, 1215-1222
(2008).

63   Jeong, J. Y. *et al.* One-step sequence- and ligation-independent cloning as a rapid and versatile
cloning method for functional genomics studies. *Appl Environ Microbiol* **78**, 5440-5443 (2012).

64   Matsui, R., Tanabe, Y. & Watanabe, D. Avian adeno-associated virus vector efficiently
transduces neurons in the embryonic and post-embryonic chicken brain. *PLoS One* **7**, e48730
(2012).

65   Piedra, J. *et al.* Development of a rapid, robust, and universal picogreen-based method to titer
adeno-associated vectors. *Hum Gene Ther Methods* **26**, 35-42 (2015).

66   Underhill, S. M. *et al.* Amphetamine modulates excitatory neurotransmission through
endocytosis of the glutamate transporter EAAT3 in dopamine neurons. *Neuron* **83**, 404-416
(2014).

67   Ovesny, M., Krizek, P., Borkovec, J., Svindrych, Z. & Hagen, G. M. ThunderSTORM: a
comprehensive ImageJ plug-in for PALM and STORM data analysis and super-resolution imaging.
Bioinformatics **30**, 2389-2390 (2014).

68   Wang, Y. *et al.* Localization events-based sample drift correction for localization microscopy
with redundant cross-correlation algorithm. *Opt Express* **22**, 15982-15991 (2014).

69   Qi, Y., Wang, J. K., McMillian, M. & Chikaraishi, D. M. Characterization of a CNS cell line, CAD, in
which morphological differentiation is initiated by serum deprivation. *J Neurosci* **17**, 1217-1225
(1997).

70   Turkowyd, B. *et al.* Establishing Live-Cell Single-Molecule Localization Microscopy Imaging and
Single-Particle Tracking in the Archaeon Haloferax volcanii. *Front Microbiol* **11**, 583010 (2020).

Acknowledgement

We thank Ralph Götz for support with expansion microscopy. The work was supported by the Lundbeck

Foundation grants R266-2017-4331 (UG), R276-2018-792 (UG), R230-2016-3154 (M.D.L.), R181-2014-

3090 (FH), R303-2018-3540 (F.H.) and R231-2016-2481-5 (ATS), Independent Research Fund Denmark –

Medical Sciences (U.G. 7016-00325B), and the NIDA-Intramural Research Program Z1A DA000610 (AHN

and DAG).

Author contributions

M.D.L. performed all the experiments and wrote the analyses not hereafter mentioned; A.L.E. devised

the cluster size algorithm and identified mRNA expression data; A.T.S. performed electrophysiology and

generated the construct encoding Cre recombinase under control of a truncated TH promotor; N.O.H.

performed the D421N CAD cell experiments; S.H.J. cloned Kir2.1 and mNaChBac plasmids and made non-

commercially available viruses; D.A.G. and A.H.N. invented and synthesized the fluorescent cocaine

analogs; J.F.S. optimized the DA neuron culture protocol; M.D.L., F.H., and U.G. conceptualized the study,

designed the research and interpreted data; U.G., F.H., M.S., and C.W. supervised the research. M.D.L.,

U.G., and F.H. wrote the paper with contributions from M.S. and all the other authors.

Competing interests

The authors declare no competing financial interests.

Fig. 1. DAT shows nanodomain distribution in cultured neurons and striatal slices. (A-D) Visualization of mEOS2-DAT overexpressed in cultured DA neurons. Endogenous TH was labeled to identify DA neurons. (A, C) Widefield images (C is boxed region in A). (B, D) PALM/dSTORM images (D is boxed region in B). (E-H) 3D dSTORM image of DAT in DA neurons (color-coded based on depth). (E) Neuronal extension. (F)

Varicosity (boxed region in E) (XY perspective). (G) YZ perspective. (H) XZ perspective. (I-L) Mouse brain
coronal section immunolabeled for DAT. (I) brain slice with striatal DAT staining. (J) Comparison of
widefield with dSTORM. (K) Widefield view of varicosity. (L) dSTORM image. (M-P) 10 x Expansion
Microscopy (ExM) on mouse striatal slice immunolabeled for DAT. N shows region of interest (R.O.I.) in
M with the full 19 μM Z slice (color-coded based on depth); O is R.O.I. in M with just 2 μM of the Z axis.
(P) Single varicosity identified in O. (Q-T) Cluster metrics demonstrated on a varicosity immunolabeled for
DAT. Q, dSTORM of single varicosity. (R, S) Identification of clusters by DBSCAN; clusters with diameter
>75 nm in green, clusters <75 nm in magenta; dense clusters (>80 localizations within a radius of 50 nm)
in red. (T) DAT localizations segmented by Voronoi tessellation.
914
915
916

917
918
Fig. 2. Nanodomain localized DAT is preferentially in an inward-facing conformation. (A-D) dSTORM on
DA neurons and DAT expressing CAD cells labeled with the fluorescent cocaine analogue DG3-63 (10 nM).
(A) Axonal extension. (B) varicosity (box in A). (C) CAD cell. (D) R.O.I. from C. (E-H) The density-based
association of DG3-63 to mEOS2-DAT. (E) Cartoon of labeling strategy. (F) dSTORM/PALM image of CAD

cell expressing mEOS2-DAT (red) and labeled with DG3-63 (10 nM)(blue). (G) Close-up dSTORM/PALM
image of CAD cell expressing mEOS2-DAT (red) and labeled with DG3-63 (10 nM) (blue). (H) The DG3-63
signal is blocked by 100 µM cocaine. (I-K) Voronoi tessellation based colocalization algorithm. Tessellation
is performed on the mEOS2-DAT data and areas are marked for when the mEOS2-DAT localization has a
DG3-63 localization within 25 nm distance. (L) Left, Total mEOS2-DAT Voronoi tesselated area distribution
(red) (probability density function (P.D.F.) and tessellated areas of mEOS2-DAT associated with DG3-63
(blue). Right, tessellated areas for mEOS2-DAT associated with DG3-63 (blue) compared to that associated
in presence of saturating cocaine. (M-O) TIRF/dSTORM of WT DAT and inward-facing mutant (D421N).
(M) Example image of DAT expressed in CAD cells. (N) Example images of clusters with diameter >75 nm
(green) in WT DAT and D421N Dcells. (O) Normalized fraction of clusters with diameter >75 nm (in %),
means ± S.E., from 38 WT DAT cells and 42 D421N cells, 3 transfections, ***p<0.001, unpaired t-test.
(P-O)dSTORM of WT DAT and inward-facing mutant (D421N). (P) Number of localizations per μm^2 forWT
DAT and D421N, means ± S.E., unpaired t test, p= 0.4. (Q) Example image of dense clusters (red) of WT
DAT and D421N (radius 100 nm, >100 localizations). (R) Fraction of localizations in dense clusters (in % of
WT DAT), means ± S.E., from 38 WT DAT cells and 42 D421N cells, 3 transfections, **p<0.01, unpaired t-
test.
939

Lycas et al.

940
941
Fig. 3. NMDA receptor activation declusters DAT in mouse brain slices. (A) Workflow diagram. (B) Data
processing steps to isolate single varicosities from dSTORM images (see Methods). (C) Representative DAT
dSTORM image before, during, and after varicosity identification. (D)Example varicosity from control,

NMDA, and AP5 + NMDA treated slices (clusters with diameter >75 nm in green, clusters <75 nm in
magenta). Right, Normalized fraction of DAT clusters >75 nm (in %), means ± S.E., from 3 experiments
with analysis of 1793 control, 1102 NMDA and 1033 NMDA + AP5 varicosities; one-way ANOVA,
****$p<0.0001$, *$p<0.05$. (E) Example varicosity from control, NMDA (20 µM), and AP5 + NMDA (20 µM)
treated slices (dense clusters in red, >80 localizations, radius 50 nm). Right, Normalized fraction of DAT
localizations in dense clusters (in %), means ± S.E.; one-way ANOVA, ****$p<0.0001$, ***$p<0.001$,
**$p<0.01$. (F) The probability density functions (P.D.F.) for the Voronoi tessellated areas averaged by
varicosity.
953

Fig. 4. DAT nanodomain localization is regulated by neuronal activity in a Ca²⁺ dependent manner. (A-D) Expression of the Na⁺ channel mNaChBac or the K⁺ channel Kir2.1 in DA neurons. Example varicosities from DAT dSTORM of control, mNaChBac or Kir2.1 expressing DA neurons. (A) DAT clusters with diameter

>75 nm in green, clusters <75 nm in magenta. (B) Dense DAT clusters in red (>80 localizations, radius 50
nm). (C) Normalized fraction of clusters >75 nm (in %). (D) Normalized fraction of DAT localizations in
dense clusters (in %), means ± S.E.; data from 3 cultures and 80 control, 230 mNaChBac and 161 Kir2.1
varicosities; one-way ANOVA, ****$p<0.0001$, **$p<0.01$, *$p<0.05$, n.s., not significant. (E-J) The effect of
NMDA on DAT nanodomains is blocked by the N-type VGCC blocker ω-conotoxin. Example varicosities
from dSTORM of DA neurons subject to 5 min treatment with vehicle, NMDA (20 μM), AP5 (100 μM) +
NMDA (20 μM), ω-Conotoxin (1 μM) + NMDA (20 μM) or ω-Agatoxin (1 μM) + NMDA (20 μM). (E) DAT
clusters with diameter >75 nm in green, clusters <75 nm in magenta. (F) Dense DAT clusters in red (>80
localizations, radius 50 nm). (G) localizations segmented by Voronoi tessellation. (H, I) Normalized fraction
of DAT clusters >75 nm (in %) (H) and normalized fraction of DAT localizations in dense clusters (in %) (I),
means ± S.E.; one-way ANOVA, ****$p< <0.0001$, *$p<0.05$, n.s., not significant. (J) Probability density
functions (P.D.F.) (curves color-coded as in the bar diagrams) for the Voronoi tessellated areas averaged
by varicosity. Data are from 3 cultures and 190 control, 147 NMDA, 170 AP5, 180 ω-Conotoxin and 154
ω-Agatoxin varicosities. (K, L) Pretreatment (30 in) with BAPTA-AM inhibits the dispersing effect of NMDA
on DAT nanodomains. (K) Normalized fraction of DAT clusters >75 nm (in %) and (L) normalized fraction
of DAT localizations in dense clusters (in %) for control, NMDA (20 μM) and NMDA (20 μM) + BAPTA-AM
(25 μM), means ± S.E.; data from 3 cultures and includes analysis of 66 control, 72 NMDA and 70 BAPTA-
AM + NMDA varicosities, one-way ANOVA, **$p<0.01$, *$p<0.05$, n.s., not significant. (M) mRNA expression
for P, R, and N-type VGCC in DA neurons based on single cell RNAseq data (see Methods).
978
979

Fig. 5. DAT nanodomain localization is regulated by the D2R and involves changes in lateral mobility. A-
F) Dispersal of DAT nanodomains is promoted by the D2R antagonist haloperidol and blocked by the
agonist quinpirole and by the N-type VGCC blocker ω-conotoxin. Example varicosities from dSTORM of

DA neurons subject 5 min treatment with haloperidol (10 nM), haloperidol (10 nM) + quinpirole (50 µM),
haloperidol (10 nM) + ω-conotoxin (1 µM) or haloperidol (10 nM) + ω-agatoxin (1 µM). (A) DAT clusters
with diameter >75 nm in green, clusters <75 nm in magenta. (B) Dense DAT clusters in red (>80
localizations, radius 50 nm). (C) localizations segmented by Voronoi tessellation. (D, E) Normalized
fraction of DAT clusters >75 nm (in %) (D) and normalized fraction of DAT localizations in dense clusters
(in %) (E), means ± S.E.; one-way ANOVA, ****$p<0.0001$, ***$p<0.001$, **$p<0.01$, *$p<0.05$, n.s., not
significant. (F) Probability density functions (P.D.F.) (curves color-coded as in the bar diagrams) for the
Voronoi tessellated areas averaged by varicosity. Probability density functions (P.D.F.) (curves color-
coded as in the bar diagrams) for the Voronoi tessellated areas averaged by varicosity. Data are from 3
cultures and 153 control, 166 haloperidol, 167 quinpirole, 178 ω-conotoxin and 176 ω-agatoxin
varicosities. (G-R) Single particle tracking dSTORM (SPT-dSTORM) identifies changes in DAT mobility in
response to NMDA and D2R antagonism. (G) Example live dSTORM images of DA neurons labeled with
the fluorescent cocaine analogue, DG3-80. (H) Boxed region in B. (I) Detected tracks and (J) Distribution
of segment lengths in varicosity from control SPT-dSTORM experiment (16 ms frame rate). (K) Example
traces of a confined and an unconstrained freely moving particle (DG3-80 labeled DAT). (L) Fraction of
unconstrained tracks out of the total number of tracks. Data are means of 23 tracks ± S.E. (M) Probability
Density Function (P.D.F.) for total mean jump distances (M.J. D) (i.e. the distance an individual
molecule/particle has traveled between each image) for control experiments. (N) P.D.F. when separated
into static and freely moving particles. Among the freely moving particles, some moved very fast with
M.J.D. of 200-350 nm. (O) NMDA (20 µM, 5 min) increases population of DAT molecules with M.J.D. > 200
nm, which is blunted by AP5 (100 µM). (P) Fraction of fast moving particles (> 200 nm), means ± S.E., data
from 13 control, 4 NMDA and 5 NMDA+AP5 experiments; (Q, R) The D2R antagonists haloperidol (10 nM,
5 min) and UH-232 (1 µM) increase the population of DAT molecules with M.J.D. > 200 nm > 200 nm; M,
Fraction of fast moving particles (> 200 nm), means ± S.E., data are from 10 control, 6 haloperidol and 5
UH-232 experiments; one-way ANOVA, *$p<0.05$; **$p<0.01$, n.s., not significant.
1010
1011

Fig. 6. DAT nanodomains overlap with PIP2 nanodomains. Example dual-color dSTORM images of phosphatidylinositol 4,5-bisphosphate (PIP2) (red, CF568) and DAT (blue, Alexa647) in a neuronal extension from cultured DA neurons (left) with close-up image of varicosity (right) corresponding to boxed region in the left image. Example dual-color dSTORM images of varicosities from DA neurons showing localizations for PIP2 (red) and DAT (blue). Left, Control; Right, neurons exposed to haloperidol (10 nM, 5 min). (C-F) Effect of NMDA and haloperidol on PIP2 clustering. Upper panels, Control, NMDA (20 µM) and NMDA (20 µM) + AP5 (100 µM); Lower panels; Control, haloperidol (10 nM) and haloperidol (10 nM) + quinpirole (50 µM). (C) Normalized fraction of PIP2 localizations in large clusters (>75 nm in diameter) (in %) and (D), normalized fraction of PIP2 localizations in dense clusters (>80 localizations, radius 50 nm) (in %), means ±S.E., one-way ANOVA, ****$p<0.0001$, **$p<0.01$, n.s., not significant. (E) Probability density functions (P.D.F.) (color-coded as in the bar diagrams) for the Voronoi tessellated areas averaged by varicosity. (F) Probability density functions (P.D.F.) (color-coded as in the bar diagrams) of the association of PIP2 with DAT clusters determined by Voronoi tessellation-based association (see Fig. 2) showing strong association PIP2 with DAT. Data are from 3 cultures and 137 control, 126 NMDA, 121 NMDA + AP5 varicosities; and 140 control, 102 haloperidol, 67 haloperidol + quinpirole varicosities,

Fig. 7. The D2R is also distributed into nanodomains. (A) Example dual-color dSTORM images of the D2R (red, CF568) and DAT (blue, Alexa647) in DA neuronal extension (left) with close-up image of varicosity (right) corresponding to boxed region in the left image. (B) Example dual color dSTORM images of varicosities from DA neurons showing localizations for D2R (red) and DAT (blue). Left, Control; Right, neurons exposed to haloperidol (10 nM, 5 min). (C-F) Effect of NMDA and haloperidol on D2R clustering. Upper panels, Control, NMDA (20 µM) and NMDA (20 µM) + AP5 (100 µM); Lower panels; Control, haloperidol (10 nM) and haloperidol (10 nM) + quinpirole (50 µM). (C) Normalized fraction of D2R localizations in large clusters (>75 nm in diameter) (in %) and (D), normalized fraction of D2R localizations in dense clusters (>80 localizations, radius 50 nm) (in %), means ±S.E., one-way ANOVA, ***$p<0.001$, **$p<0.01$, *$p<0.05$, n.s., not significant. (E) Probability density functions (P.D.F.) (color-coded as in the bar diagrams) for the Voronoi tessellated areas averaged by varicosity. (F) Probability density functions (P.D.F.) (color-coded as in the bar diagrams) of the association of D2R with DAT clusters determined by Voronoi tessellation-based association (see Fig. 2). Data are from 3 cultures and data from 3 cultures and 112 control, 157 NMDA, 75 NMDA + AP5; and 179 control, 195 haloperidol and 150 quinpirole varicosities.