A Scalable Screening of *E. coli* Strains for Recombinant Protein Expression

Luana G. Morão[a¶], Lívia R. Manzine[a¶], Angélica Luana C. Barra[a,b], Lívia Oliveira D. Clementino[a], Raíssa F. Gutierrez[a], Carsten Wrenger[c] and Alessandro S. Nascimento[a*].

a. Pólo TerRa, São Carlos Institute of Physics, University of São Paulo. Av. João Dagnone, 1100, Jd. Santa Angelina, 13563-120. São Carlos, SP. Brazil.

b. Institute of Biochemistry and Molecular Biology, Laboratory for Structural Biology of Infection and Inflammation, University of Hamburg, c/o DESY, Build. 22a. Notkestr. 85, 22603 Hamburg, Germany.

c. Unit for Drug Discovery, Department of Parasitology, Institute of Biomedical Sciences, University of São Paulo, Av. Prof. Lineu Prestes 1374, 05508-000 São Paulo-SP, Brazil.

¶ These authors contributed equally to this work.

* Corresponding author: Phone: +55-16-3373-8709. Email: asnascimento@ifsc.usp.br.

Abstract

Structural biology projects are highly dependent on the large-scale expression of
soluble protein and, for this purpose, heterologous expression using bacteria or yeast as
host systems are usually employed. In this scenario, some of the parameters to be
optimized include (*i*) those related to the protein construct, such as the use of a fusion
protein, the choice for an N-terminus fusion/tag or a C-terminus fusion/tag; (*ii*) those
related to the expression stage, such as the concentration and selection of inducer agent
and temperature expression and (*iii*) the choice of the host system, which includes the
selection of a prokaryotic or eukaryotic cell and the adoption of a strain. The optimization
of some of the parameters related to protein expression, stage (*ii*), is straightforward. On
the other hand, the determination of the most suitable parameters related to protein
construction requires a new cycle of gene cloning, while the optimization of the host cell
is less straightforward. Here, we evaluated a scalable approach for the screening of host
cells for protein expression in a structural biology pipeline. We evaluated four
Escherichia coli strains looking for the best yield in soluble protein expression using the
same strategy for protein construction and gene cloning and comparing to our standard
strain, Rosetta (DE3). Using a liquid handling device (robot), *E. coli* pT_GroE,
Lemo21(DE3), Arctics Express (DE3), and Rosetta Gami 2 (DE3) strains were screened
for the maximal yield in soluble protein recovery. For the genes used in this experiment,
the Arctic Express (DE3) strain resulted in better yields of soluble proteins. We propose
that screening of host cell/strain is feasible, even for smaller laboratories and the
experiment as proposed can easily be scalable to a high-throughput approach.

Keywords: Recombinant protein expression, *E. coli* strains, scalable protein expression.

1. Introduction

Many drug discovery projects rely on the structure determination of the drug targets. Indeed, to the best of our knowledge, the so-called 'structure-based rational design' of drug candidates dates to 1976, a year before the Protein Data Bank (PDB) was founded, when Beddell and coworkers designed a series of compounds to fit the human haemoglobin site [1]. Typical structural biology pipelines require pure and soluble protein in high amounts, which is usually achieved by heterologous expression in *Escherichia coli* [2,3] or in yeast cells, such as *Pichia pastoris* or *Saccharomyces cerevisae*, or even in fungus, such as *Aspergillus niger* (e.g., [4]).

E. coli is the host cell of choice in most structural biology approaches, due to the ease in handling, rapid doubling time, and very well-established protocols for expression. In our experience, protein expression in *E. coli* for high-throughput (HT) pipelines resulted in yields up to 946 mg of protein expressed per liter of culture, with an average of about 60-70 mg.L^{-1} for enzymes of the glycoside hydrolase (GH) superfamily [5]. Although this pipeline works very well for most cases, some '*difficult cases*' require additional efforts, that include new cloning strategies, such as the addition of fusion tags or even change of tags from C-terminus to N-terminus and vice versa [6]; the exchange of expression vector, suiting of the expression protocol or replacing the host cell.

In this context, several HT strategies have been developed for parallel cloning of genes of interest in different constructions, which explore different fusion proteins, different tags, and the location of the tag [6–8]. Also, protocols for increasing the yield of soluble expression of target proteins are now abundant in the literature (e.g., [3]). Finally, a myriad of commercial *E. coli* strains is available for efficient heterologous protein expression, promising high yield and enhanced solubility, especially for insoluble protein, making the choice for a particular strain a non-straightforward task.

Here we show the results for a scalable parallel screen for the expression of difficult cases using four *E. coli* strains. The assay can be readily scalable for an HT format and also include more strains. We show that the screening for expression efficiency in different strains can be a feasible and rapid way to rescue target genes that failed to be soluble expressed in the first HT approach, avoiding going back to the cloning stage of

74 the structural biology pipeline. An automated liquid handling device was central to the
75 screening, allowing rapid analysis of the expression in 96-well plates.

2. Materials and methods

All the genes tested in this work were cloned in the pET-Trx1A/LIC using the
strategy previously described by Camilo & Polikarpov [5]. Besides, for a particular target
(target **T16**) the pET-NESG vector was used. The pET-Trx1A/LIC is a ligation
independent cloning-based vector where the target gene is expressed with an N-terminal
His-tag fused to thioredoxin and a cleavage site for the Tobacco Etch Virus (TEV)
protease.

2.1. Gene Cloning

The LIC cloning protocol using pET-Trx1A/LIC vector was performed as described
previously [5]. Briefly, the plasmid was linearized by PCR using the following
oligonucleotides: Fw- 5´ TGGCGCCCTGAAAATAAAG; Rv-
5´CCGCGTCGGGTCAC and, for targets, genomic DNAs were used as templates in PCR
amplification using Phusion High-fidelity DNA Polymerase (New England Biolabs).
PCR products were individually treated for 16 h at 37 °C with 20 units of DpnI enzyme
(New England Biolabs). Following gel extraction using Wizard SV Gel and PCR
CleanUp System (Promega), 500 ng of purified and linearized vector and 200 ng of each
specific gene were treated with T4 DNA polymerase enzyme (Fermentas) in the presence
of 2.5 mM dTTP (vector) and 2.5 mM dATP (gene) for 30 min at 22 °C. Annealing of
LIC vectors and insert was performed after incubation of 3 µL of T4 Polymerase-treated
PCR fragments with 1 µL of T4 Polymerase-treated vector at 25 °C for 30 min. Then, the
mixture was used for *E. coli* DH5α competent cells transformation and positive clone
selection achieved in culture plates containing LB agar medium with 50 µg mL^{-1}
kanamycin after overnight incubation at 37 °C.

2.2. Protein Expression and Analysis

Sixteen target genes were cloned following the procedures above. The resulting vectors were used for *E. coli* expression in four DE3 strains: Lemo21, pT-GroE, Rosetta-gami 2, and Arctic Express. Transformations were performed in 24-well-LB-agar plates containing: kanamycin (50 µg mL^{-1}) and chloramphenicol (34 ug mL^{-1}) for Lemo21, pTGroE, and Rosetta-Gami 2 cells; kanamycin (50 µg mL^{-1}) and gentamycin (20 µg mL^{-1}) for Arctic Express. Colonies were inoculated in DW96 preculture plates containing 1 mL of LB medium supplied with cells specific antibiotics, sealed, and incubated overnight at 37 °C with shaking. Then, 100 µL of preculture was inoculated in 5 DW24 plates containing 4 mL of LB with specific antibiotics (plus 0.5 mM rhamnose for Lemo21 cells) and grown at 37 °C under 150 rpm agitation during 5h. Afterward, aliquots were collected and IPTG induction was performed using 0.4 mM final concentration for Lemo21 and 1 mM for the other strains. The temperature was reduced to 18 °C (pTGroE and Rosetta-Gami 2), 30 °C (Lemo21), or 11 °C (Arctic Express), and expression was conducted for 16 h at 150 rpm. In parallel, 100 µL of pre-culture was inoculated in 5 DW24 plates containing 4 mL of ZYP5052 [9] auto-induction medium, and the expression was performed as LB-conditions without IPTG. The plates were centrifuged for 10 min at 2000xg and each pellet was resuspended with 1 mL of lysis buffer (Tris–HCl 50 mM, NaCl 300 mM, lysozyme 0.25 mg mL^{-1}, 0.1 mM PMSF, at pH 8). Cells were lysed by freezing at -80 °C overnight and thawing at 17 °C for 1 h under agitation. DNAseI (Sigma) was added at 10 µg mL^{-1} final concentration following incubation at 17° C for 15 min. The lysate purification was carried out in an automated platform Freedom EVO 200 (Tecan) using Ni-Sepharose 6 Fast Flow Resin (GE) in 96-well Receiver Plates 20 µm (Macherey–Nagel). The lysate passed through the resin by vacuum filtration followed by resin washing with 1 mL of buffer A (Tris–HCl 50 mM, NaCl 300 mM, pH 8) and buffer B (Tris-HCl 50 mM, NaCl 300 mM, imidazole 20 mM, pH 8) for removal of unspecific and unbounded proteins. Elution of interest proteins was performed with 150 µL of buffer C (Tris-HCl 50 mM, NaCl 300 mM, imidazole 300 mM, pH 8) and samples were analyzed by gel electrophoresis for expression protein screening evaluation. The expression yield of the soluble proteins expressed was assessed by Coomassie-stained SDS-PAGE 15%.

137   An overall view of the entire pipeline, from cloning to the analysis of heterologous

138 expressed protein is shown in Figure 1.

Figure 1. Overview of the pipeline used in this work. Selected genes are cloned in the pET-Trx1A/LIC vector. After plasmid propagation in E.coli DH5α cells, the purified vector is used for transformation of a set of E. coli strains, the Expression Strain Bank. The cells are cultured in 96-well plates in auto-induction medium and, after cell lysis by freezing, a liquid handling robot separates the soluble fraction, incubates with the affinity resin, washes, and elutes the purified targets, which are further analyzed by SDS-PAGE 15%. Created with BioRender.com.

3. Results

Sixteen 'difficult cases' genes were identified in our typical protein expression

pipeline. In this pipeline, the genes are cloned in the pET-Trx1A/LIC vector and

expressed in *E. coli* Rosetta 2 (DE3) strain. In 13 out of the 16 cases, no soluble protein

expression could be observed, or the expression yield was very low for a structural

biology pipeline. These targets include bacterial (*Staphylococcus aureus and*

Enterococcus faecalis, Mycobacterium tuberculosis) enzymes, as well as plasmodial

enzymes (*Plasmodium falciparum* and *Plasmodium vivax*). The genes were initially

synthesized with codon optimization for *E. coli* expression. The only exception was the
E. faecalis genes, in which the genes were cloned directly from the genomic DNA. All
the constructs present a molecular weight among 30-50 kDa and some of them are under-
represented in the PDB, suggesting that they are difficult cases for protein expression in
the structural biology pipeline in general. A summary of the genes selected for the
screening of soluble expression in *E. coli* strains is shown in Table 1.

The protein purification procedures, as described in the Materials and Methods
section, are assisted by a liquid-handling device. This automation device is central to the
protocol used in this work, allowing a rapid evaluation of the amount of soluble protein
obtained in cell lysis. Also, the automated procedure is readily scalable for an increased
number of targets or the number of strains tested. The procedure followed the same HT
strategy previously established [5]. The actual bottleneck was the protein expression
since every *E. coli* strain has its typical resistance marker and its optimal temperature
for expression. In our assay, the growing temperature was set to 37 °C for all strains and
different temperatures were used in the induction stage, by allocating each strain in a
different shaker. Although this step requires increased human effort, it is easily scalable
to include more strains, in particular, if the additional strains can be used in the same
expression temperatures, or even for more genes, since they can be used in the same
plate in each shaker. After the protein expression step, the samples were collected in a
single plate for the purification procedure. The process can be further optimized for an
HT format, but it becomes clear that the current format is readily scalable for additional
genes/strains.

178
Figure 2. SDS-PAGE (15%) analysis of soluble protein recovery in the four strains used in this work. The red arrows
indicate the recovered protein with the expected molecular weight. The black arrows indicate the molecular weight of
the proteins that were not revered or only minimally recovered.

After purification, the obtained expression levels were compared by SDS-PAGE

analysis. Figure 2 shows the electrophoresis analysis obtained for the expression of the

target genes and the results are also summarized in Table 2. Targets **T1**, **T2** and **T7** are

orthologous enzymes from *E. faecalis*, *S. aureus,* and *P. falciparum*, respectively, and

all of them showed negligible soluble expression in *E. coli* Rosetta 2 strain. When tested

for the expression in Rosetta Gami 2, pT-GroE, LEMO21, and Arctic Express, we

observed soluble recovery of **T1** in Rosetta Gami 2, pT-GroE, Lemo21, and Arctic

Express. Additionally, **T2** was recovered in minimal amounts in Rosetta Gami 2 and

Arctic Express, while **T7** was recovered in higher amounts in Arctic Express and small

amounts in the other strains tested.

The targets **T3**, **T4**, **T5**, **T6,** and **T10** are five enzymes of the same pathway in *M.*

tuberculosis and *Plasmodium* species. Out of these five targets, the Arctic strain

recovered three of them in the soluble form and a fourth target (**T5**) to a lesser extent,

while pT-GroE recovered two of them with a very good expression pattern (**T3** and

T10). Lemo21 strain recovered **T5**, **T6**, and **T10**, and Rosetta Gami 2 recovered targets

T6 and **T10**. The targets **T8** and **T9** were recovered in very low yields in the pT-GroE

strain. The Arctic Express strain also recovered slow amounts of **T8**, while Rosetta Gami

2 recovered small amounts of **T9**.

The targets **T11** and **T14** are glycosyltransferases, which are targets known to be

insoluble in many cases. **T11** was also tested for its expression in two truncated forms

(**T12** and **T13**) and also in the full-length protein in the pET-NESG vector (**T16**). A

truncated form of **T14** was also tested in the pET-Trx1A/LIC vector and is listed as **T15**.

The Lemo21 strain was not able to recover any of the targets, while Rosetta Gami

recovered only **T13** to a very small extent. The pT-GroE strain recovered two of the

truncated forms for these targets (**T13** and **T15**) and, to a lesser extent, the third truncated

form (**T12**). Finally, the Arctic strain recovered the same three truncated forms with very

good contrast to the background proteins observed in the SDS-PAGE.

4. Discussion

After screening the strains available in our group, we found, in general, that 9 out of

the 16 targets, or 56% were recovered in reasonable amounts. In at least five cases, the

recovery was observed in multiple strains, while leastwise in three cases, a single strain

was found to be more efficient in recovering a target in the soluble form (**T7**, **T13**, and

T14). In the other five cases, including the two full-length glycosyltransferases, none of

the strains used here was found to recover the target in the soluble form.

To be able to test several different targets in a batch experiment, we missed some
optimization procedures that could enhance our recovering rate. For example, the
Lemo21 strain can fine-tune the expression level of toxic protein by the titratable
regulation of the T7 lysozyme (T7L) expression using L-rhamnose [10,11].

pT-GroE, a strain that co-expresses a molecular chaperone to assist the proper
protein folding, was found to result in a good recovering rate as well as a good contrast
from the target protein to the basal protein expression, making it very easy to follow the
target protein being expressed. Similar to pT-GroE, the Arctic Express strain combines
low temperature-adapted chaperones Cpn10 and Cpn60, similar to *E. coli* GroEL and
GroES, to result in chaperon assisted and low-temperature protein expression [12]. This
strategy was successful in several targets tested here and was also the strain
demonstrating a higher recovering rate.

The recovering rate in any structural biology pipeline will always be highly
dependent on the targets chosen in the project. In this sense, we have no aim to claim
that a given commercial strain is better or worse than each other. Rather than that, we
show that it might be worth having a suitable and scalable assay to screen different host
strains for the expression of soluble protein in difficult cases. We easily foresee a
scenario where proteins that were not recovered in the soluble fraction after a '*standard*'
protocol can be re-assayed for a '*salvage*' pipeline, where different strains are tested for
the expression. The experimental cost of this effort includes the transformation of the
existing vector in different cell strains and the implementation of a protocol for
expression and purification that is scalable and may be implemented in HT format.
Protein targets that cannot be recovered in this salvage pipeline are still suitable for
changing the construction or even going for a eukaryotic cell factory as a host for
heterologous expression. In these cases, the experimental costs are slightly higher and
involve going back to the construct and primer design, gene cloning, and testing for
expression again.

5. Conclusion

In conclusion, here we demonstrated that nine out of 16 'difficult targets' could be
recovered in the soluble fraction after testing a set of different *E. coli* strains. This easy-
to-implement assay is scalable for a different number of genes and/or strains. We found
that the screening may represent an affordable salvage pipeline for the recovery of target
proteins which complement a typical pipeline for protein expression and purification in
structural biology efforts.

Declaration of Competing Interest
256

257   The authors declare that they have no competing financial interests.

259   **Acknowledgments**
260

The authors thank to the members of the Molecular Biotecnology Group @
IFSC/USP and, in particular, for our technical staff: Maria Auxiliadora Santos, João
Possatto and Josimar Sartori. The authors also thank the financial support from the
Fundação de Apoio À Pesquisa do Estado de São Paulo (FAPESP) through grants,
2015/26722-8, 2017/03966-4, 2017/24901-8, 2018/21213-6, 2019/20219-3 and
2020/03983-9 and from the Conselho Nacional de Desenvolvimento Científico e
Tecnológico (CNPq) through grants 303165/2018-9, 485950/2013-8 and 476606/2010-
1. This study was also financed in part by the Coordenação de Aperfeiçoamento de
Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

References

1. BEDDELL CR, GOODFORD PJ, NORRINGTON FE, WILKINSON S,
WOOTTON R. COMPOUNDS DESIGNED TO FIT A SITE OF KNOWN
STRUCTURE IN HUMAN HAEMOGLOBIN. Br J Pharmacol. 1976;57: 201–
209. doi:10.1111/j.1476-5381.1976.tb07468.x

2. Rosano GL, Ceccarelli EA. Microbial Cell Factories Rare codon content affects
the solubility of recombinant proteins in a codon bias-adjusted Escherichia coli
strain. 2009; doi:10.1186/1475-2859-8-41

3. Kaur J, Kumar A, Kaur J. Strategies for optimization of heterologous protein
expression in E. coli: Roadblocks and reinforcements. International Journal of
Biological   Macromolecules.   Elsevier   B.V.;   2018.   pp.   803–822.
doi:10.1016/j.ijbiomac.2017.08.080

4. Sonoda MT, Godoy AS, Pellegrini VOA, Kadowaki MAS, Nascimento AS,
Polikarpov I. Structure and dynamics of Trichoderma harzianum Cel7B suggest
molecular architecture adaptations required for a wide spectrum of activities on
plant cell wall polysaccharides. Biochim Biophys Acta - Gen Subj. Elsevier;
2019;1863: 1015–1026. doi:10.1016/j.bbagen.2019.03.013

5. Camilo CM, Polikarpov I. High-throughput cloning, expression and purification
of glycoside hydrolases using Ligation-Independent Cloning (LIC). Protein Expr
Purif. Elsevier Inc.; 2014;99: 35–42. doi:10.1016/j.pep.2014.03.008

6. Eschenfeldt WH, Maltseva N, Stols L, Donnelly MI, Gu M, Nocek B, et al.
Cleavable C-terminal His-tag vectors for structure determination. J Struct Funct
Genomics. 2010;11: 31–39. doi:10.1007/s10969-010-9082-y

7. Stols L, Gu M, Dieckman L, Raffen R, Collart FR, Donnelly MI. A new vector for
high-throughput, ligation-independent cloning encoding a tobacco etch virus
protease   cleavage   site.   Protein   Expr   Purif.   2002;25:   8–15.
doi:10.1006/prep.2001.1603

8. Eschenfeldt WH, Lucy S, Millard CS, Joachimiak A, Mark ID. A family of LIC
vectors for high-throughput cloning and purification of proteins. Methods Mol
Biol. 2009;498: 105–15. doi:10.1007/978-1-59745-196-3_7

9.   Studier FW. Protein production by auto-induction in high-density shaking cultures.
2005; doi:10.1016/j.pep.2005.01.016

10.   Schlegel S, Löfblom J, Lee C, Hjelm A, Klepsch M, Strous M, et al. Optimizing
membrane protein overexpression in the Escherichia coli strain Lemo21(DE3). J
Mol Biol. Academic Press; 2012;423: 648–659. doi:10.1016/j.jmb.2012.07.019

11.   Schlegel S, Rujas E, Ytterberg AJ, Zubarev RA, Luirink J, de Gier JW. Optimizing
heterologous protein production in the periplasm of E. coli by regulating gene
expression levels. Microb Cell Fact. BioMed Central; 2013;12: 24.
doi:10.1186/1475-2859-12-24

12.   Ferrer M, Chernikova TN, Yakimov MM, Golyshin PN, Timmis KN. Chaperonins
govern growth of Escherichia coli at low temperatures. Nat Biotechnol. Springer
Science and Business Media LLC; 2003;21: 1267–1267. doi:10.1038/nbt1103-
1266b

13.   Gasteiger E, Hoogland C, Gattiker A, Duvaud S, Wilkins MR, Appel RD, et al.
Protein Identification and Analysis Tools on the ExPASy Server. The Proteomics
Protocols Handbook. Totowa, NJ: Humana Press; 2005. pp. 571–607.
doi:10.1385/1-59259-890-0:571

Table captions

Table 1. Target genes used for protein expression. The isoelectric point and molecular

mass were computed with the ProtParam server [13]. TBP = thiamine biosynthesis

pathway. PBP = pyridoxal 5'-phosphate biosynthesis pathway. GT= glycosyltransferase.

Table 2. Summary of the soluble protein recovery using different *E. coli* strains culture

medium. The expression levels, as shown in Figure 1, were quantified on a scale from

(+) to (++++). Here, (-) indicates no expression.

Figure Captions

Figure 1. Overview of the pipeline used in this work. Selected genes are cloned in the

pET-Trx1A/LIC vector. After plasmid propagation in *E.coli* DH5α cells, the purified

vector is used for transformation of a set of E. coli strains, the Expression Strain Bank.

The cells are cultured in 96-well plates in auto-induction medium and, after cell lysis by

freezing, a liquid handling robot separates the soluble fraction, incubates with the affinity

resin, washes, and elutes the purified targets, which are further analyzed by SDS-PAGE

15%. Created with BioRender.com.

Figure 2. SDS-PAGE (15%) analysis of soluble protein recovery in the four strains used

in this work. The red arrows indicate the recovered protein with the correct molecular

weight. The black arrows indicate the molecular weight of the proteins that were not

recovered or only minimally recovered.

352 **Table 1**. Target genes used for protein expression. The isoelectric point and molecular
353 mass were computed with the ProtParam server. TBP = thiamine biosynthesis pathway.
354 PBP = pyridoxal 5'-phosphate biosynthesis pathway. GT= glycosyltransferase.

Target	Pathway	Organism	N_{res}[a]	N_{Cys}[a]	pI[a]	MW[b] (kDa)
T1	TBP	*E. faecalis*	276	1	5.47	43.3
T2	TBP	*S. aureus*	276	2	5.78	44.2
T3	PBP	*M. tuberculosis*	299	1	5.24	45.4
T4	PBP	*M. tuberculosis*	199	2	5.33	35.2
T5	PBP	*P. vivax*	302	7	6.62	47.2
T6	PBP	*P. vivax*	219	7	6.54	38.5
T7	TBP	*P. falciparum*	310	16	6.18	49.0
T8	TBP	*P. falciparum*	302	11	8.88	47.9
T9	TBP	*P. falciparum*	400	13	8.89	60.7
T10	PBP	*P. falciparum*	219	9	6.43	38.6
T11	GT	*E. faecalis*	241	2	6.23	40.9
T12	GT	*E. faecalis*	210	2	6.22	37.7
T13	GT	*E. faecalis*	130	1	4.39	28.4
T14	GT	*E. faecalis*	325	0	9.08	51.2
T15	GT	*E. faecalis*	220	0	5.59	39.1
T16	GT	*E. faecalis*	241	2	6.23	26.9

355 [a] Applies to the target sequence, without considering the fusion protein sequence.

356 [b] Molecular mass computed with the fusion tag.

358 **Table 2.** Summary of the soluble protein recovery using different *E. coli* strains culture

359 medium. The expression levels, as shown in Figure 1, were quantified on a scale from

360 (+) to (++++). Here, (-) indicates no expression.

	E. coli strains			
Target	*Arctic Express*	*Rosetta Gami 2*	*pT-GroE*	*Lemo 21*
T1	++	++	++	++
T2	+	+	-	-
T3	+++	+	+++	+
T4	-	-	-	-
T5	++	+	-	++
T6	++++	++	-	++
T7	++	+	+	+
T8	+	-	+	-
T9	-	+	+	-
T10	++	++	++	++
T11	-	-	+	-
T12	++	+	+	-
T13	+++	+	++	-
T14	+	-	+	-
T15	+++	-	++	-
T16	-	-	+	-

362 * The recovery yield levels are ranked from no expression (-) to high yield (++++).

- Genes of interest for cloning;

- Cloning via LIC;

- Recombinant protein expression

- Cell lysis

- Separation of soluble and insoluble fractions

- DNA clean-up

- Transformation into DH5-Alpha Cells

- Liquid handler

- MiniPrep

Expression Strain Bank

- Selection of expression strains according to the desired objective (transformation)

Arctic Express (DE3)

Rosetta Gami 2 (DE3)

pT-GROE (DE3)

Lemo 21 (DE3)