High frequency rTMS modulates functional connectivity nodes and boundaries differently in the human brain

Vladimir Belov[1,#], Vladislav Kozyrev[1,2,3#], Aditya Singh[1], Matthew D. Sacchet[4], Roberto Goya-Maldonado[1,*]

Affiliations:

[1] Laboratory of Systems Neuroscience and Imaging in Psychiatry (SNIPLab), Department of Psychiatry and Psychotherapy, University Medical Center Göttingen (UMG), Göttingen, Germany

[2] Functional Imaging Laboratory, German Primate Center – Leibniz Institute for Primate Research, Göttingen, Germany

[3] Institute of Molecular and Clinical Ophthalmology Basel, Basel, Switzerland

[4] Center for Depression, Anxiety, and Stress Research, McLean Hospital, Harvard Medical School, Belmont, MA, USA

 # equal contribution

*Corresponding author:
PD Dr. Roberto Goya-Maldonado
Laboratory of Systems Neuroscience and Imaging in Psychiatry (SNIP-Lab)
Department of Psychiatry and Psychotherapy
University Medical Center Göttingen (UMG)
Von-Siebold Str. 5, 37075 Göttingen
e-mail: roberto.goya@med.uni-goettingen.de

Abstract words:	177	Figures:	6	Suppl Figures:	3
Text words (incl M&M):	6067	Tables:	0	Suppl Tables:	2
References:	68				

Abstract

Repetitive transcranial magnetic stimulation (rTMS) has gained considerable importance in the treatment of disorders, e.g. depression. However, it is not yet understood how rTMS alters brain's functional connectivity. Here we report the changes captured by resting state functional magnetic resonance imaging (rsfMRI) within the first hour after 10Hz rTMS in (1) nodes, where the strongest functional connectivity of regions is seen, and (2) boundaries, where functional transitions between regions occur. We use support vector machines (SVM), a widely used machine learning algorithm that has been proven to be robust and effective, for the classification and characterization of time intervals of major changes in node and boundary maps, while respecting the variability between subjects. Particularly in the posterior cingulate cortex and precuneus, our results reveal that the changes in connectivity at the boundaries are slower and more complex than in those observed in the nodes, but of similar magnitude according to accuracy confidence intervals. As the network boundaries are under studied in comparison to nodes in connectomics research, our results highlight their contribution for functional adjustments to rTMS.

Keywords: rTMS, individualized parcellation, fMRI, machine learning classification, functional nodes, functional boundaries

1. Introduction

Repetitive transcranial magnetic stimulation (rTMS) has become a popular method for the non-invasive modulation of brain function (George et al., 1999). Recent neuroimaging studies have shown that functional changes induced by rTMS in a localized cortical region lead to selective and distinct modulation of activity and functional connectivity both within and between large-scale brain networks (Cocchi et al., 2015; Eldaief et al., 2011; Sale et al., 2015; Singh et al., 2020, 2019; Valchev et al., 2015; Watanabe et al., 2014). The mechanisms by which rTMS induces network modulations are still not well understood. Mapping whole-brain effects caused by local neural perturbations, including by rTMS, is a growing field of research. Well-established methods now allow for the assessment of connectome-level functional adjustments to high frequency rTMS in both node and boundary maps in sequential time intervals (Cohen et al., 2008; Wig et al., 2014).

Functional magnetic resonance imaging (fMRI) data obtained while participants are not engaged in any specific task is called resting state fMRI (rsfMRI). RsfMRI has been instrumental for advancing our understanding of the brain's macroscopic functional network architecture (Power et al., 2011; Smith et al., 2013; Yeo et al., 2011) as well as which regions might be most functionally altered in psychiatric disorders (Goya-Maldonado et al., 2016; Wei et al., 2018). However, fMRI data typically consists of functional time-courses in thousands of voxels, which on the one hand allows for accurate inference of correlations or "functional connectivities" between regions, but on the other hand has high dimensionality. Different approaches have been proposed to reduce data dimensionality of the data and to identify the most relevant patterns of spatiotemporal organization in fMRI data. This is the case for whole-brain functional regions that will be represented in our study as nodes (Stanley et al., 2013) and boundaries (Cohen et al., 2008; Gordon et al., 2016; Tourville et al., 2019). Nodes are defined as the greatest strength of local or global connectivity, also known as concepts of modularity and integration respectively, which enabled many insights into dimensional organization of the healthy and diseased brain (van den Heuvel and Sporns, 2019). Boundaries are the counterpart of the nodes, identified where the connectivity strength is the lowest or absent, usually in the transition between neighboring functional regions. In contrast to the investigation of boundaries, the scientific community has given disproportionate attention to the nodes of functional networks. In node clustering approaches, spatiotemporal elements (i.e., voxels) may be grouped on the basis of the similarity versus

dissimilarity of their functional connectivity (Stanley et al., 2013). An example of a node

clustering approach is independent component analysis (ICA), a brain mapping method that

efficiently segregates functional components based on their corresponding spatiotemporal

distribution. ICA has been widely applied to the identification of large-scale brain networks

(Agosta et al., 2012; Peraza et al., 2014; Singh et al., 2020, 2019).

Group-based parcellation schemes use fMRI data from multiple individuals to map large-scale

functional brain networks, that is, collections of widespread regions showing functional

connectivity (Eickhoff et al., 2018; Glasser et al., 2016; Tourville et al., 2019). While group-

based parcellation captures major features of functional brain organization that are evident

across individuals, such approaches may obscure certain person-specific features of brain

organization. In contrast, subject-specific parcellation methods have been shown to

effectively map aspects of functional organization that differ for particular individuals (e.g.

Kong et al., 2019; Saxe et al., 2006). Several recent studies have demonstrated that extensive

rsfMRI data collected across multiple sessions from the same individual can be used to

delineate high-quality cortical parcellations at the individual level (Gordon et al., 2017;

Laumann et al., 2015; Tavor et al., 2016). Subject-specific parcellation may enable

increasingly precise planning and delivery of rTMS interventions.

Machine learning includes the development of algorithms that can detect spatially complex

and often subtle patterns in a highly dimensional data, and has been applied to neuroimaging.

Machine learning may thus be useful for individual-level predictions that could ultimately be

used in clinical contexts (Cao et al., 2020; Drysdale et al., 2017; Redlich et al., 2016). The

support vector machine (SVM) is a machine learning algorithm that constructs hyperplanes in

multidimensional space to optimally separate data classes (Evgeniou and Pontil, 2001). Taken

together, in the current study SVM was used to identify the strongest rTMS-related changes in

brain functional connectivity patterns in both nodes and boundary maps. In the domain of

"big" datasets other highly non-linear algorithms often outperform SVM (Liu et al., 2017).

Nevertheless, SVM often yields better classification in applications with smaller datasets, that

is, datasets in which the number of features exceeds greatly the number of training data

samples (Chaplot et al., 2006). The classification task can be simplified by using unbiased

feature selection approaches. Features selection involves the identification of the most useful

data features in the training dataset (Miao and Niu, 2016). These features are then solely used

for classification. Features selection has been shown to improve accuracy while also

increasing the interpretability of identified multivariate models.

Recent neuroimaging and predictive modeling findings suggest that a locally generated brain
stimulation-induced perturbation in neural activity is gradually integrated by selective
alterations of within- and between-network dynamics (Cocchi et al., 2015; Gollo et al., 2017).
In addition, animal studies have shown that 10 Hz rTMS creates a transient cortical functional
state that is characterized by increased excitability and increased response variability
(Kozyrev et al., 2018, 2014). The latter finding implicates a reduction of specificity
(decorrelation) of the functionally distinct regions and the widening of boundaries between
them. In the current project, we endeavored to understand 10 Hz rTMS modulation of whole-
brain connectivity patterns by employing complementary approaches including assessment of
both nodes and boundaries of functional brain networks. Our approach allowed for the data-
driven identification of the most substantial functional changes across time. Analyzed data
include timely precise rsfMRI measurements at baseline (R0) as well as following 10 Hz
rTMS in the left dorsolateral prefrontal cortex (DLPFC) in continuous time intervals
occurring 10-15 min, 27-32 min, and 45-50 min after stimulation (intervals named R1, R2,
R3). We used individual-level comparisons to create sham-corrected maps for nodes and
boundaries. This approach enhanced the sensitivity of detecting individual-level rTMS-
induced variations in functional connectivity. Given the multiple measurements of resting
state functional connectivity patterns from R0 to R3 time intervals, we were able to
sequentially map spatiotemporal changes related to stimulation. Based on the prior studies
described above, we hypothesized that 10 Hz rTMS would affect functional connectivity in
the nodes and in the boundaries of distant regions that interact with the DLPFC.

2. **Material and methods**

2.1 Participants and study design

23 participants were recruited from a university environment to participate in a double-blind,
sham-controlled, cross-over design study that investigated the neural effects of 10 Hz rTMS
using rsfMRI. Participants were screened with a self-report clinical questionnaire and the
Symptom Checklist 90 Revised (SCL-90-R) to ensure that they had no current or previous
history of neurological or psychiatric disorders. Additional exclusion criteria included
recreational drug use in the past month, current or history of substance abuse or addiction, any
contraindications to MRI or TMS (e.g., epilepsy, pregnancy, history of traumatic brain injury,

current use of anticonvulsant drugs), participation in any TMS or ECT study in the past 8

weeks, and unwillingness to consent or to be informed of incidental findings. Informed

consent was obtained from all subjects before their inclusion in the study. The study protocol

was approved by the Ethics Committee of the University Medical Center Göttingen.

For each participant, experiments were conducted over the course of 3 days with

approximately one week in between each day. On day 1, a structural T1-weighted volume and

rsfMRI were acquired for the identification of a subject-specific DLPFC site for rTMS

stimulation (**Supplementary Figure 1**). This target was then used for real and sham

stimulation protocols on day 2 and day 3. At the beginning of day 2 and day 3, an rsfMRI

scan (R0) was obtained pre-rTMS. Next the resting motor threshold (RMT) for each subject

and session was determined, which was then used to set the stimulation intensity (i.e., 110%

of the RMT). Thereafter, a 10 Hz rTMS clinical protocol of 3000 pulses was delivered over

37.5 min. This procedure was additionally controlled by sham rTMS in a double-blind

counterbalanced crossover design. RTMS was precisely delivered to each subject at the pre-

selected DLFPC target, guided by online neuronavigation. Three additional rsfMRI scans (R1,

R2, and R3) were obtained post- rTMS at 10-15 min, 27-32 min, and 45-50 min after the end

of stimulation. These acquisitions allowed for the assessment of functional connectivity

changes induced by rTMS.

2.2 Neuroimaging data acquisition and preprocessing.

Structural T1-weighted scans with 1-mm isotropic resolution and functional data were

obtained using a 32-channel head coil and 3T MR scanner (Magnetom TRIO, Siemens

Healthcare, Erlangen, Germany). For rsfMRI, 125 volumes were acquired in approximately

5.5 minutes using a gradient EPI sequence with the following parameters: TR of 2.5 s, TE of

33 ms, 60 slices with a multiband factor of 3, FOV of 210 mm $\times$ 210 mm, 2$\times$2$\times$2 mm, with

10% gap between slices and anterior to posterior phase encoding.

RsfMRI preprocessing was conducted in Data Processing Assistant for Resting-State fMRI

software (DPARSF V4.4, (Chao-Gan and Yu-Feng, 2010)). Initial steps of preprocessing

included the slice timing and head motion correction (Satterthwaite et al., 2013; Yan et al.,

2013). Afterwards corrected images were analyzed using the SPM12 gradient echo field map

unwarping tool (Togo et al., 2017). White matter, CSF, and global signal were then regressed

out to additionally reduce nuisance effects (Power et al., 2017). Corresponding T1-weighted

images were adjusted to the standard Montreal Neurological Institute (MNI) template and

then used for spatial normalization of rsfMRI with a resampled voxel size of 3x3x3 mm^3. The

preprocessed images were then spatially smoothed with a 4x4x4 mm full width at half

maximum (FWHM) Gaussian kernel. Data were then detrended and band-pass (0.01–0.1 Hz)

filtered to remove bias from low-frequency drift and high frequency noise.

2.3 Parcellation of individual-level resting state functional correlations

Individualized cortical parcellations were obtained using two methods: (1) the "Snowballing"

algorithm provided a connectivity node density map (Wig et al., 2014) and (2) Boundary

Mapping generated an average spatial boundary map (Cohen et al., 2008; Wig et al., 2014).

Both methods use whole-brain resting state functional connectivity (RSFC) to create

individualized three-dimensional node and boundary maps. Briefly, the Snowballing

algorithm uses seed-based RSFC to identify locations that are correlated (the "neighbors")

with a starting seed region of interest (ROI) location. These neighbors are then used as the

new seed regions and the procedure is repeated iteratively. Each iteration of identified

neighbors in this procedure is referred as a "zone". The neighbors correlated with a given seed

region are spatially clustered with their distinct local maxima. Averaged peak voxel locations

over multiple zones results in a node density map that represents the number of times a voxel

was identified as a node across all ROI correlation zones from starting seed locations. As in

(Wig et al., 2014), nodes were identified as the peak distribution from three zones of

Snowballing. In the original study the snowballing parcellation was performed on a cortical

surface, here we extended it to the whole brain volume space to include subcortical regions

and to match the dimensionality of our feature selection methods. We have validated 3D node

density maps by showing that inter-individual variability exceeds intra-individual variability

both in the current dataset as well as in an independent dataset (**Supplementary Figure 2**).

These node distribution maps have been shown to be invariant to the locations of starting

seeds as well as the seed sizes (Wig et al., 2014).   We used 278 initiation points that

corresponded to 278 ROIs of 5mm radius (Power et al., 2011) to obtain a global effect of the

rTMS. The correlation maps were thresholded at *r*>0.3.

In contrast to the definition of nodes, RSFC-Boundary Mapping identifies locations where the

patterns of RSFC correlations exhibit abrupt transitions, and therefore provides estimates of

putative boundaries between functional regions (Cohen et al., 2008). In contrast to the initially

proposed technique that flattens a given subject's cortex into a 2D surface (Wig et al., 2014),

we performed all computations directly on the subject's full-volume 3D Cartesian grid. This

allowed us to simultaneously overlay, for each subject, the 3D Snowballing node density
images with the RSFC-Boundary Mapping output (**Figure 1**). Nevertheless, similarly to the
original report, we also found that boundaries resulting from RSFC-Boundary Mapping
(**Figure 1, in green**) typically surrounded the nodes of high peak count identified by the
Snowballing algorithm (**Figure 1, in red**). Aside from this difference, we followed the RSFC-
Boundary Mapping method as previously described (Wig et al., 2014). To speed-up
computation time, the full-resolution preprocessed fMRI data were first resampled to a
coarser grid, resulting in 19,973 voxels (4.5x4.5x4.5 mm^3 in size). The whole-brain RSFC
map was then computed for each voxel by correlating the time series of the given voxel with
every other voxel. Then, the similarity between each voxel's temporal correlation map and
every other voxel's temporal correlation map was computed as the spatial correlation between
them. That resulted in 19,973x19,973 symmetrical spatial correlation matrices. Every row of
this matrix corresponds to a reference voxel and can be displayed in the volume of the brain
where the value in every voxel is the similarity between the temporal correlation map at that
position and the reference voxel. Every row was then mapped to the brain mask back,
representing voxel's similarity map. Next the spatial gradient was computed and then
averaged across all similarity maps. The average of those spatial gradient maps represents the
probability with which each location is identified as a point of rapid change in the RSFC maps
between two neighboring areas.

Figure 1: An example of individual-level node and boundary maps. It is noteworthy that
calculated RSFC-Snowballing nodes (red areas, thresholded r=0,3) are often surrounded by
transitional areas represented by RSFC-Boundary mapping boundaries (green areas
thresholded r=0,08), revealing the complementary nature of both parcellation methods.

2.4 Feature selection

Machine learning models can be directly applied to both node and boundary maps. However,
due to a large number of features (10^4-10^5 voxels) compared to the number of subjects in this
study (n=23), that would lead to a substantial overfitting. Subsequently, overfitted
classification models suffer from unsatisfactory interpretability and accuracy (Khosla et al.,
2019; Kohoutová et al., 2020). In order to select a smaller number of physiologically plausible
voxels located in node and boundary maps, we employed a novel data-driven feature selection
approach based on the resting state networks (RSNs). RSNs were accessed in the same group
of subjects but from an independent dataset (Day 1) from the classification dataset (Day 2 and
3). The RSNs were identified by group ICA (Khosla et al., 2019; Smith et al., 2013) using
MELODIC tool of FSL 5.0.7. We temporally concatenated the fMRI data of 23 subjects
recorded on Day 1. Based on visual inspection and the power spectrum of the MELODIC
output, we selected the nine best-fitting spatiotemporal independent components (IC). While
the node regions were assumed to lie close to maxima of the selected ICs, the boundary areas
were rather identified at the intersections of the ICs.

Figure 2: ICA-based binary masks of node (left) and boundary masks (right) from a dataset

not used in machine learning classification. ICs node regions grow in size when one increases

the threshold value. At the moment two or more ICs meet in one particular voxel, that voxel is

identified as the boundary. Changing the threshold value allows for the optimization of the

number of voxels corresponding to both masks.

As the different ICs showed different strengths (distributions of values across the brain) and

we intended to have multiple networks represented rather than a single "winner", the node

areas were defined by overlaying of all ICs. The percentage of voxels belonging to every IC

was controlled by the threshold; voxels with lower strength were discarded. Next, the

components were merged together resulting in an array of nodal voxels. The value of the node

density (Snowballing) map of each subject/time interval was then extracted for each of those

voxels. This process was designed to select the most important features that would then be

modeled using the classification algorithm. An example set of node masks for a specific

subject is presented in **Figure 2** (left; thresholded at 97.5% and 99% of discarded voxels).

In case of the boundary mask, we have developed an algorithm closely related to watershed

transform (Kolade et al., 2014). Underlying idea of watershed transform is finding an optimal

position for dams to be built, where the water coming from different basins, according to the

surface shape, would meet. In our case the surface is represented by all ICs and the basins are

the strongest voxels according to ICs. Starting from the maximum of each component (100%

threshold), and then reducing the threshold in 1% steps, the ICs increase in size. A voxel is

identified as a boundary when two or more ICs intersect each other (i.e., the same voxel is

included in multiple ICs). Single voxels, i.e. not surrounded by the voxels from the same IC,

were not treated as boundaries to avoid spurious findings. The number of included voxels

were dependent on the percentage level step that was used to descend from the maximum. An

example set of boundary masks for a specific subject is presented in the **Figure 2** (right;

threshold values of 75% and 85%). The resulting array of voxels was then used with the

gradient maps for machine learning classification.

2.4 SVM classification

The capacity of SVM to predict outcomes of the rTMS intervention was first assessed using

leave-one-group-out stratified cross-validation (CV), where the group was defined as all of

stimulation conditions (time intervals) of the same subject. Our main analysis employed SVM for multiclass classification with "one-versus-one" approach of all sham-subtracted conditions, i.e. R0 vs R1 vs R2 vs R3. SVM classification was performed using a linear kernel and the default scaling factor of 1. Sex and age were not regressed out because in every comparison both groups were represented by the same subjects, and thus automatically balanced.

The number of features (voxels) used as an input for SVM varied by changing the threshold value of the extracted ICA-based node and gradient masks. SVM assigns weight to every voxel, which can be interpreted as an importance of the voxel to separate conditions. The SVM was trained for every pair of conditions starting with a mask threshold corresponding to 10,000 voxels. By changing the threshold for both ICA-based node and gradient masks, we removed the "weakest" voxels and trained the SVM again from scratch. This procedure was repeated until the total number of voxels surviving thresholding reached zero. This recursive process allowed for the assessment of a model accuracy curve that was defined by the percentage of included voxels. To access the confidence interval of the accuracy curve, we ran SVM on 1000 bootstrap samples. The global maximum of this model mean across bootstraps accuracy curve represents the most informative set of voxels for classifying all of the conditions. In case the accuracy curve (or a certain part of it) is consistently higher than the chance level ($>25\%$), we perform pairwise classification, i.e. R0vsR1, R0vsR2, R0vsR3, R1vsR2, R1vsR3 and R2vsR3 to identify the time and the direction of the most significant changes happening after 10 Hz rTMS. The same algorithm was applied to both the node density and the gradient maps across voxel thresholds. This procedure is schematized in **Figure 3**.

CV is well known as an effective method in machine learning for testing generalizability of a trained model. Model performance was further tested using leave-two and leave-three-out stratified CV (7 fold and 11 fold CV) in which 2 or 3 subjects were withheld from training and assigned to the test sample. Our goal with this approach was to investigate the effect of the amount of data provided to a classifier, and thereby assess the impact of CV strategy. By providing more training data, as in leave-one-out CV, the model has more generalizable performance compared to leave-two-out and leave-three-out CV (Wottschel et al., 2019). The computations were completed in Python using custom-written scripts that used functions from the Nilearn v0.7.0 (https://nilearn.github.io/) and Sklearn v0.23.2 (https://scikit-learn.org/stable/) libraries.

Figure 3: Schematic diagram of the analysis pipeline. RsfMRI data from 4 sessions (R0, R1, R2, R3). Both rsfMRI from Day 2 (real rTMS) and Day 3 (sham rTMS) were used to compute RSFC-Snowballing density nodes and RSFC-Boundary Mapping gradients. Next sham stimulation maps were subtracted from the corresponding real stimulation maps. The extracted ICA-based masks derived from the Day 1 measurements were then applied to the corresponding node and gradient maps for feature selection. Finally, the remaining voxels were used for machine learning classification.

2.5 Effect of head motion on the separation of conditions

To exclude the possibility that condition-related differences are caused by head motion (Van Dijk et al., 2012), we performed pairwise classification of mean frame-to-frame head displacement across real, sham, and real-sham conditions (R0, R1, R2, R3) . The resulting machine learning performance accuracies are presented in **Supplementary Table 2**. As the performance of algorithms based on head motion are close to chance, this analysis confirmed no influence of head motion on the results of classification based on RSFC nodes and boundary maps. Additionally, we decided to censor out the high-motion frames according to the voxel size threshold, yet no session contained such frames.

3. **Results**

3.1 Analysis based on RSFC nodes density maps

We performed SVM classification on RSFC nodes density maps across all conditions and applied an iterative ICA-based feature selection step to spatially identify nodes that were strongly modulated by 10 Hz rTMS. The threshold for the ICA-based node mask, corresponding to the percentage of voxels removed, varied between 99.9% (154 voxels) to 94.5% (10,799 voxels) in 0.1% increments. The highest accuracy of 33.2±0.8% was achieved for a threshold of 99.1%, yielding 1690 voxels fed to the SVM (**Figure 4a**). Results indicated that by increasing the threshold, many informative for the classification voxels were discarded, yielding the massive accuracy drop between 99.9% and 99.7%. Lower thresholds led to a decrease in accuracy, which may be due to model overfitting (i.e., inclusion of uninformative voxels). Additionally, we compared the performance of the SVM model with a different number of CV folds. All three CV schemes showed similar accuracies across thresholds ranging from 99.9% to 98.7%. Lower thresholds for leave-three-out CV resulted in higher accuracy compared to leave-one-out and leave-two-out CV. The most informative voxels were located in the posterior cingulate cortex, angular gyrus, anterior insula, and visual cortex (**Figure 5**).

Since the accuracy was consistently higher than by chance, we aimed to discriminate the sessions in which connectivity was most strongly modulated by 10 Hz rTMS by performing pairwise classification of conditions. The highest accuracy of 74.2±1.1% was achieved for R1 vs. R2 comparison with a threshold value of 99.7%, yielding 764 voxels used in SVM-based classification (**Figure 6a**). All three CV schemes yielded similar accuracies for R1 vs R2 classification across thresholds ranging from 99.9% to 98.7% (**Supplementary Figure 3**). Majorly, the same set of voxels with both positive and negative weights was found modulated in R1 vs. R2 classification as in multiclass classification (**Figure 7**).

3.2 Analysis based on RSFC Boundary maps

For boundaries extracted from RSFC Boundary Mapping, the threshold for ICA-based boundary masks was varied from 50% (23 voxels) to 94% (10861 voxels) in 1% increments. The highest accuracy of 32.4±0.9% was achieved by the threshold of 59% corresponding to

113 voxels being fed into the SVM (**Figure 4b**). These voxels were predominantly located in

the ventral posterior cingulate cortex, precuneus, angular gyrus, and fusiform gyrus (**Figure**

5). Consistent with the analysis on Snowballing nodes described above, increasing the number

of voxels extracted from the ICA-based boundary mask caused a drop in the overall accuracy

of the SVM. All three CV routines showed a similar pattern in accuracy across the threshold

range of 50%-67%.

Similar to the analysis based on RSFC nodes density maps, we performed pairwise

classification of conditions to detect the timing of the strongest changes in boundaries. The

highest SVM accuracy of 74.5±1.1% was achieved for R1 vs. R3 (**Figure 6b**) for the

threshold value of 63% (174 voxels, **Figure 7**), Using leave-one-out CV resulted in the

performance improvement from 72.3% to 74.5% compared to leave-two-out CV for the ICA-

based boundary mask threshold of 63% (**Supplementary Figure 4**). The learning curve of R1

vs. R3 across CV approaches (**Supplementary Figure 4**) indicates the highest masking

threshold at which leave-one-out CV strategy yields the highest (or at least equal accuracy

compared to leave-two and leave-three-out CV) was 76%. This masking threshold

corresponded to 1194 voxels. For higher threshold values, leave-two-out CV exhibited

slightly higher accuracy than the other approaches.

The second highest accuracy of 68.5±1.3% was achieved by R1 vs. R2 classification (**Figure**

6b) using a threshold value of 67% that corresponds to 346 included voxels. For threshold

values from 60% to 86%, both leave-one-out and leave-two-out CV strategies were associated

with higher accuracy compared to leave-three-out CV (**Supplementary Figure 4**). The sign

of feature weights in the majority of identified voxels was in the opposite direction to the R1

vs. R3 comparison (**Figure 8**), suggesting fluctuations in the functional connectivity along

time.

Figure 4: Multiclass classification accuracy of SVM using node density map (**a**) and boundaries (**b**) across all threshold levels. The error bar indicates 95% confidence intervals over 1000 bootstraps. To test the stability of classification results, classification was also performed for leave-one-, -two- and -three-out CV routines. The highest performance for peaks (Acc =33.2%) was achieved for the threshold value of 99.1% corresponding to 1690 informative voxels. For boundaries, threshold of 59% yielded the highest performance (Acc=32.4%) resulting in 113 voxels to be strongly modulated via 10Hz rTMS.

Figure 5: Most strongly modulated voxels corresponding to the highest multiclass classification accuracy of SVM for both nodes and boundaries. The strength of modulation is color-coded by warm colormap. The numbers refer to axial plane z-coordinates in MNI space.

Figure 6: Pairwise classification accuracy of SVM using nodes (**a**) and boundaries (**b**) across all threshold levels. The highest accuracy for nodes was achieved by R1 vs. R2 comparison for the threshold value of 99.7%, corresponding to 764 voxels. In case of boundaries, R1 vs. R3 yielded the highest accuracy for the threshold value of 63% (174 voxels). The second highest accuracy was obtained by R1 vs. R2 comparison for the threshold value of 67% resulted in 346 voxels being highly modulated by 10 Hz rTMS.

Figure 7: Voxels corresponding to the highest performance of the SVM using the
Snowballing nodes for R1 vs. R2 comparison (top) and the Boundary-Mapping boundaries for
R1 vs. R3 comparison (bottom). The sign and strength of modulation is color-coded by warm
colormap (connectivity increase) and cold colormap (connectivity reduction). The numbers
refer to axial plane z-coordinates in MNI space.

Figure 8: Two highest classification accuracies based on the Boundary-Mapping boundaries
were achieved for comparisons R1 vs. R2 (top) and R1 vs. R3 (bottom) with 329 and 174
stable voxels respectively. The sign and strength of modulation is color-coded by a warm
colormap for increased connectivity and cold colormap for decreased connectivity. The
zoomed-in image shows that the majority of discriminative voxels in both comparisons were
located in the posterior cingulate cortex (PCC) and precuneus. The numbers refer to axial
plane z-coordinates in MNI space.

4. **Discussion**

There is considerable interest in both the neurosciences and clinical scenes to understand
changes in functional connectivity driven by 10 Hz rTMS. Here we have used a predictive
SVM model approach to identify locations and time intervals after stimulation that exhibit the

most substantial whole-brain functional changes in large-scale network nodes and boundaries. To overcome the issue of high-dimensional data, that is, the number of voxels in node and boundary maps is much greater than the number of subjects in the study, we have applied feature selection using multiple threshold ICA-based masks created from separate fMRI data. Confirming our hypothesis, we have identified connectivity changes related to 10 Hz rTMS in both network node and boundary maps. Of note, as indicated by cross-validated SVM, changes majorly comprise the PCC, angular gyrus, insula and fusiform gyrus. Accuracy confidence intervals in the classification of boundaries are similarly as substantial as those occurring in nodes. A complex pattern of changes was observed in boundaries alternating between decreases and increases in functional connectivity, which was particularly evident but not limited to, the PCC and precuneus (**Figure 8**). Such a pattern is reminiscent of the neuronal population dynamics in the cat visual cortex after 10 Hz rTMS (Kozyrev et al. 2014). The stimulation induced initial suppression of on-going cortical activity, followed by an increase in cortical excitability (long-term potentiation) that lasted about 2-3 hours, but was prone to slow activity fluctuations.

4.1 Individual nodes and boundaries

This work extends to the concept of modularity and integration frequently addressed in the field of connectomics (van den Heuvel and Sporns, 2019), both of which focus on the perspective of nodes in the formation and reorganization of networks in cognitive functions and in clinical disorders. Most connectomics studies to date have been focused on the strongest points of functional connectivity, "hubs", and the way that such hubs are organized to efficiently propagate information across regions (Crossley et al., 2013; Heuvel and Sporns, 2013). Transition gradients have recently received more attention in the literature (Hirose et al., 2012; Huntenburg et al., 2018; Jbabdi et al., 2013). To the best of our knowledge, the current study is the first to show which changes occur in functional connectivity boundaries and nodes after 10 Hz rTMS. Indeed, based on their accuracy and confidence intervals, we see that rTMS effects related to boundaries are similarly as substantial as those seen in nodes. The exact function of boundaries is not yet well understood, which might have discouraged its systematic evaluation. It may be speculated that boundaries may act to segregate information within functional regions, or that they may support network stability (Radicchi and Arenas, 2013). Another possible role of boundaries may be supporting functional adaptation of a given region during plastic changes in the mature primate brain. Evidence for this has been shown in primary motor and sensory regions (for review see Florence et al., 1997). In support

of this theory are several studies that have shown that rTMS-induced changes in neuronal

inhibition can prime cortical networks for the expression of subsequent experience-dependent

plasticity (Holtmaat and Svoboda, 2009; Kozyrev et al., 2018; Lenz et al., 2016). The high

frequency rTMS potentially creates a cortical state with enhanced plasticity, opening a time

window for targeted re-learning of connectivity patterns (Kozyrev et al. 2018).

Causal effects of 1 and 5 Hz rTMS on global functional connectivity have been explored

recently using fMRI (Castrillon et al., 2020; Davis et al., 2017). These studies have found that

distant effects, that is, effects relatively far from DLPFC, are determined by connectivity

profiles of the stimulation target with macroscopic networks. Excitatory 10 Hz rTMS in the

left DLPFC, as we used in the current study, resulted in multivariate patterns of increases and

decreases in functional connectivity. These fluctuations occurred primarily in the PCC,

angular gyrus, fusiform gyrus, and insula regions. Each of these regions have been previously

shown to be functionally related to the DLPFC (Bokde et al., 2006; Chang and Glover, 2009;

Jiang et al., 2015; Taren et al., 2017), reinforcing the notion of distant effects of this

stimulation protocol. In the current study, the SVM most substantially identified functional

connectivity changes in nodes that occur about 30 min after stimulation. In contrast, effects

related to boundaries were temporally extended up to 45 min. Previous studies that

investigated the effects of 10 Hz rTMS with rsfMRI constrained their analyses to particular

seeds or networks of interest. Another important advancement provided by the current study

is an assessment that started from a global evaluation of 10 Hz rTMS effects, considering

maps of nodes and boundaries at the subject level at different intervals. Cytoarchitectural

divisions of the PCC into dorsal and ventral parts have been previously shown to exhibit

distinct functional connectivity patterns (Vogt et al., 2006). According to findings from a

recent study (Salehi et al., 2020), both PCC and precuneus contain predominantly transient

nodes - an entropy class that the authors used to classify changes in network assignments

across subjects and brain states. Our findings are consistent with this notion, highlighting that

further functional heterogeneity is largely elicited in these regions, particularly in boundaries,

after 10 Hz rTMS.

The importance of individualized characterization of functional brain networks has been

highlighted in the literature (Gordon et al., 2017; Laumann et al., 2015; Tavor et al., 2016).

These developments may inform future clinical applications based in "precision" or "systems"

medicine. Toward this goal, methods applied in the current study might have been

advantageous, considering that individual subject analysis of node and boundary maps closely

correspond to the original study (Wig et al., 2014). In our study though functional boundaries
were not only extracted for the cortical surface. To match the resolution and dimensionality of
the node maps, we calculated boundaries on the whole brain. Furthermore, our study is unique
in that we used pairwise comparisons of each individualized map type, that is, nodes and
boundaries, in the context of a double-blind design controlling for placebo effects. This may
have allowed us to identify the contribution of boundaries to the functional changes caused by
10 Hz rTMS. This individualized approach was followed by SVM, which might have
contributed to the identification of the most important effects related to the time intervals and
areas.

4.2 SVM classification and feature selection

SVM is a machine learning approach with clear advantages over univariate models. With that
said, important preconditions had to be fulfilled to avoid potential biases. One of the main
concerns when we applied SVM was to prevent overfitting. The full-resolution rsfMRI was
first resampled to 19,973 voxels to reduce the computation time of the spatial correlation
matrix – the most computationally demanding step in the algorithm. Splitting the rsfMRI
signal into two complementary maps also had the practical benefit of enabling independent
assessment using SVM. This also reduced the number of input variables to the model, and
thus further prevented overfitting. Secondly, the large number of voxels in both individualized
node and boundary maps required additional feature selection. While there are several
masking algorithms that can locate functional nodes of the brain, to our knowledge, there is
no complementary boundary parcellation method associated with any of them. For this
reason, we have used group ICs to create both nodal and boundary masks.

Voxels included to the masks are controlled by the threshold. While these masks were built in
a complementary manner, these masks included common voxels in several regions, including
ventral PCC, the boundary area between angular gyrus and fusiform gyrus, and the boundary
area between fusiform gyrus and associative visual cortex. These regions are consistent with
findings from a recent study that reported on low stability of connectivity regions (Salehi et
al., 2020). Some of these regions have been associated with myelin content, particularly the
ones that myelinate latest during development (Becker, 1953; Glasser and Essen, 2011). As
some of the implicated regions overlap with those with late myelination, this suggests a role
for flexible connectivity in learning (Davis et al., 2017).

Our approach is closely related to the recursive feature elimination technique that has been
previously applied in neuroimaging (Lu et al., 2016; Wottschel et al., 2019). In our method,
voxels/features are not removed based on their rank according to SVM, but rather based on
the information obtained from the independent dataset (not included in SVM modeling).
Therefore, our approach is less prone to overfitting and inflating performance accuracy. On
the other hand, a drawback of our approach is that some fraction of the voxels not included
might be potentially informative for classification.

Considering our small sample size, the performance of the classifier was low – close to
chance - when all the features were used. Since in the multiclass classification SVM has
shown the accuracy consistently higher than by chance, we performed post-hoc pairwise
classification of conditions to pinpoint the timing of the most significant changes. For most of
the pairwise comparisons, there was a substantial drop in the accuracy, when more than 1000
voxels were used in the model. This observation may be explained by the presence of too
many insignificant voxels and a small sample size. After the optimal number of voxels was
reached from unbiased feature selection, further removal of voxels resulted in a drop of the
accuracy. This is consistent with previous studies that have used similar machine learning
approaches (Wottschel et al., 2019, 2015). Moreover, we investigated estimated accuracy
curves for different CV methods as the number of CV folds may influence the accuracy of
machine learning. That is, threshold values that yielded higher accuracy for leave-three-out, in
contrast to leave-one-out CV, are more likely to have inflated accuracy as the training set is
smaller. In general, the trained model should underperform or at least yield similar accuracy
when trained with less data, otherwise the performance of the model is ambiguous. In this is
also seen in the lower accuracy of the leave-three-out in contrast to the leave-one-out CV in
our data.

4.4 Limitations

Our work has important limitations that should be considered. As mentioned above, with
feature selection not all information derived from both node and boundary maps was
considered for classification. Therefore, we cannot exclude that additional areas might have
been modulated by 10 Hz rTMS and, due to the feature selection procedure, are not identified
in our results. While it is noteworthy that systematic evaluation of thresholds is a common
approach in machine learning, we acknowledge that nodes and boundary masks behave
differently when changing threshold values. This may have contributed to the exclusion of

regions as part of boundaries at a faster rate than nodes. One possible explanation for this is
non-uniform distance between networks which may cause proximal areas in boundaries to
interact/overlap at lower thresholds compared to more distant regions in nodes. In addition, a
larger sample size may have enabled further methodological improvements such as: (1)
increasing the resolution in the masking procedure; (2) consideration of more voxels from the
masks; (3) application of non-linear predictive algorithms; (4) better fine-tuning of classifier
hyperparameters; and (5) an independent validation dataset to strengthen the reliability of our
results.

5. Conclusion

Our findings provide evidence that SVM classifiers using ICA-based feature selection can
identify different spatial definition, direction, and timing in the pattern of fMRI-based brain
connectivity changes in functional nodes and boundaries derived from 10 Hz rTMS applied to
the left DLPFC. Identified nodes and boundaries are located predominantly in the ventral
PCC, precuneus, insula, and fusiform gyrus and appear approximately 30 minutes after the
stimulation is performed. By dividing the signal into two complementary parts (nodes and
boundaries), we highlight the contribution of boundaries to modulatory effects of high
frequency rTMS. These findings provide new insights into the currently unknown role of
boundaries in network organization, motivating future, related investigations for the
advancement of connectomics.

Declaration of Competing Interest

603 The authors declare no conflicts of interest.

Acknowledgements

Funding: This work was supported by the German Federal Ministry of Education and
Research (Bundesministerium fuer Bildung und Forschung, BMBF: 01 ZX 1507, ''PreNeSt -
e:Med''). Author contributions: Vladimir Belov1,#, Vladislav Kozyrev1,2,3#, Aditya Singh1,
Matthew D. Sacchet4, Roberto Goya-Maldonado1,* Conceptualization: RGM; Methodology:
VB, VK; Data curation: AS; Investigation: RGM; Supervision: RGM; Writing (original

draft): RGM, VB, VK; Writing (review and editing): RGM, MDS. Data and materials

availability: Raw data analysed during the current study are not publicly available due to

absence of written consent from the participants of the study. Scripts and codes used in the

analyses needed to evaluate the conclusions of the study will be deposited in a public

database.

References

Agosta, F., Pievani, M., Geroldi, C., Copetti, M., Frisoni, G.B., Filippi, M., 2012. Resting state fMRI in
Alzheimer's disease: beyond the default mode network. Neurobiol. Aging 33, 1564–1578.
https://doi.org/10.1016/j.neurobiolaging.2011.06.007
Becker, R.F., 1953. Essay on the cerebral cortex. By Gerhardt von Bonin. Charles C Thomas,
Springfield, Ill. 1950. 150 pp. Am. J. Phys. Anthropol. 11, 441–444.
https://doi.org/10.1002/ajpa.1330110317
Bokde, A.L.W., Lopez-Bayo, P., Meindl, T., Pechler, S., Born, C., Faltraco, F., Teipel, S.J., Möller, H.-J.,
Hampel, H., 2006. Functional connectivity of the fusiform gyrus during a face-matching task
in subjects with mild cognitive impairment. Brain 129, 1113–1124.
https://doi.org/10.1093/brain/awl051
Cao, B., Cho, R.Y., Chen, D., Xiu, M., Wang, L., Soares, J.C., Zhang, X.Y., 2020. Treatment response
prediction and individualized identification of first-episode drug-naïve schizophrenia using
brain functional connectivity. Mol. Psychiatry 25, 906–913. https://doi.org/10.1038/s41380-
018-0106-5
Castrillon, G., Sollmann, N., Kurcyus (Bieńkowska), K., Razi, A., Krieg, S., Riedl, V., 2020. The
physiological effects of noninvasive brain stimulation fundamentally differ across the human
cortex. Sci. Adv. 6, eaay2739. https://doi.org/10.1126/sciadv.aay2739
Chang, C., Glover, G.H., 2009. Effects of model-based physiological noise correction on default mode
network anti-correlations and correlations. NeuroImage 47, 1448–1459.
https://doi.org/10.1016/j.neuroimage.2009.05.012
Chao-Gan, Y., Yu-Feng, Z., 2010. DPARSF: A MATLAB Toolbox for "Pipeline" Data Analysis of Resting-
State fMRI. Front. Syst. Neurosci. 4. https://doi.org/10.3389/fnsys.2010.00013
Chaplot, S., Patnaik, L.M., Jagannathan, N.R., 2006. Classification of magnetic resonance brain images
using wavelets as input to support vector machine and neural network. Biomed. Signal
Process. Control 1, 86–92. https://doi.org/10.1016/j.bspc.2006.05.002
Cocchi, L., Sale, M.V., Lord, A., Zalesky, A., Breakspear, M., Mattingley, J.B., 2015. Dissociable effects
of local inhibitory and excitatory theta-burst stimulation on large-scale brain dynamics. J.
Neurophysiol. 113, 3375–3385. https://doi.org/10.1152/jn.00850.2014
Cohen, A.L., Fair, D.A., Dosenbach, N.U.F., Miezin, F.M., Dierker, D., Van Essen, D.C., Schlaggar, B.L.,
Petersen, S.E., 2008. Defining functional areas in individual human brains using resting
functional connectivity MRI. NeuroImage 41, 45–57.
https://doi.org/10.1016/j.neuroimage.2008.01.066
Crossley, N.A., Mechelli, A., Vértes, P.E., Winton-Brown, T.T., Patel, A.X., Ginestet, C.E., McGuire, P.,
Bullmore, E.T., 2013. Cognitive relevance of the community structure of the human brain
functional coactivation network. Proc. Natl. Acad. Sci. U. S. A. 110, 11583–11588.
https://doi.org/10.1073/pnas.1220826110
Davis, S.W., Luber, B., Murphy, D.L.K., Lisanby, S.H., Cabeza, R., 2017. Frequency-specific
neuromodulation of local and distant connectivity in aging and episodic memory function.
Hum. Brain Mapp. 38, 5987–6004. https://doi.org/10.1002/hbm.23803
Drysdale, A.T., Grosenick, L., Downar, J., Dunlop, K., Mansouri, F., Meng, Y., Fetcho, R.N., Zebley, B.,
Oathes, D.J., Etkin, A., Schatzberg, A.F., Sudheimer, K., Keller, J., Mayberg, H.S., Gunning,

F.M., Alexopoulos, G.S., Fox, M.D., Pascual-Leone, A., Voss, H.U., Casey, B.J., Dubin, M.J., Liston, C., 2017. Resting-state connectivity biomarkers define neurophysiological subtypes of depression. Nat. Med. 23, 28–38. https://doi.org/10.1038/nm.4246

Eickhoff, S.B., Yeo, B.T.T., Genon, S., 2018. Imaging-based parcellations of the human brain. Nat. Rev. Neurosci. 19, 672–686. https://doi.org/10.1038/s41583-018-0071-7

Eldaief, M.C., Halko, M.A., Buckner, R.L., Pascual-Leone, A., 2011. Transcranial magnetic stimulation modulates the brain's intrinsic activity in a frequency-dependent manner. Proc. Natl. Acad. Sci. U. S. A. 108, 21229–21234. https://doi.org/10.1073/pnas.1113103109

Evgeniou, T., Pontil, M., 2001. Support Vector Machines: Theory and Applications. pp. 249–257. https://doi.org/10.1007/3-540-44673-7_12

Florence, S.L., Jain, N., Kaas, J.H., 1997. Plasticity of Somatosensory Cortex in Primates. Semin. Neurosci. 9, 3–12. https://doi.org/10.1006/smns.1997.0101

George, M.S., Stallings, L.E., Speer, A.M., Nahas, Z., Spicer, K.M., Vincent, D.J., Bohning, D.E., Cheng, K.T., Molloy, M., Teneback, C.C., Risch, S.C., 1999. Prefrontal repetitive transcranial magnetic stimulation (rTMS) changes relative perfusion locally and remotely. Hum. Psychopharmacol. Clin. Exp. 14, 161–170. https://doi.org/10.1002/(SICI)1099-1077(199904)14:3<161::AID-HUP73>3.0.CO;2-2

Glasser, M.F., Coalson, T.S., Robinson, E.C., Hacker, C.D., Harwell, J., Yacoub, E., Ugurbil, K., Andersson, J., Beckmann, C.F., Jenkinson, M., Smith, S.M., Van Essen, D.C., 2016. A multi-modal parcellation of human cerebral cortex. Nature 536, 171–178. https://doi.org/10.1038/nature18933

Glasser, M.F., Essen, D.C.V., 2011. Mapping Human Cortical Areas In Vivo Based on Myelin Content as Revealed by T1- and T2-Weighted MRI. J. Neurosci. 31, 11597–11616. https://doi.org/10.1523/JNEUROSCI.2180-11.2011

Gollo, L.L., Roberts, J.A., Cocchi, L., 2017. Mapping how local perturbations influence systems-level brain dynamics. NeuroImage 160, 97–112. https://doi.org/10.1016/j.neuroimage.2017.01.057

Gordon, E.M., Laumann, T.O., Adeyemo, B., Huckins, J.F., Kelley, W.M., Petersen, S.E., 2016. Generation and Evaluation of a Cortical Area Parcellation from Resting-State Correlations. Cereb. Cortex N. Y. N 1991 26, 288–303. https://doi.org/10.1093/cercor/bhu239

Gordon, E.M., Laumann, T.O., Gilmore, A.W., Newbold, D.J., Greene, D.J., Berg, J.J., Ortega, M., Hoyt-Drazen, C., Gratton, C., Sun, H., Hampton, J.M., Coalson, R.S., Nguyen, A.L., McDermott, K.B., Shimony, J.S., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E., Nelson, S.M., Dosenbach, N.U.F., 2017. Precision Functional Mapping of Individual Human Brains. Neuron 95, 791-807.e7. https://doi.org/10.1016/j.neuron.2017.07.011

Goya-Maldonado, R., Brodmann, K., Keil, M., Trost, S., Dechent, P., Gruber, O., 2016. Differentiating unipolar and bipolar depression by alterations in large-scale brain networks. Hum. Brain Mapp. 37, 808–818. https://doi.org/10.1002/hbm.23070

Heuvel, M.P. van den, Sporns, O., 2013. Network hubs in the human brain. Trends Cogn. Sci. 17, 683–696. https://doi.org/10.1016/j.tics.2013.09.012

Hirose, S., Watanabe, T., Jimura, K., Katsura, M., Kunimatsu, A., Abe, O., Ohtomo, K., Miyashita, Y., Konishi, S., 2012. Local Signal Time-Series during Rest Used for Areal Boundary Mapping in Individual Human Brains. PLOS ONE 7, e36496. https://doi.org/10.1371/journal.pone.0036496

Holtmaat, A., Svoboda, K., 2009. Experience-dependent structural synaptic plasticity in the mammalian brain. Nat. Rev. Neurosci. 10, 647–658. https://doi.org/10.1038/nrn2699

Huntenburg, J.M., Bazin, P.-L., Margulies, D.S., 2018. Large-Scale Gradients in Human Cortical Organization. Trends Cogn. Sci. 22, 21–31. https://doi.org/10.1016/j.tics.2017.11.002

Jbabdi, S., Sotiropoulos, S.N., Behrens, T.E., 2013. The topographic connectome. Curr. Opin. Neurobiol., Macrocircuits 23, 207–215. https://doi.org/10.1016/j.conb.2012.12.004

Jiang, J., Beck, J., Heller, K., Egner, T., 2015. An insula-frontostriatal network mediates flexible
cognitive control by adaptively predicting changing control demands. Nat. Commun. 6, 8165.
https://doi.org/10.1038/ncomms9165
Khosla, M., Jamison, K., Ngo, G.H., Kuceyeski, A., Sabuncu, M.R., 2019. Machine learning in resting-
state fMRI analysis. Magn. Reson. Imaging 64, 101–121.
https://doi.org/10.1016/j.mri.2019.05.031
Kohoutová, L., Heo, J., Cha, S., Lee, S., Moon, T., Wager, T.D., Woo, C.-W., 2020. Toward a unified
framework for interpreting machine-learning models in neuroimaging. Nat. Protoc. 15, 1399–
1435. https://doi.org/10.1038/s41596-019-0289-5
Kolade, O., Olayinka, A.A., Ovie, U., 2014. Fingerprint Database Optimization Using Watershed
Transformation Algorithm. Open J. Optim. 3, 59–67.
https://doi.org/10.4236/ojop.2014.34006
Kong, R., Li, J., Orban, C., Sabuncu, M.R., Liu, H., Schaefer, A., Sun, N., Zuo, X.-N., Holmes, A.J.,
Eickhoff, S.B., Yeo, B.T.T., 2019. Spatial Topography of Individual-Specific Cortical Networks
Predicts Human Cognition, Personality, and Emotion. Cereb. Cortex N. Y. N 1991 29, 2533–
2551. https://doi.org/10.1093/cercor/bhy123
Kozyrev, V., Eysel, U.T., Jancke, D., 2014. Voltage-sensitive dye imaging of transcranial magnetic
stimulation-induced intracortical dynamics. Proc. Natl. Acad. Sci. U. S. A. 111, 13553–13558.
https://doi.org/10.1073/pnas.1405508111
Kozyrev, V., Staadt, R., Eysel, U.T., Jancke, D., 2018. TMS-induced neuronal plasticity enables targeted
remodeling of visual cortical maps. Proc. Natl. Acad. Sci. U. S. A. 115, 6476–6481.
https://doi.org/10.1073/pnas.1802798115
Laumann, T.O., Gordon, E.M., Adeyemo, B., Snyder, A.Z., Joo, S.J., Chen, M.-Y., Gilmore, A.W.,
McDermott, K.B., Nelson, S.M., Dosenbach, N.U.F., Schlaggar, B.L., Mumford, J.A., Poldrack,
R.A., Petersen, S.E., 2015. Functional System and Areal Organization of a Highly Sampled
Individual Human Brain. Neuron 87, 657–670. https://doi.org/10.1016/j.neuron.2015.06.037
Lenz, M., Galanis, C., Müller-Dahlhaus, F., Opitz, A., Wierenga, C.J., Szabó, G., Ziemann, U., Deller, T.,
Funke, K., Vlachos, A., 2016. Repetitive magnetic stimulation induces plasticity of inhibitory
synapses. Nat. Commun. 7, 10020. https://doi.org/10.1038/ncomms10020
Liu, P., Choo, K.-K.R., Wang, L., Huang, F., 2017. SVM or deep learning? A comparative study on
remote sensing image classification. Soft Comput. 21. https://doi.org/10.1007/s00500-016-
2247-2
Lu, X., Yang, Y., Wu, F., Gao, M., Xu, Y., Zhang, Y., Yao, Y., Du, X., Li, C., Wu, L., Zhong, X., Zhou, Y., Fan,
N., Zheng, Y., Xiong, D., Peng, H., Escudero, J., Huang, B., Li, X., Ning, Y., Wu, K., 2016.
Discriminative analysis of schizophrenia using support vector machine and recursive feature
elimination on structural MRI images. Medicine (Baltimore) 95, e3973.
https://doi.org/10.1097/MD.0000000000003973
Miao, J., Niu, L., 2016. A Survey on Feature Selection. Procedia Comput. Sci. 91, 919–926.
https://doi.org/10.1016/j.procs.2016.07.111
Peraza, L.R., Kaiser, M., Firbank, M., Graziadio, S., Bonanni, L., Onofrj, M., Colloby, S.J., Blamire, A.,
O'Brien, J., Taylor, J.-P., 2014. fMRI resting state networks and their association with
cognitive fluctuations in dementia with Lewy bodies. NeuroImage Clin. 4, 558–565.
https://doi.org/10.1016/j.nicl.2014.03.013
Power, J.D., Cohen, A.L., Nelson, S.M., Wig, G.S., Barnes, K.A., Church, J.A., Vogel, A.C., Laumann,
T.O., Miezin, F.M., Schlaggar, B.L., Petersen, S.E., 2011. Functional network organization of
the human brain. Neuron 72, 665–678. https://doi.org/10.1016/j.neuron.2011.09.006
Power, J.D., Plitt, M., Laumann, T.O., Martin, A., 2017. Sources and implications of whole-brain fMRI
signals in humans. NeuroImage 146, 609–625.
https://doi.org/10.1016/j.neuroimage.2016.09.038
Radicchi, F., Arenas, A., 2013. Abrupt transition in the structural formation of interconnected
networks. Nat. Phys. 9, 717–720. https://doi.org/10.1038/nphys2761

Redlich, R., Opel, N., Grotegerd, D., Dohm, K., Zaremba, D., Bürger, C., Münker, S., Mühlmann, L.,
Wahl, P., Heindel, W., Arolt, V., Alferink, J., Zwanzger, P., Zavorotnyy, M., Kugel, H.,
Dannlowski, U., 2016. Prediction of Individual Response to Electroconvulsive Therapy via
Machine Learning on Structural Magnetic Resonance Imaging Data. JAMA Psychiatry 73, 557–
564. https://doi.org/10.1001/jamapsychiatry.2016.0316

Sale, M.V., Mattingley, J.B., Zalesky, A., Cocchi, L., 2015. Imaging human brain networks to improve
the clinical efficacy of non-invasive brain stimulation. Neurosci. Biobehav. Rev. 57, 187–198.
https://doi.org/10.1016/j.neubiorev.2015.09.010

Salehi, M., Karbasi, A., Barron, D.S., Scheinost, D., Constable, R.T., 2020. Individualized functional
networks reconfigure with cognitive state. NeuroImage 206, 116233.
https://doi.org/10.1016/j.neuroimage.2019.116233

Satterthwaite, T.D., Elliott, M.A., Gerraty, R.T., Ruparel, K., Loughead, J., Calkins, M.E., Eickhoff, S.B.,
Hakonarson, H., Gur, R.C., Gur, R.E., Wolf, D.H., 2013. An improved framework for confound
regression and filtering for control of motion artifact in the preprocessing of resting-state
functional connectivity data. NeuroImage 64, 240–256.
https://doi.org/10.1016/j.neuroimage.2012.08.052

Saxe, R., Moran, J.M., Scholz, J., Gabrieli, J., 2006. Overlapping and non-overlapping brain regions for
theory of mind and self reflection in individual subjects. Soc. Cogn. Affect. Neurosci. 1, 229–
234. https://doi.org/10.1093/scan/nsl034

Singh, A., Erwin-Grabner, T., Sutcliffe, G., Antal, A., Paulus, W., Goya-Maldonado, R., 2019.
Personalized repetitive transcranial magnetic stimulation temporarily alters default mode
network in healthy subjects. Sci. Rep. 9, 5631. https://doi.org/10.1038/s41598-019-42067-3

Singh, A., Erwin-Grabner, T., Sutcliffe, G., Paulus, W., Dechent, P., Antal, A., Goya-Maldonado, R.,
2020. Default mode network alterations after intermittent theta burst stimulation in healthy
subjects. Transl. Psychiatry 10, 75. https://doi.org/10.1038/s41398-020-0754-5

Smith, S.M., Vidaurre, D., Beckmann, C.F., Glasser, M.F., Jenkinson, M., Miller, K.L., Nichols, T.E.,
Robinson, E.C., Salimi-Khorshidi, G., Woolrich, M.W., Barch, D.M., Uğurbil, K., Van Essen,
D.C., 2013. Functional connectomics from resting-state fMRI. Trends Cogn. Sci. 17, 666–682.
https://doi.org/10.1016/j.tics.2013.09.016

Stanley, M.L., Moussa, M.N., Paolini, B., Lyday, R.G., Burdette, J.H., Laurienti, P.J., 2013. Defining
nodes in complex brain networks. Front. Comput. Neurosci. 7.
https://doi.org/10.3389/fncom.2013.00169

Taren, A.A., Gianaros, P.J., Greco, C.M., Lindsay, E.K., Fairgrieve, A., Brown, K.W., Rosen, R.K., Ferris,
J.L., Julson, E., Marsland, A.L., Creswell, J.D., 2017. Mindfulness Meditation Training and
Executive Control Network Resting State Functional Connectivity: A Randomized Controlled
Trial. Psychosom. Med. 79, 674–683. https://doi.org/10.1097/PSY.0000000000000466

Tavor, I., Parker Jones, O., Mars, R.B., Smith, S.M., Behrens, T.E., Jbabdi, S., 2016. Task-free MRI
predicts individual differences in brain activity during task performance. Science 352, 216–
220. https://doi.org/10.1126/science.aad8127

Togo, H., Rokicki, J., Yoshinaga, K., Hisatsune, T., Matsuda, H., Haga, N., Hanakawa, T., 2017. Effects
of Field-Map Distortion Correction on Resting State Functional Connectivity MRI. Front.
Neurosci. 11. https://doi.org/10.3389/fnins.2017.00656

Tourville, J., Nieto-Castañón, A., Heyne, M., Guenther, F., 2019. Functional Parcellation of the Speech
Production Cortex. J. Speech Lang. Hear. Res. JSLHR. https://doi.org/10.1044/2019_JSLHR-S-
CSMC7-18-0442

Valchev, N., Ćurčić-Blake, B., Renken, R.J., Avenanti, A., Keysers, C., Gazzola, V., Maurits, N.M., 2015.
cTBS delivered to the left somatosensory cortex changes its functional connectivity during
rest. NeuroImage 114, 386–397. https://doi.org/10.1016/j.neuroimage.2015.04.017

van den Heuvel, M.P., Sporns, O., 2019. A cross-disorder connectome landscape of brain
dysconnectivity. Nat. Rev. Neurosci. 20, 435–446. https://doi.org/10.1038/s41583-019-0177-
6

Van Dijk, K.R.A., Sabuncu, M.R., Buckner, R.L., 2012. The influence of head motion on intrinsic
functional connectivity MRI. NeuroImage 59, 431–438.
https://doi.org/10.1016/j.neuroimage.2011.07.044
Vogt, B.A., Vogt, L., Laureys, S., 2006. Cytology and functionally correlated circuits of human
posterior cingulate areas. NeuroImage 29, 452–466.
https://doi.org/10.1016/j.neuroimage.2005.07.048
Watanabe, T., Hanajima, R., Shirota, Y., Ohminami, S., Tsutsumi, R., Terao, Y., Ugawa, Y., Hirose, S.,
Miyashita, Y., Konishi, S., Kunimatsu, A., Ohtomo, K., 2014. Bidirectional effects on
interhemispheric resting-state functional connectivity induced by excitatory and inhibitory
repetitive transcranial magnetic stimulation. Hum. Brain Mapp. 35, 1896–1905.
https://doi.org/10.1002/hbm.22300
Wei, Y., Chang, M., Womer, F.Y., Zhou, Q., Yin, Z., Wei, S., Zhou, Y., Jiang, X., Yao, X., Duan, J., Xu, K.,
Zuo, X.-N., Tang, Y., Wang, F., 2018. Local functional connectivity alterations in schizophrenia,
bipolar disorder, and major depressive disorder. J. Affect. Disord. 236, 266–273.
https://doi.org/10.1016/j.jad.2018.04.069
Wig, G.S., Laumann, T.O., Cohen, A.L., Power, J.D., Nelson, S.M., Glasser, M.F., Miezin, F.M., Snyder,
A.Z., Schlaggar, B.L., Petersen, S.E., 2014. Parcellating an individual subject's cortical and
subcortical brain structures using snowball sampling of resting-state correlations. Cereb.
Cortex N. Y. N 1991 24, 2036–2054. https://doi.org/10.1093/cercor/bht056
Wottschel, V., Alexander, D.C., Kwok, P.P., Chard, D.T., Stromillo, M.L., De Stefano, N., Thompson,
A.J., Miller, D.H., Ciccarelli, O., 2015. Predicting outcome in clinically isolated syndrome using
machine learning. NeuroImage Clin. 7, 281–287. https://doi.org/10.1016/j.nicl.2014.11.021
Wottschel, V., Chard, D.T., Enzinger, C., Filippi, M., Frederiksen, J.L., Gasperini, C., Giorgio, A., Rocca,
M.A., Rovira, A., De Stefano, N., Tintoré, M., Alexander, D.C., Barkhof, F., Ciccarelli, O.,
MAGNIMS study group and the EuroPOND consortium, 2019. SVM recursive feature
elimination analyses of structural brain MRI predicts near-term relapses in patients with
clinically isolated syndromes suggestive of multiple sclerosis. NeuroImage Clin. 24, 102011.
https://doi.org/10.1016/j.nicl.2019.102011
Yan, C.-G., Cheung, B., Kelly, C., Colcombe, S., Craddock, R.C., Di Martino, A., Li, Q., Zuo, X.-N.,
Castellanos, F.X., Milham, M.P., 2013. A Comprehensive Assessment of Regional Variation in
the Impact of Head Micromovements on Functional Connectomics. NeuroImage 76, 183–
201. https://doi.org/10.1016/j.neuroimage.2013.03.004
Yeo, B.T.T., Krienen, F.M., Sepulcre, J., Sabuncu, M.R., Lashkari, D., Hollinshead, M., Roffman, J.L.,
Smoller, J.W., Zöllei, L., Polimeni, J.R., Fischl, B., Liu, H., Buckner, R.L., 2011. The organization
of the human cerebral cortex estimated by intrinsic functional connectivity. J. Neurophysiol.
106, 1125–1165. https://doi.org/10.1152/jn.00338.2011

Supplementary Material

Supplementary Figure 1: Study design – During the first day we acquired T1 and rsfMRI images that were used for personalized target selection. The found target was then located in T1 image for stimulation on second and the third day via online neuronavigation. Subjects were assigned to an arm of the study receiving both real and sham in a counterbalanced crossover manner. At the beginning of the sessions on day 2 and 3, we obtained a baseline rsfMRI scan (R0). After, the 10 Hz rTMS was delivered at the selected target. Three following scans (R1, R2 R3) were obtained at 10 minutes, 27 minutes and 45 minutes after

the stimulation.

Supplementary Figure 2: Spatial correlation within and between node density maps in
healthy control subjects. The two subsets of baseline rsfMRI for healthy controls are here
referred as "10 Hz" with 23 subjects and "iTBS" with 26 subjects (due to groups assigned in
the PreNeSt-1 cohorts), in blue and orange, respectively. The combined datasets is depicted in
green, within subjects had a correlation coefficient (mean ± standard deviation) of
r=0.31±0.09, and between subjects of r=0.16±0.05. The two groups significantly differ from
each other, p<0.001, at the significance level of 5%. Error bars represent standard deviation.

Supplementary Figure 3: Node mapping results of all comparisons for different CV
strategies.

875 **Supplementary Figure 4:** Boundary mapping results of all comparisons for different CV

876 strategies

881   **Supplementary Figure 5:** Area under the curve (AUC) of all comparisons for node (left) and

882   boundary mapping (right)

895 **Supplementary Table 1:** Results for the top comparisons under leave-one-out CV for both

896 nodes and boundaries

Column1	Top comparison	Number of voxels	SVM accuracy (95% CI) [%]	SVM AUC (95% CI) [%]
Snowballing peak density	R1 vs. R2	764	74.2 (73.0 – 75.4)	76.4 (74.6 – 78.1)
Boundary mapping	R1 vs. R2	346	68.5 (67.2 – 69.8)	74.5 (72.7 – 86.3)
	R1 vs. R3	174	74.5 (73.4 – 75.6)	80.5 (78.7 – 82.2)

899 **Supplementary Table 2:** Accuracies of the SVM trained on head mean frame-to-frame

900 displacement for real and sham rTMS

	Accuracy Real		Accuracy Sham		Accuracy Real - Sham	
Condition	Mean	Std	Mean	Std	Mean	Std
R0 vs. R1	0.50	0.15	0.50	0.29	0.57	0.27
R0 vs. R2	0.52	0.10	0,54	0.20	0.52	0.31
R0 vs. R3	0.52	0.10	0.50	0.00	0.35	0.23
R1 vs. R2	0.54	0.14	0.54	0.20	0.46	0.20
R1 vs. R3	0.57	0.17	0.54	0.14	0.52	0.23
R2 vs. R3	0.50	0.00	0.50	0.00	0.50	0.00