A conversion from slow to fast memory in response to passive motion

Mousa Javidialsaadi[1], Scott T. Albert[2], Jinsung Wang[1*]

[1]Department of Kinesiology, University of Wisconsin – Milwaukee, Milwaukee, WI, 53201
[2]Depertment of Biomedical Engineering, Johns Hopkins University – Baltimore, MD 21205

Correspondence: Jinsung Wang, 492 Enderis Hall, University of Wisconsin – Milwaukee, WI 53201, USA. Email wang34@uwm.edu. Phone: 414-229-2645

Acknowledgements: This work was supported by a grant from the National Institutes of Health (F32NS095706). We dedicate this article to Dr. Fatemeh Pasand and Dr. Majid Chahardah Cheric for all their academic efforts and teachings. They passed away on 10 April 2018 and 26 June 2020, respectively.

Abstract

When the same perturbation is experienced consecutively, learning is accelerated on the second attempt. This savings is a central property of sensorimotor adaptation. Current models suggest that these improvements in learning are due to changes in the brain's sensitivity to error. Here, we tested whether these increases in error sensitivity could be facilitated by passive movement experiences. In each experimental group, the robot moved the arm passively in the direction that solved the upcoming rotation, but no visual feedback was provided. Then, following a break in time, participants adapted to a visuomotor rotation. Prior passive movements substantially improved motor learning, increasing total compensation in each group by approximately 30%. Similar to savings, a state-space model suggested that this improvement in learning was due to an increase in error sensitivity, but not memory retention. Thus, passive memories appeared to increase the motor learning system's sensitivity to error. However, some features in the observed behavior were not captured by this model, nor by similar empirical models, which assumed that learning was due a single exponential process. When we considered the possibility that learning was supported by parallel fast and slow adaptive processes, a striking pattern emerged; whereas initial improvements in learning were driven by a slower adaptive state, increases in error sensitivity gradually transferred to a faster learning system with the passage of time.

Introduction

In visuomotor rotation experiments, participants actively move a cursor towards an intended target, but the cursor's path is perturbed by a visual rotation (Mazzoni & Krakauer, 2006; Taylor et al., 2014). With training, participants adapt to the visual error, rotating their reach angle in the opposite direction. Like many types of sensorimotor learning, visuomotor adaptation exhibits savings (Haith et al., 2015; Kitago et al., 2013; Morehead et al., 2015; Huberdeau et al., 2015; Huberdeau et al., 2019); participants adapt to a perturbation more rapidly when it has been experienced in the past. On the other hand, when two consecutive perturbations oppose one another, learning during the second period is impaired due to anterograde interference (Brashers-Krug, Shadmehr, & Bizzi, 1996; Sing & Smith, 2010; Tong & Flanagan, 2003; Wigmore, Tong, & Flanagan, 2002). Current models have suggested that these history-dependent changes in learning rate are mediated by one's sensitivity to error (Herzfeld et al., 2014; Albert et al., 2020). When two consecutive perturbations are similar, a given error causes more learning, resulting in savings. However, when two consecutive perturbations are dissimilar, error sensitivity is suppressed, resulting in slower learning (Lerner & Albert et al., 2020). Do these history-dependent changes in error sensitivity require active movement, or can they be evoked by passive experience?

Recent studies have suggested that passively experiencing a visuomotor perturbation results in
learning which transfers to active movement (Lei et al., 2016; Bao et al., 2017; Lei et al., 2017; Tays et
al., 2020). That is, even the passive experience of a sensorimotor perturbation might be effective in
altering subsequent motor skills. For example, recent studies using robot-assisted motor experiences
suggest that passive robotic interventions can facilitate the acquisition of novel motor skills
(Reinkensmeyer and Patton, 2009; Bara and Gentaz, 2011; Basteris et al., 2012; Beets et al., 2012) and
might also improve motor function in hemiparetic stroke patients (Aisen et al., 1997; Krebs et al., 1998;
Riener et al., 2005; Kahn et al., 2006; Vergaro et al., 2010). Yet, while passive experience may help to
facilitate motor recovery (Sakamoto et al., 2012; Tays et al., 2020), little is known regarding how these
passive experiences cause learning, and how this learning is expressed during active movement.

Current state-space models of learning (Smith et al., 2006; McDougle et al., 2015; Albert et al.,
2018) posit that adaption is due to two opposing forces: error-based learning and trial-to-trial forgetting.
In principle, passive training could enhance learning either by strengthening a motor memory's
retention, or by increasing one's sensitivity to error. However, changes in learning parameters such as
error sensitivity are thought to be driven by sensory prediction errors (Popa & Ebner, 2019; Shadmehr et
al., 2010) that accompany active movements; the brain predicts the sensory consequences of its motor
plan via a forward model and then compares this to the observed sensory feedback. Repeated exposure
to these consistent prediction errors appears to upregulate sensitivity to those errors (Gonzalez-Castro
et al., 2014; Herzfeld et al., 2014; Leow et al., 2016; Albert et al., 2020). But in a more general sense,
prediction errors do not necessarily require movement. For example, in classical conditioning, these
errors are driven directly by unexpected sensory stimuli (Ohmae & Medina, 2015; Kim et al., 2020; Ito,
2007; Sears & Steinmetz, 1991; Rasmussen et al, 2008). Here, we wondered if passive movements could
generate prediction errors in a similar manner, and if consistent exposure to these errors might increase
sensitivity to error during active motor learning.

To test this, we passively moved the hand to an unexpected location in the absence of visual
feedback, prior to exposure to a visuomotor rotation. This passive movement resulted in a
proprioceptive mismatch between the hand and the visual target. In addition, though this robotic
intervention created proprioceptive errors during the passive period, the ultimate hand position
provided the solution to the upcoming visual rotation. As in earlier studies, we observed that this
manipulation facilitated active motor learning (Lei et al., 2016; Bao et al., 2017; Lei et al., 2017; Tays et
al., 2020). When we applied a standard single-module state-space model to these data, the model
predicted that improvements in learning were driven by changes in error sensitivity, not retention rates.

Yet, some features in the observed behavior were not captured by this model, nor by similar empirical

models, which assumed that learning was due a single exponential process.

On the contrary, motor learning appears to be supported by at least two parallel systems: a slow

system that responds little to error but retains its adaptation over time, and a fast system that responds

strongly to error but decays rapidly (Smith et al., 2006; Sing & Smith, 2010; Albert et al., 2018; Coltman

et al., 2019; Orozco et al., 2020). When we applied this model to the measured data, the model

suggested that improvements in learning were driven by a slower adaptive state. However, when we

extended the time between the passive training and active motor learning (to 24 hr), improvements in

the slow state of adaptation appeared to transfer to a faster learning process. Interestingly, these

changes in learning were also accompanied with biases in reach angles that appeared prior to

perturbation onset. Thus, passive errors engaged both motor control and motor learning systems, but

the resulting behavioral patterns migrated between slow and fast adaptive circuits as the passive

memory consolidated.

Methods

Here we tested how passively experiencing an error alters the process of motor adaptation. Participants

were divided into several groups. Each group experienced a passive movement period, a break in time,

and then an active adaptation period. By varying the duration of the break between passive and active

movement periods, we measured how memories created by passive errors consolidated over time.

Participants

Twenty-eight healthy volunteers (17 males, 11 females; aged: 18-35) participated in this study. All

subjects were right-handed as assessed by the Edinburgh handedness inventory (Oldfield, 1971). Each

subject signed a consent form that was approved by the University of Wisconsin-Milwaukee Institutional

Review Board. Participants were randomly assigned to one of three experimental groups or a control

group.

Apparatus

Participants were seated in a robotic exoskeleton (KIMARM, BKIN Technologies Ltd, Kingston, ON,

Canada) that provided gravitational support to both arms. The exoskeleton was positioned so that the

participant's arm was hidden underneath a horizontal display (Fig. 1A). To track the hand's position, a

small cursor was projected onto the display, over the participant's index fingertip. During each trial, the

KINARM projected visual stimuli (a start position and a target) onto the display, so that they appeared to

be within the same plane as the arm. The visual stimuli consisted of a centrally located start circle (2 cm

in diameter) and one of four target circles (2cm in diameter) located 10 cm away from the start target

(Fig. 1B). We sampled the hand's position in the x-y plane at 1000 Hz. Position data were low pass

filtered at 15 Hz, and then differentiated to obtain velocity. All post-processing, analysis, and modeling

was conducted in MATLAB R2018a (The MathWorks Inc., Natick, MA).

Experimental Design

Participants were assigned to one of three experimental groups (n=7/group). Each experimental group

experienced three separate conditions: (1) a passive movement period, (2) a baseline reaching period,

and (3) a visuomotor rotation period (Fig. 1C). To determine how passive movements altered reaching

behavior (in the baseline and rotation periods), we compared behavior in the experimental groups to

that of a control group (n=7) that did not experience the passive movement period prior to active

reaching.

In the passive movement period, the KINARM moved the participant's right arm along a straight

10 cm minimum jerk trajectory. On each trial, a visual target was displayed, but the robot passively

moved the arm along a path that was rotated 30° clockwise (CW) to the visual target (Fig. 1B, left).

Critically, this CW rotation served as the "solution" to a counterclockwise (CCW) visuomotor rotation

that participants had not yet experienced (see below). Participants were told to relax their arm and not

resist the passive motion. The passive period consisted of 30 cycles (4 trials in a cycle, 1 to each target in

a pseudorandom order). No visual feedback was provided on these trials. Thus, participants received

only proprioceptive information about the error between their arm's path and the visual target. The

passive movement period terminated with a time-delay which distinguished each experimental group:

5-min delay (n=7), 1 hr delay (n=7), or 24 hr delay (n=7). Once the delay concluded, an active baseline

reaching period began.

In the baseline period, participants moved their arm to each target over a 10-cycle

(experimental groups) or 15-cycle (control group) reaching period. On each trial, continuous visual

feedback was provided via a cursor over the index fingertip. Participants were instructed to move

rapidly and accurately to the target location. The trial ended 1.5 sec after the target was presented. To

begin the next trial, the participant had to move their hand back to the central start position.

The experiment ended with an active visuomotor rotation period that lasted 20 cycles (80 trials

total). On rotation trials, the cursor was rotated 30° CCW to the hand path (Fig. 1B, right). As noted

above, the "solution" to this rotation would be to counter-rotate the reach path by 30° CW relative to

the target. Critically, this CW rotation would coincide with the movement path experienced during the

passive period. Thus, participants in each experimental had been given the proprioceptive experience of

this "solution", though this was never explicitly revealed to each participant. Participants in the control

group, however, had never experienced the passive movement period. Thus, there was no prior

memory to draw upon during the baseline and rotation periods.

Empirical Data Analysis

Data analysis was performed using MATLAB (Mathworks, Natick, MA). The pointing angle at the

reaching movement's midpoint (5 cm displacement) was used as our performance measure, calculated

as the hand's angular position relative to the line segment connecting the start and target positions.

Data were then averaged within each 4-trial cycle. Only cycled data were used in our analyses.

Here we considered how the passive movement period altered reaching behavior during both

the baseline and rotation periods. To investigate the baseline reach period, we averaged the reach angle

over the initial 10 cycles of the baseline period. To assess the rotation period, we considered both early

and late adaptation measures. To quantify early adaptation, we isolated the reach angle on the first

rotation cycle and second rotation cycle. To quantify late adaptation, we averaged the reach angle over

the last 10 rotation cycles. Lastly, we also computed each subject's overall learning rate via an

exponential function that tracked how reach angle (y) changed over each rotation cycle (t, starting at 0):

163
$$y^{(t)} = \alpha e^{-\beta t} + c \qquad (1)$$

Here α and c determine the initial reach angle and asymptotic reach angle. The parameter β represents

the participant's learning rate during the adaptation period. We fit this exponential function to each

participant's data in the least-squares sense using *fmincon* in MATLAB R2018a. We repeated the fitting

procedure 20 times, each time varying the initial parameter guess that seeded the algorithm. We

selected the model parameters which minimized squared error over all 20 repetitions.

State-space learning model: one-state

Although the exponential function closely approximates the decay of motor error during adaptation to a

perturbation, its learning rate parameter reflects a mixture of cycle-by-cycle forgetting and error-based

learning (Lerner & Albert, et al. 2020). Thus, to better understand the adaptation process, we used a

state-space model (Smith et al., 2006). The state-space model posits that learning consists of cycle-to-

cycle error-based learning as well as cycle-by-cycle memory retention (i.e., forgetting). The forgetting

process is controlled by a retention factor (a) which specifies how much adaptation is retained from one

cycle to the next. The learning process is controlled by the participant's error sensitivity (b), which

specifies how much is learned from a given error (e). These processes together determine how the

participant's internal state (x) changes over time, in the presence of internal state noise (ε_x, normal with

zero mean, SD=σ_x):

$$x^{(t+1)} = ax^{(t)} + be^{(t)} + \varepsilon_x^{(t)} \tag{2}$$

182   Eq. (2) allows us to ascribe any differences in performance during the adaptation period to meaningful

183   quantities: retention (a) and error sensitivity (b).

184   Note however, that the internal state (x) is not a measurable quantity. Rather, on each cycle, the

185   motor output (reach angle) is measured. This reach angle (y) directly reflects the subject's internal state,

186   but is corrupted by execution noise (ε_x, normal with zero mean, SD=σ_y) according to:

$$y^{(t)} = x^{(t)} + \varepsilon_y^{(t)} \tag{3}$$

Thus, Eqs. (2) and (3) represent a single module state-space model. We fit this model to each

participant's reach angles during the adaptation period using the Expectation-Maximization (EM)

algorithm (Albert & Shadmehr, 2018). EM is an algorithm the conducts maximum likelihood estimation

in an iterative process. We used EM to identify the model parameters {a, b, x_1, σ_x, σ_y, σ_1} that maximized

the likelihood of observing the data (note that x_1 and σ_1 represent the participant's initial state and

variance, respectively). We conducted 10 iterations, each time changing the initial parameter guess that

seeded the algorithm. We selected the parameter set that maximized the likelihood function across all

10 iterations.

State-space learning model: two-state

The single-module state-space model describes learning as a single adaptive process. Motor adaptation,

however, is believed to be comprised of multiple states, each with different timescales of learning.

These states appear to be accurately summarized as a parallel two-state system, with adaptation

supported by a slow adaptive process and a fast adaptive process (Smith et al., 2006; McDougle et al.,

2015; Albert & Shadmehr, 2018; Coltman et al., 2019). The slow process exhibits slower error-based

learning, but strong retention over time. The fast process exhibits faster error-based learning, but higher

rates of forgetting. We wondered how these adaptive states may contribute to the consolidation of

passive motor memory.

To answer this question, we fit a standard two-state model of learning to individual participant

behavior. In this model, slow state and fast state adaptation are controlled by slow state and fast state

retention factors (a_s and a_f, respectively), as well as slow state and fast state error sensitivities (b_s and b_f,

respectively). As in Eq. (1), both internal states exhibit cycle-by-cycle learning and forgetting, in the

presence of internal state noise ($\varepsilon_{x,s}$ and $\varepsilon_{x,f}$, normal with zero mean, SD=σ_x):

$$x_s^{(t+1)} = a_s x^{(t)} + b_s e^{(t)} + \varepsilon_{x,s}^{(t)}$$
$$x_f^{(t+1)} = a_f x^{(t)} + b_f e^{(t)} + \varepsilon_{x,f}^{(t)}$$

(4)

212   To enforce the traditional two-state dynamics (slow state has higher retention, fast state has higher

213   error sensitivity), retention factors and error sensitivities in Eq. (4) are constrained such that $a_s > a_f$ and

214    $b_f > b_s$. As with the single-module state-space model, the two adaptive are not directly measurable. But

215   together, they sum to produce the overall adapted reach angle, which is also corrupted by execution

216   noise:

$$y^{(t)} = x_s^{(t)} + x_f^{(t)} + \varepsilon_y^{(t)}$$

(5)

Together, Eqs. (4) and (5) with the inequality constraints relating a_s, a_f, b_s, and b_f constitute the

two-state model of learning. The full parameter set consists of: $\{a_s, a_f, b_s, b_f, x_{s,1}, x_{f,1}, \sigma_x, \sigma_y, \sigma_1\}$. Note

that $x_{s,1}$, and $x_{f,1}$ are the initial slow and fast state magnitudes, which are also estimated by the model.

We fit this model to individual participant data, using the EM algorithm (Albert & Shadmehr,

2018). Note that one's ability to fit the two-state model greatly benefits from multiple trial conditions,

such as set breaks, error-clamp periods, washout periods, and perturbation reversals (Albert &

Shadmehr, 2018). However, the current experiment consisted of only perturbation trials with one

orientation. Therefore, to improve our ability to robustly recover the two-state model parameters, we

used a two-tiered fitting procedure (described below).

In our single-state model, we found that error sensitivity varied across experimental and control

groups, but not retention. Therefore, to improve our ability to recover slow and fast error sensitivity, we

started by constraining the model's retention factors (hypothesizing that these were not impacted by

the passive movement period). To do this, we started by fitting the model to mean behavior in the 5

min, 1 hr, 24 hr, and control groups. Then we calculated the midpoint in the range spanned by slow and

fast retention factors identified in each group. Lastly, we then fit the two-state model to individual

participant behavior in each group, after fixing the slow and fast retention factors to these values.

When fitting the two-state model to either group behavior or individual participant behavior, we

performed 10 iterations of the EM algorithm, each time varying the initial parameter guess that seeds

the algorithm. We then selected the parameter set computed by EM with the greatest likelihood.

Statistics

Statistical tests such as one-way ANOVA were carried in MATLAB R2018a. For post-hoc testing following

ANOVA, we used Dunnett's test to assess whether the mean behavior in the 5 min, 1 hr, and 24 hr

differed from the control group. When testing for differences in slow and fast error sensitivities in our

two-state model, we used the Kruskal-Wallis test to evaluate difference in medians. This test was used

over one-way ANOVA due to the presence of a slow state error sensitivity outlier in the 1 hr group, and

the small group sizes used in each group. For post-hoc testing following Kruskal-Wallis, we used Dunn's

test in IBM SPSS 25 to test for difference in median error sensitivity in the 5 min, 1 hr, and 24 hr groups

against control.

Results

When people adapt to two similar perturbations consecutively, re-adaptation evokes a hallmark of

sensorimotor adaptation called savings: more rapid learning during the second exposure to a

perturbation (Smith et al., 2006; Zarahn et al., 2008; Leow et al., 2013; Haith et al., 2015; Day et al.,

2018; Coltman et al., 2020; Yin & Wei, 2020). These changes in learning are thought to be mediated by

error sensitivity (Gonzalez-Castro & Hadjiosif, 2014; Herzfeld et al., 2014; Coltman et al., 2020), whereby

the trial-by-trial experience of consistent errors makes the brain more sensitive to those errors in the

future (Herzfeld et al., 2014; Leow et al., 2016; Albert et al., 2020). Do these changes in the brain's

learning systems require errors to be actively experienced, or can learning be improved by passive

experiences as well?

Here we tested this possibility by exposing participants to a passive movement period, prior to

exposing them to a visuomotor rotation (Fig. 1C). In the passive period, the robot moved the

participant's arm towards a target, without any visual feedback (Fig. 1A; Fig. 1B, left). But rather than

move the arm directly to the visual target, the robot deviated along a path rotated clockwise by 30°.

Thus, consistent proprioceptive "errors" were passively delivered to the participant without them ever

producing an active movement. Following a break in time (Fig. 1C, break; 5 min, 1 hr, or 24 hr)

participants then returned and produced active movements to the same targets, first in a baseline

condition (Fig. 1C, baseline), and then under the influence of a 30° CCW visuomotor rotation (Fig. 1B,

right; Fig. 1C, rotation). The perturbation's orientation was chosen so that the passive movements

provided the "solution" to the rotation that had not yet occurred. Is it possible that this passive memory

could facilitate learning during active movement?

To answer this question, we compared reaching movements during the baseline and adaptation

periods (Fig. 1D, black), to those of a control group that never experienced passive training (Fig. 1D,

gray). By measuring how behavior changed with the time-delay between passive and active periods, we

investigated how memory acquired passively was consolidated and expressed over time.

Figure 1. Passive and active motor paradigm. **A.** Participants (n=28) were seated in a KINARM exoskeleton which
each arm supported against gravity. Their arm was placed underneath a horizontal display. **B.** During the passive
movement period (left) participants were shown a target at either 45°, 135°, 225°, 315°, but the KINARM passively
moved the hand along a path that was rotated 30° clockwise (CW) to the visual target. No visual feedback of hand
position was provided. During the active learning period (right) participants reached to the same targets, but the
cursor was rotated counterclockwise (CCW) by 30°. Thus, to solve this rotation, participants needed to move along
the same path traversed during the passive manipulation (solution shown in dashed line. **C.** The experiment
started with 30 cycles (4 trials to each target) of passive movements. Then participants took either a 5 min, 1 hr, or
24 hr break. When participants returned to the experiment, they completed 10 cycles of baseline reaching
movements, followed by 20 visuomotor rotation cycles. **D.** We calculated the reach angle between the hand and
the straight-line path between the start and target cursor. The reach angle was calculated at the movement's
midpoint (5 cm displacement). Reach angles for the control group (no passive training) are shown in gray. Reach
angles for the 5 min group (left), 1 hr group (middle), and 24 hr group (right) are shown in black. Error bars show
mean ± SEM.

Consolidating passive movement experiences biases reaching commands

Active movement periods in the experimental and control groups began with a series of baseline

reaching movements where participants reached with veridical visual feedback (Fig. 1D, baseline). Were

these initial baseline movements altered by the passive experience of proprioceptive error?

To answer this question, we isolated baseline reaching behavior in each group (Fig. 2A). Without

any prior exposure to passive movements, the control group exhibited a counterclockwise (negative)

bias in baseline reach angle, which was likely caused by the inertial properties of the arm and robotic

apparatus (Fig. 2B, control; one-sample t-test, t(6)=-5.38, p=0.002). Remarkably, this bias was gradually

lifted in the experimental groups (Fig. 2B; one-way ANOVA, $F(3,24)=3.2$, $p=0.0415$), with the 24 hr group

showing a statistically significant reduction in bias relative to the control group (post-hoc Dunnett's test,

24 hr vs. control, $p=0.019$).

Critically, this change in baseline reach angle was directed towards past locations where the arm

had been passively moved (i.e., "rotation solution" in Fig. 2). In other words, whereas subjects without

passive training produced reaching movements that deviated slightly CCW, subjects with passive

training where the robot moved their arm CW, exhibited biases that were shifted in that direction.

Because this change in reach angle was only present in the 24 hr group, it appeared that this shift in

baseline reach angle required long-term consolidation. Moreover, the data suggested a clear trend (Fig.

2B) where changes in baseline reach angle gradually strengthened with increased time-delay between

passive and active periods (though with our small group sizes, we did not have enough power to

statistically confirm these gradual changes).

310
311
Figure 2. Passive training evoked a time-dependent bias in initial reach angle. **A.** Here we show reach angles during
the 10 baseline cycles in the active movement period. The control group (no passive training) is shown in gray. The
experimental groups (5 min, 1 hr, and 24 hr) are shown in black. Note that the control group exhibited a slight
negative (counterclockwise) bias in reach angle at the movement's midpoint. This was likely due to the inertial
properties of the arm and robotic system. However, this bias appeared to change in the experimental groups. The
direction of this shift (upwards in the graph) is directed towards the solution of the upcoming rotation (which is
also the direction along which the hand was passively moved during the preceding passive movement period). **B.**
We calculated the mean reach angle during the baseline period. Statistics show post-hoc Tukey's test following
one-way ANOVA (* indicates $p<0.05$). Error bars show mean ± SEM.

In summary, the naïve assumption might be that the reach system produces commands which

are heavily dependent on visual stimuli. Surprisingly however, it appears that when the hand is passively

moved to a location, this proprioceptive experience gets paired with visual targets; later, the reach

commands actively produced to move towards the target, are computed via some weighting between
the target's visual location and the past proprioceptive memory. This proprioceptive weighting appears
to require consolidation, arising only after long periods of delay. Next, we investigated whether this
consolidated proprioceptive memory alters the process of motor adaptation.
329
Passive movement experiences enhance motor adaptation
Following the baseline reaching period, all participants were abruptly exposed to a 30° CCW rotation
(Fig. 1C, rotation). The perturbation's orientation was selected such that rotation's "solution" (counter-
rotate clockwise) mirrored the passive movement direction participants had experienced in the past
(Fig. 1B, right). Did this passive experience somehow facilitate or obstruct the process of motor
learning?
Fig. 3A shows the rotation period reach angle in the experimental and control groups (positive
values denote greater adaptation). Remarkably, passive training appeared to greatly facilitate
adaptation in all three experimental groups (Fig. 3B, mean reach angle; one-way ANOVA, $F_{(3,24)}=3.98$,
$p=0.02$; post-hoc Dunnett's test against control, 5 min: $p=0.021$, 1 hr: $p=0.021$, 24 hr: $p=0.038$),
increasing average compensation by approximately 30%.
At what point did this improvement appear? To answer this question, we analyzed the reach
angle on the first two cycles. Though the reach angle on the first cycle appeared slightly larger in the
experimental groups (Fig. 3C), this trend was not statistically significant (one-way ANOVA $F_{(3,24)}=1.38$,
$p=0.274$) and could have been biased by lingering differences in baseline reach behavior (Fig. 2).
However, by the second reaching cycle (Fig. 3D) adaptation in each group surpassed that of the control
group (one-way ANOVA, $F_{(3,24)}=4.59$, $p=0.011$; post-hoc Dunnett's test against control, 5 min: $p=0.027$,
1 hr: $p=0.039$, 24 hr: $p=0.006$). This analysis suggested that enhancements in learning developed quite
rapidly, and unlike changes in baseline behavior (Fig. 2), did not require long-term consolidation of the
passive training period (considering the improvement was present even in the 5 min group).
All evidence pointed towards the possibility that passive training increased the rate of learning
during the rotation period. To mathematically test this idea, we fit an exponential curve (Eq. (1)) to the
adaptation period reach angles (dashed lines in Fig. 3A). Indeed, the exponential model suggested that
passive training increased the rate of learning during the rotation (Fig. 3E, one-way ANOVA,
$F_{(3,24)}=3.48$, $p=0.031$). Surprisingly however, this increased rate was not exhibited by each group; only
the 24 hr group showed a statistically significant increase in learning rate over the control group (post-
hoc Dunnett's test against control: 5 min: $p=0.99$, 1 hr: $p=0.3$, 24 hr: $p=0.046$).

Figure 3. Passive training enhances active motor learning. **A.** Here we show reach angles during the active rotation learning cycles. The control group (no passive training) is shown in gray. The experimental groups (5 min, 1 hr, and 24 hr) are shown in black. Dashed black lines show an exponential fit to the data (mean across individual subjects). **B.** We quantified the mean reach angle across the 20 rotation cycles. All experimental groups showed greater adaptation than the control group. **C.** We measured the reach angle on the initial adaptation cycle. **D.** We measured the reach angle on the second adaptation cycle. All experimental groups showed greater adaptation than the control group on this cycle. **E.** We quantified each participant's learning rate with a 3-paremeter exponential model. In each panel, statistics show post-hoc Tukey's test following one-way ANOVA (* indicates p<0.05, ** indicates p<0.01). Error bars show mean ± SEM.

Thus, our analysis presented a puzzle. On one hand, passive training improved overall adaptation (Fig. 3B), with improvements evident even on the second rotation cycle (Fig. 3D) in each group. Therefore, this facilitation did not appear to require long-term consolidation. However, the rate of learning (assessed via an exponential curve) was only enhanced in the 24 hr group (Fig. 3E) suggesting that the improvement in learning did in fact require long-term consolidation. How are we to rectify this contradiction? One idea is that there are in fact improvements in learning across all experimental groups, but this improvement is only partially captured by a model-free exponential curve.

Passive movement experiences enhance sensitivity to error, but not retention

One issue with the exponential model, is that is does not parse behavior into interpretable physiologic
processes. That is, motor adaptation is known to be governed by at least two critical processes: error-
based learning and trial-by-trial forgetting. While these two processes are mixed together within an
exponential curve, they are more easily recovered by a state-space model of learning. The state-space
model of learning (Eqs. (2) & (3)) posits that adaptation is due to learning and forgetting events, which
are controlled by one's sensitivity to error (b) and retention (a), respectively. Might passive training alter
one (or both) of these properties, thus producing the facilitation in adaptation noted in Fig. 3?

To answer this question, we fit a single-module state-space model to individual participant reach
angles in the experimental and control groups. The model (Fig. 4A, dashed lines) appeared to closely
track each group's learning curve. Next, we isolated the retention factor (Fig. 4B) and error sensitivity
(Fig. 4C) predicted by the state-space model. Interestingly, passive training did not appear to alter the
retention processes in any experimental group (Fig. 4B, one-way ANOVA, $F(3,24)=2.19$, $p=0.115$). On the
other hand, passive training clearly impacted error sensitivity (Fig. 4C, one-way ANOVA, $F(3,24)=3.81$,
$p=0.023$), which appeared to grow with time-delay duration; though only the 24 hr group showed a
statistically significant difference relative to the control condition (post-hoc Dunnett's tests against
control, 5 min: $p=0.72$, 1 hr: $p=0.061$, 24 hr: $p=0.017$).

In summary, the state-space model suggested that the passive experience of proprioceptive
error increased sensitivity to visual errors during the active movement period. These changes appeared
quite similar to those observed in savings paradigms, where initial adaptation increases error sensitivity,
but not retention (Herzfeld et al., 2014; Coltman et al., 2019). But the single-module state-space model
still could not solve the puzzle that arose in our empirical analysis of behavior; why is error sensitivity
only increased in the 24 hr group (Fig. 4C), when all the experimental groups show a facilitation in
learning (Figs. 3B&D)?

401
Conversion from slow to fast memory states during passive training consolidation
Both our exponential model (Fig. 3E) and state-space model (Fig. 4C) suggested that changes in
adaptation were specific to the 24 hr group, while the empirical data (Figs. 3B&D) showed clear
improvements in adaptation across all groups (5 min, 1 hr, and 24 hr) exposed to passive errors. This
discrepancy suggested that there were some features in the adaptation curve that were altered by
passive training, which our models could not quite capture. Both the exponential and state-space
models describe learning as a single adaptive process. Could it be that adaptation patterns were truly
the result of multiple states of learning?

To investigate this possibility, we considered a standard two-state model of learning (Smith et

al., 2006). The two-state model posits that adaptation is supported by multiple adaptive states: a slow

state and a fast state which differ in their sensitivity to error and retention. The slow process learns less

due to error, but retains its state strongly over time. The fast process learns more due to error, but is

volatile and decays more rapidly over time. How do the changes in error sensitivity noted in our single-

module state-space model relate to these two parallel adaptive processes?

To answer this question, we fit the two-state model (Eqs. (4) & (5)) to individual participant

reach angles in the experimental and control groups. Using the resultant model parameters, we

simulated the slow (Fig. 4D) and fast (Fig. 4F) states predicted by the model. These states showed a

remarkable trend; immediately after the passive training period in the 5 min group, the slow state of

learning was enhanced, but this facilitation diminished over time (Fig. 4D). On the other hand, the fast

state of learning exhibited the opposite trend; as the time-delay following the passive training period

increased, so too did activity in the fast adaptive process (Fig. 4F).

These changes in the slow and fast states were due to a striking pattern in error sensitivity. The

slow state's error sensitivity increased nearly two-fold following the passive training period in the 5 min

group (Fig. 4E, Kruskal-Wallis test, $X^2(27)=8.32$, p=0.0398) but gradually returned to control levels with

the passage of time following the passive training period (post-hoc Dunn's tests against control, 5 min:

p=0.024, 1 hr: p=1, 24 hr: p=1). On the other hand, the fast-state's error sensitivity increased by 50%

(Fig. 4G, Kruskal-Wallis test, $X^2(27)=9.04$, p=0.0288) but only in the 24 hr group after the passage of time

(post-hoc Dunn's tests against control, 5 min: p=0.486, 1 hr: p=0.087, 24 hr: p=0.012).

Figure 4. Passive training improves motor learning by increasing sensitivity to error. **A.** Here we show reach angles

during the active rotation learning cycles. The control group (no passive training) is shown in gray. The

experimental groups (5 min, 1 hr, and 24 hr) are shown in black. We fit the data with a single-module state-space

model (dashed black line; mean across individual participants). The state-space model posited that adaptation was

due to both error-based learning (controlled by error sensitivity) and trial-by-trial forgetting (controlled by

retention factor). **B.** Here we show the retention factor predicted by the single-module state-space model. **C.** Here

we show the error sensitivity predicted by the single-module state-space model. **D.** We fit a two-state model to

individual participant behavior. The predicted slow state for the experimental groups (5 min, 1 hr, and 24 hr) are

shown in blue; time-delay between passive and active training increases left-to-right. The predicted slow state in

the control group is shown in gray. **E.** Here we provide the slow state error sensitivity predicted by the two-state

model. **F.** Same as in **D**, but for the fast state of learning. **G.** Same as in **E** but for the fast state error sensitivity. In

each panel, statistics show post-hoc Tukey's test (**C**) or Dunn's test (**G**) following one-way ANOVA (**C**) or Kruskal-

Wallis (**E** and **G**). Statistics: * indicates p<0.05, ** indicates p<0.01. Error bars show mean ± SEM.

In summary, the two-state model made an intriguing prediction; adaptation in each
experimental group was enhanced by the passive error experience, but the nature of this enhancement
varied over time. Initially following the passive movement period, error sensitivity had increased in the
slower adaptive process. But with the passage of time, this improvement transferred to the fast state of
adaptation. Thus, as the passive memory consolidated, it appeared that its benefits were converted
from slower learning processes, to faster learning processes.

Discussion

When the same perturbation is experienced consecutively, learning is accelerated on the second
attempt. This savings is a central property of sensorimotor adaptation which is observed across several
motor effectors: the reaching system (Smith et al., 2006; Zarahn et al., 2008; Leow et al., 2013; Haith et
al., 2015; Coltman et al., 2020; Yin & Wei, 2020), walking (Mawase et al., 2014; Day et al., 2018), and
even the oculomotor system (Kojima et al., 2004). Current models have suggested that these
improvements in learning are due to changes in the brain's sensitivity to error (Gonzalez-Castro &
Hadjiosif, 2014; Herzfeld et al., 2014; Mawase et al., 2014; Coltman et al., 2020). Here, we tested
whether these increases in error sensitivity could be facilitated by passive experiences, as opposed to
active movements.

To do this, we used a state-space model to interpret how motor adaptation is altered due to
passive proprioceptive memory. Consistent with earlier work (Lei et al., 2016; Bao et al., 2017; Lei et al.,
2017; Tays et al., 2020) we observed that passive training facilitated active motor adaptation. In each
experimental group, the robot moved the arm passively in the direction that solved the upcoming
rotation, but no visual feedback was provided. Somehow, this passive proprioceptive experience
substantially altered subsequent motor learning, increasing total compensation in each group by
approximately 30%. Similar to savings, the state-space model suggested that this improvement in
learning was due to an increase in error sensitivity. Thus, passive memories appeared to increase the
motor learning system's sensitivity to error.

However, the state-space model's prediction exhibited a puzzle; whereas all experimental
groups showed improvements in total compensation, only the 24 hr group showed a statistically
significant increase in sensitivity to error (Fig. 4C). This peculiarity was also present when we fit an
exponential curve to estimate participant learning rates (Fig. 1E). This suggested that these models were
missing a key component: multiple adaptive states. That is, converging evidence suggests that
sensorimotor learning is supported by parallel processes (Smith et al., 2006; Mawase et al., 2014;

McDougle et al., 2015; Coltman et al., 2020). The parallel adaptive states are well-approximated as a

two-state system: with one slow state the learns slowly but is decay-resistant, and one fast state which

learns rapidly but also forgets rapidly. When we applied this two-state model to our data (Albert &

Shadmehr, 2018), a new pattern emerged; passive training initially facilitated improvements in the slow

state's sensitivity to error (Fig. 4E) which gradually transferred to the fast state of adaptation over time

(Fig. 4G). Thus, while proprioceptive memory immediately enhanced motor learning, these

improvements appeared to gradually consolidate from a slower system, into a faster system.

This putative conversion between learning systems mirrored other changes in behavior

exhibited during the baseline period. Prior to the rotation, participants reached towards each target

without any visual perturbation. These reaches were their very first attempts to produce active

movements with the robotic system. Curiously, previously acquired proprioceptive memory appeared to

alter the motor system during this baseline period, pulling the arm towards the path it travelled during

the passive movement period. This biasing in reach angle, however, was not present in the 5 min group

or 1 hr group, and only emerged with time (in the 24 hr group). Thus, the consolidation of passive

training led to time-dependent changes in both error sensitivity and initial reaching behavior, which may

or may not have shared a common source.

This biasing in initial reaching direction appeared qualitatively similar to use-dependent learning

(Wang et al., 2015; Bao et al., 2017; Lei et al., 2017; Verstynen & Sabes, 2011; Lei et al., 2016;

Diedrichsen et al., 2010; Jax & Rosenbaum, 2007; Scheidt et al., 2005; Diedrichsen, 2007); with use

dependent learning, active (Verstynen & Sabes, 2011) as well as passive (Diedrichsen et al., 2010)

motion induces persistent biases in active movement towards past motor actions or proprioceptive

states. But there is one substantial difference in the reaching patterns we measured and those exhibited

in these earlier studies. Namely, use-dependent biases in motor commands are immediately

precipitated by past active or passive motion: they do not require time to emerge. However, here we

observed that when active movement quickly (5 min break) followed passive training, no use-dependent

biases in reach angle occurred. On the other hand, once considerable time had elapsed (24 hr), passive

training biased initial reach angles during active movement. Why were use dependent changes absent in

the 5 min and 1 hr experimental groups? One possibility is that because the passive movements were

uniformly spread across the unit circle (4 movement directions, with 90° spacing between them),

movements produced a type of motor or proprioceptive averaging, qualitatively similar to the uniformly

distributed reach condition studied by Verstynen & Sabes (2011). Thus, our data propose two possible

ways that passive movements may bias active reaching: one through traditional use-dependent
mechanisms, and another that involves some process of consolidation which emerges with time.

Might these changes in initial reaching bias be linked to the changes in error sensitivity we
measured during the rotation period? Both passive movements (Carel et al., 2000) and active motor
learning (Della-Maggiore & McIntosh, 2005; Herzfeld et al., 2015; Popa et al., 2012; Izawa et al., 2012;
Tseng et al., 2007; Gibo et al., 2013; Maschke et al., 2004) are known to involve both the primary motor
cortex (M1) and the cerebellum. Might it be that interactions between these two areas lead to changes
in error sensitivity and reaching biases? While we can only speculate, one possibility might be that
prediction errors experienced during passive training cause changes in Purkinje cells (P-cells) in the
cerebellar cortex, which rapidly transfer to slower and more robust learning units in the deep cerebellar
nucleus (Herzfeld et al., 2020; Lisberger et al., 1994; McCormick & Thompson, 1984; Perret et al., 1993).
Thus, interplay between P-cells and the deep nuclei, might be responsible for facilitating the changes in
slow state error sensitivity observed early (5 min) after passive training. Next, with passage of time,
these changes may somehow be transferred through an unidentified mechanism to, and consolidated
within, M1. Perhaps it is these delayed changes in M1 that subsequently bias motor commands during
baseline reaching movements, and facilitate the faster learning state observed in the 24 hr group.

But if this is true, what are the errors that engage the cerebellar cortex during passive
movement? While normally it is thought that errors arise with the sensory outcome that follows an
active movement (Herzfeld et al., 2015; Herzfeld et al., 2018), associative learning mechanisms such as
eye-blink condition, lead to sensory-evoked complex spikes within the cerebellar cortex without any
active movement at all (Ohmae & Medina, 2015; Kim et al., 2020; Ito, 201; Sears & Steinmetz, 1991;
Rasmussen et al, 2008). Thus, it is interesting to wonder whether the discrepancy between the visual
(target) and proprioceptive (arm position) state created by our passive intervention, might too have
somehow engaged the cerebellar cortex with a prediction error (Popa & Ebner, 2019; Shadmehr et al.,
2010). If true, the repeated experience of consistent prediction errors may have upregulated error
sensitivity in the cerebellar cortex (Herzfeld et al., 2014; Albert et al., 2020), thus facilitating learning in
the active condition. Alternatively, lingering changes in the somatosensory and motor cortices induced
by passive movements may have indirectly altered cerebellar state throughout adaptation (as opposed
to passive errors driving the cerebellar cortex directly). In either case, rTMS and tDCS over the
cerebellum and M1 may provide a path to elucidating the various timescales of memory that are
engaged by passive training.

Lastly, there is another dichotomy in motor adaptation, which also may relate to the conversion

between slow and fast learning: implicit and explicit adaptation. Motor learning is known to be

supported by two parallel learning systems: a strategic explicit system that can be guided by instruction

(Taylor et al., 2014; Mazzoni & Krakauer, 2006; McDougle & Taylor, 2019), as well as an implicit system

that adapts without our conscious awareness (Avraham et al., 2020; Miyamoto et al., 2020; Javidialsaadi

and Wang., 2020). Might interplay between these two systems relate to the time-dependent error

sensitivity patterns observed during rotation learning? Because we did not measure implicit or explicit

learning, we cannot know the answer to this question. However, we did observe that reaction time (Fig.

S1A) appeared elevated after passive training in the experimental groups, which is known to commonly

accompany cognitive operations (Fernandez-Ruiz et al., 2011; Sakamoto and Kondo, 2012; Anguera et

al., 2010; Georgopoulos and Massey, 1987; McDougle et al., 2019; Albert et al., 2020).

550
551
Figure S1. Changes in reaction time. **A.** We measured reaction time as the difference between target appearance
and reach onset. Here we show reaction time during the perturbation period in the experimental (black) and
control (gray) groups. **B.** We investigated early differences in reaction time by calculating the mean reaction time
over the first three cycles. Statistics show post-hoc Tukey's test following one-way ANOVA (** indicates p<0.01).
Error bars show mean ± SEM.

However, this trend was only statistically significant in the 5 min group (Fig. S1B) and appeared

to dissipate quite rapidly in the 1 hr and 24 hr groups. If explicit strategies did contribute to increases in

error sensitivity, it is also puzzling why reaction time was greatest in the 5 min group (Fig. S1B); this

group was primarily enhanced by a slow learning state (Figs. 4D&E) which is believed to reflect implicit

learning processes (McDougle et al., 2015). Nevertheless, the idea that passive training may evoke

changes in both implicit and explicit processes is a fascinating possibility which remains to be formally

tested.

References

Albert, S. T., & Shadmehr, R. (2018). Estimating properties of the fast and slow adaptive processes during sensorimotor adaptation. *Journal of neurophysiology*, *119*(4), 1367-1393.

Albert, Scott T., Jihoon Jang, Adrian M. Haith, Gonzalo Lerner, Valeria Della-Maggiore, John W. Krakauer, and Reza Shadmehr. "Competition between parallel sensorimotor learning systems." *bioRxiv* (2020).

Albert, S. T. (2020). *The role of error-based learning in movement and stillness* (Doctoral dissertation, Johns Hopkins University).

Aisen, M. L., Krebs, H. I., Hogan, N., McDowell, F., & Volpe, B. T. (1997). The effect of robot-assisted therapy and rehabilitative training on motor recovery following stroke. *Archives of neurology*, *54*(4), 443-446.

Anguera, J. A., Reuter-Lorenz, P. A., Willingham, D. T., & Seidler, R. D. (2010). Contributions of spatial working memory to visuomotor learning. *Journal of cognitive neuroscience*, *22*(9), 1917-1930.

Avraham, G., Morehead, J. R., Kim, H. E., & Ivry, R. B. (2021). Reexposure to a sensorimotor perturbation produces opposite effects on explicit and implicit learning processes. *PLoS biology*, *19*(3), e3001147.

Bao, S., Lei, Y., & Wang, J. (2017). Experiencing a reaching task passively with one arm while adapting to a visuomotor rotation with the other can lead to substantial transfer of motor learning across the arms. *Neuroscience letters*, *638*, 109-113.

Bara, F., & Gentaz, E. (2011). Haptics in teaching handwriting: The role of perceptual and visuo-motor skills. *Human movement science*, *30*(4), 745-759.

Basteris, A., Bracco, L., & Sanguineti, V. (2012). Robot-assisted intermanual transfer of handwriting skills. *Human movement science*, *31*(5), 1175-1190.

Beets, I. A., Macé, M., Meesen, R. L., Cuypers, K., Levin, O., & Swinnen, S. P. (2012). Active versus passive training of a complex bimanual task: is prescriptive proprioceptive information sufficient for inducing motor learning?. *PLoS One*, *7*(5), e37687.

Brashers-Krug, T., Shadmehr, R., & Bizzi, E. (1996). Consolidation in human motor memory. *Nature*, *382*(6588), 252-255.

Carel, C., Loubinoux, I., Boulanouar, K., Manelfe, C., Rascol, O., Celsis, P., & Chollet, F. (2000). Neural substrate for the effects of passive training on sensorimotor cortical representation: a study with functional magnetic resonance imaging in healthy subjects. *Journal of Cerebral Blood Flow & Metabolism*, *20*(3), 478-484.

Castro, L. N. G., Hadjiosif, A. M., Hemphill, M. A., & Smith, M. A. (2014). Environmental consistency
determines the rate of motor adaptation. *Current Biology*, *24*(10), 1050-1061.

Coltman, S. K., Cashaback, J. G., & Gribble, P. L. (2019). Both fast and slow learning processes contribute
to savings following sensorimotor adaptation. *Journal of neurophysiology*, *121*(4), 1575-1583.

Coltman, S. K., & Gribble, P. L. (2020). Time course of changes in the long-latency feedback response
parallels the fast process of short-term motor adaptation. *Journal of Neurophysiology*, *124*(2), 388-399.

Day, K. A., Leech, K. A., Roemmich, R. T., & Bastian, A. J. (2018). Accelerating locomotor savings in
learning: compressing four training days to one. *Journal of neurophysiology*, *119*(6), 2100-2113.

Della-Maggiore, V., & McIntosh, A. R. (2005). Time course of changes in brain activity and functional
connectivity associated with long-term adaptation to a rotational transformation. *Journal of*
neurophysiology, *93*(4), 2254-2262.

Diedrichsen, J., Shadmehr, R., & Ivry, R. B. (2010). The coordination of movement: optimal feedback
control and beyond. *Trends in cognitive sciences*, *14*(1), 31-39.

Diedrichsen, J., Criscimagna-Hemminger, S. E., & Shadmehr, R. (2007). Dissociating timing and
coordination as functions of the cerebellum. *Journal of Neuroscience*, *27*(23), 6291-6301.

Fernandez-Ruiz, J., Wong, W., Armstrong, I. T., & Flanagan, J. R. (2011). Relation between reaction time
and reach errors during visuomotor adaptation. *Behavioural brain research*, *219*(1), 8-14.

Georgopoulos, A. P., & Massey, J. T. (1987). Cognitive spatial-motor processes. *Experimental brain*
research, *65*(2), 361-370.

Gibo, T. L., Criscimagna-Hemminger, S. E., Okamura, A. M., & Bastian, A. J. (2013). Cerebellar motor
learning: are environment dynamics more important than error size?. *Journal of*
neurophysiology, *110*(2), 322-333.

Haith AM, Huberdeau DM, Krakauer JW. The influence of movement preparation time on the expression
of visuomotor learning and savings. J Neurosci 35: 5109–5117, 2015.

Herzfeld, D. J., Hall, N. J., Tringides, M., & Lisberger, S. G. (2020). Principles of operation of a cerebellar
learning circuit. *Elife*, *9*, e55217.

Herzfeld, D. J., Kojima, Y., Soetedjo, R., & Shadmehr, R. (2018). Encoding of error and learning to correct
that error by the Purkinje cells of the cerebellum. *Nature neuroscience*, *21*(5), 736-743.

Herzfeld DJ, Vaswani PA, Marko MK, Shadmehr R. A memory of errors in sensorimotor learning. Science
345: 1349–1353, 2014.

Herzfeld, D. J., Kojima, Y., Soetedjo, R., & Shadmehr, R. (2015). Encoding of action by the Purkinje cells of
the cerebellum. *Nature*, *526*(7573), 439-442.

Huberdeau, D. M., Haith, A. M., & Krakauer, J. W. (2015). Formation of a long-term memory for
visuomotor adaptation following only a few trials of practice. *Journal of neurophysiology*, *114*(2), 969-
977.

Huberdeau, D. M., Krakauer, J. W., & Haith, A. M. (2019). Practice induces a qualitative change in the
memory representation for visuomotor learning. *Journal of neurophysiology*, *122*(3), 1050-1059.

Ito, K., Doi, M., Kondo, T.: Feedforward Adaptation to a Varying Dynamical Environment during Reaching
Movements, Journal of Robotics Mechatronics, 19, 4, 474–481 (2007).

Izawa, J., Criscimagna-Hemminger, S. E., & Shadmehr, R. (2012). Cerebellar contributions to reach
adaptation and learning sensory consequences of action. *Journal of Neuroscience*, *32*(12), 4230-4239.

Javidialsaadi, M., & Wang, J. (2021). Lack of interlimb transfer following visuomotor adaptation in a
person with congenital mirror movements despite the awareness of the visuomotor perturbation. *Brain
and Cognition*, *147*, 105653.

Jax, S. A., & Rosenbaum, D. A. (2007). Hand path priming in manual obstacle avoidance: evidence that
the dorsal stream does not only control visually guided actions in real time. *Journal of Experimental
Psychology: Human Perception and Performance*, *33*(2), 425.

Kahn, L. E., Zygman, M. L., Rymer, W. Z., & Reinkensmeyer, D. J. (2006). Robot-assisted reaching exercise
promotes arm movement recovery in chronic hemiparetic stroke: a randomized controlled pilot
study. *Journal of neuroengineering and rehabilitation*, *3*(1), 1-13.

Kim, H. E., Avraham, G., & Ivry, R. B. (2020). The psychology of reaching: action selection, movement
implementation, and sensorimotor learning. *Annual review of psychology*, *72*.

Kitago T, Ryan SL, Mazzoni P, Krakauer JW, Haith AM. Unlearning versus savings in visuomotor
adaptation: comparing effects of washout, passage of time, and removal of errors on motor memory.
Front Hum Neurosci 7: 307, 2013.

Kojima Y, Iwamoto Y, Yoshida K. Memory of learning facilitates saccadic adaptation in the monkey. *J
Neurosci* 24: 7531–7539, 2004.

Krebs, H. I., Hogan, N., Aisen, M. L., & Volpe, B. T. (1998). Robot-aided neurorehabilitation. *IEEE
transactions on rehabilitation engineering*, *6*(1), 75-87.

Lei, Y., Bao, S., & Wang, J. (2016). The combined effects of action observation and passive proprioceptive
training on adaptive motor learning. *Neuroscience*, *331*, 91-98.

Lei, Y., Bao, S., Perez, M. A., & Wang, J. (2017). Enhancing generalization of visuomotor adaptation by
inducing use-dependent learning. *Neuroscience*, *366*, 184-195.

Leow LA, de Rugy A, Marinovic W, Riek S, Carroll TJ. Savings for visuomotor adaptation require prior
history of error, not prior repetition of successful actions. *J Neurophysiol* 116: 1603–1614, 2016.

Leow, L. A., De Rugy, A., Loftus, A. M., & Hammond, G. (2013). Different mechanisms contributing to
savings and anterograde interference are impaired in Parkinson's disease. *Frontiers in human*
neuroscience, *7*, 55.

Lerner, G., Albert, S., Caffaro, P. A., Villalta, J. I., Jacobacci, F., Shadmehr, R., & Della-Maggiore, V. (2020).
The origins of anterograde interference in visuomotor adaptation. *Cerebral Cortex*, *30*(7), 4000-4010.

Lisberger, S. G. (1994). Neural basis for motor learning in the vestibuloocular reflex of primates. III.
Computational and behavioral analysis of the sites of learning. *Journal of Neurophysiology*, *72*(2), 974-
998.

Maschke, M., Gomez, C. M., Ebner, T. J., & Konczak, J. (2004). Hereditary cerebellar ataxia progressively
impairs force adaptation during goal-directed arm movements. *Journal of neurophysiology*, *91*(1), 230-
238.

Mawase, F., Shmuelof, L., Bar-Haim, S., & Karniel, A. (2014). Savings in locomotor adaptation explained
by changes in learning parameters following initial adaptation. *Journal of neurophysiology*, *111*(7), 1444-
1454.

Mazzoni, P., & Krakauer, J. W., (2006). An implicit plan overrides an explicit strategy during visuomotor
adaptation. Journal of neuroscience, 26(14), 3642-3645.

McCormick, D. A., & Thompson, R. F. (1984). Neuronal responses of the rabbit cerebellum during
acquisition and performance of a classically conditioned nictitating membrane-eyelid response. *Journal*
of Neuroscience, *4*(11), 2811-2822.

McDougle, S. D., Bond, K. M., & Taylor, J. A., (2015). Explicit and implicit processes constitute the fast
and slow processes of sensorimotor learning. Journal of Neuroscience, 35(26), 9568-9579.

McDougle, S. D., & Taylor, J. A. (2019). Dissociable cognitive strategies for sensorimotor learning. *Nature*
communications, *10*(1), 1-13.

Miyamoto, Y. R., Wang, S., & Smith, M. A. (2020). Implicit adaptation compensates for erratic explicit
strategy in human motor learning. *Nature neuroscience*, *23*(3), 443-455.

Morehead, J. Ryan, Salman E. Qasim, Matthew J. Crossley, and Richard Ivry. "Savings upon re-aiming in
visuomotor adaptation." *Journal of neuroscience* 35, no. 42 (2015): 14386-14396.

Ohmae, S., & Medina, J. F. (2015). Climbing fibers encode a temporal-difference prediction error during
cerebellar learning in mice. *Nature neuroscience*, *18*(12), 1798-1803.

Orozco, S. P., Albert, S. T., & Shadmehr, R. (2020). Spontaneous recovery and the multiple timescales of
human motor memory. *bioRxiv*.

Perrett, S. P., Ruiz, B. P., & Mauk, M. D. (1993). Cerebellar cortex lesions disrupt learning-dependent
timing of conditioned eyelid responses. *Journal of Neuroscience*, *13*(4), 1708-1718.

Popa, L. S., & Ebner, T. J. (2019). Cerebellum, predictions and errors. *Frontiers in cellular*
neuroscience, *12*, 524.

Popa, L. S., Hewitt, A. L., & Ebner, T. J. (2012). Predictive and feedback performance errors are signaled
in the simple spike discharge of individual Purkinje cells. *Journal of Neuroscience*, *32*(44), 15345-15358.

Reinkensmeyer, D. J., & Patton, J. L. (2009). Can robots help the learning of skilled actions?. *Exercise and*
sport sciences reviews, *37*(1), 43.

Riener, R., Nef, T., & Colombo, G. (2005). Robot-aided neurorehabilitation of the upper
extremities. *Medical and biological engineering and computing*, *43*(1), 2-10.

Rasmussen, A., Jirenhed, D. A., & Hesslow, G. (2008). Simple and complex spike firing patterns in
Purkinje cells during classical conditioning. *The Cerebellum*, *7*(4), 563.

Sakamoto, T., & Kondo, T. (2012, November). Can passive arm movement affect adaptation to
visuomotor rotation?. In *2012 IEEE International Conference on Development and Learning and*
Epigenetic Robotics (ICDL) (pp. 1-6). IEEE.

Shadmehr, R., Smith, M. A., & Krakauer, J. W. (2010). Error correction, sensory prediction, and
adaptation in motor control. *Annual review of neuroscience*, *33*, 89-108.

Scheidt, R. A., Conditt, M. A., Secco, E. L., & Mussa-Ivaldi, F. A. (2005). Interaction of visual and
proprioceptive feedback during adaptation of human reaching movements. *Journal of*
neurophysiology, *93*(6), 3200-3213.

Sing, G. C., & Smith, M. A. (2010). Reduction in learning rates associated with anterograde interference
results from interactions between different timescales in motor adaptation. *PLoS Comput Biol*, *6*(8),
e1000893.

Smith MA, Ghazizadeh A, Shadmehr R. Interacting adaptive processes with different timescales underlie
short-term motor learning. *PLoS Biol* 4: e179, 2006.

Steinmetz, J. E., Sears, L. L., Gabriel, M., Kubota, Y., & Poremba, A. (1991). Cerebellar interpositus
nucleus lesions disrupt classical nictitating membrane conditioning but not discriminative avoidance
learning in rabbits. *Behavioural brain research*, *45*(1), 71-80.

Taylor, J. A., Krakauer, J. W., & Ivry, R. B., (2014). Explicit and implicit contributions to learning in a
sensorimotor adaptation task. Journal of Neuroscience, 34(8), 3023-3032.

Tays, G., Bao, S., Javidialsaadi, M., & Wang, J. (2020). Consolidation of use-dependent motor memories
induced by passive movement training. *Neuroscience Letters*, *732*, 135080.

Tong, C., & Flanagan, J. R. (2003). Task-specific internal models for kinematic transformations. *Journal of*
Neurophysiology, *90*(2), 578-585.

Tseng, Y. W., Diedrichsen, J., Krakauer, J. W., Shadmehr, R., & Bastian, A. J. (2007). Sensory prediction
errors drive cerebellum-dependent adaptation of reaching. *Journal of neurophysiology*, *98*(1), 54-62.

Vergaro, E., Casadio, M., Squeri, V., Giannoni, P., Morasso, P., & Sanguineti, V. (2010). Self-adaptive
robot training of stroke survivors for continuous tracking movements. *Journal of neuroengineering and*
rehabilitation, *7*(1), 1-12.

Verstynen, T., & Sabes, P. N. (2011). How each movement changes the next: an experimental and
theoretical study of fast adaptive priors in reaching. *Journal of Neuroscience*, *31*(27), 10050-10059.

Wang, J., & Lei, Y. (2015). Direct-effects and after-effects of visuomotor adaptation with one arm on
subsequent performance with the other arm. *Journal of neurophysiology*, *114*(1), 468-473.

Wigmore, V., Tong, C., & Flanagan, J. R. (2002). Visuomotor rotations of varying size and direction
compete for a single internal model in a motor working memory. *Journal of Experimental Psychology:*
Human Perception and Performance, *28*(2), 447.

Yin, C., & Wei, K. (2020). Savings in sensorimotor adaptation without an explicit strategy. *Journal of*
neurophysiology, *123*(3), 1180-1192.

Zarahn, E., Weston, G. D., Liang, J., Mazzoni, P., & Krakauer, J. W. (2008). Explaining savings for
visuomotor adaptation: linear time-invariant state-space models are not sufficient. *Journal of*
neurophysiology, *100*(5), 2537-2548.

742