A conversion from slow to fast memory in response to passive motion

Mousa Javidialsaadi[1i], Scott T. Albert[2i], Jinsung Wang[1*]

[1]Department of Kinesiology, University of Wisconsin – Milwaukee, Milwaukee, WI, 53201

[2]Depertment of Biomedical Engineering, Johns Hopkins University – Baltimore, MD 21205

*Correspondence: Jinsung Wang, 492 Enderis Hall, University of Wisconsin – Milwaukee, WI

53201, USA.

Email: wang34@uwm.edu.

Acknowledgements: This work was supported by a grant from the National Institutes of Health

(F32NS095706). We dedicate this article to Dr. Fatemeh Pasand and Dr. Majid Chahardah Cheric

for all their academic efforts and teachings. They passed away on 10 April 2018 and 26 June 2020,

respectively.

Data availability statement: The data that support the findings of this study are available from

the corresponding author upon reasonable request.

17
18

---

[i] The first and second authors contributed equally to this work.

Abstract

When the same perturbation is experienced consecutively, learning is accelerated on the second

attempt. This savings is a central property of sensorimotor adaptation. Current models suggest

that these improvements in learning are due to changes in the brain's sensitivity to error. Here,

we tested whether these increases in error sensitivity could be facilitated by passive movement

experiences. In each experimental group, a robot moved the arm passively in the direction that

solved the upcoming rotation, with no visual feedback provided. Following that, participants

adapted to a visuomotor rotation. Prior passive movements substantially improved motor

learning, increasing total compensation in each group by approximately 30%. Similar to savings,

a state-space model suggested that this improvement in learning was due to an increase in error

sensitivity, but not memory retention. When we considered the possibility that learning was

supported by parallel fast and slow adaptive processes, a striking pattern emerged: whereas

initial improvements in learning were driven by a slower adaptive state, increases in error

sensitivity gradually transferred to a faster learning system with the passage of time. These

findings suggest that passive errors engage motor learning systems, but the resulting behavioral

patterns migrate between slow and fast adaptive circuits as the passive memory is consolidated.

Introduction

When we make a reaching movement, our brain predicts how our motor commands will change our sensory state. Perturbations to our movements create an error between the predicted sensory state and the observed sensory state (Popa & Ebner, 2019; Shadmehr et al., 2010). These sensory prediction errors cause learning, or changes to our motor commands on the next movement (Krakauer et al., 2000; Mazzoni & Krakauer, 2006). One hallmark of sensorimotor learning is savings: when people are exposed to the same perturbation twice in a row, they learn more rapidly during the second exposure (Smith et al., 2006; Kitago et al., 2013). Some models suggest that savings occurs because the brain recalls the movements it made in the past (Haith et al., 2015; Morehead et al., 2015). Other models suggest that savings occurs because the brain becomes more sensitive to errors it has experienced in the past (Gonzalez-Castro & Hadjiosif et al., 2014; Herzfeld et al., 2014; Albert et al., 2020). A common assumption across both models, however, is that savings is driven by active movement experiences.

But changes in learning rate can also occur even when people have not actively moved in the past. For example, recent studies have shown that passive sensorimotor experiences can improve active motor learning (Lei et al., 2016; Bao et al., 2017; Lei et al., 2017; Tays et al., 2020). Robot-assisted experiences enhance our ability to acquire novel motor skills (Reinkensmeyer and Patton, 2009; Bara and Gentaz, 2011; Basteris et al., 2012; Beets et al., 2012), facilitate motor recovery (Sakamoto et al., 2012; Tays et al., 2020), and improve motor function in hemiparetic stroke patients (Aisen et al., 1997; Krebs et al., 1998; Riener et al., 2005; Kahn et al., 2006; Vergaro et al., 2010).

For active movements, current state-space models (Smith et al., 2006; McDougle et al.,

2015; Albert et al., 2018; Coltman et al., 2019; Orozco et al., 2020) posit that adaption is due to

an interplay between at least two parallel adaptive processes: a slowly-adapting system (slow

process) which responds weakly to error, but strongly retains its memory, and a rapidly-adapting

system (fast process) which responds strongly to error, but weakly retains its memory. An

increase in either system's sensitivity to error is widely associated with savings in sensorimotor

learning paradigms (Gonzalez-Castro et al., 2014; Herzfeld et al., 2014; Leow et al., 2016; Albert

et al., 2020). Does passive training, then, enhance active motor learning by increasing error

sensitivity in slow and/or fast learning processes?

Here we investigated this question in a visuomotor rotation experiment. In this paradigm,

participants move a cursor towards a target, but the cursor's position is rotated relative to the

hand's path. We used a robot to passively create a memory that could help solve a visuomotor

rotation. Following this passive training, participants actively moved a rotated cursor to various

visual targets. We varied the time between passive and active training to allow the passive

memory to consolidate, and investigated how this consolidation altered the error sensitivities in

the slow and fast adaptive processes (Shadmehr & Brashers-Krug, 1997; Caithness et al., 2004;

Krakauer et al., 2005; Lerner & Albert et al., 2020).

Materials & Methods

Participants

Twenty-eight healthy volunteers (17 males, 11 females; aged: 18-35) participated in this study.

All subjects were right-handed as assessed by the Edinburgh handedness inventory (Oldfield,

1971). Each subject signed a consent form that was approved by the University of Wisconsin-

Milwaukee Institutional Review Board. Participants were randomly assigned to one of three

experimental groups or a control group.

Apparatus

Participants were seated in a robotic exoskeleton (KIMARM, BKIN Technologies Ltd, Kingston,

ON, Canada) that provided gravitational support to both arms. The exoskeleton was positioned

so that the participant's arm was hidden underneath a horizontal display (Fig. 1A). To track the

hand's position, a small cursor was projected onto the display, over the participant's index

fingertip. During each trial, the KINARM projected visual stimuli (a start position and a target)

onto the display, so that they appeared to be within the same plane as the arm. The visual stimuli

consisted of a centrally located start circle (2 cm in diameter) and one of four target circles (2cm

in diameter) located 10 cm away from the start target (Fig. 1B). We sampled the hand's position

in the x-y plane at 1000 Hz. Position data were low pass filtered at 15 Hz, and then differentiated

to obtain velocity. Post-processing, analysis, and modeling were conducted in MATLAB R2018a

(The MathWorks Inc., Natick, MA).

Figure 1. Passive movement experiences facilitate active motor adaptation. **A.** Participants were

seated in a KINARM exoskeleton. The arm was placed underneath a horizontal display. **B.** The

experiment began with a baseline phase, where the cursor moved with the hand. During the

passive movement period, participants were shown a target in the 45°, 135°, 225° or 315°

direction, but the KINARM passively moved the hand 30° CW relative to the visual target. No

visual feedback was provided. Then participants took a break (time passage). When participants

returned to the task, they started an active rotation period where the cursor was rotated 30° to

the hand. Thus, to "solve" the rotation, participants needed to reach along the same path

traversed during the passive period. **C.** Complete experiment paradigm. Experimental groups

were divided into a 5 min, 1 hr, or 24 hr time passage condition. **D.** Average reach trajectories over the 20 rotation cycles. The control group (no passive training) is shown in gray. Reach angles for 5 min group (left), 1 hr group (middle), and 24 hr group (right) are shown in black. We calculated the reach angle between the hand and target cursor, at the movement's midpoint (5 cm displacement). **E.** Cycle-by-cycle hand angles at reach midpoint. **F.** We calculated the mean reach angle across the rotation period. Statistics show post-hoc Tukey's test following one-way ANOVA (* indicates $p < 0.05$). Error bars show mean ± SEM.

Experimental Design

Participants were assigned to one of three experimental groups (n=7/group). Each experimental group experienced four separate periods: (1) a baseline reaching period, (2) a passive movement period, (3) a break in time, and (4) a visuomotor rotation period (Fig. 1C). To determine how passive movements altered reaching behavior, we compared behavior in the experimental groups to that of a control group (n=7) that did not experience the passive movement period prior to rotation learning.

In the baseline period (Fig. 1B, baseline), participants moved their arm to each target over a 10-cycle reaching period. On each trial, continuous visual feedback was provided via a cursor over the index fingertip. Participants were instructed to move rapidly and accurately to the target location. The trial ended 1.5 sec after the target was presented. To begin the next trial, the participant had to move their hand back to the central start position.

In the passive movement period, the KINARM moved the participant's right arm along a straight 10 cm minimum jerk trajectory. On each trial, a visual target was displayed, but the robot

passively moved the arm along a path that was rotated 30° clockwise (CW) to the visual target

(Fig. 1B, passive period). Critically, this CW rotation served as the "solution" to a counterclockwise

(CCW) visuomotor rotation that participants had not yet experienced. Participants were told to

relax their arm and not resist the passive motion. The passive period consisted of 30 cycles (4

trials in a cycle, 1 to each target in a pseudorandom order). No visual feedback was provided on

these trials. Thus, participants received only proprioceptive information about the error between

their arm's path and the visual target. The passive movement period terminated with a time-

delay which distinguished each experimental group: 5-min, 1-hr, or 24-hr delay. Once the delay

concluded, the rotation period began.

The experiment ended with an active visuomotor rotation period that lasted 20 cycles (80

trials total). On rotation trials, the cursor was rotated 30° CCW relative to the hand path (Fig. 1B,

rotation). As noted above, the "solution" to this rotation would be to counter-rotate the reach

path by 30° CW relative to the target. Critically, this CW rotation would coincide with the

movement path experienced during the passive period. Thus, participants in each experimental

had been given the proprioceptive experience of this "solution", though this was never explicitly

revealed to each participant. Participants in the control group, however, had never experienced

the passive movement period. Thus, there was no prior memory to draw upon during the rotation

period.

Empirical Data Analysis

Data analysis was performed using MATLAB (Mathworks, Natick, MA). The pointing angle at the

reaching movement's midpoint (5 cm displacement) was used as our performance measure,

calculated as the hand's angular position relative to the line segment connecting the start and

target positions. Data were then averaged within each 4-trial cycle.

Here we considered how passive training altered adaptation during the rotation period.

Fig. 1D illustrates the average reach trajectory over the 20-cycle rotation period. To calculate

these mean hand paths, we artificially rotated each reach trajectory to create a 90° angle

between the start location and the target (i.e., movement directed along the y-axis alone). Next,

we interpolated the hand's lateral deviation (the x-axis in Fig. 1D) at 25 points evenly spaced

between the start location and an 8 cm displacement towards the target (the y-axis in Fig. 1D).

Fig. 1E shows hand angles on the last two baseline cycles, as well as the learning curve during the

20-cycle rotation period. To assess overall adaptation, we averaged each participant's midpoint

hand angle over the 20-cycle rotation period (Figs. 1F).

To aid visual inspection, we fit an exponential model to the mean learning curve in each

group (Fig. 1E, dashed line, exponential fit). This exponential curve tracked how reach angle (y)

changed over each cycle (t, starting at 0):

$$y^{(t)} = \alpha e^{-\beta t} + c \qquad (1)$$

Here α and c determine the initial reach angle and asymptotic reach angle. The parameter β

represents the participant's learning rate during the adaptation period. We fit this exponential

function to each group's mean data in the least-squares sense using *fmincon* in MATLAB R2018a.

We repeated the fitting procedure 20 times, each time varying the initial parameter guess that

seeded the algorithm. We selected the model parameters which minimized squared error over

all 20 repetitions.

We observed that participants who completed passive training exhibited improved

adaptation (Fig. 1F). These improvements could have been due to 3 non-exclusive causes. First,

the differences in adapted reach angles may have been due to biases in reach angles that lingered

from the baseline reach period (Hypothesis 1). Second, improvements in adaptation may have

been due to use-dependent biases in reach angles induced by passive training (Hypothesis 2).

Third, improved adaptation may have been due to an enhancement in trial-to-trial learning

induced by passive training (Hypothesis 3). To evaluate each hypothesis, we analyzed separate

reach periods during the experiment protocol.

For Hypothesis 1 (lingering baseline biases), we calculated the reach angles at the very

end of the baseline period in each participant. We calculated these angles both on the last 2

baseline trials (Figs. 2A-C) and the last 4 baseline trials. For Hypothesis 2 (use-dependent bias at

the start of the rotation period), we calculated the reach angles at the very start of the rotation

period in each participant (Figs. 2D-F). We calculated these angles both on the initial 2 rotation

trials and the initial 4 rotation trials.

Figure 2. Improvements in adaptation were unrelated to biases in reach angle. **A.** We measured

reach angles at the very end of the baseline period (end of baseline, last 2 baseline trials). **B.**

Average trajectory on the end of baseline trials. **C.** Mean reach angle over the end of baseline

trials. **D.** We measured reach angles at the start of the rotation period (initial adaptation, first 2

rotation trials). **E.** Average trajectory on the initial adaptation trials. **F.** Mean reach angle over the

initial adaptation trials. Statistics indicate the group main effect in a one-way ANOVA (n.s.

denotes p>0.05). Error bars show mean ± SEM.

For Hypothesis 3 (passive training improves learning), we noted that adaptation is

supported by slow-learning processes and fast-learning processes (Smith et al., 2006). Fast

processes are dominant early during adaptation. Slow processes are dominant late in adaptation.

Thus, we measured adaptation both early and late during the learning process. To determine the

point at which fast adaptation switched to slow adaptation, we calculated a cycle-by-cycle

learning rate measure, i.e., the adaptation curve's slope on each cycle. To do this, we used a

second-order finite difference (Fig. 3A) to approximate the derivative of the learning curves in

Fig. 1E on each cycle. We observed that adaptation exhibited rapid learning on adaptation cycles

1 and 2, intermediate learning on cycle 3, and slow learning on cycle 4 and beyond. Thus, we

divided the adaptation period into a rapid early learning period (Figs. 3B&C; cycles 1 and 2), as

well as a slow late learning period (Figs. 3D&E; rotation cycles 3-20). We calculated the mean

reach angle over these two periods in Figs. 3C&E. In addition, we calculated a normalized early

learning metric by subtracting off the baseline reaching angle exhibited on the last baseline cycle.

Figure 3. Passive training enhances early and late motor adaptation. **A.** We calculated an

instantaneous adaptation rate measure on each cycle (using a second-order finite difference,

Bozorgpour., 2023; Bozorgpour et al., 2023). Here data are averaged across all experimental

groups. The adaptation rate was large during an initial 2-cycle period (rapid early learning). The

adaptation rate was small late during adaptation (slow late learning). **B.** Mean reach trajectory

over the rapid early learning period. Control group (no passive training) is shown in gray.

Experimental groups (5 min, 1 hr, and 24 hr) are shown in black. **C.** Mean reach angle over the

rapid early learning period. **D.** Mean reach trajectory over the slow late learning period. **E.** Mean

reach angle over the late slow learning period. Statistics: in each panel, statistics show post-hoc

Dunnett's test against the control group following one-way ANOVA (*p<0.05). Error bars show

mean ± SEM.

State-space learning model: one-state

To better understand the adaptation process, we used a state-space model (Smith et al., 2006).

The state-space model posits that learning consists of cycle-to-cycle error-based learning as well

as cycle-by-cycle memory retention (i.e., forgetting). The forgetting process is controlled by a

retention factor (a), which specifies how much adaptation is retained from one cycle to the next.

The learning process is controlled by the participant's error sensitivity (b), which specifies how

much is learned from a given error (e). These processes together determine how the participant's

internal state (x) changes over time, in the presence of internal state noise (ε_x, normal with zero

mean, SD=σ_x):

$$x^{(t+1)} = ax^{(t)} + be^{(t)} + \varepsilon_x^{(t)} \tag{2}$$

Eq. (2) allows us to ascribe any differences in performance during the adaptation period to

meaningful quantities: retention (a) and error sensitivity (b).

Note, however, that the internal state (x) is not a measurable quantity. Rather, on each

cycle, the motor output (reach angle) is measured. This reach angle (y) directly reflects the

subject's internal state, but is corrupted by execution noise (ε_x, normal with zero mean, SD=σ_y)

according to:

$$y^{(t)} = x^{(t)} + \varepsilon_y^{(t)} \qquad\qquad (3)$$

Thus, Eqs. (2) and (3) represent a single module state-space model. We fit this model to each

participant's reach angles during the adaptation period using the Expectation-Maximization (EM)

algorithm (Albert & Shadmehr, 2018). EM is an algorithm that conducts maximum likelihood

estimation in an iterative process. We used EM to identify the model parameters {a, b, x_1, σ_x, σ_y,

σ_1} that maximized the likelihood of observing the data (note that x_1 and σ_1 represent the

participant's initial state and variance, respectively). We conducted 10 iterations, each time

changing the initial parameter guess that seeded the algorithm. We selected the parameter set

that maximized the likelihood function across all 10 iterations. We compared the retention

factors and error sensitivities predicted by the model in Figs. 4B&C.

Figure 4. Passive training improves motor learning by increasing sensitivity to error. **A.** Here we

show reach angles during the active rotation learning cycles. Control group (no passive training)

is shown in gray. Experimental groups (5 min, 1 hr, and 24 hr) are shown in black. We fit the data

with a single-module state-space model (dashed black line; mean across individual participants).

The state-space model posited that adaptation was due to both error-based learning (controlled

by error sensitivity) and trial-by-trial forgetting (controlled by retention factor). **B.** Here we show

retention factor predicted by the single-module state-space model. **C.** Here we show error

sensitivity predicted by the single-module state-space model. **D.** We fit a two-state model to

individual participant behavior. Predicted slow state for experimental groups (5 min, 1 hr, and 24

hr) are shown in blue; time-delay between passive and active training increases left-to-right.

Predicted slow state in control group is shown in gray. **E.** Here we provide slow state error

sensitivity predicted by the two-state model. **F.** Same as in **D**, but for the fast state of learning. **G.**

Same as in **E** but for the fast state error sensitivity. In each panel, statistics show post-hoc Tukey's

test (**C**) or Dunn's test (**G**) following one-way ANOVA (**C**) or Kruskal-Wallis (**E** and **G**). Statistics: *

indicates p<0.05, ** indicates p<0.01). Error bars show mean ± SEM.

State-space learning model: two-state

The single-module state-space model describes learning as a single adaptive process. Motor

adaptation, however, is believed to be comprised of multiple states, each with different

timescales of learning. These states appear to be accurately summarized as a parallel two-state

system, with adaptation supported by a slow adaptive process and a fast adaptive process (Smith

et al., 2006; McDougle et al., 2015; Albert & Shadmehr, 2018; Coltman et al., 2019). The slow

process exhibits slower error-based learning, but strong retention over time. The fast process

exhibits faster error-based learning, but higher rates of forgetting. We wondered how these

adaptive states are altered by consolidation of passive motor memory.

To answer this question, we fit a standard two-state model of learning to individual

participant behavior. In this model, slow state and fast state adaptation are controlled by slow

state and fast state retention factors (a_s and a_f, respectively), as well as slow state and fast state

error sensitivities (b_s and b_f, respectively). As in Eq. (1), both internal states exhibit cycle-by-cycle

learning and forgetting, in the presence of internal state noise ($\varepsilon_{x,s}$ and $\varepsilon_{x,f}$, normal with zero

mean, SD=σ_x):

$$\begin{aligned} x_s^{(t+1)} &= a_s x^{(t)} + b_s e^{(t)} + \varepsilon_{x,s}^{(t)} \\ x_f^{(t+1)} &= a_f x^{(t)} + b_f e^{(t)} + \varepsilon_{x,f}^{(t)} \end{aligned} \tag{4}$$

To enforce the traditional two-state dynamics (slow state has higher retention, fast state has higher error sensitivity), retention factors and error sensitivities in Eq. (4) are constrained such that $a_s > a_f$ and $b_f > b_s$. As with the single-module state-space model, the two adaptive are not directly measurable. But together, they sum to produce the overall adapted reach angle, which is also corrupted by execution noise:

$$y^{(t)} = x_s^{(t)} + x_f^{(t)} + \varepsilon_y^{(t)} \tag{5}$$

Together, Eqs. (4) and (5), along with the inequality constraints relating a_s, a_f, b_s, and b_f, constitute the two-state model of learning. The full parameter set consists of: $\{a_s, a_f, b_s, b_f, x_{s,1}, x_{f,1}, \sigma_x, \sigma_y, \sigma_1\}$. Note that $x_{s,1}$, and $x_{f,1}$ are the initial slow and fast state magnitudes, which are also estimated by the model.

We fit this model to individual participant data, using the EM algorithm (Albert & Shadmehr, 2018). Note that one's ability to fit the two-state model greatly benefits from multiple trial conditions, such as set breaks, error-clamp periods, washout periods, and perturbation reversals (Albert & Shadmehr, 2018). However, the current experiment consisted of only perturbation trials with one orientation. Therefore, to improve our ability to robustly recover the two-state model parameters, we used a two-tiered fitting procedure (described below).

In our single-state model, we found that error sensitivity varied across experimental and control groups, but not retention. Therefore, to improve our ability to recover slow and fast error sensitivity (Albert & Shadmehr, 2018), we started by constraining the model's retention factors (hypothesizing that these were not impacted by the passive movement period). To do this, we

started by fitting the model to mean behavior in the 5 min, 1 hr, 24 hr, and control groups. Then

we calculated the midpoint in the range spanned by slow and fast retention factors identified in

each group. Lastly, we then fit the two-state model to individual participant behavior in each

group, after fixing the slow and fast retention factors to these values.

When fitting the two-state model to either group behavior or individual participant

behavior, we performed 10 iterations of the EM algorithm, each time varying the initial

parameter guess that seeds the algorithm. We then selected the parameter set computed by EM

with the greatest likelihood. We used these parameter sets to simulate the model's slow and fast

learning states in Figs. 4D&F. We compared the error sensitivities predicted by the model in Figs.

4E&G.

Statistics

To compare the control and experimental groups, we conducted one-way ANOVAs in MATLAB

R2018a. For post-hoc testing, we used Dunnett's test to assess whether the behavior in the 5-

min, 1-hr, and 24-hr delay groups differed from the control group. However, we used a non-

parametric test to analyze our two-state model parameters, which are prone to outlying

parameter estimates (Albert & Shadmehr, 2018; Coltman & Gribble, 2019). Thus, we used a

Kruskal-Wallis test to evaluate differences in median error sensitivity (one outlier observed in the

1-hr slow state error sensitivity). For post-hoc testing following Kruskal-Wallis, we used Dunn's

test in IBM SPSS 25 (5-min, 1-hr, and 24-hr groups tested against control).

Results

1 *Passive proprioceptive training facilitates active reach adaptation*

2 The experiment began with a baseline period where participants reached to targets with veridical

3 hand position feedback (Fig. 1B, baseline). Next, a passive training period ensued, where the

4 robot moved the participant's arm, without any visual feedback (Fig. 1B, passive period). On

5 these trials, the robot moved the hand along a path rotated 30° CW relative to the target. Thus,

6 each passive movement terminated in a proprioceptive "error" between the visual target and

7 the hand. Following a break in time (Fig. 1B, time passage; 5 min, 1 hr, or 24 hr) participants

8 returned to the task, but now actively moved to each target in the presence of a visuomotor

9 rotation (Fig. 1B, rotation). How was adaptation altered by passive training?

10 To answer this question, we compared reaching movements during the rotation period

11 (Fig. 1D, black) to those of a control group that never experienced passive training (Fig. 1D, gray).

12 We calculated the angle of the hand's path midway through the movement (Fig. 1D, midpoint).

13 Hand angles rapidly adapted over the 20 rotation cycles (Fig. 1E) to reduce errors between the

14 cursor and target. Remarkably, we observed that passive training greatly improved motor

15 adaptation, increasing average compensation during the rotation period by 30% in all 3

16 experimental groups (Fig. 1F; one-way ANOVA, $F_{(3,24)}=3.98$, $p=0.02$; post-hoc Dunnett's test

17 against control, 5 min: $p=0.021$, 1 hr: $p=0.021$, 24 hr: $p=0.038$).

18 It is possible that this improved adaptation was partially due to biases in reach angles that

19 had developed during the baseline period. To assess whether baseline biases skewed our findings

20 during the adaptation period (Hypothesis 1), we measured participant reach angles at the very

21 end of the baseline period (Fig. 2A, end of baseline). At the end of baseline training, active

22 reaching movements exhibited small biases ranging between about -3° and 0°. Critically,

however, we did not detect any statistically significant differences in these initial biases between

the experimental and control groups on the last two baseline trials (Fig. 2C; one-way ANOVA,

F(3,24)=0.97, p=0.42), nor on the last four baseline trials (one-way ANOVA, F(3,24)=0.82,

p=0.496). Thus, improvements in adaptation (Fig. 1F) were not due to any biases in reach angles

that were carried over from the baseline period.

Instead, it was passive training that improved active motor adaptation. These

improvements could have had two non-exclusive causes. First, passive training may have created

a use-dependent bias in the initial reach angles during the adaptation period (Hypothesis 2).

Second, passive training may have primed the motor learning system, increasing the amount

participants adapted to visual errors (Hypothesis 3). Had passive training caused a use-dependent

bias in reach angle (Hypothesis 2), this bias might have been present on the very initial rotation

cycles (Fig. 2D, initial adaptation). However, when we assessed the initial reach angles during the

rotation period (Figs. 2D&E), we did not detect any statistically significant differences across the

experimental and control groups on the first two rotation trials (Fig. 2F; one-way ANOVA,

F(3,24)=0.48, p=0.7), nor on the first four rotation trials (one-way ANVOA, F(3,24)=1.38, p=0.274).

Thus, improvements in adaptation were not observed in the initial reach angles during

the rotation period; rather, they emerged as participants actively adapted to the rotation

(Hypothesis 3). At what point did these changes emerge during rotation training? To investigate

this, we first noted that adaptation is supported by multiple processes, some fast and others slow

(Smith et al., 2006; Albert & Shadmehr, 2018), with fast processes dominant early during

adaptation, and slow processes dominant late in adaptation. Indeed, trial-to-trial learning was

very rapid over an initial 3-cycle period (Fig. 3A, gray region, rapid early learning), and then

decreased abruptly over the remaining training epochs (Fig. 3A, slow late learning). Did passive

training improve adaptation during the rapid initial learning phase, or during the late slow

learning phase?

To answer this question, we compared each experimental group's mean adaptation level

over the early learning period with the control group (Fig. 3B). Interestingly, passive training

strongly improved adaptation over the rapid early learning period (Fig. 3C; one-way ANOVA,

$F(3,24)=6.53$, $p=0.002$), but the facilitation appeared stronger when more time had passed

between passive and active training (post-hoc Dunnett's test against control: 5 min: $p=0.061$, 1

hr: $p=0.003$, 24 hr, $p=0.001$). The same trend was observed when we normalized initial

adaptation levels to each group's baseline reach angles (one-way ANOVA, $F(3,24)=3.2$, $p=0.041$;

post-hoc Dunnett's test against control: 5 min: $p=0.277$, 1 hr: $p=0.048$, 24 hr, $p=0.026$). In sum,

early learning was enhanced in the 1 hr and 24 hr groups.

Lastly, we measured each group's mean adaptation over the late slowly adapting period

(Fig. 3D). Here, we observed that less time passage between passive and active training produced

stronger improvements in learning over the control group (Fig. 3E; one-way ANOVA,

$F(3,24)=3.54$, $p=0.03$; post-hoc Dunnett's test against control: 5 min: $p=0.021$, 1 hr: $p=0.037$, 24

hr: $p=0.075$).

In summary, we observed that passive training enhanced adaptation. These

improvements were not due to a use-dependent bias in initial rotation reach angles. Rather,

passive training appeared to improve active motor learning. But adaptation patterns varied with

time passage between passive and active training. With short delays, improvements emerged

later during rotation training. With long delays, improvements emerged earlier during rotation

training.

Passive movement experiences enhance sensitivity to error, but not retention

To better understand how passive training improved motor learning, we used a state-space

model. In a state-space model, adaptation is an interplay between two processes: error-based

learning and trial-by-trial forgetting. Error-based learning is controlled by the participant's

sensitivity to error (b in Eq. 2). Forgetting is controlled by the participant's retention factor (a in

Eq. 2). How did passive training alter each participant's error sensitivity and retention?

To answer this question, we fit a single-module state-space model to individual

participant reach angles in the experimental and control groups. The model (Fig. 4A, dashed lines)

appeared to closely track each group's learning curve. Next, we isolated the retention factor (Fig.

4B) and error sensitivity (Fig. 4C) predicted by the state-space model. Interestingly, passive

training did not appear to alter the retention processes in any experimental group (Fig. 4B, one-

way ANOVA, $F(3,24)=2.19$, $p=0.115$). On the other hand, passive training clearly impacted error

sensitivity (Fig. 4C, one-way ANOVA, $F(3,24)=3.81$, $p=0.023$), which appeared to grow with time-

delay duration; though only the 24 hr group showed a statistically significant difference relative

to the control condition (post-hoc Dunnett's tests against control, 5 min: $p=0.72$, 1 hr: $p=0.061$,

24 hr: $p=0.017$).

In sum, the state-space model predicted that passive training increased how sensitive

participants were to errors experienced during the rotation period. But a puzzle remained: why

is error sensitivity only increased in the 24 hr group (Fig. 4C), when all experimental groups

showed a facilitation in learning (Fig. 1F)? We speculated that our state-space model was limited

because it described adaptation as a single learning process. However, motor adaptation appears

to be supported by at least two parallel adaptive processes: a slow state which has low error

sensitivity but strong retention, and a fast state that has high error sensitivity but weak retention

(Smith et al., 2006).

To better understand the error sensitivity patterns in Fig. 4C, we next applied a two-state

learning model (Eqs. (4) & (5)) to the data. The two-state model predicted that adaptation was

supported by the slow-adapting processes shown in Fig. 4D and the rapid-adapting processes

shown in Fig. 4F. These states showed a striking trend: the slow-adapting process was more active

with short delays between passive and active training (i.e., the 5 min group). The fast state of

learning exhibited the opposite trend; it was more active with long delays between passive and

active training (i.e., the 24 hr group).

These changes in the slow and fast states were due to a clear pattern in error sensitivity.

The slow state's error sensitivity increased nearly two-fold following the passive training period

in the 5 min group (Fig. 4E, Kruskal-Wallis test, $X^2(27)=8.32$, p=0.0398) but gradually returned to

control levels as time elapsed between passive and active training (post-hoc Dunn's tests against

control, 5 min: p=0.024, 1 hr: p=1, 24 hr: p=1). On the other hand, the fast-state's error sensitivity

increased by 50% (Fig. 4G, Kruskal-Wallis test, $X^2(27)=9.04$, p=0.0288) but only in the 24 hr group

after the passage of time (post-hoc Dunn's tests against control, 5 min: p=0.486, 1 hr: p=0.087,

24 hr: p=0.012). …

In sum, the two-state model indicated that passive training increased error sensitivity.

Increases in error sensitivity improved visuomotor adaptation. With little time passage between

passive and active training, error sensitivity was mostly improved in a slower adaptive state (Figs. 4D&E). With greater time passage, error sensitivity was mostly improved in a faster adaptive state (Figs. 4F&G).

Discussion

When the same perturbation is experienced consecutively, learning is accelerated on the second attempt. This savings is a central property of sensorimotor adaptation that is observed across several motor effectors: the reaching system (Smith et al., 2006; Zarahn et al., 2008; Leow et al., 2013; Haith et al., 2015; Coltman et al., 2020; Yin & Wei, 2020), walking (Mawase et al., 2014; Day et al., 2018), and even the oculomotor system (Kojima et al., 2004). Current models have suggested that these improvements in learning are due to changes in the brain's sensitivity to error (Gonzalez-Castro & Hadjiosif, 2014; Herzfeld et al., 2014; Mawase et al., 2014; Coltman et al., 2020). Here, we tested whether these increases in error sensitivity could be facilitated by passive experiences, as opposed to active movements.

Consistent with earlier work (Lei et al., 2016; Bao et al., 2017; Lei et al., 2017; Tays et al., 2020), we observed that passive training facilitated active motor adaptation. In each experimental group, a robot moved the arm passively in the direction that provided a solution to the upcoming rotation perturbation, with no visual feedback provided. This passive proprioceptive experience substantially altered subsequent motor learning, increasing total compensation in each group by approximately 30%. Similar to savings, the state-space model suggested that this improvement in learning was due to an increase in error sensitivity. Thus, passive memories appeared to increase the motor learning system's sensitivity to error.

However, the state-space model's prediction exhibited a puzzle: whereas all experimental

groups showed improvements in total compensation, only the 24-hr group showed a statistically

significant increase in sensitivity to error (Fig. 4C). This suggested that this model was missing a

key component: multiple adaptive states. That is, converging evidence suggests that

sensorimotor learning is supported by parallel processes (Smith et al., 2006; Mawase et al., 2014;

McDougle et al., 2015; Coltman et al., 2020). The parallel adaptive states are well-approximated

as a two-state system: one slow state that learns slowly but is decay-resistant, and one fast state

that learns rapidly but also forgets rapidly. When we applied this two-state model to our data

(Albert & Shadmehr, 2018), a new pattern emerged: passive training initially facilitated

improvements in the slow state's sensitivity to error (Fig. 4E). As time passed between passive

and active training, these changes in the slow state's error sensitivity dissipated; simultaneously,

the fast process increased its error sensitivity (Fig. 4G).

How might passive training increase the brain's sensitivity to error during active

movements? Both passive movements (Carel et al., 2000) and active motor learning (Della-

Maggiore & McIntosh, 2005; Herzfeld et al., 2015; Popa et al., 2012; Izawa et al., 2012; Tseng et

al., 2007; Gibo et al., 2013; Maschke et al., 2004) are known to involve both the primary motor

cortex (M1) and the cerebellum. One hypothesis is that prediction errors experienced during

passive training engage Purkinje cells (P-cells) in the cerebellar cortex as well neurons in the deep

cerebellar nuclei (Herzfeld et al., 2020; Lisberger et al., 1994; McCormick & Thompson, 1984;

Perret et al., 1993). An interplay between P-cells and nuclear neurons may produce the changes

in slow-state error sensitivity we observed early (5 min) after passive training. However, with

time passage, plasticity in the cerebellum may be transferred to, and consolidated within M1 and

S1 neurons via an unidentified mechanism (Kumar et al., 2019; Herzfeld et al., 2020; Orozco et

al., 2020). These changes in the sensorimotor cortices may produce the increases in fast-state

error sensitivity we observed with longer delays between passive and active training (24-hr

group).

This first hypothesis draws upon the idea that passive training creates sensory prediction

errors (SPEs). Current models have proposed that these SPEs engage the inferior olive, which in

turn generate complex spikes in cerebellar P-cells (Herzfeld et al., 2015; Herzfeld et al., 2018).

Associative learning paradigms such as eye-blink conditioning demonstrate that these SPEs and

complex spikes are evoked even without active movement (Ohmae & Medina, 2015; Kim et al.,

2020; Ito, 201; Sears & Steinmetz, 1991; Rasmussen et al, 2008). Thus, in our passive training

paradigm, it is possible that the discrepancy between the visual target and the arm's

proprioceptive state (position) may have created an SPE and corresponding complex spikes in the

cerebellar cortex (Popa & Ebner, 2019; Shadmehr et al., 2010). Repeated exposure to consistent

prediction errors may have upregulated error sensitivity in the cerebellar cortex (Herzfeld et al.,

2014; Albert et al., 2020), thus facilitating learning in the active condition. An alternate possibility

is that passive training engaged the primary motor and somatosensory cortices, which then

indirectly altered sensitivity to SPEs in the cerebellum via descending input. In either case, rTMS

and tDCS over the cerebellum and M1 may provide a path to elucidating the various timescales

of memory that are engaged by passive training.

A second hypothesis is that passive movements altered error sensitivity through use-

dependent learning (Wang et al., 2015; Bao et al., 2017; Lei et al., 2017; Verstynen & Sabes, 2011;

Lei et al., 2016; Diedrichsen et al., 2010; Jax & Rosenbaum, 2007; Scheidt et al., 2005;

Diedrichsen, 2007), as a process by which motor plans are biased towards past movements.

These biases arise in response to past active training (Verstynen & Sabes, 2011) as well as passive

training (Diedrichsen et al., 2010). In our paradigm, it did not appear that passive training

improved learning via a simple biasing in reach direction; when we measured participant reach

angles on the initial rotation trials which followed passive training, we did not observe any biases

in reach angle (Figs. 2D-F). Thus, while we have previously observed that passive training induces

use-dependent biases in reach angle (Wang et al., 2015; Bao et al., 2017; Lei et al., 2017), it did

not appear that these biases were strong enough to "pull" the arm toward the experienced

movement direction in our experimental conditions. This may have been due to the uniform

target arrangement in our experiment (Verstynen & Sabes, 2011) which could "average out" use-

dependent learning across the 4 symmetric targets in the workspace.

A third intriguing hypothesis is that passive training engaged both a strategic explicit

system that can be guided by instruction (Taylor et al., 2014; Mazzoni & Krakauer, 2006;

McDougle & Taylor, 2019), and an implicit system that adapts without our conscious awareness

(Avraham et al., 2020; Miyamoto et al., 2020; Javidialsaadi and Wang, 2020; Wang et al., 2019;

Jo et al., 2022). Might an interplay between these two systems be related to the time-dependent

error sensitivity patterns observed during rotation learning? Because we did not measure implicit

or explicit learning, we cannot know the answer to this question. However, we did observe that

reaction time (Fig. 5A) appeared elevated after passive training in the experimental groups, which

commonly accompanies cognitive operations involved in explicit strategy use (Fernandez-Ruiz et

al., 2011; Sakamoto and Kondo, 2012; Anguera et al., 2010; Georgopoulos and Massey, 1987;

McDougle et al., 2019; Albert et al., 2020).

Figure 5. Changes in reaction time. **A.** We measured reaction time as the difference between target appearance and reach onset. Here we show reaction time during the perturbation period in experimental (black) and control (gray) groups. **B.** We investigated early differences in reaction time by calculating the mean reaction time over the first three cycles. Statistics show post-hoc Tukey's test following one-way ANOVA (** indicates p<0.01). Error bars show mean ± SEM.

With that said, changes in reaction time were only statistically significant in the 5 min group (Fig. 5B) and appeared to dissipate quite rapidly in the 1-hr and 24-hr groups. If explicit strategies did contribute to increases in error sensitivity, it is also puzzling why reaction time was greatest in the 5 min group when this group was primarily enhanced by a slow learning state (Figs. 4D&E), which is thought to reflect implicit learning processes (McDougle et al., 2015). Nevertheless, the idea that passive training may evoke changes in both implicit and explicit processes is a fascinating possibility, which remains to be more formally tested.

Conclusion

In summary, passive motor training led to improvements in active motor learning due to increases in error sensitivity. Fast and slow processes appeared to mediate these changes in learning but were each modulated over different timescales. With little time-passage between

passive and active training, error sensitivity was enhanced in the slow adaptive state. With

greater time-passage, error sensitivity was enhanced in the fast adaptive state. These changes in

learning may have been mediated by interactions between the cerebral and cerebellar cortices,

or potentially through implicit and explicit learning systems.

References

Albert, S. T., & Shadmehr, R. (2018). Estimating properties of the fast and slow adaptive processes

during sensorimotor adaptation. *Journal of neurophysiology*, *119*(4), 1367-1393.

Albert, Scott T., Jihoon Jang, Adrian M. Haith, Gonzalo Lerner, Valeria Della-Maggiore, John W.

Krakauer, and Reza Shadmehr. "Competition between parallel sensorimotor learning

systems." *bioRxiv* (2020).

Albert, S. T. (2020). *The role of error-based learning in movement and stillness* (Doctoral

dissertation, Johns Hopkins University).

Aisen, M. L., Krebs, H. I., Hogan, N., McDowell, F., & Volpe, B. T. (1997). The effect of robot-

assisted therapy and rehabilitative training on motor recovery following stroke. *Archives of*

neurology, *54*(4), 443-446.

Anguera, J. A., Reuter-Lorenz, P. A., Willingham, D. T., & Seidler, R. D. (2010). Contributions of

spatial working memory to visuomotor learning. *Journal of cognitive neuroscience*, *22*(9), 1917-

1930.

Avraham, G., Morehead, J. R., Kim, H. E., & Ivry, R. B. (2021). Reexposure to a sensorimotor

perturbation produces opposite effects on explicit and implicit learning processes. *PLoS*

biology, *19*(3), e3001147.

Bao, S., Lei, Y., & Wang, J. (2017). Experiencing a reaching task passively with one arm while

adapting to a visuomotor rotation with the other can lead to substantial transfer of motor

learning across the arms. *Neuroscience letters*, *638*, 109-113.

Bara, F., & Gentaz, E. (2011). Haptics in teaching handwriting: The role of perceptual and visuo-

motor skills. *Human movement science*, *30*(4), 745-759.

Basteris, A., Bracco, L., & Sanguineti, V. (2012). Robot-assisted intermanual transfer of

handwriting skills. *Human movement science*, *31*(5), 1175-1190.

Beets, I. A., Macé, M., Meesen, R. L., Cuypers, K., Levin, O., & Swinnen, S. P. (2012). Active versus

passive training of a complex bimanual task: is prescriptive proprioceptive information sufficient

for inducing motor learning?. *PLoS One*, *7*(5), e37687.

Bozorgpour, R. (2023). Computational Explorations in Biomedicine: Unraveling Molecular

Dynamics for Cancer, Drug Delivery, and Biomolecular Insights using LAMMPS Simulations. *arXiv*

preprint arXiv:2311.13000.

Bozorgpour, R., Sheybanikashani, S., & Mohebi, M. (2023). Exploring the Role of Molecular

Dynamics Simulations in Most Recent Cancer Research: Insights into Treatment Strategies. *arXiv*

preprint arXiv:2310.19950.

Brashers-Krug, T., Shadmehr, R., & Bizzi, E. (1996). Consolidation in human motor

memory. *Nature*, *382*(6588), 252-255.

Carel, C., Loubinoux, I., Boulanouar, K., Manelfe, C., Rascol, O., Celsis, P., & Chollet, F. (2000).

Neural substrate for the effects of passive training on sensorimotor cortical representation: a

study with functional magnetic resonance imaging in healthy subjects. *Journal of Cerebral Blood*

Flow & Metabolism, *20*(3), 478-484.

Castro, L. N. G., Hadjiosif, A. M., Hemphill, M. A., & Smith, M. A. (2014). Environmental

consistency determines the rate of motor adaptation. *Current Biology*, *24*(10), 1050-1061.

Coltman, S. K., Cashaback, J. G., & Gribble, P. L. (2019). Both fast and slow learning processes

contribute to savings following sensorimotor adaptation. *Journal of neurophysiology*, *121*(4),

1575-1583.

Coltman, S. K., & Gribble, P. L. (2020). Time course of changes in the long-latency feedback

response parallels the fast process of short-term motor adaptation. *Journal of*

Neurophysiology, *124*(2), 388-399.

Day, K. A., Leech, K. A., Roemmich, R. T., & Bastian, A. J. (2018). Accelerating locomotor savings

in learning: compressing four training days to one. *Journal of neurophysiology*, *119*(6), 2100-

2113.

Della-Maggiore, V., & McIntosh, A. R. (2005). Time course of changes in brain activity and

functional connectivity associated with long-term adaptation to a rotational

transformation. *Journal of neurophysiology*, *93*(4), 2254-2262.

Diedrichsen, J., Shadmehr, R., & Ivry, R. B. (2010). The coordination of movement: optimal

feedback control and beyond. *Trends in cognitive sciences*, *14*(1), 31-39.

Diedrichsen, J., Criscimagna-Hemminger, S. E., & Shadmehr, R. (2007). Dissociating timing and

coordination as functions of the cerebellum. *Journal of Neuroscience*, *27*(23), 6291-6301.

Fernandez-Ruiz, J., Wong, W., Armstrong, I. T., & Flanagan, J. R. (2011). Relation between reaction

time and reach errors during visuomotor adaptation. *Behavioural brain research*, *219*(1), 8-14.

Georgopoulos, A. P., & Massey, J. T. (1987). Cognitive spatial-motor processes. *Experimental*

brain research, *65*(2), 361-370.

Gibo, T. L., Criscimagna-Hemminger, S. E., Okamura, A. M., & Bastian, A. J. (2013). Cerebellar

motor learning: are environment dynamics more important than error size?. *Journal of*

neurophysiology, *110*(2), 322-333.

Haith AM, Huberdeau DM, Krakauer JW. The influence of movement preparation time on the

expression of visuomotor learning and savings. J Neurosci 35: 5109–5117, 2015.

Herzfeld, D. J., Hall, N. J., Tringides, M., & Lisberger, S. G. (2020). Principles of operation of a

cerebellar learning circuit. *Elife*, *9*, e55217.

Herzfeld, D. J., Kojima, Y., Soetedjo, R., & Shadmehr, R. (2018). Encoding of error and learning to

correct that error by the Purkinje cells of the cerebellum. *Nature neuroscience*, *21*(5), 736-743.

Herzfeld DJ, Vaswani PA, Marko MK, Shadmehr R. A memory of errors in sensorimotor learning.

Science 345: 1349–1353, 2014.

Herzfeld, D. J., Kojima, Y., Soetedjo, R., & Shadmehr, R. (2015). Encoding of action by the Purkinje

cells of the cerebellum. *Nature*, *526*(7573), 439-442.

Huberdeau, D. M., Haith, A. M., & Krakauer, J. W. (2015). Formation of a long-term memory for

visuomotor adaptation following only a few trials of practice. *Journal of neurophysiology*, *114*(2),

969-977.

Huberdeau, D. M., Krakauer, J. W., & Haith, A. M. (2019). Practice induces a qualitative change in

the memory representation for visuomotor learning. *Journal of neurophysiology*, *122*(3), 1050-

1059.

Ito, K., Doi, M., Kondo, T.: Feedforward Adaptation to a Varying Dynamical Environment during

Reaching Movements, Journal of Robotics Mechatronics, 19, 4, 474–481 (2007).

Izawa, J., Criscimagna-Hemminger, S. E., & Shadmehr, R. (2012). Cerebellar contributions to reach

adaptation and learning sensory consequences of action. *Journal of Neuroscience*, *32*(12), 4230-

4239.

Javidialsaadi, M., & Wang, J. (2021). Lack of interlimb transfer following visuomotor adaptation

in a person with congenital mirror movements despite the awareness of the visuomotor

perturbation. *Brain and Cognition*, *147*, 105653.

Jax, S. A., & Rosenbaum, D. A. (2007). Hand path priming in manual obstacle avoidance: evidence

that the dorsal stream does not only control visually guided actions in real time. *Journal of*

Experimental Psychology: Human Perception and Performance, *33*(2), 425.

Jo, Y., Javidialsaadi, M., & Wang, J. (2022). Facilitative effects of use-dependent learning on

interlimb transfer of visuomotor adaptation in a person with congenital mirror

movements. *Human Movement Science*, *84*, 102973.

Kahn, L. E., Zygman, M. L., Rymer, W. Z., & Reinkensmeyer, D. J. (2006). Robot-assisted reaching

exercise promotes arm movement recovery in chronic hemiparetic stroke: a randomized

controlled pilot study. *Journal of neuroengineering and rehabilitation*, *3*(1), 1-13.

Kim, H. E., Avraham, G., & Ivry, R. B. (2020). The psychology of reaching: action selection,

movement implementation, and sensorimotor learning. *Annual review of psychology*, *72*.

Kitago T, Ryan SL, Mazzoni P, Krakauer JW, Haith AM. Unlearning versus savings in visuomotor

adaptation: comparing effects of washout, passage of time, and removal of errors on motor

memory. Front Hum Neurosci 7: 307, 2013.

Kojima Y, Iwamoto Y, Yoshida K. Memory of learning facilitates saccadic adaptation in the

monkey. *J Neurosci* 24: 7531–7539, 2004.

Krebs, H. I., Hogan, N., Aisen, M. L., & Volpe, B. T. (1998). Robot-aided neurorehabilitation. *IEEE*

transactions on rehabilitation engineering, *6*(1), 75-87.

Kumar, N., Manning, T. F., Ostry, D. G. (2019). Somatosensory cortex participates in the

consolidation of human motor memory. *PLoS Biol* 17(10): e3000469.

Lei, Y., Bao, S., & Wang, J. (2016). The combined effects of action observation and passive

proprioceptive training on adaptive motor learning. *Neuroscience*, *331*, 91-98.

Lei, Y., Bao, S., Perez, M. A., & Wang, J. (2017). Enhancing generalization of visuomotor

adaptation by inducing use-dependent learning. *Neuroscience*, *366*, 184-195.

Leow LA, de Rugy A, Marinovic W, Riek S, Carroll TJ. Savings for visuomotor adaptation require

prior history of error, not prior repetition of successful actions. *J Neurophysiol* 116: 1603–1614,

2016.

Leow, L. A., De Rugy, A., Loftus, A. M., & Hammond, G. (2013). Different mechanisms contributing

to savings and anterograde interference are impaired in Parkinson's disease. *Frontiers in human*

neuroscience, *7*, 55.

Lerner, G., Albert, S., Caffaro, P. A., Villalta, J. I., Jacobacci, F., Shadmehr, R., & Della-Maggiore, V.

(2020). The origins of anterograde interference in visuomotor adaptation. *Cerebral Cortex*, *30*(7),

4000-4010.

Lisberger, S. G. (1994). Neural basis for motor learning in the vestibuloocular reflex of primates.

III. Computational and behavioral analysis of the sites of learning. *Journal of*

Neurophysiology, *72*(2), 974-998.

130 Maschke, M., Gomez, C. M., Ebner, T. J., & Konczak, J. (2004). Hereditary cerebellar ataxia

131 progressively impairs force adaptation during goal-directed arm movements. *Journal of*

132 *neurophysiology*, *91*(1), 230-238.

133 Mawase, F., Shmuelof, L., Bar-Haim, S., & Karniel, A. (2014). Savings in locomotor adaptation

134 explained by changes in learning parameters following initial adaptation. *Journal of*

135 *neurophysiology*, *111*(7), 1444-1454.

136 Mazzoni, P., & Krakauer, J. W., (2006). An implicit plan overrides an explicit strategy during

137 visuomotor adaptation. Journal of neuroscience, 26(14), 3642-3645.

138 McCormick, D. A., & Thompson, R. F. (1984). Neuronal responses of the rabbit cerebellum during

139 acquisition and performance of a classically conditioned nictitating membrane-eyelid

140 response. *Journal of Neuroscience*, *4*(11), 2811-2822.

141 McDougle, S. D., Bond, K. M., & Taylor, J. A., (2015). Explicit and implicit processes constitute the

142 fast and slow processes of sensorimotor learning. Journal of Neuroscience, 35(26), 9568-9579.

143 McDougle, S. D., & Taylor, J. A. (2019). Dissociable cognitive strategies for sensorimotor

144 learning. *Nature communications*, *10*(1), 1-13.

145 Miyamoto, Y. R., Wang, S., & Smith, M. A. (2020). Implicit adaptation compensates for erratic

146 explicit strategy in human motor learning. *Nature neuroscience*, *23*(3), 443-455.

Morehead, J. Ryan, Salman E. Qasim, Matthew J. Crossley, and Richard Ivry. "Savings upon re-

aiming in visuomotor adaptation." *Journal of neuroscience* 35, no. 42 (2015): 14386-14396.

Ohmae, S., & Medina, J. F. (2015). Climbing fibers encode a temporal-difference prediction error

during cerebellar learning in mice. *Nature neuroscience*, *18*(12), 1798-1803.

Orozco, S. P., Albert, S. T., & Shadmehr, R. (2020). Spontaneous recovery and the multiple

timescales of human motor memory. *bioRxiv*.

Perrett, S. P., Ruiz, B. P., & Mauk, M. D. (1993). Cerebellar cortex lesions disrupt learning-

dependent timing of conditioned eyelid responses. *Journal of Neuroscience*, *13*(4), 1708-1718.

Popa, L. S., & Ebner, T. J. (2019). Cerebellum, predictions and errors. *Frontiers in cellular*

neuroscience, *12*, 524.

Popa, L. S., Hewitt, A. L., & Ebner, T. J. (2012). Predictive and feedback performance errors are

signaled in the simple spike discharge of individual Purkinje cells. *Journal of Neuroscience*, *32*(44),

15345-15358.

Reinkensmeyer, D. J., & Patton, J. L. (2009). Can robots help the learning of skilled

actions?. *Exercise and sport sciences reviews*, *37*(1), 43.

Riener, R., Nef, T., & Colombo, G. (2005). Robot-aided neurorehabilitation of the upper

extremities. *Medical and biological engineering and computing*, *43*(1), 2-10.

Rasmussen, A., Jirenhed, D. A., & Hesslow, G. (2008). Simple and complex spike firing patterns in
Purkinje cells during classical conditioning. *The Cerebellum*, *7*(4), 563.

Sakamoto, T., & Kondo, T. (2012, November). Can passive arm movement affect adaptation to
visuomotor rotation?. In *2012 IEEE International Conference on Development and Learning and*
Epigenetic Robotics (ICDL) (pp. 1-6). IEEE.

Shadmehr, R., Smith, M. A., & Krakauer, J. W. (2010). Error correction, sensory prediction, and
adaptation in motor control. *Annual review of neuroscience*, *33*, 89-108.

Scheidt, R. A., Conditt, M. A., Secco, E. L., & Mussa-Ivaldi, F. A. (2005). Interaction of visual and
proprioceptive feedback during adaptation of human reaching movements. *Journal of*
neurophysiology, *93*(6), 3200-3213.

Sing, G. C., & Smith, M. A. (2010). Reduction in learning rates associated with anterograde
interference results from interactions between different timescales in motor adaptation. *PLoS*
Comput Biol, *6*(8), e1000893.

Smith MA, Ghazizadeh A, Shadmehr R. Interacting adaptive processes with different timescales
underlie short-term motor learning. *PLoS Biol* 4: e179, 2006.

Steinmetz, J. E., Sears, L. L., Gabriel, M., Kubota, Y., & Poremba, A. (1991). Cerebellar interpositus
nucleus lesions disrupt classical nictitating membrane conditioning but not discriminative
avoidance learning in rabbits. *Behavioural brain research*, *45*(1), 71-80.

Taylor, J. A., Krakauer, J. W., & Ivry, R. B., (2014). Explicit and implicit contributions to learning in

a sensorimotor adaptation task. Journal of Neuroscience, 34(8), 3023-3032.

Tays, G., Bao, S., Javidialsaadi, M., & Wang, J. (2020). Consolidation of use-dependent motor

memories induced by passive movement training. *Neuroscience Letters*, *732*, 135080.

Tong, C., & Flanagan, J. R. (2003). Task-specific internal models for kinematic

transformations. *Journal of Neurophysiology*, *90*(2), 578-585.

Tseng, Y. W., Diedrichsen, J., Krakauer, J. W., Shadmehr, R., & Bastian, A. J. (2007). Sensory

prediction errors drive cerebellum-dependent adaptation of reaching. *Journal of*

neurophysiology, *98*(1), 54-62.

Vergaro, E., Casadio, M., Squeri, V., Giannoni, P., Morasso, P., & Sanguineti, V. (2010). Self-

adaptive robot training of stroke survivors for continuous tracking movements. *Journal of*

neuroengineering and rehabilitation, *7*(1), 1-12.

Verstynen, T., & Sabes, P. N. (2011). How each movement changes the next: an experimental and

theoretical study of fast adaptive priors in reaching. *Journal of Neuroscience*, *31*(27), 10050-

10059.

Wang, J., & Lei, Y. (2015). Direct-effects and after-effects of visuomotor adaptation with one arm

on subsequent performance with the other arm. *Journal of neurophysiology*, *114*(1), 468-473.

Wang, J., Bao, S., & Tays, G. D. (2019). Lack of generalization between explicit and implicit

visuomotor learning. *PloS one*, *14*(10), e0224099.

Wigmore, V., Tong, C., & Flanagan, J. R. (2002). Visuomotor rotations of varying size and direction

compete for a single internal model in a motor working memory. *Journal of Experimental*

Psychology: Human Perception and Performance, *28*(2), 447.

Yin, C., & Wei, K. (2020). Savings in sensorimotor adaptation without an explicit strategy. *Journal*

of neurophysiology, *123*(3), 1180-1192.

Zarahn, E., Weston, G. D., Liang, J., Mazzoni, P., & Krakauer, J. W. (2008). Explaining savings for

visuomotor adaptation: linear time-invariant state-space models are not sufficient. *Journal of*

neurophysiology, *100*(5), 2537-2548.

209