1

The dual function monoclonal antibodies VIR-7831 and VIR-7832 demonstrate potent in vitro and

in vivo activity against SARS-CoV-2

Andrea L. Cathcart[1], Colin Havenar-Daughton[1], Florian A. Lempp[1], Daphne Ma[1], Michael A. Schmid[2],

Maria L. Agostini[1], Barbara Guarino[2], Julia Di iulio[1], Laura E. Rosen[1], Heather Tucker[1], Joshua Dillen[1],

Sambhavi Subramanian[1], Barbara Sloan[1], Siro Bianchi[2], Dora Pinto[2], Christian Saliba[2], Jason A

Wojcechowskyj[1], Julia Noack[1], Jiayi Zhou[1], Hannah Kaiser[1], Arthur Chase[1], Martin Montiel-Ruiz[1],

Exequiel Dellota Jr.[1], Arnold Park[1], Roberto Spreafico[1], Anna Sahakyan[1], Elvin J. Lauron[1], Nadine

Czudnochowski[1], Elisabetta Cameroni[1], Sarah Ledoux[1], Adam Werts[3], Christophe Colas[1], Leah Soriaga[1],

Amalio Telenti[1], Lisa A. Purcell[1], Seungmin Hwang[1], Gyorgy Snell[1], Herbert W. Virgin[1], Davide Corti[2],

Christy M. Hebner[1*]

[1]Vir Biotechnology, San Francisco, California 94158, USA

[2]Humabs Biomed SA, a subsidiary of Vir Biotechnology, 6500 Bellinzona, Switzerland

[3]Lovelace Biomedical, Albuquerque, New Mexico 87108, USA

*corresponding author: chebner@vir.bio

ABSTRACT

VIR-7831 (sotrovimab) and VIR-7832 are dual action monoclonal antibodies (mAbs) targeting the

spike glycoprotein of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). VIR-7831

and VIR-7832 were derived from a parent antibody (S309) isolated from memory B cells of a 2003

severe acute respiratory syndrome coronavirus (SARS-CoV) survivor. Both mAbs contain an "LS"

mutation in the Fc region to prolong serum half-life. In addition, VIR-7832 encodes an Fc GAALIE

mutation that has been shown previously to evoke CD8+ T-cells in the context of an in vivo viral

respiratory infection. VIR-7831 and VIR-7832 potently neutralize wild-type and variant authentic

virus in vitro as well as variant pseudotyped viruses. In addition, they retain activity against

monoclonal antibody resistance mutations conferring reduced susceptibility to currently authorized

mAbs. The VIR-7831/VIR-7832 epitope does not overlap with mutational sites in current variants

of concern and continues to be highly conserved among circulating sequences consistent with the

high barrier to resistance observed in vitro. Furthermore, both mAbs can recruit effector

mechanisms in vitro that may contribute to clinical efficacy via elimination of infected host cells. In

vitro studies with these mAbs demonstrated no enhancement of infection. In a Syrian Golden

hamster proof-of concept wildtype SARS-CoV-2 infection model, animals treated with VIR-7831

had less weight loss, and significantly decreased total viral load and infectious virus levels in the

lung compared to a control mAb. Taken together, these data indicate that VIR-7831 and VIR-7832

are key agents in the fight against COVID-19.

INTRODUCTION

The coronavirus disease (COVID-19) pandemic caused by severe acute respiratory syndrome coronavirus

2 (SARS-CoV-2) has resulted in more than 260 million confirmed cases and over 5.2 million deaths

worldwide[1]. SARS-CoV-2 infection results in a broad range of disease severity[2]. Infection fatality rates

increase significantly with age, with 28.3% of COVID-19 patients over the age of 85 succumbing to

disease[2]. However, even in mild-to-moderate COVID-19 patients, significant post-infection sequelae can

affect overall health and cause long-term disability[3]. While multiple SARS-CoV-2 vaccines are now

available for use, issues of supply, vaccine hesitancy and emergence of variants may prevent rapid

attainment of herd immunity[4–9]. In addition, there are individuals who remain at risk despite vaccination

due to disease or underlying immunodeficiency. Thus, additional interventions and potential prophylactic

agents are needed to reduce morbidity and mortality due to COVID-19.

Several monoclonal antibodies (mAbs) targeting the SARS-CoV-2 spike protein have recently been

authorized or approved for use in early treatment of COVID-19 patients[10–13] and clinical data have shown

promising results in treatment and prophylactic studies[12–16]. However, rapidly spreading variants exhibit

reduced susceptibility in vitro to some currently authorized antibodies that target the receptor binding

motif (RBM) of the viral spike (S) glycoprotein [10,11,17,18]. Therefore, mAbs targeting unique S epitopes are

needed for use alone or in combination with current agents for the treatment and prevention of COVID-

19. Furthermore, in addition to viral neutralization, antibodies possessing potent effector function to aid in

the killing of virally infected cells and the elicitation of T cell immunity could significantly assist in

halting disease progression[19–21].

VIR-7831 and VIR-7832 are dual action mAbs derived from the parent antibody S309 identified from a

2003 SARS-CoV survivor[22]. These mAbs target an epitope containing a glycan (at position N343) that is

highly conserved within the Sarbecovirus subgenus in a region of the S receptor binding domain (RBD)

that does not compete with angiotensin converting enzyme 2 (ACE2) binding[23]. [10,11,17,18]The variable

region of VIR-7831 and VIR-7832 have been engineered for enhanced developability. In addition, both

antibodies possess an Fc "LS" mutation that confers extended half-life by binding to the neonatal Fc

receptor[24–26]. VIR-7832 is identical to VIR-7831 with the exception of the addition of a 3 amino acid

GAALIE (G236A, A330L, I332E) modification to the Fc domain[27]. The GAALIE modification has

previously been shown in vitro to enhance binding to FcγIIa and FcγIIIa receptors, decrease affinity for

FcγIIb compared to typical IgG1 and evoke protective CD8+ T-cells in the context of viral respiratory

infection in vivo [28,29].

Here we characterize the antiviral activity of VIR-7831 and VIR-7832. These mAbs effectively neutralize

SARS-CoV-2 live virus in vitro as well as in pseudotyped virus assays against emerging variants of

concern and variants that confer resistance to currently authorized or approved mAbs[30]. In addition to the

neutralizing capacity, both antibodies demonstrate potent effector function and mediate antibody

dependent cellular cytotoxicity (ADCC) and antibody dependent cellular phagocytosis (ADCP) in vitro.

Furthermore, resistance selection experiments and epitope conservation analyses indicate the potential for

a high barrier to resistance. Data derived from the Syrian golden hamster model demonstrates efficacy in

a proof-of-concept in vivo model. Taken together, these data indicate that VIR-7831 and VIR-7832 are

promising key components of the arsenal in the fight against COVID-19.

RESULTS

VIR-7831 and VIR-7832 bind SARS-CoV-2 spike and effectively neutralize live virus in vitro.

Previously published work showed that S309 bound SARS-CoV-2 recombinant and cell surface-

associated S and neutralized live virus in vitro[22]. We initiated these studies by repeating and extending

these earlier results. To determine the binding activity of VIR-7831 and VIR-7832 to the SARS-CoV-2 S,

enzyme-linked immunosorbent assay (ELISA), surface plasmon resonance (SPR) and flow cytometry

assays were utilized. VIR-7831 and VIR-7832 bound to recombinant S RBD (amino acids 331-541) with

EC_{50} values of 20.40 ng/mL and 14.9 ng/mL, respectively, by ELISA (**Figure 1a**). Using SPR, both

antibodies demonstrated potent binding to recombinant S RBD with an equilibrium constant (Kd) of 0.21

nM (**Figure 1b**). As antibody recognition of cell surface-bound S could mediate killing of virally infected

cells, flow cytometry-based studies using cells transiently transfected with a S-encoding plasmid were

used to examine antibody binding to cell surface-expressed S trimer. By this method, both VIR-7831 and

VIR-7832 bound efficiently to surface-expressed S (**Figure 1c**).

To examine neutralization capacity, VIR-7831 and VIR-7832 were tested in a VeroE6 cell-based live

SARS-CoV-2 virus system against the Washington 2019 (wild-type) virus as well as against the Alpha

(B.1.1.7), Beta (B.1.351), Gamma (P.1), Delta (B.1.617.2) and Kappa (B.1.617.1) variants.

Concentration-dependent viral neutralization of the Washington 2019 strain was observed for both

antibodies, with geometric mean $IC_{50/90}$ values of 100.1/186.3 ng/mL and 78.3/253.1 ng/mL, respectively

(**Figure 1d**). $IC_{50/90}$ values observed for VIR-7831 and VIR-7832 against the Beta, Gamma, Delta and

Kappa variant viruses were similar to those against the wild-type strain. A slight shift in the VIR-

7831/VIR-7832 $IC_{50/90}$ compared to wild-type was observed for the Alpha variant. VIR-7831 showed a 3-

fold and 4.1-fold shift in IC_{50} and IC_{90}, respectively, against the Alpha variant compared to wild-type

while VIR-7832 had a 3.1-fold shift in IC_{50} and 3.7-fold shift in IC_{90} versus wild-type (**Figure 1d, Table**

1). As variant evolution is a natural part of SARS-CoV-2 biology and emerging live virus variants are not

always readily accessible for testing, a vesicular stomatitis virus (VSV)-based pseudotyped virus system

targeting Vero E6 cells was used to examine VIR-7831 and VIR-7832 neutralization against emergent

variants (**Table 2**). Fold-changes in VIR-7831 and VIR-7832 IC_{50} values compared to wild-type against

pseudotyped virus expressing spike from the Alpha, Beta, Gamma, Delta or Kappa variants were similar

to those observed in the authentic virus system. VIR-7831 was tested against an extended panel of

pseudotyped viruses incorporating emerging variants as well as variants deemed as Variants of Concern

(VOC) or Variants of Interest (VOI) by the World Health Organization (WHO). VIR-7831 retained

activity in vitro against all variants tested with fold changes in IC_{50}s ranging from 0.4- to 2.3-fold (**Table**

2).

Neutralization of pseudotyped virus encoding individual omicron substitutions. On November 26,

2021, the WHO designated Omicron (B.1.1.529) as an official VOC. Omicron is reported to encode ~30

substitutions in the spike, including substitutions that may render some currently authorized or approved

COVID-19 mAbs inactive [31–33]. VIR-7831 was tested against a panel of available pseudotyped viruses

encoding a subset of individual mutations found in the Omicron spike (Figure 2a). In vitro pseudotyped

virus data demonstrate that VIR-7831 retains neutralizing activity against all tested individual Omicron

substitutions including the G339D, K417N, N440K, S477N, T478K, Q493R and N501Y substitutions

found in the RBD of the Omicron spike. While data for G446S and E484A are not yet available, G446V

and E484K substitutions have been tested in a pseudotyped virus assay and VIR-7831 retains activity

against these mutations. The spike amino acids G339 and N440 are both part of the VIR-7831 epitope. In

vitro studies using pseudotyped virus encoding either the G339D or N440K substitutions show that VIR-

7831 retains activity against these spike substitutions (with IC_{50} fold changes of 1.2 and 0.5, respectively)

(Figure 2a, 2b). In addition, a lysine substitution has been detected at the VIR-7831 epitope position R346

in a small proportion of available Omicron sequences as of November 29, 2021. VIR-7831 also retains

activity in vitro against pseudotyped virus carrying the R346K substitution with a 0.7-fold-change in IC_{50}.

While the effect of other individual Omicron substitutions in the RBD on VIR-7831 activity have not yet

been studied in the pseudotyped virus system (S371L, S373P, S375F, G446S, E484A, G496S, Q498R,

Y505H), notably none of these amino acid positions are part of the sotrovimab epitope. These data for the

individual RBD substitutions in the Omicron spike suggest that VIR-7831 is likely to retain activity

against the Omicron variant.

VIR-7831 and VIR-7832 exhibit potent effector function in vitro. Although direct antiviral

mechanisms are crucial to provide protection, Fc-dependent mechanisms mediated by interaction with Fc

gamma receptors (FcγRs) on immune cells or with complement, can contribute to overall potency in vivo.

The potential for VIR-7831 and VIR-7832 to mediate effector functions were assessed in vitro by

measuring binding to FcγRs and C1q and in assays designed to demonstrate antibody-dependent cellular

cytotoxicity (ADCC) or antibody-dependent cellular phagocytosis (ADCP)[34–37].

Antibody binding to the human activating FcγRIIa (low-affinity R131 and high affinity H131 alleles),

FcγRIIIa (low-affinity F158 and high-affinity V158 alleles), and to the inhibitory FcγRIIb were examined

using SPR (**Supplemental figure 1a**). VIR-7831 similarly bound both the H131 and R131 alleles of

FcγRIIa and binds FcγRIIb. VIR-7831 bound both FcγRIIIa alleles, with reduced binding to the F158

allele compared to V158, as is characteristic for human IgG1[38]. Binding of VIR-7831 to C1q was similar

to the parental antibody (S309-LS) (**Supplemental figure 1b**). As previously reported for antibodies

encoding the GAALIE mutation[27,39], VIR-7832 bound with comparatively higher affinity to activating

FcγRIIa and FcγRIIIa than VIR-7831 (**Supplemental figure 1a**). Conversely, VIR-7832 showed reduced

affinity for FcγIIb and abrogation of binding to C1q (**Supplemental figure 1b**).

The antibodies were also assessed for the ability to activate human FcγRIIa, FcγRIIb or FcγRIIIa, using a

Jurkat cell reporter assay[40] (**Figure 3a-d**). S309-GRLR, which contains the effector function-abrogating

G236R, L328R mutations was used as a negative control. Cells stably transfected with the SARS-CoV-2

spike protein (CHO-CoV-2-Spike) served as target cells. Both VIR-7831 and the parental S309-LS

activated signaling of the higher-affinity allele FcγRIIa (H131) but did so less efficiently than the

GAALIE-containing antibody VIR-7832 (**Figure 3a**) while VIR-7831, VIR-7832 and S309-LS induced

similar low-level activation of the inhibitory receptor FcγRIIb (**Figure 3b**). VIR-7831 demonstrated

substantially lower activation of FcγRIIIa F158 versus V158 as expected while VIR-7832 showed

increased activation of both alleles of FcγRIIIa (F158 and V158) (**Figures 3c,d**).

To further elucidate the effector function potential of the antibodies, ADCC and ADCP assays were

performed using donor PBMCs or NK cells as effector cells and CHO cells stably expressing S (CHO-

CoV-2-Spike) as target cells (**Figure 3e-g**). The ability of antibodies to activate NK cell-mediated killing

was measured in vitro using two genotyped donors expressing homozygous low-affinity (F/F158) or high-

affinity (V/V158) (**Figure 3e-f**). Compared to the parental mAb S309-LS, VIR-7831 had slightly

increased capacity to induce NK cell-mediated ADCC when using cells from either F/F158 or V/V158

donors. As expected, VIR-7832 induced NK cell-mediated ADCC in cells from donors expressing the

low-affinity F/F158 allele of FcγIIIa more efficiently than VIR-7831. These results were confirmed with

NK cells from a heterozygous donor (F/V158).

The ability of VIR-7831 and VIR-7832 to facilitate ADCP by primary CD14[+] monocytes was measured in

vitro by exposing freshly isolated human PBMCs to CHO-CoV-2-Spike cells that were pre-incubated

with antibody (**Figure 3g**). VIR-7831, VIR-7832 and S309-LS induced similar levels of ADCP by CD14[+]

monocytes. These results indicate that VIR-7831 and VIR-7832 have the potential to trigger ADCC and

ADCP of cells displaying SARS CoV-2 S protein.

Subneutralizing levels of VIR-7831 and VIR-7832 do not enhance virus uptake, replication or

cytokine production in vitro. One potential concern with any antibody therapeutic targeting a viral agent

is the possibility of antibody-dependent enhancement (ADE). ADE is an in vivo phenomenon in which

the presence of an antibody worsens disease. There are several in vitro assays that may provide plausible

correlates for ADE in vivo, though none of these have been proven relevant to COVID-19 as to date ADE

has not been observed in trials of monoclonal antibodies or plasma[13,15,41,42]. ADE can occur by several

potential mechanisms[43]. Poorly neutralizing antibodies or subneutralizing levels of antibody could

theoretically facilitate enhanced virus entry and infection through Fc receptor interactions. A second

theoretical mechanism involves antibody-antigen complex formation leading to enhanced cytokine

production. A third mechanism of ADE has been observed in a porcine model of influenza where the

kinetics of viral fusion to the target cell was enhanced in a Fab-dependent manner by fusion-enhancing

non-neutralizing antibodies[44,45].

To explore whether VIR-7831 and VIR-7832 exhibit in vitro activities that might be related to ADE in

vivo, we evaluated SARS-CoV-2 replication in human cells that express FcγRs: monocyte-derived

dendritic cells (moDCs), peripheral blood mononuclear cells (PBMCs) and the human U937 macrophage

cell line (**Supplemental Figure 2a-b**). Subneutralizing concentrations of VIR-7831 and VIR-7832 were

precomplexed with SARS-CoV-2 (MOI =0.01) and added to target cells. Using immunostaining methods,

at 24 hours post-infection no productive entry of SARS-CoV-2 into moDCs, PBMCs, or U937 cells was

observed in the presence or absence of either mAb, while VeroE6 control cells demonstrated

internalization in all conditions evaluated. Reduced internalization of SARS-CoV-2 in VeroE6 cells was

observed at the highest concentration of VIR-7831 and VIR-7832 (p-value <0.05), indicating effective

virus neutralization prevented virus entry. Using a focus forming assay, virus replication and secretion of

infectious virus were detectable by 48 hours post-infection in VeroE6 cells, with comparable levels of

replication in the presence or absence of VIR-7831 or VIR-7832. However, no replication of SARS-CoV-

2 was detected in moDCs, PBMCs or U937 cells regardless of antibody treatment, indicating lack of

productive SARS-CoV-2 infection of these cells, consistent with previously published data[46].

To evaluate the potential for VIR-7831 and VIR-7832 to enhance cytokine release upon SARS-CoV-2

infection in FcγR-expressing cells, cytokines and chemokines were measured in the supernatants from

cells infected with SARS-CoV-2-in the presence of VIR-7831 or VIR-7832 (**Supplemental figure 2c**).

Levels of IFN-γ, IL-10, IL-6, IL 8, IP-10, MCP-1, and TNF-α in the supernatant were quantified by MSD

at 24- and 48-hours post-infection. For all cell types evaluated, cytokine/chemokine production was

similar between all antibody concentrations tested and the no antibody control at both 24- and 48-hours

post-infection. Taken together, these in vitro data indicate that neither VIR-7831 nor VIR-7832 exhibit in

vitro activities that have been proposed to possibly correlate with ADE in vivo.

VIR-7831 and VIR-7832 have a high barrier to resistance in vitro and do not display cross-

resistance with other SARS-CoV-2 mAbs. We next determined whether resistant variants could be

elicited by serial passage of SARS-CoV-2 in the presence of VIR-7832. As VIR-7831 and VIR-7832

differ only in the Fc region of the antibody, resistance selection experiments were conducted with VIR-

7832 as a proxy for both antibodies. SARS-CoV-2 was subjected to 10 passages in the presence of VIR-

7832 at fixed concentrations of ~10x, 20x, 50x or 100x IC_{50} (1, 2, 5, or 10 μg/mL) in VeroE6 cells. No

CPE was detected in wells passaged with antibody through 10 passages, while CPE was observed in the

no antibody control in all passages. Similarly, no virus was detected by focus forming assay at any

concentration of VIR-7832 through all 10 passages even at the lowest concentration tested.

As no viral breakthrough was observed in the fixed concentration resistance selection, a second method

was employed wherein SARS-CoV-2 virus was passaged in sub-IC_{50} concentrations of antibody followed

by subsequent passaging in the presence of increasing concentrations of mAb in an attempt to force

resistance emergence (**Supplemental Figure 3**). Passaging was performed in duplicate wells to account

for founder effects, and concentration increases for each well were based on CPE observations. Five

sequential passages were conducted using increasing concentrations of VIR-7832 at 0.5, 1, 2, 5 and ~10x

IC_{50} (0.05, 0.1, 0.2, 0.5, 1 µg/mL; **Supplemental Figure 3a**), though no CPE was observed by passages 4

and 5 (0.5 and 1 µg/mL, respectively) indicating that variants originally selected at the lower

concentrations were either unfit or susceptible to the higher concentrations of antibody. To further assess

whether resistance mutations could be generated, selection was restarted using passage 3 virus generated

with ~2x IC_{50} (0.2 µg/mL) of VIR-7832 in duplicate wells at ~2x and ~5x IC_{50} (0.2, 0.5 µg/mL),

generating two passage lineages (**Supplemental Figure 3b-c**).

Supernatants were evaluated for detectable virus at each passage by focus forming assay and cell

supernatants from viral passages containing detectable virus were tested in SARS-CoV-2 neutralization

assays to evaluate IC_{50} shifts as a marker of reduced susceptibility (**Supplemental Table 1**). With the

exception of passage 8, modest fold changes were observed, with shifts in IC_{50} values ranging from 5.4-

to 6.5-fold compared to the wild-type SARS-CoV-2 stock virus. In lineage 1, the passage 8 virus

displayed a >10-fold shift in IC_{50} (greater than highest concentration tested). Sequence analysis detected

an identical 4 amino acid insertion in the N-terminal domain (215-216insKLRS) and 5 amino acid

deletion in correspondence of the furin cleavage site (675-679del) in both lineages at all passages

sequenced, as well as the amino acid substitution E340A in lineage 1, and R682W, and V1128F in

lineage 2. The deletion at amino acids 675-679 has been previously described during passaging of SARS-

CoV-2 in tissue culture suggesting enrichment to be a result of cell culture adaptation[47] while the 215-

216insKLRS was detected in the input virus. Neither 215-216insKLRS nor R682W variants were highly

enriched with passaging (**Supplemental Table 1**) and enrichment of 675-679del and V1128F did not

profoundly alter the VIR-7832 IC_{50}. However, appearance of the E340A variant at 98.7% did correlate

with a >10-fold shift in IC_{50} suggesting this variant may confer resistance.

To evaluate whether amino acid variants identified in the resistance selection conferred reduced

susceptibility to VIR-7831 and VIR-7832, neutralization of pseudotyped viruses encoding the S variants

was assessed (**Supplemental Table 2**). VIR-7831 and VIR-7832 neutralized R682W and V1128F SARS-

CoV-2 pseudotyped virus spike variants with IC_{50} values similar to wild type (< 2-fold change in IC_{50})

indicating that these variants do not alter susceptibility. In contrast, E340A conferred reduced

susceptibility to VIR-7831 and VIR-7832 (> 100-fold change in IC_{50}) indicating that E340A is a VIR-

7831/VIR-7832 monoclonal antibody resistance mutation (MARM).

As VIR-7831/VIR-7832 demonstrated a unique in vitro resistance profile, we investigated the potential

for cross-resistance to MARMs that confer reduced susceptibility to the authorized monoclonal antibodies

bamlanivimab, imdevimab and casirivimab [10,11,48–50] using pseudotyped virus. Notably, some of these

mutations are found in highly prevalent variants of concern[18,51,52]. VIR-7831 effectively neutralized

pseudotyped viruses expressing spike MARMs that alter bamlanivimab, casirivimab and/or imdevimab

activity (**Table 3**). Fold changes in IC_{50} values compared to wild-type were <3-fold for 18/19 variants

tested. A modest 3.4-fold shift in the VIR-7831 IC_{50} was observed for the V445A variant that confers

reduced susceptibility to imdevimab. These data indicate that VIR-7831/VIR-7832 does not display cross-

resistance with currently authorized mAbs and supports the potential combination use of VIR-7831/VIR-

7832 with other mAb therapeutics.

The VIR-7831/VIR-7832 epitope is highly conserved among SARS-CoV-2 sequences. The parental

antibody of VIR-7831 and VIR-7832 (S309) binds to a highly conserved sarbecovirus epitope that is

potentially intolerant of variation. To investigate the current state of epitope conservation, $>4,500,000$

spike sequences from SARS-CoV-2 deposited in the GISAID database as of November 11, 2021 were

examined for epitope variation. More than 99.6% conservation is seen for those amino acids comprising

the epitope among currently available sequences for all positions including 15/23 amino acid positions

that were ≥ 99.99 conserved (**Table 4**).

VIR-7831 activity against viral mutants carrying single substitutions in the epitope was assessed in

pseudotyped virus assays. VIR-7831 effectively neutralized epitope variants at most amino acid positions

tested (Supplemental **Table 5**). A moderate shift in activity was observed for the K356T variant (5.9-fold

shift in IC_{50}). Variants at two positions, E340 and P337, resulted in significant IC_{50} shifts indicating

reduced susceptibility to VIR-7831. Moderate shifts in potency were observed for P337H and P337T

variants (5.13- and 10.62-fold, respective) while more significant shifts in potency were observed for

P337L/R/K and E340A/K/G/Q (18.21-fold to >300-fold). Notably, these variants are detected in a low

number of sequences and do not have a pattern that suggest emergence in the GISAID database (296 and

223 variant counts out of >4,500,000 sequences for P337 and E340, respectively). This observation is

consistent with the possibility that substitutions at these positions come with a fitness cost to the virus.

VIR-7831 reduces weight loss, total viral load and infectious virus levels in a hamster model of

SARS-CoV-2 infection. To evaluate the efficacy of VIR-7831 in vivo, the hamster model was utilized.

As it was unknown what effect the LS mutation would have in the hamster, a non-LS version of VIR-

7831 (SGHmAb-no-LS) was used for these experiments. Hamsters were administered SGHmAb-no-LS

intraperitoneally at Day -1 (30, 5, 0.5 or 0.05 mg/kg) or Day -2 (15, 5, 0.5 or 0.05 mg/kg) prior to

intranasal SARS-CoV-2 inoculation (**Figure 4a**). Using body weight as a marker of degree of clinical

disease, doses of ≥5mg/kg resulted in significantly reduced weight loss at Day 4 compared to controls.

(**Figure 4b-e**). Significant decreases in lung viral load were also observed at ≥5mg/kg as measured by

RT-qPCR (**Figures 4f-g**). Day 4 $TCID_{50}$ measurements indicated that antibody administered at ≥0.5

mg/kg resulted in significantly lower levels of infectious virus in lung tissue compared to controls

(**Figure 4h-i**). Notably, across these experiments, no enhancement of disease was observed in animals

receiving SGHmAb-no-LS based on changes in weight, viral RNA in the lungs, or $TCID_{50}$ infectious

virus levels. Collectively, these data indicate that VIR-7831 prevented in a dose-dependent fashion virus

replication and morbidity in SARS-CoV-2 challenged hamsters without signs of ADE at any dose tested.

DISCUSSION

Here we show the in vitro and in vivo preclinical characterization of VIR-7831 and VIR-7832[53–55]. Both

antibodies demonstrate high-affinity binding to S in vitro, including on the surface of cells, and

effectively neutralize wildtype SARS-CoV-2 in a live virus assay. VIR-7831 and VIR-7832 retain activity

against the Alpha (B.1.1.7), Beta (B.1.351), Gamma (P.1), Delta (B.1.617.2) and Kappa (B.1.617.1)

variants in an authentic virus system, consistent with data using pseudotyped viruses. For the Omicron

variant, VIR-7831 activity against individual spike substitutions was examined in a pseudotyped virus

system and demonstrated that VIR-7831 retains activity in vitro against the substitutions tested. The

single mutation neutralization data were consistent with previously published data examining the effects

of RBD amino acid substitutions on VIR-7831 using deep mutational scanning[56]. Using this method,

substitutions at amino acid positions 339, 371, 373, 375, 417, 440, 446, 477, 478, 484, 493, 496, 498, 501

and 505 pertinent to the Omicron spike substitutions did not alter binding of the VIR-7831 parent

antibody, S309, to the RBD. While in vitro testing to determine VIR-7831 activity against the full set of

substitutions detected in the Omicron (B.1.1.529) spike is underway, pseudotyped virus data for

individual spike substitutions as well as previously published RBD binding data evaluating individual

substitutions indicate that VIR-7831 is likely to retain activity against Omicron. VIR-7831 also binds

C1q, activate FcγRs and demonstrate ADCC and ADCP in vitro. Experiments in the hamster model of

SARS-CoV-2 infection show proof-of-concept efficacy in vivo. Notably, in vitro and in vivo data did not

provide any supporting evidence that these antibodies would demonstrate ADE in a clinical setting.

That VIR-7831 and VIR-7832 retain activity against spike variants in authentic virus and pseudotyped

virus assays is a key finding at this stage of the pandemic. With the increased transmissibility and

potential for more severe disease observed with these viruses, the availability of therapeutic or

prophylactic mAbs that remains active against variants is essential. In addition to retaining activity

against circulating WHO VOCs and VOIs, in pseudotyped virus experiments VIR-7831 showed no

significant cross-resistance with variants that reduce the activity of other currently authorized or approved

mAbs. These data additionally demonstrate the uniqueness of VIR-7831 and VIR-7832 and further

highlight the utility VIR-7831 and VIR-7832 have, alone or in combination, as clinical agents.

Notably, even at this stage of the pandemic, the VIR-7831/VIR-7832 epitope remains highly conserved

among available sequences of circulating virus with ≥99.66% conservation of epitope amino acids. This is

consistent with the value of the strategy used for isolation of monoclonal antibodies that neutralize both

SARS-CoV and SARS-CoV-2 based on the idea that these two virulent human viruses are

phylogenetically divergent within the sarbecovirus subgenus. Furthermore, MARMs identified at

positions P337 and E340 are present at very low levels among current sequences. That amino acids P337

and E340 remain ≥99.99% conserved at this stage of the pandemic indicates that variants at these

positions may confer disadvantageous effects on the virus, consistent with the conservation of this epitope

across the sarbecovirus family[22].

Viral variants of concern for RBM-targeting mAbs are quickly spreading[58]. The vaccines presently being

deployed around the world generate high-titer neutralizing antibodies that target the S protein RBM.

Importantly, the RBM is highly immunodominant for responses to natural infection[59]. Vaccine-induced

and convalescent immunity may therefore potentially put further mutational pressure on the RBM

sequence to evade such antibody responses. In contrast, antibody responses overlapping with the VIR-

7831/VIR-7832 epitope are limited after infection [59], possibly because of the shielding effect of the highly

conserved N343 glycan. In this regard the epitope may face less vaccine- or infection-generated immune

pressure, potentially preserving this conserved epitope long-term.

Recent data have indicated that the cells used to generate live virus stocks and overexpression of

ACE2/TMPRSS2 in target cells used for assays can affect mAb activity in vitro[30,60]. The VIR-7831/VIR-

7832 parental antibody S309 seems particularly sensitive to in vitro methods using ACE2 overexpressing

cells[60]. It is therefore notable that VIR-7831 displays significant efficacy in an in vivo proof-of-concept

SARS-CoV-2 infection experiment using hamsters despite the fact that patterns of engagement of hamster

FcRs by human IgG1 antibodies may not reflect patterns of human IgG1 antibodies with their cognate

human FcRs. These findings argue that in vitro data derived from such ACE2 and/or TMPRSS2

overexpression cell lines do not accurately reflect the in vivo antiviral capacity of tested mAbs.

Furthermore, that the significant in vivo effects of VIR-7831 in the hamster model likely occurred in the

absence of full effector functions due to species-specific interactions between antibodies and FcRs, argues

that effects in COVID-19 patients incorporating both the neutralization capacity of the antibody plus the

ability to harness the strength of the immune system could lead to positive clinical outcomes.

The clinical potential of VIR-7832, with the inclusion of the GAALIE Fc mutation, is of special interest

in the context of SARS-CoV-2 infection. Previously published data by the Ravetch laboratory comparing

the in vivo efficacy of a hemagglutinin-targeting mAb with and without inclusion of the GAALIE

mutation in a transgenic humanized FcγR mouse model of influenza infection demonstrated superior

efficacy of the GAALIE-containing antibody in both therapeutic and prophylactic experiments[27]. These

effects were mediated by protective CD8$^+$ T cell responses elicited by the GAALIE antibody. Clinical

data examining the contribution of the adaptive immune response in SARS-CoV-2 infection indicate that

poor T cell induction correlates with severe disease (reviewed in [61]). Thus, the potential for VIR-7832 to

augment the T cell response to SARS-CoV-2 infection could conceivably play a crucial role in limiting

progression to severe COVID-19 disease or in treatment of severe established disease. This latter

possibility is supported by recent publications showing that monoclonal antibodies with effector functions

are especially effective in the therapeutic setting via recruitment of tissue-protective monocyte functions

[20], and that potency of antibodies in the pre-clinical mouse model does not correlate with in vitro

neutralizing activity of antibodies[19].

Taken together, these data indicate that VIR-7831 and VIR-7832 canplay a powerful role in the fight

against COVID-19 through the dual action of broadly neutralizing activity paired with engagement of the

immune system through effector function capabilities.

METHODS

Cells. Vero E6 cells (ATCC) and Lenti-X 293T cells (Takara) were cultured in Dulbecco's Modified

Eagle's medium (DMEM), 10% FBS, 1x Penicillin-Streptomycin at 37°C, 5% CO_2.

Monoclonal Antibodies. VIR-7831 and VIR-7832 were produced at WuXi Biologics (China). SGHmAb-no-LS, S309-LS, and S309-GRLR were produced at Humabs Biomed SA, a subsidiary of Vir Biotechnology (Bellinzona, Switzerland) in expiCHO cells transiently co-transfected with plasmids expressing the heavy and light chain, as previously described [62]

Virus. SARS-CoV-2 isolates USA-WA1/2020, UK/VUI/3/2020, hCoV-19/South Africa/KRISP-K005325/2020, hCoV-19/Japan/TY7-503/2021, hCoV-19/USA/CA-SU-15-S02/2021 and hCoV-19/USA/PHC658/2021 were obtained from BEI Resources. To propagate SARS-CoV-2, VeroE6 or VeroE6-TMPRSS2 cells were seeded at 10×10^6 cells in T175 flasks in growth media and infected the next day at a MOI of 0.001 in virus propagation media. Virus was adsorbed for 1 hour at 37°C. Virus inoculum was removed, flasks were washed once with PBS, 25 mL of infection media was added to the cells and flasks were incubated at 37°C. Supernatants were collected at 48 hours post-infection once cytopathic effect was visible, centrifuged at 500 x g for 5 minutes, followed by a second centrifugation at 1000 x g for 5 minutes. Clarified supernatants were then aliquoted and stored at -80°C. Virus titers were determined using a plaque assay on VeroE6 cells, using standard methods. Briefly, 10-fold dilutions of virus stock were incubated in 6 well plates with 2.4% colloidal cellulose overlay for 24 hours. Cells were fixed with 4% PFA for 30 minutes at room temperature (RT), permeabilized with 0.125% Triton X-100, stained with anti-SARS-CoV-2 nucleocapsid antibody at 1:5000 and goat anti-rabbit IgG HRP at 1:5000. Plaque forming units (PFU) were visualized with TrueBlue reagent.

In vitro binding ELISA. For the ELISA assay, 96-well plates were coated with 100 μl/well recombinant SARS-CoV2 RBD diluted in assay diluent (1% BSA/PBS) at a final concentration of 2 μg/mL and incubated overnight at 4°C. Plates were washed three times with 300 μl/well wash buffer using an automated washer. Assay diluent (100 μl/well) was added to block the plates and incubated for 1 hour at room temperature (RT) with shaking. Assay diluent was removed, and plates washed three times with wash buffer. Serial 1:3 dilutions of mAb (concentration range from 6 μg/mL to 0.33 ng/mL) in assay diluent were dispensed at 100 μl/well and incubated 1 hour at RT with shaking, then washed three times

with wash buffer. The HRP-conjugated secondary antibody reagent (1:5,000 dilution in assay diluent)

was added to each well (100 µl/well) and incubated for 1 hour at RT with shaking. After three washes

with wash buffer, 100 µl/well of 2-component TMB peroxidase substrate solution was dispensed in each

well and developed for 5 minutes at RT. The reaction was stopped with 100 µL/well 1M H_2SO_4 and the

OD was read immediately at 450 nm on a SpectraMax M5 Microplate reader. EC_{50} values were calculated

using non-linear regression of log (agonist) versus response in Graph Pad Prism.

Spike binding affinity quantification by SPR. Antibody was diluted to 2 µg/mL (1 mL) in HBS-EP+

buffer and injected at 10 µL/min for 30 seconds across one flow cell of a CM5 sensor chip immobilized

with anti-human Fc antibody docked in a Biacore T200. SARS-CoV2-RBD diluted in HBS-EP+ buffer

was then injected at a single concentration, 1:3 dilutions from 100 nM to 3.7 nM, across both the flow cell

containing captured the antibody as well as a reference flow cell containing only anti-human Fc antibody.

Binding was measured with a flow rate of 30 µL/min and an injection time of 600 seconds; dissociation

was monitored for 1800 seconds after injection. Data were collected at 10 Hz. After each binding

measurement, regeneration reagent was injected to prepare the surface for a new cycle. Experiments were

performed at 25°C, with the samples held at 15 °C in the instrument prior to injection.

Measurement of Binding to Human Fcγ Receptors by SPR. Binding of VIR-7831 and VIR-7832 to

human recombinant FcγRs was measured by surface plasmon resonance (SPR) on a Biacore T200.

Briefly, Biotin CAPture Reagent (modified streptavidin) was injected across all flow cells of a CAP

sensor chip docked in a Biacore T200. Biotinylated Fc receptors at 1 µg/mL were injected across a single

flow cell at 10 µL/min for 60 seconds (one receptor per flow cell), with one flow cell reserved as a

reference surface. VIR 7831 or VIR-7832 at 100 µg/mL (diluted in HBS-EP+) were injected across all

flow cells for 200 seconds using a flow rate of 30 µL/min and association was monitored. Dissociation

was monitored for another 200 seconds after injection. Data was collected at 10 Hz. After each binding

measurement, CAP Regeneration reagent was injected to prepare the surface for a new cycle.

Experiments were performed at 25°C, with the samples held at 15°C in the instrument prior to injection.

Measurement of Binding to Human Complement Protein C1q. Binding of VIR-7831 and VIR-7832 to

human complement was measured by biolayer interferometry (BLI) using an Octet Red96 instrument

(FortéBio). Briefly, anti-human Fab (CH1-specific) sensors were used to capture VIR-7831 and VIR-

7832 at 10 µg/ml for 10 minutes. The IgG-loaded sensors were then exposed to kinetics buffer containing

3 µg/ml of purified human C1q for 4 minutes, followed by a dissociation step in the same buffer for

additional 4 minutes. Association and dissociation profiles were measured in real time as changes in the

interference pattern.

Binding to Cell Surface Expressed SARS-CoV-2 Spike Protein. The SARS-CoV-2 spike protein

coding sequence (YP_009724390.1, Wuhan-Hu-1 strain) was cloned into a cell expression plasmid under

the control of the human CMV promoter (phCMV1) to generate phCMV1 WT spike. ExpiCHO-S cells

were seeded the day before transfection at 3×10^6 cells/mL in ExpiCHO Expression Medium.

Immediately before transfection, the cells were seeded at 6×10^6 cells cells/mL in a volume of 15 mL in

125 mL shake flasks. Six µg of phCMV1 WT spike plasmid or vector control were diluted in 1.2 mL of

iced OptiPRO SFM., followed by addition of 48 µL of ExpiFectamine CHO Reagent and complexing for

1 minute at RT. The transfection mixture was added dropwise to cells with gentle swirling. Cells were

then incubated at 37°C, 8% CO2 with shaking for 42 hours. At 42 hours post-transfection, ExpiCHO-S

cells were harvested, washed twice with FACS buffer and resuspended at a concentration of 1.0×10^6

cell/mL in PBS. Cells (5×10^4 cells in 50 µl/wells) were dispensed into a 96-well V-bottom plate.

Antibody was serially diluted (1:4, 10 points) starting at a concentration of 10 µg/mL. Cells were pelleted

at 300 x g for 5 minutes and resuspended in 50 µL/well of antibody serial dilutions and plates were

incubated for 45 mins on ice. Cells were washed twice in FACS buffer. Alexa Fluor 647-labelled Goat

Anti-Human IgG secondary Ab was diluted 1:750 in FACS buffer and 50 µL was added to the cell pellet

for 15 min on ice. Cells were washed twice with FACS buffer, resuspended in 1% PFA. Data was

acquired by flow cytometry (CytoFlex LX).

Pseudotyped virus production. Lenti-X™ 293T cells were seeded in 10-cm dishes for 80% next day

confluency. The next day, cells were transfected with the plasmid pcDNA3.1(+)-spike-D19 (encoding the

SARS-CoV-2 spike protein) or pcDNA3.1(+)-spike-D19 variants using the transfection reagent TransIT-

Lenti according to the manufacturer's instructions. One day post-transfection, cells were infected with

VSV-luc (rVSVΔG; Kerafast) at an MOI of 3. The cell supernatant containing SARS-CoV-2 pseudotyped

virus was collected at day 2 post-transfection, centrifuged at 1000 x g for 5 minutes to remove cellular

debris, aliquoted and frozen at -80°C.

In Vitro Neutralization of SARS-CoV-2 Pseudotyped Virus. VeroE6 cells were seeded into flat

bottom tissue culture 96-well plates at 20,000 cells/well and cultured overnight at 37°C. Twenty-four

hours later, 9-point 1:4 serial dilutions of VIR-7831 were prepared in infection medium and each dilution

was tested in triplicate per plate (range: 20,000 to 0.3 ng/mL final concentration). SARS-CoV-2 virus

stock was diluted in infection media for a final concentration of 2000 plaque forming units per well (MOI

0.1). Antibody dilutions were added to virus and incubated for 30 minutes at 37°C. Media was removed

from the VeroE6 cells, mAb-virus complexes were added, and cells were incubated at 37°C. At 6 hours

post-infection, cells were fixed with 250 μL 4% PFA, incubated for 30 minutes at RT, then washed 3

times with PBS to remove residual PFA. The cells were permeabilized with 50 μL of 0.125% Triton X-

100 in PBS for 30 minutes at RT. The blocking buffer was removed, 50 μL of SARS-CoV-2 nucleocapsid

antibody at 1:2,000 in blocking buffer was added, and plate was incubated for 1 hour at RT. Plates were

washed three times with PBS and then incubated for 1 hour at RT with 50 μL/well of goat anti-rabbit-

Alexa647 secondary antibody at a final dilution of 1:1,000 mixed with 2 ug/mL Hoechst dye in blocking

buffer. After washing 5 times with PBS, 100 μL of fresh PBS was added for imaging. Plates were imaged

on a Cytation5 plate reader. Whole well images were acquired (12 images at 4X magnification per well)

and nucleocapsid-positive cells were counted using the manufacturer's software.

Live virus neutralization. VeroE6 cells were seeded into flat bottom tissue culture 96-well plates at

20,000 cells/well and cultured overnight at 37°C. Twenty-four hours later, 9-point 1:4 serial dilutions of

VIR-7831 were prepared in infection medium and each dilution was tested in triplicate per plate (range:

20,000 to 0.3 ng/mL final concentration). SARS-CoV-2 virus stock was diluted in infection media for a

final concentration of 2000 plaque forming units per well (MOI 0.1). Antibody dilutions were added to

virus and incubated for 30 minutes at 37°C. Media was removed from the VeroE6 cells, mAb-virus

complexes were added, and cells were incubated at 37°C. At 6 hours post-infection, cells were fixed with

250 µL 4% PFA, incubated for 30 minutes at RT, then washed 3 times with PBS to remove residual PFA.

The cells were permeabilized with 50 µL of 0.125% Triton X-100 in PBS for 30 minutes at RT. The

blocking buffer was removed, 50 µL of SARS-CoV-2 nucleocapsid antibody at 1:2,000 in blocking buffer

was added, and plate was incubated for 1 hour at RT. Plates were washed three times with PBS and then

incubated for 1 hour at RT with 50 µL/well of goat anti-rabbit-Alexa647 secondary antibody at a final

dilution of 1:1,000 mixed with 2 ug/mL Hoechst dye in blocking buffer. After washing 5 times with PBS,

100 µL of fresh PBS was added for imaging. Plates were imaged on a Cytation5 plate reader. Whole well

images were acquired (12 images at 4X magnification per well) and nucleocapsid-positive cells were

counted using the manufacturer's software..

Determination of Viral Titer by Focus-Forming Assay. One day prior to infection, $1.2X10^4$ VeroE6

cells were plated in black-walled, clear bottomed 96-well plates. Virus samples were diluted 1:5 in

infection media and adsorbed onto VeroE6 cells for one hour at 37°C. The cells were washed once and

overlaid with 1% methylcellulose/serum-containing media. At 24 hours post-infection, the

methylcellulose overlay was removed, and cells were washed with PBS. Cells were fixed with 4% PFA,

incubated for 30 minutes at RT, then washed with PBS to remove residual PFA. The cells were

permeabilized with 50 µL of 0.25% Triton X-100 in PBS for 30 minutes at RT. The Triton X-100 was

removed, cells were washed twice with PBS, and incubated with 50 µL of SARS-CoV-2 nucleocapsid

antibody at 1:2,000 in blocking buffer for one hour at RT. Plates were washed three times with PBS and

then incubated for one hour at RT with 50 µL/well of goat anti-rabbit-Alexa647 secondary antibody at

1:1,000 in blocking buffer. After washing three times with PBS, 50 µL of Hoechst dye at 1:1,000 in PBS

was added for imaging. Plates were imaged on a Cytation5 plate reader. Whole well images were

acquired (12 images at 4X magnification per well) and nucleocapsid-positive foci were counted using the

manufacturer's software and used to determine focus-forming units/mL supernatant (FFU/mL).

Determination of mAb-Dependent Activation of Human FcγRIIa, FcγRIIIa or FcγRIIb. VIR-7831,

VIR-7832, S309-LS, and a control mAb with abrogated FcγR binding, S309-GRLR, were serially diluted

6-fold in assay buffer from 10,000 ng/ml to 0.006 ng/ml. Nine-point serial dilutions of mAbs were

incubated with 12,500 (for FcγRIIIa and FcγRIIb) or 10,000 (for FcγRIIa) CHO-CoV-2-Spike cells per

96-plate well in a white, flat-bottom plate for 15 minutes at room temperature. Jurkat effector cells

expressing indicated FcγRs and stably transfected with an NFAT-driven luciferase gene were thawed,

diluted in assay buffer, and added to the plate at an effector to target cell ratio of 6:1 for FcRγIIIa and

FcγRIIb or 5:1 for FcγIIa. Control wells were also included that were used to measure antibody-

independent activation (containing target cells and effector cells but no antibody) and background

luminescence of the plate (wells containing assay buffer only). Plates were incubated for 18 hours at 37°C

with 5% CO2. Activation of human FcγRs in this bioassay results in the NFAT-mediated expression of

the luciferase reporter gene. Luminescence was measured with a luminometer after adding the Bio

GloTM Luciferase Assay Reagent according to the manufacturer's instructions. To control for

background, the mean of the relative luminescence units (RLU) values in wells containing only Assay

Buffer was calculated and subtracted from all data points. Data were expressed as the average of RLUs

over the background

Determination of NK-Cell Mediated Antibody-Dependent Cellular Cytotoxicity. Primary NK cell

activation was tested using freshly isolated cells from two previously genotyped donors expressing

homozygous low affinity (F158) or high affinity (V158) FcγRIIIa. Serial dilutions of mAbs (serially

diluted 10-fold in AIM-V Medium from 40,000 ng/ml to 0.075 ng/ml) were incubated with 7,500 CHO-

CoV-2 Spike cells per well of a 96 well round-bottom plate for 10 minutes. Target cell and antibody

mixtures were then incubated with primary human NK cells as effectors at an effector-to-target ratio of

10:1. ADCC was measured using lactate dehydrogenase (LDH) release as a readout according to the

manufacturer's instructions (Cytotoxicity Detection Kit (LDH), Roche) after 4 hours of incubation at

37°C. In brief, plates were centrifuged for 4 minutes at 400 x g, and 35 µl of supernatant was transferred

to a flat 384 well plate. LDH reagent was prepared and 35 µl were added to each well. Using a kinetic

protocol, the absorbance at 490 nm and 650 nm was measured once every 2 minutes for 8 minutes, and

the slope of the kinetics curve was used as result. The percent specific lysis was determined by applying

the following formula: (specific release – spontaneous release) / (maximum release - spontaneous release)

x 100.

Determination of Monocyte-Mediated Antibody-Dependent Cellular Phagocytosis. ADCP assays

were performed using human PBMCs freshly isolated from whole blood. CHO CoV-2-Spike cells were

used as target cells and were fluorescently labeled with PKH67 Fluorescent Cell Linker Kit (Sigma

Aldrich) prior to incubation with mAbs, according to manufacturer's instructions. Serial dilutions of

mAbs (serially diluted 5-fold from 5,000 ng/ml to 0.32 ng/ml in RPMI-1640 + L-glutamine supplemented

with 10% Hyclone FBS + 2x anti-anti (antibiotic-antimycotic)) were incubated with 10,000 CHO-CoV-2-

Spike cells per well of a 96 well polypropylene plate for 10 minutes. Primary PBMCs were fluorescently

labeled with Cell Trace Violet according to the manufacturer's instructions. Target cell and antibody

mixtures were then incubated with labeled PBMCs at an effector-to-target ratio of 16:1. After an

overnight incubation at 37°C, monocytes were stained with anti-human CD14-APC antibody (BD

Pharmingen). Antibody-mediated phagocytosis was determined by flow cytometry, gating on CD14+

cells that were double positive for cell trace violet and PKH67. Raw data were exported from the flow

cytometer into the flow cytometry analysis software FlowJo v10 (Becton Dickinson) for gating and

determination of the percentage of $CD14^+$ cells that were also double positive for cell trace violet and

PKH67. Cells expressing only cell trace violet or only PKH67 were used to set the positive staining gates.

In vitro resistance selection. The selection of variants in the presence of increasing concentrations of

VIR-7832 was conducted in VeroE6 cells. The day before infection, 6 x 10^4 VeroE6 cells were seeded in

24 well plates and incubated overnight at 37°C. The next day, 600 focus forming units (FFU) of SARS-

CoV-2 virus (MOI = 0.01) was incubated with 0.5X IC_{50} of VIR-7832 (0.05 μg/mL) at 37°C for one hour

in infection media. The mAb-virus complexes were adsorbed on VeroE6 cells for one hour at 37°C in

duplicate wells. After adsorption, cells were washed with DMEM and overlaid with infection media

containing 0.05 μg/mL VIR-7832. Control wells infected without antibody were included with each

passage. Infected cells were monitored visually for CPE daily. In general, when infected cells exhibited ≥

50% CPE, the culture supernatants were harvested, diluted 1:200, and added to fresh VeroE6 cells in 24-

well plates with equivalent or increasing concentrations of VIR-7832. At each passage, supernatant was

aliquoted and frozen at -80°C for titer and neutralization analyses.

In vitro assessment of potential for ADE. VeroE6 cells were plated at $1.25X10^4$ cells/well one day prior

to infection. For each independent experiment, moDCs and PBMCs from five unique moDC donors and

six unique PBMC donors were used, with three unique donors used for each independent experiment.

Cryopreserved monocytes from unique donors were differentiated into moDCs for six days using human

moDC differentiation media according to the manufacturer's protocol. Cryopreserved PBMCs from

unique donors are thawed in the presence 0.3 mg/mL DNase and cultured in media for one day prior to

infection. On the day of infection, moDCs, PBMCs, and U937 cells were counted and plated at $7.5X10^4$

cells/well.

To examine viral entry, 24 hours post-infection, cells were fixed with 4% PFA, incubated for 30 minutes

at RT, then washed with PBS to remove residual PFA. The cells were permeabilized with 50 μL of 0.25%

Triton X-100 in PBS for 30 minutes at RT. The Triton X-100 was removed, cells were washed twice with

PBS, and incubated with 50 μL of SARS-CoV-2 nucleocapsid antibody at 1:2,000 in blocking buffer for

one hour at RT. Plates were washed three times with PBS and then incubated for one hour at RT with 50

μL/well of goat anti-rabbit-Alexa647 secondary antibody at 1:1,000 in blocking buffer. After washing

three times with PBS, 50 μL of Hoechst dye at 1:1,000 in PBS was added for imaging. Plates were

imaged on a Cytation5 plate reader. Whole well images were acquired (12 images at 4X magnification

per well) and nucleocapsid-positive cells were counted using the manufacturer's software. The percent of

nucleocapsid+ cells was quantified using the Gen5 Imager software (Biotek, Vermont) as number of

Cy5+ cells, [(nucleocapsid+ cells)/number of Hoechst+ cells (total cells)]x100. Data was analyzed using

Prism v8.00 (GraphPad Software, La Jolla California USA, www.graphpad.com).

In order to quantify chemokines and cytokines from supernatants in a BSL2 laboratory, supernatants were

inactivated by 10 minutes exposure to UVC light at 5,000 µJ/cm2. Supernatants were diluted 1:5 in

infection media and levels of cytokines/chemokines were quantified using the U-plex 96-well assay

according to the manufacturer's protocol (Meso Scale Diagnostics, Rockville, MD). Quantification of

cytokines and chemokines were determined based on an 8-point standard curve in duplicate, provided by

the manufacturer. Cytokine data was analyzed using the Discovery Workbench v4.0.13 software (Meso

Scale Diagnostics). Data was graphed and statistical analyses were conducted using Prism software.

Sequencing of SARS-CoV-2 Spike Gene. To isolate nucleic acid from the supernatant of viral passages,

120 µL of cell supernatant was added to 360 µL of Trizol and stored at -80°C for further analysis. Trizol

collected samples from viral passages where a shift in neutralization > 2-fold relative to wild type was

detected were subjected to RNA isolation using PureLink RNA Mini Kit with the incorporation of on-

column PureLink DNase Treatment, following manufacturer's instructions. Reverse transcription

reactions were performed with 6 µL of purified RNA and oligoT primers using the NEB ProtoScript II

First Strand cDNA Synthesis kit, according to manufacturer's instructions. The resulting cDNA was used

as a template for PCR amplification of the spike gene using the KapaBiosystems polymerase (KAPA HiFi

HotStart ReadyMix) with primers 5' aattatcttggcaaaccacg-3' and 5' tgaggcttgtatcggtatcg-3'.

Amplification conditions included an initial 3 minutes at 95°C, followed by 28 cycles with 20 seconds at

98°C, 15 seconds at 62°C and 72°C for 2 minutes, with a final 4 minutes at 72°C. PCR products were

purified using AMPure XP beads following manufacturer's instructions. The size of the amplicon was

confirmed by analyzing 2 µL of PCR products using the Agilent D5000 ScreenTape System. Products

were quantified by analyzing 1 µL with the Quant-iT dsDNA High-Sensitivity Assay Kit. Twenty ng of

purified PCR product was used as input for library construction using the NEBNext Ultra II FS DNA

Library Prep kit following manufacturer's instructions. DNA fragmentation was performed for 13

minutes. NEBNext Multiplex Oligos for Illumina Dual Index Primer Set 1 was used for library

construction, with a total of 6 PCR cycles. Libraries size was determined using the Agilent D1000

ScreenTape System and quantified with the Quant iT dsDNA High-Sensitivity Assay Kit. Equal amounts

of each library were pooled together for multiplexing and 'Protocol A: Standard Normalization Method'

of the Illumina library preparation guide was used to prepare 8 pM final multiplexed libraries with 1%

PhiX spike-in for sequencing. The MiSeq Reagent Kit v3 (600-cycle) was used for sequencing the

libraries on the Illumina MiSeq platform, with 300 cycles for Read 1, 300 cycles for Read 2, 8 cycles for

Index 1, and 8 cycles for Index 2.

Bioinformatics Analysis of Conservation. Available genome sequences for SARS-CoV-2 were

downloaded from Global Initiative on Sharing All Influenza Data (GISAID; https://www.gisaid.org/) on

June 4, 2021. Bat and pangolin sequences were removed to yield human-only sequences. The spike open

reading frame was localized by aligning the reference protein sequence (NCBI reference sequence:

YP_009724390.1) to the genomic sequence of isolates with Exonerate v.2.4.0. Coding nucleotide

sequences were translated in silico using seqkit v.0.12.0. Multiple sequence alignment was performed

using MAFFT v.7.455. Variants were determined by comparison of aligned sequences to the reference

sequence using the R v3.6.3/Bioconductor v.3.10 package Biostrings v.2.54.0.

In vivo studies. Syrian golden hamster studies were conducted at Lovelace Biomedical (Albuquerque,

NM). Twelve- to sixteen-week-old male hamsters were interperitoneally administered a non-LS version

of VIR-7831 (SGHmAb-no-LS), control antibody or diluent Day -1 or Day -2 prior to virus challenge.

Animals were inoculated intranasally at Day 0 with 7.4×10^4 $TCID_{50}$ with SARS-CoV-2 (isolate USA-

WA1/2020). Animals were also weighed once daily in the morning beginning on study Day -10 and

continuing until the end of the study. Following euthanasia, RT-qPCR was performed on lung

homogenates using quantitative real-time PCR methods targeting the SARS-CoV-2 N gene and the

median tissue culture infections dose (TCID$_{50}$) was determined per Lovelace internal methodology.

Author Contributions

Conceived studies: A.L.C, C.H-D., F.A.P, D.M., M.S., L.S., A.T., L.A.P., S.H., G.S., H.W.V., D.C., C.M.H.

Designed studies and experiments: A.L.C, C.H.D., F.A.P, D.M., M.S., M.L.A., E. D., B.G., J.D., L.R.,A.C., A.S.,

R.S., J.W., N.C., E.C., S.L., C.C., D.P., C.S., J.N., A.P., A.W., L.S., A.T., L.A.P., S.H., G.S., H.W.V., D.C., C.M.H.

Performed experiments: D.M., M.S., M.L.A., B.G., J.D., E.D., A.S., L.R., H.T., J.D., S.S., D.P., C.S., J.N., B.S.,

S.B., J.W., J.Z., H.K., A.C., M.M-R., A.P., A.W., N.C., E.C. Analyzed and interpreted data: A.L.C., C.H-D., F.A.L.,

D.M., M.S., M.L.A., B.G., J.D., D.P., C.S., J.N., E.L., A.S., R.S., L.R., H.T., B.S., S.B., J.W., J.Z., H.K., A.C., M. M-

R., N.C., E.C., S.L., A.W., C.C., L.S., A.T., S.H., G.S., H.W.G, D.C., C.M.H. Prepared the manuscript with input

from all authors: A.L.C., G.S., A.T., L.P., D.C., H.W.G., C.M.H.

Competing interests

629   Some authors are current or former employees of Vir Biotechnology or Humabs BioMed SA (a fully-

630   owned subsidiary of Vir Biotechnology) and may hold shares in Vir Biotechnology. H.W.V. is a founder of

631   PierianDx and Casma Therapeutics. This research was sponsored and funded by Vir Biotechnology in

632   collaboration with GSK.

REFERENCES

1. WHO Coronavirus Disease (COVID-19) Dashboard | WHO Coronavirus Disease (COVID-19) Dashboard. https://covid19.who.int/.

2. Levin, A. T. *et al.* Assessing the age specificity of infection fatality rates for COVID-19: systematic review, meta-analysis, and public policy implications. *Eur J Epidemiol* **35**, 1123–1138 (2020).

3. Dennis, A. *et al.* Multi-organ impairment in low-risk individuals with long COVID. doi:10.1101/2020.10.14.20212555.

4. Murphy, J. *et al.* Psychological characteristics associated with COVID-19 vaccine hesitancy and resistance in Ireland and the United Kingdom. *Nat Commun* **12**, 29 (2021).

5. Khubchandani, J. *et al.* COVID-19 Vaccination Hesitancy in the United States: A Rapid National Assessment. *J Commun Health* 1–8 (2021) doi:10.1007/s10900-020-00958-x.

6. Sallam, M. *et al.* High Rates of COVID-19 Vaccine Hesitancy and Its Association with Conspiracy Beliefs: A Study in Jordan and Kuwait among Other Arab Countries. *Nato Adv Sci Inst Se* **9**, 42 (2021).

7. Tegally, H. *et al.* Emergence and rapid spread of a new severe acute respiratory syndrome-related coronavirus 2 (SARS-CoV-2) lineage with multiple spike mutations in South Africa. *medRxiv* (2020) doi:10.1101/2020.12.21.20248640.

8. Naveca, F. *et al.* Phylogenetic relationship of SARS-CoV-2 sequences from Amazonas with emerging Brazilian variants harboring mutations E484K and N501Y in the Spike protein - SARS-CoV-2 coronavirus / nCoV-2019 Genomic Epidemiology - Virological. https://virological.org/t/phylogenetic-relationship-of-sars-cov-2-sequences-from-amazonas-with-emerging-brazilian-variants-harboring-mutations-e484k-and-n501y-in-the-spike-protein/585 (2011).

9. Rambaut, A. *et al.* Preliminary genomic characterisation of an emergent SARS-CoV-2 lineage in the UK defined by a novel set of spike mutations - SARS-CoV-2 coronavirus / nCoV-2019 Genomic Epidemiology - Virological. https://virological.org/t/preliminary-genomic-characterisation-of-an-emergent-sars-cov-2-lineage-in-the-uk-defined-by-a-novel-set-of-spike-mutations/563 (2020).

10. Bamlanivimab EUA Letter of Authorization Reissue 02092021. https://www.fda.gov/media/143603/download.

11. Casirivimab and Imdevimab EUA Fact Sheet for Healthcare Providers. https://www.fda.gov/media/143892/download.

12. Lilly's neutralizing antibody bamlanivimab (LY-CoV555) prevented COVID-19 at nursing
homes in the BLAZE-2 trial, reducing risk by up to 80 percent for residents.
https://investor.lilly.com/node/44291/pdf.

13. Gottlieb, R. L. *et al.* Effect of Bamlanivimab as Monotherapy or in Combination With
Etesevimab on Viral Load in Patients With Mild to Moderate COVID-19. *Jama* **325**, 632–644
(2021).

14. Chen, P. *et al.* SARS-CoV-2 Neutralizing Antibody LY-CoV555 in Outpatients with Covid-
19. *New Engl J Med* (2020) doi:10.1056/nejmoa2029849.

15. Weinreich, D. M. *et al.* REGN-COV2, a Neutralizing Antibody Cocktail, in Outpatients with
Covid-19. *New Engl J Med* **384**, 238–251 (2020).

16. Gupta, A. *et al.* Early Treatment for Covid-19 with SARS-CoV-2 Neutralizing Antibody
Sotrovimab. *New Engl J Med* **385**, 1941–1950 (2021).

17. Starr, T. N. *et al.* Prospective mapping of viral mutations that escape antibodies used to treat
COVID-19. *Science* **371**, 850–854 (2021).

18. Wang, P. *et al.* Increased Resistance of SARS-CoV-2 Variants B.1.351 and B.1.1.7 to
Antibody Neutralization. *Biorxiv* 2021.01.25.428137 (2021) doi:10.1101/2021.01.25.428137.

19. Schäfer, A. *et al.* Antibody potency, effector function, and combinations in protection and
therapy for SARS-CoV-2 infection in vivo. *J Exp Med* **218**, (2020).

20. Winkler, E. S. *et al.* Human neutralizing antibodies against SARS-CoV-2 require intact Fc
effector functions and monocytes for optimal therapeutic protection. *Biorxiv* 2020.12.28.424554
(2020) doi:10.1101/2020.12.28.424554.

21. Bournazos, S. & Ravetch, J. V. Fcγ Receptor Function and the Design of Vaccination
Strategies. *Immunity* **47**, 224–233 (2017).

22. Pinto, D. *et al.* Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV
antibody. *Nature* **583**, 290–295 (2020).

23. Pinto, D. *et al.* Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV
antibody. *Nature* **583**, 290–295 (2020).

24. Ko, S.-Y. *et al.* Enhanced neonatal Fc receptor function improves protection against primate
SHIV infection. *Nature* **514**, 642 (2014).

25. Zalevsky, J. *et al.* Enhanced antibody half-life improves in vivo activity. *Nat Biotechnol* **28**,
157 (2010).

26. Gaudinski, M. R. *et al.* Safety and pharmacokinetics of the Fc-modified HIV-1 human
monoclonal antibody VRC01LS: A Phase 1 open-label clinical trial in healthy adults. *Plos Med*
15, e1002493 (2018).

27. Bournazos, S., Corti, D., Virgin, H. W. & Ravetch, J. V. Fc-optimized antibodies elicit CD8
immunity to viral respiratory infection. *Nature* 1–9 (2020) doi:10.1038/s41586-020-2838-z.

28. Weitzenfeld, P., Bournazos, S. & Ravetch, J. V. Antibodies targeting sialyl Lewis A mediate
tumor clearance through distinct effector pathways. *J Clin Invest* **129**, 3952–3962 (2019).

29. Bournazos, S., Corti, D., Virgin, H. W. & Ravetch, J. V. Fc-optimized antibodies elicit CD8
immunity to viral respiratory infection. *Nature* 1–9 (2020) doi:10.1038/s41586-020-2838-z.

30. Diamond, M. *et al.* SARS-CoV-2 variants show resistance to neutralization by many
monoclonal and serum-derived polyclonal antibodies. *Res Square* (2021) doi:10.21203/rs.3.rs-
228079/v1.

31. GISAID. https://www.epicov.org/epi3.

32. Stanford Covid Database. https://covdb.stanford.edu/page/susceptibility-data.

33. NCATS Database. https://opendata.ncats.nih.gov/variant/activity.

34. Kallewaard, N. L. *et al.* Structure and Function Analysis of an Antibody Recognizing All
Influenza A Subtypes. *Cell* **166**, 596–608 (2016).

35. DiLillo, D. J., Tan, G. S., Palese, P. & Ravetch, J. V. Broadly neutralizing hemagglutinin
stalk–specific antibodies require FcγR interactions for protection against influenza virus in vivo.
Nat Med **20**, 143–151 (2014).

36. Dunand, C. J. H. *et al.* Both Neutralizing and Non-Neutralizing Human H7N9 Influenza
Vaccine-Induced Monoclonal Antibodies Confer Protection. *Cell Host Microbe* **19**, 800–813
(2016).

37. Leon, P. E. *et al.* Optimal activation of Fc-mediated effector functions by influenza virus
hemagglutinin antibodies requires two points of contact. *Proc National Acad Sci* **113**, E5944–
E5951 (2016).

38. Bruhns, P. *et al.* Specificity and affinity of human Fcγ receptors and their polymorphic
variants for human IgG subclasses. *Blood* **113**, 3716–3725 (2009).

39. Weitzenfeld, P., Bournazos, S. & Ravetch, J. V. Antibodies targeting sialyl Lewis A mediate
tumor clearance through distinct effector pathways. *J Clin Invest* **129**, 3952–3962 (2019).

40. Cheng, Z. J. *et al.* Development of a robust reporter-based ADCC assay with frozen, thaw-
and-use cells to measure Fc effector function of therapeutic antibodies. *J Immunol Methods* **414**,
69–81 (2014).

41. Arvin, A. M. *et al.* A perspective on potential antibody-dependent enhancement of SARS-
CoV-2. *Nature* 1–11 (2020) doi:10.1038/s41586-020-2538-8.

42. Joyner, M. J. & Wright, R. S. Safety Update: CO VID-19 Convalescent Plasma in
20,000 Hospitalized Patients. *Mayo Clinic Proceedings* (2020).

43. Arvin, A. M. *et al.* A perspective on potential antibody-dependent enhancement of SARS-
CoV-2. *Nature* **584**, 353–363 (2020).

44. Khurana, S. *et al.* Vaccine-induced anti-HA2 antibodies promote virus fusion and enhance
influenza virus respiratory disease. *Sci Transl Med* **5**, 200ra114 (2013).

45. Winarski, K. L. *et al.* Antibody-dependent enhancement of influenza disease promoted by
increase in hemagglutinin stem flexibility and virus fusion kinetics. *Proc National Acad Sci* **116**,
15194–15199 (2019).

46. Hui, K. P. Y. *et al.* Tropism, replication competence, and innate immune responses of the
coronavirus SARS-CoV-2 in human respiratory tract and conjunctiva: an analysis in ex-vivo and
in-vitro cultures. *Lancet Respir Medicine* (2020) doi:10.1016/s2213-2600(20)30193-4.

47. Liu, Z. *et al.* Identification of Common Deletions in the Spike Protein of Severe Acute
Respiratory Syndrome Coronavirus 2. *J Virol* **94**, (2020).

48. Baum, A. *et al.* Antibody cocktail to SARS-CoV-2 spike protein prevents rapid mutational
escape seen with individual antibodies. *Science* **369**, 1014–1018 (2020).

49. Thomson, E. C. *et al.* Circulating SARS-CoV-2 spike N439K variants maintain fitness while
evading antibody-mediated immunity. *Cell* (2021) doi:10.1016/j.cell.2021.01.037.

50. Starr, T. N., Greaney, A. J., Dingens, A. S. & Bloom, J. D. Complete map of SARS-CoV-2
RBD mutations that escape the monoclonal antibody LY-CoV555 and its cocktail with LY-
CoV016. doi:10.1101/2021.02.17.431683.

51. Wise, J. Covid-19: The E484K mutation and the risks it poses. *Bmj* **372**, n359 (2021).

52. Chen, R. E. *et al.* Resistance of SARS-CoV-2 variants to neutralization by monoclonal and
serum-derived polyclonal antibodies. *Nat Med* 1–10 (2021) doi:10.1038/s41591-021-01294-w.

53. VIR-7831 for the Early Treatment of COVID-19 in Outpatients - Full Text View -
ClinicalTrials.gov. https://clinicaltrials.gov/ct2/show/NCT04545060?term=VIR-
7831&draw=2&rank=2.

54. A Study of Immune System Proteins in Participants With Mild to Moderate COVID-19
Illness - Full Text View - ClinicalTrials.gov.
https://clinicaltrials.gov/ct2/show/NCT04634409?term=VIR-7831&draw=2&rank=3.

55. Vir Biotechnology and GSK announce NHS-supported AGILE study to evaluate VIR-7832
in the early treatment of COVID-19 | GSK. https://www.gsk.com/en-gb/media/press-releases/vir-
biotechnology-and-gsk-announce-nhs-supported-agile-study-to-evaluate-vir-7832-in-the-early-
treatment-of-covid-19/.

56. Starr, T. N. *et al.* SARS-CoV-2 RBD antibodies that maximize breadth and resistance to
escape. *Nature* **597**, 97–102 (2021).

57. Starr, T. N. *et al.* SARS-CoV-2 RBD antibodies that maximize breadth and resistance to
escape. *Nature* 1–9 (2021) doi:10.1038/s41586-021-03807-6.

58. CoVariants. https://covariants.org/.

59. Piccoli, L. *et al.* Mapping neutralizing and immunodominant sites on the SARS-CoV-2 spike
receptor-binding domain by structure-guided high-resolution serology. *Cell* (2020)
doi:10.1016/j.cell.2020.09.037.

60. Rappazzo, C. G. *et al.* Broad and potent activity against SARS-like viruses by an engineered
human monoclonal antibody. *Science* **371**, 823–829 (2021).

61. Sette, A. & Crotty, S. Adaptive immunity to SARS-CoV-2 and COVID-19. *Cell* **184**, 861–
880 (2021).

62. Stettler, K. *et al.* Specificity, cross-reactivity and function of antibodies elicited by Zika virus
infection. *Science* **353**, aaf8505 (2016).

FIGURE LEGENDS

Figure 1. VIR-7831 and VIR-7831 bind S and neutralize SARS-CoV-2 virus and S variants of concern in

vitro. a) Binding of VIR-7831 (black circles) and VIR-7832 (blue squares) to SARS-CoV-2 RBD was

tested by ELISA. Shown is the average of four replicates and SD derived from three independent

experiments. b) Association and dissociation profiles of VIR-7831 to SARS-CoV-2-RBD were measured

using SPR. The double reference subtracted curves (shown for single replicates) are plotted together with

the curve fit in black (obscured by close overlay with the data). Values are from two independent

experiments. c) Binding of VIR-7831 (black circles) and VIR-7832 (blue squares) to cell-surface S

protein was determined by flow cytometry. Data are expressed as the percentage of the positive cells.

Results shown are from one experiment and representative of three independent experiments performed.

d) In vitro neutralization of live SARS-CoV-2 by different concentrations of VIR-7831 (left) and VIR-

7832 (right) measured by nucleocapsid staining 6-hours post-infection. Results shown are from one

experiment and representative of at least two independent experiments performed.

Figure 2. VIR-7831 (sotrovimab) retains activity in vitro against individual mutations found in the

Omicron variant. (a) Average fold change in VIR-7831 and VIR-7832 IC_{50} compared to relative wild-type

controls for S variants tested in a VSV/VeroE6 pseudotyped virus system. Data shown are averages of at

least two independent experiments. (b) VIR-7831 neutralization curves from experiments examining

activity against G339D, N440K or R346K pseudotyped virus.

Figure 3. VIR-7831 and VIR-7832 demonstrate effector function in vitro. In vitro effector function (a-e)

activation profiles of human FcγRIIa (a), FcγRIIb (b), FcγRIIIa low-affinity (F158) (c) or FcγRIIIa high-

affinity binding allele (V158) (d) using bioreporter assays using S-expressing CHO cells as the target

antigen. Data points show means± SD of duplicates. NK-cell mediated killing (ADCC) of S-expressing

CHO cells using freshly isolated cells from two donors previously genotyped for homozygous expression

of low-affinity (F/F158) (e) or high-affinity (V/V158) FcγRIIIa (f). Data points are means of

quadruplicates ± SD. g) Antibody-dependent cellular phagocytosis (ADCP) using S-expressing CHO cells

and freshly isolated PBMCs. Data represent the means of duplicates ± SD.

Figure 4. VIR-7831 shows in vivo efficacy in a hamster SARS-CoV-2 model of infection. a) Overview

of hamster in vivo study design. b) and c) Animal weight over time as a percent of starting weight in

animals dosed a Day -1 (b) or Day -2 (c). Medians of at least N=6 animals and interquartile range are

shown. d) and e) Day 4 terminal weights expressed as a percentage of starting weight for animals dosed at

Day -1 (d) or Day -2 (e). Bar denotes median values. f) and g) Day 4 lung viral load in Day -1 (f) or Day -

2 (g) treated animals as assessed by RT-qPCR. Bar denotes median values. h) and i) infectious virus in

lung at Day 4 for Day -1 (h) or Day -2 (i) dosed animals. Bar denotes median values. ns=not significant,

** = p<0.05, and *** = <0.005 as assessed by the Mann-Whitney U-test.

Table 1. VIR-7831 and VIR-7832 retain activity against S variants of concern in an authentic virus

system**.** Average fold change in VIR-7831 and VIR-7832 IC_{50} compared to relative wild-type controls for

S variants tested in an authentic virus system. Data shown are averages of at least two independent

experiments.

Table 2. VIR-7831 and VIR-7832 retain activity against S variants of concern in a pseudotyped virus

system**.** Average fold change in VIR-7831 and VIR-7832 IC_{50} compared to relative wild-type controls for

S variants tested in a VSV/VeroE6 pseudotyped virus system. Data shown are averages of at least two

independent experiments.

Table 3. VIR-7831 and VIR-7832 retain activity against variants that confer resistance to authorized or

approved mAbs. Activity of VIR-7831 against variants conferring reduced susceptibility to

bamlanivimab, imdevimab or casirivimab in a VSV/VeroE6 pseudotyped virus system. The geometric

mean of IC_{50}s and average fold-change versus the relative wild-type control from at least two independent

experiments are shown.

Table 4. The VIR-7831/VIR-7832 epitope is highly conserved. Conservation data comprising >4,500,000

sequences from the GISAID database and variants at each position are shown. Substitutions in green were

tested in a pseudotyped virus assay.

Supplemental Figure 1. Binding of VIR-7831 and VIR-7832 to human FcγRs and C1q as measured by

SPR. Binding of VIR-7831 and VIR-7832 to a) human FcγRIIa (H131 and R131 alleles), FcγRIIIa (F158

and V158 alleles) and FcγRIIb were measured using SPR. Biotinylated purified FcγRs were captured on

the sensor chip surface prior to injection of VIR-7831 or VIR-7832. Association and dissociation profiles

(separated by the vertical dotted line) were measured in real time as change in the SPR signal. b) Binding

of VIR-7831 and VIR-7832 to complement component C1q was measured using BLI on an Octet Red96

instrument. Association and dissociation profiles (separated by the vertical dotted line) were measured in

real time as change in the interference pattern.

Supplemental Figure 2. Sub-neutralizing concentrations of VIR-7831 and VIR-7832 do not enhance

viral entry, viral replication or cytokine production in vitro. Internalization (a) and replication (b) of

SARS-CoV-2 was evaluated in VeroE6, moDCs or PBMCs at various timepoints. Two independent

experiments with human moDCs and PBMCs from three individual donors were analyzed (5 unique

moDC donors, 6 unique PBMC donors total between two experiments). VeroE6 cells were run in

duplicate for both independent experiments. Data from each replicate well from two independent

experiments are plotted as individual points, with horizontal lines representing the median. Mann-

Whitney U-test comparison to no antibody group, *p<0.05. c) Supernatant cytokine and chemokine levels

as measured by MSD at the indicated time post infection. Data from two independent experiments (three

replicates each, five unique donors) are plotted as the mean and SD.

Supplemental Figure 3. Overview of VIR-7832 resistance selection method. All passaging was

conducted in duplicate wells. (a) VIR-7832 concentration was increased during each passage. P3 X

indicates passage 3 virus, after which virus was lost with subsequent increases in concentration. In (b) and

(c), p3X denotes where passage 3 virus from (a) was used to initiate (b) viral lineage 1 and (c) viral

lineage 2. Arrows indicate passages that were subjected to sequence analysis, and * indicate the passages

in lineage 1 with no detectable virus or CPE. Selection continued for a total of eight passages.

Supplemental Table 1. Amino acid substitutions identified in the SARS-CoV-2 S upon in vitro selection

with VIR-7832. Spike gene sequences were compared to a SARS-CoV-2 reference sequence (NCBI:

NC_045512.2) to identify variants. Fold-changes in IC_{50} were determined compared to the SARS-CoV-2

virus stock.

Supplemental Table 2. VIR-7831 and VIR-7832 activity against selected S variants. VIR-7831/VIR-

7832 epitope variants observed by in vitro resistance selection were individually tested in a VSV/VeroE6

pseudotyped virus system. The geometric mean of $IC_{50}s$ and average fold-change versus the relative wild-

type control from at least two independent experiments are shown.

Supplemental Table 3. Activity of VIR-7831 against epitope variants. VIR-7831/VIR-7832 epitope

variants detected in sequences from the GISAID database were tested in a VSV/VeroE6 pseudotyped

virus system. The geometric mean of $IC_{50}s$ and average fold-change versus the relative wild-type control

from at least two independent experiments are shown.

Figure 1

Figure 2

a)

Spike Domain	Substitution	Fold-Change in VIR-7831 IC$_{50}$ Compared to Wild-Type
N-Terminal Domain	del69-70	0.9[a]
	T95I	0.6[b]
	G142D	1.0[c]
Receptor Binding Domain	G339D	1.2
	R346K	0.7
	K417N	0.6[d]
	G446S	1.5(V)[e]
	E484A	0.3(K)[e]
	N440K	0.5
	S477N	2.0
	T478K	1.0[c]
	Q493K	0.9
	N501Y	0.4[f]
S1/S2	D614G	1.0
	H655Y	0.4[g]
	P681H	1.3[h]

a Substitution tested in the context of variant B.1.258
b Substitution tested in the context of the Iota and Kappa variants
c Substitution tested in the context of the Delta and Delta Plus variants
d Substitution tested in context of the Beta and Delta Plus variants
e Data shown are for alternative mutations (in parentheses) at the same position
f Substitutions tested in the context of the Beta and Gamma variants
g Substitution tested in the context of the Gamma variant
h Substitution tested in the context of the Mu variant

b)

Figure 3

Figure 4

Table 1

SARS-CoV-2 Variant	Geometric Mean VIR-7831 IC_{50} (ng/ml) (Average Fold Change IC_{50} vs. Wild-Type)	Geometric Mean VIR-7831 IC_{90} (ng/ml) (Average Fold Change IC_{90} vs. Wild-Type)	Geometric Mean VIR-7832 IC_{50} (ng/ml) (Average Fold Change IC_{50} vs. Wild-Type)	Geometric Mean VIR-7832 IC_{90} (ng/ml) (Average Fold Change IC_{90} vs. Wild-Type)
Alpha (B.1.1.7)	187.15 (3.0)	1246.86 (4.1)	181.12 (3.1)	1222.95 (3.7)
Beta (B.1.351)	71.89 (1.2)	385.01 (1.3)	59.67 (1.1)	309.84 (0.9)
Gamma (P.1)	73.11 (1.6)	335.79 (1.4)	48.94 (1.2)	217.02 (0.9)
Delta (B.1.617.2)	51.29 (0.4)	219.08 (0.6)	59.13 (0.5)	274.24 (0.7)
Kappa (B.1.617.1)	118.64 (0.9)	379.89 (1.0)	132.37 (1.0)	428.83 (1.0)

Table 2

Variant	Spike Mutations	Fold-Change in VIR-7831 IC$_{50}$ vs. Wild-type	Fold-Change in VIR-7832 IC$_{50}$ vs. Wild-type
Alpha (B.1.1.7)	H69-, V70-, Y144-, N501Y, A570D, D614G, P681H, T716I, S982A, D1118H	2.3	2.5
Beta (B.1.351)	L18F, D80A, D215G, R246I, K417N, E484K, N501Y, D614G, A701V	0.6	0.7
Gamma (P.1)	D138Y, D614G, E484K, H655Y, K417T, L18F, N501Y, P26S, R190S, T1027I, T20N, V1176F	0.4	0.4
Delta (B.1.617.2)	T19R, G142D, E156G, F157-, R158-, L452R, T478K, D614G, P681R, D950N	1	NT
Epsilon (B.1.427/B.1.429)	S13I, W152C, L452R, D614G	0.7	NT
Eta (B.1.525)	Q52R, A67V, H69-, V70-, Y144-, E484K, D614G, Q677H, F888L	0.9	NT
Iota (B.1.526)	L5F, T95I, D253G, E484K, D614G, A701V	0.6	NT
Kappa (B.1.617.1)	T95I, G142D, E154K, L452R, E484Q, D614G, P681R, Q1071H	0.7	NT
Lambda (C.37)	G75V, T76I, del246-252, L452Q, F490S, T859N	1.5	NT
Delta Plus (AY.1)	T19R, T95I, G142D, E156G, F157-, R158-, W258L, K417N, L452R, T478K, D614G, P681R, D950N	1.1	NT
Delta Plus (AY.2)	T19R, V70F, G142D, E156G, F157-, R158-, A222V, K417N, L452R, T478K, D614G, P681R, D950N	1.3	NT
Delta Plus (AY. 4.2)	T19R, T95I, G142D, Y145H, E156G, F157, R158-, A222V, L452R, T478K, D614G, P681R, D950N	1.6	NT
Mexico/Swiss (B.1.1.519)	T478K, D614G, P681H, T732A	0.8	NT
Scotland (B.1.258)	H69-, V70-, N439K, D614G	0.9	NT
US (R.2)	E484K, D614G, Q677H, T732S, E1202Q	0.8	NT
Liverpool (A.23.1)	R102I, F157L, V367F, E484K, Q613H, P681R	1.1	NT
Cameroon (B.1.619)	I210T, N440K, E484K, D614G, D936N, S939F, T1027I	1.3	NT
Bristol (B.1.1.7+E484K)	H69-, V70-, Y144-, E484K, N501Y, A570D, D614G, P681H, T716I, S982A, D1118H	1.7	NT
Mu (B.1.621)	T95I, Y144T, Y145S, ins146N, R346K, E484K, N501Y, D614G, P681H, D950N	1.3	NT

Table 3

Amino Acid position	Substitution / Deletion	mAb with Reduced Susceptibility	Variants in Tested Spike Sequence	VIR-7831 IC_{50} (ng/ml)	Average Fold Change in IC_{50} Compared to Relative Wild-Type
E406	W	casirivimab+ imdevimab	E406W	53.66	0.74
K417	E	casirivimab	K417E	67.71	0.89
N439	K	imdevimab	N439K, D614G	17.05	0.86
N440	D	imdevimab	N440D	80.47	1.29
N440	K	imdevimab	N440K, D614G	19.99	0.48
K444	Q	imdevimab	K444Q	79.68	1.11
V445	A	imdevimab	V445A	41.74	3.38
G446	V/I	imdevimab	G446V, D614G	18.41	1.50
Y453	F	casirivimab	G261D, Y453F	27.28	2.19
L455	F	casirivimab	L455F, D614G	21.65	0.56
G476	S	casirivimab	G476S	36.97	2.94
E484	K	bamlanivimab	E484K, D614G	12.91	0.33
F486	V/I	casirivimab	F486V	82.24	1.10
Y489	H	casirivimab	Y489H	92.29	1.48
F490	S	bamlanivimab	F490S	33.10	0.85
Q493	K	casirivimab, bamlanivimab	Q483K	69.79	0.98
S494	P	casirivimab, bamlanivimab	S494P, D614G	29.10	2.50

Table 4

Amino Acid Position	Reference Amino Acid[a]	Substitutions Identified in order of Prevalence[b]	Percent Reference AA Conservation
332	I	V, T, N, D, S	>99.99
333	T	I, K, A, S, P, R	>99.99
334	N	K, Y, S, H, D, T, -, E, I	>99.99
335	L	F, V, S, -, G, M	>99.99
336	C	S*, -, P, R	>99.99
337	P	L, S, T, H, R, -, K	>99.99
339	G	D, S, C, V, F, -, N	99.98
340	E	D, K, A, G, Q, V, -, I	>99.99
341	V	I, F, A, -, G, L, P, S	>99.99
343	N	Y, -, D, K, S	>99.99
344	A	S, T, V, D, F, G, -, P	99.97
345	T	S, I, -, N	>99.99
346	R	K, S, I, G, T, -, F	99.66
354	N	K, D, S, T, H, Y, I, -, G, P	99.95
356	K	R, N, E, Q, M, T, -, G, W, Y, I, KR	99.98
357	R	K, I, -, T, G, S, C, L, RIS	99.95
358	I	V, L, F, -, T*, A, E*	>99.99
359	S	N, T, G, R, I, -, C, F	99.98
360	N	Y, S, -, K, D, F, T, A, L	>99.99
361	C	T*, -, G*, N*, F*, R*, S*, Y	>99.99
440	N	K, S, Y, T, D, I, E, -, F, H, R	99.79
441	L	R, F, I, P, V, -, Y, H, M, S, E, N	>99.99
509	R	-, K*, S*, T*, I*, H	>99.99

AA = amino acid
[a]NCBI reference sequence: YP_009724390.1
[b]Dash indicates a stop codon. Asterisks indicate the substitution could not be evaluated due to low expression of the spike protein containing this substitution

Supplemental Figure 1

Supplemental Figure 2

Supplemental Figure 3

Supplemental Table 1

Passage	Spike Gene Amino Acid Substitution (Freq) [a,b]	EC$_{50}$ (µg/mL)	Fold Change in EC$_{50}$ to WT[c]
SARS-CoV-2 virus stock[c]	H66R (5.7%) N74K (12.6%) T76I (5.6%) 215-216insKLRS (60.9%) H655Y (3.1%)	0.06	NA
VIR-7832 Lineage 1, passage 4	215-216insKLRS (74.5%) 675-679 del (20.6%)	0.34	5.64
VIR-7832 Lineage 1, passage 5	215-216insKLRS (74.6%) 675-679del (66.0%)	0.35	5.93
VIR-7832 Lineage 1, passage 8	215-216insKLRS (74.7%) E340A (98.7%) 675-679del (84.5%)	ND	>10
VIR-7832 Lineage 2, passage 5	215-216insKLRS (73.9%) 675-679del (47.3%) R682W (4.9%) V1128F (3.5%)	0.32	5.40
VIR-7832 Lineage 2, passage 6	215-216insKLRS (75.3%) 675-679del (74.2%) R682W (4.9%) V1128F (30.9%)	0.39	6.54

Supplemental Table 2

Amino Acid Changes in Spike protein	VIR-7831		VIR-7832	
	Geomean Neutralization IC_{50} (ng/mL)	Fold Change Relative to Wild-Type	Geomean Neutralization IC_{50} (ng/mL)	Fold Change Relative to Wild-Type
Wild Type[a]	104.46	NA	100.75	NA
E340A	> 10,000	>100	> 10,000	> 107
R682W	53.96	0.52	47.78	0.49
V1128F	50.65	0.53	49.69	0.60

Supplemental Table 3

Epitope Reference Amino Acid	Amino Acid Substitutions in Spike Protein	Geomean Neutralization IC_{50} (ng/mL)	Average Fold Change IC_{50} Relative to Reference[a]
I332	I332V, D614G	53.43	1.48
	I332T, D614G	66.74	1.61
T333	T333K, D614G	22.14	0.67
	T333I, D614G	30.93	0.59
N334	N334K, D614G	45.36	1.27
	N334H, D614G	30.74	0.87
	N334S, D614G	34.37	1.13
	N334Y, D614G	32.89	0.96
L335	L335F	29.19	0.81
	L335S, D614G	29.8	0.85
	L335V, D614G	42.2	0.63
P337	P337H, D614G	185.29	5.13
	P337L, D614G	>10000	>192
	P337K, D614G	>10000	>304
	P337R, D614G	>10000	>192
	P337S, D614G	127.69	1.26
	P337T, D614G	383.27	10.62
G339	G339D, D614G	117.38	1.18
	G339S, D614G	32.67	0.86
	G339C, D614G	68.79	1.18
	G339V, D614G	78.32	1.26
E340	E340A	>10000	>100
	E340D, D614G	81.46	2.45
	E340K	>10000	>297
	E340G, D614G	640.07	18.21
	E340Q, D614G	>2500	>50
	E340V, D614G	>10000	>200
V341	V341I, D614G	14.6	0.16
N343	N343S, D614G	16.87	0.48

Supplemental Table 3 (con't)

Epitope Reference Amino Acid	Amino Acid Substitutions in Spike Protein	Geomean Neutralization IC$_{50}$ (ng/mL)	Average Fold Change IC$_{50}$ Relative to Reference[a]
A344	A344S	92.19	0.89
	A344T, D614G	40.94	0.62
	A344V, D614G	53.87	1.14
	A344D, D614G	131.53	2.57
T345	T345S, D614G	42.92	0.69
	T345N, D614G	20.29	0.57
	T345I, D614G	11.78	0.20
R346	R346I, D614G	65.39	1.72
	R346S, D614G	42.87	1.13
	R346T, D614G	89.04	1.25
	R346K, D614G	24.76	0.72
	R346G, D614G	43.73	0.85
N354	N354D	104.8	1.00
	N354K, T95I	70.62	0.76
	N354H, D614G	74.88	1.06
	N354K, T95I	70.62	0.76
	N354S, D614G	61.78	0.89
	N354I, D614G	37.09	1.05
	N354T, D614G	43.32	0.65
	N354Y, D614G	67.63	1.08
K356	K356A	111.18	1.06
	K356R, D614G	29.61	0.78
	K356N, D614G	53.22	1.12
	K356M, D614G	64.09	1.03
	K356T, D614G	281.13	5.9
	K356E, D614G	76.54	1.65
	K356Q, D614G	43.2	0.94
R357	R357K, D614G	39.13	1.02
	R357I, D614G	46.41	0.98

Supplemental Table 3 (con't)

Epitope Reference Amino Acid	Amino Acid Substitutions in Spike Protein	Geomean Neutralization IC_{50} (ng/mL)	Average Fold Change IC_{50} Relative to Reference[a]
I358	I358V, D614G	32.71	0.7
	I358L	44.01	0.8
	I358F, D614G	42.69	1.19
S359	S359N	95.55	0.96
	S359G, D614G	37.56	0.8
	S359R, D614G	62	0.89
	S359T, D614G	39.98	0.84
	S359I, D614G	38.01	0.84
N360	N360S, D614G	34.08	0.72
	N360Y, D614G	29.67	0.61
N440	N440D	80.47	1.29
	N440Y, D614G	42.65	0.68
	N440S, D614G	51.34	0.73
	N440K, D614G	19.99	0.48
	N440I, D614G	40.21	1.22
	N440T, D614G	50.81	0.82
	N440E, D614G	46.97	0.80
	N440F, D614G	43.07	0.85
L441	L441F, D614G	25.21	0.4
	L441I, D614G	32.96	0.7
	L441R, D614G	42.3	0.87
	L441V, D614G	48.24	0.98
	L441P, D614G	44.95	0.76