1   Movement patterns and activity levels are shaped by the neonatal environment

2   in Antarctic fur seal pups

4   Rebecca Nagel [1], Sina Mews [2], Timo Adam [2,3], Claire Stainfield [4], Cameron Fox-Clarke [4],

5   Camille Toscani [4], Roland Langrock [2], Jaume Forcada [4*], Joseph I. Hoffman [1,4*]

7   1. Department of Animal Behaviour, Bielefeld University, 33501 Bielefeld, Germany

8   2. Department of Business Administration and Economics, Bielefeld University, 33501

9   Bielefeld, Germany

10   3. Centre for Research into Ecological and Environmental Modelling, School of

11   Mathematics and Statistics, The Observatory, Buchanan Gardens, University of St

12   Andrews, KY16 9LZ, UK

4. British Antarctic Survey, High Cross, Madingley Road, Cambridge CB3 OET, UK

* Joint senior authors

Corresponding author: Rebecca Nagel, rebecca.nagel@uni-bielefeld.de, Department of

Animal Behaviour, Bielefeld University, 33501 Bielefeld, Germany

Abstract

Tracking studies of juveniles are rare compared to those of adults and consequently, little is

known about the influence of intrinsic and extrinsic factors on activity during this critical life

stage. We therefore collected GPS data from birth until moult from 66 Antarctic fur seal pups

in two nearby but contrasting breeding colonies. The Special Study Beach (SSB) is a high-

density colony separated from the inland tussock grass habitat by a steep gully, whereas

Freshwater Beach (FWB) is a low-density colony that is topographically more open. We

investigated the explanatory power of multiple individual-based and environmental variables

using a hidden Markov model. We found that activity was typified by central place

exploration, with directed movement away from and subsequent return to a location of low

activity. The probability of such directed exploration was unaffected by several factors known

to influence marine mammal movement including sex, body condition and temperature.

Conversely, FWB pups were more active, increased their activity with age and transitioned

earlier into the tussock grass, which offers protection from predators and extreme weather.

Our study illustrates the profound importance of extrinsic factors to early movement and

highlights the adaptive potential of movement in the face of environmental variability.

Keywords

Activity, animal movement, fur seal, GPS, hidden Markov model (HMM), pinniped

Introduction

Movement is a defining characteristic of life, and it underpins critical components of behaviour [1] and fitness [2]. Understanding how animals adapt their movement in response to external and internal stimuli is therefore fundamental to the management and conservation of wild populations [1,3]. Rapid advancements in bio-logging technology and the increased accessibility of statistical tools for drawing meaningful inference from fine-scaled observations has led to unprecedented insights into the movement of a variety of species in their natural habitats [4,5]. Marine vertebrates in particular have benefited from such improved methodologies, exemplified by a recent analysis of more than 2,600 tracked individuals documenting extraordinary convergence in movement patterns across 50 species [6].

Despite these advancements, several authors have drawn attention to age and sex biases in the literature, with datasets of adult females being particularly over-represented [3,7]. This focus on adults is problematic because the movement and distribution of neonatal and juvenile individuals is of key importance for understanding population dynamics. For example, long-term monitoring of three albatross species has shown that high rates of juvenile mortality due to interactions with fisheries are likely driving observed declines in population size [8]. Similarly, loggerhead sea turtles occupy distinct oceanic environments as juveniles, making them more susceptible to bycatch on pelagic longlines [9].

In addition to informing conservation efforts, studies of the movement of young individuals can also provide valuable insights into the ontogeny of social and survival skills. A tracking study of European shags, for example, has shown that higher rates of juvenile mortality correlate with poor foraging proficiency [10], highlighting the importance of learning and memory for successful recruitment. Studies of several different pinniped species including Antarctic fur seals [11], New Zealand sea lions [12], Steller sea lions [13], northern fur seals [14] and grey seals [15] have also shown that sex-specific differences in foraging and diving behaviour become established early in life, implying that intrinsic factors may outweigh size-related effects associated with early sexual dimorphism on habitat choice.

These and other studies represent an important step forward in the field of movement ecology because they help to close the gap between adult movement and the 'lost years' of juveniles [3]. However, with few exceptions, studies of juveniles have focused on the period after nutritional independence. While this can be justified for many birds and other species where neonates are more or less stationary until weaned, in many other species movement behaviour and social interactions among conspecifics earlier in life play a key role in

development. For example, play behaviour in pre-weaned Stellar sea lions and Galápagos fur

seals facilitates the development of muscle mass and hones fighting skills that are important

for future foraging and reproductive success [16,17].

Antarctic fur seals (*Arctocephalus gazella*) are another pinniped species where

neonatal movement may play an important role in early development. Newborn pups are

regularly left unattended in dense rookeries while their mothers, who are central place

foragers, intersperse short periods of nursing on land with longer foraging trips at sea [18].

Without their mothers to defend them, pups are at an increased risk of predation from

predatory birds [19] and traumatic injury due to crushing by territorial males [20,21].

Furthermore, the natal coat of pups lacks the water-repellent properties of adult fur [22],

which prevents them from spending prolonged time at sea, so pups can only begin to develop

the efficient swimming and diving behaviour necessary for nutritional independence after they

moult at around 60 days of age [23,24]. Anecdotal observations of pups venturing into

shallow pools and streams shortly after birth [23] and the emergence of sex-specific

differences in habitat use later in life [25] provide some insights into pup activity, but we still

know surprisingly little about the intrinsic and extrinsic factors that may shape pup movement

prior to moulting.

Here, we analysed hourly GPS data from 66 Antarctic fur seal pups tracked from birth

until moulting (Figure 1) using a hidden Markov model. Focal individuals were selected at

random from two breeding colonies, the "Special Study Beach" (SSB) and "Freshwater

Beach" (FWB), which are separated by around 200 meters. Because of their close proximity,

these colonies experience comparable climatic conditions and breeding females from both

locations likely forage in the same area [26], which is reflected by the fact that they do not

differ significantly in quality traits such as body size or condition [27]. However, the density

of animals is higher at SSB than at FWB [27]. This creates a different dynamic, first because

tightly packed males at SSB have little freedom to move, resulting in a more static system

[28], and second because avian predators are better able to penetrate low density breeding

aggregations on FWB, leading to a greater likelihood of pups being predated [27]. The two

colonies also differ in their size and topology. SSB is somewhat smaller and relatively

'closed' as a steep gully separates the beach from the tussock grass immediately inland, while

FWB is flatter and consequently more 'open'.

We used these two colonies as a 'natural experiment' to investigate the influence of

the early neonatal environment on fur seal movement, as well as to evaluate the relative

importance of extrinsic versus intrinsic factors such as sex, age and body condition.

Furthermore, as the number of breeding individuals, foraging trip durations and pup birth
weights vary appreciably from year to year depending on local food abundance [29], we
replicated our study across two consecutive seasons of contrasting food availability [27]. We
hypothesised that (i) intrinsic and extrinsic factors shown in previous studies to influence
movement in pinnipeds, including sex [30–32], body condition [33] and ambient temperature
[34–36], will also affect the movement of Antarctic fur seal neonates, (ii) activity levels and
movement patterns will be influenced by density, with FWB pups being more active to avoid
predation, and (iii) levels of pup activity will increase with age, as there is likely strong
selection for the early development of motor skills and lean muscle mass.

Methods

Field methods

This study was conducted at two breeding colonies on Bird Island, South Georgia (54°00'24.8⬜ S, 38°03'04.1⬜ W; Figure 2a) during the austral summers (December to March) of 2018 – 19 (hereafter referred to as 2019) and 2019 – 20 (hereafter referred to as 2020). The first of these years was among the worst on record in terms of breeding female numbers, pup birth weights and maternal foraging trip durations [27], implying that 2019 was a year of unusually low food abundance [29,37,38].

In both seasons, 25 Antarctic fur seal pups from each colony were captured two to three days after birth (December) and every ten days thereafter until they began to moult (March). Individuals were randomly selected with respect to sex and birth date during the pupping period. Capture, restraint and handling of pups followed protocols established over 36 consecutive years of the long-term monitoring and survey program of the British Antarctic Survey (BAS). Briefly, pups were captured with a slip noose or by hand and were restrained by hand. After handling, pups were returned to their mothers or released as close to their capture sites as possible. At every capture, weight and length measurements were taken, from which a scaled mass index was calculated according to Peig and Green [39]. This condition metric serves as a reliable indicator of overall fitness as it has been correlated with mortality [40] and reproductive success [41] in a variety of species.

To facilitate the tracking and recapture of focal individuals, VHF transmitters (Sirtrack core marine glue-on V2G 152A; dimensions: 40 x 20 x 10 mm body with a 200 mm antenna, weight: 16 g) were attached to the dorsal side of the neck between the shoulder blades with epoxy glue. Pups were similarly fitted with GPS loggers in waterproof ABS plastic enclosures (Perthold Engineering CatLog Gen2 GPS Loggers; dimensions: 50 x 40 x 17 mm, weight: 36 g), which recorded latitude and longitude positions every hour from birth until moulting of the pelage (Figure 1). Together, these two tags accounted for less than 1% of the mean focal pup mass at first capture (5.6 kg). Temporary bleach marks (Clairol Nice'n Easy Borne Blonde Permanent Hair Color) applied to the fur were used to identify pups. Unrecovered tags ($n = 12$) and bleach marks are shed naturally when the seals moult (March – April), precluding any long-term consequences for the pups.

As part of the BAS contribution to the Ecosystem Monitoring Programme of the Convention for the Conservation of Antarctic Marine Living Resources (CCAMLR), the attendance behaviour of breeding females has been monitored since 1982, with radio telemetry protocols established in 1992 to track around 25 adult females per season on FWB.

We contributed towards this ongoing effort by attaching VHF transmitters (Sirtrack core
marine glue-on V2G 154C; dimensions: 65 x 28 x 13 mm body with a 250 mm antenna,
weight: 42 g) to the mothers of our focal FWB pups. Adult females were captured with a
noosing pole and held on a restraint board. Daily attendance was monitored using a fixed-
position radio antenna (Televilt RX900) combined with visual checks of the island with a
hand-held VHF receiver (AOR Ltd., AR8200). The daily absence or presence of females
ashore was noted from first capture until the final measurement, when the pups either moulted
or died. Mothers were fitted with cattle ear tags (Dalton Supplies, Henley on Thames, UK) in
the trailing edge of each fore flipper [42] for identification.

Data analysis framework

Within our dataset, hourly GPS data were successfully collected from a total of 66
Antarctic fur seal pups (breakdown by season: $n = 40$ in 2019 and $n = 26$ in 2020; by colony:
$n = 32$ at FWB and $n = 34$ at SSB). The majority of tracked individuals ($n = 53$, or 80%)
survived until weaning. For those individuals that died, we truncated the GPS data back to the
last time point that the animal was seen alive prior to analysis.

Our GPS data contained some missing locations, which in a handful of cases led to
observation gaps of several hours. Most if not all missing values likely arose due to signal
lapse, which occurs when the satellite connection to a GPS device is interrupted due to cloud
cover, physical obstruction, etc. [43]. In an otherwise regular sequence, such missing data are
manageable, but large gaps may introduce bias into the model [44]. Therefore, for
observational gaps exceeding four hours, we split the individual's track into separate bursts or
intervals of continuous data. Bursts that were shorter than 48 hours usually contained a
considerable amount of missing observations and were therefore excluded from the analysis,
corresponding to roughly 2% of the full dataset including missing locations. Overall, the
tracks of 35 individuals were split into two to five bursts, while data from all other individuals
($n = 31$) consisted of a single burst.

Our final dataset contained 67,417 hourly observations with a median number of 1,124
observations per individual (min = 88, max = 1,916). The step lengths (in meters) between
consecutive GPS locations were calculated and screened for implausible movements. We set
all values larger than 400 meters/hour (corresponding to 297 observations) to be missing. This
is in accordance with Jones *et al.* [25], who found that 300 meters was the maximum distance
Antarctic fur seal pups younger than 50 days of age travelled.

To investigate environmental factors influencing pup movement, hourly dry bulb air
temperature and wind speed measurements for Bird Island were obtained from the BAS
Meteorology and Ozone Monitoring Programme, and missing values (85 observations) were
linearly interpolated. The binary variables sex (with female as the reference category) and
season (with 2019 as the reference category) were held constant for each individual. The
continuous variables age (measured in days since initial capture, two to three days after birth)
and scaled mass index (calculated every ten days) were kept constant between days and
measurements. All metric variables were standardised. All analyses and visualisations for this
study were implemented in R version 4.0.2 [45] using the integrated development
environment RStudio [46].

Hidden Markov models

To investigate the explanatory power of environmental and individual-based variables
on pup movement, we fitted a hidden Markov model (HMM) to the hourly GPS data. This
time series model encompasses both the observed movement of an individual and an
underlying ('hidden') state sequence, which is used to infer behavioural processes (e.g. active/
inactive) [5]. HMMs have accordingly been used to analyse animal tracking data in relation
to, among others, environmental conditions [47], anthropogenic activity [48], sex [49],
ontogeny [50] and individualised niches [51].

To distinguish between different behavioural processes in Antarctic fur seal pups, we
fitted HMMs to the hourly observed step lengths. We assumed the observation process to be
the same for all individuals and modelled the step lengths using a gamma distribution,
conditional on the states. Prior to building a final model, we conducted exploratory analyses
on the number of states and compared univariate HMMs to bivariate HMMs additionally
including the turning angle between consecutive GPS locations (see Supplementary Material
S3 for more details). We restricted our final analysis to a parsimonious univariate HMM with
two states as the inclusion of the turning angle as well as additional states resulted in a
negligible improvement of model fit while complicating the interpretation of the underlying
states [52]. To check whether the HMMs based on hourly data adequately reflect the animals'
movement behaviour, we also fitted an HMM to five-minute interval GPS data collected from
three fur seal pups, each between 30 – 50 days of age, for 13 – 19 days (mean = 3,965 data
points). The results showed that with the coarser temporal resolution (hourly data) there was
no relevant loss of information on pup activity with respect to the aims of the study (see
Supplementary Material S4 for more details).

We further investigated the effects of external and internal factors – including colony,

season, time of day, temperature, wind speed, sex, age and body condition – on the activity

level of fur seal pups by modelling state transition probabilities as a function of the covariates

using a logit link function. We allowed the effects of the covariates to differ between the two

colonies by including interaction terms with the binary covariate 'colony' (FWB/SSB). To

account for the periodic nature of time of day, its effect was modelled using trigonometric

functions. To assess the relative importance of the covariates, we calculated the differences in

AIC values between the full model and models sequentially excluding each covariate and its

interaction with colony. All HMMs were fitted by numerically maximising the likelihood

using *moveHMM* [44]. To further investigate model fit, we calculated pseudo-residuals and

checked them for normality and autocorrelation [5]. Based on the full model, we inferred the

most likely underlying state sequence using the Viterbi algorithm [5]. We also calculated the

state occupancy probabilities as a function of each covariate [53], while fixing the other

covariates to their respective means or reference categories.

Post-hoc analyses

The HMM decomposed the GPS data into two distinct states, which we inferred as

'inactive' and 'active' movement (see Results for more details). To determine whether

additional variables that we were not able to include in the HMM explain a significant

proportion of variation in activity, we built two generalised linear mixed models (GLMMs)

post-hoc. In our first model, we included maternal attendance (0 = present ashore, 1 = absent)

as the explanatory variable. As data on maternal attendance were only available for pups born

on FWB, this covariate could not be included in the full HMM, which defined an interaction

term between all covariates and 'colony'. In our second model, we used the survival status of

each pup (0 = died, 1 = survived) as the explanatory variable. Here, we wanted to test for an

association between survival status and movement without assuming the direction of

causality. In both models, pup ID was included as a random effect to account for repeated

measurements of individuals. The response variable was the inferred state of an individual (0

= inactive, 1 = active) at a given time point and was modelled with a binomial error

distribution using *lme4* [54]. The residuals of the models were visually inspected for linearity

and equality of error variances (using plots of residuals versus fits), normality (using Q – Q

plots) and over or under-dispersion (by comparing the dispersion of simulated to observed

residuals) using *DHARMa* [55].

Animal ethics and permits

Animal handling was carried out by the British Antarctic Survey under permits from

the Government of South Georgia and the South Sandwich Islands (Wildlife and Protected

Areas Ordinance (2011), RAP permit numbers 2018/024 and 2019/032). All procedures used

were approved by the British Antarctic Survey Animal Welfare and Ethics Review Body

(AWERB applications 2018/1050 and 2019/1058).

Results

Hourly GPS data were successfully collected from a total of 66 Antarctic fur seal pups

from two nearby but contrasting breeding colonies (Figure 2a). Deployment durations ranged

from 20 to 80 days (median = 51 days) for the 53 surviving pups and from 4 to 41 days

(median = 15 days) for the 13 pups that died. On average, pups travelled 43.9 meters per hour.

In general, pup movement showed a star-like pattern characterised by directed exploration

within a relatively small area around a central location of low activity (Figure 2b). However,

the spatial distribution of these 'home patches' varied by colony. Pups born at FWB remained

significantly closer to their natal colony (Wilcoxon rank sum test, p = 0.002), traveling on

average 87.6 meters into the tussock grass immediately inland. By contrast, pups born at SSB

moved an average of 205.0 meters from their natal colony in a wider variety of directions

(Figure 2c).

Hidden Markov model: Activity patterns of fur seal pups

The two states of the HMM were clearly discriminated from each other. State 1

captured smaller step lengths (mean step length = 22.5 meters) corresponding to less active

movement and is hereafter referred to as the 'inactive state'. State 2 was characterised by

more active behaviour covering longer distances (mean step length = 75.8 meters) and is

hereafter referred to as the 'active state'. Minimal differences between the marginal

distribution under the fitted model and the empirical distribution as well as the analysis of

pseudo-residuals suggest that our model provides a good fit to the data (see Supplementary

Material S5 for more details). Based on the Viterbi-decoded state sequences, pups spent on

average 65.2% (range = 45.7 – 97.3%) of their time in the inactive state and 34.8% (2.7 –

54.3%) of their time in the active state. The pups' decoded movement patterns are

summarised in Figure 2b and representative examples of individual tracks are shown in

Supplementary Figure S2.

Differences in AIC values (delta AIC) between the full model and HMMs sequentially

excluding each variable and its interaction term with colony are shown in Table 1. We

considered variables to be relevant for modelling pup activity levels if their delta AIC values

were considerably larger than zero. Based on this criterion, the covariates year, time of day

and age were determined to have the strongest effects on pup activity levels via both main

effects and interactions with colony (Table 1). By contrast, air temperature, wind speed, sex

and body condition had little to no influence on activity levels. Pups tended to be slightly

more active in the first year of the study, particularly at FWB where the probability of

occupying the active state fell from 43.8% in 2019 to 34.0% in 2020. Activity levels also

showed a clear diurnal pattern, peaking in the middle of the day in both colonies (Figure 3a

and b). Finally, marked differences were observed in colony-specific developmental

trajectories, with animals from FWB exhibiting an almost linear increase in activity with age

(Figure 3c), whereas activity levels at SSB remained more or less constant throughout

ontogeny (Figure 3d). FWB pups also tended to move inland into the tussock grass earlier

than pups from SSB (Figures 3e and f).

Post-hoc analyses

To test additional hypotheses that could not be addressed in the HMM, we performed

post-hoc logistic regression analyses to evaluate whether the presence of the mother ashore

and the fate of the pup (died or survived until the end of the study) explained a significant

proportion of the variation in pup activity levels ('states'). As expected, pups were

significantly more active when their mothers were absent from Bird Island on foraging trips

(Figure 4a; GLMM, estimate = 0.09, *s.e.* = 0.04, *p*-value = 0.03). More interestingly, the

survival status of a pup was also significantly associated with activity (GLMM, estimate =

0.758, *s.e.* = 0.19, *p*-value < 0.001), with pups that survived until weaning spending on

average 34% more time in the active state (Figure 4b).

Discussion

We collected and analysed hourly GPS data from Antarctic fur seal pups tracked from birth until moulting in order to investigate the drivers of movement patterns and activity levels during this critical life stage. We found that pup movement was characterised by a star-like pattern, with individuals repeatedly returning to a central location of low activity after bouts of directed exploration. The HMM showed that the probability of such active movement was highest during the day and was mainly influenced by season, colony of birth and age. Our findings provide fresh insights into the movement patterns of pinnipeds prior to nutritional independence and highlight the importance of the life-history stage and the neonatal environment on behaviour.

Movement in fur seal pups

A recent study of movement in Antarctic fur seal pups found evidence for sex-specific differences in habitat use prior to nutritional independence and the onset of overt sexual size dimorphism. Specifically, male pups were found to exhibit more risk prone behaviour with increasing age, traveling further at sea and spending more time in exposed beach habitats [25]. While this study highlighted the importance of an intrinsic factor, sex, on neonatal movement, the contributions of other factors, both intrinsic and extrinsic, remained largely unexplored. We therefore used HMMs to conduct a formal analysis incorporating multiple environmental and individual-based variables including the time of day, air temperature, wind speed, age, body condition and sex. Rather than focusing on animals from a single colony, our study design incorporated variation in both density and terrain by tracking pups from two contrasting breeding colonies, while replication across two consecutive seasons also allowed us to account for inter-annual variation in food availability.

Our hourly GPS data revealed a distinct pattern of terrestrial pup movement that has not been reported in previous studies. Specifically, we documented a star-like pattern characterised by directed movement away from and subsequent return to a location of low activity. This appears to be a genuine behavioural pattern rather than an artefact of our temporal scale of sampling, as GPS locations collected every five minutes from a subset of individuals revealed a very similar picture (Supplementary Material Figure S4). Star-like movement patterns are typified by central place foragers, which make round trips between a central location and a foraging patch [56]. However, as fur seal pups are entirely reliant on their mothers for food until they moult, repeated movements away from a central location are unlikely to have anything to do with resource acquisition. Instead, our results appear to be

indicative of bouts of exploration within a defined radius around a suckling location, which
would imply that Antarctic fur seal pups are 'central place explorers'.

We would expect central place exploration in fur seal pups to be adaptive given that it
entails energetic costs and may even make it more difficult for pups to be located by their
mothers when they return from foraging trips. We can envisage a number of possible
explanations for this behaviour. First, high levels of activity may facilitate the development of
the muscle mass necessary for future foraging success. Second, movement both within and
beyond the confines of the beach where a pup is born may increase the scope for social
interactions, which might be important for future mating success. Third, a transition from
beach habitats to the tussock grass offers protection against harsh weather conditions and may
reduce the risk of a pup either being crushed by a territorial male or predated.

Results of the HMM

We used an HMM framework to infer the discrete behavioural states underlying pup
movement and the probability of switching between these states given certain environmental
and individual covariates. This integrated approach to decompose behavioural patterns has
been applied across a wide range of species and habitats and has emerged as a powerful tool
for animal movement modelling [5]. In our model, we uncovered a strong influence of year,
whereby the probability of occupying the active state was higher in the first breeding season
(2019), particularly for FWB pups. One possible explanation for this pattern could be that
inter-annual differences in movement are adaptive. The first breeding season in our study was
among the worst on record in terms of breeding female numbers, pup birth weights and
foraging trip durations [27], indicating low prey abundance and poor environmental
conditions [57]. Local densities at both colonies were correspondingly below average [27].
Given a greater risk of predation at low density [27], pups might increase their activity to
avoid harassment, pecking injuries and predation by generalist seabirds, such as southern and
northern giant petrels [58]. Alternatively, higher levels of activity could be a consequence of
increased stress hormone levels. As mothers spent more time at sea foraging in 2019, pups
were subjected to longer periods of food deprivation. Such extended bouts of starvation in
pinniped pups have been associated with an increase in glucocorticoids [59], which are known
to increase activity levels in some species [60]. Understanding the interrelationships among
prolonged fasting, glucocorticoids levels and activity in fur seal pups may help to shed further
light on individual responses to environmental challenges.

Another clear extrinsic determinant was the time of day, with activity levels peaking

just before solar noon. This observation is in line with a previous study of Galápagos fur seal

pups [17], but otherwise very little is known in general about diurnal patterns of activity

among fur seals while ashore. Daily meter-resolution location data from densely packed

territory holding males reveal negligible movements from day to day [28], implying that these

animals remain more or less stationary. By contrast, telemetry studies on adult female fur

seals correlate nocturnal foraging behaviour with diel variation in the time of arrival and

departure from the breeding colony [61]. Consequently, it is unclear to what extent our results

can be extrapolated across the life history, especially given the likelihood of an ontogenetic

shift in times of peak activity as pups transition from a reliance on milk to nutritional

independence.

The HMM also revealed an increase in pup activity with age, but only for pups born at

FWB. One explanation for this colony-specific difference could be that FWB pups are able to

express their full behavioural repertoire throughout the course of ontogeny, becoming more

active as they grow, whereas pups born on SSB are constrained by the high density of

conspecifics. In particular, pups that traverse tightly packed harems on SSB run a higher risk

of being crushed by a territorial male or bitten by a breeding female [20,21]. Alternatively, we

recently found that FWB pups are more likely to be predated by generalist seabirds [27]. This

might translate into an increase of predator-avoidance activity as pups mature, which would

help to explain why activity no longer increases after around 40 days of age, when the

majority of individuals have transitioned to the more sheltered tussock grass [25].

In addition to pups from FWB increasing their activity over time, we also found that

they tended to move earlier into the tussock grass than pups from SSB. While this earlier

transition could simply be a consequence of increased activity, it might also be adaptive for

FWB pups to move inland as quickly as possible given the higher risk of predation at low

density [27]. In other words, pups may move into the more sheltered tussock grass earlier in

life to avoid predatory seabirds and might thereafter display a higher propensity for active

movement to offset reduced social interactions in the dense vegetation. However, differences

in the topography of the two colonies might also influence the timing of this transition, as a

steep gully separates the beach at SSB from the tussock grass, while FWB offers a more

gradual transition between the two habitats. To disentangle the effects of density and

topography would require a larger study embracing a greater diversity of breeding colonies.

Contrary to our initial expectations, the HMM found little to no effect of air

temperature and wind speed on pup activity. This is surprising given that the pups in our study

lacked the water-repellent fur of adults and were thus poorly protected from the elements [22].
Reduced activity [34] and huddling behaviour [36] have been documented in pinniped
neonates as effective thermoregulatory behaviours to withstand the respectively hottest and
coldest daily temperatures, so we anticipated a reduction in pup activity under marginal
weather conditions. However, large variation in temperature was not observed on Bird Island
(2019 mean temperature = 3.7°C ± *s.d.* 1.3; 2020 mean temperature = 3.9°C ± *s.d.* 1.4).
Consequently, behavioural adjustments in activity for effective thermoregulation may not
have been necessary in the specific context of our study. Alternatively, other climatic
variables that we could not account for, such as precipitation or humidity, may have a
disproportionate influence on activity levels. Future studies involving the direct observation
of focal individuals under diverse climatic conditions would shed further light on this topic.

The HMM also showed that pup activity levels were largely unaffected by body
condition. This was unexpected because Antarctic fur seal pups rely on their mother's milk
for nutrition before moulting their pelage and must tolerate bouts of starvation lasting up to 11
days while their mothers forage at sea [18]. As a result, pup growth is known to decline with
prolonged maternal absence [62], implying that fewer resources should be available for
movement. Taken at face value, the lack of a relationship between body condition and
movement may suggest that the short-term benefits of high activity, such as muscle mass
development and increased social interaction, may outweigh the costs associated with
diverting resources from growth. This would be in line with a previous study that similarly
found that Galápagos fur seal pups maintained high activity levels throughout bouts of
starvation [17]. However, it is also possible that an effect of body condition could not be
detected in our study because our model assumed that condition was constant between
successive measurements, whereas in practice it will vary to some extent from day to day.

Despite a number of studies having shown that sex-specific differences in activity,
habitat use, foraging and diving behaviour are established early in life in several pinniped
species [12–16,63], including Antarctic fur seals [25], our HMM did not uncover any obvious
sex differences in activity. This is probably a consequence of the timeframe of our study.
Jones *et al.* [25], for example, were only able to detect sex-specific differences in habitat use
in Antarctic fur seal pups older than 41 days. Given that we focused on the time window from
birth until moulting at around 60 days of age, the results of the two studies are consistent and
lend support to the notion that sex-specific movement patterns take several weeks to become
established.

Results of the post-hoc *analysis*

Finally, *post-hoc* analysis showed that pups were significantly less active when their
mother was ashore, suggesting that pup activity is correlated to some extent with maternal
foraging behaviour. This association is most likely a reflection of the upmost importance of
milk consumption for pup survival and development. A mother may spend as little as 24
hours and on average only 2 days ashore during each nursing bout [61], so pups must
maximise nutrient update during this time. Moreover, adult female fur seals frequently display
aggressive behaviour towards foreign pups [20], potentially limiting a pup's opportunity for
social interactions when the mother is present [17].

We also found that pups that died tended to be less active. In our dataset, cause of
death was assigned to all but one pup as either starvation ($n = 5$) or predation ($n = 6$), closely
related determinants given that smaller, weaker pups may be more likely to be predated. One
possible explanation for the reduced activity of pups that died could therefore be related to
poor body condition. While this would appear to contradict the lack of an overall association
between activity and condition in the HMM, it is possible that activity may decline only after
body condition falls below a critical threshold [17], an effect we may not have captured due to
the small number ($n = 13$) and early occurrence (median = 15 days) of mortalities in our
dataset. Alternatively, causality could flow in the other direction, with less active pups being
more likely to be predated. This would be more in line with our HMM results as well as with
the hypothesis that increased activity is related to predator-avoidance behaviour.

Conclusions

We document the movement behaviour and activity levels of Antarctic fur seal pups
between birth and moulting in relation to a multitude of extrinsic and intrinsic explanatory
variables. Our findings suggest that during this early life stage, pup activity is mainly shaped
by extrinsic factors including year, time of day and colony of birth. In contrast to a previous
study of activity later in life [25], we found little effect of intrinsic factors such as body
condition and sex. Our study therefore highlights the importance of external influences during
this critical phase of life when pups are at their greatest risk of predation and have not yet
developed their water-repellent adult coat.

Data Availability Statement

Raw data will be uploaded to the Dryad Digital Repository upon acceptance of this manuscript. R code is available as a R Markdown file and included in the Supplementary material.

Acknowledgements

The authors would like to thank Ana Bertoldi Carneiro, Freya Blockley, Jamie Coleman, Alexandra Dodds, Vicki Foster, Derren Fox, Iain Angus Gordon, Pauline Goulet, Rosie Hall, Cary Jackson, Adam Lowndes, Elizabeth Morgan, Rachael Orben, Jessica Ann Philips, David Reid and Mark Whiffin for help in the field.

Author contributions

RN, CS, CF-C and CT collected the data. SM and TA built the HMM models. JIH, RL and JF conceived of the study and contributed funding and materials. All of the authors commented on and approved the final manuscript.

Funding

This research was funded by the German Research Foundation (DFG) as part of the SFB TRR 212 (NC³) – Project numbers 316099922, 396774617 and 396782756. It was also supported by core funding from the Natural Environment Research Council to the British Antarctic Survey's Ecosystems Program.

References

1.   Liedvogel M, Chapman BB, Muheim R, Åkesson S. 2013 The behavioural ecology of
animal movement: reflections upon potential synergies. *Anim. Migr.* , 39–46.
(doi:10.2478/ami-2013-0002)

2.   Crossin GT, Cooke SJ, Goldbogen JA, Phillips RA. 2014 Tracking fitness in marine
vertebrates: Current knowledge and opportunities for future research. *Mar. Ecol. Prog.*
Ser. **496**, 1–17. (doi:10.3354/meps10691)

3.   Hays GC *et al.* 2016 Key questions in marine megafauna movement ecology. *Trends*
Ecol. Evol. **31**, 463–475. (doi:10.1016/j.tree.2016.02.015)

4.   Hussey NE *et al.* 2015 Aquatic animal telemetry: A panoramic window into the
underwater world. *Science (80-.).* **348**, 1255642. (doi:10.1126/science.1255642)

5.   Zucchini W, MacDonald IL, Langrock R. 2016 *Hidden Markov models for time series:*
An introduction using R. Boca Raton: Chapman and Hall/CRC.

6.   Sequeira AMM *et al.* 2018 Convergence of marine megafauna movement patterns in
coastal and open oceans. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 3072–3077.
(doi:10.1073/pnas.1716137115)

7.   McIntyre T. 2014 Trends in tagging of marine mammals: A review of marine mammal
biologging studies. *African J. Mar. Sci.* **36**, 409–422.
(doi:10.2989/1814232X.2014.976655)

8.   Pardo D, Forcada J, Wood AG, Tuck GN, Ireland L, Pradel R, Croxall JP, Phillips RA.
2017 Additive effects of climate and fisheries drive ongoing declines in multiple
albatross species. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E10829–E10837.
(doi:10.1073/pnas.1618819114)

9.   Watson JW, Epperly SP, Shah AK, Foster DG. 2005 Fishing methods to reduce sea
turtle mortality associated with pelagic longlines. *Can. J. Fish. Aquat. Sci.* **62**, 965–
981. (doi:10.1139/f05-004)

10.   Daunt F, Afanasyev V, Adam A, Croxall JP, Wanless S. 2007 From cradle to early
grave: Juvenile mortality in European shags *Phalacrocorax aristotelis* results from
inadequate development of foraging proficiency. *Biol. Lett.* **3**, 371–374.
(doi:10.1098/rsbl.2007.0157)

11.   Kernaléguen L, Arnould JPY, Guinet C, Cazelles B, Richard P, Cherel Y. 2016 Early-
life sexual segregation: Ontogeny of isotopic niche differentiation in the Antarctic fur
seal. *Sci. Rep.* **6**, 33211. (doi:10.1038/srep33211)

12.   Leung ES, Chilvers BL, Nakagawa S, Robertson BC. 2014 Size and experience matter:

Diving behaviour of juvenile New Zealand sea lions (*Phocarctos hookeri*). *Polar Biol.*
37, 15–26. (doi:10.1007/s00300-013-1405-6)

13. Bishop A, Brown C, Rehberg M, Torres L, Horning M. 2018 Juvenile Steller sea lion
(*Eumetopias jubatus*) utilization distributions in the Gulf of Alaska. *Mov. Ecol.* **6**, 1–
15. (doi:10.1186/s40462-018-0124-6)

14. Zeppelin T, Pelland N, Sterling J, Brost B, Melin S, Johnson D, Lea MA, Ream R.
2019 Migratory strategies of juvenile northern fur seals (*Callorhinus ursinus*): Bridging
the gap between pups and adults. *Sci. Rep.* **9**, 1–16. (doi:10.1038/s41598-019-50230-z)

15. Carter MID, McClintock BT, Embling CB, Bennett KA, Thompson D, Russell DJF.
2019 From pup to predator: generalized hidden Markov models reveal rapid
development of movement strategies in a naïve long-lived vertebrate. *Oikos* **129**, 630–
642. (doi:10.1111/oik.06853)

16. Gentry RL. 1974 The development of social behavior through play in the Steller sea
lion. *Am. Zool.* **14**, 391–403. (doi:10.1093/icb/14.1.391)

17. Arnold W, Trillmich F. 1985 Time budget in Galápagos fur seal pups: The influence of
the mother's presence and absence on pup activity and play. *Behaviour* **92**, 302–321.

18. Forcada J, Staniland IJ. 2018 Antarctic Fur Seal. In *Encyclopedia of Marine Mammals*
(eds B Würsig, JGM Thewissen, KM Kovacs), pp. 25–27. Academic Press.
(doi:10.1016/b978-0-12-804327-1.00046-7)

19. Hunter S. 1983 The food and feeding ecology of the giant petrels *Macronectes halli*
and *M. giganteus* at South Georgia. *J. Zool.* **200**, 521–538. (doi:10.1111/j.1469-
7998.1983.tb02813.x)

20. Doidge DW, Croxall JP, Baker JR. 1984 Density-dependent pup mortality in the
Antarctic fur seal *Arctocephalus gazella* at South Georgia. *J. Zool. London* **202**, 449–
460. (doi:10.1111/j.1469-7998.1984.tb05095.x)

21. Reid K, Forcada J. 2005 Causes of offspring mortality in the Antarctic fur seal,
Arctocephalus gazella: The interaction of density dependence and ecosystem
variability. *Can. J. Zool.* **83**, 604–609. (doi:10.1139/z05-045)

22. Irving L, Peyton LJ, Bahn CH, Peterson RS. 1962 Regulation of temperature in fur
seals. *Physiol. Zool.* **35**, 275–284.

23. McCafferty DJ, Boyd IL, Taylor RI. 1998 Diving behaviour of Antarctic fur seal
(*Arctocephalus gazella*) pups. *Can. J. Zool.* **76**, 513–520. (doi:10.1139/z97-219)

24. Payne MR. 1979 Growth in the Antarctic fur seal *Arctocephalus gazella*. *J. Zool.*
London **187**, 1–20.

25.   Jones KA, Wood H, Jonathan P, Ratcliffe N, Votier SC, Staniland IJ. 2020 Risk

exposure trade-offs in the ontogeny of sexual segregation in Antarctic fur seal pups.

Behav. Ecol. , 1–12. (doi:10.1093/beheco/araa018)

26.   Hunt GL, Heinemann D, Everson I. 1992 Distributions and predator-prey interactions

of macaroni penguins, Antarctic fur seals, and Antarctic krill near Bird Island, South

Georgia. *Mar. Ecol. Prog. Ser.* **86**, 15–30. (doi:10.3354/meps086015)

27.   Nagel R, Stainfield C, Fox-Clarke C, Toscani C, Forcada J, Hoffman JI. 2021 Evidence

for an Allee effect in a declining fur seal population. *Proc. R. Soc. B*

(doi:10.1098/rspb.2020-2882)

28.   Hoffman JI, Trathan PN, Amos W. 2006 Genetic tagging reveals extreme site fidelity

in territorial male Antarctic fur seals Arctocephalus gazella. *Mol. Ecol.* **15**, 3841–3847.

(doi:10.1111/j.1365-294X.2006.03053.x)

29.   Forcada J, Hoffman JI. 2014 Climate change selects for heterozygosity in a declining

fur seal population. *Nature* **511**, 462–465. (doi:10.1038/nature13542)

30.   Photopoulou T, Heerah K, Pohle J, Boehme L. 2020 Sex-specific variation in the use of

vertical habitat by a resident Antarctic top predator. *Proc. R. Soc. B* **287**, 20201447.

(doi:10.1098/rspb.2020.1447)

31.   Jones KA, Ratcliffe N, Votier SC, Newton J, Forcada J, Dickens J, Stowasser G,

Staniland IJ. 2020 Intra-specific niche partitioning in Antarctic fur seals, *Arctocephalus*

gazella. Sci. Rep. **10**, 3238. (doi:10.1038/s41598-020-59992-3)

32.   Le Boeuf BJ, Crocker DE, Blackwell SB, Morris P a, Thorson PH. 1993 Sex

differences in diving and foraging behaviour of northern elephant seals. *Symp. Zool.*

Soc. London **66**, 149–178.

33.   Hassrick JL, Crocker DE, Teutschel NM, McDonald BI, Robinson RW, Simmons SE,

Costa DR. 2010 Condition and mass impact oxygen stores and dive duration in adult

female northern elephant seals. *J. Exp. Biol.* **213**, 585–592. (doi:10.1242/jeb.037168)

34.   Limberger D, Trillmich F, Biebach H, Stevenson RD. 1986 Temperature regulation and

microhabitat choice by free-ranging Galapagos fur seal pups (*Arctocephalus*

galapagoensis). *Oecologia* **69**, 53–59. (doi:10.1007/BF00399037)

35.   Campagna C, Le Boeuf BJ. 1988 Thermoregulatory behaviour of southern sea lions

and its effect on mating strategies. *Behaviour* **107**, 72–90.

(doi:10.1163/156853988X00205)

36.   Gentry RL. 1973 Thermoregulatory behavior of eared seals. *Behaviour* **46**, 73–93.

37.   Croxall JP, McCann TS, Prince PA, Rothery P. 1988 Reproductive performance of

seabirds and seals at South Georgia and Signy Island, South Orkney Islands, 1976–
1987: Implications for Southern Ocean monitoring studies. In *Antarctic Ocean and*
Resources Variability (ed D Sahrhage), pp. 261–285. Springer, Berlin, Heidelberg.
(doi:10.1007/978-3-642-73724-4_23)

38.   Boyd IL, Arnould JPY, Barton T, Croxall JP. 1994 Foraging behaviour of Antarctic fur
seals during periods of contrasting prey abundance. *J. Anim. Ecol.* **63**, 703–713.
(doi:10.2307/5235)

39.   Peig J, Green AJ. 2009 New perspectives for estimating body condition from
mass/length data: The scaled mass index as an alternative method. *Oikos* **118**, 1883–
1891. (doi:10.1111/j.1600-0706.2009.17643.x)

40.   Trites AW, Jonker RAH. 2000 Morphometric measurements and body conditions of
healthy and starveling Steller sea lion pups (*Eumetopias jubatus*). *Aquat. Mamm.* **26.2**,
151–157.

41.   Guinet C, Roux JP, Bonnet M, Mison V. 1998 Effect of body size, body mass, and
body condition on reproduction of female South African fur seals (*Arctocephalus*
pusillus) in Namibia. *Can. J. Zool.* **76**, 1418–1424. (doi:10.1139/z98-082)

42.   Gentry RL, Holt JR. 1982 Equipment and techniques for handling northern fur seals.
NOAA Tech. Rep. NMFS SSRF-758.

43.   Meseck K, Jankowska MM, Schipperijn J, Natarajan L, Godbole S, Carlson J,
Takemoto M, Crist K, Kerr J. 2016 Is missing geographic positioning system data in
accelerometry studies a problem, and is imputation the solution? *Geospat. Health* **11**,
157–163. (doi:10.4081/gh.2016.403)

44.   Michelot T, Langrock R, Patterson TA. 2016 moveHMM: An R package for the
statistical modelling of animal movement data using hidden Markov models. *Methods*
Ecol. Evol. **7**, 1308–1315. (doi:10.1111/2041-210X.12578)

45.   R Core Team. 2019 R: A language and environment for statistical computing.

46.   RStudio Team. 2020 RStudio: Integrated development environment for R.

47.   Roeleke M, Blohm T, Hoffmeister U, Marggraf L, Schlägel UE, Teige T, Voigt CC.
2020 Landscape structure influences the use of social information in an insectivorous
bat. *Oikos* **129**, 912–923. (doi:10.1111/oik.07158)

48.   van Beest FM *et al.* 2019 Classifying grey seal behaviour in relation to environmental
variability and commercial fishing activity - a multivariate hidden Markov model. *Sci.*
Rep. **9**, 5642. (doi:10.1038/s41598-019-42109-w)

49.   Towner A V., Leos-Barajas V, Langrock R, Schick RS, Smale MJ, Kaschke T, Jewell

630   OJD, Papastamatiou YP. 2016 Sex-specific and individual preferences for hunting

631   strategies in white sharks. *Funct. Ecol.* **30**, 1397–1407. (doi:10.1111/1365-2435.12613)

632 50. Grecian WJ, Lane J V., Michelot T, Wade HM, Hamer KC. 2018 Understanding the

633   ontogeny of foraging behaviour: Insights from combining marine predator bio-logging

634   with satellite-derived oceanography in hidden Markov models. *J. R. Soc. Interface* **15**,

635   20180084. (doi:10.1098/rsif.2018.0084)

636 51. Schwarz JFL, Mews S, DeRango EJ, Langrock R, Piedrahita P, Páez-Rosas D, Krüger

637   O. 2021 Individuality counts: A new comprehensive approach to foraging strategies of

638   a tropical marine predator. *Oecologia* (doi:10.1007/s00442-021-04850-w)

639 52. Pohle J, Langrock R, van Beest FM, Schmidt NM. 2017 Selecting the number of states

640   in hidden Markov models: Pragmatic solutions illustrated using animal movement. *J.*

641   *Agric. Biol. Environ. Stat.* **22**, 270–293. (doi:10.1007/s13253-017-0283-8)

642 53. Patterson TA, Basson M, Bravington M V., Gunn JS. 2009 Classifying movement

643   behaviour in relation to environmental conditions using hidden Markov models. *J.*

644   *Anim. Ecol.* **78**, 1113–1123. (doi:10.1111/j.1365-2656.2009.01583.x)

645 54. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting linear mixed-effects models

646   using lme4. *J. Stat. Softw.* **67**, 1–48. (doi:10.18637/jss.v067.i01)

647 55. Hartig F. 2020 DHARMa: Residual diagnostics for hierarchical (multi-level/ mixed)

648   regression models. , http://florianhartig.github.io/DHARMa/.

649 56. Bell WJ. 1990 Central place foraging. In *Searching Behaviour*, pp. 171–187. Springer,

650   Dordrecht. (doi:10.1007/978-94-011-3098-1_12)

651 57. Forcada J, Trathan PN, Reid K, Murphy EJ. 2005 The effects of global climate

652   variability in pup production of Antarctic fur seals. *Ecology* **86**, 2408–2417.

653   (doi:10.2307/3451030)

654 58. Gianuca D *et al.* 2019 Sex-specific effects of fisheries and climate on the demography

655   of sexually dimorphic seabirds. *J. Anim. Ecol.* **88**, 1366–1378. (doi:10.1111/1365-

656   2656.13009)

657 59. Ortiz RM, Wade CE, Ortiz CL. 2001 Effects of prolonged fasting on plasma cortisol

658   and TH in postweaned northern elephant seal pups. *Am. J. Physiol. - Regul. Integr.*

659   *Comp. Physiol.* **280**, R790–R795. (doi:10.1006/gcen.2000.7514)

660 60. Crossin GT, Trathan PN, Phillips RA, Gorman KB, Dawson A, Sakamoto KQ,

661   Williams TD. 2012 Corticosterone predicts foraging behavior and parental care in

662   macaroni penguins. *Am. Nat.* **180**, E31–E41. (doi:10.1086/666001)

663 61. Boyd IL, Lunn NJ, Barton T. 1991 Time budgets and foraging characteristics of

664   lactating Antarctic fur seals. *J. Anim. Ecol.* **60**, 577–592. (doi:10.2307/5299)

665   62.   Lunn NJ, Boyd IL, Barton T, Croxall JP. 1993 Factors affecting the growth rate and

666   mass at weaning of Antarctic fur seals at Bird Island, South Georgia. *J. Mammal.* **74**,

667   908–919. (doi:10.2307/1382429)

668   63.   Warren NL, Trathan PN, Forcada J, Fleming A, Jessopp MJ. 2006 Distribution of post-

669   weaning Antarctic fur seal *Arctocephalus gazella* pups at South Georgia. *Polar Biol.*

670   **29**, 179–188. (doi:10.1007/s00300-005-0037-x)

671 **Tables**

672 Table 1: Differences in AIC values (ΔAIC) between the full model and HMMs sequentially

673 excluding each variable and its interaction term with colony.

Covariate	Exclude	AIC	ΔAIC	
all		592365		676
sex	Interaction with colony	592368.9	3.9	677
	all terms	592366.3	1.3	678
age	interaction with colony	592443.7	78.7	679
	all terms	592534.6	169.6	680
body condition	interaction with colony	592370.4	5.4	681
	all terms	592386.5	21.5	682
time of day	interaction with colony	592377.7	12.7	683
	all terms	592535.4	170.4	684
season	interaction with colony	592408.9	43.9	685
	all terms	592418.4	53.4	686
temperature	interaction with colony	592367.5	2.5	687
	all terms	592367.8	2.8	688
wind speed	interaction with colony	592362.1	-2.9	689
	all terms	592360.1	-4.9	690
colony	all terms	592557.5	192.5	691

675 (header row)
692

Figures

Figure 1: VHF monitoring and GPS tracking. (a) VHF transmitters for pups (top) and mothers

(bottom) were temporarily attached to the dorsal side of the neck between the shoulder blades

with epoxy glue. GPS loggers in waterproof ABS plastic enclosures were similarly fitted to

pups. Tag specifications: Sirtrack core marine glue-on V2G 152A, 40 x 20 x 10 mm body

with 200 mm antenna, weight 16 g; Sirtrack core marine glue-on V2G 154C, 65 x 28 x 13

mm body with 250 mm antenna, weight 42 g; Perthold Engineering CatLog Gen2 GPS

Loggers, 50 x 40 x 17 mm, weight 36 g. (b) Antarctic fur seal mother-pup pair. VHF

transmitters are visible on both individuals, while the pup has additionally been fitted with a

GPS logger and given a temporary bleach mark for identification. Unrecovered tags and

bleach marks shed naturally with the pelage moult during March to April, precluding any

long-term consequences for the animals. Photo credit: Claire Stainfield.

Figure 2: Movement patterns of Antarctic fur seal pups on Bird Island, South Georgia (see

inset). (a) Location of the two fur seal breeding colonies from which pups were tagged with

GPS loggers. Freshwater Beach (FWB) and Special Study Beach (SSB) are separated by

approximately 200 meters. (b) Decoded states from the HMM showing the inferred activity

patterns of 66 pups throughout ontogeny. Pups tended to exhibit star-like movement patterns,

with directed movement away from and subsequent return to a central location of low activity.

(c) GPS locations recorded throughout ontogeny for 32 pups born on FWB and 34 pups born

on SSB. Animals from FWB stayed relatively close to their colony of birth, whereas SSB

pups often moved farther inland. To aid visibility, each panel is a magnified view of Bird

Island (longitude range = -38.060: -38.045, latitude range= -54.014: -54.005) and

consequently, some decoded states and GPS points recorded outside this frame are not visible.

For a complete map of the entire island showing all recorded states and positions, see

Supplementary Material Figure S1. For representative examples of individual GPS tracks,

including one pup from each breeding colony and season, see Supplementary Material Figure

S2.

Figure 3: (a–d) Mean probability (± 95% confidence interval) of being in an 'active'

behavioural state as a function of the covariates included in the HMM. Probabilities were

calculated for each covariate and state by fixing the other covariates to their respective means

or reference categories. Confidence intervals for the probabilities were obtained based on

Monte Carlo simulation from the estimators' approximate distribution as implied by

maximum likelihood theory. (a, b) Pups from both breeding colonies exhibit diurnal
movement patterns characterised by a higher probability of being active during the day. (c, d)
The probability of being in an active state gradually increased from birth to nutritional
independence for FWB pups, while SSB pups showed similar levels of activity throughout
ontogeny. (e, f) FWB pups moved away from their colony of birth and inland into the tussock
grass behind the beach habitat earlier, whereas pups at SSB tended to transition to the tussock
grass later in ontogeny.
735
Figure 4: Results of post-hoc analyses. (a) Pups were less active when their mothers were
present on the island; (b) Pups that died spent less time overall in the active state. Sample
sizes (n), GLMM slope estimates and p-values are provided.

Supplementary material

Figure S1: Map of Bird Island, South Georgia, showing all recorded GPS positions and states.

Figure S2: Representative examples of individual GPS tracks, including one pup from each

breading colony and season.

Figure S3: The number of states in an HMM must be specified before running the model and

only models with a biologically meaningful numbers of states should be tested. We expected

Antarctic fur seal pup movement to include a resting state (short step-lengths), an active state

(long step-lengths with more directed movement) and potentially a local exploration state

(moderate step-lengths with high degrees of turning). Thus, we evaluated HMMs with two

and three movement states, respectively. We also compared univariate HMMs to bivariate

HMMs that included the turning angle between consecutive GPS locations. The marginal

distribution under the fitted model and the empirical distribution is shown. In the 3-state

HMM, the states' distributions of both the step length and especially the turning angle

overlapped greatly and did not allow us to adequately distinguish the active from the local

exploration state. The distribution of the turning angle in the 2-state bivariate HMM was the

same for both states and therefore did not add any information to the model. Overall, neither

the additional state (three rather than two) nor the inclusion of the turning angle resulted in a

significant improvement of model fit. We therefore restricted our final analysis to a

parsimonious univariate HMM with two states.

Figure S4: HMMs fitted to 5-minute interval GPS data collected from three fur seal pups,

each between 30 – 50 days of age, for 13 – 19 days (mean = 3,965 data points). We evaluated

HMMs with two and three movement states. With respect to the aims of our study, there is no

relevant loss of information on pup activity by using the coarser temporal resolution (hourly

data).

Figure S5: Pseudo-residuals of the final HMM.

R Markdown

a

b

a

FWB

SSB

FWB
SSB
Inactive state
Active state

200 m

b

c