Trophic transfer of microplastics enhances tissue accumulation of chemical additives in fish

Takaaki Hasegawa[a], Kaoruko Mizukawa[b], Yeo Bee Geok[b], Tomonori Sekioka[c], Hideshige Takada[b], Masahiro Nakaoka[d,*]

[a] Fisheries Resources Institute, Japan Fisheries Research and Education Agency; Yokohama, Kanagawa, 236-8648, Japan

[b] Laboratory of Organic Geochemistry, Tokyo University of Agriculture and Technology; Fuchu, Tokyo, 183-8538, Japan.

[c] Graduate School of Environmental Science, Hokkaido University; Sapporo, Hokkaido, 183-8538, Japan.

[d] Field Science Center for Northern Biosphere, Hokkaido University; Akkeshi, Hokkaido, 088-1113, Japan.

*Corresponding author: thasegawa9@affrc.go.jp

Abstract

A variety of chemical additives are incorporated into plastics during their production

process to give desirable properties. Although studies have suggested that microplastic

ingestion can lead to accumulation of these chemicals in marine organisms, none provided the

direct evidence in fish species. Here, we demonstrated the tissue accumulation of chemical

additives in fish following microplastic ingestion. We exposed fish (*Myoxocephalus brandti*)

to polyethylene microplastics compounded with brominated flame retardants (BFRs; BDE209,

DBDPE) and ultraviolet stabilizers (UVs; UV-234, UV-327, BP-12) suspended in the water

column, or to mysids (*Neomysis* spp.) pre-exposed to the same microplastics. Our results

showed a maximum of 345-fold higher concentrations of additives in fish exposed to

microplastics than the ambient sample. Also, fish fed plastic-exposed mysids accumulated

significantly greater concentrations of BFRs in muscle than fish exposed to microplastics

suspended in the water column ($p < 0.001$). This indicates that trophic transfer of microplastics

has greater contribution to tissue accumulation of BFRs in fish than the waterborne ingestion.

In contrast, no significant difference in the accumulation of UVs was found between the

treatments, except for UV-327 in liver. These results suggests that the relative contribution

between direct ingestion of microplastics from the water column and indirect ingestion via

trophic transfer on tissue accumulation of additives in fish varies among compounds.

Compounds with high molecular weight and high hydrophobicity like BDE209 and DBDPE

were considered difficult to leach out from plastics, however, our study showed that trophic

transfer of microplastics can facilitate the accumulation of such compounds from plastics to

fish.

Keywords: Plastic ingestion, Brominated flame retardants, Ultraviolet stabilizers, *Neomysis*

spp. *Myoxocephalus brandti*

1. Introduction

As plastic production has been increasing over the years, plastic debris have become

ubiquitous pollutants found throughout marine environments (Auta et al., 2017; Burns and

Boxall, 2018; Desforges et al., 2014; Duis and Coors, 2016). Especially, plastic particles

smaller than 5 mm called microplastics are concerned as an emerging pollutant in marine

ecosystems (NOAA, 2020). Their small size allows a variety of organisms to ingest such as

whales, seabirds, fish, and invertebrates (Auta et al., 2017; Cole et al., 2013; Rummel et al.,

2016; Zhang et al., 2019). Many studies have shown that ingesting microplastics can cause

negative physical impacts on organisms such as blockage and laceration of the digestive tract

(Derraik, 2002; Khosrovyan et al., 2012; Wright et al., 2013). In most cases, however, these

negative effects were observed with unrealistically high exposure concentration in laboratory

experiments (Cunningham and Sigwart, 2019; Lenz et al., 2016), and the current monitored

concentration in the ocean is considered to have negligible impacts on organisms because

microplastics can be egested from the body without long-term accumulation (Beiras and

Schönemann, 2020; A. Dawson et al., 2018).

Aside from their physical effects, microplastics also have chemical effects on

organisms. Plastics contain an array of chemicals that are either adsorbed to the polymer

surface from the surrounding water or incorporated during the manufacturing process, the latter

called plastic additives (Andrady, 2011; Cole et al., 2011; Ziccardi et al., 2016). These

chemicals have been shown to transfer to biological tissues following the ingestion of

microplastics and are concerned about their impacts on ecosystems because some of them have

persistent and toxic natures (Hermabessiere et al., 2017a; Wardrop et al., 2016; Ziccardi et al.,

2016). However, some model studies suggested that plastics are not important for the

accumulation of these pollutants in organisms because their concentrations in plastics would

be at equilibrium with those in organisms and the chemical diffusion from plastics to organisms

would not occur (Gouin et al., 2011; Koelmans et al., 2016). In contrast, microplastics may

play a more relevant role in the transfer of chemical additives as they are added to plastic

polymers at high concentrations (~80% by weight (Gangi, 1999)), whereas the information

about the tissue accumulation of additives "compounded" in plastics is still far limited

compared to that of adsorbed pollutants. A recent study by Tanaka et al. (2020) demonstrated

that the ingestion of plastics resulted in the tissue accumulation of chemical additives in seabird.

While no study has so far provided the direct evidence that microplastic ingestion can lead to

the tissue accumulation of chemicals additives in fish species, Coffin et al. (2019) demonstrated

that *in vivo* fish gut simulated condition using the digestive enzyme enhanced the leaching of

additives from plastics, which suggests that tissue accumulation in fish species is plausible.

Studies using fish species are especially important considering the food web structure of marine

ecosystems and the potential impact on human health through direct consumption (Campanale

et al., 2020; Huang et al., 2020).

81   There are two possible pathways for predatory fish to ingest microplastics, either from

the water column or their prey (Walkinshaw et al., 2020). Most microplastic exposure studies

focus either on waterborne or dietary exposure, but none of them compared their relative

contribution to the total ingestion. Our prior work compared the waterborne ingestion of

microplastics and the trophic transfer using fish and mysids, and revealed that the trophic

transfer is more important than the waterborne ingestion (Hasegawa and Nakaoka, 2021). If

microplastic ingestion by fish leads to tissue transfer of plastic-derived chemicals, fish may

receive more impacts on the exposure of those chemicals via trophic transfer of microplastics

than the waterborne exposure.

90   The present study aims to examine the tissue accumulation of chemical additives in

fish following microplastic exposure and the relative contribution of the waterborne and trophic

transfer exposure on the accumulation. We used a crustacean mysid (*Neomysis* spp.) and a

benthic fish (*Myoxocephalus brandti*) as a model prey-predator system that was also used in

our prior work (Hasegawa and Nakaoka, 2021). We conducted the exposure experiments using

polyethylene microplastics compounded with two brominated flame retardants (BFRs;

BDE209, DBDPE) and three ultraviolet stabilizers (UVs; UV-234, UV327, BP-12). To

examine the relative importance of the two exposure pathways, we exposed the fish to

polyethylene microplastics suspended in the water column or to mysids pre-exposed to

polyethylene microplastics, and analyzed the concentration of each additive in fish muscle and

liver. We hypothesized that the fish would accumulate the additives following microplastic

ingestion and that trophic transfer of microplastics would yield the greater accumulation of the

additives than the waterborne ingestion.

2. Materials & Methods

2.1 Animal collection and acclimation

We collected organisms for the experiment on September 7, 2020 at a seagrass bed in

Akkeshi-ko estuary (43°02′ N, 144°52′ E), which is located in eastern Hokkaido, northern

Japan. Mysids (*Neomysis* spp., body length: 7.70 ± 1.53 mm standard deviation, wet weight:

8.20 ± 3.59 mg standard deviation) and juvenile *M. brandti* (body length: 7.40 ± 0.52 cm

standard deviation, wet weight: 4.35 ± 0.81 g standard deviation) were collected using an

epibenthic sled. For the measurement of contaminant concentrations in the ambient

environment, 90 mysids and three fish were immediately frozen and stored until the later

chemical analysis, and the other fresh animals were brought to the laboratory for the

experiments. Mysids were acclimated in 30-L aquaria overnight with flow-through seawater

that were filtered by fine sand to allow them clear possible microplastics from their guts. Ten

glass aquaria (245×165×160 mm) were prepared and one randomly selected fish was assigned

to each aquarium, and then acclimated for 7 days prior to the start of the experiment. During

the acclimation, we daily fed the mysids microalgae (Shellfish diet 1800; Reed Mariculture)

and the fish fresh mysids. Each aquarium contained approximately 100 mysids or 20 fish.

2.2 Preparation of plastics with additives

We used cylindrical low-density polyethylene pellets (diameter 5 mm, length 5 mm;

DJK Corporation, Chiba, Japan) compounded with five plastic additives. Polyethylene is one

of the most produced plastic polymers and is commonly found in marine environments

(Andrady, 2011; Beiras et al., 2018; Burns and Boxall, 2018). Following five additives were

industrially compounded in the polyethylene pellets: 1,2,3,4,5-pentabromo-6-(2,3,4,5,6-

pentabromophenoxy)benzene (BDE209, CAS no. 1163-19-5), 1,2,3,4,5-pentabromo-6-[2-

(2,3,4,5,6-pentabromophenyl)ethyl]benzene (DBDPE, CAS no. 84852-53-9), 2-(benzotriazol-

2-yl)-4,6-bis(2-phenylpropan-2-yl)phenol (UV-234, CAS no.70321-86-7), 2,4-ditert-butyl-6-

(5-chlorobenzotriazol-2-yl)phenol (UV-327, CAS no. 3864-99-1), and (2-hydroxy-4-

octoxyphenyl)-phenylmethanone (BP-12, CAS no. 1843-05-6). BDE209 and DBDPE are

widely used in plastic products and electrical appliances as flame retardants and are suspected

to be non-readily degradable and have long-term toxicity (Hermabessiere et al., 2017b; NITE,

2017). UV-234, UV-324 and BP-12 are also commonly added to plastic products as UV-

stabilizers, which have the potential for bioaccumulation and toxicity on organisms (Kim et al.,

2011; Nakata et al., 2009). These compounds were selected as previous studies reported their

frequent occurrences in plastic debris collected in natural environments and marine organisms

(Rani et al., 2015; Rochman et al., 2014; Tanaka et al., 2019). The concentration of each

additive in the pellets was set below 0.04 % by weight, which is of the same order of magnitude

as those observed in these studies. Polyethylene powder (Flo-Thene, FG701N, Sumitomo Seika

Chemicals Co., Ltd., Osaka, Japan) and authentic standards of the additives in powder form

(Wako Pure Chemical Industries, Ltd., Osaka, Japan) were mixed well and molded into pellets

in a co-rotating twin-screw kneading extruder (HK-25D; Parker Corporation, Inc., Tokyo,

Japan). The pellets were then melted and re-extruded twice to obtain a uniform distribution of

the constituents. The pellets were further grinded into powders by a plastic cutting mill (PLC-

2M, Osaka Chemical Co., Ltd., Osaka, Japan) and the powders were sieved over a 63 μm and

32 μm mesh. Particles left on the 32 μm sieve were collected and used for the experiments.

For measurement of the particle size, the quantification method with nile red was

adapted from Erni-Cassola et al. (2017) using a nylon mesh filter with 8 μm pore size. Particle

area was measured, and the square root was calculated as particle size. The targeted

polyethylene microplastics had an average size of 29.90 ± 13.40 μm and their size distribution

is shown in Fig.1. The concentrations of five additives in polyethylene microplastics were

quantified (Table 1) and the composition of each additive is shown in Fig. 2.

To prepare the microplastic-suspended solution, a glass beaker was filled with 100 mL

of distilled water and boiled in a microwave. Ten microliters of surfactant (Tween 80;

polyethylene sorbitol ester, Cospheric LLC) were added and stirred with a glass rod for 30 s.

Next, 5 mL of the solution was transferred to another grass beaker filled with 95 mL of boiled

distilled water to make 0.005 % surfactant solution. The solution was left at room temperature

for 1 h. Before starting the exposure experiment, 8 mg of polyethylene microplastics were

added to a plastic tube containing 8 mL of 0.005 % surfactant solution. The tube was vortexed

and left at room temperature for 1 h to immerse into the solution. The suspension solution was

prepared each time for the experiments to minimize the leaching of the chemicals from

polyethylene microplastics during the immersion process.

2.3 Leaching experiment

To determine the amount of chemicals leaching from microplastics to seawater, three

beakers filled with 1-L filtered seawater (< 1 μm) containing 2–mg/L microplastics were

prepared and left for 24 h. We constantly circulated the seawater by aeration during the test.

After 24 h, the seawater was transferred to a 1-L stainless container and stored in the freezer

until the chemical analysis.

2.4 Ingestion experiment for mysids

Thirty 1-L glass beakers were prepared and divided into the following two groups;

fifteen bottles filled with 1-L filtered seawater (< 1 μm) containing 500-μg/L microplastics,

and 15 bottles filled with 1-L leachate seawater. The microplastic concentration was chosen as

an environmentally relevant concentration as it is within the same order of magnitude of the

microplastic concentration reported in the North Pacific Subtropical Gyre, which is among the

most heavily contaminated area around the world (Goldstein et al., 2012). Leachate seawater

was prepared by filtering 1-L filtered seawater (< 1 μm) containing 500-μg/L microplastics to

determine the uptake of the chemicals leaching from microplastics during the experiment by

mysids. We randomly selected mysids from the acclimation aquaria and placed three in each

bottle. We then fed the mysids 2 mg (dry weight) of microalgae. We constantly circulated the

seawater by aeration to ensure that the microplastics were evenly distributed in the bottles.

After 24 h of microplastic exposure, the mysids were flushed gently with filtered seawater to

remove beads from the exoskeleton and their body length and wet weight were measured. Then,

the mysids were frozen at - 30℃ and stored until the later analysis. The same procedure was

repeated 4 days to collect 150 mysids in total from each group except that only 15 mysids in

each group were used at 4th day.

2.5 Trophic transfer experiment

After the acclimation for fish, ten fish aquaria were divided into the following two

groups: Five aquaria contained 2-L filtered seawater (< 1 μm) with a 500-μg/L microplastic

suspension and nine plastic-free mysids each, so that the fish would ingest polyethylene

microplastics only from the water column. Five other aquaria each contained nine mysids that

had been pre-exposed to 500-µg/L microplastics. The five aquaria containing pre-exposed

mysids were filled with 2-L filtered seawater (1 µm) without a microplastic suspension so that

uptake of polyethylene microplastics would be solely from the food source. The number of

mysids was determined from a preliminary experiment to ensure that all mysids were consumed

by the fish within 24 h. Pre-exposure of the mysids followed the protocol for the first

experiment. We carried out the exposure for 10 days and exchanged the seawater once a day.

We constantly circulated the seawater by aeration throughout the experiment to ensure that

microplastics were evenly distributed in the aquaria. We did not observe any microplastics on

the bottom of the tanks throughout the experiment. After 10 days of exposure, the fish were

gently flushed with filtered seawater to remove beads from the body surface. Following

measurement of their body length and wet weight, the fish were frozen at - 30°C and stored

until the later analysis.

2.6 Sample analysis

2.6.1 Chemical and reagents

Acetone, n-hexane, 2,2,4-trimethylpentane (iso-octane), methanol (MeOH), pyridine,

acetic anhydride, hydrochloric acid, anhydrous sodium sulfate, Wakogel Q-22 (75 µm, 200

mesh) and Wakogel Q-23 (75-150 µm, 100–200 mesh) were purchased from Wako Pure

Chemical Ltd. (Osaka, Japan). Dichloromethane (DCM) were purchased from Kanto Chemical

Co., Inc. (Tokyo, Japan). Analytical standards for BDE209 and DBDPE were purchased from

Wellington Laboratories Inc. (Guelph, Ontario, Canada). Hexa-nona BDE individual congeners

were obtained from Wellington Laboratories Inc. (BDE179, 183, 188 and 202), Cambridge

Isotope Laboratories Inc. (BDE207; Tewksbury, MA, US) and Accustandard Inc. (BDE155,

184, 196, 197, 203, 206 and 208; New Haven, CT, US) respectively. UV327 and BP12 were

also obtained from Accustandard Inc. Analytical surrogates were used the followings; f-

BDE208   (4'-fluoro-2,2',3,3',4,5,5',6,6'-nonabromo-diphenylether)   from   Chiron   AS

(Trondheim, Norway) for BFRs, UV327-d20 were from Toronto Research Chemicals, Inc.

(North York, ON Canada) for UV327 and UV234 and BP12-d17 were from Hayashi Pure

Chemical Ind., Ltd. (Osaka, Japan) for BP12. As internal injection standard (IIS), chrysene-

d12 was purchased from Sigma–Aldrich (St. Louis, MO, USA). All glassware was rinsed with

methanol, acetone, and distilled n-hexane three times respectively or pre-baked at 550 ℃ for 4

hours.

2.6.2   Biological samples

For chemical analysis, the liver and whole-body (mainly muscle tissues, excluding the

intestines, head and fins) were used. Each freeze-dried sample was extracted by accelerated

solvent extractor (ASE200; Dionex, Sunnyvale, CA, USA) with acetone/DCM (1:3, v/v). The

samples were extracted in 11 mL stainless steel cells under the following conditions: preheat

time, 0 min; heat time, 5 min; static time, 5 min; flush%, 100%; purge, 60 s; 2 cycles; pressure,

1500 psi; temperature, 100 ℃.

To determine lipid contents of whole-body sample from the weight of the dried

extracts, aliquot of the extract was taken into another weighed-4 mL vial. After injection the

surrogates (f-BDE208, UV327-d20 and BP12-d17) into another aliquot, it was concentrated

just to dryness in a rotary evaporator and transferred into a 10 mL centrifugation tube. After

reducing the solvent less than 100 mL by gentle nitrogen stream, we added 50 mL each of

pyridine and acetic anhydride into each tube. After holding at room temperature for >8 h, the

reaction was stopped with the addition of 0.2 mL 4 M HCl. Acetylates were extracted with n-

hexane 5 times. The n-hexane extract was passed through anhydrous sodium sulfate for

dehydration and collected into pair-shaped flask.

After a solvent substitution from n-hexane to DCM, the acetylated sample was purified

by gel permeation chromatography (GPC). About 2 mL of sample was subjected to GPC (2 cm

i.d. ×30 cm, CLNpak EV-2000; Showa-denko, Tokyo, Japan) to separate target compounds

from biolipids in DCM at an eluent at flow rate of 4 mL/min. The fraction with a retention time

of 12 to 25 min was collected for further purification.

The GPC eluent was concentrated just to dryness in a rotary evaporator and placed in

a 10% H2O-deactivated silica gel column (Wakogel Q-22, 1 cm i.d. × 9 cm). The first fraction

including BFRs was eluted with 35 mL n-hexane/DCM (3:1, v/v). The second and third fraction

including UVs were eluted with 15 mL MeOH/DCM (1:99, v/v) and 20 mL (1%

MeOH/DCM)/Acetone (1:4, v/v) respectively. The second and third fraction were combined

and evaporated the solvents and transferred into a 1 mL amber ampoule for instrumental

analysis.

The first eluent containing BFRs was concentrated just to dryness in a rotary

evaporator and placed in a fully activated silica gel column (Wakogel Q-23, 0.47 cm i.d. ×18

cm). The first fraction, containing aliphatic hydrocarbons, was eluted with 5 mL n-hexane. The

second fraction, containing BFRs was eluted with 15 mL n-hexane/DCM (85:15, v/v) or 10 mL

n-hexane/DCM (3:1, v/v). The BFR fraction was concentrated and transferred into a 1 mL

amber ampoule for instrumental analyses.

2.6.3　Water samples

To analyze the water samples from the leaching experiment, the water was filtered

through a glass fiber filter (GF/F), and f-BDE208, UV327-d20, and BP12-d17 dissolved in

acetone were added to the filtrate as surrogates. Then, liquid-liquid extraction was carried out

with dichloromethane (DCM) at 10% of the volume of the water sample, and the organic phase

was collected while dehydrating with anhydrous sodium sulfate. This process was repeated

three times. The extracts were concentrated using a rotary evaporator and fractionated and

purified using acetylation and 10 % water-inactivated silica gel column chromatography

(Wakogel Q-22) as with the biological samples.

2.6.4 Instrumental analysis

For analysis of BFRs, the solvent was evaporated just to dryness under a nitrogen

stream, and the residue was redissolved in 100 mL iso-octane.

UVs were analyzed on a 5977 quadrupole mass spectrometer fitted with a 7890 gas

chromatograph (GC–MS; Agilent technologies, Inc., Santa Clara, CA, US). An Agilent HP-

5MS 30-m fused silica capillary column (0.25 mm i.d., 0.25 mm film thickness) was used with

helium as the carrier gas at 100 kPa. GC–MS operating conditions were set at 70 eV ionization

potential with the source at 240 °C and the electron multiplier voltage at ~2000 eV. The

injection port was maintained at 300 °C, and the sample was injected in splitless mode, then

purged 2 min after injection. The column temperature was held at 100 °C for 1 min, rose at

30 °C/min to 160 °C, at 10 °C/min to 310 °C and was then held for 10 min. After a solvent

delay of 4 min, selected ion monitoring mode was used. We monitored UV327 at m/z = 342

UV234 at 432, BP-12 at 213. Surrogate standards and IIS were monitored at m/z = 359

(UV327-d20), m/z = 343 (BP12-d17), at m/z = 240 (chrysene-d12) respectively. These were

quantified by comparing the integrated peak area of the quantification ion with that of the IIS

as derived from calibration lines drawn for individual UVs.

BFRs were determined by a micro electron capture detector fitted with a HP 7890 gas

chromatograph (GC-mECD, Agilent Technologies Inc.). An Agilent DB-5 15-m fused silica

capillary column (0.25 mm i.d., 0.25 mm film thickness) was used with helium as the carrier

gas. To separate BDE208 and f-BDE208 clearly and gain higher peaks of BDE209 and DBDPE,

we used ramped pressure mode for the analysis. The column pressure was held at 20.19 psi for

31 min, rose at 15 psi/min to 35 psi, and was then held for 6 min. The injection port was

maintained at 250 °C, and the sample was injected in splitless mode, then purged 0.75 min after

injection. The column temperature was held at 80 °C for 2 min, rose at 30 °C/min to 240 °C, at

3 °C/min to 300 °C and was then held for 15 min. BFRs were identified and quantified against

the standards.

2.6.5   Analytical quality control and quality assurance

A procedural blank was run in every set analyzed. Analytical values less than three

times of the corresponding blank are expressed as "below the limit of quantification", <LOQ.

To certificate analytical quality control and quality assurance, quadruplicate mussel samples

which were spiked the standard solution of target compounds were analyzed. The relative

standard deviations of the concentrations of BDE209, DBDPE, UV327, UV234 and BP12 were

2%, 2%, 2%, 10% and 12% respectively. The averaged recoveries of BDE209, DBDPE,

UV327, UV234, and BP12 were 90%, 114%, 92%, 100%, and 83% respectively.

2.7 Statistical Analysis

All analysis were conducted in R (R Core Team, 2020). To compare the differences in

additive accumulation by mysids, fish muscle, and fish liver among treatments, we used

generalized linear models (GLMs) with log link functions. A gamma distribution was assumed

to account for the positive continuous values because the Shapiro-Wilk normality tests showed

that the response variable had a non-normal distribution. We used log-transformed sample dry

weight as an offset in the models. To test the effect of treatment, we performed a likelihood

ratio test using the Anova function in the "car" package (Fox and Weisberg, 2019). For pairwise

comparisons, we used Tukey's HSD test as post-hoc analysis.

3. Results

3.1 Leaching experiment

Overall, 2.33% of the amount of additives were leached out from polyethylene

microplastics for 24 h. The leaching rate varied among different additives (Table 2) with the

mean of 3.16 %. BP-12 showed the higher leaching rate (8.79 %) than other additives (<3 %).

3.2 Ingestion experiment for mysids

Concentration of five additives accumulated in mysids significantly differed among

treatments (Table 3). Mysids contained approximately 4-2450 times higher concentrations of

additives by ingesting microplastics than those from the ambient environment (Fig. 3). Mysids

treated with the leachate seawater also showed slightly elevated concentrations of additives

than those from the ambient environment. For BP-12, the concentration in mysids treated with

the leachate seawater was not significantly different from those exposed to microplastics.

3.3 Trophic transfer experiment

Concentration of the five additives in fish varied significantly among treatments

(Table 5) although the patterns of variation greatly differed among additives and between body

tissues (Fig. 4).

For BFRs, the treatment had significant effects on their concentrations both in muscle

and liver. In muscle, fish fed plastic-exposed mysids showed the higher concentrations than

those exposed to microplastics suspended in the water column. The mean effect was greater in

BDE209 (71-fold on dry weight basis, 161-fold on lipid weight basis) than DBDPE (22-fold

on dry weight basis, 52-fold on lipid weight basis). Although the lipid weight basis analysis

showed the significantly higher concentration in the water treatment than the ambient control,

no significant differences were observed between them on dry weight basis. Notably, the

within-treatment variation in the mysid treatment was also greater than the other two treatments.

In contrast, liver showed no significant difference in the concentration of BDE209 between the

mysid and water treatments although they were more than 3 fold greater than the ambient

control. The same patterns of significant variation was also found for DBDPE except that the

concentration in liver in the mysid treatment was not different from the ambient controls. Liver

showed different accumulation patterns of BFRs between the treatments. In the water treatment,

the mean concentrations were higher in liver (BDE209: 5.83 ng/g-dry, DBDPE: 9.79 ng/g-dry)

than muscle (BDE209: 0.47 ng/g-dry, DBDPE: 0.60 ng/g-dry) while the mysid treatment

showed the lower concentrations in liver (BDE209: 3.99 ng/g-dry, DBDPE: 2.52 ng/g-dry) than

muscle (BDE209: 4.30 ng/g-dry, DBDPE: 5.67 ng/g-dry).

For UVs, the treatment had significant effects on their concentration except for UV-

234 in muscle where no treatment effect was observed. In muscle, UV-327 and BP-12 showed

3-5 fold higher concentrations in the treatment groups than the ambient control (7-12 fold on

lipid weight basis). Unlike BDE209 and DBDPE, however, no significant differences were

found between the mysid and water treatments. In liver, UV-234 and UV-327 respectively

showed 9-13 fold and 35-89 fold higher concentration in the treatment groups than the ambient

controls, and also the concentration of UV-327 in the mysid treatment was significantly higher

than that in the water treatment. While BP-12 showed 3-fold higher concentration in the mysid

treatment than the ambient control, the concentration in the water treatment was not

significantly different from both the mysid treatment and the ambient control. In general, liver

showed the higher concentrations of UVs in liver than muscle in both treatments, whereas the

accumulation tendency between the tissues greatly differed among the chemicals. For example,

UV-234 showed the greatest increase in liver compared to muscle among UVs (Water: 29-fold,

Mysid: 53-fold), whereas BP-12 only showed 7-fold increase in the water treatment and 13-

fold increase in the mysid treatment.

4. Discussion

This study provides the first evidence that microplastic ingestion leads to the tissue

accumulation of chemical additives in fish. We also revealed that trophic transfer of

microplastics enhanced the accumulation of BFRs in fish muscle compared to the waterborne

ingestion. However, as we observed the equal contribution to the accumulation of UVs between

direct ingestion of microplastics from the water column and indirect ingestion via trophic

transfer, their relative contribution varies among additives.

Previous studies showed that microplastics can act as a vector for the transfer of

plastic-associated chemicals to marine organisms, but mostly targeted at the environmental

contaminants adsorbed onto plastics, such as PCBs and PAHs (Besseling et al., 2013; Rochman

et al., 2013). However, some model prediction studies pointed out that microplastics would not

increase the body burden of these contaminants in organisms because other input sources like

water and food would play a more important role in their accumulation (Bakir et al., 2016;

Gouin et al., 2011; Koelmans et al., 2016). In contrast, plastic additives are different from

pollutants adsorbed to plastics with respect that they are not attached to the polymer surface,

but incorporated into plastics at high concentrations (Gangi, 1999). Several field observation

studies found a positive relationship between plastic ingestion and chemical additives detected

from marine fish and seabirds (Gassel et al., 2013; Tanaka et al., 2013), suggesting the potential

occurrence of the tissue transfer of plastic-derived additives in natural environments. Our

results support these evidence along with the previous finding by Tanaka et al. (2020) where

they demonstrated that plastic ingestion led to the tissue accumulation of additives in seabird

(*Calonectris leucomelas*) by the laboratory experiment. Gut fluid in both seabird and fish have

been shown to play a key role in the leaching of chemicals from plastics (Coffin et al., 2019;

Tanaka et al., 2015), which explains the tissue accumulation of chemical additives following

plastic ingestion. Given the generality of plastic ingestion in fish and detection of additives in

plastics found in natural environments (Granek et al., 2020; Wang et al., 2020), our study raises

a concern about the exposure of these additives to fish.

We found that trophic transfer of microplastics has greater effects on the accumulation

of BDE209 and DBDPE in fish than waterborne ingestion. There are three possible

explanations for this result which are not exclusive. First, fish was exposed to more chemical

additives by ingesting microplastics via trophic transfer. Our previous study demonstrated that

fish ingested greater amount of microplastics via trophic transfer than the waterborne ingestion

(Hasegawa and Nakaoka, 2021). Higher ingestion rate of microplastics via trophic transfer

should have resulted in the greater exposure intensity of chemical additives in fish stomach.

Secondly, greater secretion of gut fluid may have elevated the leaching rate of additives from

plastics in stomach. Gut fluid plays a key role in leaching hydrophobic chemicals from ingested

plastics (Coffin et al., 2019; Tanaka et al., 2015). Digesting plastic-contaminated prey could

facilitated the secretion of gut fluid in stomach, which may have enhanced the leaching of

chemicals from plastics. Thirdly, mysids fragmented microplastics into smaller particles and

helped the release of chemicals from plastics. Some crustacean species can fragment plastics

into smaller particles, including mysids (A. L. Dawson et al., 2018; Hasegawa and Nakaoka,

2021; Mateos-Cárdenas et al., 2020). Generally, smaller plastics can release chemicals more

easily due to their larger surface-to-volume ratio (Wright and Kelly, 2017). Ingesting

microplastics from mysids, therefore, changed the size composition of plastics in fish stomach,

and may have enhanced the leaching of chemicals. Compounds with high molecular weight

and high hydrophobicity such as BDE209 and DBDPE (Table 1) were considered difficult to

leach out from plastics and to transfer into biological tissues (Braekevelt et al., 2003; Cheng et

al., 2020; Sun et al., 2019). However, we demonstrated the elevated accumulation of these

compounds in the fish by ingesting microplastics from mysids. Thus, our study highlights the

importance of indirect exposure via trophic transfer in chemical effects of microplastics on

organisms.

421   Although we observed the greater accumulation of BDE209 and DBDPE via trophic

422 transfer of microplastics, the waterborne exposure equally contributed the accumulation of UV-

423 327, UV-234 and BP-12. This suggests that the trend in accumulation of chemical additives

424 vary depending both on the properties of the chemicals and on exposure pathway. Given the

425 ingestion of microplastics from the water column in fish are much lower than the trophic

426 transfer (Hasegawa and Nakaoka, 2021), there could be other accumulation routes in the

427 waterborne exposure of chemical additives. In marine fish, there are three possible routes for

428 chemical uptake from the environment: gill uptake, dermal absorption and gut absorption by

429 drinking water (Qiao et al., 2000). Although microplastic ingestion is assumed to only

430 contribute to gut absorption, gill uptake could also be important because microplastics can also

431 accumulate in gill. Water flux in the gill of marine benthic fish *Oligocottus maculosus*, which

432 shares the same family of our target fish *M. brandti*, by ventilation is 300-fold higher than that

433 in stomach by drinking water (Somo et al., 2020; Webb et al., 2001). Also, UV-234, UV-324

434 and BP-12 are likely to leach out from plastics more easily than BDE209 and DBDPE due to

435 their lower molecular weight and less hydrophobic nature (Table 1). Considering the more

436 frequent contact with microplastics and the lower hydrophobic nature, gill may have also

437 played a role in the transfer of the UVs into fish body. Given that mysids accumulated additives

438 leached out from plastics to the water column, fish was also likely to accumulate the additives

439 from the water column in the same manner, especially for BP-12 that showed the higher

440 leaching rate (10%) and the higher accumulation in mysids. Although we need further

441 investigation on the mechanism that differentiates the behavior of additives depending on each

442 exposure pathway, both waterborne exposure and trophic transfer of microplastics are

443 important for the accumulation of plastic additives in fish.

444   Overall, we observed the preferential accumulation of the chemicals in liver over

445 muscle, which is consistent with the previous findings targeted at the same class of compounds

446 (Nakata et al. 2009; Tanaka et al. 2020). However, the effect size was much smaller in BFRs

(0.4-16 fold) than UVs (7-53 fold). Tanaka et al. (2020) observed the totally opposite trend in

seabird that showed the considerably high level of accumulation for BDE209, but less for UV-

327 and BP-12 in the liver. This indicates the species-specific accumulation patterns even in

the same compounds. Tanaka et al. (2020) explained the lower accumulation of BP-12 by the

degradation in liver due to the susceptible nature of BP-12 to the metabolization. In contrast,

BDE209 is known to be resilient to biodegradation because of their high molecular weight. As

several studies showed that BDE209 and DBDPE can be readily metabolized to lower

brominated congeners in fish species (Chen et al., 2012; X. Wang et al., 2019), lower

accumulation level for these two compounds could be explained by the metabolization. Still,

we cannot fully exclude this possibility without conducting a depuration experiment to examine

in vivo kinetics of these chemicals. Future studies should investigate the temporal variation of

chemical concentrations in different tissues to understand the fate of plastic additives in

organisms.

Studies have demonstrated that both BFRs and UVs have negative impacts on

organisms (Kim et al., 2011; Morohoshi et al., 2005; X. Wang et al., 2019). Especially,

extensive research have been conducted for BDE209 and have shown that the compound can

act as an endocrine disrupter and cause neurotoxicity on fish (Sun et al., 2020; X. Wang et al.,

2019; Y. Wang et al., 2019). BDE209 is also concerned for the accumulation in a human body

because of the potential adverse health effects (Darnerud, 2008). As global seafood

consumption has been increasing over the years, humans may be more frequently exposed to

such additives through seafood consumption. A simple risk assessment of plastic additives on

humans through consumptions of fish can be made as follows. First, BDE209 accumulation in

fish, dairy intake of BDE209 through seafood consumption was estimated based on the result

of BDE209 concentration in fish muscle. Assuming annual seafood consumption of 24.6 kg

year^{-1} and the average body weight of 60 kg for adult Japanese (Ministry of Agriculture, 2016;

Ministry of Health, 2018) and using our observed data on the mean BDE209 concentration of

4.70 ng/g.wet weight in fish muscle considering the both microplastic exposure pathways, dairy

intake of BDE209 can be estimated as 0.46 ng/kg/day. According to the report from National

Institute of Technology and Evaluation, lowest observed adverse effects levels (LOAEL) of

BDE209 is estimated as 50 ng/kg/day to induce chronic toxicity for a human (NITE, 2017),

which is approximately 110-fold higher than the estimated dairy intake value through seafood

consumption. As our study used the environmentally relevant concentration of additives,

accumulation of plastic additives in fish through microplastic exposure should be negligible

for a human body in the current plastic pollution level of the world's ocean. However, current

study only shows the marginal results with 10-day microplastic exposure to fish. Considering

a long-term exposure of microplastics to fish throughout the life, fish would accumulate far

more plastic-derived additives. A study predicted the abundance of plastics in the ocean would

increase 50-fold by 2100 from the present condition (Everaert et al., 2018), suggesting that

plastic exposure through seafood consumption would provide a relevant pathway for BDE209

to have adverse health effects on a human in the future scenario. We are also being exposed to

these additives from the daily use of plastic products. The risk assessment study estimated a

daily exposure amount of BDE209 to humans from different exposure routes such as inhalation

of inside dusts or elution from plastic products, but did not include the exposure through

seafood consumption (NITE, 2017). This study clearly showed the chemical accumulation in

fish following microplastic exposure. Thus, these risk assessment studies should consider

seafood consumption as a factor for the exposure of plastic-derived chemicals to a human body.

Conclusion

This study is the first to demonstrate that microplastic ingestion can lead to tissue

accumulation of chemical additives in fish. We also revealed that indirect exposure of

microplastics via trophic transfer can enhance the accumulation of BDE209 and DBDPE that

were assumed difficult to accumulate in organisms because of their low leachability from

plastics. Some studies have suggested that microplastics themselves do not accumulate in the

body and have negligible effects on organisms because they can be excreted from the body.

Whereas, our findings indicate that microplastics play an important role in chemical

bioaccumulation. As many chemical additives are potentially toxic, this study raises the

concern about their negative impacts on marine ecosystems. To better understand the fates and

the effects of chemical additives accumulated in organisms, further research should examine

their *in vivo* kinetics and toxicity.

References

Andrady, A.L., 2011. Microplastics in the marine environment. Mar. Pollut. Bull. 62, 1596–

1605. https://doi.org/10.1016/j.marpolbul.2011.05.030

Auta, H.S., Emenike, C.U., Fauziah, S.H., 2017. Distribution and importance of microplastics

in the marine environment: A review of the sources, fate, effects, and potential solutions.

Environ. Int. 102, 165–176. https://doi.org/10.1016/j.envint.2017.02.013

Bakir, A., O'Connor, I.A., Rowland, S.J., Hendriks, A.J., Thompson, R.C., 2016. Relative

importance of microplastics as a pathway for the transfer of hydrophobic organic

chemicals to marine life. Environ. Pollut. 219, 56–65.

https://doi.org/10.1016/j.envpol.2016.09.046

Beiras, R., Bellas, J., Cachot, J., Cormier, B., Cousin, X., Engwall, M., Gambardella, C.,

Garaventa, F., Keiter, S., Le Bihanic, F., López-Ibáñez, S., Piazza, V., Rial, D., Tato, T.,

Vidal-Liñán, L., 2018. Ingestion and contact with polyethylene microplastics does not

cause acute toxicity on marine zooplankton. J. Hazard. Mater. 360, 452–460.

https://doi.org/10.1016/j.jhazmat.2018.07.101

Beiras, R., Schönemann, A.M., 2020. Currently monitored microplastics pose negligible

ecological risk to the global ocean. Sci. Rep. 10, 1–9. https://doi.org/10.1038/s41598-

020-79304-z

Besseling, E., Wegner, A., Foekema, E.M., Van Den Heuvel-Greve, M.J., Koelmans, A.A.,

2013. Effects of microplastic on fitness and PCB bioaccumulation by the lugworm

Arenicola marina (L.). Environ. Sci. Technol. 47, 593–600.

https://doi.org/10.1021/es302763x

Braekevelt, E., Tittlemier, S.A., Tomy, G.T., 2003. Direct measurement of octanol-water

partition coefficients of some environmentally relevant brominated diphenyl ether

congeners. Chemosphere 51, 563–567. https://doi.org/10.1016/S0045-6535(02)00841-X

Burns, E.E., Boxall, A.B.A., 2018. Microplastics in the aquatic environment: Evidence for or

against adverse impacts and major knowledge gaps. Environ. Toxicol. Chem. 37, 2776–

2796. https://doi.org/10.1002/etc.4268

Campanale, C., Massarelli, C., Savino, I., Locaputo, V., 2020. A detailed review study on

potential effects of microplastics and additives of concern on human health.

Chen, Q., Yu, L., Yang, L., Zhou, B., 2012. Bioconcentration and metabolism of

decabromodiphenyl ether (BDE-209) result in thyroid endocrine disruption in zebrafish

larvae. Aquat. Toxicol. 110–111, 141–148.

https://doi.org/10.1016/j.aquatox.2012.01.008

Cheng, H., Luo, H., Hu, Y., Tao, S., 2020. Release kinetics as a key linkage between the

occurrence of flame retardants in microplastics and their risk to the environment and

ecosystem : A critical review. Water Res. 185, 116253.

https://doi.org/10.1016/j.watres.2020.116253

Coffin, S., Huang, G.Y., Lee, I., Schlenk, D., 2019. Fish and seabird gut conditions enhance

desorption of estrogenic chemicals from commonly-ingested plastic items. Environ. Sci.

Technol. 53, 4588–4599. https://doi.org/10.1021/acs.est.8b07140

Cole, M., Lindeque, P., Fileman, E., Halsband, C., Goodhead, R., Moger, J., Galloway, T.S.,

2013. Microplastic ingestion by zooplankton. Environ. Sci. Technol. 47, 6646–6655.

https://doi.org/10.1021/es400663f

Cole, M., Lindeque, P., Halsband, C., Galloway, T.S., 2011. Microplastics as contaminants in

the marine environment: A review. Mar. Pollut. Bull. 62, 2588–2597.

https://doi.org/10.1016/j.marpolbul.2011.09.025

Cunningham, E.M., Sigwart, J.D., 2019. Environmentally accurate microplastic levels and

their absence from exposure studies. Integr. Comp. Biol. 59, 1485–1496.

https://doi.org/10.1093/icb/icz068

Darnerud, P.O., 2008. Brominated flame retardants as possible endocrine disrupters 152–160.

https://doi.org/10.1111/j.1365-2605.2008.00869.x

Dawson, A., Huston, W., Kawaguchi, S., King, C., Cropp, R., Wild, S., Eisenmann, P.,

Townsend, K., Bengtson Nash, S.M., 2018. Uptake and depuration kinetics influence

microplastic bioaccumulation and toxicity in antarctic krill (*Euphausia superba*).

Environ. Sci. Technol. 52, 3195–3201. https://doi.org/10.1021/acs.est.7b05759

Dawson, A.L., Kawaguchi, S., King, C.K., Townsend, K.A., King, R., Huston, W.M.,

Bengtson Nash, S.M., 2018. Turning microplastics into nanoplastics through digestive

fragmentation by Antarctic krill. Nat. Commun. 9, 1–8. https://doi.org/10.1038/s41467-

018-03465-9

Derraik, J.G.B., 2002. The pollution of the marine environment by plastic debris: A review.

Mar. Pollut. Bull. 44, 842–852. https://doi.org/10.1016/S0025-326X(02)00220-5

Desforges, J.P.W., Galbraith, M., Dangerfield, N., Ross, P.S., 2014. Widespread distribution

of microplastics in subsurface seawater in the NE Pacific Ocean. Mar. Pollut. Bull. 79,

94–99. https://doi.org/10.1016/j.marpolbul.2013.12.035

Duis, K., Coors, A., 2016. Microplastics in the aquatic and terrestrial environment: sources

(with a specific focus on personal care products), fate and effects. Environ. Sci. Eur. 28,

1–25. https://doi.org/10.1186/s12302-015-0069-y

Erni-Cassola, G., Gibson, M.I., Thompson, R.C., Christie-Oleza, J.A., 2017. Lost, but found

with nile red: A novel method for detecting and quantifying small microplastics (1 mm

577  to 20 μm) in environmental samples. Environ. Sci. Technol. 51, 13641–13648.

578  https://doi.org/10.1021/acs.est.7b04512

579 Everaert, G., Van Cauwenberghe, L., De Rijcke, M., Koelmans, A.A., Mees, J.,

580  Vandegehuchte, M., Janssen, C.R., 2018. Risk assessment of microplastics in the ocean:

581  Modelling approach and first conclusions. Environ. Pollut. 242, 1930–1938.

582  https://doi.org/10.1016/j.envpol.2018.07.069

583 Fox, J., Weisberg, S., 2019. An {R} Companion to Applied Regression, Third. ed. Sage,

584  Thousand Oaks {CA}.

585 Gangi, J. Di, 1999. Phthalates in PVC medical products from 12 countries. Greenpeacee,

586  Washington, DC.

587 Gassel, M., Harwani, S., Park, J., Jahn, A., 2013. Detection of nonylphenol and persistent

588  organic pollutants in fish from the North Pacific Central Gyre. Mar. Pollut. Bull. 73,

589  231–242. https://doi.org/10.1016/j.marpolbul.2013.05.014

590 Gouin, T., Roche, N., Lohmann, R., Hodges, G., 2011. A thermodynamic approach for

591  assessing the environmental exposure of chemicals absorbed to microplastic. Environ.

592  Sci. Technol. 45, 1466–1472. https://doi.org/10.1021/es1032025

593 Granek, E.F., Brander, S.M., Holland, E.B., 2020. Microplastics in aquatic organisms:

594  Improving understanding and identifying research directions for the next decade.

595  Limnol. Oceanogr. Lett. 5, 1–4. https://doi.org/10.1002/lol2.10145

596 Hasegawa, T., Nakaoka, M., 2021. Trophic transfer of microplastics from mysids to fish

597  greatly exceeds direct ingestion from the water column. Environ. Pollut. 273, 116468.

598  https://doi.org/10.1016/j.envpol.2021.116468

599 Hermabessiere, L., Dehaut, A., Paul-Pont, I., Lacroix, C., Jezequel, R., Soudant, P., Duflos,

600  G., 2017a. Occurrence and effects of plastic additives on marine environments and

601  organisms: A review. Chemosphere 182, 781–793.

602  https://doi.org/10.1016/j.chemosphere.2017.05.096

Hermabessiere, L., Dehaut, A., Paul-Pont, I., Lacroix, C., Jezequel, R., Soudant, P., Duflos,

G., 2017b. Occurrence and effects of plastic additives on marine environments and

organisms: A review. Chemosphere 182, 781–793.

https://doi.org/10.1016/j.chemosphere.2017.05.096

Huang, W., Song, B., Liang, J., Niu, Q., Zeng, G., 2020. Microplastics and associated

contaminants in the aquatic environment : A review on their ecotoxicological effects ,

trophic transfer , and potential impacts to human health. J. Hazard. Mater. 124187.

https://doi.org/10.1016/j.jhazmat.2020.124187

Khosrovyan, A.., Marin, M.J.S.., Peret, A.M.., De Almeida, D.F.., Angel DelValls, T.., Riba,

I.., 2012. Weight-of-evidence-based assessment of sediment quality of the São Francisco

River (Brazil) with the help of software tool SQA. iEMSs 2012 - Manag. Resour. a Ltd.

Planet Proc. 6th Bienn. Meet. Int. Environ. Model. Softw. Soc. 2958–2965.

Kim, J.W., Chang, K.H., Isobe, T., Tanabe, S., 2011. Acute toxicity of benzotriazole

ultraviolet stabilizers on freshwater crustacean (*Daphnia pulex*). J. Toxicol. Sci. 36,

247–251. https://doi.org/10.2131/jts.36.247

Koelmans, A.A., Bakir, A., Burton, G.A., Janssen, C.R., 2016. Microplastic as a vector for

chemicals in the aquatic environment: critical review and model-supported

reinterpretation of empirical studies. https://doi.org/10.1021/acs.est.5b06069

Lenz, R., Enders, K., Nielsen, T.G., 2016. Microplastic exposure studies should be

environmentally realistic. Proc. Natl. Acad. Sci. 113, E4121–E4122.

https://doi.org/10.1073/pnas.1606615113

Mateos-Cárdenas, A., O'Halloran, J., van Pelt, F.N.A.M., Jansen, M.A.K., 2020. Rapid

fragmentation of microplastics by the freshwater amphipod *Gammarus duebeni* (Lillj.).

Sci. Rep. 10, 1–12. https://doi.org/10.1038/s41598-020-69635-2

Ministry of Agriculture, F. and F., 2016. Status of seafood consumption in Japan [WWW

Document]. URL

https://www.jfa.maff.go.jp/j/kikaku/wpaper/h29_h/trend/1/t1_2_4_2.html

Ministry of Health, L. and W., 2018. Average height and weight of Japanse by age and sex

[WWW Document]. URL https://www.mhlw.go.jp/toukei/youran/indexyk_2_1.html

(accessed 12.27.20)

Morohoshi, K., Yamamoto, H., Kamata, R., Shiraishi, F., Koda, T., Morita, M., 2005.

Estrogenic activity of 37 components of commercial sunscreen lotions evaluated by in

vitro assays. Toxicol. Vitr. 19, 457–469. https://doi.org/10.1016/j.tiv.2005.01.004

Nakata, H., Murata, S., Filatreau, J., 2009. Occurrence and concentrations of benzotriazole

UV stabilizers in marine organisms and sediments from the Ariake Sea, Japan. Environ.

Sci. Technol. 43, 6920–6926. https://doi.org/10.1021/es900939j

NITE, 2017. Risk assessment for chemical substances contained in products:

Decabromodiphenyl ether 1–12.

NOAA, 2020. What are microplastics? [WWW Document]. URL

https://oceanservice.noaa.gov/facts/microplastics.html (accessed 7.23.20).

Qiao, P., Gobas, F.A.P.C., Farrell, A.P., 2000. Relative contributions of aqueous and dietary

uptake of hydrophobic chemicals to the body burden in juvenile rainbow trout. Arch.

Environ. Contam. Toxicol. 39, 369–377. https://doi.org/10.1007/s002440010117

R Core Team, 2020. R: A Language and Environment for Statistical Computing.

Rani, M., Shim, W.J., Han, G.M., Jang, M., Al-Odaini, N.A., Song, Y.K., Hong, S.H., 2015.

Qualitative Analysis of Additives in Plastic Marine Debris and Its New Products. Arch.

Environ. Contam. Toxicol. 69, 352–366. https://doi.org/10.1007/s00244-015-0224-x

Rochman, C.M., Hoh, E., Kurobe, T., Teh, S.J., 2013. Ingested plastic transfers hazardous

chemicals to fish and induces hepatic stress. Sci. Rep. 3, 1–7.

https://doi.org/10.1038/srep03263

Rochman, C.M., Lewison, R.L., Eriksen, M., Allen, H., Cook, A.M., Teh, S.J., 2014.

Polybrominated diphenyl ethers (PBDEs) in fish tissue may be an indicator of plastic

contamination in marine habitats. Sci. Total Environ. 476–477, 622–633.

https://doi.org/10.1016/j.scitotenv.2014.01.058

Rummel, C.D., Löder, M.G.J., Fricke, N.F., Lang, T., Griebeler, E., Janke, M., Gerdts, G.,

2016. Plastic ingestion by pelagic and demersal fish from the North Sea and Baltic Sea.

MPB 102, 134–141. https://doi.org/10.1016/j.marpolbul.2015.11.043

Somo, D.A., Onukwufor, J.O., Wood, C.M., Richards, J.G., 2020. Interactive effects of

temperature and hypoxia on diffusive water flux and oxygen uptake rate in the tidepool

sculpin, Oligocottus maculosus. Comp. Biochem. Physiol. -Part A　Mol. Integr.

Physiol. 250. https://doi.org/10.1016/j.cbpa.2020.110781

Sun, B., Hu, Y., Cheng, H., Tao, S., 2019. Releases of brominated flame retardants (BFRs)

from microplastics in aqueous medium: Kinetics and molecular-size dependence of

diffusion. Water Res. 151, 215–225. https://doi.org/10.1016/j.watres.2018.12.017

Sun, Y., Wang, Y., Liang, B., Chen, T., Zheng, D., Zhao, X., Jing, L., Zhou, X., Sun, Z., Shi,

Z., 2020. Hepatotoxicity of decabromodiphenyl ethane (DBDPE) and

decabromodiphenyl ether (BDE-209) in 28-day exposed Sprague- Dawley rats. Sci.

Total Environ. 705, 135783. https://doi.org/10.1016/j.scitotenv.2019.135783

Tanaka, K., Takada, H., Yamashita, R., Mizukawa, K., Fukuwaka, M. aki, Watanuki, Y.,

2013. Accumulation of plastic-derived chemicals in tissues of seabirds ingesting marine

plastics. Mar. Pollut. Bull. 69, 219–222. https://doi.org/10.1016/j.marpolbul.2012.12.010

Tanaka, K., Takada, H., Yamashita, R., Mizukawa, K., Fukuwaka, M.A., Watanuki, Y., 2015.

Facilitated leaching of additive-derived PBDEs from plastic by seabirds' stomach oil

and accumulation in tissues. Environ. Sci. Technol. 49, 11799–11807.

https://doi.org/10.1021/acs.est.5b01376

Tanaka, K., van Franeker, J.A., Deguchi, T., Takada, H., 2019. Piece-by-piece analysis of

additives and manufacturing byproducts in plastics ingested by seabirds: Implication for

risk of exposure to seabirds. Mar. Pollut. Bull. 145, 36–41.

https://doi.org/10.1016/j.marpolbul.2019.05.028

Tanaka, K., Watanuki, Y., Takada, H., Ishizuka, M., Yamashita, R., Kazama, M., Hiki, N.,

Kashiwada, F., Mizukawa, K., Mizukawa, H., Hyrenbach, D., Hester, M., Ikenaka, Y.,

Nakayama, S.M.M., 2020. *In vivo* accumulation of plastic-derived chemicals into

seabird tissues. Curr. Biol. 30, 723-728.e3. https://doi.org/10.1016/j.cub.2019.12.037

Walkinshaw, C., Lindeque, P.K., Thompson, R., Tolhurst, T., Cole, M., 2020. Microplastics

and seafood: lower trophic organisms at highest risk of contamination. Ecotoxicol.

Environ. Saf. 190, 110066. https://doi.org/10.1016/j.ecoenv.2019.110066

Wang, W., Ge, J., Yu, X., 2020. Bioavailability and toxicity of microplastics to fish species:

A review. Ecotoxicol. Environ. Saf. 189, 109913.

https://doi.org/10.1016/j.ecoenv.2019.109913

Wang, X., Ling, S., Guan, K., Luo, X., Chen, L., Han, J., Zhang, W., Mai, B., Zhou, B., 2019.

Bioconcentration, biotransformation, and thyroid endocrine disruption of

decabromodiphenyl ethane (DBDPE), a novel brominated flame retardant, in zebrafish

larvae. Environ. Sci. Technol. 53, 8437–8446. https://doi.org/10.1021/acs.est.9b02831

Wang, Y., Chen, T., Sun, Y., Zhao, X., Zheng, D., Jing, L., Zhou, X., Sun, Z., Shi, Z., 2019.

A comparison of the thyroid disruption induced by decabrominated diphenyl ethers

(BDE-209) and decabromodiphenyl ethane (DBDPE) in rats. Ecotoxicol. Environ. Saf.

174, 224–235. https://doi.org/10.1016/j.ecoenv.2019.02.080

Wardrop, P., Shimeta, J., Nugegoda, D., Morrison, P.D., Miranda, A., Tang, M., Clarke,

B.O., 2016. Chemical pollutants sorbed to ingested microbeads from personal care

products accumulate in fish. Environ. Sci. Technol. 50, 4037–4044.

https://doi.org/10.1021/acs.est.5b06280

Webb, N.A., Shaw, J.R., Morgan, J., Hogstrand, C., Wood, C.M., 2001. Acute and chronic

physiological effects of silver exposure in three marine teleosts. Aquat. Toxicol. 54,

161–178. https://doi.org/10.1016/S0166-445X(01)00150-3

Wright, S.L., Kelly, F.J., 2017. Plastic and human health: A micro issue? Environ. Sci.

Technol. 51, 6634–6647. https://doi.org/10.1021/acs.est.7b00423

Wright, S.L., Thompson, R.C., Galloway, T.S., 2013. The physical impacts of microplastics

on marine organisms: A review. Environ. Pollut.

https://doi.org/10.1016/j.envpol.2013.02.031

Zhang, F., Wang, X., Xu, J., Zhu, L., Peng, G., Xu, P., Li, D., 2019. Food-web transfer of

microplastics between wild caught fish and crustaceans in East China Sea. Mar. Pollut.

Bull. 146, 173–182. https://doi.org/10.1016/j.marpolbul.2019.05.061

Ziccardi, L.M., Edgington, A., Hentz, K., Kulacki, K.J., Kane Driscoll, S., 2016.

Microplastics as vectors for bioaccumulation of hydrophobic organic chemicals in the

marine environment: A state-of-the-science review. Environ. Toxicol. Chem.

https://doi.org/10.1002/etc.3461

Tables

Table 1 Profile of additives compounded in the plastics used for the experiment. Octanol-water

partition coefficient (K_{ow}) is represented as a hydrophobicity parameter of each chemical.

Compounds	Concentration (µg/g-plastic)	Molecular weight (g/mol)	logK$_{ow}$
BDE209	357	959	12.1
DBDPE	74.2	971	13.6
UV-234	361	448	7.67
UV-327	397	358	6.91
BP-12	123	326	6.96

Table 2 Result of the leaching experiment with 2 mg of microplastics exposed to 1-L seawater for 24 h. SE represents standard error. Grey colored cells represent values under the limit of quantifications (LOQ). All DBDPE values are technically lower than LOQ because the blank value was very high.

Additive	Blank	Initial amount in plastics (ng)		Final amount in seawater (ng)		Leaching rate (%)	
		Mean	SE	Mean	SE	Mean	SE
BDE 209	0.127	713	60951	6.27	5.27	0.86%	0.72%
DBDPE	60.0	148	21930	4.13	3.53	2.61%	2.20%
UV 327	0.208	722	66262	5.12	1.34	0.70%	0.18%
UV 234	0.327	794	9662	23.2	5.94	2.83%	0.71%
BP 12	4.83	246	12082	23.8	3.14	8.79%	1.07%

731 **Table 3** Results of likelihood ratio test on the generalized linear model for the concentrations

732 of additives in mysid as functions of treatment (ambient control, leachate treatment, and

733 microplastic treatment).

Additive	Fixed factor	LR Chisq	df	*p*-value
BDE209	Treatment	253	2	< 0.001
DBDPE	Treatment	186	2	< 0.001
UV234	Treatment	134	2	< 0.001
UV327	Treatment	280	2	< 0.001
BP12	Treatment	10.8	2	0.004

736 **Table 4** Results of likelihood ratio test on the generalized linear model for the concentrations

737 of additives in fish muscle and liver as functions of treatment (ambient control, water treatment,

738 and mysid treatment).

Body tissue	Additive	Fixed factor	LR Chisq	df	*p*-value
Muscle (Dry weight)	BDE209	Treatment	27.9	2	< 0.001
	DBDPE	Treatment	27.2	2	< 0.001
	UV234	Treatment	0.441	2	0.802
	UV327	Treatment	9.72	2	0.00775
	BP12	Treatment	99.5	2	< 0.001
Muscle (Lipid weight)	BDE209	Treatment	38.2	2	< 0.001
	DBDPE	Treatment	36.4	2	< 0.001
	UV234	Treatment	5.15	2	0.0763
	UV327	Treatment	17.6	2	< 0.001
	BP12	Treatment	91.0	2	< 0.001
Liver (Dry weight)	BDE209	Treatment	18.5	2	< 0.001
	DBDPE	Treatment	11.9	2	0.00257
	UV234	Treatment	33.4	2	< 0.001
	UV327	Treatment	40.4	2	< 0.001
	BP12	Treatment	7.01	2	0.0301

741 **Figures**

743   **Fig. 1.** Size distribution of polyethylene microplastics used for the experiment. Red line

744   denotes the mean particle size.

746   **Fig. 2.** Composition of additives in polyethylene microplastics used for the experiment.

749 **Fig. 3** Concentration of the five additives in mysids per individual, collected from the ambient

750 environment (ABT), treated with the leachate seawater (LCT), and exposed to microplastics

751 (MPs). Results are shown as box and whisker plots with median (solid horizontal line),

752 interquartile range (25th and 75th percentiles; box), and the 10th and 90th percentiles

753 (whiskers). Different letters denote significant differences by post-hoc comparison ($p < 0.05$

754 GLM with post-hoc Tukey's HSD).

Fig. 4. Concentration of the five additives in fish body and liver, collected from the ambient

environment (ABT), exposed to microplastics suspended in the water column (Water), and fed

plastic-exposed mysid (Mysid). Both dry weight and lipid weight base concentrations are

shown for body, but only the dry weight base concentration is shown for liver because the

available sample volume was limited. Results are shown as box and whisker plots with median

(solid horizontal line), interquartile range (25th and 75th percentiles; box), and the 10th and

90th percentiles (whiskers). Different letters denote significant differences by post-hoc

comparison ($p < 0.05$ GLM with post-hoc Tukey's HSD).

765