Gene expression shapes the patterns of parallel evolution of herbicide resistance in the

agricultural weed *Monochoria vaginalis*

Shinji Tanigaki[1], Akira Uchino[2], Shigenori Okawa[3], Chikako Miura[4], Kenshiro Hamamura[5], Mitsuhiro Matsuo[6], Namiko Yoshino[7], Naoya Ueno[8], Yusuke Toyama[9], Naoya Fukumi[10], Eiji Kijima[11], Taro Masuda[12], Yoshiko Shimono[1], Tohru Tominaga[1], Satoshi Iwakami[1*]

[1]Graduate school of agriculture, Kyoto University, Kitashirakawa-Oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan

[2]Central Region Agricultural Research Center, National Agriculture and Food Research Organization, Tsu 514-2392, Japan

[3]Miyagi Prefectural Furukawa Agricultural Experiment Station, Fukoku 88, Furukawa-Osaki, Osaki 989-6227, Japan

[4]Akita Prefectural Agricultural Experiment Station, 34-1 Yuwaaikawa-azagenpachizawa, Akita 010-1231, Japan

[5]Japan Association for Advancement of Phyto-Regulators (JAPR), 860 Kashiwada-cho, Ushiku 300-1211, Japan

[6]Field Science Center, Faculty of Agriculture, University of Miyazaki, 1-1 Gakuen-kibanadai-nishi, Miyazaki 889-2192, Japan

[7]Tohoku Agricultural Research Center, National Agriculture and Food Research Organization, 50 Harajukuminami, Arai, Fukushima 960-2156, Japan

[8]Yamanashi Prefectural Agritechnology Center, 1100 Shimoimai, Kai 400-0105, Japan

[9]Shizuoka Prefectural Research Institute of Agriculture and Forestry, 678-1 Tomigaoka, Iwata 438-0803, Japan

[10]Tottori Agricultural Experiment Station, 260 Hashimoto, Tottori 680-1142, Japan

[11]Yamaguchi Prefectural Agriculture and Forestry General Technology Center, 1-1-1 Ouchihikami, Yamaguchi 753-0231, Japan

[12]Faculty of Agriculture, Setsunan University, 45-1 Nagaotoge-cho, Hirakata 573-0101, Japan

*Corresponding Author, E-mail: iwakami.satoshi.2v@kyoto-u.ac.jp, Phone: +81-75-753-6064

Abstract

The evolution of herbicide resistance in weeds is an example of parallel evolution, through which genes encoding herbicide target proteins are repeatedly represented as evolutionary targets. The number of herbicide target-site genes differs among species, and little is known regarding the effects of duplicate gene copies on the evolution of herbicide resistance. We investigated the evolution of herbicide resistance in *Monochoria vaginalis*, which carries five copies of sulfonylurea target-site acetolactate synthase (*ALS*) genes. Suspected resistant populations collected across Japan were investigated for herbicide sensitivity and *ALS* gene sequences, followed by functional characterisation and *ALS* gene expression analysis. We identified over 60 resistant populations, all of which carried resistance-conferring amino acid substitutions exclusively in *MvALS1* or *MvALS3*. All *MvALS4* alleles carried a loss-of-function mutation. Although the enzymatic properties of ALS encoded by these genes were not markedly different, the expression of *MvALS1* and *MvALS3* was prominently higher among all *ALS* genes. The higher expression of *MvALS1* and *MvALS3* is the driving force of the biased representation of genes during the evolution of herbicide resistance in *M. vaginalis*. Our findings highlight that gene expression is a key factor in creating evolutionary hotspots.

Keywords: acetohydroxy acid synthase, acetolactate synthase, convergent evolution, evolutionary constraint, target-site resistance, weed evolution

INTRODUCTION

Agricultural weeds are often subject to artificial lethal stresses employed in an effort to increase crop yield. Such weeding stresses, in turn, act as selection pressures driving the evolution of weed populations. While multiple evolved traits related to weed adaptation in agricultural ecosystems have been documented, such as flowering phenology, mimicry morphology, seed shattering, seed dormancy, and competitiveness (Fukano, Guo, Uchida, Tachiki, & Cornelissen, 2020; Waselkov, Regenold, Lum, & Olsen, 2020; Wedger & Olsen, 2018; Ye et al., 2019), herbicide resistance remains among the best studied and characterised trait (Powles & Yu, 2010). Evolved herbicide resistance has been documented worldwide in as many as 263 weed species (Heap, 2021), posing a great threat to current global food security.

While weed resistance has often been studied from an agricultural viewpoint as a threat to food security, it has also attracted the attention of evolutionary biologists as a classic example of an 'evolution in action' (Baucom, 2019). An important insight from studies on weed resistance is that nature often mimics itself: certain genes have repeatedly acted as the major components in the evolution of resistance (Martin & Orgogozo, 2013). Research so far has observed that weed populations subjected to herbicide applications evolved resistance via an identical amino acid substitution (AAS) in the genes encoding herbicide target sites, irrespective of evolutionary lineage (Baucom, 2016). These mutations result in conformational alterations in the binding site of each herbicide, leading to the herbicide-insensitive form of the protein (Powles & Yu, 2010). Although this parallelism has been observed in many herbicides with diverse modes of action, resistance to herbicides inhibiting acetolactate synthase (ALS, or acetohydroxy acid synthase), in particular,

represents a case of extreme parallelism at the molecular level (Yu & Powles, 2014). Due to

their low toxicity to animals, high crop selectivity, and high efficacy at low doses, ALS

herbicides are among the most frequently used herbicides in many countries, including Japan

(Hamamura, 2018; Peters & Strek, 2018). The ALS residues where AASs induce resistance

have been reported in as many as eight from natural weed populations, and most of these

mutations were not significantly associated with fitness cost (Yu & Powles, 2014). This nature

is considered as the driving force to render *ALS* loci as an evolutionary hotspot for resistance

to ALS herbicides (Baucom, 2019).

Little attention has been paid to the effects of duplicated gene copies on the

evolution of herbicide resistance. Having undergone whole genome duplications during the

long course of evolution (Clark & Donoghue, 2018), many plants possess duplicated copies of

target-site genes even in diploid species, e.g. *ALS* genes in *Senecio vulgaris* and *Zea*

mays(Délye, Causse, & Michel, 2016; Svitashev et al., 2015) and acetyl-CoA carboxylase

genes in *Echinochloa* spp. (Iwakami et al., 2015; Iwakami, Uchino, Watanabe, Yamasue, &

Inamura, 2012). Genetic redundancy by duplicated copies fosters the evolutionary

differentiation of each copy, such as gene expression and/or function of the encoded protein,

including pseudogenisation (Flagel & Wendel, 2009). Although not much is understood in the

differentiation of herbicide target-site genes in weeds, the differentiation, if any, may provide

distinct competency for resistance evolution in each "potentially hot" locus, influencing the

evolutionary dynamics of resistance mechanisms.

Monochoria vaginalis (syn. *Pontederia vaginalis*) is a noxious allohexaploid ($2n = 4x = 52$)

weed in temperate and tropical Asian paddy fields (Holm, Plucknett, Pancho, & Herberger,

1977). It carries five *ALS* loci, with all genes transcribed at the seedling stage (Iwakami et al.,

2020) when plants are exposed to herbicides in agricultural fields. Among the limited number

of populations tested to date, mutations known to confer ALS herbicide resistance have been

reported exclusively in *MvALS1* or *MvALS3* (e.g. Iwakami et al., 2020; Ohsako & Tominaga,

2007), despite the presence of multiple potential evolutionary hotspots for resistance. In the

present study, we confirmed the occurrence of resistance-conferring mutations exclusively in

MvALS1 and *MvALS3* among 68 resistant populations across Japan. Enzyme kinetics and

expression analysis of each *ALS* copy and its product unveiled the potential mechanisms that

make these two loci the hotspots of resistance evolution.

MATERIALS AND METHODS

M. vaginalis

M. vaginalis was collected from paddy fields across Japan (Figure 1a-b; Table S1). Most of

the paddy fields were small, measuring 0.1 to 0.3 ha an average. Plants from a single paddy

field are hereafter referred to as a population. Adjacent fields were avoided during sampling.

The plants were either directly used for resistance diagnosis using the rooting assay or

cultivated for the collection of selfed seeds for the following pot assay (Figure 1c). The

phenotyped plants were subjected to *ALS* gene sequencing (see section 2.4).

Resistance diagnosis in *M. vaginalis*

Resistance to ALS herbicides was diagnosed using the rooting assay with 45 $\mu g \cdot L^{-1}$ of

bensulfuron-methyl, as described by Hamamura et al. (2003). In some populations, the selfed

seeds were used for resistance diagnosis using a recommended dose of imazosulfuron [90 g

active ingredient (a.i.)·ha^{-1}], as described previously (Iwakami et al., 2020). Briefly, seeds

were germinated in distilled water in a 1.5 mL tube at 25°C in a climate chamber with a 12 h

photoperiod. The germinated seeds were transplanted into autoclaved soil under flooded

conditions. When plants reached the three-leaf stage, a commercial formulation of

imazosulfuron (75% a.i.) was applied to the surface of standing water. The survival rate of

plants was evaluated at 3 weeks post application.

Imazosulfuron dose–response analysis in *M. vaginalis*

KST1, KOM1, SMM23, AKT3, NRS2, and SMM13 populations were used in the assay. The

dose–response analysis was performed as described by Iwakami et al. (2020), with some

modifications. Briefly, the seeds were germinated as above, and five seedlings were

transplanted into autoclaved soil in 100 cm^2 plastic pots. The plants were cultivated under

flooded conditions. At the two- to three-leaf stage, the commercial formulation of

imazosulfuron was applied to the surface of standing water in individual pots at a rate

calculated for the area of each pot (0, 0.09, 0.9, 9, 90, 900, 9,000, and 90,000 g a.i.·ha^{-1}). At 3

weeks post application, the shoots were harvested and dry weight was measured. The dose

that inhibited growth to 50% (GR$_{50}$) and 90% (GR$_{90}$) compared with the control was

calculated by fitting a log-logistic curve (two-parameter model) using the drc package (ver.

3.0.1) (Ritz, Baty, Streibig, & Gerhard, 2015).

Sequencing of *ALS* genes in *M. vaginalis*

DNA was extracted from the leaves of plants subjected to the herbicide sensitivity assay using

the cetyltrimethylammonium bromide (CTAB) method (Murray & Thompson, 1980). For

some old populations, DNA was extracted from dead seeds using the Plant Genomic DNA Kit

(Tiangen, Beijing, China). The entire coding sequence of *MvALS1* to *MvALS5* was

PCR-amplified as described by Iwakami et al. (2020). The amplicons were treated with

ExoSAP-IT (Affymetrix, CA, USA) to remove dNTPs and primers, and sent for Sanger

sequencing to Eurofins Genomics (Tokyo, Japan).

To evaluate the genetic diversity of the whole coding sequence in each *ALS* gene, the

nucleotide diversity of each *ALS* gene was calculated using DnaSP 6 (Rozas et al., 2017). The

nucleotide diversity at each nucleotide position was also computed to evaluate the difference

of the diversity patterns among *ALS* genes. Previously reported full-length coding sequences

of three populations (SMM13, SMM23, and TOT1) were also used in the analysis (Iwakami

et al., 2020).

Expression and *in vitro* assay of ALS

The pGEX6P-1 vector (Cytiva, Tokyo, Japan) was digested with SalI-HF (New England

Biolabs, Tokyo, Japan). Alleles of *MvALS1/2/3/5* from the sensitive population (SMM13)

were individually ligated at the SalI site of pGEX-6 via SLiCE reaction following the

protocol described by Motohashi (2015). Each *ALS* gene was expressed without the estimated

chloroplast transit peptide (CTP) region corresponding to 1-86 residues of *Arabidopsis*

thaliana ALS (Chang & Duggleby, 1997; Singh, Schmitt, Lillis, Hand, & Misra, 1991). The

primers used for the SLiCE reactions are listed in Table S2. The resultant expression vector

was transformed into *Escherichia coli* BL21 (DE3) cells (Sigma-Aldrich, MO, USA).

The transformed cells were inoculated in 200 mL of LB medium containing 100

$\mu g \cdot ml^{-1}$ of ampicillin and cultured at 37°C. When OD_{600} reached 0.4-0.7, isopropyl

β-D-thiogalactopyranoside was added to the medium at a final concentration of 1 mM, and

the culture was continued at 16°C for 24 h. The cells were harvested by centrifugation at

14,000 $\times g$ for 10 min. The cells were resuspended in phosphate-buffered saline with 1%

Triton X-100, 1 mM phenylmethylsulfonyl fluoride, 5 mM dithiothreitol, and 0.5 mM $MgCl_2$

and sonicated for 5 min on ice. GST-fused ALS was batch-purified with glutathione

Sepharose 4B (Cytiva) according to the manufacturer's protocol. ALS was eluted in 400 μL

of 50 mM Tris–HCl buffer (pH 8.0) containing 10 mM glutathione, followed by the addition

of glycerol at a final concentration of 10% (v/v).

ALS activity was evaluated as described by Chang and Duggleby (1997), with some

modifications. The reaction mixture (300 μL) containing 20 mM K_2HPO_4/KH_2PO_4 (pH 7.0),

50 mM sodium pyruvate, 10 mM $MgCl_2$, 1 mM thiamine diphosphate, 20 μM FAD, 0.5 ng

enzyme, and varying concentrations of sodium pyruvate (0, 0.78, 1.56, 3.12, 6.25, 12.5, and

25 mM) was incubated at 30°C for 60 min, and the reaction was terminated by adding 30 μL

of 6N H_2SO_4. The resultant solution was incubated at 60°C for 15 min, followed by the

addition of 300 μL creatine (0.5%, w/v) and α-naphthol (0.5%, w/v, in 2.5N NaOH).

Absorbance was measured at 525 nm. The data were plotted and curve-fitted to the

Michaelis–Menten equation using the drc package (ver. 3.0.1) (Ritz et al., 2015).The

Michaelis–Menten constant (K_m) and turnover number (k_{cat}) was determined for each ALS.

The experiments were conducted in triplicate and repeated twice.

Protein concentration was determined using Bio-Rad Protein Assay Dey Reagent

Concentrate (Bio-Rad, CA) with bovine serum albumin as the standard, following

manufacturer's instructions.

Targeted mutagenesis of *ALS* genes

Alleles of *MvALS1*/2/3/5 from the SMM13 population and a deleted allele of *MvALS5* from

the SMM3 population were cloned into the pCRblunt vector (Thermo Fisher Scientific, Tokyo,

Japan). The codons for Pro197 of the *ALS* genes were mutated to carry a resistance-conferring

Ser codon. Targeted mutagenesis was performed according to the protocol in QuikChange

Site-Directed Mutagenesis System (Stratagene, CA, USA), with some modifications. Briefly,

the clones were used as a template for PCR using PrimeSTAR MAX (TaKaRa, Kusatsu,

Japan); the primers are listed in Table S2. The PCR products were treated with DpnI-HF

(New England Biolabs). The resultant solution was used for *E. coli* DH5α transformation

(ECOS Competent *E. coli* DH5 α, Nippon Gene, Tokyo, Japan) according to the procedures

described in the manual. Plasmids were extracted from a culture solution inoculated with a

single colony of transformed *E. coli*. The success of the targeted mutagenesis was confirmed

by sequencing the extracted plasmids.

Transformation and phenotyping of transgenic *A. thaliana*

The Pro197Ser-mutated *ALS* genes and the native form of *MvALS1* gene were inserted into

the pCAMBIA1390 vector under the cauliflower mosaic virus 35S promoter using the

In-Fusion DH Cloning Kit (TaKaRa), as described previously (Iwakami et al., 2019). The

primers used in this reaction are listed in Table S2. *A. thaliana* transformation was performed

using the agrobacterium-mediated floral dip method as described previously (Guo et al.,

2019). The transformants were selected with hygromycin B (20 mg· L^{-1}) .

To quantify transgene expression, real-time RT-PCR was performed. Briefly, RNA

from 10-day-old T3 homozygous lines was extracted using the Plant Total RNA Mini Kit

(Favorgen, Ping-Tung, Taiwan). Contaminated DNA in the extracted RNA was removed using

the Turbo DNA-*free* kit (Thermo Fisher Scientific). The resultant RNA was reverse

transcribed according to the method described by Iwakami et al. (2019). PCR was run using

THUNDERBIRD SYBR qPCR Mix (TOYOBO, Osaka, Japan) on QuantStudi 12K Flex

Real-Time PCR System (Thermo Fisher Scientific). The primers for transgene expression

were designed using the terminator sequence of pCAMBIA1390 (Table S2). *GAPDH* was

used as the internal control; primers developed by Czechowski, Stitt, Altmann, Udvardi, and

Scheible (2005) were used. Data were analysed using the $\Delta\Delta$CT method (Schmittgen & Livak,

2008).

Imazosulfuron sensitivity of the transformants was evaluated as described

previously (Dimaano, Yamaguchi, Fukunishi, Tominaga, & Iwakami, 2020) with some

modifications. Seeds were sown on Murashige and Skoog medium (Murashige & Skoog,

1962) containing various concentrations of imazosulfuron. Plants were grown for 10 days at

22°C. Sensitivity was evaluated based on the emergence of true leaves. GR_{50} was computed

as described above.

Gene expression analysis in *M. vaginalis*

SMM13 and SMM23 were used as representatives of the sensitive and resistant populations, respectively. Germinated seeds were transplanted into autoclaved soils. Six plants from each population were grown in water until the three-leaf stage. Half of the plants were treated with imazosulfuron at a concentration of 90 g a.i.·ha^{-1}. At 24 h post application, the shoots of all plants were snap frozen in liquid nitrogen and stored at -80°C until further use. RNA was extracted using the RNeasy Plant Mini Kit (Qiagen, Venlo, Netherlands), followed by DNase treatment using the Turbo DNA-*free* kit (Thermo Fisher Scientific).

To estimate approximate transcript abundance, PCR products generated by the primers designed based on the completely conserved regions among the five copies of *ALS* genes (Table S2) were cloned into the pCRblunt vector (Thermo Fisher Scientific). Over 50 clones were sent to Eurofins Genomics for sequencing. cDNA was prepared from the RNA of an imazosulfuron-treated sensitive plant. The experiment was performed twice.

For RNA-Seq analysis, 12 RNA samples were shipped to GENEWIZ (Kawaguchi, Japan) for stranded RNA-Seq using Illumina HiSeq 4000 (150-bp paired-end reads). Contaminated adaptor sequences were trimmed using 'cutadapt' (ver. 2.5) (Martin, 2011). Low-quality bases were removed using Trimmomatic (ver. 0.38) (Bolger, Lohse, & Usadel, 2014) with the following options: 'LEADING:25 TRAILING:25 SLIDINGWINDOW:4:15 MINLEN:100'. Reads of a non-treated sensitive plant were used for contig assembly. The assembly was performed using Trinity (ver. 2.4.0) (Grabherr et al., 2011) with a k-mer size of 32. The biological function of the contigs was inferred using BLASTX (Blast+ ver. 2.9.0) (Camacho et al., 2009) against rice protein sequences (*Oryza sativa* MSU release 7). Only one

contig under the identical gene identifier annotated by Trinity was used, as described by Ono

et al. (2015). Briefly, from contigs with hits in the BLAST analysis, those with the highest bit

score were selected, and from contigs without hits, the longest ones were selected. Contigs

that were annotated as *ALS* genes were replaced with full-length *ALS* genes of *M. vaginalis*

(DDBJ accession numbers: LC488950 to LC488953 and LC488956). The reads were mapped

using Bowtie2 (ver. 2.3.4.1) (Langmead & Salzberg, 2012) with the following options: '-X

900 --very-sensitive --no-discordant --no-mixed --dpad 0 --gbar 99999999'. The concordantly

mapped reads were counted using RSEM (ver. 1.3.2) (Li & Dewey, 2011). Multiple

comparisons were performed using the GLM approach with the quasi-likelihood F-test using

the edgeR package (ver. 3.28.1.) (Robinson, McCarthy, & Smyth, 2010). Contigs with

fold-changes of more or less than 2 and *q*-values less than 0.05 were evaluated as

differentially expressed genes (DEGs).

RESULTS

Herbicide resistance of *M. vaginalis* collected across Japan

In this study, we sampled 113 suspected resistant populations across Japan and tested for ALS

herbicide sensitivity using the rooting or pot assays (Figure 1). Sixty-six populations (58.4%)

were confirmed to be resistant to ALS herbicides. All the resistant populations carried one or

two homozygous mutations in the codons of *ALS* genes that are known to confer resistance to

the ALS herbicides, namely Pro197, Ala205, and Asp376, implying that *M. vaginalis* tend to

evolve resistance to ALS inhibitors via target-site resistance (TSR) mechanisms as reported in

many non-Poaceae weeds (Yu & Powles, 2014).

274   Analysis of the codons involved in ALS resistance in the 66 populations tested in

this study and 2 previously reported resistant populations (Iwakami et al., 2020) revealed 9

types of resistance-conferring AASs. Intriguingly, the identified resistance-conferring

mutations were exclusively found in *MvALS1* and *MvALS3*, and not in other three genes

(Figure 2a). Mutations at Pro197 dominated the others (Figure 2b), probably because

sulfonylureas, a chemical class known to select for Pro197 mutation (Yu & Powles, 2014),

have been the major ALS herbicides used in Japan (Hamamura, 2018). There was no clear

association between mutation type (i.e. AASs and genes) and geographical region (Figure 1b;

Table S1). As relatively rare cases, populations with two independent mutations were also

found. AIZ2 and KR1 populations carried two resistance-conferring mutations, namely

Pro197Ser and Asp376Glu, in a single gene *(MvALS1)*, while YNZ1 population carried

Asp376Glu in *MvALS1* and Pro197Ser in *MvALS3*. In addition, mutations that have not been

reported in natural weed populations, including Asp376Asn in NDA1 and Ala205Thr in

ANP1, were also detected. Both mutations were confirmed to confer reduced sensitivity to

sulfonylurea herbicides in yeast (Bedbrook et al., 1995; Duggleby, Pang, Yu, & Guddat, 2003).

Thus, AASs may be the causal mutations for resistance in these weed populations, although it

should be noted that we did not actively screen for additional non-target site mechanisms.

291   Transition was the dominant type of resistance-conferring mutation (Figure 2c),

consistent with the results of a previous genome-wide mutation analysis of *A. thaliana* (Weng

et al., 2019). The highest frequency of the Pro197Ser transition mutation (CCT→<u>T</u>CT) may

be partly attributed to the nature of mutation patterns, although other factors such as high

level resistance caused by this mutation (Sada, Ikeda, & Kizawa, 2013) may play a more

important role in the observed bias.

To gain insights into the mystery of the exclusive presence of the TSR allele in only

two *ALS* loci, we further analysed the coding sequence of each *ALS* gene in 66 resistant and

30 sensitive populations, along with the previously reported sequences of 2 resistant and 1

sensitive populations (Iwakami et al., 2020) (Table S1). Coding sequences of all *ALS* genes in

M. vaginalis were homozygous, contrary to the observation in obligate outcrossing weeds

such as *Lolium rigidum* and *Papaver rhoeas* (Délye, Boucansaud, Pernin, & Le Corre, 2009;

Délye, Pernin, & Scarabel, 2011). This result highlights the predominant self-pollinating

nature of *M. vaginalis*. Consistent with previous reports (Iwakami et al., 2020; Ohsako &

Tominaga, 2007), a T insertion causing a frameshift was detected in all *MvALS4* alleles

investigated in this study (Figure 2a-d). The products encoded by these mutant alleles lacked

domains with pivotal roles in ALS activity, indicating that this gene does not encode the

functional form of ALS. Thus, the prevalence of frame-shifted alleles in *MvALS4* among *M.*

vaginalis populations in Japan explains the absence of resistance-conferring mutations in this

gene. While all alleles were pseudogenised in *MvALS4*, variations were observed in *MvALS1*

and *MvALS5*; as such a 13 bp deletion was detected in *MvALS1* in 1 population and a 40 bp

deletion in *Mv*ALS5 in 12 populations (Figure 2a, TableS1). The reading frame of these

alleles also shifted, resulting in premature stop codons (Figure 2d). The putative

loss-of-function alleles of *MvALS5* were found in populations sampled from various areas

without any geographical association. Contrary to that in *MvALS4*, majority of the alleles

were not pseudogenised in *MvALS1* and *MvALS5*.

The analysis of coding sequences revealed that genes with resistance-conferring

alleles (*MvALS1* and *MvALS3*) showed a higher nucleotide diversity and greater variation in

haplotype number (Figure 2e). This trend held true even when naturally selected

resistance-conferring alleles were excluded from the analysis: a greater nucleotide diversity

(per site) was observed at three and four positions, irrespective of the resistance selection, in

MvALS1 and *MvALS3*, respectively. Thus, *MvALS1* and *MvALS3* might be more prone to

spontaneous mutations, some of which may occur by chance at the resistance-conferring

codons. However, the difference in the rate of spontaneous mutations among genes can hardly

explain the all-or-nothing resistance-conferring alleles of the *ALS* gene family in *M. vaginalis*.

Other more pronounced factors may have played key roles in shaping these biased patterns.

Functional characterisation of each ALS

To elucidate the mechanism underlying the disproportionate distribution of TSR mutations

among *ALS* copies, we next compared the enzymatic functions of the protein encoded by the

canonical alleles of each *ALS* gene. ALS activity of the purified GST-tagged protein was

assessed (Figure S1). The activity of the recombinant proteins was evaluated based on the

conversion of the produced acetolactate to a red insoluble complex. The reaction solutions of

the recombinant proteins of MvALS1/2/3/5 developed a red colour, indicating that all

isozymes produced acetolactate (Figure 3a). The enzymatic properties of the respective

isozymes were similar, although the affinity to the substrate pyruvate (*kcat/Km*) was relatively

high in MvALS1.

We examined the potential of TSR mutation to lead to herbicide resistance in all

loci under consideration by transforming *A. thaliana.* In this study, we artificially generated

Pro197Ser alleles (Figure 3b), which are known to endow extreme resistance to some ALS

herbicides (such as imazosulfuron) in other species (Sada et al., 2013). First, we investigated

the impact of the TSR mutation by comparing the imazosulfuron sensitivity of *A. thaliana*

lines carrying the *MvALS1* gene with or without the TSR mutation. We generated five and

four single-copy lines of the native allele and the TSR allele with varying transcript levels

(Figure 3c). Lines with the TSR allele survived the lethal dose of 10 nM, and even 100 nM

(Figure 3d). On the other hand, lines with the wild-type allele exhibited only a marginal

resistance level, with plants in the highest expression lines dead at 100 nM. The results

indicate that the qualitative, not quantitative, alteration in the ALS target site plays a key role

in the evolution of resistance to ALS herbicides.

Next, we investigated the sensitivity of *A. thaliana* lines with the TSR allele of the

respective *MvALS* gene. Interestingly, all the TSR alleles conferred imazosulfuron resistance

in *A. thaliana* when overexpressed (Figure 3e-f). Resistance indices of all transgenic lines

were correlated with the transcript level of transgenes (*ALS* genes of *M. vaginalis*), and a

similar resistance level was observed when the transcripts of the respective *ALS* genes were

accumulated at a similar level. Notably, even a small difference in transcript level drastically

affected herbicide sensitivity when the transcript levels were within a specific range. For

instance, although the transcript levels in three *MvALS1* transgenic lines (lines indicated in

pink area in Figure 3b) differed by only 5.1-fold at the maximum, their resistance levels

differed by as much as 2,500-fold. Meanwhile, very low or high transcript levels did not

affect sensitivity.

We also tested the putative loss-of-function allele of *MvALS5*, as it only partially

lacks the γ domain—one of the three ALS domains (Figure 2d). All T2 generations in which

the allele of *MvALS5* carrying the Pro197Ser mutation was deleted were sensitive to

imazosulfuron, while at least seven lines carrying the canonical allele of *MvALS5* showed

decreased imazosulfuron sensitivity (Figure S2). This result implies that the deleted form of

MvALS5 does not encode functional ALS.

Collectively, the results of recombinant protein assay and ectopic expression in *A.*

thaliana suggest that the activities of the four isozymes were near identical, although a

slightly higher substrate specificity was observed in MvALS1 in the recombinant assay.

Notably, the recombinant ALS assay is affected by multiple factors, such as the prediction of

the CTP cleavage site and presence of a GST tag, which may have influenced the results for

MvALS1. Our data suggest that *MvALS2* and *MvALS5* could potentially be involved in the

evolution of ALS resistance in *M. vaginalis* contrary to the expectation from the nation-wide

survey (Figure 2).

Transcript levels of *MvALS* genes

Since no significant differences were observed in recombinant protein assays and ectopic

expression in *A. thaliana*, we compared the transcript levels of *ALS* genes in *M. vaginalis*

seedlings in the presence or absence of imazosulfuron using mRNA-Seq. *De novo* assembly

followed by the curations of *ALS* contigs, we obtained 84,649 contigs with average length of

802.5 bp.

The transcriptome of herbicide-treated sensitive plants formed a distinct cluster

(Figure 4a), indicating that herbicide treatment triggered massive reprogramming of gene

expression within 24 h in these plants. Statistical analysis revealed 12,845 differentially

expressed contigs between the treated and untreated sensitive populations (Figure S3). In

contrast, gene expression in the resistant population remained almost unchanged following

herbicide treatment, which was expected based on a previous report that herbicide treatment

only slightly suppressed growth in this resistant population (Iwakami et al., 2020). Moreover,

the difference between the untreated sensitive and resistant populations was minor, with only

528 differentially expressed contigs (Figure S3).

391   The transcription of *ALS* genes was almost stable among all the samples (Figure 4b).

The only exception was *MvALS2*, which was slightly downregulated (2.36-fold, $q<0.01$) in

herbicide-treated sensitive plants. As the differences among the populations were small, we

averaged the transcripts per kilobase million (TPM) data of 12 libraries and examined the

transcript level of each *ALS* gene. The TPM value of *MvALS1* and *MvALS3* ranked 2,487 and

3,062, respectively, and these values were much higher than the TPM values of *MvALS2*,

MvALS5, and *MvALS4* (Figure 4c-d). Moreover, respectively 45.3% and 36.9% of the reads

mapped to the *ALS* genes were from *MvALS1* and *MvALS3*, followed by 9.6% from *MvALS2*.

These findings were consistent with the results of gene cloning (Figure S4). Therefore, the

two genes carrying resistance-conferring mutations (*MvALS1* and *MvALS3*) in natural *M.*

vaginalis populations were among the most highly expressed *ALS* genes. The absence of a

significant difference in the transcript levels of each copy between the sensitive and resistant

populations under non-treated conditions implies that the observed diversity of transcription

among *ALS* genes is not an evolutionary response to the selection of ALS herbicides.

Roles of *MvALS1* and *MvALS3* in resistance evolution

We compared the roles of *MvALS1* and *MvALS3* in the resistant phenotype of *M. vaginalis*. Imazosulfuron sensitivity of populations carrying the same mutation in *MvALS1* or *MvALS3* was compared. Three populations with the Pro197Ser mutation showed similar GR_{50} and GR_{90} values, irrespective of the mutated genes (Figure 5). Moreover, similar GR_{50} and GR_{90} values were observed in another set of populations carrying the Asp376Glu mutation. Therefore, the mutated alleles of *MvALS1* or *MvALS3* equally contribute to resistance in *M. vaginalis*.

DISCUSSION

Discerning the evolutionary genetic paths available for adaptation to a particular environment opens the door to predicting evolutionary dynamics and controlling the evolution of the species. In the evolution of weed resistance to herbicides, evolutionary paths are often restricted. In particular, in resistance to ALS herbicides in broadleaf weeds, extreme parallelism in *ALS* genes is often observed, implying that almost no genetic solutions are available to adapt to the herbicides except for the alteration of *ALS* genes. Meanwhile, it remains unclear whether all *ALS* loci represent potential evolutionary paths for herbicide resistance when multiple copies of the *ALS* gene exist in the species of interest. To investigate whether duplicated genes generate additional chances for resistance evolution, we studied the molecular mechanisms underlying the evolution of herbicide resistance in the noxious paddy weed *M. vaginalis*, taking advantage of the highest copy number of *ALS* genes reported so far

in a weed. Through the study of >100 populations, we revealed that the Japanese populations

of *M. vaginalis* utilised only two *ALS* loci, *MvALS1* and *MvALS3*, for TSR evolution among

the five *ALS* loci in the genome. The subsequent molecular characterisations of each *ALS*

gene strongly suggest that the exclusivity was underpinned by the large difference in the

expression of each gene: the transcripts of *MvALS1* and *MvALS3* are more abundant than

those of the other *ALS* genes. This study revealed that among the multiple potential hotspots

in the genome of *M. vaginalis*, expression level confers evolutionary competency in two *ALS*

loci, shaping the biased pattern of resistance-conferring alleles in resistant populations of *M.*

vaginalis in Japan.

The generation of evolutionary innovations derived from gene duplication is often

addressed in evolutionary biology (Panchy, Lehti-Shiu, & Shiu, 2016; Xu, Wang, Guo, He, &

Shi, 2020). While attention has been paid to the acquisition of novel functions or increased

protein abundance by duplicate genes, little is known about their roles in the expansion of

evolutionary paths. When the molecular function of the encoded protein is conserved, all

duplicates can be considered to have potential for evolutionary innovation. In fact, in weeds

with multiple copies of *ALS* genes such as *Schoenoplectiella juncoides* (Sada & Uchino,

2017), *Descurainia sophia* (Xu, Xu, Li, & Zheng, 2020), *Senecio valgaris* (Délye et al., 2016),

and *Echinochloa crus-galli* (Löbmann, Schulte, Runge, Christen, & Petersen, 2021), all the

loci have been used for resistance evolution, suggesting that the duplicate copies increase the

chances of resistance evolution. On the other hand, this study revealed that the independent

selective pressure exerted in many Japanese agricultural fields exclusively selected only two

ALS loci for evolutionary alteration, suggesting that duplicated loci do not necessarily

generate available evolutionary tools, at least in the context of microevolution, even if the

molecular function of the encoded protein is conserved (see discussion below).

Functional characterisation of *MvALS1/2/3/5* revealed no marked differences in

their functions, indicating that *MvALS2* and *MvALS5*, which did not carry

resistance-conferring non-synonymous mutations in natural populations, are the 'potential'

hotspots for resistance evolution, similar to *MvALS1* and *MvALS3*. However, since the two

loci are transcriptionally not very active in seedlings, resistance-conferring mutations in these

ALS genes would not confer herbicide resistance in agricultural fields under the current

expression profiles of the *ALS* genes in *M. vaginalis*. Assuming that the transcript abundance

of *ALS* genes is proportional to the amount of ALS protein, MvALS2 and MvALS5 accounted

for <10% of the ALS pool, whereas MvALS1 and MvALS3 accounted for nearly 40%. This

result indicates that over 90% ALS would be inhibited by herbicides if a resistance-conferring

mutation is present in *MvALS2* or *MvALS5*; this proportion would further decrease in the first

generation of plants carrying a spontaneous mutation due to the heterozygous status, and such

plants would likely not survive herbicide stress in agricultural fields. In addition to relative

abundance, absolute transcript abundance may also play important roles. We observed that

MvALS1 and *MvALS3* were among the most transcriptionally active genes in *M. vaginalis*,

positioned close to the first inflexion point on the sigmoid curve of the transcription ranking

plot, while the other three genes were in the middle or lower parts of the curve (Figure 4c).

Interestingly, three *ALS* genes in hexaploid wheat, each of which confers resistance when

Pro197 is substituted via targeted mutagenesis (Zhang et al., 2019), were also placed close to

the first inflexion point (Figure S5a). These observations, together with the mRNA–GR$_{50}$

relationships in transformed *A. thaliana* (Figure 3b), suggest that a certain degree of transcript

abundance may be required for protein expression. Thus, in order for *MvALS2* and *MvALS5* to

be involved in resistance, an exceptionally large increase in gene expression or two sequential

mutations, a gene up-regulation followed by a non-synonymous substitution, would be

required. The impact of the former, as evident from the *A. thaliana* transformation experiment

(Figure 3c-d), would be only marginal, which would hardly endow appreciable resistance in

the fields, considering that a >10 times higher resistance level is required for wild-type *M.*

vaginalis to tolerate the field dose of imazosulfuron (90 g a.i.·ha^{-1}) (Figure 5b). The latter is

theoretically possible, but would require two independent mutations, one qualitative and one

quantitative, in the same individual, implying that this rarely occurs. The most likely scenario

of *MvALS2/5*-based TSR evolution would be a non-synonymous TSR substitution in

populations with a defect in a highly expressed *ALS* gene, such as the HG1 population (Figure

2a), where the relative dominance of *MvALS2/5* to the whole ALS pool is higher. Currently,

individuals with loss-of-function alleles of *MvALS1* are extremely rare in Japan. If this

genotype increases in the future, individuals with TSR mutations in *MvALS2/5* may arise. It

will be important to study the effects of *MvALS1* deficiency on fitness to estimate the

evolutionary dynamics of this genotype.

Although our data strongly underline the role of gene expression in the biased

representation of mutated genes inducing resistance, we cannot rule out the role of other

factors. Firstly, given the observed higher sequence variations in *MvALS1* and *MvALS3*

(Figure 2e), the spontaneous mutation rate might be higher for these genes, leading to their

overrepresentation. According to the classic evolutionary theory, the spontaneous mutation

rate is random across a genome, and the apparent bias is the consequence of fitness costs.

However, a recent sequencing study revealed that cytogenetic features influenced the

mutation rate itself to a certain degree (Monroe et al., 2020). Thus, the spontaneous mutation

rate may drive the biased representation of genes. Secondly, multi-layered DNA repair

systems (Kimura & Sakaguchi, 2006) may also play a role in the bias. These repair

mechanisms are affected by the flanking sequences, local DNA structure, and gene expression,

among other factors, and do not act uniformly throughout the genome. The local DNA

environment of each *ALS* locus may have affected the manifestation of spontaneous DNA

mutations. Thirdly, our gene expression results must be interpreted with caution, as we

analysed only two populations. There may be natural variations in *ALS* gene expression

among *M. vaginalis* populations. However, similar herbicide responses observed among

populations carrying the same mutations in this study (Figure 5) may alternatively indicate

relatively conserved gene expression among *M. vaginalis* populations in Japan. Large-scale

analyses, such as population transcriptomics, combined with comprehensive investigations of

differential mutation rates in the *ALS* gene family would provide a more holistic view of

evolutionary patterns in *M. vaginalis*.

509   Notably, resistance-conferring mutations were more frequent in *MvALS1* than in

MvALS3 (Figure 2a). This may be because a mutation in *MvALS1* may confer higher

resistance than that in *MvALS3*, which is beneficial to plants, particularly those carrying

mutations conferring low resistance, such as Pro197Leu (e.g. Sada et al., 2013), in the

heterozygous status. However, comparison of plants carrying the identical mutations in

different genes did not reveal significant differences in herbicide sensitivity (Figure 5), which

is in agreement with an independent study (K. Ohta, Y. Fujino & Y. Sada, unpublished).

Furthermore, in transformed *A. thaliana*, *MvALS1* and *MvALS3* conferred a similar level of

resistance when expressed equally (Figure 3b). Thus, although higher substrate affinity was

detected for MvALS1 in our recombinant enzyme assay (Figure 3a), other factors (e.g. the

bias of material sampling) were likely responsible for the bias between *MvALS1* and *MvALS3*.

In conclusion, comprehensive analysis of *ALS* gene sequences in Japanese

populations of *M. vaginalis* revealed that variable transcript levels of these genes are

associated with the biased patterns of mutations for the evolution of herbicide resistance. This

study provides a novel insight that gene expression quantitatively provides evolutionary

competency to potential hotspots across genome for resistance evolution. In other words, gene

expression level can act as a genomic constraint for the parallel evolution of herbicide

resistance. Of note, the Japanese populations of *M. vaginalis* may only reflect a particular

genetic group of this globally distributed weed (mainly in eastern and southern Asia),

considering that Japan is a small island country. Thus, further large-scale studies of the global

populations would provide a more comprehensive understanding of the evolutionary

strategies of herbicide resistance in *M. vaginalis*, which may not be limited to the two

hotspots (*MvALS1* and *MvALS3*) detected in this study. In the era of genomics, it is

anticipated that numerous genomes with more complex DNA structures, such as polyploid

genomes, will be compiled, which will inevitably impose more attention to the roles of

redundant copies during the course of evolution. The perspective and framework of this study

are applicable to research on TSR evolution in agricultural weeds as well as broader studies

focusing on the process of bias for genetic loci in parallel evolution.

ACKNOWLEDGEMENTS

We thank Nihon Nohyaku for providing seeds of *Monochoria vaginalis*, and members of the weed science lab in Kyoto University for the cooperation to collect *M. vaginalis*, Yuuri Nakamura for technical assistance. Computations were partially performed on the NIG supercomputer at ROIS National Institute of Genetics. This work was partly supported by Japan Association for Advancement of Phyto-Regulators (JAPR) grant no. 2017-4, 2018-4 (to A.U. and S.I.), 2019-2 (to S.O. and S.I.).

AUTHOR CONTRIBUTIONS

Project Conception, S.I.; Experimental Design, S.T., T.M., and S.I.; Performance of Experiments, S.T., A.U., S.O., S.I.; Bioinformatics Analysis, S.T. and S.I.; Material Collection and/or Maintenance, S.T., A.U., S.O., C.M., K.H., M.M., N.Y., N.U., Y.T., N.F., E.K., Y.S., T.T., and S.I.; Visualization, S.T. and S.I.; Writing, S.I.; Supervision, T.T., S.I.

DATA ACCESSIBILITY:

mRNA-Seq data have been deposited in the DDBJ Sequence Read Archive (DRA) database with BioProject ID PRJDB11326 (BioSample: SSUB017551).

REFERENCES

Baucom, R. S. (2016). The remarkable repeated evolution of herbicide resistance. *American*
Journal of Botany, 103, 181-183.

Baucom, R. S. (2019). Evolutionary and ecological insights from herbicide-resistant weeds:
what have we learned about plant adaptation, and what is left to uncover? *New*
Phytologist, 223, 68-82.

Bedbrook, J. R., Chaleff, R. S., Falco, S. C., Mazur, B. J., Somerville, C. R., & Yadav, N. S.
(1995).

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics, 30*, 2114-2120.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T.
L. (2009). BLAST+: architecture and applications. *BMC Bioinformatics, 10*, 421.

Chang, A. K., & Duggleby, R. G. (1997). Expression, purification and characterization of
Arabidopsis thaliana acetohydroxyacid synthase. *Biochemical Journal, 327*, 161-169.

Clark, J. W., & Donoghue, P. C. J. (2018). Whole-genome duplication and plant
macroevolution. *Trends in Plant Science, 23*, 933-945.

Czechowski, T., Stitt, M., Altmann, T., Udvardi, M. K., & Scheible, W. R. (2005).
Genome-wide identification and testing of superior reference genes for transcript
normalization in Arabidopsis. *Plant Physiology, 139*, 5-17.

Délye, C., Boucansaud, K., Pernin, F., & Le Corre, V. (2009). Variation in the gene encoding
acetolactate-synthase in *Lolium* species and proactive detection of mutant,
herbicide-resistant alleles. *Weed Research, 49*, 326-336.

Délye, C., Causse, R., & Michel, S. (2016). Genetic basis, evolutionary origin and spread of
resistance to herbicides inhibiting acetolactate synthase in common groundsel
(*Senecio vulgaris*). *Pest Management Science, 72*, 89-102.

Délye, C., Pernin, F., & Scarabel, L. (2011). Evolution and diversity of the mechanisms
endowing resistance to herbicides inhibiting acetolactate-synthase (ALS) in corn
poppy (*Papaver rhoeas* L.). *Plant Science, 180*, 333-342.

Dimaano, N. G., Yamaguchi, T., Fukunishi, K., Tominaga, T., & Iwakami, S. (2020).
Functional characterization of cytochrome P450 CYP81A subfamily to disclose the
pattern of cross-resistance in *Echinochloa phyllopogon*. *Plant Molecular Biology, 102*,
403-416.

Duggleby, R. G., Pang, S. S., Yu, H., & Guddat, L. W. (2003). Systematic characterization of

mutations in yeast acetohydroxyacid synthase. Interpretation of herbicide-resistance

data. *European Journal of Biochemistry, 270*, 2895-2904.

Flagel, L. E., & Wendel, J. F. (2009). Gene duplication and evolutionary novelty in plants.

New Phytologist, 183, 557-564.

Fukano, Y., Guo, W., Uchida, K., Tachiki, Y., & Cornelissen, H. (2020). Contemporary

adaptive divergence of plant competitive traits in urban and rural populations and its

implication for weed management. *Journal of Ecology, 108*, 2521-2530.

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., . . . Regev,

A. (2011). Full-length transcriptome assembly from RNA-Seq data without a

reference genome. *Nature Biotechnology, 29*, 644-652.

Guo, F., Iwakami, S., Yamaguchi, T., Uchino, A., Sunohara, Y., & Matsumoto, H. (2019).

Role of CYP81A cytochrome P450s in clomazone metabolism in *Echinochloa*

phyllopogon. *Plant Science, 283*, 321-328.

Hamamura, K. (2018). Development of herbicides for paddy rice in Japan. *Weed Biology and*

Management, 18, 75-91.

Hamamura, K., Muraoka, T., Hashimoto, J., Tsuruya, A., Takahashi, H., Takeshita, T., &

Noritake, K. (2003). Identification of sulfonylurea-resistant biotypes of paddy field

weeds using a novel method based on their rooting responses. *Weed Biology and*

Management, 3, 242-246.

Heap, I. (2021). The international survey of herbicide resistant weeds. Online. Internet.

Thursday, February 11, 2021. Available www.weedscience.com.

Holm, L. G., Plucknett, D. L., Pancho, J. V., & Herberger, J. P. (1977). *The world's worst*

weeds. Distribution and biology. Honolulu: University of Hawaii.

Iwakami, S., Hashimoto, M., Matsushima, K.-i., Watanabe, H., Hamamura, K., & Uchino, A.

(2015). Multiple-herbicide resistance in *Echinochloa crus-galli* var. *formosensis*, an

allohexaploid weed species, in dry-seeded rice. *Pesticide Biochemistry and Physiology,*

119, 1-8.

Iwakami, S., Kamidate, Y., Yamaguchi, T., Ishizaka, M., Endo, M., Suda, H., . . . Matsumoto,

H. (2019). CYP81A P450s are involved in concomitant cross-resistance to acetolactate

synthase and acetyl-CoA carboxylase herbicides in *Echinochloa phyllopogon*. *New*

Phytologist, 221, 2112-2122.

Iwakami, S., Tanigaki, S., Uchino, A., Ozawa, Y., Tominaga, T., & Wang, G. X. (2020).

Characterization of the acetolactate synthase gene family in sensitive and resistant

622   biotypes of two tetraploid *Monochoria* weeds, *M. vaginalis* and *M. korsakowii*.
623   *Pesticide Biochemistry and Physiology, 165*, 104506.

624   Iwakami, S., Uchino, A., Watanabe, H., Yamasue, Y., & Inamura, T. (2012). Isolation and
625   expression of genes for acetolactate synthase and acetyl-CoA carboxylase in
626   *Echinochloa phyllopogon*, a polyploid weed species. *Pest Management Science, 68*,
627   1098-1106.

628   Kimura, S., & Sakaguchi, K. (2006). DNA repair in plants. *Chemical Reviews, 106*, 753-766.

629   Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature
630   Methods, 9*, 357-359.

631   Li, B., & Dewey, C. N. (2011). RSEM: accurate transcript quantification from RNA-Seq data
632   with or without a reference genome. *BMC Bioinformatics, 12*, 323.

633   Löbmann, A., Schulte, M., Runge, F., Christen, O., & Petersen, J. (2021). Occurrence,
634   resistance factors and cross-resistance patterns to herbicides inhibiting acetolactate
635   synthase (ALS) of *Echinochloa crus-galli* (L.) Pal. Beauv. in Central Europe. *Journal
636   of Plant Diseases and Protection*. Retrieved from   doi:10.1007/s41348-021-00434-1

637   Martin, A., & Orgogozo, V. (2013). The loci of repeated evolution: a catalog of genetic
638   hotspots of phenotypic variation. *Evolution, 67*, 1235-1250.

639   Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing
640   reads. *EMBnet.journal, 17*.

641   Monroe, J. G., Srikant, T., Carbonell-Bejerano, P., Exposito-Alonso, M., Weng, M.-L., Rutter,
642   M. T., . . . Weigel, D. (2020). Mutation bias shapes gene evolution in *Arabidopsis
643   thaliana*. *bioRxiv*. Retrieved from   doi:https://doi.org/10.1101/2020.06.17.156752

644   Motohashi, K. (2015). A simple and efficient seamless DNA cloning method using SLiCE
645   from *Escherichia coli* laboratory strains and its application to SLiP site-directed
646   mutagenesis. *BMC Biotechnology, 15*, 47.

647   Murashige, T., & Skoog, F. (1962). A revised medium for rapid growth and bio assays with
648   tobacco tissue cultures. *Physiologia Plantarum, 15*, 473-497.

649   Murray, M., & Thompson, W. F. (1980). Rapid isolation of high molecular weight plant DNA.
650   *Nucleic Acids Research, 8*, 4321-4326.

651   Ohsako, T., & Tominaga, T. (2007). Nucleotide substitutions in the acetolactate synthase
652   genes of sulfonylurea-resistant biotypes of *Monochoria vaginalis* (Pontederiaceae).
653   *Genes & Genetic Systems, 82*, 207-215.

Ono, H., Ishii, K., Kozaki, T., Ogiwara, I., Kanekatsu, M., & Yamada, T. (2015). Removal of

redundant contigs from *de novo* RNA-Seq assemblies via homology search improves
accurate detection of differentially expressed genes. *BMC Genomics, 16*, 1031.

Panchy, N., Lehti-Shiu, M., & Shiu, S. H. (2016). Evolution of gene duplication in plants.
Plant Physiology, 171, 2294-2316.

Peters, B., & Strek, H. J. (2018). Herbicide discovery in light of rapidly spreading resistance
and ever-increasing regulatory hurdles. *Pest Management Science, 74*, 2211-2215.

Powles, S. B., & Yu, Q. (2010). Evolution in action: plants resistant to herbicides. *Annual*
Review of Plant Biology, 61, 317-347.

Ritz, C., Baty, F., Streibig, J. C., & Gerhard, D. (2015). Dose-response analysis using R. *PLoS*
One, 10, e0146021.

Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010). edgeR: a Bioconductor package
for differential expression analysis of digital gene expression data. *Bioinformatics, 26*,
139-140.

Rozas, J., Ferrer-Mata, A., Sanchez-DelBarrio, J. C., Guirao-Rico, S., Librado, P.,
Ramos-Onsins, S. E., & Sanchez-Gracia, A. (2017). DnaSP 6: DNA sequence
polymorphism analysis of large data sets. *Molecular Biology and Evolution, 34*,
3299-3302.

Sada, Y., Ikeda, H., & Kizawa, S. (2013). Resistance levels of sulfonylurea‑resistant
Schoenoplectus juncoides (Roxb.) Palla with various Pro197 mutations in acetolactate
synthase to imazosulfuron, bensulfuron‑methyl, metsulfuron‑methyl and
imazaquin‑ammonium. *Weed Biology and Management, 13*, 53-61.

Sada, Y., & Uchino, A. (2017). Biology and mechanisms of sulfonylurea resistance in
Schoenoplectiella juncoides, a noxious sedge in the rice paddy fields of Japan. *Weed*
Biology and Management, 17, 125-135.

Schmittgen, T. D., & Livak, K. J. (2008). Analyzing real-time PCR data by the comparative
C(T) method. *Nature Protocols, 3*, 1101-1108.

Singh, B., Schmitt, G., Lillis, M., Hand, J. M., & Misra, R. (1991). Overexpression of
acetohydroxyacid synthase from *Arabidopsis* as an inducible fusion protein in
Escherichia coli: production of polyclonal antibodies, and immunological
characterization of the enzyme. *Plant Physiology, 97*, 657-662.

Svitashev, S., Young, J. K., Schwartz, C., Gao, H., Falco, S. C., & Cigan, A. M. (2015).
Targeted mutagenesis, precise gene editing, and site-specific gene insertion in maize
using Cas9 and guide RNA. *Plant Physiology, 169*, 931-945.

Waselkov, K. E., Regenold, N. D., Lum, R. C., & Olsen, K. M. (2020). Agricultural
adaptation in the native North American weed waterhemp, *Amaranthus tuberculatus*
(Amaranthaceae). *PLoS One, 15*, e0238861.
Wedger, M. J., & Olsen, K. M. (2018). Evolving insights on weedy rice. *Ecological Genetics*
and Genomics, 7-8, 23-26.
Weng, M. L., Becker, C., Hildebrandt, J., Neumann, M., Rutter, M. T., Shaw, R. G., . . .
Fenster, C. B. (2019). Fine-grained analysis of spontaneous mutation spectrum and
frequency in *Arabidopsis thaliana. Genetics, 211*, 703-714.
Xu, S., Wang, J., Guo, Z., He, Z., & Shi, S. (2020). Genomic convergence in the adaptation to
extreme environments. *Plant Commun, 1*, 100117.
Xu, Y., Xu, L., Li, X., & Zheng, M. (2020). Investigation of resistant level to
tribenuron-methyl, diversity and regional difference of the resistant mutations on
acetolactate synthase (ALS) isozymes in *Descurainia sophia* L. from China. *Pesticide*
Biochemistry and Physiology, 104653.
Ye, C. Y., Tang, W., Wu, D., Jia, L., Qiu, J., Chen, M., . . . Fan, L. (2019). Genomic evidence
of human selection on Vavilovian mimicry. *Nature Ecology & Evolution, 3*,
1474-1482.
Yu, Q., & Powles, S. B. (2014). Resistance to AHAS inhibitor herbicides: current
understanding. *Pest Management Science, 70*, 1340-1350.
Zhang, R., Liu, J., Chai, Z., Chen, S., Bai, Y., Zong, Y., . . . Gao, C. (2019). Generation of
herbicide tolerance traits and a new selectable marker in wheat using base editing.
Nature Plants, 5, 480-485.

 Figure legends

 Figure 1. *Monochoria vaginalis* studied in this work

 (a) Infestation of *M. vaginalis* in paddy fields in Japan. (b) Collection sites of *M. vaginalis*.

 MvALS1, populations carrying resistance-conferring mutations in *MvALS1*; *MvALS3*,

 populations carrying resistance-conferring mutations in *MvALS3*; *MvALS1&3*, populations

 carrying resistance-conferring mutations in *MvALS1* and *MvALS3*; N.D., populations with no

 resistance-conferring mutations in *ALS* genes. (c) Methods for resistance diagnosis. In rooting

 assay, roots were cut at 1 cm from the base of shoots and placed in bensulfuron-methyl

 solution for 7-10 days. Plants with emerging roots were diagnosed as resistant. In the pot

 assay, seeds were germinated in a tube, followed by transplantation to pots filled with

 autoclaved soil. The recommended dose of imazosulfuron (90 g a.i.·ha^{-1}) was applied at the

 three-leaf stage. Plant survival was evaluated at 3 weeks post application. DNA extracted

 from the plants subjected to sensitivity assays were used for *ALS* gene sequencing.

 Figure 2. *ALS* genes of *Monochoria vaginalis*

 (a) Status of *ALS* genes in 68 resistant populations analysed in this study. YNZ1 carries

 mutations both in *MvALS1* and *MvALS3*. AIZ2 and KR1 populations carry two independent

 mutations in *MvALS1*. F.S., frameshift mutation. (b) Patterns of resistance-conferring

 mutations. (c) Patterns of DNA conversions of resistance-conferring mutations. (d) Schematic

 representation of frameshifted alleles found in *MvALS1*, *MvASL4*, and *MvALS5*. *MvALS4*

 alleles encoding the canonical form of ALS were not detected in the present study. Triangles

 represent a premature stop codon. (e) Nucleotide variations in the five *ALS* genes. The coding

sequences of 68 resistant and 31 sensitive populations were analysed. The plot shows

nucleotide diversity calculated at each nucleotide position. Positions causing an amino acid

replacement that confers resistance are indicated in red. Nucleotide diversity of the entire

genes (π) and haplotype numbers were calculated.

Figure 3. Enzymatic functions of ALS encoded by each gene of *Monochoria vaginalis*

(a) Enzymatic properties of recombinant ALS. (b) Transformation of *Arabidopsis thaliana.*

ALS genes were isolated from a sensitive plant, and were mutagenised to encode the

Pro197Ser form of ALS. The native form of *MvALS1* was also transformed. (c) *MvALS1*

expression in transgenic *A. thaliana* lines. Bars represent standard error (n=3). (d) Response

of transgenic *A. thaliana* to 10 and 100 nM imazosulfuron. (e) Relationship of transcript level

and imazosulfuron sensitivity in transgenic *Arabidopsis thaliana* expressing each *ALS* gene

with Ser at Pro197 codon. (f) Response of transgenic *A. thaliana* to 10 nM imazosulfuron.

The plants correspond to the lines indicated by circles in Figure 3b.

Figure 4. Expression of *ALS* genes in *Monochoria vaginalis*

(a) Transcriptome of sensitive (S) and resistant (R) *M. vaginalis* with or without herbicide

imazosulfuruon (H) application. Transcriptome of the R plants was almost unchanged, while

that of the S plants was drastically affected. (b) Transcripts per kilobase million (TPM) of

each *ALS* gene. *$q<0.05$. (c) Expression rank in *M. vaginalis* transcriptome. Data represent

the average of 12 mRNA-Seq samples. (d) Relative transcript abundance of *ALS* genes in *M.*

vaginalis.

Figure 5. Herbicide sensitivity of *Monochoria vaginalis* carrying a mutation in *MvALS1*

and *MvALS3*

(a) *M. vaginalis* 3 weeks post imazosulfuron application. (b) Dose response to imazosulfuron.

Bars represent standard error (n=3). (c) Doses that inhibited the growth to 50% (GR_{50}) and

90% (GR_{90}).

(a)

(b)

(c)

1. Rooting assay

ALS gene sequencing

2. Recommended rate pot assay

(a)

Population	MvALS1	MvALS2	MvALS3	MvALS4	MvALS5
YOK1	D376E	–	–	F.S.	–
DSN1	P197H	–	–	F.S.	–
DSN2	P197S	–	–	F.S.	–
DSN3	D376E	–	–	F.S.	–
ODT1	P197S	–	–	F.S.	–
AKT1	P197S	–	–	F.S.	–
AKT2	P197S	–	–	F.S.	–
AKT3	D376E	–	–	F.S.	–
KUR1	P197H	–	–	F.S.	–
KUR2	P197A	–	–	F.S.	–
MSM1	P197S	–	–	F.S.	–
TKU1	P197L	–	–	F.S.	F.S.
TKU2	D376E	–	–	F.S.	–
TKU3	P197S	–	–	F.S.	–
TKU4	P197S	–	–	F.S.	–
TKU5	P197H	–	–	F.S.	–
NDA1	D376N	–	–	F.S.	F.S.
ETZ1	P197H	–	–	F.S.	–
KTK1	P197T	–	–	F.S.	–
TRG1	D376E	–	–	F.S.	F.S.
TRG3	D376E	–	–	F.S.	F.S.
NRS1	P197S	–	–	F.S.	–
ANP1	A205T	–	–	F.S.	–
AIS1	P197H	–	–	F.S.	F.S.
KST1	P197S	–	–	F.S.	–
TKS1	P197H	–	–	F.S.	–
TKS4	P197H	–	–	F.S.	–
TKS8	P197H	–	–	F.S.	–
TNB1	D376E	–	–	F.S.	–
KU1	D376E	–	–	F.S.	–
KMO1	P197S	–	–	F.S.	–
ISE1	P197S	–	–	F.S.	–
IGA1	A205V	–	–	F.S.	–
IGA2	P197L	–	–	F.S.	F.S.
IGA3	P197S	–	–	F.S.	–
TSU3	P197S	–	–	F.S.	–
TOT1	P197L	–	–	F.S.	–
YMG2	P197S	–	–	F.S.	–
FUK1	P197S	–	–	F.S.	–
TKE1	P197S	–	–	F.S.	–
TKD1	P197H	–	–	F.S.	–
SGA1	D376E	–	–	F.S.	–
AIZ2	P197S, D376E	–	–	F.S.	F.S.
KR1	P197S, D376E	–	–	F.S.	F.S.
MRK2	P197L	–	–	F.S.	F.S.
AGN1	P197S	–	–	F.S.	–
NGT1	P197S	–	–	F.S.	–
YNZ1	D376E	–	P197S	F.S.	–
KTG1	–	–	P197S	F.S.	–
TYM2	–	–	P197A	F.S.	–
USK1	–	–	P197S	F.S.	–
HIT1	–	–	P197S	F.S.	–
MIU1	–	–	P197S	F.S.	–
SMM23	–	–	P197S	F.S.	–
SMM3	–	–	P197S	F.S.	F.S.
SMM4	–	–	P197S	F.S.	–
NRS2	–	–	D376E	F.S.	–
IWT1	–	–	D376E	F.S.	–
TKS6	–	–	P197H	F.S.	–
NTN1	–	–	P197L	F.S.	–
MIM1	–	–	P197S	F.S.	–
KRM1	–	–	P197S	F.S.	–
FK1	–	–	D376E	F.S.	–
AIZ1	–	–	P197T	F.S.	–
OMH1	–	–	P197S	F.S.	–
OMH2	–	–	P197S	F.S.	–
YMG1	–	–	D376E	F.S.	–
HG1	F.S.	–	P197A	F.S.	–

(b) *MvALS1*, *MvALS3*

Number of mutation — categories: P197S, P197H, P197A, P197L, P197T, A205T, A205V, D376E, D376N

(c) Number of mutation at P197, A205, D376

Legend:
- GC→AT
- GC→TA
- GC→CG
- AT→TA
- AT→CG

(d)

MvALS1
ATG — STOP
13-bp deletion
(+798 – +810)

MvALS4
ATG — STOP
Not found
1-bp insertion
(+974 – +976)

MvALS5
ATG — STOP
40-bp deletion
(+1856 – +1898)

(e)

MvALS1 — π = 0.00136, Haplotype = 33
MvALS2 — π = 0.00024, Haplotype = 4
MvALS3 — π = 0.00123, Haplotype = 11
MvALS4 — π = 0.00050, Haplotype = 2
MvALS5 — π = 0.00027, Haplotype = 4

Nucleotide diversity (site) vs Nucleotide position

(a)

Gene	Reaction	k_{cat} (min^{-1})	K_m (mM)	k_{cat}/K_m (min^{-1}mM^{-1})
(control)		—	—	—
MvALS1		0.95 ± 0.06	8.53 ± 1.51	0.111 ± 0.021
MvALS2		0.83 ± 0.07	13.13 ± 3.03	0.063 ± 0.016
MvALS3		0.78 ± 0.05	17.25 ± 2.42	0.045 ± 0.007
MvALS5		0.81 ± 0.04	14.51 ± 1.70	0.056 ± 0.007

(b)

(c)

(d)

(e)

(f)

(a)

(b)

(c)

(d)

(a)

	S	P197S			D376E	
(g a.i. ha⁻¹)		ALS1	ALS1	ALS3	ALS1	ALS3

(b) Field dose

(c)

Population	Mutation	Gene	GR50		GR90	
			g a.i. ha⁻¹	R/S	g a.i. ha⁻¹	R/S
KST1	P197S	*MvALS1*	15799	2000	308232	30219
KMO1	P197S	*MvALS1*	5029.9	637	211830	20768
SMM23	P197S	*MvALS3*	5553.9	703	630419	61806
AKT3	D376E	*MvALS1*	197.5	25	29757	2917
NRS2	D376E	*MvALS3*	323	41	17118	1678
SMM13	-	-	7.9	1	10.2	1