Title

Telencephalic outputs from the medial entorhinal cortex are copied directly to the

hippocampus

Authors and Affiliations

Sau Yee Tsoi, Merve Öncül, Ella Svahn, Mark Robertson, Zuzanna Bogdanowicz, Christina

McClure and Gülşen Sürmeli*

*Corresponding Author

University of Edinburgh, Centre for Discovery Brain Sciences, Edinburgh, UK

G.S., S.Y.T., M.Ö., E.S., M.R., Z.B., C.M.

Simons Initiative for the Developing Brain

G.S.

Current addresses

University College London, UK

E.S.

NHS Greater Glasgow and Clyde, UK

M.R.

VectorBuilder Inc., 1010 W 35th Street, Suite 515 Chicago, IL 60609, USA

C.M.

Abstract

Complementary actions of the neocortex and the hippocampus enable encoding and long-term storage of experience-dependent memories. Standard models for memory storage assume that sensory signals reach the hippocampus from superficial layers of the entorhinal cortex (EC). Deep layers of the EC, on the other hand, relay hippocampal outputs to telencephalic structures including many parts of the neocortex. Here we show that cells in Layer 5a of the medial EC send a copy of their telencephalic outputs back to the CA1 region of the hippocampus. Combining cell-type specific anatomical tracing, high-throughput RNA-sequencing based projection mapping and optogenetics-aided circuit mapping, we show that these projections have a unique topography and target pyramidal cells and interneurons in CA1. Our results suggest that rather than serving as a relay, deep EC may coordinate hippocampal-neocortical interactions in memory consolidation.

Introduction

Interplay between the hippocampus and the neocortex is a central tenet of theories for systems memory consolidation (Marr, 1971; McClelland et al., 1995). The entorhinal cortex (EC) mediates the two-way communication between the neocortex and the hippocampus by channelling inputs to and outputs from the hippocampus. During a new experience the superficial layers relay convergent inputs from the cortex to the hippocampus where initial conjunctive representations of our everyday experiences are generated. Subsequently during rest or sleep, hippocampal output is redistributed to the brain (Squire et al., 2015; Frankland and Bontempi 2005). Leading theoretical models consider the deep layers of EC as a relay of hippocampal output that reinstates neocortical activation patterns representing memories enabling neocortex-wide synaptic changes required for long-term storage (Koster et al., 2018; Kumaran et al., 2016; McClelland et al., 1995; Schapiro et al., 2017).

Recent discoveries revealed complex and distinct input-output interactions of molecularly defined subtypes of neurons in the deep layers of medial EC (MEC), suggesting their functions extend beyond passing on hippocampal outputs. Layers 5a (L5a) and 5b (L5b) of the EC are genetically distinct with differential input-output organisations. L5a but not L5b is the sole origin of the long-range telencephalic outputs of the EC (Sürmeli et al., 2015; Ohara et al., 2018, Kitamura et al., 2017), whereas L5b, but not L5a, is directly influenced by superficial layers of EC (Beed et al., 2020; Sürmeli et al., 2015). Inputs from the dorsal hippocampus preferentially target L5b over L5a (Rozov et al., 2020; Sürmeli et al., 2015; Wozny et al., 2018) but ventral hippocampus preferentially target L5a (Rozov et al. 2020). Moreover, the capacity of deep layers to integrate neocortical inputs with hippocampal output (Czajkowski et al. 2013; Beed et al. 2020; Sürmeli et al. 2015) suggest more complex roles.

An intriguing possibility, noted in experiments with classic retrograde tracers, is that deep

layers of MEC might also project to the hippocampal CA1 region, which provides the deep

EC with hippocampal output (Köhler, 1985; Witter and Amaral, 1991). Such recurrent

connectivity between CA1 and the deep MEC could be important in coordinating

hippocampal-entorhinal activity during memory consolidation (Ólafsdóttir et al. 2016),

provide a shortcut for hippocampal outputs to re-enter the hippocampus (Kumaran and

McClelland, 2012; Kloosterman et al., 2003) or have a more global coordination function

encompassing the hippocampus and the neocortex (Khodagholy et al., 2017). Despite its

potential functional significance, the key properties of this projection pathway including

whether it originates from a particular subset of neurons and which neurons it targets in the

hippocampus remain unknown.

Here we reveal that neurons in L5a, but not L5b, of the MEC project to hippocampal CA1.

We establish that the telencephalon-projecting neurons copy their outputs directly to

pyramidal and subclasses of interneuron populations. Our results suggest that Layer 5a of

the MEC has a dual function in coordinating neocortical and hippocampal networks.

Results

To find out whether neurons in either L5a or L5b project to the hippocampus we injected a

retrograde adeno-associated viral vector (AAV) expressing green fluorescent protein (GFP)

into intermediate CA1. As well as finding retrogradely-labelled cell bodies in layer 3 of MEC,

which contains neurons that are well-established as a critical input to stratum lacunosum of

CA1 (Naber et al., 2001), we also found labelled cell bodies in L5a, which we identified by

immunostaining against the ETS variant transcription factor-1 (Etv1) protein. Retrogradely-

labelled neurons were not found in L5b (Fig. 1a; Sürmeli et al., 2015).

Figure 1. Origin and distinct topography of entorhinal deep layer projections to CA1
a, Left: A retrograde AAV virus was injected in the intermediate hippocampus. Middle: Entorhinal area on a horizontal brain section. Retrogradely-labelled neurons were found in L3 as well as L5a, marked by Etv1 expression (Scale bar: 250 µm). Inset: high-magnification image of boxed region in left panel showing co-expression of GFP and Etv1 (Scale bar: 50 µm).
b, Left: *Rbp4-Cre* mice were injected in the deep MEC with a Cre-dependent virus expressing GFP. Middle: expression of GFP overlaps with expression of Etv1 (88.6%, n = 3 mice, 1866 cells; Scale bar: 250 µm). Inset: high-magnification of boxed region in middle panel (Scale bar: 100µm).
c, Coronal section showing the distribution of axons of MEC L5a cells in CA1 (Scale bar: 500 µm). Inset: high-magnification image of boxed region in left panel (Scale bar: 100 µm).
d, Quantification of fluorescence intensity across the layers of CA1. The area between 0 and 1 in **c** was divided into 20 bins, and the mean fluorescence intensity was calculated for each bin. Values shown are normalised mean fluorescence intensity values (n = 6 mice, 2 sections per brain).
e. Left: Horizontal section showing the injection site at low exposure. Right: A high exposure image from the hippocampal area of the same tissue showing the distribution of L5a axons across the proximal-distal axis of CA1 (Scale bar: 250 µm).
f. Normalised mean fluorescence density values in proximal versus distal CA1 (n = 5 mice).
g. Normalised mean fluorescence density values in deep versus superficial SP (n = 6 mice). Within SP, axon density in deep SP was significantly higher than density in superficial SP (paired one-tailed Wilcoxon signed rank test, p = 0.0156, effect size = 0.899, n = 6).

To further investigate this projection, we identified and validated a transgenic mouse line that

enables specific genetic access to L5a. In this mouse line, Cre expression is driven by the

promoter of the retinol binding protein 4 gene (*Rbp4-Cre*; Gerfen et al., 2013). Following

injection of a Cre-dependent AAV expressing GFP or mCherry into the deep MEC of *Rbp4-*

Cre mice, fluorescent signal was detected mainly in L5a (Fig. 1b and Extended Data Fig.

1a). The majority of labelled L5a neurons expressed Etv1 (88.6%, n = 3 mice, 1866 cells;

Fig. 1b), similar to L5a neurons that were retrogradely labelled from the hippocampus

(78.8%, n = 3 mice, 749 cells) or from cortical (89.2%, n = 8 mice, 2175 cells) and

subcortical target areas (89.0%, n = 7 mice, 1771 cells; Extended Data Fig. 1b). Thus, like

telencephalon-projecting L5a neurons, deep MEC neurons that project to the hippocampus

are distinguished by expression of Etv1 and can be genetically accessed using *Rbp4-Cre*

mice.

To what extent does the projection from L5a to CA1 differ from previously-identified

projections arising from superficial layers of MEC? In contrast to projections from superficial

MEC, which target either stratum lacunosum (SL) or stratum moleculare (SM) of CA1

(Kitamura et al., 2014), axons from L5a neurons were primarily found in deep stratum

pyramidale (SP) and stratum lacunosum (SL) but were sparse in stratum moleculare (SM;

Fig. 1c,d,e,g). Axons from L5a also differed in their proximo-distal organisation within CA1.

Whereas projections from superficial layers of MEC favour proximal CA1 (Tamamaki and

Nojyo, 1995), axons from L5a had highest density in distal CA1 (paired one-tailed Wilcoxon

signed rank test, p = 0.0312, n = 5 mice; Fig. 1e,f and Extended Data Fig. 1c). We observed

the same topography when axon terminals were labeled using fluorescently-tagged

presynaptic proteins confirming that the topography is not the result of labelling of passing

axons (Extended Data Fig. 1d,e). This distinct topography suggests a unique functional role

for the projections from L5a of MEC to CA1.

Our findings suggest two intriguing possibilities for the organisation of output neurons in L5a

of the MEC. The hippocampus-projecting neurons in L5a may be distinct from the

telencephalon-projecting neurons, implying separate processing within the deep MEC layers

of signals to the hippocampus and telencephalon. Alternatively, the same neurons in L5a

may project to both the telencephalon and the hippocampus, implying that telencephalic

outputs from MEC are copied back to the hippocampus. To distinguish between these

possibilities we used a combinatorial viral labelling strategy. We restricted reporter gene

expression to subpopulations of L5a cells that project to specific targets by injecting a

retrograde AAV expressing Cre recombinase in either the retrosplenial cortex (RSC) or

nucleus accumbens (NucAcb), and a Cre-dependent fluorescent reporter virus into the MEC

(Fig. 2a,b and Extended Data Fig. 2a,b,c,d). With this approach we detected in CA1

fluorescently-labelled axons originating from both RSC-projecting and NucAcb-projecting

L5a neurons (Fig. 2c,d). The axon distribution across layers was similar to when L5a

neurons were labelled in bulk using the *Rbp4-Cre* mouse line (Fig. 2e,f). These results

establish the principle that axon projections from L5a of MEC to the hippocampus are

collaterals of projections to telencephalic targets. However, it is unclear from these

experiments whether this principle applies to projections from L5a to all of its many

telencephalic targets.

Entorhinal cortex outputs have diverse targets, including the entire cortical mantle as well as

parts of the basal ganglia and amygdala (Swanson and Köhler, 1986). To test whether axon

collateralisation to the hippocampus is a general feature of all telencephalon-targeting

neurons, we used Multiplex Analysis of Projections by Sequencing (MAPseq; Kebschull et

al., 2016), a high-throughput method of mapping single-neuron projections by tagging them

with unique RNA sequences or barcode RNAs. We injected the MAPseq barcode RNA virus

library into the full dorso-ventral extent of the deep MEC (Extended Data Fig. 3a) and

quantified barcode RNA expression in tissue collected from dorsal and ventral MEC, dorsal

and ventral hippocampus and three major divisions of the brain (isocortex, cerebral nuclei

(CNU), olfactory areas and cortical subplate (Olf/Ctxsp), (Fig 3a, see methods for detailed

description). The majority of barcodes that were detected in any one

Figure 2. Retrosplenial cortex and Nucleus accumbens outputs of the MEC is copied to CA1
a, b, Experimental strategy. In wild-type mice, a retrogradely-transported virus expressing Cre was injected in either the retrosplenial cortex (RSC; **a**) or nucleus accumbens (NucAcb; **b**), while a Cre-dependent reporter virus was injected in the deep MEC.
c, d, Hippocampal axon collaterals of RSC-projecting **(c)** or NucAcb-projecting MEC neurons (**d**; Scale bar: 250 µm). Inset: high-magnification image of boxed region in left panels showing distribution of axons across CA1 layers (Scale bar: 100 µm).
e, f, Quantification of mean fluorescence intensity across the layers of CA1 when only RSC-projecting (**e**) or NucAcb-projecting (**f**) MEC neurons were labelled (n = 3 mice for each group). Axon topography is similar to when L5a neurons are marked globally in the *Rbp4-Cre* line.

of the three target divisions (isocortex, CNU or Olf/Ctxsp) were also detected in the

hippocampus, suggesting that collateral axons to the hippocampus are a common feature of

telencephalon-projecting neurons (Fig. 3b). The results were similar regardless of whether

neurons were located in dorsal or ventral MEC (Extended Data Fig. 3b), suggesting that

collateralisation does not depend on the neuron's position in the dorso-ventral axis of the

MEC. Using barcode counts as a measure of projection strength (Kebschull et al., 2016) we

further found that dorsal MEC neurons preferentially project to dorsal aspects of the

hippocampus and ventral MEC neurons to the ventral aspects, indicating that the back-

projections to the hippocampus are topographically organised (Fig. 3c). Together these

results reveal general organising principles by which all projections from MEC to the

telencephalon are copied back to the hippocampus.

Figure 3. Telencephalic outputs are widely copied onto the hippocampus
a, Experimental strategy. A MAPseq virus library was injected into deep MEC (n = 3). 44 hrs later, mice were sacrificed and brains were sectioned sagittally at 400 μm thickness. Tissue from 5 regions (Isocortex, CNU, Olf/Ctxsp, dHip and vHip), as well as dorsal and ventral MEC were then dissected and collected in tubes. Brainstem tissue was collected as a negative control. Tissue was further processed for RNA extraction and next generation sequencing.
b, Nearly all barcodes detected in CNU, Olf/Ctxsp and Isocortex were also detected in the hippocampus (Isocortex: 98.6% ± 0.06%; CNU: 99.2% ± 0.08%; Olf/Ctxsp: 99.0% ± 0.4%; n = 3 mice, 5358 barcodes).
c, Dorso-ventral topography of projections revealed by relative barcode counts. For each barcode, counts were normalised to show the relative projection strength of each neuron to dorsal and ventral hippocampus (n = 3 mice).

Which cell types within the hippocampus receive signals from L5a of MEC? While the

compartmentalised arrangement of axons from L5a of MEC suggests selective targeting of

CA1 layers, axonal topography does not necessarily reflect functional connectivity.

Therefore, we targeted viral vectors expressing a channelrhodopsin2-mCherry conjugate

to L5a neurons in MEC and tested for connectivity using whole-cell ex-vivo patch clamp

recordings from CA1 neurons.

We focused initially on responses of pyramidal cells. Brief light pulses evoked depolarising

sub-threshold postsynaptic potentials (PSPs) in 64% of SP pyramidal neurons recorded at

their resting membrane potential (n = 64 cells, 27 mice, Fig. 4a,b,c, and Extended Data Fig.

4e). The response probability of neurons in distal CA1 was not significantly different than

neurons in proximal CA1 (Fig. 4d). The majority of PSPs (20 out of 26) maintained their

polarity when the membrane potential was adjusted from rest (Vm = -66.7 ± 0.6) to -50 mV,

indicating that they were glutamatergic. In these neurons EPSPs were maintained when

$GABA_A$ receptors were blocked with Gabazine, but were abolished by the AMPA receptor

antagonist NBQX, demonstrating that they are glutamatergic (Fig. 4e). The responses had

short latencies (3.04 ± 0.26 ms, n = 20 cells, 15 mice) that were relatively invariant from trial

to trial (Fig. 4f) and were independent of the response's position within a train of stimuli,

indicating that they are monosynaptic (Extended Data Fig. 4d). Consistent with this,

responses were sensitive to bath application of tetrodotoxin (TTX) but recoverable after

application of 4-Aminopyridine (4-AP; Fig. 4g; Petreanu et al., 2007). Together, these data

demonstrate that projections from L5a of the MEC provide glutamatergic excitatory inputs

directly to pyramidal neurons in proximal and distal CA1.

A smaller population of pyramidal neurons showed PSPs with characteristics of indirect

inhibitory connections (6 out of 26). These responses either reversed polarity when the cell

was held above the chloride reversal potential (Fig. 4h) or had an early depolarising and a

late hyperpolarising component (Extended data Fig. 4b). Application of Gabazine either

completely blocked the PSPs (Fig. 4i) or revealed a larger excitatory component that was

sensitive to application of NBQX (Extended data Fig. 4b). Thus, inputs from L5a of the MEC

also recruit local interneurons that provide inhibitory input to SP pyramidal neurons.

Figure 4. L5a neurons of the MEC provide direct excitatory and indirect inhibitory inputs to pyramidal cells in CA1

a, Experimental design showing viral expression, placement of patch-clamp electrode and light delivery over CA1.

b, Examples of biocytin-filled pyramidal neurons recorded in CA1.

c, Top: an example electrical recording of a pyramidal neuron at rest (-67 mV) showing depolarizing responses upon 3 ms light stimulation (top) (blue line, Scale bar: 1 mV, 10 ms). Bottom: train of action potentials upon 200pA step-current injection (bottom). (Scale bar: 20 mV, 100 ms)

d, Proportion of responsive pyramidal neurons located in the distal and proximal halves of CA1. The difference in proportions was not significant (X-squared = 1.0932, df = 1, p-value = 0.2958, Chi-squared test, n = 12 cells in proximal and n = 39 cells in distal CA1).

e, Effects of bath application of Gabazine (orange, n = 10 cells, 9 mice) and NBQX (green, n = 5 cells) on PSPs recorded from a pyramidal neuron (Scale bar: 0.5 mV, 10 ms) and a summary plot of PSP amplitude measurements for all tested pyramidal neurons. Note that some neurons were only treated with Gabazine which did not cause a significant change in amplitudes (p = 0.97, 2-tailed Student's t-test, n = 10 cells, 9 mice).

f, An example of ten consecutive PSP responses recorded from a single pyramidal neuron illustrates the short and invariant latency of PSPs (Scale bar: 0.5 mV, 10 ms) and a cumulative probability plot of standard deviation of latencies for neurons with PSP responses that were > 1 mV in amplitude (n = 20 cells).

g, Effects of bath application of TTX (orange) and 4-AP (green) on PSPs recorded from a pyramidal neuron (Scale bar: 0.5 mV, 10 ms) and a summary plot of changes in PSP amplitudes for all tested pyramidal neurons. TTX application abolished responses (n = 5 cells, 5 mice, p = 0.01, 2-tailed Student's t-test).

h. An example inhibitory PSP response recorded from a pyramidal neuron upon 10 Hz light stimulation (blue bars). Response polarity reversed when the neuron's membrane potential was adjusted to -50 mV and was abolished after application of Gabazine (Scale bars: 0.2 mV, 100 ms). Inset shows the long latency (> 10 ms) of PSP onset indicating polysynaptic connectivity.

i. Effects of bath application of Gabazine on the response amplitude of long latency PSPs (n = 3 cells).

To identify which interneurons in CA1 were targets of L5a projections, we tested responses

of interneurons in all layers. In SP we distinguished interneurons from principal cells either

by classifying them based on biophysical properties (fast-spiking: SP_{FS} and non-fast spiking:

SP_{NFS}; Extended Data Fig. 4a and see methods), or by recording from fluorescently-labelled

GABAergic or parvalbumin-expressing (PV+) inhibitory neurons in double transgenic mice

(Fig 5a,b, and Extended Data Fig. 4c). Stimulation of L5a axons elicited both sub-threshold

and supra-threshold responses primarily in SP, SR and SL interneurons (Fig. 5c,d and

Extended Data Fig. 4e). Depolarising PSPs were observed in 64% of the SP_{FS} including PV+

interneurons, 46% of the SP_{NFS}, 64% of SR and lower proportions of recorded neurons in SL

(31%), SM (23%) and SO (16%). Except for interneurons in SO, PSPs in all layers showed

characteristics of monosynaptic connectivity (Fig. 5g,h,i,j and Extended Data Fig. 4g,h) and

AMPA receptor-mediated glutamatergic synaptic transmission (Fig. 5e,f and Extended Data

Fig. 4f). Therefore, projections from L5a of MEC recruit CA1 interneurons through

monosynaptic inputs driven by excitatory glutamatergic synapses.

Figure 5. L5a neurons of the MEC provide direct excitatory inputs to interneurons in CA1.

a, Experimental design showing viral expression, patch pipette and light stimulation. Interneurons are marked by the expression of venus fluorescent protein driven by the vesicular GABA transporter (*Vgat*) gene's promoter.

b, Biocytin-filled neurons in the *Rbp4-Cre X Vgat-Venus* double transgenic mouse line. Fluorescent labelling facilitated patch-clamp recordings made from GABAergic interneurons and distinguishing SR, SL and SM layers.

c, Representative examples of depolarising responses recorded at resting membrane potential following 3 ms blue light stimulation (blue line) of L5a axons in CA1 (left) (Scale bar: 1 mV, 10 ms). Neurons in SO and SM were typically not responsive (< 30%). Responses from SR neurons were on average larger than the responses from neurons in other layers (Scale bar: 1 mV, 10 ms). Neurons' spiking response to injecting 200 pA current (right) (Scale bar: 10 mV, 100 ms).

d, Proportion of responsive interneurons in all layers of CA1 recorded at resting membrane potential. Green highlighted segment corresponds to the proportion of cells with depolarising membrane potentials; grey highlighted segment corresponds to neurons with no change in their membrane potential.
SO: n = 18 cells, 10 mice;
SP_{FSint} : n = 33 cells, 8 *Rbp4-Cre X Pvalb-Flp mice*, 5 *Rbp4-Cre X Vgat-Venus mice*, 8 *Rbp4-Cre* mice;
SP_{NSFint}: n = 13 cells, 11 *Rbp4-Cre* mice, 1 *Rbp4-Cre X Vgat-Venus* mouse;
SR: n = 14 cells, 6 mice; SL: n = 60 cells, 15 mice; SM: n = 21 cells, 10 mice

e, f, Effects of bath application of Gabazine (orange) and NBQX (green) on PSPs recorded from a fast-spiking pyramidal layer interneuron (left) (Scale bar: 1 mV, 5 ms)**.** Summary quantification of PSP amplitudes for multiple cells (right). The PSP amplitudes were largely unaffected by application of Gabazine (SP_{int} : n = 6 cells, p = 0.99 2-tailed Student's t test) but were largely blocked by NBQX (SP_{int} : n = 5 cells, p = 0.03, 2-tailed Student's t test) indicating AMPA receptor-mediated glutamatergic synaptic transmission.

g, h, Effects of bath application of TTX (orange) and 4-AP (green) on the response amplitude PSPs recorded from pyramidal layer interneurons. Example traces from a fast-spiking pyramidal layer interneuron (Scale bar: 1 mV, 5 ms). Summary plots of response amplitude measurements from multiple recordings (SP_{int}: n = 4 cells).

i, j, An example of ten consecutive PSP responses recorded from a single fast-spiking interneuron in SP illustrates the short and invariant latency of PSPs. Cumulative probability plots of standard deviation of latencies for neurons with PSP responses that were > 1 mV in amplitude (mean latency SP: 2.08 ± 0.08 ms, n = 21 cells).

Discussion

Together our results suggest a substantial revision to the idea that the entorhinal deep layers

unidirectionally convey hippocampal output messages to the neocortex. Using two lines of

evidence, combinatorial viral labelling and MAPseq, we show that L5a axons bifurcate to

target both the telencephalon and the hippocampus. Our data shows that nearly all

telencephalic output is copied back to the hippocampus regardless of its destination in the

telencephalon following distinct topographical organization rules.

Although cells in deep entorhinal layers have been previously observed when retrograde

dyes were placed at the CA1-subiculum border (Köhler, 1985; Witter and Amaral, 1991) the

tools we have introduced in this study made it possible to investigate the distinct source of

these projections and their targets within CA1. Our observation that L5b does not participate

in this pathway reinforces the complex input-output organisation of the deep MEC. L5b is an

integrator of local and neocortical input with hippocampal output and distributes its signals

locally within MEC. It provides an indirect feedback message to the hippocampus via its

projections to L3 of the MEC. L5a on the other hand appears not to be a focus of local

computations. Instead, it is suited to coordinate the wider hippocampal-neocortical loop by

providing a rapid copy of entorhinal output onto CA1.

In understanding the respective roles of projections from the superficial versus deep EC to

CA1, elucidating the topographical organization of the two projection pathways is crucial.

Our results show that in the transverse axis of CA1, the density of projections originating

from L5a of the MEC has a gradient descending towards proximal CA1 (towards CA2). This

is in contrast to previous reports where traditional (non-cell type specific) tracers targeted to

MEC revealed dense projections in proximal but not in distal CA1 (Tamamaki and Nojyo,

1995; Naber et al., 2001). The topographical organization of projections originating

specifically in L5a of the MEC that we reveal here suggests that this pathway serves a

function that is distinct from the ones that originate in the superficial layers of MEC instead of

merely supplementing them (Witter et al., 2017). It also opens up a possibility for the lateral

and medial entorhinal cortex input streams to converge in CA1.

Our high-throughput sequencing based projection mapping approach allowed us to also

reveal the topography of the deep layer projections to the hippocampus in the dorso-ventral

axis. The dorso-ventral target zone in the hippocampus of axons originating from L5a of the

MEC is in register with the dorso-ventral position of the originator cells in L5a. This is similar

to the topography of perforant-path projections of the superficial EC targeting the dentate

gyrus supporting the view that dorso-ventral compartments of the hippocampus might

operate independently (Ruth et al., 1982; Dolorfo and Amaral 1998).

Our findings raise new questions about the functional roles of outputs from the entorhinal

cortex, and about the nature of interactions between the entorhinal cortex and the

hippocampus. Telencephalic projections of L5a cells have been shown to be necessary for

long-term memory formation and retrieval (Kitamura et al., 2017) likely by distributing

compressed hippocampal outputs to the neocortex (McClelland et al., 1995). Returning a

copy of these outputs back to CA1 might create a self-reinforcing or a self-correcting loop.

They might also provide a gating signal to modulate the impact in CA1 of inputs from CA3

(Dudman et al., 2007) and/or neocortical return inputs via L3 of EC. It is also conceivable

that they influence mechanisms of learning and memory by affecting rhythmic network

activity through the inputs on the pyramidal cells and interneuron populations that we

described here (Colgin 2016; Ognjanovski et al. 2017; Xia et al. 2017). In addition, L5a might

also relay cortical input directly to CA1 however sources of cortical afferents to entorhinal

L5a remain to be established.

Methods

Mice

All animal experiments were approved by the University of Edinburgh animal welfare committee and were performed under a UK Home Office project license. The *Rbp4-Cre* mouse line was generated by GenSat and obtained from MMRRC (*Tg(Rbp4-cre)KL100Gsat/Mmucd*). The *Vgat-Venus* mouse line was generated by Dr. Atsushi Miyawaki at RIKEN, Wako, Japan (Wang et al., 2009). The *Pvalb-Flp* mouse line was obtained from Jackson Laboratories *(B6.Cg-Pvalb-tm4.1(FlPo)Hze/J; Jax 022730)*. Wild-type mice were obtained from Charles River Laboratories C57Bl6J stock. Double transgenics were generated by crossing the *Rbp4-Cre* line to the *Vgat-Venus* or the *Pvalb-Flp* lines. *Rbp4-Cre* and *Pvalb-Flp* lines were maintained as heterozygous and all mice were on C57Bl6 background.

Viral constructs and injection strategy

8 to 14 week-old male and female mice were used in all experiments. For targeting deep MEC a craniotomy was made 3.4 to 3.65 mm lateral to the bregma (X) between the transverse sinus and lambdoid suture (Y = 4.5 to 5 mm caudal to bregma). The injection pipette was at a 9 degree angle caudal to the dorsoventral plane (See Fig. 1b). 100 nl of virus was slowly released at four Z-depths 3.0, 2.8, 2.6 and 2.4 mm from the surface of the brain. 3 mins after delivering the virus the injection needle was retracted to the next injection depth until the most dorsal location where 10 mins past before the needle was fully retracted. For anterograde tracing of L5a axons and terminals *AAV-EF1a-DIO-mCherry* (serotypes 2 and 5, titer 5.3 x 10^{12}, lot #AV4375H, K., Deisseroth, UNC Vector Core), *AAV-CAG-FLEX-GFP* (titer 3.7 x 10^{12}, lot #AV4530B, E. Boyden, UNC Vector Core) and AAV-hSyn-FLEx-mGFP-2A-Synaptophysin-mRuby (Addgene:71760) were used.

For injections in CA1 a craniotomy was made 3.5 mm lateral and 3.30 mm caudal to the

bregma in C57BL6J mice. 100nl virus was delivered at 3.0, 2.8, 2.6, 2.4 and 2.2 mm from

the surface. For retrograde labelling of hippocampus-projecting EC neurons, 100nl of a

rAAV2retro-CAG-GFP (titre 4.3 x 10^{12}, lot #AV6951B, E. Boyden, UNC Vector Core) was

injected at each depth.

To assess whether telencephalic projection neurons in L5a of MEC co-express Etv1

(Extended Data Fig. 1b), fluorophore coupled choleratoxin beta subunit (CTB-Alexa 488 and

CTB-Alexa 555) was injected at various sites in wild-type C57BL6J mice as previously

described(Sürmeli et al., 2015).

To achieve Cre expression in target specific subpopulation of L5a neurons in the MEC (Fig.

2) a *rAAVretro-EF1a-Cre-WPRE* (viral construct was a gift from Karl Deisseroth, titre 2.4 x

10^{12}) was made using an AAV-retro helper plasmid (Addgene plasmid ID 81070) as

described previously(McClure et al., 2011). A cocktail of *rAAVretro-EF1a-Cre-WPRE* and

AAV-CAG-GFP (titer 4.3 x 10^{12}, lot #AV6951B, Boyden, UNC Vector Core) viruses were

injected either in NucAcb (X: +1.0 mm; Y: +1.2 mm; Z: -3.8, -4.0 mm) or RSC (X: +0.4 mm;

Y: -2.8, -3.2 mm; Z: -0.9 mm) in wild-type mice. A Cre inducible reporter virus was also

injected in the MEC as described above. GFP expression driven by the *CAG* promoter at the

target site was used for verification of injection location.

To achieve expression of channelrhodopsin-2 in L5a axons, we injected *AAV2-EF1a-DIO-*

hChR2(H134R)-mCherry-WPRE-pA (titer 5.1 x 10^{12}, lot #AV4319J, Deisseroth) in the MEC

of the *Rbp4-Cre* or the *Rbp4-Cre X Vgat-Venus* or the *Rbp4-Cre X Pvalb-FLP* double

transgenic mice. Additionally, in order to fluorescently label parvalbumin expressing

interneurons in CA1 a Flp recombinase-dependent AAV vector was constructed in house

and injected in the hippocampus (X: +3.0 mm; Y: -3.5 mm; Z: -2.0, -3.0 mm, 200 nl virus was

injected at each Z-depth). To produce the virus, the viral construct pAM-FLEX-GFP(Murray

et al., 2011) (a gift from Peer Wulff) was used as the viral backbone. The eGFP, loxP and

lox2272 sites were removed and replaced with a synthesised cassette (GenScript)

containing a multiple cloning site flanked by two sets of heterotypic (FRT and F3),

antiparallel FRT sites to produce pAM FLPX. For pAM FLPX eGFP and pAM FLPX mCherry,

the fluorescent proteins were amplified from pAM FLEX eGFP and pmCherry-C1 (Clontech)

respectively, with EcoRI and SacI overhangs and cloned into pAM FLPX. All AAV preps

were generated in house as previously described(McClure et al., 2011) and were titered by

qPCR following expression in HEK cells (> 10^{12} genome copies (GC)/ml).

327 Tissue processing and immunohistochemistry

3 to 4 weeks after the virus injection surgery, mice were transcardially perfused with 4%

paraformaldehyde as previously described(Sürmeli et al., 2015). Brains were sectioned at 60

µm using a cryostat (Leica CM3050 S) either coronally or horizontally and sections were

stored in PBS at 4°C. All subsequent incubation and washing steps were carried out at 4°C.

Prior to antibody staining, sections were blocked with either 5% normal goat serum (NGS;

Abcam: ab13970) or 2% bovine serum albumin (BSA; Sigma: A9418) in 0.3% PBST (PBS

with 0.3% Triton-X100) for 2hrs. Sections were then transferred to primary antibody solutions

made with either 5% NGS or 2% BSA in 0.3% PBST and incubated overnight. After 3

washes, each 20 min, in 0.3% PBST, sections were transferred to secondary antibody

solutions made with 0.3% PBST and, if required, NeuroTrace 640/660 (1:800; Life

Technologies: N21483) and incubated overnight. After 3 washes, sections were incubated in

DAPI (1:2,000; Sigma Aldrich: D9542) solution made in PBS for 20 min at room temperature,

where required, and then mounted on microscope slides using Mowiol® 4-88 (Aldrich:

81381). Slides were covered with glass coverslips and left to dry overnight at 4°C in the dark.

The following primary antibodies were used: rat anti-mCherry (1:2,000; ThermoFisher:

M11217), chicken anti-GFP (1:10,000; Abcam: ab13970), rabbit anti-Etv1(Arber et al., 2000)

(1:1,000), chicken anti-NeuN (1:1,000; Sigma Aldrich: ABN91). The following secondary

antibodies from Thermo Fisher Scientific were used at a concentration of 1:800: goat anti-

rabbit A488 (A11034), goat anti-chicken A488 (A11039), goat anti-rat A555 (A21434) and

goat anti-rabbit (A21244).

Cell counting

Every 4th brain section was imaged as a Z-stack (1 µm step size) using either the Zeiss

LSM 800 (Zen, v2.6.76) or Leica SP8 confocal microscope (Leica Application Suite X,

v3.5.6.21594) at 20x magnification. Regions of interest were drawn around L5a and cell

counting was carried out either manually using Fiji software, or using cell-counting tools in

Vision4D (Arivis, v3.2.0).

The boundaries for L5a were determined using either a DAPI or Neurotrace counterstain.

The border between medial and lateral divisions of EC was determined in each section using

layer 2 as a guide – in lateral EC, layer 2 is separated into two distinct layers, while this

separation is not seen in medial EC(Cappaert et al., 2015).

Quantification of the distribution of fluorescent signal

Quantification of fluorescence across the radial axis of the CA1

The distribution of axons of L5a neurons in the radial axis of the hippocampal CA1 was

quantified in slide-scanner images (Fig. 1) or confocal images (Fig. 2) using Fiji software.

Coronal brain sections between 3.28 to 3.58 mm posterior to Bregma, which contain distal

CA1, were selected for analysis. 2 sections from each brain in which the fluorescent labeling

was representative of all the sections were used. DAPI was used as a counterstain. In each

section, a 400 µm-wide rectangular ROI was drawn across the radial axes, covering all

layers of a randomly selected region in intermediate CA1 (Fig. 1c). This ROI was then

divided into 20 equal bins and the mean fluorescence intensity was calculated for each bin.

The mean intensity values were then normalised in each brain by scaling the values such

that the highest value is 1.

Quantification of fluorescence in the transverse axis of the CA1

A Zeiss Axioscan slidescanner was used to image every second brain section at 10x

magnification. The projection strengths of MEC L5a neurons to proximal and distal halves of

CA1 were quantified in horizontal sections located at depths between 2.56 mm to 4.12 mm

from the surface of the brain, with injection sites located across medial, mid and lateral MEC

(15-18 brain sections per mouse, n = 5 mice). GFP and mCherry fluorescence signals were

amplified with immunostaining as described previously. The borders of CA1 were drawn on

brain section images using a customisable digital microscopy analysis platform (Visiopharm),

and the proximo-distal border was defined as the border equidistant from the proximal and

distal ends of CA1. The mean fluorescence density, defined as the total number of pixels

above a set threshold in an area divided by the total number of pixels in the area, was

measured in proximal and distal CA1. The threshold was determined manually by ensuring

that only pixels representing axons were detected as signal. A median unsharp background

correction was used to remove background noise from axons outside of the focus plane of

the image. The mean fluorescence density values were then normalised within each brain by

scaling the values such that the total fluorescence density value (proximal + distal) in each

brain is equal to 1.

The spread of viral expression in MEC was assessed from the fluorescent signal in all

sections for each brain. Brains in which labelled neurons were found outside L5 were

excluded; this was a result of occasional and sparse labelling in L2 or L3 or the

parasubiculum in the *Rbp4-Cre* line.

MAPseq

5 C57Bl6J adult male and female mice were injected in the deep MEC with the MAPseq

Sindbis viral barcode library (3 x 10^{10} GC/ml) provided by the MAPseq facility (Cold Spring

Harbor Laboratories). To cover the whole mediolateral and dorsoventral extent of the MEC

virus was injected in two locations: 3.4 mm and 3.6 mm lateral to bregma. 100 nl virus was

delivered to 3.0, 2.8, 2.6, 2.4 mm below the surface of the brain. After 44 hrs, mice were

sacrificed, brains were extracted and immediately immersed in oxygenated cold ACSF

prepared in ultrapure water. All surgical tools and surfaces were treated with RNaseZAP

(Invitrogen) prior to the start of the experiment and in-between samples. 400 µm fresh

sagittal brain sections were cut using a vibratome(Pastoll et al., 2012). Immediately after

this, GFP expression in the deep EC was verified under a fluorescent microscope and tissue

was dissected in cold oxygenated ACSF using microdissection blades on sylgard covered

petri dishes which were kept on ice. Each slice was dissected on a previously unused

surface of the plate with fresh ACSF and a dedicated microblade was used for the dissection

of each brain division to prevent contamination. Tissue pieces were collected into bead

containing (Qiagen 69989) collection tubes (Qiagen 19560) on dry ice. After tissue

collection, 400 µl Trizol (Thermo Fisher Scientific #15596026) was added to each collection

tube. The procedure was repeated for all 5 brains. 3 brains with the largest coverage of MEC

deep layers were selected to proceed to the next steps. Tissue was stored at -80 degrees

before it was shipped on dry ice to CSHL MAPseq core facility for further processing. RNA

extraction, production of double stranded cDNA, library preparation and Illumina sequencing

and preprocessing of sequencing data was performed by the MAPseq core facility as

described in(Kebschull et al., 2016).

For the tissue dissections, identification of brain areas was done by using Allen Brain

Reference Atlas (https://mouse.brain-map.org/). The isocortex division included the

somatosensory, motor, visual, posterior parietal, anterior cingulate and retrosplenial areas

combined. The cerebral nuclei (CNU) division was restricted to striatum. The

olfactory/cortical subplate division (Olf/Ctxsp) was a combination of olfactory areas and

cortical subplate including amygdalar nuclei. The remaining two divisions were dorsal (dHip)

and ventral hippocampus (vHip) including subiculum. Some brain areas were excluded from

the study because of the difficulty in dissecting or identifying brain areas in the sagittal plane.

All sections > 3.7 mm lateral to bregma which are not annotated in Allen Brain Reference

Atlas were excluded. Therefore, neocortical areas in the most lateral sections such as

perirhinal cortex, ectorhinal cortex, temporal association areas were not included in the

study. The claustrum and adjacent neocortical areas (visceral, agranular insular, gustatory)

were excluded as it was not possible to separate these areas precisely to prevent

contamination between the assigned divisions. Since borders between the brain divisions

CNU and OLF/Ctxsp were not always clear, dissections avoided these areas hence the brain

areas in these divisions are partially included. White matter between the hippocampus and

the neocortex carrying axon tracts were also excluded. Brainstem and cervical spinal cord

tissue were used as control. When L5a neurons in the MEC were labelled using strategies

explained in Fig. 1, no axonal projections to these areas were observed (unpublished

experiments). Consistent with this, a low number of barcodes were identified in the negative

control samples (on average 1.0% of barcodes had counts in control areas, n = 3).

A limitation of the barcode-based single-cell tracing method comes from the possibility of

multiple neurons being represented by the same barcode. This is prevented by adjusting the

barcode diversity and the size of the target population as explained in detail in(Kebschull et

al., 2016). In order to assess the expected fraction of uniquely labelled neurons in our study

we counted the total number of neurons in L5a of the MEC. A total of 2314 ± 131, (n = 2

mice) NeuN-labelled neurons were counted. Using the formula $F = (1-(1/N))^{k-1}$ where N is

the barcode diversity (2×10^6)(Kebschull et al., 2016) and k is the number of infected

neurons the predicted ratio of uniquely labelled neurons is 99.9%. Multiple representation of

neurons due to a neuron being infected by several different barcode RNA-expressing viral

particles was not corrected for since the projection patterns are not affected by

overrepresentation of neurons(Kebschull et al., 2016).

MAPseq data analysis

Barcode counts were first normalised in each area by the relative number of spike-in RNAs

for each sample. Orphan barcodes, barcodes which did not have counts in the injection site,

(dMEC or vMEC) were removed. We then calculated the 95th percentile of the barcode

counts in our negative controls and based on this set all barcode counts of 1 to 0. A small

number of barcodes had a higher count in any target area compared to the injection site

which might be a result of incomplete dissection of the injection site or viral expression in

areas that the virus spilled into as we retracted the pipette. These barcodes were removed.

Since our goal was to find whether neurons projecting to the telencephalon also project to

the hippocampus we removed the barcodes that had no counts in any of the telencephalic

target areas. For the same reason, barcodes that were detected only in the dHip or vHip

hippocampus were also removed. Finally, we excluded all barcodes with counts of less than

400 in the injection site to minimise the possibility of incomplete transport of RNA barcodes

to axons in target areas due to weak expression at the cell bodies or low counts due to PCR

or polymerase errors. Barcodes with low counts in all target areas (< 10) were also excluded

to account for potential false positives.

To quantify the proportion of barcodes that were present in both a target division (Fig. 2h;

Isocortex, CNU or Olf/Ctxsp) and the hippocampus, the following formula was used: (total

number of barcodes with counts detected in both hippocampus and target division / total

number of barcodes with counts detected in target division). To compare the difference

between barcodes with the injection site in dMEC vs vMEC, the barcodes were divided into

two groups – "dorsal MEC", where counts in dMEC were higher than in vMEC, and vice

versa for "ventral MEC". A paired two-tailed Wilcoxon signed rank test was used to check if

there was a significant difference between the proportion of collateralising neurons in dMEC

and vMEC.

To visualise the distribution of barcode counts in dHip and vHip from "dorsal MEC" or

"ventral MEC" groups, we generated heat maps showing normalised projection strengths

between dHip and vHip for all barcodes. The normalised projection strengths were

calculated by the formula: projection strength$_{dHip}$ = (counts$_{dHip}$) / (counts$_{dHip+vHip}$) * 100, and

likewise for vHip. Barcodes were sorted by maximum projection site.

Ex-vivo electrophysiology

Slice preparation

Slice preparation and subsequent data acquisition was done as previously

described(Sürmeli et al., 2015). 3 to 4 weeks after the injection of the viral vectors ex-vivo

brain slices were prepared. Following decapitation, brains were immersed for 2 mins in 4°C

artificial cerebrospinal fluid (ACSF) of the following composition (mM): 86 NaCl, 1.2

NaH_2PO_4, 2.5 KCl, 25 $NaHCO_3$, 25 Glucose, 75 Sucrose, 0.5 $CaCl_2$, 7 $MgCl_2$, bubbled with

95% O_2 / 5% CO_2. They were then sectioned horizontally (400 μm) using a vibratome (Leica

VT1200) ACSF. Tissue was collected and maintained in extracellular solution of the

following composition (mM): 124 NaCl, 1.2 NaH_2PO_4, 2.5 KCl, 25 $NaHCO_3$, 20 Glucose, 2

$CaCl_2$ and 1 $MgCl_2$, continuously supplied with 95% O_2 / 5% CO_2. Slices were allowed to rest

for 15 mins at 35°C followed by a minimum of 30 mins recovery time at room temperature

before the start of the experiment.

Electrophysiological recordings

Whole-cell patch-clamp recordings were made in pyramidal neurons and interneurons in all

layers of hippocampal CA1. Typically, 2 to 3 slices from the intermediate hippocampus were

used where morphologies of the pyramidal cells and interneurons were confirmed to be

intact with post hoc staining and imaging of biocytin filled neurons. Data was collected using

AxoGraph (v1.7.6) software.

Pipettes with 4 to 6 MΩ resistance were pulled from borosilicate glass (Sutter Instruments)

and filled with an intracellular solution of following composition (mM): 130 K Gluconate, 10

KCl, 10 HEPES, 2 $MgCl_2$, 0.1 EGTA, 2 Na_2ATP, 0.3 Na_2GTP, 10 NaPhosphocreatine, 5.4

Biocytin. The intracellular solution was adjusted to a pH of 7.2 to 7.4 with KOH and

osmolarity 290-300 mOsm. All recordings were made in current clamp mode with pipette

capacitance neutralisation and bridge balance applied. Subthreshold membrane properties

were measured from the changes in membrane potential upon depolarising and

hyperpolarising current injections (typically -40 pA to +40 pA, in 20 pA). Rheobase was

established from responses to a depolarizing current ramp (50 pA/sec, maximum current

400 pA).

For optogenetic stimulation of ChR2, an LED of wavelength 470 nm (ThorLabs) was

attached to the epifluorescence port of the microscope. Where necessary, the irradiance of

the LED (max 9 mW) was controlled by voltage commands. Pharmacological tests were

done by bath application of the following reagents with the indicated final concentrations in

standard extracellular solution: Gabazine (10 µM, Hello Bio, Cat. No. HB0901), NBQX (10

µM, Tocris, Cat. No. 0373), D-AP5 (50 µM, Tocris, Cat. No. 0106), 4-AP (200 µM, Tocris,

Cat. No. 0940), TTX (1 µM, Hello Bio). To allow for morphological reconstructions cells were

filled with Biocytin (5.4 mM in intracellular solution, Sigma, Cat. No. B4261) during

electrophysiological recordings.

Analysis of electrophysiological recordings

Electrophysiological properties were analysed using built-in and custom routines in

IGORpro8 (WaveMetrics) and Matlab (MathWorks). All basic properties were established

from the I-V protocol described above. Input resistance was determined from the largest

depolarising current step injected. Sag ratio was calculated as $V_{steady\ state}/ V_{min}$ from the largest

hyperpolarising current step. Rheobase and action potential (AP) threshold were determined

from the first AP during injection of steadily increasing current in a ramp. From the same

protocol, after-hyperpolarisation (AHP) was calculated as the difference between AP

threshold and the most negative peak of the hyperpolarisation following the first spike. Half-

width was measured as the width of the AP at half its maximum spike amplitude. Firing

frequency was measured from a 1 s current injection of 200 pA. Maximum firing frequency

was measured between the first two APs, and base frequency between the last two APs in a

train.

For optogenetic stimulations, responsiveness was confirmed for each cell by a significant

difference between the detected peak of change in membrane potential after light stimulation

and the average baseline using a 2-tailed, type 1 student's t-test. The PSP amplitude and

latency were measured using the Neuromatic toolbox in IGORPro8.

Cell types were determined based on biophysical properties extracted from

electrophysiology data. Pyramidal cells were identified from a max firing frequency of < 50

Hz, AHP < 10 mV, AP half-width ≥ 0.7 ms and sag ≤ 0.9, with descending hierarchical order.

Fast spiking interneurons were categorised from a max firing frequency ≥ 100 Hz, AHP ≥ 12

mV, AP half-width < 0.5 ms and sag > 0.9. Non-fast spiking interneurons were classified

from a max firing frequency 50-100 Hz, AHP ≥ 12 mV, AP half-width 0.5 to 0.9 ms, sag ratio

> 0.9. and max firing frequency 50-100 Hz.

Neurons with a resting membrane potential less negative than -50 mV or a bridge balance

higher than 40 MΩ were excluded from the analysis, as were neurons that did not fit within a

class from above criteria.

Immunostaining of electrophysiology slices

For identification of cells following recordings, tissue was fixed in 4% PFA overnight at 4∘C.

Slices were washed with PBS three times, 20 min each, and incubated in Streptavidin-Alexa

488 (S11223) or Alexa 647 (1:500, S32357, Invitrogen) and DAPI (1:1000, in 0.3% PBST)

overnight at 4°C. Slices were washed in 0.3% PBST four times and mounted on glass slides

with Mowiol® 4-88. Mcherry fluorescence in the MEC was examined with confocal imaging

and those animals where virus injection missed MEC L5 were excluded.

For staining of the interneuron marker parvalbumin (PV), slices were prepared as described

above, then incubated in primary antibody solution containing mouse anti-PV (1:1000, PV

235, Swant) and 5% NGS in 0.3% PBST for 48 hrs at 4°C. Slices were then washed and

incubated in secondary antibody solution with Alexa-conjugated Streptavidin and DAPI

prepared in 0.3% PBST and mounted on glass slides after overnight incubation.

A Zeiss LSM800 microscope was used for image acquisition of the slices. Pinhole was set to

1 Airy Unit. Objectives used include x10 (air), x20 (air) and x40 (oil) to image the

hippocampal formation, morphology of biocytin filled cells, and immunolabelling of

interneurons, respectively.

Statistics

Statistical tests (Student's t-test, Chi-squared test, Wilcoxon signed rank test) were

performed using R (www.r-project.org) 3.6.0, 4.0.0 and 4.0.3. Normality was tested for using

both the Shapiro-Wilk test and QQ plots. All data is presented in the format of mean ± SEM

unless otherwise stated. A p-value < 0.05 was regarded as significant.

Data availability

On publication data and analysis scripts will be made publicly available via University of

Edinburgh's Datashare service (http://datashare.is.ed.ac.uk/). This is an online data

repository maintained by the University. MAPseq data will be made available at NLM

Sequence Read Archive BioProject.

Code availability

Analysis script will be made publicly available upon publication via University of Edinburgh's
Datashare service

Acknowledgements

We thank Matt Nolan and Nolan Lab for discussions and sharing reagents and equipment.
We thank Justus Kebschull for sharing his insight in MAPseq technique, Maria Doitsidou for
sharing her equipment and lab space. We also thank Matt Nolan, Klara Gerlei, Brianna
Vandrey, Christina Brown and our lab members for critical reading of the manuscript and
Gamze Şener for her help with early experiments. C.M. was supported by a BBSRC project
grant BB/M025454/1. This research was supported by a Royal Society and Wellcome Trust
Sir Henry Dale fellowship 211236/Z/18/Z.

Contributions

S.Y.T. and M.Ö. equally contributed to the study.
G.S. conceptualised the study, acquired funding and administered the project. G.S. and
S.Y.T. performed MAPseq experiments and analysed the data. M.Ö., G.S. and E.S.
performed electrophysiology experiments and data analysis. Anatomy and histology
experiments were performed and analysed by S.Y.T., G.S., M.Ö., E.S., and M.R.. C.M.
developed reagents. Z.B. developed analysis tools and contributed to early discussions.
G.S., S.Y.T., E.S. and M.Ö. prepared the original manuscript.

References

Arber, S., Ladle, D.R., Lin, J.H., Frank, E., and Jessell, T.M. (2000). ETS gene Er81 controls the formation of functional connections between group Ia sensory afferents and motor neurons. Cell *101*, 485–498.

Beed, P., de Filippo, R., Holman, C., Johenning, F.W., Leibold, C., Caputi, A., Monyer, H., and Schmitz, D. (2020). Layer 3 Pyramidal Cells in the Medial Entorhinal Cortex Orchestrate Up-Down States and Entrain the Deep Layers Differentially. Cell Rep. *33*, 108470.

Cappaert, N.L.M., Van Strien, N.M., and Witter, M.P. (2015). Chapter 20 - Hippocampal Formation. In The Rat Nervous System (Fourth Edition), G. Paxinos, ed. (San Diego: Academic Press), pp. 511–573.

Colgin, L.L. (2016). Rhythms of the hippocampal network. Nature Reviews Neuroscience *17*, 239–249.

Czajkowski, R., Sugar, J., Zhang, S.-J., Couey, J.J., Ye, J., and Witter, M.P. (2013). Superficially projecting principal neurons in layer V of medial entorhinal cortex in the rat receive excitatory retrosplenial input. J. Neurosci. *33*, 15779–15792.

Dolorfo, C. L., and D. G. Amaral. 1998. Entorhinal Cortex of the Rat: Topographic Organization of the Cells of Origin of the Perforant Path Projection to the Dentate Gyrus. The Journal of Comparative Neurology 398 (1): 25–48.

Dudman, J.T., Tsay, D., and Siegelbaum, S.A. (2007). A role for synaptic inputs at distal dendrites: instructive signals for hippocampal long-term plasticity. Neuron *56*, 866–879.

Frankland, P. W. & Bontempi, B. The organization of recent and remote memories. *Nat. Rev. Neurosci.* **6**, 119–130 (2005).

Gerfen, C.R., Paletzki, R., and Heintz, N. (2013). GENSAT BAC cre-recombinase driver lines to study the functional organization of cerebral cortical and basal ganglia circuits. Neuron *80*, 1368–1383.

Kebschull, J.M., da Silva, P.G., Reid, A.P., Peikon, I.D., Albeanu, D.F., and Zador, A.M. (2016). High-Throughput Mapping of Single-Neuron Projections by Sequencing of Barcoded RNA. Neuron *91*, 975–987.

Khodagholy, D., Gelinas, J.N., and Buzsáki, G. (2017). Learning-enhanced coupling between ripple oscillations in association cortices and hippocampus. Science *358*, 369–372.

Kitamura, T., Pignatelli, M., Suh, J., Kohara, K., Yoshiki, A., Abe, K., and Tonegawa, S. (2014). Island Cells Control Temporal Association Memory. Science *343*, 896–901.

Kitamura, T., Ogawa, S.K., Roy, D.S., Okuyama, T., Morrissey, M.D., Smith, L.M., Redondo, R.L., and Tonegawa, S. (2017). Engrams and circuits crucial for systems consolidation of a memory. Science *356*, 73–78.

Kloosterman, F., Van Haeften, T., Witter, M.P., and Lopes Da Silva, F.H. (2003). Electrophysiological characterization of interlaminar entorhinal connections: an essential link for re-entrance in the hippocampal-entorhinal system. Eur. J. Neurosci. *18*, 3037–3052.

Köhler, C. (1985). A projection from the deep layers of the entorhinal area to the
hippocampal formation in the rat brain. Neurosci. Lett. *56*, 13–19.

Koster, R., Chadwick, M.J., Chen, Y., Berron, D., Banino, A., Düzel, E., Hassabis, D., and
Kumaran, D. (2018). Big-Loop Recurrence within the Hippocampal System Supports
Integration of Information across Episodes. Neuron *99*, 1342–1354.e6.

Kumaran, D., Hassabis, D., and McClelland, J.L. (2016). What Learning Systems do
Intelligent Agents Need? Complementary Learning Systems Theory Updated. Trends Cogn.
Sci. *20*, 512–534.

Kumaran, D., and McClelland, J.L. (2012). Generalization through the recurrent interaction of
episodic memories: a model of the hippocampal system. Psychol. Rev. *119*, 573–616.
647
Marr, D. (1971). Simple memory: a theory for archicortex. Philos. Trans. R. Soc. Lond. B
Biol. Sci. *262*, 23–81.

McClelland, J.L., McNaughton, B.L., and O'Reilly, R.C. (1995). Why there are
complementary learning systems in the hippocampus and neocortex: insights from the
successes and failures of connectionist models of learning and memory. Psychol. Rev. *102*,
419.

McClure, C., Cole, K.L.H., Wulff, P., Klugmann, M., and Murray, A.J. (2011). Production and
titering of recombinant adeno-associated viral vectors. J. Vis. Exp. e3348.

Murray, A.J., Sauer, J.-F., Riedel, G., McClure, C., Ansel, L., Cheyne, L., Bartos, M.,
Wisden, W., and Wulff, P. (2011). Parvalbumin-positive CA1 interneurons are required for
spatial working but not for reference memory. Nat. Neurosci. *14*, 297–299.

Naber, P.A., Lopes da Silva, F.H., and Witter, M.P. (2001). Reciprocal connections between
the entorhinal cortex and hippocampal fields CA1 and the subiculum are in register with the
projections from CA1 to the subiculum. Hippocampus *11*, 99–104.

Ognjanovski, N., Schaeffer, S., Wu, J., Mofakham, S., Maruyama, D., Zochowski, M., and
Aton, S.J. (2017). Parvalbumin-expressing interneurons coordinate hippocampal network
dynamics required for memory consolidation. Nat. Commun. *8*, 15039.

Ohara, S., Onodera, M., Simonsen, Ø.W., Yoshino, R., Hioki, H., Iijima, T., Tsutsui, K.-I., and
Witter, M.P. (2018). Intrinsic Projections of Layer Vb Neurons to Layers Va, III, and II in the
Lateral and Medial Entorhinal Cortex of the Rat. Cell Reports *24*, 107–116.
668
669   Ólafsdóttir, H.F., Carpenter, F., and Barry, C. (2016). Coordinated grid and place cell replay
670   during rest. Nat. Neurosci. *19*, 792–794.
671
Pastoll, H., White, M., and Nolan, M. (2012). Preparation of parasagittal slices for the
investigation of dorsal-ventral organization of the rodent medial entorhinal cortex. J. Vis.
Exp.

Petreanu, L., Huber, D., Sobczyk, A., and Svoboda, K. (2007). Channelrhodopsin-2–assisted
circuit mapping of long-range callosal projections. Nat. Neurosci. *10*, 663–668.

Rozov, A., Rannap, M., Lorenz, F., Nasretdinov, A., Draguhn, A., and Egorov, A.V. (2020).
Processing of Hippocampal Network Activity in the Receiver Network of the Medial
Entorhinal Cortex Layer V. The Journal of Neuroscience *40*, 8413–8425.

Ruth, R. E., Collier, T. J. & Routtenberg, A. Topography between the entorhinal cortex and
the dentate septotemporal axis in rats: I. Medial and intermediate entorhinal projecting cells.
J. Comp. Neurol. **209**, 69–78 (1982).

Schapiro, A.C., Turk-Browne, N.B., Botvinick, M.M., and Norman, K.A. (2017).
Complementary learning systems within the hippocampus: a neural network modelling
approach to reconciling episodic memory with statistical learning. Philos. Trans. R. Soc.
Lond. B Biol. Sci. *372*.

Squire, L. R., Genzel, L., Wixted, J. T. & Morris, R. G. Memory consolidation. Cold Spring
Harb. Perspect. Biol. **7**, a021766 (2015).

Sürmeli, G., Marcu, D.C., McClure, C., Garden, D.L.F., Pastoll, H., and Nolan, M.F. (2015).
Molecularly Defined Circuitry Reveals Input-Output Segregation in Deep Layers of the
Medial Entorhinal Cortex. Neuron *88*, 1040–1053.

Swanson, L.W., and Köhler, C. (1986). Anatomical evidence for direct projections from the
entorhinal area to the entire cortical mantle in the rat. J. Neurosci. *6*, 3010–3023.

Tamamaki, N., and Nojyo, Y. (1995). Preservation of topography in the connections between
the subiculum, field CA1, and the entorhinal cortex in rats. J. Comp. Neurol. *353*, 379–390.

Wang, Y., Kakizaki, T., Sakagami, H., Saito, K., Ebihara, S., Kato, M., Hirabayashi, M.,
Saito, Y., Furuya, N., and Yanagawa, Y. (2009). Fluorescent labeling of both GABAergic and
glycinergic neurons in vesicular GABA transporter (VGAT)–Venus transgenic mouse.
Neuroscience *164*, 1031–1043.

Witter, M.P., and Amaral, D.G. (1991). Entorhinal cortex of the monkey: V. Projections to the
dentate gyrus, hippocampus, and subicular complex. J. Comp. Neurol. *307*, 437–459.

Witter, M.P., Doan, T.P., Jacobsen, B., Nilssen, E.S., and Ohara, S. (2017). Architecture of
the Entorhinal Cortex A Review of Entorhinal Anatomy in Rodents with Some Comparative
Notes. Front. Syst. Neurosci. *11*, 46.

Wozny, C., Beed, P., Nitzan, N., Pössnecker, Y., Rost, B.R., and Schmitz, D. (2018).
VGLUT2 Functions as a Differential Marker for Hippocampal Output Neurons. Front. Cell.
Neurosci. *12*, 337.

Xia, F., Richards, B.A., Tran, M.M., Josselyn, S.A., Takehara-Nishiuchi, K., and Frankland,
P.W. (2017). Parvalbumin-positive interneurons mediate neocortical-hippocampal
interactions that are necessary for memory consolidation. Elife *6*.

717
718

Extended Data Figures

Extended Data Figure 1. Injection site images, quantification of co-localisation between retrograde markers and Etv1, synaptic terminal imaging in CA1.
a, Horizontal brain sections from a single brain show the dorso-ventral spread of virus-mediated reporter gene expression (Scale bar: 500 µm).
b, Quantification of overlap between retrograde labelling from cortical and subcortical areas and Etv1 expression in L5a of the MEC. The number of mice, mean proportion and total cells counted for each area are shown in the table.
c, Representative injection sites for brains used for quantifying the difference in axon density between proximal and distal CA1 show the medio-lateral extent of injections in the MEC
(Scale bar: 500 µm).
d, Horizontal view of hippocampal CA1 showing axon terminals of entorhinal L5a neurons (cyan).
e, A high contrast high magnification image showing synaptic puncta labelling in stratum pyramidale**.** (Scale bar: 10 µm).
Hip, hippocampus; Amg, amygdala; AOm, medial anterior olfactory area; Cg, cingulate area; NAc, nucleus accumbens; V1, primary visual cortex; V2, secondary visual areas; RSC, retrosplenial cortex.

Extended Data Figure 2. Images of injection sites.
a, b, Coronal brain sections showing labelled neurons in MEC with the combinatorial viral strategy (Scale bar: 500 μm). Inset: high-magnification confocal image of boxed region in left panel showing cell bodies in MEC expressing mCherry.
c, d, Coronal brain sections showing GFP expression at the injection sites in retrosplenial cortex (**c**) or nucleus accumbens (**d**) (Scale bar: 500 μm). Inset: high-magnification of boxed region in left panel.

RSC, retrosplenial cortex; NucAcb, nucleus accumbens.

Extended Data Figure 3. Verification of viral expression and further MAPseq data visualisation.
a, Brightfield and epifluorescence microscope images of the injection site on a sagittal brain section showing the spread of neurons infected by the MAPseq virus library.
b, A neuron's position in the dorso-ventral axis does not affect the likelihood of collateralisation of its axons to the hippocampus (Paired two-tailed Wilcoxon signed rank test, p = 0.75, median proportion of collateralizing barcodes in dEC = 98.4%, vEC = 98.9%, n = 3).

Extended Data Figure 4. Experimental design and properties of responses to optogenetic stimulation

a, Summary table for biophysical properties of neurons recorded in the study.

b, An example of a biphasic response recorded from a pyramidal neuron upon 3ms stimulation of axons of MEC L5a neurons in CA1. A late hyperpolarising (red) response was abolished by application of Gabazine to reveal a larger depolarising PSP (black) which was abolished by application of NBQX (grey) (Scale bar: 10 ms, 0.5 mV).

c, Experimental strategy for fluorescence guided patch-clamp recordings made from Parvalbumin expressing neurons. Left: A FlpX-dependent GFP expressing virus was injected in the hippocampus in *Rbp4-CreX Pvalb-Flp* mice. Right: Confocal image of horizontal slice showing labelled interneurons. Bottom: An example fast-spiking interneuron marked as described in **c**, that showed depolarising PSP upon light stimulation. During recording, the neuron was filled with biocytin and later stained with Streptavidin and Anti-Parvalbumin antibody.

d, (i) Traces of responses to a train of 10 light stimulation (blue lines), showing relatively stable EPSP amplitude across simulations. (Scale bar: 30 ms, 1 mV) (ii) Magnified images of 1st and 10th pulse illustrate the invariant latency in one example cell (Scale bars: 0.5 ms, 0.5 mV) (iii) Quantification of latency across the train in all cells. Error bars represent SEM.

e, Plots showing the frequency distribution of PSP amplitudes in pyramidal neurons and interneurons. PSPs over 12.5 mV were detected in interneurons and often resulted in spiking responses. (SP_Pyr : n = 19, SP_int : n = 19, SR : n = 9, SL&SM : n = 22 cells).

719 **f,** Effects of bath application of Gabazine and NBQX on PSPs recorded from interneurons in SR and SL. The
720 PSP amplitudes were largely unaffected by application of Gabazine (SR: n = 2 cells & SL n = 2 cells, p = 0.28,
      2-tailed Student's t test) but were largely blocked by NBQX (SR: n = 2 cells & SL n = 2 cells, p = 0.04, 2-tailed
721 Student's t test) indicating AMPA receptor mediated glutamatergic synaptic transmission.
      g, Effects of bath application of TTX and 4-AP on the response amplitude PSPs recorded from interneurons in
722 SR, SL and SM (SR: n = 2, SL: n = 3, SM: n = 1 cells).
      h, Cumulative probability plots of standard deviation of latencies for neurons with PSP responses that were > 1
723 mV in amplitude. EPSPs of interneurons in all layers except SO had a short latency after light stimulation
      (mean latency SR: 2.00 ± 0.08 ms, n = 6 cells; SL & SM: 2.12 ± 0.21 ms, n = 23 cells). In SO 2 out of 3 of the
724 evoked PSPs had a larger latency of onset (mean latency = 8.5 ± 0.63 ms, n = 2 cells, 2 mice, data not
      shown).

725
726

728
729

734
735

736