Single-Cell Spatial Proteomics Analyses of Head and Neck Squamous Cell Carcinoma Reveal Tumor Heterogeneity and Immune Architectures Associated with Clinical Outcome

Katie E. Blise[a], Shamilene Sivagnanam[b], Grace L. Banik[c,*], Lisa M. Coussens[d], and Jeremy Goecks[a,**]

[a]Department of Biomedical Engineering, Oregon Health & Science University, Portland, OR USA
[b]Program in Computational Biology, Department of Cell, Developmental & Cancer Biology, Oregon Health & Science University, Portland, OR USA
[c]Departments of Cell, Developmental & Cancer Biology, Otolaryngology-Head & Neck Surgery, Oregon Health & Science University, Portland, OR USA
[d]Department of Cell, Developmental & Cancer Biology, and Knight Cancer Institute, Oregon Health & Science University, Portland, OR USA
*Current affiliation: Division of Otolaryngology, Children's Hospital of Philadelphia, Philadelphia, PA USA
**Corresponding author

Correspondence:
Jeremy Goecks, Ph.D.
Associate Professor, Department of Biomedical Engineering
Oregon Health & Science University
3181 SW Sam Jackson Park Road, Mail Code: KR-COM, Portland, OR 97239
Tel: 503-494-0490, Email: goecksj@ohsu.edu

Running Title: Single-cell spatial analysis of head and neck cancer

Keywords: head and neck squamous cell carcinoma, spatial proteomics, tumor microenvironment, immune compartmentalization, tumor heterogeneity

Funding: LMC acknowledges funding from the National Cancer Institute (NCI) of the National Institutes of Health under award numbers U01CA224012, R01CA223150, R01CA226909, and R21HD099367 and funding from the Brenden-Colson Center for Pancreatic Care at Oregon Health & Science University (OHSU). JG acknowledges funding under award number U24CA231877. LMC and JG acknowledge funding from the OHSU Knight Cancer Institute, NCI award number U2CCA233280, and the Prospect Creek Foundation to the OHSU SMMART (Serial Measurement of Molecular and Architectural Responses to Therapy) Program. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health, OHSU, or the Knight Cancer Institute.

Conflict of Interests: L.M. Coussens is a paid consultant for Cell Signaling
Technologies, AbbVie Inc., and Shasqi Inc., received reagent and/or research support
from Plexxikon Inc., Pharmacyclics, Inc., Acerta Pharma, LLC, Deciphera
Pharmaceuticals, LLC, Genentech, Inc., Roche Glycart AG, Syndax Pharmaceuticals
Inc., Innate Pharma, NanoString Technologies, and Cell Signaling Technologies, is a
member of the Scientific Advisory Boards of Syndax Pharmaceuticals, Carisma
Therapeutics, Zymeworks, Inc, Verseau Therapeutics, Cytomix Therapeutics, Inc., and
Kineta Inc, and is a member of the Lustgarten Therapeutics Advisory working group. No
potential conflicts of interest were disclosed by the other authors.
53
54 **Word count, abstract**: 160
55 **Word count, text:** 4216
56 **Figure count:** 4
57 **Table count:** 2
58

ABSTRACT

Recent research has provided compelling evidence that the spatial organization of cells within the tumor-immune microenvironment (TiME) of solid tumors correlates with survival and response to therapy in numerous cancer types. Here, we report results of a quantitative single-cell spatial analysis of the TiME of primary and recurrent human head and neck squamous cell carcinoma (HNSCC) tumors, that builds upon our initial longitudinal study of these same HNSCCs that annotated immune complexity at near single cell resolution. Herein, we extended multiple spatial algorithms to quantify spatial landscapes of immune cells within TiMEs. Most notably, we report that spatial compartmentalization, rather than mixing, between neoplastic tumor cells and immune cells is associated with longer patient survival, as well as revealing mesenchymal spatial cellular neighborhoods and their association with improved patient outcomes. Results reported herein are concordant with studies in other tumor types, thus indicating that cellular heterogeneity within tumors trends with spatial TiME features, and are likely prognostic for patient survival.

INTRODUCTION

Tumor microenvironments, comprising both neoplastic tumor cells and recruited stromal cells of various lineages, including a diverse assemblage of immune, mesenchymal, and vascular cells, play a key role in both de novo progression of tumors and regulating response to therapies (1-3). Numerous studies have reported that the spatial organization of the tumor-immune microenvironment (TiME), beyond just the types and quantities of cells present in the TiME, is prognostic for survival and response to therapy in multiple cancer types (4-13). Measures that quantify this spatial organization can range from simple density ratios within specific tumor regions (14), such as the Immunoscore (15), a now commonly used biomarker for tumor staging, to more complex measures that account for the precise locations of specific cells relative to other cells, such as mixing scores(4) and cellular neighborhood measures (7). These more advanced spatial quantifications are a result of emerging single-cell multiplex tissue imaging modalities (4, 6, 7, 16-18), that provide detailed phenotypic and effector statuses for each cell, while maintaining the spatial context of the tissue assayed. Knowing the precise locations of cells in the TiME enables a deeper understanding of how cells interact within the tumor, as both direct and indirect cell signaling mechanisms require cells to be near, if not directly adjacent to one another (19). This understanding can then aid treatment decisions, as many therapies require spatial proximity of specific cell types for efficacy (10). Given that single-cell imaging technologies are still quite new, there is much to be discovered regarding how the spatial organization of cells within the TiME relates to clinical outcome, and may be used for patient stratification decisions for therapy.

We previously developed a multiplex immunohistochemistry (mIHC) imaging platform to aid studies investigating the immune contexture of solid tumors and their response to therapies, at the single-cell level (16, 17). Using a sequential antibody staining protocol, detection of 25-50 proteins can be enumerated at single-cell resolution across a single

formalin-fixed paraffin-embedded (FFPE) tissue section. This enables single-cell
phenotyping of discrete leukocyte lineages, capturing their phenotypic effector statuses,
and importantly, reveals their spatial relationships to other cells in the tissue section.
Utilizing this mIHC approach on a small cohort of primary and recurrent human head
and neck squamous cell carcinomas (HNSCCs), we found immune contextures
associated with disease recurrence, most notable that macrophage to CD8[+] T cell ratios
within tumor tissues correlate with outcomes (16, 17). HNSCC is the sixth leading form
of cancer worldwide (20), and it accounts for more than 10,000 deaths per year in the
US alone (21). While patients harboring human papilloma virus (HPV) within neoplastic
cells tend to exhibit a better prognosis, their HPV-negative [HPV(-)] counterparts
typically exhibit T cell suppressive TiMEs and have a significantly greater risk of
recurrence and shorter 3-year survival (22-24). There is a critical need to improve the
understanding of HNSCC TiMEs to enable better patient stratification for therapy, as
well as identify new targets that could be leveraged for therapeutic intervention to
improve outcome, particularly for patients with HPV(-) tumors who currently lack
promising therapeutic options.
121
Here, we built upon our initial study that annotated FFPE sections of nine HPV(-)
matched primary newly diagnosed versus their recurrent HNSCCs via our mIHC
proteomics imaging platform for TiME leukocyte lineage contexture (17). We measured
tumor heterogeneity and compositional changes from primary to recurrent tumors and,
importantly, deployed multiple spatial algorithms to quantify the spatial organization of
the TiMEs. Finally, we correlated these spatial features with survival and identified TiME
architectures associated with survival that could be used to stratify future patients for
therapy. Overall, our results are concordant with those from other studies, indicating
that the features identified herein are likely shared and prognostic across cancer types.
131

METHODS

133
Multiplex immunohistochemistry data generation

135
mIHC is an immunohistochemical-based imaging platform that evaluates sequentially
stained immune lineage epitope-specific antibodies for immunodetection on FFPE
tissue sections (16, 17). Using our computational image processing workflow along with
a hierarchical gating strategy, we are able to phenotype single cells while maintaining
the Cartesian coordinates of each cell on the tissue section. We applied the mIHC
pipeline to analyze matched primary and recurrent FFPE tissue specimens from nine
HPV(-) HNSCC patients. We sequentially stained for 29 antibodies and gated cells as
neoplastic tumor, stromal (mesenchymal), or one of seven different leukocyte subtypes
spanning both lymphoid and myeloid lineages (**Table 1**). One to three regions of 2500^2
μm^2 each were assayed with mIHC per tumor resection. Patient characteristics and
parameters of patient consent are as reported in Banik et al (17). Briefly, study subjects
included nine patients with recurrent HNSCC after definitive surgery, chemotherapy,
and/or radiation therapy. Matched FFPE tissue specimens from the newly diagnosed
primary and recurrent primary tumors for each patient were obtained from the Oregon
Health and Science University Knight Biolibrary. Tissue specimens were deidentified

and coded with a unique identifier prior to analysis. Demographic and clinical data
including HPV status, tobacco and alcohol use, treatment regimens, and survival
outcomes was collected. All HNSCC tumors were staged according to the 8th edition
AJCC/UIC TNM classification and cohort characteristics are shown in **Table 2** (17). All
studies involving human tissue were approved by institutional IRB (protocol #809 and
#3609), and written informed consent was obtained.
157
158 **Tumor heterogeneity analyses**
159
The Kullback-Leibler divergence was calculated using the entropy function from the
Scipy Python package (25). The distribution of each individual tumor region's cellular
composition was compared to the tumor's average cellular composition distribution. Log
base 2 was used for the calculation. Unsupervised hierarchical clustering of each tumor
region was performed using the normalized percent contribution of each cell type
present as the features. Euclidean distance was used to determine distances between
observations.
167
168 **Mixing score analysis**
169
Keren et al. developed a mixing score to quantify the ratio of neoplastic and immune cell
spatial interactions (4). This score is defined as the number of interactions between
neoplastic tumor cells and immune cells divided by the number of interactions between
immune cells and another immune cell within a tumor region. We defined there to be an
interaction between two cells if their centers [the (x,y) coordinates provided by mIHC]
were within 15 µm from one another. We used the median mixing score value (0.1034)
for all primary tumors as the threshold to distinguish between mixed and
compartmentalized spatial organization groups. Tumor regions with a mixing score of
greater than 0.1034 were defined as mixed. Tumor regions with a mixing score of less
than 0.1034 were defined as compartmentalized. We used a density threshold of less
than 250 immune cells per 800^2 $µm^2$ to define tumor regions as cold. We chose this
threshold to match that used by Keren et al.
182
183 **Cellular neighborhood clustering analyses**
184
Cellular neighborhoods were defined by drawing a circle of a specified radius around all
seed cells of a given phenotype. Cells whose centers were inside the circle were
considered neighbors of that seed cell and contributed to that cell's neighborhood. All
neighborhoods across all tumor regions were then clustered using scikit-learn's
MiniBatchKMeans function (26) to perform K-means clustering according to their
normalized cellular composition. The elbow method was used to determine the number
of clusters to form. Each resulting cluster was comprised of neighborhoods with a
similar cellular makeup.
193
194 **Statistics**
195

Independent t-tests were used to determine statistically significant differences for
independent samples whose differences followed a normal distribution. Mann-Whitney
U tests were used to determine statistically significant differences for independent
samples whose differences did not follow a normal distribution. Paired t-tests were used
to determine statistically significant differences for paired samples whose differences
followed a normal distribution. Wilcoxon signed rank tests were used to determine
statistically significant differences for paired samples whose differences did not follow a
normal distribution. P-values less than 0.05 were considered statistically significant. All
statistical calculations were performed with the Scipy package using Python software
(25).
206
207 **Survival analyses**
208
209 Kaplan-Meier curves were generated and a log-rank test was performed using the
210 lifelines package with Python software (27). P-values less than 0.05 were considered
211 statistically significant.
212
213 **Data and software availability**
214
All of the data produced by our mIHC computational image processing pipeline,
including protein abundance, cell phenotype, and cell location information saved in the
form of a matrix, in addition to survival data, is available for download at Zenodo
(https://doi.org/10.5281/zenodo.4584441). All computational analyses in this study were
performed using Python software, version 3.6.5. The code created to produce the
results of this study is available at https://github.com/kblise/HNSCC_mIHC_paper.
221
222 # RESULTS
223
224 **Immune contexture of HNSCC**
225
All mIHC data for this study was generated using the protocol described in Banik et al
(17). Single cells were phenotyped in 47 tumor regions from nine primary and nine
matched recurrent tumors according to the gating strategy shown in **Table 1**, and their
locations within the tumor region were reported as Cartesian coordinates. Patient
demographic and clinical data is shown in **Table 2** (17). With this data, we investigated
tumor heterogeneity both within and across patient samples, quantified the spatial
architecture of the TiME using a mixing score and a neighborhood clustering method,
associated spatial architectures with clinical outcome, and performed additional
analyses to explain these prognostic features.
235
236 **Single-cell analyses reveal varying degrees of tumor heterogeneity**
237
To address the question of how cellular composition varies across tumor regions, we
assessed tumor heterogeneity at multiple levels, including intra-tumoral, inter-tumoral,
and inter-patient cellular heterogeneity by calculating the Kullback-Leibler (KL)
divergence for each region, performing hierarchical clustering, and conducting a

principal component analysis (PCA). The fraction of each cell type present per region
was first computed across all eighteen tumors (**Figure 1A,B**). Within individual tumors,
there was less variability in the number of immune cells present across regions as
compared to the number of neoplastic and non-immune cells present. Our interpretation
of this finding was that the neoplastic cells and non-immune cell populations are
generally the main drivers of intra-tumoral heterogeneity. To further assess this intra-
tumoral heterogeneity in a more quantitative way, we calculated the KL divergence of
each region's distribution of cell types present, as compared to its tumor's average cell
type distribution. KL divergence is a measure of how similar two distributions are; this
approach has been used previously to quantify tumor heterogeneity (6). We found that
all regions had minimal KL divergence values (KL < 0.35; **Figure 1C**), signifying a low
degree of intra-tumoral heterogeneity.
254
To investigate inter-tumoral heterogeneity, we performed unsupervised hierarchical
clustering on the 47 tumor regions based on their normalized cellular composition
(**Figure 1D**). We found that only one patient (in green at the bottom of the heatmap)
contained all five tumor regions clustering together, independent of primary or recurrent
state. All other patients' tumors exhibited a greater degree of inter-tumoral
heterogeneity, as demonstrated by the distance between primary and recurrent tumor
regions in the heatmap's accompanying dendrogram (**Figure 1D**). Finally, we assessed
inter-patient tumor heterogeneity, and found that regions evaluated from the same
patient tended to cluster together more than regions evaluated from different patients
(**Figure 1D**), indicating increased tumor heterogeneity between patients as compared to
between samples from the same patient. PCA results also supported these
observations (**Figure 1E**).
267
268   **TiME cellular composition altered by therapy**
269
Multiple studies have reported differences in TiME cellular makeup (28, 29) and tumor
clonal diversity (30) between primary and recurrent tumors. To detect whether any
immune contexture changes existed following therapy in our cohort, we analyzed the
cellular composition of primary tumors as compared to their recurrent tumors. All
patients received combination therapy of cisplatin or cetuximab accompanied by
radiation following surgical resections of their primary tumors. We used the average
fraction of each cell type present across regions for a given tumor and compared
primary tumor composition to their matched recurrent tumor composition. We observed
a significant difference in the fraction of B cells present in recurrent tumors following
therapy for all patients (p=0.004, **Figure 2A**), with no other common differences found
across patients for any other immune cell type (**Figure 2B**). Further, hierarchical
clustering and PCA results did not yield discrete clusters of regions grouped by their
tumor status (primary or recurrent) (**Figure 1D, 2C**). We then questioned whether
subsets of patients contained similar compositional changes to their TiMEs. To address
this question, we performed unsupervised hierarchical clustering on the normalized
average change in cellular composition across all cell types. This resulted in two distinct
groups of patients; these appear to be most different from one another in the relative
degree of change in their CD8[+] T cell, CD4[+] T helper, and dendritic cell populations

(**Figure 2D**). Interestingly, the two patients that received cetuximab (orange and purple),
rather than cisplatin, clustered together within one of these groups and had similar
overall changes to their tumor compositions following therapy. Together, these results
indicate that there were shared trends in TiME composition changes for subsets of
patients specific to therapy, and regardless of therapy, all patients saw a significant
decrease in B cells from primary to recurrent tumors.
294
295 **TiME cellular composition associated with survival**
296
Prior research has revealed that increased $CD8^+$ T cell abundance is associated with
better outcomes in HNSCC (31-34). Using the average fraction of $CD8^+$ T cells present
in primary tumors, no significant association with survival was found (p=0.423, **Figure**
2E). However, patients with a smaller fraction of macrophages present in their primary
tumors had better progression free survival (PFS) (p=0.017, **Figure 2F**). These findings
are supported by other studies that also identified higher levels of macrophages
associated with more aggressive tumors and decreased survival (35, 36). These
findings expand on the results of our prior study, which reported that patients with
myeloid-inflamed tumors demonstrated decreased PFS than those with lymphoid
inflamed tumors (17). Here we report that this difference can be attributed to a single
tumor feature—the fraction of macrophages present. We also found that patients with a
greater fraction of other non-immune cells ($PANCK^-$, $CD45^-$) correlated with improved
PFS (p=0.003, **Figure 2G**), with no other cell types evaluated demonstrating an
association with survival.
311
312 **Quantifying the spatial organization of tumor and immune cells**
313
To investigate if TiME cellular spatial relationships were prognostic in our cohort of
primary and recurrent HNSCC as has been reported in other cancer types (5-13), we
deployed a mixing score, which was used previously to analyze immune cell spatial
compartmentalization in triple negative breast cancers (4). To do this, we quantified the
number of immune and neoplastic tumor cells within 15 μm of each other, then divided
by the number of immune cells within 15 μm from another immune cell. We labeled
each region as mixed or compartmentalized using the median mixing score value for all
primary tumors as the threshold (**Figure 3A,B**; see Methods). This threshold
categorized tumor regions as mixed when at least one neoplastic tumor cell was within
15 μm from an immune cell for approximately every ten immune cells within 15 μm from
another immune cell. Tumor regions were compartmentalized when this ratio was
smaller. Regions with fewer than 250 $CD45^+$ immune cells per 800^2 $μm^2$ present were
labeled as cold. Of the 47 total tumor regions, 23 were mixed, 20 were
compartmentalized, and four were cold.
328
After calculating the mixing score for each region, we explored how cellular composition
related to spatial organization. Regions with more mixing between neoplastic cells and
immune cells often demonstrated higher presence of neoplastic cells than those with
higher compartmentalization (**Figure 3C**). This is unsurprising given the equation for the
mixing score, as more neoplastic cells will inevitably only contribute to the numerator of

the score, and thus will increase the overall score. However, regions with more
compartmentalization between neoplastic cells and immune cells often also contained
increased counts of alpha smooth muscle actin (αSMA)$^+$ mesenchymal cells as
compared to those with higher mixing (**Figure 3D**). This result was intriguing because
these cells have no effect on the mixing score, yet there is a clear association between
composition and organization. This association is further demonstrated by PCA, where
regions of similar cellular composition cluster together and also tend to contain similar
spatial organizations (**Figure 3E**).
342
We next examined associations between biomarker expression reflecting on bioeffector
phenotypes of leukocytes and mixing scores for each tumor region. Mixed regions
contained a greater proportion of neoplastic cells expressing immunoregulatory
proteins, such as PD-1 and PD-L1, than compartmentalized regions (p=0.0003,
p=0.015, **Figure 3F**). We also found that compartmentalized regions contained a
greater proportion of lymphocytes (CD8$^+$ T cells, CD4$^+$ T cells, and B cells) expressing
the proliferation marker Ki-67 (p=0.020, **Figure 3G**).
350
351 **Spatial compartmentalization associated with longer survival**
352
353 To investigate how spatial mixing correlated with PFS or patient outcome, we averaged
354 the mixing score across regions and assigned a final mixing score and spatial label for
355 each tumor. Patients with more compartmentalization between neoplastic cells and
356 immune cells in their primary tumors exhibited longer PFS than those with more mixing
357 between these cell types (p=0.027, **Figure 3H**).
358
In addition to quantifying TiME spatial organization using a mixing score, we also
computed cellular neighborhoods, that were then clustered to identify common spatial
features across tumors. Neighborhoods represent physical groupings of cells within a
set distance threshold from a seed cell (**Figure 4A**). Each cell within the distance
threshold was deemed a neighbor of the seed cell, contributing to that seed cell's
neighborhood's composition. After identifying all neighborhoods for each seed cell
present across all tumor regions, neighborhoods were grouped with K-means clustering,
using the fraction of each cell type present in the neighborhoods as the clustering
features. This revealed clusters of neighborhoods with similar cellular makeups. We
then used the average neighbor composition of each cluster to perform survival
analyses.
370
Given the relationship between mixing score and αSMA$^+$ mesenchymal cell abundance,
we applied our neighborhood clustering analysis with αSMA$^+$ cells as the seed cells. We
used a distance threshold of 22.5 µm, as this produced neighborhoods with an average
of approximately ten neighbors present. A recent study involving cellular neighborhood
analyses employed a method that selected the ten nearest spatial neighbors of the seed
cell, regardless of the distance between the seed cell and its neighbors (7). By setting a
distance threshold of 22.5 µm, our method required the cells be close, if not directly
touching, while still capturing enough neighbors to cluster on. Clustering results yielded
six groups, each different in their composition (**Figure 4B**). αSMA$^+$ cell neighborhoods

in cluster 1 contained a majority of αSMA$^+$ cells as the neighbors. Cluster 2 contained
both αSMA$^+$ cells as well as other non-immune (CD45$^-$), non-neoplastic cells (panCK$^-$)
as neighbors. Cluster 3 was defined by its large proportion of panCK$^+$ neoplastic cells,
whereas Cluster 4 consisted of roughly half CD45$^+$ immune cells and half other non-
immune, non-neoplastic cells. Cluster 5 contained the highest proportion of CD45$^+$
immune cells, and cluster 6 was dominated by other non-immune and non-neoplastic
cells.
387
We found that patients with a greater proportion of their αSMA$^+$ cells in clusters 2 and 6
in their primary tumors demonstrated longer PFS (cluster 2: p=0.059, cluster 6:
p=0.003, **Figure 4C,D**), thus indicating that patients whose αSMA$^+$ cells were
predominantly surrounded by αSMA$^+$ cells or other non-immune, non-neoplastic cells in
their TiMEs were associated with better outcome. In addition, patients with a greater
proportion of their αSMA$^+$ cells belonging to cluster 3, which included a large number of
neoplastic cells, in their primary tumors demonstrated shorter PFS (p=0.047, **Figure
4E**).
396
397 # DISCUSSION
398
The significant role that TiME cellular composition plays in tumor progression and
response to therapy has been accepted for over a decade (37). However, recent
findings powered by single-cell proteomics imaging technologies have found that the
spatial organization of the cells present in the TiME also plays a critical role (4-13). It is
still unclear exactly which spatial features are important in dictating response to therapy
or clinical outcome, as well as whether these features are shared across cancer types,
and how they could be leveraged for therapeutic decisions and patient stratification.
Here, we leveraged single-cell spatial proteomics data generated by our mIHC
immunoassay-based imaging platform to quantitatively assess the TiME of nine
matched primary and recurrent HPV(-) HNSCCs in order to identify possible prognostic
spatial features. Our results provide novel insights into the heterogeneity and spatial
landscape of HPV(-) HNSCCs, and highlight possible TiME spatial layouts that may
impact clinical outcome across cancer types.
412
Despite the fact that HPV(-) HNSCCs often contain limited immune infiltrates (22), our
analyses revealed significant differences in the immune cell spatial organization within
these tumors that were associated with survival, highlighting the importance of
considering the spatial context of the TiME. Most strikingly, patients whose primary
tumors contained more compartmentalization between their tumor cell and immune cell
populations demonstrated longer PFS. This correlation was also identified in a similar
analysis of triple negative breast cancer patients (4), indicating that a
compartmentalized spatial architecture may play a favorable role in survival across
cancer types. Our analyses of functional marker expression indicate that the immune
cells in regions of compartmentalization are proliferating and are perhaps subject to
reduced direct regulation by neoplastic tumor cells (38-42) than those in regions of
mixing. This increased proliferation and decreased regulation of immune cells in
compartmentalized tumors may contribute to improved survival. We also found that the

presence of spatial cellular neighborhoods consisting of αSMA$^+$ cells and other non-immune and non-neoplastic cells surrounding αSMA$^+$ cells was associated with longer survival. On the contrary, the presence of cellular neighborhoods consisting of neoplastic tumor cells surrounding αSMA$^+$ cells was associated with shorter survival. Given that αSMA is a common marker for cancer-associated fibroblasts (20, 43), whose presence in tumors tends to be associated with tumor growth, metastasis, and resistance to therapy (44-47), our results indicate that beyond just the abundance of these cells, their spatial organization may play a crucial role in determining how a tumor progresses, and/or how it responds to standard therapy. Our cellular neighborhood analyses highlight the importance of considering how multiple cells can work in tandem with one another, beyond single cell-cell interactions, to impact TiME organization, and ultimately, clinical outcome.

We also found concordance between the results from our heterogeneity and composition analyses and those from several other studies. Our results are similar to a study that clustered HNSCC biopsies based on their neoplastic and immune gene signatures as determined by RNA-sequencing, and reported that samples from the same patient were more similar to each other than samples from different patients (48). This trend has also been observed across cancer types, including melanoma (49), hepatocellular carcinoma (50), pancreatic ductal adenocarcinoma (51), and breast cancer (52). Finally, we found that decreased macrophage abundance was associated with longer survival, which has previously been reported in head and neck cancers (35, 36). Collectively, these results demonstrate concordance across multiomic analyses and cancer types.

This concordance provides orthogonal support for our conclusions and indicates that single-cell heterogeneity and spatial organization of tumors may share similarities across different types of cancer and across different molecular assays. Our spatial analyses demonstrate the use of various algorithms to quantify the spatial landscape of tumors using single-cell imaging data. Further, our computational methods provide a framework for future single-cell imaging analyses, as they are applicable to any multiplex imaging assay, including mIHC, co-detection by indexing technology (CODEX) (7), cyclic immunofluorescence (18), and imaging mass cytometry (6). Our results highlight several spatial architectures that may help guide precision medicine approaches for HPV(-) HNSCC patients, including architectures that may help predict which patients will have shorter survival and thus warrant more aggressive therapy and clinical follow-up versus those that predict which patients have an improved survival outlook, where therapy and follow-up could be modulated. While our results provide evidence that the spatial organization of HPV(-) HNSCC tumors plays a role in clinical outcome, future studies on larger cohorts are needed to assess the full prognostic capabilities of the spatial features identified in our analyses.

ACKNOWLEDGEMENTS

The authors thank Dr. Courtney Betts, Justin Tibbitts, Teresa Beechwood, and Meghan Lavoie for regulatory and technical assistance. The authors thank the OHSU Knight

472 Biostatistics Shared Resource for guidance with statistical calculations. The authors
473 also acknowledge and thank the patients who donated tissue samples for this study.

REFERENCES

1. Binnewies M, Roberts EW, Kersten K, Chan V, Fearon DF, Merad M, *et al.* Understanding the tumor immune microenvironment (TIME) for effective therapy. Nat Med **2018**;24(5):541-50.
2. Junttila MR, de Sauvage FJ. Influence of tumour micro-environment heterogeneity on therapeutic response. Nature **2013**;501(7467):346-54.
3. Petitprez F, de Reynies A, Keung EZ, Chen TW, Sun CM, Calderaro J*, et al.* B cells are associated with survival and immunotherapy response in sarcoma. Nature **2020**;577(7791):556-60.
4. Keren L, Bosse M, Marquez D, Angoshtari R, Jain S, Varma S*, et al.* A Structured Tumor-Immune Microenvironment in Triple Negative Breast Cancer Revealed by Multiplexed Ion Beam Imaging. Cell **2018**;174(6):1373-87 e19.
5. Ali HR, Jackson HW, Zanotelli VRT, Danenberg E, Fischer JR, Bardwell H*, et al.* Imaging mass cytometry and multiplatform genomics define the phenogenomic landscape of breast cancer. Nature Cancer **2020**;1(2):163-75.
6. Jackson HW, Fischer JR, Zanotelli VRT, Ali HR, Mechera R, Soysal SD*, et al.* The single-cell pathology landscape of breast cancer. Nature **2020**;578(7796):615-20.
7. Schurch CM, Bhate SS, Barlow GL, Phillips DJ, Noti L, Zlobec I*, et al.* Coordinated Cellular Neighborhoods Orchestrate Antitumoral Immunity at the Colorectal Cancer Invasive Front. Cell **2020**.
8. Farkkila A, Gulhan DC, Casado J, Jacobson CA, Nguyen H, Kochupurakkal B*, et al.* Immunogenomic profiling determines responses to combined PARP and PD-1 inhibition in ovarian cancer. Nat Commun **2020**;11(1):1459.
9. Barua S, Fang P, Sharma A, Fujimoto J, Wistuba I, Rao AUK*, et al.* Spatial interaction of tumor cells and regulatory T cells correlates with survival in non-small cell lung cancer. Lung Cancer **2018**;117:73-9.
10. Gide TN, Silva IP, Quek C, Ahmed T, Menzies AM, Carlino MS*, et al.* Close proximity of immune and tumor cells underlies response to anti-PD-1 based therapies in metastatic melanoma patients. Oncoimmunology **2020**;9(1):1659093.
11. Giraldo NA, Nguyen P, Engle EL, Kaunitz GJ, Cottrell TR, Berry S*, et al.* Multidimensional, quantitative assessment of PD-1/PD-L1 expression in patients with Merkel cell carcinoma and association with response to pembrolizumab. J Immunother Cancer **2018**;6(1):99.
12. Johnson DB, Bordeaux J, Kim JY, Vaupel C, Rimm DL, Ho TH*, et al.* Quantitative Spatial Profiling of PD-1/PD-L1 Interaction and HLA-DR/IDO-1 Predicts Improved Outcomes of Anti-PD-1 Therapies in Metastatic Melanoma. Clin Cancer Res **2018**;24(21):5250-60.
13. Enfield KSS, Martin SD, Marshall EA, Kung SHY, Gallagher P, Milne K*, et al.* Hyperspectral cell sociology reveals spatial tumor-immune cell interactions associated with lung cancer recurrence. J Immunother Cancer **2019**;7(1):13.
14. Galon J, Costes A, Sanchez-Cabo F, Kirilovsky A, Mlecnik B, Lagorce-Pages C*, et al.* Type, density, and location of immune cells within human colorectal tumors predict clinical outcome. Science **2006**;313(5795):1960-4.

15. Galon J, Pages F, Marincola FM, Angell HK, Thurin M, Lugli A, *et al.* Cancer
classification using the Immunoscore: a worldwide task force. J Transl Med
2012;10:205.
16. Tsujikawa T, Kumar S, Borkar RN, Azimi V, Thibault G, Chang YH, *et al.*
Quantitative Multiplex Immunohistochemistry Reveals Myeloid-Inflamed Tumor-
Immune Complexity Associated with Poor Prognosis. Cell Rep **2017**;19(1):203-
17.
17. Banik G, Betts CB, Liudahl SM, Sivagnanam S, Kawashima R, Cotechini T, *et al.*
High-dimensional multiplexed immunohistochemical characterization of immune
contexture in human cancers. Methods Enzymol **2020**;635:1-20.
18. Eng J, Thibault G, Luoh SW, Gray JW, Chang YH, Chin K. Cyclic Multiplexed-
Immunofluorescence (cmIF), a Highly Multiplexed Method for Single-Cell
Analysis. Methods Mol Biol **2020**;2055:521-62.
19. Brucher BL, Jamall IS. Cell-cell communication in the tumor microenvironment,
carcinogenesis, and anticancer treatment. Cell Physiol Biochem **2014**;34(2):213-
43.
20. Alsahafi E, Begg K, Amelio I, Raulf N, Lucarelli P, Sauter T, *et al.* Clinical update
on head and neck cancer: molecular biology and ongoing challenges. Cell Death
Dis **2019**;10(8):540.
21. Cohen EEW, Bell RB, Bifulco CB, Burtness B, Gillison ML, Harrington KJ, *et al.*
The Society for Immunotherapy of Cancer consensus statement on
immunotherapy for the treatment of squamous cell carcinoma of the head and
neck (HNSCC). J Immunother Cancer **2019**;7(1):184.
22. Canning M, Guo G, Yu M, Myint C, Groves MW, Byrd JK, *et al.* Heterogeneity of
the Head and Neck Squamous Cell Carcinoma Immune Landscape and Its
Impact on Immunotherapy. Front Cell Dev Biol **2019**;7:52.
23. Fakhry C, Westra WH, Li S, Cmelak A, Ridge JA, Pinto H, *et al.* Improved
survival of patients with human papillomavirus-positive head and neck squamous
cell carcinoma in a prospective clinical trial. J Natl Cancer Inst **2008**;100(4):261-
9.
24. Ang KK, Harris J, Wheeler R, Weber R, Rosenthal DI, Nguyen-Tan PF, *et al.*
Human papillomavirus and survival of patients with oropharyngeal cancer. N Engl
J Med **2010**;363(1):24-35.
25. Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, *et*
al. SciPy 1.0: fundamental algorithms for scientific computing in Python. Nat
Methods **2020**;17(3):261-72.
26. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, *et al.*
Scikit-learn: Machine Learning in Python. JMLR **2011**;12:2825-30.
27. Davidson-Pilon C, Kalderstam J, Jacobson N, sean-reed, Kuhn B, Zivich P, *et al.*
CamDavidsonPilon/lifelines: v0.21.0. Zenodo **2020**.
28. Predina J, Eruslanov E, Judy B, Kapoor V, Cheng G, Wang LC, *et al.* Changes in
the local tumor microenvironment in recurrent cancers may explain the failure of
vaccines after surgery. Proc Natl Acad Sci U S A **2013**;110(5):E415-24.
29. Zheng B, Wang J, Cai W, Lao I, Shi Y, Luo X, *et al.* Changes in the tumor
immune microenvironment in resected recurrent soft tissue sarcomas. Ann
Transl Med **2019**;7(16):387.

30. Walens A, Lin J, Damrauer JS, McKinney B, Lupo R, Newcomb R, *et al.*
Adaptation and selection shape clonal evolution of tumors during residual
disease and recurrence. Nat Commun **2020**;11(1):5017.
31. Hartman DJ, Ahmad F, Ferris RL, Rimm DL, Pantanowitz L. Utility of CD8 score
by automated quantitative image analysis in head and neck squamous cell
carcinoma. Oral Oncol **2018**;86:278-87.
32. Balermpas P, Rodel F, Rodel C, Krause M, Linge A, Lohaus F, *et al.* CD8+
tumour-infiltrating lymphocytes in relation to HPV status and clinical outcome in
patients with head and neck cancer after postoperative chemoradiotherapy: A
multicentre study of the German cancer consortium radiation oncology group
(DKTK-ROG). Int J Cancer **2016**;138(1):171-81.
33. Nasman A, Romanitan M, Nordfors C, Grun N, Johansson H, Hammarstedt L, *et*
al. Tumor infiltrating CD8+ and Foxp3+ lymphocytes correlate to clinical outcome
and human papillomavirus (HPV) status in tonsillar cancer. PLoS One
2012;7(6):e38711.
34. Nordfors C, Grun N, Tertipis N, Ahrlund-Richter A, Haeggblom L, Sivars L, *et al.*
CD8+ and CD4+ tumour infiltrating lymphocytes in relation to human
papillomavirus status and clinical outcome in tonsillar and base of tongue
squamous cell carcinoma. Eur J Cancer **2013**;49(11):2522-30.
35. Marcus B, Arenberg D, Lee J, Kleer C, Chepeha DB, Schmalbach CE, *et al.*
Prognostic factors in oral cavity and oropharyngeal squamous cell carcinoma.
Cancer **2004**;101(12):2779-87.
36. Wolf GT, Chepeha DB, Bellile E, Nguyen A, Thomas D, McHugh J, *et al.* Tumor
infiltrating lymphocytes (TIL) and prognosis in oral cavity squamous carcinoma: a
preliminary study. Oral Oncol **2015**;51(1):90-5.
37. Fridman WH, Pages F, Sautes-Fridman C, Galon J. The immune contexture in
human tumours: impact on clinical outcome. Nat Rev Cancer **2012**;12(4):298-
306.
38. Freeman GJ, Long AJ, Iwai Y, Bourque K, Chernova T, Nishimura H, *et al.*
Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family member
leads to negative regulation of lymphocyte activation. J Exp Med
2000;192(7):1027-34.
39. Akinleye A, Rasool Z. Immune checkpoint inhibitors of PD-L1 as cancer
therapeutics. J Hematol Oncol **2019**;12(1):92.
40. Kleffel S, Posch C, Barthel SR, Mueller H, Schlapbach C, Guenova E, *et al.*
Melanoma Cell-Intrinsic PD-1 Receptor Functions Promote Tumor Growth. Cell
2015;162(6):1242-56.
41. Du S, McCall N, Park K, Guan Q, Fontina P, Ertel A, *et al.* Blockade of Tumor-
Expressed PD-1 promotes lung cancer growth. Oncoimmunology
2018;7(4):e1408747.
42. Li H, Li X, Liu S, Guo L, Zhang B, Zhang J, *et al.* Programmed cell death-1 (PD-
1) checkpoint blockade in combination with a mammalian target of rapamycin
inhibitor restrains hepatocellular carcinoma growth induced by hepatoma cell-
intrinsic PD-1. Hepatology **2017**;66(6):1920-33.
43. Parajuli H, Teh MT, Abrahamsen S, Christoffersen I, Neppelberg E, Lybak S, *et*
al. Integrin alpha11 is overexpressed by tumour stroma of head and neck

squamous cell carcinoma and correlates positively with alpha smooth muscle
actin expression. J Oral Pathol Med **2017**;46(4):267-75.

44.   Shiga K, Hara M, Nagasaki T, Sato T, Takahashi H, Takeyama H. Cancer-
Associated Fibroblasts: Their Characteristics and Their Roles in Tumor Growth.
Cancers (Basel) **2015**;7(4):2443-58.

45.   Tao L, Huang G, Song H, Chen Y, Chen L. Cancer associated fibroblasts: An
essential role in the tumor microenvironment. Oncol Lett **2017**;14(3):2611-20.

46.   Sahai E, Astsaturov I, Cukierman E, DeNardo DG, Egeblad M, Evans RM*, et al.*
A framework for advancing our understanding of cancer-associated fibroblasts.
Nat Rev Cancer **2020**;20(3):174-86.

47.   Johansson AC, Ansell A, Jerhammar F, Lindh MB, Grenman R, Munck-Wikland
E*, et al.* Cancer-associated fibroblasts induce matrix metalloproteinase-mediated
cetuximab resistance in head and neck squamous cell carcinoma cells. Mol
Cancer Res **2012**;10(9):1158-68.

48.   Wood O, Clarke J, Woo J, Mirza AH, Woelk CH, Thomas GJ*, et al.* Head and
Neck Squamous Cell Carcinomas Are Characterized by a Stable Immune
Signature Within the Primary Tumor Over Time and Space. Clin Cancer Res
2017;23(24):7641-9.

49.   Reuben A, Spencer CN, Prieto PA, Gopalakrishnan V, Reddy SM, Miller JP*, et*
al. Genomic and immune heterogeneity are associated with differential
responses to therapy in melanoma. NPJ Genom Med **2017**;2.

50.   Zhang Q, Lou Y, Yang J, Wang J, Feng J, Zhao Y*, et al.* Integrated multiomic
analysis reveals comprehensive tumour heterogeneity and novel
immunophenotypic classification in hepatocellular carcinomas. Gut
2019;68(11):2019-31.

51.   Lin W, Noel P, Borazanci EH, Lee J, Amini A, Han IW*, et al.* Single-cell
transcriptome analysis of tumor and stromal compartments of pancreatic ductal
adenocarcinoma primary tumors and metastatic lesions. Genome Med
2020;12(1):80.

52.   Gyanchandani R, Lin Y, Lin HM, Cooper K, Normolle DP, Brufsky A*, et al.*
Intratumor Heterogeneity Affects Gene Expression Profile Test Prognostic Risk
Stratification in Early Breast Cancer. Clin Cancer Res **2016**;22(21):5362-9.

Cell Phenotype	Antibody Markers
CD8$^+$ T Cell	CD45$^+$ CD20$^-$ CD3$^+$ CD8$^+$
CD4$^+$ T Helper	CD45$^+$ CD20$^-$ CD3$^+$ CD8$^-$ FOXP3$^-$
CD4$^+$ Regulatory T Cell	CD45$^+$ CD20$^-$ CD3$^+$ CD8$^-$ FOXP3$^+$
B Cell	CD45$^+$ CD20$^+$
Macrophage	CD45$^+$ CD20$^-$ CD3$^-$ CD66B$^-$ CD68$^+$
Granulocyte	CD45$^+$ CD20$^-$ CD3$^-$ CD66B$^+$
Dendritic Cell	CD45$^+$ CD20$^-$ CD3$^-$ CD66B$^-$ CD68$^-$ MHCII$^+$
Other Immune	CD45$^+$ CD20$^-$ CD3$^-$ CD66B$^-$ CD68$^-$ MHCII$^-$
αSMA$^+$ Mesenchymal	CD45$^-$ PANCK$^-$ αSMA$^+$
Other Non-Immune	CD45$^-$ PANCK$^-$ αSMA$^-$
Neoplastic Tumor	CD45$^-$ PANCK$^+$
Additional Functional Markers: PD-1, PD-L1, Ki-67	

Table 1. mIHC cell phenotype classification. Antibody markers used to hierarchically phenotype cells into one of eleven phenotype classifications. Additional functional markers stained are listed at the bottom.

Characteristic	n (%)
Gender	Female: 7 (78%) Male: 2 (22%)
Race	Asian: 1 (11%) White: 8 (89%)
Average Age at Primary Surgery	61 years
Alcohol History	Yes: 7 (78%) No: 2 (22%)
Smoking History	Yes: 5 (56%) No: 4 (44%)
HPV Status	Positive: 0 (0%) Negative: 9 (100%)
Tumor Site	Larynx: 3 (33%) Oral Cavity: 4 (44%) Oropharynx: 2 (22%)
TNM Stage	Stage 1: 2 (22%) Stage 2: 1 (11%) Stage 3: 1 (11%) Stage 4: 5 (56%)
Therapy Following Primary Resection	Cisplatin + Radiation: 6 (67%) Cetuximab + Radiation: 3 (33%)
Average Progression Free Survival	504 days (1.4 years)

Table 2. Patient and tumor characteristics. Adapted from Banik et al (17).

Figure 1. Heterogeneity across patients and tumor regions. A, Fraction (out of 1.0) of each cell type present per individual primary (P) and recurrent (R) tumor. Each dot represents a single tumor region (n=47). **B**, Fraction (out of 1.0) of each immune cell type present per individual primary (P) and recurrent (R) tumor. Each dot represents a single tumor region (n=47). **C**, Dot plot of the Kullback-Leibler divergence from a single tumor region's cellular distribution compared to the tumor's average cellular distribution. **D**, Heatmap of cellular composition across tumor regions. Rows are individual tumor regions that are ordered by the hierarchical clustering of their cellular composition. Columns are the cell types used as clustering features. Composition was normalized [0,1] before clustering. Leftmost column is color coded by patient. Next column is color coded by tumor type. **E**, Principal component analysis on cellular composition. Each dot (n=47) represents one tumor region and is colored by patient.

Figure 2. Tumor cellular composition changes following therapy and composition associations with survival. A, Box plot showing the percent that each cell type contributes to each tumor's cellular makeup. Each dot represents the percent contribution of that cell type for one tumor, averaged across regions (n=9 primary tumors, n=9 recurrent tumors). Statistical significance calculated using parametric paired two-tailed t-tests for cell types whose differences follow a normal distribution and non-parametric paired two-tailed t-tests (Wilcoxon signed rank test) for cell types whose differences do not follow a normal distribution. **B**, Line plots showing the change in cellular composition from primary to recurrent tumors for each cell type. Each dot represents the percent contribution of that cell type for one tumor, averaged across regions. Lines are colored by patient. **C**, Principal component analysis on cellular composition. Each dot (n=47) represents one tumor region and is colored by its primary or recurrent designation. **D**, Heatmap of average change in cellular composition from primary to recurrent tumors for each patient. Rows are individual patients that are ordered by the hierarchical clustering of their change in TiME cellular composition (averaged across regions). Columns are the cell types used as clustering features. Compositional change was normalized [0,1] before clustering. Leftmost column is color coded by patient. **E**, Kaplan-Meier curve of progression free survival for patients split by the percentage of CD8$^+$ T cells in their primary tumors. Patients were split on the median value. P-value calculated using the log-rank test. **F**, Kaplan-Meier curve of progression free survival for patients split by the percentage of macrophages in their primary tumors. Patients were split on the median value. P-value calculated using the log-rank test. **G**, Kaplan-Meier curve of progression free survival for patients split by the percentage of other non-immune, non-neoplastic cells in their primary tumors. Patients were split on the median value. P-value calculated using the log-rank test.

Figure 3. Mixing score quantifies the spatial organization of tumors. A, Mixed tumor region (top) versus compartmentalized tumor region (bottom). Leftmost panel is the mIHC image of the tumor region (red = neoplastic tumor; blue = non-neoplastic, non-immune; other colors = immune). Remaining panels show scatterplot reconstructions of the mIHC image, excluding the non-immune, non-neoplastic cells. Each dot represents one cell, and is colored by cell type. **B**, Box plot showing the mixing scores across all primary and recurrent tumors (n=18). Each dot (n=47) represents the mixing score for one tumor region and is colored according to its spatial architecture. The average spatial architecture designation for the overall tumor is printed above each box. **C**, Bar chart showing the raw count of neoplastic tumor cells present per tumor region. Bars are ordered by neoplastic cell count and are colored according to the region's spatial architecture. **D**, Bar chart showing the raw count of αSMA^+ mesenchymal cells present per tumor region. Bars are ordered by αSMA^+ cell count and are colored according to the region's spatial architecture. **E**, Principal component analysis on cellular composition. Each dot (n=47) represents one tumor region and is colored according to the region's spatial architecture. **F**, PD-1 and PD-L1 expression on neoplastic tumor cells by spatial architecture. Each dot represents the percentage of neoplastic cells positive for that immunoregulatory protein for a single tumor region (n=43, excluding cold regions). P-values calculated using a one-tailed Mann-Whitney U test. **G**, Ki-67 expression on lymphocytes ($CD8^+$ T cells, $CD4^+$ Regulatory T cells, $CD4^+$ T Helper, and B cells) by spatial architecture. Each dot represents the percentage of lymphocytes positive for Ki-67 for a single tumor region (n=43, excluding cold regions). P-value calculated using a one-tailed t-test. **H**, Kaplan-Meier curve of progression free survival for patients split by the mixing score of their primary tumors. Patients were split on the median value. Results exclude cold tumor regions. P-value calculated using the log-rank test.

Figure 4. αSMA⁺ mesenchymal cellular neighborhood clustering. **A**, Cellular neighborhoods were defined by drawing a circle of a specified radius around each seed cell (green) of a designated phenotype. Cells whose centers were inside the circle were considered neighbors of that seed cell. This figure was created using BioRender.com. **B**, Stacked bar chart showing the average cellular composition of each αSMA⁺ mesenchymal cell neighborhood cluster (n=6). Bars are colored by cell type and represent the average fraction (out of 1.0) of each cell type present in the neighborhoods belonging to each cluster. **C**, Kaplan-Meier curve of progression free survival for patients split by the percentage of αSMA⁺ mesenchymal cell neighborhoods belonging to cluster 2 in their primary tumors. Patients were split on the median value. P-value calculated using the log-rank test. **D**, Kaplan-Meier curve of progression free survival for patients split by the percentage of αSMA⁺ mesenchymal cell neighborhoods belonging to cluster 6 in their primary tumors. Patients were split on the median value. P-value calculated using the log-rank test. **E**, Kaplan-Meier curve of progression free survival for patients split by the percentage of αSMA⁺ mesenchymal cell neighborhoods belonging to cluster 3 in their primary tumors. Patients were split on the median value. P-value calculated using the log-rank test.