Single-Cell Spatial Proteomics Analyses of Head and Neck Squamous Cell Carcinoma Reveal Tumor Heterogeneity and Immune Architectures Associated with Clinical Outcome

Katie E. Blise[a,b], Shamilene Sivagnanam[b,c], Grace L. Banik[c,d*], Lisa M. Coussens[b,c], and Jeremy Goecks[a,b,**]

[a]Department of Biomedical Engineering, Oregon Health & Science University, Portland, OR USA
[b]The Knight Cancer Institute, Oregon Health & Science University, Portland, OR USA
[c]Department of Cell, Developmental & Cancer Biology, Oregon Health & Science University, Portland, OR USA
[d]Otolaryngology-Head & Neck Surgery, Oregon Health & Science University, Portland, OR USA
*Current affiliation: Division of Otolaryngology, Children's Hospital of Philadelphia, Philadelphia, PA USA
**Corresponding author

Correspondence:
Jeremy Goecks, Ph.D.
Associate Professor, Department of Biomedical Engineering
Oregon Health & Science University
3181 SW Sam Jackson Park Road, Mail Code: KR-COM, Portland, OR 97239
Tel: 503-494-0490, Email: goecksj@ohsu.edu

Running Title: Single-cell spatial analysis of head and neck cancer

Keywords: head and neck squamous cell carcinoma, spatial proteomics, tumor microenvironment, immune compartmentalization, tumor heterogeneity

Funding: KEB acknowledges funding from the National Cancer Institute (NCI) of the National Institutes of Health under award number 5T32CA254888. LMC acknowledges funding from the NCI awards U01CA224012, R01CA223150, R01CA226909, and R21HD099367 and funding from the Brenden-Colson Center for Pancreatic Care at Oregon Health & Science University (OHSU). JG acknowledges funding under award number U24CA231877. LMC and JG acknowledge funding from the OHSU Knight Cancer Institute, NCI award number U2CCA233280, and the Prospect Creek Foundation to the OHSU SMMART (Serial Measurement of Molecular and Architectural Responses to Therapy) Program. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health, OHSU, or the Knight Cancer Institute.

Conflict of Interests: L.M. Coussens is a paid consultant for Cell Signaling Technologies, AbbVie Inc., and Shasqi Inc., received reagent and/or research support from Plexxikon Inc., Pharmacyclics, Inc., Acerta Pharma, LLC, Deciphera Pharmaceuticals, LLC, Genentech, Inc., Roche Glycart AG, Syndax Pharmaceuticals Inc., Innate Pharma, NanoString Technologies, and Cell Signaling Technologies, is a member of the Scientific Advisory Boards of Syndax Pharmaceuticals, Carisma Therapeutics, Zymeworks, Inc, Verseau Therapeutics, Cytomix Therapeutics, Inc., Hibercell., Inc., Alkermes, Inc., and Kineta Inc, and is a member of the Lustgarten Therapeutics Advisory working group. No potential conflicts of interest were disclosed by the other authors.

Word count, abstract: 180
Word count, main text: 6001
Figure count: 5 main, 3 supplemental
Table count: 3

ABSTRACT

There is increasing evidence that the spatial organization of cells within the tumor-

immune microenvironment (TiME) of solid tumors influences survival and response to

therapy in numerous cancer types. Here, we report results and demonstrate the

applicability of quantitative single-cell spatial proteomics analyses in the TiME of

primary and recurrent human papillomavirus (HPV)-negative head and neck squamous

cell carcinoma (HNSCC) tumors. Single-cell compositions of a nine patient, primary and

recurrent (n=18), HNSCC cohort is presented, followed by deeper investigation into the

spatial architecture of the TiME and its relationship with clinical variables and

progression free survival (PFS). Multiple spatial algorithms were used to quantify the

spatial landscapes of immune cells within TiMEs and demonstrate that neoplastic

tumor-immune cell spatial compartmentalization, rather than mixing, is associated with

longer PFS. Mesenchymal (αSMA$^+$) cellular neighborhoods describe distinct immune

landscapes associated with neoplastic tumor-immune compartmentalization and

improved patient outcomes. Results from this investigation are concordant with studies

in other tumor types, suggesting that trends in TiME cellular heterogeneity and spatial

organization may be shared across cancers and may provide prognostic value in

multiple cancer types.

INTRODUCTION

Tumor microenvironments, comprising both neoplastic tumor cells and recruited stromal

cells of various lineages, including a diverse assemblage of immune, mesenchymal,

and vascular cells, play a key role in both de novo progression of tumors and regulating

response to therapies[1-3]. Numerous studies have reported that, in addition to the types

and quantities of cells present in the tumor immune-microenvironment (TiME), the

spatial organization of the TiME is prognostic for survival and response to therapy in

multiple cancer types[4-13]. Metrics that quantify this spatial organization can range from

simple density ratios within specific tumor regions[14], such as the Immunoscore[15], a now

commonly used biomarker for colorectal tumor staging, to more complex measures that

account for the precise locations of specific cells relative to other cells, such as mixing

scores[4] and cellular neighborhood measures[7]. These more advanced spatial

quantifications are a result of emerging single-cell multiplex tissue imaging

modalities[4,6,7,16-18], which provide detailed phenotypic and effector proteomic markers

for each cell, while maintaining the spatial architecture of the tissue assayed. Knowing

the precise locations of cells in the TiME enables a deeper understanding of how cells

interact within the tumor, as both direct and indirect cell signaling mechanisms require

cells to be near, if not directly adjacent to one another[19]. This understanding can aid

treatment decisions, as many therapies require spatial proximity of specific cell types for

efficacy[10]. Given that single-cell imaging technologies are still relatively new, there is

much to be discovered regarding how the spatial organization of cells within the TiME

relates to clinical outcome and may be used for patient stratification decisions for

therapy.

Head and neck squamous cell carcinoma (HNSCC) is the sixth leading form of cancer

worldwide[20], and it accounts for more than 10,000 deaths per year in the US alone[21].

While patients harboring human papillomavirus (HPV) within neoplastic cells tend to

exhibit a better prognosis, their HPV-negative [HPV(-)] counterparts typically exhibit T

cell suppressive TiMEs and have a significantly greater risk of recurrence and shorter 3-

year survival[22-24]. There is a critical need to improve understanding of HNSCC TiMEs to

enable better patient stratification for therapy, as well as identify new targets that could

be leveraged for therapeutic intervention to improve outcome, particularly for patients

with HPV(-) tumors who currently lack promising therapeutic options. We previously

developed a multiplex immunohistochemistry (mIHC) imaging platform to aid studies

investigating the immune contexture of solid tumors and their response to therapies at

the single-cell level[16,17]. Using a sequential antibody staining protocol, detection of 12-

30 proteins can be enumerated at single-cell resolution across a single formalin-fixed

paraffin-embedded (FFPE) tissue section. This enables single-cell phenotyping of

discrete leukocyte lineages, and importantly, reveals their spatial relationships with

other cells in the tissue section. Utilizing this mIHC approach on a small cohort of

eighteen HPV(-) primary and matched recurrent HNSCC tumor samples collected from

nine patients, we previously reported immune contextures associated with disease

recurrence, most notably that myeloid inflamed profiles in primary tumors exhibited

shorter progression free survival (PFS) compared to lymphoid inflamed profiles[16,17].

In this study, we have significantly extended our prior analysis of this cohort, focusing

on tumor heterogeneity and compositional changes from primary to recurrent tumors, in

addition to using multiple spatial algorithms to quantify the spatial organization of the

TiMEs. We then correlated these spatial features with PFS and identified TiME

architectures that may be important for therapeutic decision making. Overall, we found

increased neoplastic tumor-immune cell spatial compartmentalization in primary tumors

to be associated with longer PFS. These tumors also contained alpha smooth muscle

actin (αSMA$^+$) cells with more organized structure located near T and B cells, as well as

near cells involved in antigen presentation. Our results are concordant with those from

other studies, indicating that the features identified herein are likely shared and

prognostic across cancer types.

RESULTS

One to three regions of 2500^2 μm^2 from each of the nine patients primary and matched

recurrent tumor resections (n=18) were analyzed, for a total of 47 regions (**Table 1**,

Figure 1a). For this study, we utilized a gating strategy with thirteen lineage or

functional protein biomarkers to classify cells as neoplastic tumor cells, stromal cells

(mesenchymal), or one of seven different leukocyte subtypes spanning lymphoid and

myeloid lineages (**Table 2, Supplemental Figure 1a**). We investigated tumor

heterogeneity both within and across patient samples, quantified the cellular spatial

relationships within the TiME using a mixing score and performed a neighborhood

clustering method to describe the association between TiME spatial architecture, clinical

features, and PFS.

Single-cell proteomic analyses reveal varying degrees of tumor heterogeneity

To quantify how cellular composition varied across tumor regions, we assessed tumor

heterogeneity at multiple levels, including intra-tumoral, intra-patient, and inter-patient

cellular heterogeneity by calculating Kullback-Leibler (KL) divergences for each region,

performing hierarchical clustering, and conducting a principal component analysis

(PCA). The density of each cell type per region was measured for all eighteen tumor

specimens by taking the count of each cell type divided by the measured tissue area in

mm^2 (**Figure 1b,c**). We then calculated the coefficient of variation per cell type for each

tumor, and averaged these values to quantitatively describe the cell types contributing

most to intra-tumoral heterogeneity within the cohort. The coefficient of variation is

defined as the standard deviation divided by the mean, and it provides a normalized

measure of variability for comparison across cell types with large differences in

densities. On average, B cells exhibited the greatest coefficient of variation across the

cohort relative to other cell types (**Table 3**). This is likely due to the fact that B cells were

frequently observed to be spatially clustered together, resulting in regions of either high

B cell density or low B cell density despite being collected from the same tumor

(**Supplemental Figure 1b**).

To further quantify and assess tumor heterogeneity both within and across patients, we

calculated the KL divergence of each tumor region from five average cell type

distributions. KL divergence is a relative measure of how similar two distributions are,

with larger values reflecting less similarity between the distributions and smaller values

reflecting more similarity between the distributions. This measure has been used

previously to quantify tumor heterogeneity[6]. By calculating and comparing the

divergences of each tumor region from multiple average cell type distributions, we were

able to assess heterogeneity within and across tumors and patients. Overall, we

observed that heterogeneity was lower across regions from the same tumor and tumors

from the same patient (primary or recurrent), while higher across tumors from different

patients. This is evidenced by smaller intra-tumoral and intra-patient KL divergence

values for the majority of tumor regions (**Figure 1d**).

The cellular distributions used to calculate the five KL divergence values per tumor

region were (1) the average cellular distribution across all regions sampled from the

same tumor ["Intra-Tumor (P or R only)"]; (2) the average cellular distribution across the

patient's primary and recurrent tumors ["Intra-Patient (P and R)"]; (3) the average

cellular distribution across all tumors in the cohort collected from the same timepoint

["Inter-Patient (P or R only)"]; (4) the average cellular distribution across all tumors in

the cohort resected from the same anatomic site ["Inter-Patient (Same Anatomic Site)"],

and; (5) the average cellular distribution across all tumors collected from all patients in

the cohort, regardless of primary or recurrent status or anatomic site ["Inter-Patient

(all)"]. By comparing the relative KL divergence values to each other, we found tumor

regions to be more similar to regions sampled from the same tumor and patient than

regions collected from tumors of other patients. Notably, we found no significant

difference between "Inter-Patient (Same Anatomic Site)" and "Inter-Patient (all),"

indicating that tumor regions diverged by the same degree from regions sampled at the

same anatomic site as they did from regions sampled at all three anatomic sites (oral

cavity, oropharynx, larynx) of the head and neck region. Finally, given the large

proportion of neoplastic tumor cells comprising the TiME for many of the tumor regions,

we assessed KL divergence using only the distribution of immune cells present and

found similar results (**Supplemental Figure 1c**). This indicates that immune cell

composition is more similar within regions from the same patient than across regions

collected from different patients.

To further investigate intra-patient heterogeneity, we performed unsupervised

hierarchical clustering on the 47 tumor regions based on their normalized density

composition (**Figure 1e**). We found that two patients (3, green; 7, pink) contained all

tumor regions clustering together, independent of primary or recurrent state. These

patients also had the smallest intra-patient KL divergence values (**Figure 1d**), indicating

that the cell densities of these patients' primary and recurrent tumors were similar to

each other. Three patients (2, orange; 4, red; 9, yellow) contained nearly all regions

clustered together. The remaining four patients' tumors exhibited greater degrees of

intra-patient heterogeneity, as demonstrated by the distance between primary and

recurrent tumor regions on the clustered heatmap (**Figure 1e**). Overall, we found that

regions evaluated from the same patient tended to cluster together more than regions

evaluated from different patients (**Figure 1e**), indicating increased heterogeneity

between patients as compared to between samples from the same patient. We also

examined whether tumor regions clustered by the anatomic resection site and found

that the clusters formed did not group by site. These results, in addition to those of our

KL divergence analyses, indicate that anatomic site was likely not the main contributor

of cellular heterogeneity in this cohort. PCA results also supported these observations

(**Figure 1f,g**).

TiME cellular composition altered by therapy

Multiple studies have reported differences in TiME cellular makeup[25,26] and tumor clonal

diversity[27] between primary and recurrent tumors. To assess whether any immune

contexture changes occurred following post-operative therapy in our cohort, we

analyzed the cellular composition of primary tumors as compared to their recurrent

tumors. All patients received a combination therapy of cisplatin and/or cetuximab

accompanied by radiation following surgical resections of their primary tumors. We used

the average density of each cell type present across regions for a given tumor and

compared primary tumor composition to their matched recurrent tumor composition.

While we did not observe any significant differences in cell density between primary and

recurrent tumors (p>0.112), we did find that all patients experienced a decrease in the

density of B cells from their primary to recurrent tumors (**Figure 2a,b**). This result is

supported by a recent study that found that a large cohort of HNSCC patients

experienced a decline in B cells from primary to recurrent tumors, and this reduction

was dependent upon patients receiving adjuvant chemoradiotherapy[28]. We found no

common differences across all patients for any other cell type, although eight of the nine

patients saw a decrease in the density of neoplastic tumor cells from their primary to

recurrent tumors (**Figure 2b**).

We then assessed whether patients exhibited similar compositional differences in

primary and recurrent TiMEs by performing unsupervised hierarchical clustering on the

normalized average difference in cellular composition for each cell type. This resulted in

two groups of patients (**Figure 2c**). Interestingly, the two patients that received

cetuximab (2, orange; 5, purple), rather than cisplatin, clustered together within one of

these groups and were the only two patients to experience a decrease in the density of

every cell type following therapy. Across the cohort these patients had the greatest

decrease in the density of macrophages, granulocytes, and other $CD45^+$ immune cells

present from their primary tumors to their recurrent tumors following therapy (**Figure**

2b,c). Interestingly, the one patient that received both cisplatin and cetuximab (7, pink)

was present in the other cluster from the two patients that received only cetuximab. This

was the only patient to experience an increase in the density of neoplastic tumor cells

(**Figure 2b,c**). This patient also experienced the second largest increase in $CD8^+$ T cells

as well as the greatest decrease in $CD4^+$ regulatory T cells, potentially indicating a pro-

inflammatory response to—or despite—increased neoplastic tumor cell density.

Altogether, these results indicate that shared trends in TiME composition changes from

primary to recurrent tumors specific to therapy exist, and regardless of therapy, all

patients exhibited a significant decrease in B cells from primary to recurrent tumors.

Approximately half (n=4) of the patients in the cohort experienced an increase in $CD8^+$ T

cell density while the other half (n=5) experienced a decrease in $CD8^+$ T cell density

following therapy. This was the only cell type that increased in density for nearly half of

the cohort and decreased for the other half. To determine if there was a survival

advantage for patients that experienced this increase, we split our cohort into short-term

or long-term survivor groups using median PFS and observed that all patients who

experienced an increase in $CD8^+$ T cell density from their primary to recurrent tumors

were long-term survivors (**Figure 2d**). Interestingly, the density of $CD8^+$ T cells in the

primary tumor alone did not associate with PFS (p=0.829). Prior research has revealed

that increased CD8[+] T cell abundance in the primary tumor is associated with better

outcomes in HNSCC[29-33]. However, these studies largely included HPV-positive

HNSCCs, which is more often associated with greater densities of CD8[+] T cells and

improved survival[22-24], thus unsurprising that our results differ. However, our results are

concordant with a recent study in HNSCC that reported longer survival was associated

with patients who had experienced an increase in CD8[+] tumor-infiltrating lymphocytes

from their primary to recurrent tumor[34]. Another study in HNSCC found a similar trend

between increased CD8[+] T cell infiltration, longer survival, presence of specific

neoantigens, and increased cytolytic activity in recurrent tumors[35]. Notably, the four

patients in our cohort that experienced the greatest decrease in CD8[+] T cell density in

recurrence had TNM stage 4 primary tumors, while patients that experienced an

increase in CD8[+] T cell density in recurrence included TNM stages 1, 2, 3, and 4.

Quantifying the spatial organization of neoplastic and immune cells

Prior studies reported the abundance of various cell types, including CD8[+] T cells[33],

CD4[+] regulatory T cells[36], and macrophages[37] to be associated with survival in HNSCC.

We analyzed the average density of each cell type across primary tumors for their

correlations with PFS, but found no significant association with PFS for any single cell

type (p>0.159). Given the prognostic potential of TiME cellular spatial organization as

has been reported in other cancer types[5-13], we quantified the spatial organization of

cells within tumor regions and examined the association of the spatial features with

clinical outcome.

We first deployed a mixing score, used previously to analyze immune cell spatial

compartmentalization in triple negative breast cancers[4]. The mixing score measures the

enrichment of neoplastic tumor-immune cell proximity relative to immune-immune cell

proximity within a set distance. We quantified the number of immune and neoplastic

tumor cells within 15 μm of each other, divided by the number of immune cells within 15

μm from another immune cell. Each region was labeled as mixed or compartmentalized

using the median mixing score value for all primary tumors as the threshold (**Figure**

3a,b; see Methods). This threshold classified tumor regions as mixed if at least one

neoplastic tumor cell was within 15 μm from an immune cell for approximately every ten

immune cells within 15 μm from another immune cell. Tumor regions were considered

compartmentalized when this ratio was smaller. Regions with fewer than 250 CD45[+]

immune cells per 800^2 $μm^2$ present were labeled as cold, utilizing the same immune cell

density threshold from the original study[4]. Of the 47 total tumor regions, 25 were mixed,

20 were compartmentalized, and two were cold.

After calculating the mixing score for each region, we examined the spatial

heterogeneity of our cohort. Our tumor compositional heterogeneity analyses revealed

that intra-tumoral and intra-patient heterogeneity was less than inter-patient

heterogeneity. To determine whether this observation held for spatial organization

heterogeneity, we compared each region's mixing score to five groups of average

mixing scores, which were computed from the same five groups as the analysis in

Figure 1d: Intra-Tumor (P or R only), Intra-Patient (P and R), Inter-Patient (P or R only),

Inter-Patient (Same Anatomic Site), and Inter-Patient (all). Contrary to our analysis in

Figure 1d, the mixing score is only one feature, not a distribution of features, thus we

used the difference in mixing scores, rather than the KL divergence. In addition, we

calculated the absolute values of these differences as a way to normalize the data in

order to capture the degree of difference in spatial organization, allowing us to

subsequently test for differences across the five levels of heterogeneity. We found there

to be less intra-tumoral heterogeneity than inter-patient heterogeneity (**Supplemental**

Figure 2a). This result indicates that, in terms of neoplastic tumor-immune cell mixing,

tumor regions resemble regions sampled from the same tumor more than regions

sampled from tumors of other patients.

We then considered whether tumor regions sampled from the same anatomic site

differed in their spatial organization and found that of the three anatomic sites, tumor

regions from the oral cavity contained significantly different average mixing scores than

tumor regions from the larynx (p=0.016, **Figure 3c**). No significant differences were

found in average mixing scores between the oral cavity and the oropharynx or the

larynx and the oropharynx. Regions from larynx tumors did exhibit a greater range of

mixing scores than oral cavity or oropharynx (**Figure 3c**) indicating that larynx tumors

exhibit greater spatial heterogeneity in terms of neoplastic tumor-immune cell proximity.

We found no significant difference in the mixing scores of primary versus recurrent

tumors (Wilcoxon signed rank test, p=0.441).

Spatial compartmentalization associated with longer progression free survival

To investigate how spatial mixing correlated with patient outcome, we averaged the

mixing scores across regions and assigned a final mixing score and spatial label for

each tumor. Patients with more compartmentalization between neoplastic cells and

immune cells in their primary tumors exhibited significantly longer PFS than those with

greater mixing between these cell types (p=0.032, **Figure 3d**). We then examined how

the average mixing score of the tumors related to the TNM stage and anatomic site of

each primary tumor. Four of the five mixed primary tumors were TNM stage 4, and one

was TNM stage 1. The four compartmentalized primary tumors were TNM stages 1, 2,

3, and 4. All anatomic sites were present in both mixed and compartmentalized spatial

architecture groups (mixed: 2 oral cavity, 1 oropharynx, 2 larynx; compartmentalized: 2

oral cavity, 1 oropharynx, 1 larynx), indicating no single anatomic site had

predominantly mixed or compartmentalized spatial architecture.

Spatial architecture associated with cellular composition

We next explored how cellular composition related to spatial organization, in an effort to

explain the association found between mixing score and PFS. By coloring the initial

PCA on TiME composition by mixing score, we found that tumor regions from the same

mixing group clustered together (**Figure 3e**). Due to this association between cellular

composition and spatial organization, we wondered whether certain cell types would be

more frequent in mixed or compartmentalized tumors. Given how the mixing score is

calculated, we hypothesized that compartmentalized tumor regions would have greater

densities of immune cells than mixed regions, while mixed tumor regions would have

greater densities of neoplastic tumor cells than compartmentalized regions, and found

this to be the case (**Figure 3f**). Namely, compartmentalized tumor regions had greater

densities of CD4$^+$ T helper cells (p=0.016), B cells (p=0.048), antigen presenting cells

(APCs) (p=0.037), and other CD45$^+$ immune cells (p=0.048) than mixed tumor regions.

Given the role of CD4$^+$ T helper cells, B cells, and other MHCII$^+$ immune cells in antigen

presentation, these results could indicate enhanced antigen presentation in

compartmentalized tumor regions as compared to mixed regions. Conversely, mixed

tumor regions contained greater densities of neoplastic tumor cells than

compartmentalized regions (p<0.001).

We further examined associations between functional phenotypes of leukocytes and

mixing scores for each tumor region. Mixed regions contained a greater proportion of

lymphocytes, including CD8$^+$ T cells (p=0.023), CD4$^+$ T helper cells (p=0.017), and B

cells (p=0.016), expressing the immunoregulatory protein programmed death ligand

(PD)-1 than compartmentalized regions (**Figure 3g**). PD-1 is recognized as an indicator

of T cell antigen experience, whereas its expression on B cells has been reported to

suppress T cell effector function[38,39], indicating a suppressive and potentially

dysfunctional immune environment in mixed tumors. On the contrary,

compartmentalized regions contained a greater proportion of APCs expressing the

proliferation marker Ki-67 than mixed regions (p=0.032, **Figure 3h**), supporting the

notion that antigen presentation is a key feature of compartmentalized regions. We

performed a bootstrapping analysis to confirm the robustness of these results and

demonstrate that no one tumor was biasing the functional marker results

(**Supplemental Figure 2b,c,d,e**).

In addition to identifying differences between spatial architectures and their respective

immune and neoplastic tumor cell densities and functional marker expressions, we

found that tumor regions with more compartmentalization between neoplastic cells and

immune cells also contained greater densities of αSMA+ mesenchymal cells, as

compared to those with higher mixing (p=0.002, **Figure 3f**). This result was intriguing

because these cells were not included when computing the mixing score, yet there is a

clear association between αSMA+ cell density and the tumor's spatial organization

(**Figure 3i**).

αSMA+ mesenchymal spatial cellular neighborhoods reveal spatial landscapes

associated with progression free survival advantage

Given the relationship between mixing score and αSMA+ mesenchymal cell density, we

deployed a cellular neighborhood clustering analysis to identify which cell types were

spatially proximal to αSMA+ cells across tumors in order to gain a better understanding

of whether these cells were contributing to the neoplastic tumor-immune spatial

compartmentalization observed. This analysis first involved calculating neighborhoods,

which were defined as physical groupings of cells within a set distance threshold from a

seed cell (**Figure 4a**). Each cell within the distance threshold was deemed a neighbor of

the seed cell, contributing to that seed cell's neighborhood's composition. After

identifying neighborhoods for each seed cell present across all tumor regions,

neighborhoods were grouped with K-means clustering, using the fraction of each cell

type present in the neighborhoods as the clustering features. This revealed clusters of

αSMA⁺ cell neighborhoods with similar cellular makeups across all tumor regions.

We applied a neighborhood clustering analysis with αSMA⁺ cells as the seed cells with

a distance threshold of 30 μm, as this produced neighborhoods with an average of

approximately ten neighbor cells. A recent study involving cellular neighborhood

analyses employed a method that selected the ten nearest spatial neighbors of the seed

cell, regardless of the distance between the seed cell and its neighbors[7]. By setting a

distance threshold of 30 μm, our method required the cells be close, if not directly

touching, while still capturing enough neighbors to cluster on. Any αSMA⁺ cell that did

not have any neighbors was removed from downstream clustering analyses. Clustering

results yielded seven groups, each different in their average composition of αSMA⁺

neighborhoods (**Figure 4b**). Clusters 1 and 2 contained mostly other αSMA⁺ cells

comprising the neighborhoods; in fact, cluster 1 was almost exclusively made up of

αSMA⁺ cells. Cluster 3 contained the greatest proportion of neoplastic tumor cell

neighbors. Clusters 4, 5, 6, and 7 were all comprised of roughly 75% immune cells as

neighbors, although they differed in the types of immune cells present. Cluster 4 was

defined by a majority of other CD45⁺ immune cells not explicitly defined within our

gating strategy. To elucidate marker expression within these other CD45⁺ cells, we

performed a post-hoc t-distributed stochastic neighbor embedding (t-SNE) analysis on

cells classified as 'other immune,' and found them to contain a large population of

CD163⁺ cells and a smaller population of mast cells (tryptase⁺) (**Supplemental Figure**

3a). Cluster 5 contained primarily CD8⁺ T cells as the dominant immune population.

Cluster 6 consisted of CD4$^+$ T helper cells, B cells, and the greatest proportion of APCs

of any cluster. Finally, cluster 7 was defined by its large proportion of granulocytes.

We confirmed that no single tumor region, entire tumor, or patient dominated any of the

αSMA$^+$ neighborhood clusters by examining the percent contribution of each of the

seven clusters per 47 tumor regions, eighteen tumors, and nine patients. While clusters

were present in varying degrees across tumor regions, our results indicated that no

region (**Supplemental Figure 3b**), tumor (**Supplemental Figure 3c**), or patient

(**Supplemental Figure 3d**) was solely responsible for giving rise to any of the clusters.

We also confirmed that all seven clusters were present in tumors collected from each of

the three anatomic sites (**Supplemental Figure 3e**), as well as each of the four TNM

stages (**Supplemental Figure 3f**).

To identify groups of patients with primary tumors of similar αSMA$^+$ cellular

neighborhoods, we performed unsupervised hierarchical clustering on the normalized

average αSMA$^+$ cellular neighborhood composition across the nine primary tumors. This

resulted in two groups of patients, differing in proportional compositions of αSMA$^+$

cellular neighborhoods (**Figure 4c**). On average, both groups had roughly 20% of their

αSMA$^+$ cells assigned to cluster 2 and roughly 10% of their αSMA$^+$ cells assigned to

cluster 5 (**Figure 4d**). However, the two groups differed in that group 1 (blue) included

patients with αSMA$^+$ cells predominantly assigned to clusters 1 and 6, meaning their

αSMA$^+$ cells were primarily surrounded by CD4$^+$ T helper cells, B cells, and other

αSMA$^+$ cells. On the contrary, group 2 (orange) included patients with more of their

αSMA⁺ cells assigned to clusters 3, 4, and 7, meaning their αSMA⁺ cells were mostly

surrounded by neoplastic tumor cells, other immune cells, and granulocytes. Consistent

with our results, visualization of tissue regions illustrates primary tumors in group 1

containing more αSMA⁺ stromal cells overall, frequently neighboring CD4⁺ T helper cells

and B cells, with greater MHCII positivity (**Figure 4e**). Primary tumors in group 2

contained less structured αSMA⁺ stromal cells, fewer neighboring immune cells, and

less MHCII positivity, differences that were strikingly apparent in the mIHC stained

tissue images (**Figure 4e**). Despite both groups containing nearly equal proportions of

CD8⁺ T cells neighboring αSMA⁺ cells, group 1 contained higher densities of CD4⁺ T

cells and increased MHCII positivity as compared to group 2 (**Figure 4e**).

To determine if the composition of αSMA⁺ cellular neighborhood groups was correlated

with clinical outcome, we performed a survival analysis on the two groups of patients.

We found that patients in group 1 had significantly longer PFS than patients in group 2

(p=0.047, **Figure 4f**). Patients in group 1 had primary tumors annotated as TNM stages

1, 2, 3, and 4, while patients in group 2 had four primary tumors annotated as TNM

stage 4 and one annotated as TNM stage 1. Tumors from all three anatomic sites were

represented in both groups. An analysis of the proportions of αSMA⁺ neighborhood

clusters present in each of the nine primary tumors revealed the varying degrees to

which each of the seven clusters were present in each of the tumors (**Figure 4g**).

Notably, we found there to be a positive correlation between the presence of clusters 1

and 6 (r = +0.69) as well as clusters 4 and 7 (r = +0.66). We found negative correlations

between the presence of clusters 1 and 4 (r = -0.88), clusters 3 and 6 (r = -0.69), and

clusters 6 and 7 (r = -0.65). Finally, we found the two groups resulting from hierarchical

clustering to be associated with mixing status. Specifically, group 1 consisted of 75%

compartmentalized tumors, and group 2 consisted of 80% mixed tumors. Overall, these

results describe interesting spatial relationships between immune cells, mesenchymal

stroma, and neoplastic tumor cells, indicating increased antigen presentation and

immune activity associated with compartmentalization and progression free survival.

DISCUSSION

The significant role that TiME cellular composition plays in tumor progression and

response to therapy has been accepted for over a decade[40]. However, recent findings

powered by single-cell proteomics imaging technologies have found that the spatial

organization of the cells present in the TiME also plays a critical role[4-13]. TiME spatial

quantifications are just beginning to provide novel insights into tumor biology, and thus it

is still unclear exactly which spatial features are important in dictating response to

therapy or clinical outcome, as well as whether these features are shared across cancer

types, and how they could be leveraged for therapeutic decisions and patient

stratification. Here, we leveraged single-cell spatial proteomics data generated by our

mIHC immunoassay-based imaging platform to quantitatively assess the TiME of nine

matched primary and recurrent HPV(-) HNSCCs in order to demonstrate the use of

spatial features in disease prognosis. Our results provide novel insights into the

heterogeneity and spatial landscape of HPV(-) HNSCCs, and highlight possible TiME

spatial landscapes that may impact clinical outcome across cancer types.

We found concordance between the results from our heterogeneity and composition

analyses and those from several other studies. Our results are similar to a study that

clustered HNSCC biopsies based on their neoplastic and immune gene signatures as

determined by RNA-sequencing, and reported that samples from the same patient were

more similar to each other than samples from different patients[41]. This trend has also

been observed across cancer types, including melanoma[42], hepatocellular carcinoma[43],

pancreatic ductal adenocarcinoma[44], and breast cancer[45]. Moreover, we found that

patients who experienced an increase in CD8$^+$ T cell density from their primary to

recurrent tumors were associated with improved PFS, which has previously been

reported in head and neck cancer[34,35]. Conversely, patients experiencing the greatest

decrease in CD8$^+$ T cell density in recurrence had TNM stage 4 primary tumors. This

could indicate that later staged primary tumors are better equipped to evade immune

attack in recurrence, and a therapy to elicit an anti-tumor immune response may be

beneficial.

Despite the fact that HPV(-) HNSCCs often contain limited immune infiltrates[22], our

analyses revealed significant differences in the immune cell spatial organization within

these tumors that were associated with progression free survival, highlighting the

importance of considering the spatial context of the TiME. Most strikingly, patients

whose primary tumors contained more compartmentalization between their neoplastic

tumor cell and immune cell populations demonstrated longer PFS. This correlation was

also identified in a similar analysis of triple negative breast cancer patients[4], indicating

that a compartmentalized spatial architecture may play a favorable role in survival

across cancer types. Our analyses of TiME composition and functional marker

expression indicate there is likely more antigen presentation and less immunoregulation

in regions of compartmentalization rather than in regions of mixing. This favorable

immune landscape in compartmentalized tumors may contribute to improved survival.

On the contrary, four of the five of the mixed primary tumors were classified as TNM

stage 4, which may contribute to shortened survival for these patients. However, given

that compartmentalized tumors were not all classified as an early TNM stage and

instead ranged from TNM stage 1 to 4, it is difficult to determine the exact association

between neoplastic tumor and immune cell mixing versus TNM stage, and further

investigation is warranted to understand these correlations and impact on prognosis.

Prior studies have identified a mesenchymal HNSCC subtype[46,47]; our αSMA[+]

mesenchymal cellular neighborhood analyses highlight the importance of considering

how these cells are organized within the TiME, beyond simply considering their

presence in the tumor. This could reveal more precise mesenchymal subtypes for

improved stratification for patient care. αSMA is a common marker for cancer-

associated fibroblasts (CAFs)[20,48], whose presence in tumors, including HNSCC, tends

to be associated with tumor progression, metastasis, and resistance to therapy[49-54].

However, despite compartmentalized tumors having increased αSMA[+] cell density,

patients with compartmentalized primary tumors demonstrated longer PFS, which

indicates that the spatial organization of αSMA[+] cells may be related to their function.

We found αSMA[+] cells neighboring immune cells in both mixed and compartmentalized

primary tumors, but the types of immune cells near αSMA[+] cells differed. This is

interesting, as emerging research has found that CAFs, which are often defined by their

expression of αSMA, can modulate immune cell function within the TiME. With our

mIHC platform we were able to identify the differences in the types of immune cells

neighboring αSMA[+] cells and relate these differences to survival, supporting recent

research on the impact of CAFs on various immune cell populations. A recent study in

melanoma found CAFs to be instrumental in aiding tertiary lymphoid structure (TLS)

development[55]; TLS are defined by their large concentration of B cells surrounded by T

cells, and their presence in tumors is associated with improved patient outcome[3,56,57]. In

compartmentalized tumors, we noticed the αSMA[+] cells to be more structured and

located neighboring dense pockets of T cells and B cells, perhaps indicating the

formation of TLS. Although this antibody panels did not include a biomarker for high

endothelial venules to confirm presence of TLS, our results support the notion that CAF-

lymphocyte interactions can be clinically beneficial.

Conversely, patients with predominantly mixed primary tumors and shorter PFS

contained αSMA[+] cells primarily located near granulocytes and other CD45[+] immune

cells, many of which were likely CD163[+] myelomonocytic cells or mast cells. CD163 is a

marker for scavenger receptor activity and is commonly used to demark pro-tumor type

tumor-associated macrophages and monocytes. Supporting these findings, a study in

oral squamous cell carcinoma found the presence of CAFs to be associated with

increased presence of CD163[+] macrophages and worse survival[58]. Another study

reported CAFs to be correlated with an increase in monocyte expression of CD163,

which in turn suppressed T cell proliferation and increased neoplastic cell proliferation in

breast cancer[59]. Mast cells mediate innate and acquired immune response as a part of

the myeloid lineage. They have been reported to facilitate neo-vascularization and

tumor dissemination in HNSCC, and found to be correlated with increased angiogenesis

in advanced HNSCC[60]. However, interactions between mast cells and CAFs is largely

unknown; our results indicate this interaction may be associated with increased

neoplastic density and worse PFS. Of note, a recent study in melanoma and pancreatic

adenocarcinoma found neutrophils, a subclass of granulocytes, to exert pro-tumor

effects when in the presence of CAFs[61], and another study found the combined

presence of neutrophils and CAFs to be associated with shortened survival in gastric

adenocarcinoma[62].

Finally, patients with predominantly mixed primary tumors and shorter PFS were found

to contain fewer overall stromal cells and greater proportions of those αSMA$^+$ cells

neighboring neoplastic tumor cells. It has been reported that neoplastic tumor cells in

direct contact with CAFs move along tracks laid by the CAFs in the extracellular matrix,

promoting tumor growth[63,64]. It is possible that αSMA$^+$ cells in mixed tumors may provide

avenues for neoplastic tumor cells to transit, thus leading to more advanced tumor

progression. Overall, spatial analyses herein deepened our understanding of neoplastic

tumor and immune cell organization relative to each other, and how this organization is

related to αSMA$^+$ cells working in tandem with many cells, beyond single cell-cell

interactions, to impact TiME organization, and ultimately, clinical outcome. The

conclusions from our spatial analyses and their relationship to clinical variables are

summarized in **Figure 5**.

The concordance found among results herein and those of the aforementioned studies

provides orthogonal support for our conclusions and indicates that single-cell

heterogeneity and spatial organization of tumors may share similarities across different

types of cancer and across different molecular assays. Our spatial analyses

demonstrate the use of various algorithms to quantify the spatial landscape of tumors

using single-cell imaging data. Further, our computational methods provide a framework

for future single-cell imaging analyses, as they are applicable to any multiplex imaging

assay, including mIHC, co-detection by indexing technology (CODEX)[7], cyclic

immunofluorescence[18,65,66], and imaging mass cytometry[6]. Our results highlight several

spatial architectures that may help guide precision medicine approaches for HPV(-)

HNSCC patients, including architectures that may help stratify patients who may have

shorter PFS, and thus warrant more aggressive therapy or clinical follow-ups. While our

results provide evidence that the spatial organization of HPV(-) HNSCC tumors

correlates with clinical outcome, future studies with larger cohorts will be needed to

evaluate the strength and validity of our observations. Studies with greater

representation of anatomic site and stage are also needed to assess the prognostic

value of the spatial features identified in this HNSCC cohort. Despite these limitations,

this study demonstrates practical analysis strategies that elucidate spatial architecture

features for potential use in precision medicine.

METHODS

Multiplex immunohistochemistry data generation

mIHC is an immunohistochemical-based imaging platform that evaluates sequentially

stained immune lineage epitope-specific antibodies for immunodetection on FFPE

tissue sections[16,17]. Images were stained and processed as described in our previous

report[17]. A table of antibodies, species, vendor, and concentration used in staining are

previously reported in Table 1 of Banik et al[17]. Briefly, sequentially stained images were

co-registered in MATLAB. AEC signal from each antibody stain was extracted and

normalized, and the mean intensity of each single cell for each marker was quantified in

Cell Profiler. Watershed based nuclei segmentation on hematoxylin staining was used

to identify single cells in FIJI. Using a hierarchical gating strategy, single cells were

phenotyped using image gating cytometry in FCS Express 7 Image Cytometry RUO

(**Supplemental Figure 1a**). A threshold was set on the scatterplot of mean intensity for

each marker within the gating strategy, validated by visual live rendering of masked cell

objects within the selected gate on extracted marker signals. Cartesian coordinates of

each phenotyped cell were maintained relative to the tissue region.

We applied the mIHC pipeline to analyze matched primary and recurrent FFPE tissue

specimens from nine HPV(-) HNSCC patients. Each patient underwent surgical

resection of their primary tumor prior to beginning a regimen of chemotherapy and

radiation therapy. Upon recurrence, the patient underwent another surgical resection of

their recurrent tumor. Tissue specimens for each patient were obtained from the Oregon

Health & Science University Knight Biolibrary and were deidentified and coded with a

unique identifier prior to analysis. Patient demographic and clinical data including HPV

status, tobacco and alcohol use, treatment regimens, and survival outcomes were

collected. All HNSCC tumors were staged according to the 8th edition AJCC/UIC TNM

classification and cohort characteristics are shown in **Table 1** as reported in Banik et

al[17]. All studies involving human tissue were approved by institutional IRB (protocol

#809 and #3609), and written informed consent was obtained.

Tumor heterogeneity analyses

The Kullback-Leibler divergence was calculated using the entropy function from the

Scipy Python package[67]. The distribution of each individual tumor region's cellular

composition was compared to the average of five different cellular composition

distributions: the average distribution for the region's tumor, the average distribution for

the patient's primary and recurrent tumors, the average distribution across all tumors of

the same timepoint in the cohort (primary or recurrent), the average distribution across

all tumors resected from the same anatomic site in the cohort, and the average

distribution across all tumors from all patients in the cohort (primary and recurrent

combined). Log base 2 was used for the calculation. Unsupervised hierarchical

clustering of each tumor region was performed using the log10+1 normalized density of

each cell type present as the features. Euclidean distance was used to determine

distances between observations, and the Ward method was used for the linkage.

TiME compositional change clustering analysis

Unsupervised hierarchical clustering of each patient was performed using the

normalized change in density of each cell type as the features. Euclidean distance was

used to determine distances between observations, and the Ward method was used for

the linkage. Normalization in the change in density was computing by first calculating

the absolute value of the raw change in density for each cell type. These values were

then normalized to a range of [0,1]. Finally, the values that were originally negative

(decreasing change), were flipped to be negative values again, such that all values

ranged from [-1,1] with zero representing no change.

Mixing score analysis

Keren et al. developed a mixing score to quantify the ratio of neoplastic and immune cell

spatial interactions[4]. This score is defined as the number of interactions between

neoplastic tumor cells and immune cells divided by the number of interactions between

immune cells and another immune cell within a tumor region. We defined there to be an

interaction between two cells if their centers [the (x,y) coordinates provided by mIHC]

were within 15 µm from one another. We used the median mixing score value (0.107)

for all primary tumors as the threshold to distinguish between mixed and

compartmentalized spatial organization groups. Tumor regions with a mixing score of

greater than 0.107 were defined as mixed. Tumor regions with a mixing score of less

than 0.107 were defined as compartmentalized. We used a density threshold of less

than 250 immune cells per 800^2 $µm^2$ to define tumor regions as cold. We chose this

threshold to match that used by Keren et al.

Functional marker bootstrapping analyses

Bootstrapping analyses involved the following steps. First, one tumor region per

eighteen tumor samples was randomly selected. The regions were then split into two

groups based on their mixed or compartmentalized spatial architecture designation, and

the average proportion of the specified cell population expressing the specified

functional marker was calculated for each group. Finally, this process was repeated 100

times, yielding 100 values representing the proportion of the specified cell population

expressing the specified functional marker for mixed tumor regions and 100 values

representing the proportion of the specified cell population expressing the specified

functional marker for compartmentalized tumor regions. These values were then

compared between the mixed and compartmentalized groups for differences.

Cellular neighborhood clustering analyses

Cellular neighborhoods were defined by drawing a circle of a specified radius around all

seed cells of a given phenotype. Cells whose centers were inside the circle were

considered neighbors of that seed cell and contributed to that cell's neighborhood. All

neighborhoods across all tumor regions were then clustered using scikit-learn's

MiniBatchKMeans function[68] to perform K-means clustering according to their

normalized cellular composition. The elbow method was used to determine the number

of clusters to form. Each resulting cluster was comprised of neighborhoods with a

similar cellular makeup. Unsupervised hierarchical clustering of each primary tumor was

performed using the log10+1 normalized proportion (out of 100) of αSMA+ cell

neighborhood clusters present in the tumor as the features. Euclidean distance was

used to determine distances between observations, and the Ward method was used for

the linkage.

Statistics

Independent t-tests were used to determine statistically significant differences for

independent samples whose differences followed a normal distribution. Mann-Whitney

U tests were used to determine statistically significant differences for independent

samples whose differences did not follow a normal distribution. Paired t-tests were used

to determine statistically significant differences for paired samples whose differences

followed a normal distribution. Wilcoxon signed rank tests were used to determine

statistically significant differences for paired samples whose differences did not follow a

normal distribution. One-way ANOVA tests were used to determine statistically

significant differences for multi-group comparisons. If the ANOVA result was significant,

a Tukey honestly significant difference post-hoc test was conducted to determine which

groups were significantly different from one another. A Benjamini-Hochberg correction

was used to account for multiple hypothesis testing in analyses that involved

systematically testing multiple variables. P-values less than 0.05 were considered

statistically significant. All statistical calculations were performed with the Scipy and

statsmodels packages using Python software[67,69].

Survival analyses

Kaplan-Meier curves were generated and a log-rank test was performed using the

lifelines package with Python software[70]. P-values less than 0.05 were considered

statistically significant.

Data and software availability

All of the data produced by our mIHC computational image processing pipeline,

including protein abundance, cell phenotype, and cell location information saved in the

form of a matrix, in addition to survival data, is available for download at Zenodo

(https://doi.org/10.5281/zenodo.5540356). All computational analyses in this study were

performed using Python software, version 3.6.5. The code created to produce the

results of this study is available at https://github.com/kblise/HNSCC_mIHC_paper.

ACKNOWLEDGEMENTS

The authors thank Dr. Courtney Betts, Justin Tibbitts, Teresa Beechwood, and Meghan

Lavoie for regulatory and technical assistance. The authors thank the OHSU Knight

Biostatistics Shared Resource for guidance with statistical calculations. The authors

also acknowledge and thank the patients who donated tissue samples for this study.

REFERENCES

Binnewies, M. *et al.* Understanding the tumor immune microenvironment (TIME) for effective therapy. *Nat Med* **24**, 541-550 doi:10.1038/s41591-018-0014-x (2018).

Junttila, M. R. & de Sauvage, F. J. Influence of tumour micro-environment heterogeneity on therapeutic response. *Nature* **501**, 346-354 doi:10.1038/nature12626 (2013).

Petitprez, F. *et al.* B cells are associated with survival and immunotherapy response in sarcoma. *Nature* **577**, 556-560 doi:10.1038/s41586-019-1906-8 (2020).

Keren, L. *et al.* A Structured Tumor-Immune Microenvironment in Triple Negative Breast Cancer Revealed by Multiplexed Ion Beam Imaging. *Cell* **174**, 1373-1387 e1319 doi:10.1016/j.cell.2018.08.039 (2018).

Ali, H. R. *et al.* Imaging mass cytometry and multiplatform genomics define the phenogenomic landscape of breast cancer. *Nature Cancer* **1**, 163-175 doi:10.1038/s43018-020-0026-6 (2020).

Jackson, H. W. *et al.* The single-cell pathology landscape of breast cancer. *Nature* **578**, 615-620 doi:10.1038/s41586-019-1876-x (2020).

Schurch, C. M. *et al.* Coordinated Cellular Neighborhoods Orchestrate Antitumoral Immunity at the Colorectal Cancer Invasive Front. *Cell* doi:10.1016/j.cell.2020.07.005 (2020).

Farkkila, A. *et al.* Immunogenomic profiling determines responses to combined PARP and PD-1 inhibition in ovarian cancer. *Nat Commun* **11**, 1459 doi:10.1038/s41467-020-15315-8 (2020).

Barua, S. *et al.* Spatial interaction of tumor cells and regulatory T cells correlates with survival in non-small cell lung cancer. *Lung Cancer* **117**, 73-79 doi:10.1016/j.lungcan.2018.01.022 (2018).

Gide, T. N. *et al.* Close proximity of immune and tumor cells underlies response to anti-PD-1 based therapies in metastatic melanoma patients. *Oncoimmunology* **9**, 1659093 doi:10.1080/2162402X.2019.1659093 (2020).

Giraldo, N. A. *et al.* Multidimensional, quantitative assessment of PD-1/PD-L1 expression in patients with Merkel cell carcinoma and association with response to pembrolizumab. *J Immunother Cancer* **6**, 99 doi:10.1186/s40425-018-0404-0 (2018).

Johnson, D. B. *et al.* Quantitative Spatial Profiling of PD-1/PD-L1 Interaction and HLA-DR/IDO-1 Predicts Improved Outcomes of Anti-PD-1 Therapies in Metastatic Melanoma. *Clin Cancer Res* **24**, 5250-5260 doi:10.1158/1078-0432.CCR-18-0309 (2018).

Enfield, K. S. S. *et al.* Hyperspectral cell sociology reveals spatial tumor-immune cell interactions associated with lung cancer recurrence. *J Immunother Cancer* **7**, 13 doi:10.1186/s40425-018-0488-6 (2019).

Galon, J. *et al.* Type, density, and location of immune cells within human colorectal tumors predict clinical outcome. *Science* **313**, 1960-1964 doi:10.1126/science.1129139 (2006).

15   Galon, J. *et al.* Cancer classification using the Immunoscore: a worldwide task
force. *J. Transl. Med.* **10**, 205 doi:10.1186/1479-5876-10-205 (2012).

16   Tsujikawa, T. *et al.* Quantitative Multiplex Immunohistochemistry Reveals
Myeloid-Inflamed Tumor-Immune Complexity Associated with Poor Prognosis.
Cell Rep **19**, 203-217 doi:10.1016/j.celrep.2017.03.037 (2017).

17   Banik, G. *et al.* High-dimensional multiplexed immunohistochemical
characterization of immune contexture in human cancers. *Methods Enzymol* **635**,
1-20 doi:10.1016/bs.mie.2019.05.039 (2020).

18   Eng, J. *et al.* Cyclic Multiplexed-Immunofluorescence (cmIF), a Highly
Multiplexed Method for Single-Cell Analysis. *Methods Mol. Biol.* **2055**, 521-562
doi:10.1007/978-1-4939-9773-2_24 (2020).

19   Brucher, B. L. & Jamall, I. S. Cell-cell communication in the tumor
microenvironment, carcinogenesis, and anticancer treatment. *Cell. Physiol.*
Biochem. **34**, 213-243 doi:10.1159/000362978 (2014).

20   Alsahafi, E. *et al.* Clinical update on head and neck cancer: molecular biology
and ongoing challenges. *Cell Death Dis.* **10**, 540 doi:10.1038/s41419-019-1769-9
(2019).

21   Cohen, E. E. W. *et al.* The Society for Immunotherapy of Cancer consensus
statement on immunotherapy for the treatment of squamous cell carcinoma of
the head and neck (HNSCC). *J Immunother Cancer* **7**, 184 doi:10.1186/s40425-
019-0662-5 (2019).

22   Canning, M. *et al.* Heterogeneity of the Head and Neck Squamous Cell
Carcinoma Immune Landscape and Its Impact on Immunotherapy. *Front Cell*
Dev Biol **7**, 52 doi:10.3389/fcell.2019.00052 (2019).

23   Fakhry, C. *et al.* Improved survival of patients with human papillomavirus-positive
head and neck squamous cell carcinoma in a prospective clinical trial. *J. Natl.*
Cancer Inst. **100**, 261-269 doi:10.1093/jnci/djn011 (2008).

24   Ang, K. K. *et al.* Human papillomavirus and survival of patients with
oropharyngeal cancer. *N. Engl. J. Med.* **363**, 24-35 doi:10.1056/NEJMoa0912217
(2010).

25   Predina, J. *et al.* Changes in the local tumor microenvironment in recurrent
cancers may explain the failure of vaccines after surgery. *Proc. Natl. Acad. Sci.*
U. S. A. **110**, E415-424 doi:10.1073/pnas.1211850110 (2013).

26   Zheng, B. *et al.* Changes in the tumor immune microenvironment in resected
recurrent soft tissue sarcomas. *Ann Transl Med* **7**, 387
doi:10.21037/atm.2019.07.43 (2019).

27   Walens, A. *et al.* Adaptation and selection shape clonal evolution of tumors
during residual disease and recurrence. *Nat Commun* **11**, 5017
doi:10.1038/s41467-020-18730-z (2020).

28   Watermann, C. *et al.* Recurrent HNSCC Harbor an Immunosuppressive Tumor
Immune Microenvironment Suggesting Successful Tumor Immune Evasion. *Clin*
Cancer Res **27**, 632-644 doi:10.1158/1078-0432.CCR-20-0197 (2021).

29   Hartman, D. J., Ahmad, F., Ferris, R. L., Rimm, D. L. & Pantanowitz, L. Utility of
CD8 score by automated quantitative image analysis in head and neck
squamous cell carcinoma. *Oral Oncol.* **86**, 278-287
doi:10.1016/j.oraloncology.2018.10.005 (2018).

30   Balermpas, P. *et al.* CD8+ tumour-infiltrating lymphocytes in relation to HPV
status and clinical outcome in patients with head and neck cancer after
postoperative chemoradiotherapy: A multicentre study of the German cancer
consortium radiation oncology group (DKTK-ROG). *Int. J. Cancer* **138**, 171-181
doi:10.1002/ijc.29683 (2016).
31   Nasman, A. *et al.* Tumor infiltrating CD8+ and Foxp3+ lymphocytes correlate to
clinical outcome and human papillomavirus (HPV) status in tonsillar cancer.
PLoS One **7**, e38711 doi:10.1371/journal.pone.0038711 (2012).
32   Nordfors, C. *et al.* CD8+ and CD4+ tumour infiltrating lymphocytes in relation to
human papillomavirus status and clinical outcome in tonsillar and base of tongue
squamous cell carcinoma. *Eur. J. Cancer* **49**, 2522-2530
doi:10.1016/j.ejca.2013.03.019 (2013).
33   Balermpas, P. *et al.* Tumour-infiltrating lymphocytes predict response to definitive
chemoradiotherapy in head and neck cancer. *Br. J. Cancer* **110**, 501-509
doi:10.1038/bjc.2013.640 (2014).
34   So, Y. K. *et al.* An increase of CD8(+) T cell infiltration following recurrence is a
good prognosticator in HNSCC. *Sci. Rep.* **10**, 20059 doi:10.1038/s41598-020-
77036-8 (2020).
35   Schutt, C. R. *et al.* Genomic and neoantigen evolution from primary tumor to first
metastases in head and neck squamous cell carcinoma. *Oncotarget* **12**, 534-548
doi:10.18632/oncotarget.27907 (2021).
36   Russell, S. *et al.* Immune cell infiltration patterns and survival in head and neck
squamous cell carcinoma. *Head Neck Oncol.* **5**, 24 (2013).
37   Wolf, G. T. *et al.* Tumor infiltrating lymphocytes (TIL) and prognosis in oral cavity
squamous carcinoma: a preliminary study. *Oral Oncol.* **51**, 90-95
doi:10.1016/j.oraloncology.2014.09.006 (2015).
38   Wang, X. *et al.* PD-1-expressing B cells suppress CD4(+) and CD8(+) T cells via
PD-1/PD-L1-dependent pathway. *Mol. Immunol.* **109**, 20-26
doi:10.1016/j.molimm.2019.02.009 (2019).
39   Thibult, M. L. *et al.* PD-1 is a novel regulator of human B-cell activation. *Int.*
Immunol. **25**, 129-137 doi:10.1093/intimm/dxs098 (2013).
40   Fridman, W. H., Pages, F., Sautes-Fridman, C. & Galon, J. The immune
contexture in human tumours: impact on clinical outcome. *Nat. Rev. Cancer* **12**,
298-306 doi:10.1038/nrc3245 (2012).
41   Wood, O. *et al.* Head and Neck Squamous Cell Carcinomas Are Characterized
by a Stable Immune Signature Within the Primary Tumor Over Time and Space.
Clin Cancer Res **23**, 7641-7649 doi:10.1158/1078-0432.CCR-17-0373 (2017).
42   Reuben, A. *et al.* Genomic and immune heterogeneity are associated with
differential responses to therapy in melanoma. *NPJ Genom Med* **2**
doi:10.1038/s41525-017-0013-8 (2017).
43   Zhang, Q. *et al.* Integrated multiomic analysis reveals comprehensive tumour
heterogeneity and novel immunophenotypic classification in hepatocellular
carcinomas. *Gut* **68**, 2019-2031 doi:10.1136/gutjnl-2019-318912 (2019).
44   Lin, W. *et al.* Single-cell transcriptome analysis of tumor and stromal
compartments of pancreatic ductal adenocarcinoma primary tumors and

metastatic lesions. *Genome Med.* **12**, 80 doi:10.1186/s13073-020-00776-9
(2020).
45 Gyanchandani, R. *et al.* Intratumor Heterogeneity Affects Gene Expression
Profile Test Prognostic Risk Stratification in Early Breast Cancer. *Clin Cancer*
Res **22**, 5362-5369 doi:10.1158/1078-0432.CCR-15-2889 (2016).
46 Chung, C. H. *et al.* Molecular classification of head and neck squamous cell
carcinomas using patterns of gene expression. *Cancer Cell* **5**, 489-500
doi:10.1016/s1535-6108(04)00112-6 (2004).
47 Walter, V. *et al.* Molecular subtypes in head and neck cancer exhibit distinct
patterns of chromosomal gain and loss of canonical cancer genes. *PLoS One* **8**,
e56823 doi:10.1371/journal.pone.0056823 (2013).
48 Parajuli, H. *et al.* Integrin alpha11 is overexpressed by tumour stroma of head
and neck squamous cell carcinoma and correlates positively with alpha smooth
muscle actin expression. *J. Oral Pathol. Med.* **46**, 267-275 doi:10.1111/jop.12493
(2017).
49 Zhang, Q. *et al.* Integrated analysis of single-cell RNA-seq and bulk RNA-seq
reveals distinct cancer-associated fibroblasts in head and neck squamous cell
carcinoma. *Ann Transl Med* **9**, 1017 doi:10.21037/atm-21-2767 (2021).
50 Lim, K. P. *et al.* Fibroblast gene expression profile reflects the stage of tumour
progression in oral squamous cell carcinoma. *J. Pathol.* **223**, 459-469
doi:10.1002/path.2841 (2011).
51 Shiga, K. *et al.* Cancer-Associated Fibroblasts: Their Characteristics and Their
Roles in Tumor Growth. *Cancers (Basel)* **7**, 2443-2458
doi:10.3390/cancers7040902 (2015).
52 Tao, L., Huang, G., Song, H., Chen, Y. & Chen, L. Cancer associated fibroblasts:
An essential role in the tumor microenvironment. *Oncol. Lett.* **14**, 2611-2620
doi:10.3892/ol.2017.6497 (2017).
53 Sahai, E. *et al.* A framework for advancing our understanding of cancer-
associated fibroblasts. *Nat. Rev. Cancer* **20**, 174-186 doi:10.1038/s41568-019-
0238-1 (2020).
54 Johansson, A. C. *et al.* Cancer-associated fibroblasts induce matrix
metalloproteinase-mediated cetuximab resistance in head and neck squamous
cell carcinoma cells. *Mol. Cancer Res.* **10**, 1158-1168 doi:10.1158/1541-
7786.MCR-12-0030 (2012).
55 Rodriguez, A. B. *et al.* Immune mechanisms orchestrate tertiary lymphoid
structures in tumors via cancer-associated fibroblasts. *Cell Rep.* **36**, 109422
doi:10.1016/j.celrep.2021.109422 (2021).
56 Cabrita, R. *et al.* Tertiary lymphoid structures improve immunotherapy and
survival in melanoma. *Nature* **577**, 561-565 doi:10.1038/s41586-019-1914-8
(2020).
57 Helmink, B. A. *et al.* B cells and tertiary lymphoid structures promote
immunotherapy response. *Nature* **577**, 549-555 doi:10.1038/s41586-019-1922-8
(2020).
58 Fujii, N. *et al.* Cancer-associated fibroblasts and CD163-positive macrophages in
oral squamous cell carcinoma: their clinicopathological and prognostic

significance. *J. Oral Pathol. Med.* **41**, 444-451 doi:10.1111/j.1600-
0714.2012.01127.x (2012).
59 Gok Yavuz, B. *et al.* Cancer associated fibroblasts sculpt tumour
microenvironment by recruiting monocytes and inducing immunosuppressive PD-
1(+) TAMs. *Sci. Rep.* **9**, 3172 doi:10.1038/s41598-019-39553-z (2019).
60 Markwell, S. M. & Weed, S. A. Tumor and stromal-based contributions to head
and neck squamous cell carcinoma invasion. *Cancers (Basel)* **7**, 382-406
doi:10.3390/cancers7010382 (2015).
61 Munir, H. *et al.* Stromal-driven and Amyloid beta-dependent induction of
neutrophil extracellular traps modulates tumor growth. *Nat Commun* **12**, 683
doi:10.1038/s41467-021-20982-2 (2021).
62 Cong, X. *et al.* CD66b(+) neutrophils and alpha-SMA(+) fibroblasts predict clinical
outcomes and benefits from postoperative chemotherapy in gastric
adenocarcinoma. *Cancer Med* **9**, 2761-2773 doi:10.1002/cam4.2939 (2020).
63 Gaggioli, C. *et al.* Fibroblast-led collective invasion of carcinoma cells with
differing roles for RhoGTPases in leading and following cells. *Nat. Cell Biol.* **9**,
1392-1400 doi:10.1038/ncb1658 (2007).
64 Goetz, J. G. *et al.* Biomechanical remodeling of the microenvironment by stromal
caveolin-1 favors tumor invasion and metastasis. *Cell* **146**, 148-163
doi:10.1016/j.cell.2011.05.040 (2011).
65 Lin, J. R., Fallahi-Sichani, M. & Sorger, P. K. Highly multiplexed imaging of single
cells using a high-throughput cyclic immunofluorescence method. *Nat Commun*
6, 8390 doi:10.1038/ncomms9390 (2015).
66 Lin, J. R. *et al.* Highly multiplexed immunofluorescence imaging of human tissues
and tumors using t-CyCIF and conventional optical microscopes. *Elife* **7**
doi:10.7554/eLife.31657 (2018).
67 Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in
Python. *Nat. Methods* **17**, 261-272 doi:10.1038/s41592-019-0686-2 (2020).
68 Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *JMLR* **12**, 2825-
2830 (2011).
69 Seabold, S. & Perktold, J. statsmodels: Econometric and statistical modeling with
python. *Proceedings of the 9th Python in Science Conference* (2010).
70 Davidson-Pilon, C. *et al.* CamDavidsonPilon/lifelines: v0.21.0. *Zenodo*
doi:10.5281/zenodo.805993 (2020).
953

TABLE LEGENDS

Table 1. Patient and tumor characteristics. Adapted from Banik et al[17] to show de-identified patient characteristics.

Table 2. mIHC cell phenotype classification. Antibody panels used to hierarchically phenotype cells into one of ten phenotype classifications. Additional functional markers detected are listed at the bottom.

Table 3. Coefficient of variation. The coefficient of variation was calculated per tumor per cell type and then averaged across all tumors (n=16). Tumors with only one region (n=2) were removed from this analysis. Cell types are ordered from largest coefficient of variation to smallest coefficient of variation.

FIGURE LEGENDS

Figure 1. Heterogeneity across patients and tumor regions. a, Cohort and tissue region selection overview. One to three regions of 2500^2 μm^2 were assayed with mIHC per eighteen tumor resections and are represented by the green boxes in the tissue images. The black line corresponds to 2500 μm for each image. **b**, Density (cells/mm^2) of each cell type present per individual primary (P) and recurrent (R) tumor. Each dot represents a single tumor region (n=47). **c**, Density (cells/mm^2) of each immune cell type present per individual primary (P) and recurrent (R) tumor. Each dot represents a single tumor region (n=47). **d**, Box plot of the Kullback-Leibler divergences from a single tumor region's cellular distribution compared to: the tumor's average cellular distribution

[Intra-Tumor (P or R only)], the patient's average cellular distribution [Intra-Patient (P

and R)], the cohort's average cellular distribution across tumors of the same timepoint

[Inter-Patient (P or R only)], the cohort's average cellular distribution across tumors of

the same anatomic site [Inter-Patient (Same Anatomic Site)], the cohort's average

cellular distribution across all tumors from all patients [Inter-Patient (all)]. P-values

calculated using a one-way ANOVA multi-group significance test followed by a Tukey

honestly significant difference post-hoc test. **e**, Heatmap of cellular composition across

tumor regions. Rows are individual tumor regions that are ordered by the hierarchical

clustering of their cellular composition. Columns are the cell types used as clustering

features. Composition was normalized using a log10+1 transformation before clustering.

Leftmost three columns are color coded by patient, tumor timepoint, and anatomic site.

f, Principal component analysis on cellular density following a log10+1 transformation.

Each point (n=47) represents one tumor region and is colored by patient. The shape of

each point denotes primary or recurrent status. **g**, Principal component analysis on

cellular density following a log10+1 transformation. Each dot (n=47) represents one

tumor region and is colored by the anatomic resection site.

Figure 2. Tumor cellular composition changes following therapy. a, Box plot

showing the average density of each cell type split by primary and recurrent status.

Each dot represents the density of that cell type for one tumor, averaged across regions

(n=9 primary tumors, n=9 recurrent tumors). **b**, Line plots showing the change in cellular

composition from primary to recurrent tumors for each cell type. Each dot represents the

density of that cell type for one tumor, averaged across regions. Lines are colored by

patient. **c**, Heatmap of average change in cellular composition from primary to recurrent

tumors for each patient. Rows are individual patients that are ordered by the

hierarchical clustering of their change in TiME cellular composition (averaged across

regions). Columns are the cell types used as clustering features. Compositional change

was normalized [-1,1] before clustering (see Methods). Leftmost two columns are color

coded by patient followed by therapy administered. **d**, Box plot showing the average

change in density of each cell type for each patient (n=9) colored by short-term or long-

term progression free survival, splitting on median progression free survival.

1008

Figure 3. Mixing score quantifies the spatial organization of tumors. a, mIHC

images of a representative mixed tumor region (top) versus a compartmentalized tumor

region (bottom). Leftmost panel shows tumor regions with neoplastic tumor cells

(purple) and $CD45^+$ immune cells (green). Remaining panels show zoomed in areas of

mixing (top) and compartmentalization (bottom), first with both cell populations present

and then separated out. **b**, Box plot showing the mixing scores across all primary and

recurrent tumors (n=18). Each dot (n=47) represents the mixing score for one tumor

region and is colored according to its spatial architecture. The average spatial

architecture designation for the overall tumor is printed above each box. **c**, Box plot

showing the mixing score of each tumor region split by the anatomic site of its resection.

P-value calculated using a one-way ANOVA multi-group significance test followed by a

Tukey honestly significant difference post-hoc test. **d**, Kaplan-Meier curve of

progression free survival for patients split by the mixing score of their primary tumors.

Patients were split on the median value. P-value calculated using the log-rank test. **e**,

Principal component analysis on cellular density following a log10+1 transformation.

Each dot (n=47) represents one tumor region and is colored according to the region's

spatial architecture. **f**, Box plot showing the density of each cell type split by the tumor

region's spatial architecture. Each dot represents the density of that cell type for one

region (n=47 per cell type). Statistical significance calculated using independent one-

tailed t-tests for cell types whose differences follow a normal distribution and non-

parametric one-tailed t-tests (Mann-Whitney U Test) for cell types whose differences do

not follow a normal distribution. P-values were corrected using the Benjamini-Hochberg

procedure. **g**, PD-1 expression on $CD8^+$ T cells, $CD4^+$ T helper cells, and B cells by

spatial architecture. Each dot represents the percentage of each cell type positive for

PD-1 for a single tumor region (n=45, excluding cold regions). P-values calculated using

a one-tailed Mann-Whitney U test and corrected using the Benjamini-Hochberg

procedure. **h**, Ki-67 expression on APCs by spatial architecture. Each dot represents

the percentage of cells positive for Ki-67 for a single tumor region (n=45, excluding cold

regions). P-values calculated using a Mann-Whitney U test and corrected using the

Benjamini-Hochberg procedure. **i**, Bar chart showing the density of αSMA^+

mesenchymal cells present per tumor region. Bars are ordered by αSMA^+ cell density

and are colored according to the region's spatial architecture.

1041

Figure 4. αSMA⁺ mesenchymal cellular neighborhood clustering. **a**, Cellular

neighborhoods were defined by drawing a circle of a specified radius around each seed

cell (green) of a designated phenotype. Cells whose centers were inside the circle were

considered neighbors of that seed cell. This figure was created using BioRender.com.

b, Stacked bar chart showing the average cellular composition of each αSMA$^+$

mesenchymal cell neighborhood cluster (n=7). Bars are colored by cell type and

represent the average fraction (out of 1.0) of each cell type present in the

neighborhoods belonging to each cluster. **c**, Heatmap of αSMA$^+$ cell neighborhood

clusters present averaged across primary tumors. Rows are primary tumors that are

ordered by the hierarchical clustering of their average of αSMA$^+$ neighborhood cluster

presence. Columns are the αSMA$^+$ cell neighborhood clusters used as clustering

features. Percent (out of 100) of αSMA$^+$ neighborhood clusters was normalized using a

log10+1 transformation before clustering. Leftmost column is color coded by patient. **d**,

Stacked bar chart showing the average proportion (out of 1.0) of αSMA$^+$ cell

neighborhood clusters present in each of the two hierarchically clustered groups of

primary tumors. **e**, Panel of mIHC images containing two merged pseudo-colored

images of one region from 4 patients visually illustrates representative regions from

patients in group 1 and group 2. The top panel of images are merged pseudo-colored

stains containing CD3, CD8, CD20, αSMA, and PANCK. The bottom panel of images

are the same regions as the top panel with merged pseudo-colored stains containing

MHCII, CD66B, Ki-67, αSMA, and PANCK. White arrows point out representative

αSMA$^+$ cells. **f**, Kaplan-Meier curve of progression free survival for patients grouped

together in the hierarchical clustering of their primary tumor αSMA$^+$ mesenchymal cell

neighborhood abundance. P-value calculated using the log-rank test. **g**, Stacked bar

chart showing the average proportion (out of 1.0) of αSMA$^+$ cell neighborhood clusters

present in each of the primary tumors (n=9). Tumors are ordered by descending

proportions of cluster 1.

1069

Figure 5. Proposed model of primary HPV(-) HNSCC tumor-immune

microenvironments. a, Depiction of a tumor with a compartmentalized spatial

architecture. These tumors have decreased mixing between immune cells and

neoplastic tumor cells and tend to contain greater immune cell density, specifically

those involved in antigen presentation, as well as increased density of αSMA$^+$

mesenchymal cells. The αSMA$^+$ cells present tend to be neighbored by CD4$^+$ T helper

cells and B cells, as well as other αSMA$^+$ cells. Compartmentalized primary tumors were

found to be associated with longer progression free survival. **b**, Depiction of a tumor

with a mixed spatial architecture. These tumors contain increased mixing between

immune cells and neoplastic tumor cells and tend to contain increased neoplastic tumor

cells, as well as increased PD-1-positive lymphocytes. The αSMA$^+$ cells present tend to

be neighbored by neoplastic tumor cells, other immune cells, many of which are

CD163$^+$ or are mast cells, and granulocytes. Mixed primary tumors were found to be

associated with shorter progression free survival and later TNM stage. This figure was

created using BioRender.com.

1085

Supplemental Figure 1a. Gating template used to determine cell phenotypes and

classification of ten populations using image gating cytometry in FCS Image Cytometry

7 RUO. Scatter plots show single cell mean intensity distributions rescaled from 0-1 on

a log10 scale with gated populations in colored boxes. Image plot shows a semi-

transparent pseudo-colored mask of each cell object colored by the classified

population overlaid on signal extracted nuclei image. The markers used for identification

of cell phenotypes are shown in Table 1.

1093

Supplemental Figure 1b. mIHC images of six tumor regions selected from two different

tumor resections (patient 9 primary and patient 8 recurrent). Blue pseudo color indicates

all cell nuclei present in the region (Hematoxylin$^+$) and yellow indicates B cells. In both

tumors, two regions have low densities of B cells, whereas one region has a high

density and spatial concentration of B cells, despite the three regions being sampled

from the same tumor.

1100

Supplemental Figure 1c. Box plot of the Kullback-Leibler divergences from a single

tumor region's immune cell distribution compared to: the tumor's average immune cell

distribution [Intra-Tumor (P or R only)], the patient's average immune cell distribution

[Intra-Patient (P and R)], the cohort's average immune cell distribution across tumors of

the same timepoint [Inter-Patient (P or R only)], the cohort's average immune cell

distribution across tumors of the same anatomic site [Inter-Patient (Same Anatomic

Site)], the cohort's average immune cell distribution across all tumors from all patients

[Inter-Patient (all)]. P-values calculated using a one-way ANOVA multi-group

significance test followed by a Tukey honestly significant difference post-hoc test.

1110

Supplemental Figure 2a. Box plot of the absolute difference from a single tumor

region's mixing score compared to: the tumor's average mixing score [Intra-Tumor (P or

R only)], the patient's average mixing score [Intra-Patient (P and R)], the cohort's

average mixing score across tumors of the same timepoint [Inter-Patient (P or R only)],

the cohort's average mixing score across tumors of the same anatomic site [Inter-

Patient (Same Anatomic Site)], the cohort's average mixing score across all tumors

from all patients [Inter-Patient (all)]. P-value calculated using a one-way ANOVA multi-

group significance test.

1119

1120   **Supplemental Figure 2b**. Histogram showing the results of a bootstrapping analysis on

1121   the percentage of $CD8^+$ T cells expressing PD-1. Bars are colored by spatial

1122   architecture.

1123

1124   **Supplemental Figure 2c**. Histogram showing the results of a bootstrapping analysis on

1125   the percentage of $CD4^+$ T helper cells expressing PD-1. Bars are colored by spatial

1126   architecture.

1127

1128   **Supplemental Figure 2d**. Histogram showing the results of a bootstrapping analysis on

1129   the percentage of B cells expressing PD-1. Bars are colored by spatial architecture.

1130

1131   **Supplemental Figure 2e**. Histogram showing the results of a bootstrapping analysis on

1132   the percentage of APCs expressing Ki-67. Bars are colored by spatial architecture.

1133

1134   **Supplemental Figure 3a**. t-SNE representation of single cells from the 'Other Immune'

1135   class colored by heatmap of mean intensity for CD163 and tryptase (mast cells) show

1136 distinct populations of cells expressing high levels of CD163 and tryptase, indicating

1137 these cell types are likely present in the 'Other Immune' class.

1138

1139 **Supplemental Figure 3b**. Stacked bar chart showing the proportion (out of 1.0) of

1140 αSMA^+ cell neighborhood clusters present in each tumor region (n=47).

1141

1142 **Supplemental Figure 3c**. Stacked bar chart showing the average proportion (out of

1143 1.0) of αSMA^+ cell neighborhood clusters present in each tumor (n=18).

1144

1145 **Supplemental Figure 3d**. Stacked bar chart showing the proportion (out of 1.0) of

1146 αSMA^+ cell neighborhood clusters present in each patient's primary and recurrent

1147 tumors averaged (n=9).

1148

1149 **Supplemental Figure 3e**. Stacked bar chart showing the average proportion (out of

1150 1.0) of αSMA^+ cell neighborhood clusters present in tumors collected from the three

1151 anatomic sites.

1152

1153 **Supplemental Figure 3f**. Stacked bar chart showing the average proportion (out of 1.0)

1154 of αSMA^+ cell neighborhood clusters present in the primary tumors by their TNM stage.

Patient ID	Anatomic Site of Resection	Primary Tumor TNM Stage	Therapy Following Primary Resection	Gender	Race	Alcohol History	Smoking History	HPV Status	Progression Free Survival (days)
1	Oral Cavity	2	Cisplatin + Radiation	Male	White	Yes	Yes	Negative	804
2	Oral Cavity	4	Cetuximab + Radiation	Male	White	Yes	Yes	Negative	123
3	Oral Cavity	1	Cisplatin + Radiation	Female	White	Yes	No	Negative	1447
4	Oral Cavity	1	Cisplatin + Radiation	Female	White	No	No	Negative	246
5	Oropharynx	4	Cetuximab + Radiation	Female	White	Yes	No	Negative	202
6	Oropharynx	4	Cisplatin + Radiation	Female	Asian	No	No	Negative	188
7	Larynx	4	Cisplatin + Cetuximab + Radiation	Female	White	Yes	Yes	Negative	409
8	Larynx	3	Cisplatin + Radiation	Female	White	Yes	Yes	Negative	1033
9	Larynx	4	Cisplatin + Radiation	Female	White	Yes	Yes	Negative	83

Cell Phenotype	Antibody Markers
CD8$^+$ T Cell	CD45$^+$ CD20$^-$ CD3$^+$ CD8$^+$
CD4$^+$ T Helper	CD45$^+$ CD20$^-$ CD3$^+$ CD8$^-$ FOXP3$^-$
CD4$^+$ Regulatory T Cell	CD45$^+$ CD20$^-$ CD3$^+$ CD8$^-$ FOXP3$^+$
B Cell	CD45$^+$ CD20$^+$
Macrophage	CD45$^+$ CD20$^-$ CD3$^-$ CD66B$^-$ CD68$^+$
Granulocyte	CD45$^+$ CD20$^-$ CD3$^-$ CD66B$^+$
Antigen Presenting Cell	CD45$^+$ CD20$^-$ CD3$^-$ CD66B$^-$ CD68$^-$ MHCII$^+$
Other Immune	CD45$^+$ CD20$^-$ CD3$^-$ CD66B$^-$ CD68$^-$ MHCII$^-$ CD8$^-$ FOXP3$^-$
αSMA$^+$ Mesenchymal	CD45$^-$ PANCK$^-$ αSMA$^+$
Neoplastic Tumor	CD45$^-$ PANCK$^+$
Additional Functional Markers: PD-1, PD-L1, Ki-67	

Cell Phenotype	Average Coefficient of Variation
B Cell	0.658
Macrophage	0.563
αSMA$^+$ Mesenchymal	0.545
Neoplastic Tumor	0.541
Granulocyte	0.523
Antigen Presenting Cell	0.519
Other Immune	0.402
CD8$^+$ T Cell	0.384
CD4$^+$ T Helper	0.376
CD4$^+$ Regulatory T Cell	0.311

Blise et al. Figure 1

a Compartmentalized

More immune cells involved in antigen presentation

More Ki-67⁺ APCs

More αSMA⁺ cells

αSMA⁺ cells neighboring CD4⁺ T helper cells and B cells

αSMA⁺ cells neighboring αSMA⁺ cells

Longer PFS

b Mixed

More PD-1⁺ lymphocytes

More neoplastic tumor cells

αSMA⁺ cells neighboring neoplastic tumor cells

αSMA⁺ cells neighboring other immune cells

αSMA⁺ cells neighboring granulocytes

Shorter PFS
Later TNM stage

a

b

	Region 1	Region 2	Region 3
Patient 9 Primary			
Patient 8 Recurrent			

B cells — yellow
Nuclei — blue
500 μm scale bars

c

Legend:
- Intra-Tumor (P or R only)
- Intra-Patient (P and R)
- Inter-Patient (P or R only)
- Inter-Patient (Same Anatomic Site)
- Inter-Patient (all)

p = 0.001
p = 0.020
p = 0.001
p = 0.012
p = 0.001
p = 0.006

Y-axis: KL Divergence (0, 0.5, 1.0, 1.5, 2.0, 2.5)
X-axis: Tumor (1P, 1R, 2P, 2R, 3P, 3R, 4P, 4R, 5P, 5R, 6P, 6R, 7P, 7R, 8P, 8R, 9P, 9R)

a

b

c

d

e