Low intensity repetitive transcranial magnetic stimulation drives structural synaptic plasticity in the young and aged motor cortex.

Alexander D Tang[1,2*], William Bennett[3*], Aidan D Bindoff[3], Jessica Collins[3], Michael I Garry[4], Jeffery J Summers[4,5], Mark R Hinder[4], Jennifer Rodger[1,2], Alison J Canty[3]

1. Experimental and Regenerative Neurosciences, School of Biological Sciences, University of Western Australia, Perth, Australia.
2. Perron Institute for Neurological and Translational Sciences, Perth, Australia.
3. Wicking Dementia Research and Education Centre, College of Health and Medicine, University of Tasmania, Hobart, Australia.
4. School of Psychological Sciences, College of Health and Medicine, University of Tasmania, Hobart, Australia.
5. Research Institute for Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, United Kingdom

* Equal contribution

Author for correspondence: Alexander D Tang - Adtang.research@gmail.com

Abstract

Repetitive transcranial magnetic stimulation (rTMS) is a non-invasive tool commonly used to drive neural plasticity in the young adult and aged brain. Recent data from mouse models have shown that even at low intensities (0.12 Tesla), rTMS can drive neuronal and glial plasticity in the motor cortex. However, the physiological mechanisms underlying low intensity rTMS (LI-rTMS) induced plasticity and whether these are altered with normal ageing are unclear. Using longitudinal *in vivo* 2-photon microscopy, we investigated the effect of LI-rTMS on the structural plasticity of pyramidal neuron dendritic spines in the motor cortex following a single train of LI-rTMS (in young adult and aged animals) or the same LI-rTMS train administered on 4 consecutive days (in young adult animals only). We found that LI-rTMS altered the rate of dendritic spine losses and gains, dependent on the number of stimulation sessions and that a single session of LI-rTMS was effective in driving structural synaptic plasticity in both young adult and aged mice. To our knowledge, these results provide the first *in vivo* evidence that rTMS drives synaptic plasticity in the brain and uncovers structural synaptic plasticity as a key mechanism of LI-rTMS induced plasticity.

Introduction

For over three decades, repetitive transcranial magnetic stimulation (rTMS) has been used to drive neural plasticity in the human brain. Traditionally, this has been achieved by delivering transient pulses of magnetic fields over the scalp at high stimulation intensities (>1 Tesla) to drive activity within neural networks (i.e., suprathreshold stimulation). However, using rodent models, we have shown that even at low subthreshold intensities (e.g. 0.12 Tesla) that do not directly induce neuronal activity, rTMS can still induce neural plasticity. For example, we have shown that a single session of low intensity rTMS (LI-rTMS), alters neuronal excitability at the single cell (layer 5 pyramidal neurons) [1] and network level (motor evoked potentials) [2] whereas multiple sessions of LI-rTMS have been shown to promote neuronal (e.g. skilled motor learning [3]) and glial plasticity [4, 5]. However, the effect of LI-rTMS on synaptic plasticity has not, to date, been explored.

Irrespective of intensity, synaptic plasticity has long thought to be the main physiological mechanism underlying rTMS induced plasticity given the ability of rTMS to alter learning and motor evoked potential amplitudes in humans and rodents [6, 7]. In addition, synaptic plasticity is impaired in the aged brain [8-10] which in part, may explain the reports of reduced rTMS-induced plasticity in older adults [11-13]. However, despite its widespread use, the physiological mechanisms underlying rTMS induced neural plasticity remain unclear, making it difficult to determine which neurological diseases or disorders are suited for rTMS intervention. While high intensity repetitive magnetic stimulation has been shown to induce synaptic plasticity *in vitro* (organotypic hippocampal slices) [14-16], it is unknown whether high or low intensity rTMS induces synaptic plasticity *in vivo* and whether this is decreased in the aged brain.

Rodent rTMS models are a useful adjunct to human studies as they allow for direct measures of neural plasticity, including measurements of synaptic plasticity and connectivity in the living brain with longitudinal *in vivo* microscopy. Using this technique, structural plasticity of dendritic spines (the post-synaptic structure) can be quantified through changes to the density of dendritic spines and to the rates of

dendritic spine gains and losses. For example, in the motor cortex, structural

synaptic plasticity is a fundamental process that facilitates the learning of skilled

motor behaviours, with increases to the rate of dendritic spine gains and losses on

the apical dendrites of layer 5 pyramidal neurons during the learning period [17, 18].

Using a similar longitudinal *in vivo* imaging approach, here we characterised the

changes to structural synaptic plasticity in the motor cortex following a single session

of LI-rTMS in young adult and "aged" mice. Additionally, we investigated whether the

effects of LI-rTMS are cumulative by comparing the effects of a single day of LI-

rTMS to that of multiple days of stimulation in young adult mice.

Materials and methods

Animals

Thy1-GFP-M mice (maintained on a C57BL/6J background) were used throughout,

which express enhanced green fluorescent protein (EGFP) in a sparse subset of

cortical excitatory neurons in layers 2/3 and 5 [19]. Mice were group housed on a 12-

hour light/dark cycle and given food and water *ad libitum*. All animal experimentation

was performed under the guidelines stipulated by the University of Tasmania Animal

Ethics Committee (approval number: A0013168), which is in accordance with the

Australian code of practice for the care and use of animals for scientific purposes.

Mice were classified as "young adult" aged 3-6 months, or "aged" at 23-24 months.

The single session of LI-rTMS group (total n=10 animals) had both young adult (4

male, 1 female) and aged adult animals (n=5 all male). The multiple sessions of LI-

rTMS group (total n=5 animals) consisted of young adult animals only (3 males, 2

female).

Cranial window implantation

Animals were habituated to the animal facility for at least 7 days prior to cranial

window implantation surgery. For cranial window imaging, we used a modified

protocol from [20]. Briefly, animals were given pre-operative analgesia

(buprenorphine, 0.1mg/kg, subcutaneous), anaesthetised with isoflurane and placed

in a stereotaxic frame. Local anaesthetic (5mg/kg bupivacaine) was infiltrated under

the scalp prior to incision and skin removal. A high-speed dental drill was used to

perform a 3mm craniotomy over the right primary motor cortex region (M1) likely

blending into the SS1 somatosensory region (window centred over +1mm anterior

from bregma, +2.5mm lateral to the midline). Artificial cerebrospinal fluid (ACSF) was

applied regularly to cool the bone during drilling. The exposed dura was left intact

and was gently cleaned with sterile gelfoam (Pfizer), soaked in ACSF.

Dexamethasone was applied topically to the dura (approximately 20µl of 4mg/ml

solution), as this has been shown to enhance window clearance [21]. A 5mm circular

glass coverslip was placed over the craniotomy, and the perimeter sealed with

Loctite 454 (a low-volatile, low-temperature curing cyanoacrylate gel). A titanium bar

with a threaded hole was glued onto the left side, opposite the window. The exposed

scalp and skin margin was sealed with dental acrylic (Heraeus). Animals recovered

over a 2–3-week period post-surgery before undergoing imaging.

Multiphoton in vivo imaging

In vivo two-photon laser scanning microscopy (2PLSM) was performed using a

custom-built upright laser scanning microscope (Scientifica) equipped with galvo-

galvo mirrors, a 20X water immersion lens (NA 1.0, Zeiss Plan-Apo) and a high-

sensitivity GaAsP non-descanned photomultiplier detector

(Hamamatsu). Femtosecond-pulsed infrared excitation was from a mode-locked Ti-

sapphire laser tuned to 910nm and equipped with group velocity dispersion

compensation (Mai Tai DeepSee, Spectra-Physics). Power delivered to the back

aperture was 20-90mW, depending on depth. This setup enabled us to image a good

cranial window to >1000µm into the cortex. Images were acquired with ScanImage

3.8.1 software (Vidrio Technologies, LLC) [22] incorporating the Navigator plugin.

Animals were anaesthetised during imaging using either isoflurane or injectable

anaesthetic (100mg/kg ketamine, 10mg/kg xylazine, intraperitoneal injection). The

titanium bar embedded in the acrylic cap during surgery was attached to a

customised brass clamp to maintain stability during imaging. Images of dendritic

arbours were captured from the upper 100µm of the cortex for the purposes of

quantification, and the neuron type (layer 2/3 or layer 5) was identified with deeper

imaging. Images were captured at 0.2μm/pixel resolution in the XY plane, with 1μm

Z-steps. Dendritic arbours were selected on the basis of clarity and were not

restricted to any particular position within the window. Repeat imaging of dendritic

segments was carried out using vascular landmarks to locate the regions of interest.

Low intensity rTMS

Purpose built rodent specific coils (circular coil 8mm in diameter and height-see [2])

controlled by an arbitrary waveform generator (Agilent 335141B) connected to a

bipolar power supply (Kepco BOP 100-4M) were used to administer the rTMS trains.

The rTMS pulses were monophasic in shape (400μs rise time, dB/dT of 300T/s)

which induced a peak magnetic field of 0.12T at the base of the coil. At this intensity,

the peak sound intensity of the rTMS "clicks" are ~26dB SPL at the base of the coil

[2] which is below the hearing thresholds previously reported for adult C57Bl6J mice

[23]. Each stimulation session delivered 600 pulses of intermittent theta-burst

stimulation (iTBS) [24] which contained pulse trains consisting of 3 pulses at 50Hz,

repeated at 5Hz and delivered for 2s, with an inter-train interval of 8s (total

stimulation time 192s).

LI-rTMS was delivered 3 hours after the end of each imaging session to allow for

complete recovery from the anaesthesia, as anaesthesia during the delivery of rTMS

has been shown to alter the effect of rTMS in the rodent brain [25, 26]. Stimulation

was delivered to lightly restrained mice that were placed in a restraint bag with a

breathing hole at one end (Able Scientific). The rTMS coil was positioned over the

mouse cranium, such that the coil windings overlaid the cranial window, with the coil

offset laterally to minimise direct stimulation of the motor cortex in the other

hemisphere (i.e., the motor cortex without a cranial window). During stimulation, the

coil was held directly on the restraint bag (~1mm distance from the base of the coil to

the mouse cranium).

Imaging timeline

Imaging was separated into 3 time periods (summarised in Figure 1), (i) pre-

stimulation, (ii) +21 hours post-stimulation and (iii) +45 hours post-stimulation. For

both the single and multiple stimulation groups, pre-stimulation imaging consisted of

three imaging sessions conducted over 3 consecutive days separated by 24 hours to

capture baseline spine density and dynamics (rate of spine losses and gains). The

+21 hours post-stimulation timepoints were conducted 21 hours after each LI-rTMS

(i.e. once for the single session of LI-rTMS group or a total of four +21 hours post-

stimulation timepoints for the multiple stimulation group). A final imaging session was

conducted 24 hours after the last +21 hours post-stimulation period and represents

the +45 hours post-stimulation timepoint for both the single and multiple stimulation

groups.

Figure 1. Imaging timeline for the single and multiple sessions of LI-rTMS

groups.

Image annotation and spine analysis

Captured stacks were screened for dendrite lengths that were suitable for spine

analysis. Although no minimum length was prescribed, only dendritic segments that

had at least 11 spines at the final pre-stimulation timepoint were included in the

analysis. This resulted in a total of 29 analysable dendritic segments for the single

stimulation group (pooled from 10 mice) and 15 analysable dendritic segments for

the multiple stimulation group (pooled from 5 mice). Overlapping image stacks were

montaged together using XUVTools software [27]. Image stacks were false coloured

for depth in Fiji [28] and rotated in 3D to ensure that unique dendritic segments were

identified. Where segments joined at a branch point, they were combined and

considered to be a single segment. Segments were cropped and then annotated

using Matlab scripts included with ScanImage (scim_spineAnalysis.m) to correlate

spines across longitudinal time series [20, 22]. Spines were annotated by drawing a

line from the centre of the dendritic backbone to the tip of the spine and were only

annotated if present across two consecutive z-planes within the stack. Spines were

then correlated across the timeseries and exported as spreadsheets of spine lengths

where each column was a timepoint and each row corresponded to a single spine

across the timeseries. For each dendritic segment, the width of the dendritic

backbone was measured at five evenly spaced points along the segment and spine

lengths were corrected by subtracting half the mean backbone width.

A custom R script (R core team, 2014) was used to analyse gains and losses, where

the criterion for a spine being counted as present was set to 0.4µm (corresponding to

two pixels protruding from the edge of the backbone). In addition, a spine had to

reach a minimum of 0.6µm (three pixels) at some point along the timeseries, or the

entire row was censored. Gains were therefore classified as when a spine went from

absent (<0.4µm) to present (≥0.4µm), with loss being the inverse. Transient losses

where the spine was absent for only a single timepoint before reappearing at the

same location were checked manually. In all except a single case (out of more than

90 apparent transient loss events), these were found to be spurious, and were

caused by poor image quality, or rotation of the spine around the backbone to project

along the z-axis (detectable by an increase in the brightness of the backbone). All

such transient loss events were reclassified as being present throughout. Following

data curation and removal of dendritic arbours with insufficient spines, the dendritic

arbours which were analysed statistically tallied (across all timepoints) 17 segments

(total length 1807µm) with 2423 annotated spines in the young adult single

stimulation group; 12 segments (total length 1691µm) with 1852 annotated spines in

the aged adult single stimulation group and 15 segments (total length 1586µm) with
2865 spines annotated in the young adult multiple stimulation group.

Figure 2. Representative images of an analysed dendritic arbour from the
single session of LI-rTMS group.

(A) Portion of a dendritic arbour, shown as a maximum projection of a 120µm
stack, false colour coded for depth. The colour scale along the bottom shows the
correspondence between hue and depth within the stack. Depth of an arbour within
the stack does not necessarily correspond to depth from the cortical surface, due to
the angle of capture. However, the colours do indicate the relative depths of
adjacent or overlapping dendrites. The rectangular box indicates a portion of the
annotated subregion shown in panel B).

(B) 60µm subregion of a single annotated dendrite from the single stimulation group
(total annotated length was 242µm), showing five sequential timepoints at 24hr
intervals. The red asterisk indicates LI-rTMS stimulation, occurring 3hrs after the
Pre-stim 3 image. Spines were annotated in MATLAB and quantitatively analysed
using custom R scripts, with the threshold for spine presence set to projection of
≥0.4µm from the edge of the dendritic backbone. Spines over the time series were
classified as stable, lost or gained. Blue arrowheads indicate spines present
throughout (stable).

Filled red arrowheads indicate spines present that will be lost, with the point of loss

shown as a hollow red arrowhead. Hollow green arrowheads indicate nascent

spines that have not yet exceeded the threshold, with the solid green arrowhead

indicating the point of gain.

Statistics

For statistical comparison, we compared the 3 time periods (pre-stimulation, +21hrs

post-stimulation and +45hrs post-stimulation). Spine density represents the number

of spines per μm measured at each imaging session/timepoint. The rate of spine

losses and gains represents the number of new spines (per μm) gained or lost (per

spine) *since* the last imaging session. The rate of spine losses cannot be calculated

for the first pre-stimulation timepoint, as this timepoint provides the baseline counts

of spines for each dendritic arbour. To facilitate comparison between gains and

losses, we similarly excluded this timepoint for the analysis of gains.

Bayesian hierarchical generalized linear regression models were used to estimate

($\log_e$-) spine density and the rate of spine losses and gains at each 24-hour

observation and the average rate (per 24-hour) within the 3 imaging periods (pre-

stimulation, +21hrs post-stimulation, +45hrs post stimulation with LI-rTMS) for each

stimulation group. Data were counts, therefore a Poisson distribution was assumed,

with log-link function. The rate of spine losses per 24-hours was adjusted for the

($\log_e$-) spine count at the previous 24-hour observation of the same dendrite. The

rate of spine gain per 24-hours was adjusted for dendritic length ($\log_e$-μm), as longer

dendrites are likely to have more spines. Weakly regularizing student-t priors with 3

degrees of freedom (fat tails) were specified, expressing some scepticism that there

was any difference in rate of gains or losses between observations or stimulation

periods.

Given that the data in the single stimulation and multiple stimulation groups are

identical in the way they are collected from the pre-stimulation period up to the first

+21hrs post-stimulation observation (see Figure 1), a pooled analysis was run on this

combined dataset to supplement our single stimulation analysis.

All statistical analysis was conducted in the R statistical computing environment (R Core Team, 2014). We used the R package 'brms' [29] to specify models using Hamiltonian Monte Carlo sampling in Stan [30]. Chain mixing and convergence was assessed using trace plots and R-hat statistics, and model fit was assessed using (graphical) posterior predictive checks. Estimated marginal means (averaged over age and/or layer) were computed using the 'emmeans' R package [31]. Marginal likelihoods were computed over 27×10^3 posterior samples (after a 3×10^3 iteration warmup). All code and data to reproduce the analysis will be accessible through online repositories (https://research-repository.uwa.edu.au/en/persons/alex-tang/datasets/ and https://eprints.utas.edu.au/).

We fitted 3-level models with random intercepts for dendritic segments nested within animal, with age and cell body location (layers 2/3 or 5) included as fixed effects. We report rate of gains per day, per micron, and rate of losses per day, per spine (lagged to the previous observation as described above). Uncertainty is expressed as 95% (posterior) credible intervals (CI). As we used an internal control design, our analysis (including hypothesis testing) estimates the dendritic spine dynamics at the population-level at the pre-, +21hrs post-stimulation and +45hrs post-stimulation time periods. Bayes Factors (BF- evidence ratios) are reported to aid in the interpretation of our results and we used the following BF cut-offs as evidence in favour of the alternative hypothesis [32].

- **BF = 1** - No evidence
- **1 < BF <= 3** - Anecdotal
- **3 < BF <= 10** - Moderate
- **10 < BF <= 30** - Strong
- **BF>100** – Extreme

Results

Single session of LI-rTMS (young adult and aged mice)

Density

Compared to pre-stimulation (0.216 spines μm^{-1}, 95%CI 0.149, 0.284, averaged over

layer and age, n=29 dendritic segments from 10 mice) there was anecdotal evidence

for a decrease in spine density at +21hrs post-stimulation (0.213 spines μm^{-1}, 95%CI

0.146, 0.282, BF=1.78). In contrast, there was strong evidence for a decrease in the

mean dendritic spine density at +45hrs post-stimulation (0.201 spines μm^{-1}, 95%CI

0.137, 0.265, BF =18.2, see Figure 3A). Further analysis showed anecdotal evidence

for a greater effect of LI-rTMS on dendritic spine density in young adults (n=5)

compared to aged animals (n=5) (BF=1.5).

Rate of spine loss

Compared to pre-stimulation (0.228 $spine^{-1}$, 95%CI 0.156, 0.315, averaged over

layer and age), there was strong evidence for an increase in the mean rate of spine

losses following LI-rTMS at +21hrs post-stimulation (0.262 $spine^{-1}$, 95%CI 0.173,

0.361, BF =13.4) but not at +45hrs post-stimulation (0.227 $spine^{-1}$, 95%CI 0.151,

0.313, BF=1.1, see Figure 3C). In addition, there was strong evidence for a greater

rate of spine losses at +21 and compared to +45hrs post-stimulation (BF=10.3),

further suggesting that there was an increase in the rate of spine loss following LI-

rTMS that resolves by +45hrs post-stimulation. Analysis of age showed moderate

evidence for a greater increase in the rate of spine losses in young adult animals

compared to aged adult animals (BF=4.2).

Rate of spine gain

Compared to pre-stimulation (0.046 spines μm^{-1}, 95%CI 0.035, 0.058, averaged over

layer and age), there was anecdotal evidence of an increase in the rate of spine gain

following LI-rTMS +21hrs post-stimulation (0.048 spines μm^{-1}, 95%CI 0.035, 0.06,

averaged over layer and age, BF = 2.0, see Figure 3E). In contrast, there was

extreme evidence for a decrease in the rate of spine gain at +45hrs post-stimulation

(0.034 spines μm^{-1}, 95%CI 0.025, 0.044, averaged over layer and age, pre-

stimulation> +45hrs post-stimulation: BF = 564.4, +45hrs post-stimulation< +21hrs

post-stimulation: BF = 661.6). There was anecdotal evidence for a greater change in

the rate of spine gain between young adult and aged animals (BF=0.9).

Multiple sessions of LI-rTMS (young adult mice)

Density

Compared to the pre-stimulation period (0.216 spines μm^{-1}, 95%CI 0.149, 0.284, n=15 dendritic segments from 5 mice, averaged over layer), there was anecdotal evidence for a decrease in the mean density of dendritic spines following LI-rTMS at the +21hrs post-stimulation timepoints (averaged over the 4 +21hrs post-stimulation timepoints, 0.213 spines μm^{-1}, 95%CI 0.146, 0.282, BF = 1.8, see Figure 3B) and at +45hrs post-stimulation (0.220 spines μm^{-1}, 95%CI 0.137, 0.265, BF = 2.4). In addition, there was anecdotal evidence for an increase in density between the +21hrs post-stimulation timepoints and +45hrs post-stimulation (BF=4.2).

Rate of spine loss

Compared to pre-stimulation (0.166 $spine^{-1}$, 95%CI 0.116, 0.219, averaged over layer), there was strong evidence for an increase in the mean rate of spine losses at the +21hrs post-stimulation timepoints (averaged over the 4 +21hrs post-stimulation timepoints 0.199 $spine^{-1}$, 95%CI 0.146, 0.255, BF= 20.4, see Figure 3D) but not at +45hrs post-stimulation (0.175 $spine^{-1}$, 95%CI 0.111, 0.240, BF=1.6).

Rate of spine gain

Compared to pre-stimulation (0.057 spines μm^{-1}, 95%CI 0.014, 0.109, averaged over layer), there was anecdotal evidence for a decrease in the mean rate of spine gain at the +21hrs post-stimulation timepoints (averaged over the 4 +21hrs post-stimulation timepoints +0.059 spines μm^{-1}, 95%CI 0.016, 0.115, BF= 1.9, see Figure 3F). Following LI-rTMS, there was moderate evidence for a decrease in the rate of spine gain at +45hrs post-stimulation (0.050 spines μm^{-1} 95%CI 0.013, 0.098, BF = 3.5).

Figure 3: Single (left side) and multiple sessions of LI-rTMS (right side) drive structural synaptic plasticity in pyramidal neurons of the motor cortex.

(A and B) Analysis of dendritic spine density showed strong evidence for a decrease in the mean dendritic spine density +45hrs post a single session of LI-rTMS, compared to pre-stimulation.

(C and D) Analysis of the rate of dendritic spine losses showed strong evidence for an increase in the mean rate of spine losses +21hrs post-stimulation in both the single and multiple sessions of LI-rTMS groups compared to pre-stimulation.

(E and F) Analysis of the rate of dendritic spine gains showed strong evidence for a decrease in the mean rate of dendritic spine gains +45hrs post a single session of LI-rTMS compared to pre-stimulation.

Data are shown as the aggregate means (solid-coloured lines) for each time period
(pre-stimulation=blue, +21hrs post-stimulation=green, +45hrs post-
stimulation=orange) overlaid on the mean (•) at each imaging observation. Error
bars represent the 95% **credible intervals** for each timepoint, whereas the shaded
boxes represent the 95% credible intervals for each time period. Each data point
represents data from an individual dendritic arbour with data from young (o) and
aged (x) animals.

Pooled analysis of single and multiple stimulation data (from pre-stimulation
to the first +21hrs post-stimulation timepoint)

As the single and multiple stimulation groups are identical experimentally, to the first
+21hrs post-stimulation imaging timepoint (i.e., up to +21hrs post a single
stimulation), we pooled the data from both groups up until this point to supplement
our single stimulation analysis.

Density

Increasing the sample size in the pooled analysis (total n=44 dendritic segments
from 26 arbours, from 15 mice) showed anecdotal evidence for an increase in the
mean dendritic spine density +21hrs post a single session of LI-rTMS (0.229 spines
μm^{-1}, 95%CI 0.173, 0.288) compared to pre-stimulation (0.223 spines μm^{-1}, 95%CI
0.171, 0.284, BF = 2.1, see Supplementary Figure 1A).

Rate of spine loss

Pooled analysis showed strong evidence of an increase in the rate of spine losses at
+21hrs post a single session of LI-rTMS (0.238 $spine^{-1}$, 95% CI 0.150, 0.263)
compared to pre-stimulation (0.204 $spine^{-1}$, 95% CI 0.174, 0.307, BF=34.9, see
Supplementary Figure 1B).

Rate of spine gains

Pooled analysis showed strong evidence (BF=35.0, see Supplementary Figure 1C)

for an increase in the rate of spine gain at the +21hrs post a single session of LI-

rTMS (0.053 spines μm^{-1} 95%CI 0.035, 0.057) compared to pre-stimulation (0.045

spines μm^{-1} 95%CI 0.039, 0.066).Given that this result was not evident in the single

stimulation analysis, this suggests that this effect is not as robust as the change

seen in spine losses, as it could only be detected by increasing the sample size

Further analysis of the pooled data did not indicate that animals in the multiple

stimulation group had a greater increase in gains during the first +21hrs post-

stimulation period compared to the animals in the single stimulation group (see

Supplementary Material).

Table 1. Summary of results after LI-rTMS (i.e., post-stimulation time points compared to pre-stimulation). The direction of change (increased/decreased) is indicated only for results with a BF≥10 (i.e., with ≥strong evidence of a change). NOTE: "Age" is not included in the table as it was investigated in the single stimulation group only and did not show strong evidence for any spine measure. Pooled analysis encompasses +21hrs post-stimulation time points only.

	Dendritic spine density		Rate of dendritic spine losses		Rate of dendritic spine gains	
	+21 hours post-stim	+45 hours post-stim	+21 hours post-stim	+45 hours post-stim	+21 hours post-stim	+45 hours post-stim
Single session of LI-rTMS	- BF=1.78	↓ BF=18.2	↑ BF=13.4	- BF=1.1	- BF=2.0	↓ BF=564.4
Multiple sessions of LI-rTMS	- BF=1.8	- BF=2.4	↑ BF=20.4	- BF=1.6	- BF=1.9	- BF=3.5
Pooled analysis up to first +21hrs post-stimulation timepoint	- BF=2.1	■	↑ BF=34.9	■	↑ BF=35.0	■

Discussion

To our knowledge, this is the first *in vivo* study to provide direct evidence that rTMS, irrespective of intensity, drives structural synaptic plasticity in the motor cortex. Following a single session of LI-rTMS, there was an increase to the rate of dendritic spine losses (present at +21hrs post-stimulation) and a long-lasting decrease in

dendritic spine density and to the rate of dendritic spine formation (present at +45hrs
post-stimulation) which did not differ between young and aged animals (see Figure
3). Surprisingly, multiple days/sessions of LI-rTMS did not have a cumulative effect
or longer lasting effect on structural plasticity compared to a single session of LI-
rTMS but instead maintained the more acute effects induced by a single LI-rTMS
session (increased rate of spine losses at +21hrs post-stimulation) which returned to
baseline values by +45hrs after the last LI-rTMS session.

Given that previous high intensity rTMS studies in humans have shown age-related
differences in rTMS-induced plasticity in other measures of neural plasticity (motor
evoked potentials and the default mode network) with the iTBS protocol [12, 13], we
might have expected similar results in our study. However, we failed to find strong
evidence of a difference between young adult and aged animals in the structural
plasticity of dendritic spines induced by a single session of LI-rTMS. This may be
explained by the absence of age-dependent differences in dendritic spine properties
pre-stimulation, such that LI-rTMS modulated both the young and aged brain as they
did not differ in the amounts of synaptic connectivity and remodelling capacity prior
to stimulation. Interestingly, these results are in direct contrast to the work of
Davidson et al., who reported age-dependent differences in pyramidal neuron
dendritic spine density and dynamics in the forelimb region of the motor cortex in
Thy1-GFP mice [10]. Specifically, in both our young and aged animals, the mean
dendritic spine density and rate of spine *gains* in the pre-stimulation period was
approximately half of that reported by Davidson et al., whereas the rate of spine
losses from our young adults was almost twice as large. In addition to several
methodological differences that may explain these contrasting findings, a key point of
difference between our studies was that our cranial windows were centred more
anterior relative to bregma to that used in Davidson et al. Therefore, while we did not
find age-dependent effects on baseline or LI-rTMS induced spine plasticity in the
area of the motor cortex we imaged, it's possible that such differences do occur,
depending on the anatomical region imaged or circuit stimulated.

In young adult mice, we have previously shown that 10 daily sessions of LI-rTMS

(with the same 600 pulses of iTBS protocol) to the motor cortex paired with skilled

motor training improves motor behaviour and the rate of learning [3]. In that study,

our behavioural analysis suggested that multiple sessions of LI-rTMS maintained the

transient effects induced by a single stimulation. Similarly, repeated blocks of high

intensity iTBS within a single stimulation session to the rat barrel cortex was shown

to have a cumulative effect on cortical activity [33]. Interestingly, multiple sessions of

LI-rTMS maintained the initial structural plasticity induced by LI-rTMS, with both

groups showing an increase in the rate spine losses at the +21hrs post-stimulation

timepoint (pooled analysis- see supplementary Figure 1). However, the effects of

multiple sessions were shorter lasting (+21hrs post-stimulation) compared to a single

session which also showed decreased spine density and rate of spine gains at the

+45hrs post-stimulation timepoint. While it is unclear why a single-session of LI-rTMS

had longer lasting effects on structural plasticity than multiple-sessions, the net

changes observed with both groups suggest a common outcome of refinement in

synaptic connectivity as an underlying mechanism of LI-rTMS-induced plasticity.

This is the first demonstration, to our knowledge, that LI-rTMS has delayed effects

on neural plasticity, as our previous work has shown that LI-rTMS (using the iTBS

protocol) drives acute changes to motor behaviour [3] and neuronal excitability [1].

While further investigation is needed to determine the functional consequences of

structural plasticity induced with LI-rTMS, we suggest that the refinement in synaptic

connectivity induced may promote neural plasticity, by removing or preventing

redundant synaptic connections, thereby producing a more efficient neural circuit.

This is consistent with the use of LI-rTMS (0.012T) in ephrin-A2A5-/- mice showing

that LI-rTMS reorganises abnormal circuitry by removing aberrant connections [34,

35].This delayed form of LI-rTMS induced structural plasticity in the motor cortex

may work in tandem with, or be a result of the immediate effects of LI-rTMS, such

that the acute changes in neuronal excitability enhance motor behaviour which is

later supported by the delayed refinement/pruning of the neural circuitry. However,

future studies combining LI-rTMS with motor training and *in vivo* imaging is needed

to confirm our theory.

The use of rodent-specific coils offered us the advantage of delivering focal

stimulation (e.g. restricting stimulation to one motor cortex) but at the expense of

stimulation intensity [2]. Therefore, while we have shown that LI-rTMS (i.e.

subthreshold stimulation) induces synaptic plasticity, it remains unknown what effect

high intensity/suprathreshold rTMS has on structural synaptic plasticity *in vivo*.

Investigating such changes in a mouse model is technically challenging as "small"

commercial coils (e.g. 25mm figure of 8 coils) stimulate the entire brain [36]. In

addition, despite promising improvements in rodent-specific TMS coils [37], there are

currently no rTMS coils appropriately sized for the mouse brain that can deliver high

intensity rTMS, due to the thermal stress induced with each pulse [2, 38], particularly

at high stimulation frequencies such as iTBS. As a result, characterising structural

synaptic plasticity with *in vivo* imaging following high intensity rTMS may be more

practical in a rat model, where commercial coils can be used with some degree of

focality [26, 39, 40].

Despite the extremely low sound generation (<26dBL sound pressure) produced by

our rodent coils during rTMS, a limitation of our study is the absence of a sham

stimulation group. While internal controls/ comparisons to pre-stimulation measures

are commonly used in human rTMS studies (e.g., motor evoked potential properties

normalised to pre-stimulation values), we cannot rule out the possibility that our

results were influenced by factors outside of LI-rTMS such as daily anaesthesia.

However, given that we did not observe a consistent change over time (e.g., a

change in dendritic spine density that increased over time) or consistent changes

between the stimulation groups (e.g., a change in a dendritic spine property that was

seen at the 5th imaging session in both the single and multiple stimulation groups),

repeated anaesthesia is unlikely to have been a significant factor.

In conclusion, our results show that LI-rTMS drives structural synaptic plasticity in

the mouse motor cortex. These findings uncover structural synaptic plasticity as a

key mechanism of LI-rTMS induced neural plasticity and more broadly, highlights the

ability of rTMS to alter synaptic connectivity in the brain.

Acknowledgements

The authors thank Claire Hadrill for her assistance with the initial spine annotations and Dr Saied Mehrkanoon for assistance with the modification and re-generation of the MATLAB scripts used for spine annotations. This project was funded by the National Health and Medical Research Council (NHMRC - APP1050261). JR was supported by research fellowships from the NHMRC (APP1002258) and Multiple Sclerosis Western Australia. MRH was supported by research fellowships from the Australian Research Council (DE120100729 and FT150100406). AT was supported by doctoral scholarship from the Australian Government, the Bruce and Betty Green Foundation and post-doctoral funding from the Raine Medical Research Foundation (RPG06-20).

References

1.	Tang AD, Hong I, Boddington LJ, Garrett AR, Etherington S, Reynolds JN, et al. Low-intensity repetitive magnetic stimulation lowers action potential threshold and increases spike firing in layer 5 pyramidal neurons in vitro. Neuroscience. 2016;335:64-71.

2.	Tang AD, Lowe AS, Garrett AR, Woodward R, Bennett W, Canty AJ, et al. Construction and evaluation of rodent-specific rTMS coils. Front Neural Circ. 2016;10:47.

3.	Tang AD, Bennett W, Hadrill C, Collins J, Fulopova B, Wills K, et al. Low intensity repetitive transcranial magnetic stimulation modulates skilled motor learning in adult mice. Sci Rep. 2018;8(4016).

4.	Cullen CL, Senesi M, Tang AD, Clutterbuck MT, Auderset L, O'Rourke ME, et al. Low-intensity transcranial magnetic stimulation promotes the survival and maturation of newborn oligodendrocytes in the adult mouse brain. Glia. 2019;67(8):1462-77.

5.	Cullen CL, Pepper RE, Clutterbuck MT, Pitman KA, Oorschot V, Auderset L, et al. Periaxonal and nodal plasticities modulate action potential conduction in the adult mouse brain. Cell Rep. 2021;34(3):108641.

6.	Thickbroom GW. Transcranial magnetic stimulation and synaptic plasticity: experimental framework and human models. Exp Brain Res. 2007;180(4):583-93.

7.   Tang A, Thickbroom G, Rodger J. Repetitive transcranial magnetic stimulation
of the brain: mechanisms from animal and experimental models. Neuroscientist.
2017;23(1):82-94.

8.   Landfield PW, McGaugh JL, Lynch G. Impaired synaptic potentiation
processes in the hippocampus of aged, memory-deficient rats. Brain Res.
1978;150(1):85-101.

9.   Pinho J, Vale R, Batalha VL, Costenla AR, Dias R, Rombo D, et al. Enhanced
LTP in aged rats: Detrimental or compensatory? Neuropharmacology. 2017;114:12-
9.

10.   Davidson AM, Mejía-Gómez H, Jacobowitz M, Mostany R. Dendritic spine
density and dynamics of layer 5 pyramidal neurons of the primary motor cortex are
elevated with aging. Cereb Cortex. 2020;30(2):767-77.

11.   Todd G, Kimber TE, Ridding MC, Semmler JG. Reduced motor cortex
plasticity following inhibitory rTMS in older adults. Clin Neurophysiol.
2010;121(3):441-7.

12.   Opie GM, Vosnakis E, Ridding MC, Ziemann U, Semmler JG. Priming theta
burst stimulation enhances motor cortex plasticity in young but not old adults. Brain
Stimul. 2017;10(2):298-304.

13.   Abellaneda-Pérez K, Vaqué-Alcázar L, Vidal-Piñeiro D, Jannati A, Solana E,
Bargalló N, et al. Age-related differences in default-mode network connectivity in
response to intermittent theta-burst stimulation and its relationships with maintained
cognition and brain integrity in healthy aging. NeuroImage. 2019;188:794-806.

14.   Vlachos A, Müller-Dahlhaus F, Rosskopp J, Lenz M, Ziemann U, Deller T.
Repetitive magnetic stimulation induces functional and structural plasticity of
excitatory postsynapses in mouse organotypic hippocampal slice cultures. J
Neurosci. 2012;32(48):17514-23.

15.   Lenz M, Platschek S, Priesemann V, Becker D, Willems LM, Ziemann U, et al.
Repetitive magnetic stimulation induces plasticity of excitatory postsynapses on
proximal dendrites of cultured mouse CA1 pyramidal neurons. Brain Struct Funct.
2015;220(6):3323-37.

16.   Lenz M, Galanis C, Müller-Dahlhaus F, Opitz A, Wierenga CJ, Szabó G, et al.
Repetitive magnetic stimulation induces plasticity of inhibitory synapses. Nat
Commun. 2016;7:10020.

17.   Xu T, Yu X, Perlik AJ, Tobin WF, Zweig JA, Tennant K, et al. Rapid formation

and selective stabilization of synapses for enduring motor memories. Nature.

2009;462(7275):915-9.

18.   Fu M, Yu X, Lu J, Zuo Y. Repetitive motor learning induces coordinated

formation of clustered dendritic spines in vivo. Nature. 2012;483(7387):92-5.

19.   Feng G, Mellor RH, Bernstein M, Keller-Peck C, Nguyen QT, Wallace M, et al.

Imaging neuronal subsets in transgenic mice expressing multiple spectral variants of

GFP. Neuron. 2000;28(1):41-51.

20.   Holtmaat A, Bonhoeffer T, Chow DK, Chuckowree J, De Paola V, Hofer SB, et

al. Long-term, high-resolution imaging in the mouse neocortex through a chronic

cranial window. Nat Protoc. 2009;4(8):1128-44.

21.   Park K, You J, Du C, Pan Y. Cranial window implantation on mouse cortex to

study microvascular change induced by cocaine. Quant Imaging Med Surg.

2014;5(1):97-107.

22.   Pologruto TA, Sabatini BL, Svoboda K. ScanImage: Flexible software for

operating laser scanning microscopes. Biomed Eng Online. 2003;2(1):13.

23.   Yu Y-F, Zhai F, Dai C-F, Hu J-J. The relationship between age-related

hearing loss and synaptic changes in the hippocampus of C57BL/6J mice. Exp

Gerentol. 2011;46(9):716-22.

24.   Huang Y-Z, Edwards MJ, Rounis E, Bhatia KP, Rothwell JC. Theta burst

stimulation of the human motor cortex. Neuron. 2005;45(2):201-6.

25.   Gersner R, Kravetz E, Feil J, Pell G, Zangen A. Long-term effects of repetitive

transcranial magnetic stimulation on markers for neuroplasticity: differential

outcomes in anesthetized and awake animals. J Neurosci. 2011;31(20):7521. doi:

10.1523/JNEUROSCI.6751-10.2011.

26.   Sykes M, Matheson NA, Brownjohn PW, Tang AD, Rodger J, Shemmell JBH,

et al. Differences in motor evoked potentials induced in rats by transcranial magnetic

stimulation under two separate anesthetics: implications for plasticity studies. Front

Neural Circ. 2016;10(80).

27.   Emmenlauer M, Ronneberger O, Ponti A, Schwarb P, Griffa A, Filippi A, et al.

XuvTools: free, fast and reliable stitching of large 3D datasets. J Microsc.

2009;233(1):42-60.

28.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et

al. Fiji: an open-source platform for biological-image analysis. Nat Methods.

2012;9(7):676-82.

29.   Bürkner P-C. Advanced Bayesian multilevel modeling with the R package

brms. R J. 2018;10(1).

30.   Stan-development-team. RStan: The R interface to Stan. R package version

2.19.3. http://mc-stan.org/ ed2020.

31.   Lenth R, Singmann H, Love J, Buerkner P, Herve M. Emmeans: Estimated

marginal means, aka least-squares means. R package version. 2018;1(1):3.

32.   Jeffreys H. The Theory of Probability. Third Edition ed: Oxford University

Press; 1961.

33.   Thimm A, Funke K. Multiple blocks of intermittent and continuous theta-burst

stimulation applied via transcranial magnetic stimulation differently affect sensory

responses in rat barrel cortex. J Phsyiol. 2015;593(4):967-85.

34.   Rodger J, Mo C, Wilks T, Dunlop SA, Sherrard RM. Transcranial pulsed

magnetic field stimulation facilitates reorganization of abnormal neural circuits and

corrects behavioral deficits without disrupting normal connectivity. FASEB J.

2012;26(4):1593-606.

35.   Makowiecki K, Harvey AR, Sherrard RM, Rodger J. Low-intensity repetitive

transcranial magnetic stimulation improves abnormal visual cortical circuit

topography and upregulates BDNF in mice. J Neurosci. 2014;34(32):10780.

36.   Alekseichuk I, Mantell K, Shirinpour S, Opitz A. Comparative modeling of

transcranial magnetic and electric stimulation in mouse, monkey, and human.

Neuroimage. 2019;194:136-48.

37.   Meng Q, Jing L, Badjo JP, Du X, Hong E, Yang Y, et al. A novel transcranial

magnetic stimulator for focal stimulation of rodent brain.

38.   Weissman JD, Epstein CM, Davey KR. Magnetic brain stimulation and brain

size: relevance to animal studies. Electroencephalogr Clin Neurophysiol.

1992;85(3):215-9.

39.   Rotenberg A, Muller PA, Vahabzadeh-Hagh AM, Navarro X, López-Vales R,

Pascual-Leone A, et al. Lateralization of forelimb motor evoked potentials by

transcranial magnetic stimulation in rats. Clin Neurophysiol. 2010;121(1):104-8.

40.   Murphy SC, Palmer LM, Nyffeler T, Müri RM, Larkum ME. Transcranial

magnetic stimulation (TMS) inhibits cortical dendrites. eLife. 2016;5:e13598. doi:

10.7554/eLife.13598.

632