Title

The Neural Basis of Delayed Gratification

Short Title

The Neural Basis of Delayed Gratification

Authors

Zilong Gao[1, 2#], Hanqing Wang[3#], Chen Lu[4], Tiezhan Lu[1, 2], Sean Froudist-Walsh[3],

Ming Chen[4], Xiao-jing Wang[3*], Ji Hu[4, 5*], Wenzhi Sun[2, 6*]

Affiliations

[1] Academy for Advanced Interdisciplinary Studies, Peking University, Beijing

100871, China

[2] Chinese Institute for Brain Research, Beijing 102206, China

[3] Center for Neural Science, New York University, New York, NY 10003, USA.

[4] School of Life Science and Technology, ShanghaiTech University, Shanghai

201210, China

[5] Shanghai Key Laboratory of Psychotic Disorders, Shanghai Mental Health Center,

Shanghai 200030, China

[6] School of Basic Medical Sciences, Capital Medical University, Beijing 100069,

China

[#] These authors contributed equally to this work

[*] Corresponding authors. Email: xjwang@nyu.edu, huji@shanghaitech.edu.cn &

sunwenzhi@cibr.ac.cn

Abstract

Balancing instant gratification versus delayed, but better gratification is important for optimizing survival and reproductive success. Although delayed gratification has been studied through human psychological and brain activity monitoring as well as animal research, little is known about its neural basis. We successfully trained mice to perform a waiting-for-water-reward delayed gratification task and used these animals in physiological recording and optical manipulation of neuronal activity during the task to explore its neural basis. Our results showed that the activity of dopaminergic neurons in the ventral tegmental area increases steadily during the waiting period. Optical activation or silencing of these neurons, respectively, extends or reduces the duration of waiting. To interpret this data, we developed a reinforcement learning (RL) model that reproduces our experimental observations. Steady increases in DAergic activity signal the value of waiting and support the hypothesis that delayed gratification involves real-time deliberation.

KEYWORDS

Delayed Gratification, Dopaminergic, Ventral Tegmental Area, Ramping Activity, Reinforcement Learning, Continuous Deliberation

TEASER

Sustained ramping of dopaminergic activation helps individuals to resist impulsivity and wait for a later but larger reward.

INTRODUCTION

To optimize survival and reproductive success, animals need to balance instant gratification versus delayed, but better gratification. Repeated exposure to instant gratification may disrupt this balance, thereby increasing impulsive decisions. Such decisions contribute to numerous human disorders, such as addiction and obesity(*1, 2*). Delayed gratification is an important process that balances time delay with increased reward (*3*). It is influenced by strengths in patience, willpower, and self-control(*4*). Psychologists and neuroscientists have long studied this important behavior through human psychological and brain activity assessments as well as rodent-based studies. Although the dopamine system has been implicated in delayed gratification, nevertheless the precise neural activity of dopamine system that allows better gratification has not been demonstrated. Additionally, no studies to date have causally manipulated the dopamine system during delayed gratification tasks.

During a well-controlled delayed gratification task, an individual must balance the benefits vs. risks of delay in receiving an available reward. The choice to continue waiting requires suppression of the constant temptation of an immediate reward, in favor of an enhanced reward in the future (*3, 5, 6*). Midbrain dopaminergic (DAergic) neurons are well known to play central roles in reward-related and goal-directed

behaviors (*7-12*). Studies have revealed that DAergic activity signals spatial or operational proximity to distant rewards (*7, 13, 14*), which has been postulated to sustain or motivate goal-directed behaviors while resisting distractions. DAergic neurons play important roles in time judgment (*15*) and cost-benefit calculations which are necessary for value-based decision making (*13, 16-18*).

We successfully trained mice to perform a waiting-for-water-reward delayed gratification task. Recording and manipulation of neuronal activities during this task allowed us to explore the cellular regulation of delayed gratification. We found that the activity of ventral tegmental area (VTA) DAergic neurons ramped up consistently while mice were waiting in place for rewards. Transient activation of DAergic neurons extended, whereas inhibition reduced the duration of the waiting period. Then we adopted reinforcement learning (RL) computational models to predict and explain our experimental observations.

RESULTS

Mice can learn to wait for greater rewards by delayed gratification task training

First, we trained water-restricted mice to perform a one-arm foraging task (pre-training) in which delay did not result in an increased reward (*19*). The period in which the mouse remained in the waiting zone was defined as the waiting duration, and the time during which the mouse traveled from the waiting zone to reach the water reward was defined as the running duration (Fig. 1A, left panel). When the mouse exited the waiting zone and licked the water port in the reward zone, it could

receive a 10 µl of water drop regardless of the time spent in the waiting zone (Fig. 1A,

right panel, black line). During a week of training, the average waiting and running

durations both significantly decreased from Day 1 to Day 7 (Day1, Waiting: 5.58±

0.63 sec; Running: 3.46±0.28 sec; p<0.001; Day 7, Waiting: 1.99±0.19 sec; Running:

1.28±0.09 sec, p<0.001, n=7 mice, Friedman test, Figs. 1C-E, Movie S1). All mice

learned the strategy of reducing durations of both waiting and running to maximize

the reward rate (defined as µl of water per second in a given trial, Fig. S1C).

Next, we trained the same mice using a delayed gratification paradigm, where the

size of the reward increased quadratically with time spent in the waiting zone (Fig.

1A, right panel, green line). Over the next three weeks, this resulted in shifting the

distributions of the waiting duration towards a longer wait. The averaged waiting

period significantly increased from 2.76 ± 0.15s on Day 1 to 4.62 ± 0.30 s on Day 15

(p < 0.001, n = 7 mice, Friedman test, Figs. 1F, H, and Movie S2), Continuous

training steadily increased the averaged waiting duration, whereas the training did not

change the average running duration from 1.19±0.13s on Day 1 to 1.24 ± 0.10 s on

Day 15 (p = 0.97, n = 7 mice, Friedman test, Figs. 1G, H). The reward rate increased

steadily, indicating that the mice were successfully learning to successfully delay

gratification (fig. S1D).

The activity of VTA DAergic neurons increases steadily during the waiting

period

To monitor the activity of VTA DAergic neurons during the delayed gratification task, we employed fiber photometry to record the calcium signals in VTA DAergic neurons in freely moving mice for as long as one month (Fig. 2A-C, optical fiber placement illustrated in Fig S2). On the first day of pre-task training, the calcium signal rose rapidly upon reward and quickly reached a peak. A few days of training dramatically reshaped the response pattern. Once the mice re-entered the waiting zone, the activity of DAergic neurons started to rise and reached the highest level when the animal received a reward (fig. S3A).

We next analyzed the activity of these same neurons in the mice as they learned the delayed gratification task. The recording traces showed that training gradually reshaped the pattern and time course of activity (Fig. 2D). The activity started to ramp up once the mice entered the waiting zone, and then reached its highest level when they exited. To investigate carefully the dynamical properties of the ramping activity during waiting, we sorted the calcium signals from Day 15 of one mouse by their length of waiting durations and plotted them with a heat map (Fig. 2E). We divided trials according to the trial outcome (reward volume) and calculated the calcium signals while the mouse exited waiting zone with different reward volumes. Our results showed that the Z-scored calcium signals at 0.5 sec before exit were significantly increased as reward volumes increased (F=24.67, p<0.01, n=7, one-way ANOVA, Fig. 2F), but the mean signal curves followed similar trajectories regardless of trial outcome (Fig. 2G). Then, we calculated the slopes of signal curves with different outcomes over 4 time windows (0-2, 2-4, 4-6, 6-8 s) by linear regression analysis. The slopes during the same

time window did not differ significantly between reward groups (0~2s, F=0.10, p=0.96; 2~4s, F=1.03, p=0.38; 4~6s, F=1.00, p=0.34, n=7, one-way ANOVA, Fig. 2H). We pooled and plotted the slopes of different waiting periods together and found the activity curves kept rising steadily, almost saturating after 6 secs from the time the mice entered the waiting zone. The ramp-up of DAergic activity became less variable with delayed gratification task training in our experimental data (figs. S4A-D). All these results indicated that VTA DAergic neurons consistently ramp up their activity during waiting in as animals are trained in the delayed gratification task.

High-frequency spiking of VTA DAergic neurons sustains waiting in the delayed gratification task

Does tonic or phasic firing of DAergic neurons underlie the ramping calcium signal? To answer this, we conducted single unit recordings when mice were performing the delayed gratification task (Fig. 3A). A custom-made head plate was placed on the skull and affixed in place with dental cement. After recording, placements of recording electrodes were confirmed with electrolytic lesion inside VTA all 5 mice (Fig. 3B). We found 17 putative DAergic neurons displayed short bursts of firing during the waiting period (Fig. 3C). On trials in which the mice waited for a short duration the firing rate was low throughout the waiting period (Fig. 3D-E). On trials in which the mice waited for a long duration the firing rate was initially low before increasing during the later waiting period (Fig. 3D-E). We averaged Peri-stimulus time histograms (PSTHs) of all trials to obtain a response curve (Fig. 3F, n=17 cells). Similar to calcium signal, the response curve of firing rate noticeably ramped up with increased waiting

time, before reaching a plateau at around 4 seconds. To compare with the experimental calcium signal, we used a convolution algorithm to predict calcium responses trial by trial based on the firing rate of all 17 recorded cells (Fig. 3G showed the predicted calcium responses from Fig 3E). The average of predicted calcium signal (ΔF/F %, 0.5 sec before exiting from all trials) of all cells fit the measured calcium response (Z-scored ΔF/F, 0.5 sec before exit from the last week of training) curve very well (Fig. 3H, r=0.982, Pearson correlation). This analysis shows that the spiking activity of single DAergic neurons underlies the ramping calcium signal revealed by fiber photometry recording.

Optogenetic manipulation of VTA DAergic activity alters the waiting durations in the delayed gratification task

To determine whether VTA DAergic activity controls performance in the delayed gratification task, we manipulated VTA DAergic neurons temporally using optogenetic tools in 20% of pseudo-randomly chosen trials while the mice were waiting during the delayed gratification task (Figs. 4A-C). Activating the VTA DAergic neurons shifted the cumulative probability distribution towards a longer waiting duration (Fig. 4D, blue, F=12.93, p=0.002, n=6 mice, one-way ANOVA), while inhibiting these same neurons shifted the distribution significantly in the opposite direction (Fig. 4E, yellow, F=7.76, p=0.008, n=6 mice, one-way ANOVA). The effects of optogenetic stimulation or inhibition on the cumulative probability distributions for waiting duration were only observed in the Laser-ON trials. In contrast, the Laser-OFF trials, including those immediately after the Laser-ON trials treated as a single group, were not significantly

different from the trials from the previous day (p>0.5, Figs. 4D-E). The optical manipulation didn't influence the running durations in the delayed gratification task in mice that expressed ChR2 or eNpHR 3.0 (figs. S5A-B), nor did it change the waiting duration distribution of mice that expressed mCherry in DAergic neurons in delayed gratification tasks (figs. S6A-B). To rule out the possibility of optogenetic manipulation-induced memory, we performed a random place preference test with the same stimulation dosage. Neither activating nor inhibiting VTA DAergic neurons significantly changed the transient waiting duration or pattern in the location in which the laser was activated in any of the tested mice (Figs. S5E-H) or in the mCherry expressing controls (figs. S6C-F).

A reinforcement learning (RL) model suggests that ramping up VTA DAergic activity signals the value of waiting for delayed gratification

How does a mouse manage to wait longer for a larger reward while ignoring smaller but more immediate reward options? We propose two models of behavioral scenarios that exemplify possible strategies a mouse may implement to achieve extended waiting performances: (i) setting a goal of expected waiting duration before the initiation of waiting or (ii) continuously deliberating during the waiting period. According to the first hypothesis (i), we modeled a reinforcement learning (RL) agent that keeps time until a predetermined moment has passed (Fig. 5A, *Decision Ahead*); according to (ii), we modeled a second RL agent that continuously balances the values of waiting versus leaving to control the decision to wait or to leave for the reward. To implement these models, we used a version of the state-action-reward-state-action (SARSA) algorithm

with a series of states (Fig. 5A, *Continuous Deliberation*, see methods) (*20, 21*). The

behaviors of both models were able to replicate the behavioral performance we

observed in animal experiments (Figs. 5B-C). The distributions of behavioral

performances between early training days and late training days from the experimental

data were very different from each other, such that the Kullback-Leibler (KL)

divergence was large (0.39 ± 0.06). The KL divergences between the distributions of

simulated behavioral performances and experimental data were significantly small to

the large KL divergence value ($p = 0.005$, $n = 7$ mice, Friedman test), and there was no

difference ($p > 0.99$) between Decision Ahead RL model and Continuous Deliberation

RL model in either the early or late training session. (Fig. 5D). We couldn't determine

which model is better based on behavioral performance alone, given that both models

reproduced the behavioral data well.

What does the ramping up of DAergic activity mean in the delayed gratification

task? We tried to explain it with our RL models. In the *Decision Ahead* model, the agent

keeps time until the predetermined moment has passed, which suggests the ramping

DAergic activity may relate to timing in the delayed gratification task. Some studies

have proposed that the ramping activity is consistent with a role in the classical model

of timing, with the movement being initiated when the ramping activity reaches a nearly

fixed threshold value, following an adjusted slope of ramping activity(*22-25*). In

contrast, our results showed that the DAergic activity ramped up to different values

with similar trajectories on at a nearly constant slope (Figs. 2F-H). This suggests that

VTA DAergic neurons may not implement a decision variable for the *Decision Ahead*

scenario. In the *Continuous Deliberation* RL model, we compared the curves of the

value of waiting and leaving with the ramping up of DAergic activity and found the

behavioral performance of both animals and model agents reached the asymptote. The

values of waiting (Fig. 5E, purple, r = 0.99, Pearson correlation) and the leaving (Fig.

5E, green, r = 0.91, Pearson correlation) each correlated positively with the ramp of

DAergic activity during waiting (Fig. 5E, blue, Z-scored $\Delta F/F$, 0.5 sec before exit from

the last week of training). This detailed analysis suggested that the *Continuous*

Deliberation RL model agreed with previous studies (*13, 26-28*) and that ramping

DAergic activity signals the value of actions, either waiting or leaving, in the delayed

gratification task.

In the *Decision Ahead* RL model, if the agent keeps timing during the waiting

period through ramping DAergic activity to encode the elapsed of time (*29-31*), extra

VTA DAergic activation should represent a longer time, thus leading to earlier

cessation of waiting. This prediction is contrary to our optogenetics result that DAergic

activation led to a longer waiting (Fig. 4D). Instead, we reproduced the optogenetic

manipulations in the *Continuous Deliberation* RL model by either increasing or

decreasing the value of waiting (Q_{wait}) in a pseudo-random 20% of trials. The increase

or decrease in waiting durations only occurred in the Q_{wait} − manipulated trials (p<0.001,

Friedman test, n=10), whereas the remaining trials, including the next trials after value

manipulation, did not differ significantly from controls (p>0.999, Friedman test, n=10,

Figs. 5F-G). Importantly, manipulating the value of leaving (Q_{leave}) in a pseudo-random

20% of trials induced the opposite results, inconsistent with the experimental data (figs.

S8A-B). Manipulating the reward prediction error (RPE) signal in the model led to persistent changes to the waiting time on subsequent trials (figs. S8C-D). This is inconsistent with the experimental finding that optogenetic manipulation of dopamine neuron firing only affects waiting time on the current trial (Figs 4D-E). Thus the ramping dopamine signal does is not consistent with an RPE signal in the delayed gratification task. Our experimental data and the *Continuous Deliberation* RL model together indicated that the ramping up of VTA DAergic activity profoundly influenced the waiting behavioral performance in the delayed gratification task, which suggested that this activity signals the value of waiting, rather than the value of leaving or reward prediction error. Conceptually, our analysis revealed that delayed gratification involves real-time deliberation.

VTA DAergic activity during waiting predicts the behavioral performance in the delayed gratification task

Our optogenetic manipulation experiments and *Continuous Deliberation* RL model indicated that VTA DAergic activity during waiting influenced the waiting durations while the mouse performed the delayed gratification task (Figs. 4D-E and Figs. 5F-G). Although the activity of VTA DAergic neurons ramped up consistently during waiting (Figs. 2G-H), it still fluctuated to a certain extent moment by moment. Therefore, we sought next to determine whether this fluctuation of DAergic activities influences the waiting behavior in the delayed gratification task. A strong prediction made by the *Continuous Deliberation* model is that, if DAergic activity signals the value of waiting at each specific moment, then a momentary increase in the DA signal

will make the agent more likely to keep waiting in the next "time bin," but not in

subsequent time bins (fig. S8E-F). That is to say, the value of waiting is only positively

correlated with the behavior of the immediately following time bin, which indicates the

Markovian nature of the model (*32*). We thus aimed to test the relationship between the

amplitude of the momentary VTA DAergic signal and the behavior (i.e., waiting or

leaving) within each time bin to determine how the momentary DAergic activity (the

calcium signal amplitude in 0-1 sec, 1-2 sec, 2-3 sec, or 3-4 sec after waiting onset,

shown as each cluster of bars in Fig. 6B) affects the waiting performance in the

subsequent periods (behavior from 1-2 sec, 2-3 sec, 3-4 sec, and 4-5 sec for DAergic

activity from 0-1 sec; behavior within 2-3 sec, 3-4 sec and 4-5 sec for DAergic from in

1-2 sec, and so on; Fig. 6A). To integrate data from multiple sessions as well as multiple

animals, we took the advantage of the linear mixed model analysis (LMM, see Method)

(*33-35*). We examined the correlation between momentary DAergic activity and

behavior (i.e. the momentary binary waiting decision). For the behavior, we coded

sustained waiting as 1 and leaving in that time bin as 0 (trials that stopped before the

examined time window were not taken into account) for different pairs of time bins.

Ten independent LMM analyses were done for each activity-behavior pair, as indicated

by the dash lines in Fig. 6A. The regression coefficients, as well as the confidence

intervals of each of the ten pairs, are shown by corresponding bars in Fig. 6B. The

correlation is only significantly positive between adjacent DAergic and waiting bins

(p<0.001, n=7 mouse, black lines, regressed coefficient median; boxes, 50% confidence

interval; whisker, 95% confidence interval, Fig. 6B, the first bar of each cluster, where

the DA activity bin is 1sec ahead of the waiting bin). There was no significant correlation between DAergic activity at 3-4 sec and behavior at 4-5 sec (r=0.007, p=0.61, n=7), which may result from insufficient data for those long trials. Apart from these pairs with adjacent bins, other pairs of DAergic activity and waiting behavior didn't show any significant correlation (Fig. 6B, the remaining bars of each cluster). These results indicate that the waiting decision of the current moment is only influenced by the most recent DAergic signal but not by DAergic signal further in the past, which suggests that deliberation for waiting in delayed gratification may be a Markov process as we formalized in the *Continuous Deliberation* RL model(*32*).

In the *Continuous Deliberation* RL model, the probability of waiting (P_w) positively correlates with the value of waiting (Q_{wait}). To explore the impact of DAergic activity on the probability of waiting in our experimental data, we binned DAergic activity of every trial and normalized data points (V_{DA}) in each momentary DAergic period (with each period lasting 1 sec and starting from 0 to 9 sec, as shown in northeast figure of Fig. 6C). Then, we divided the trials into two groups by setting a series of arbitrary thresholds (red, High DAergic activity, $V_{DA-Z} \geq Th$; green, Low DAergic activity, $V_{DA-Z} \leq -Th$, where *Th* was the threshold for the analysis of high/low DAergic activity) from these trials (Fig. 6C, *Th* was set to 0.9). We analyzed the P_w of low or high DAergic activity trials for the adjacent waiting period with different thresholds. In doing so, we found that the probability of waiting increased rapidly as the absolute value of the threshold increased. The Pw of High DA and Low DA activity trials fit well with a fifth-degree polynomial function (R^2=0.93, -2.1≤ threshold≤2.1). When the absolute

values of the threshold are large enough (|Th|≥1.7), the Pw of the High DA activity

trials is significantly (p=0.04, F(1,12)=5.483, Two-way ANOVA) higher than the Pwait

of the Low DA-ramping activity trials in adjacent waiting periods (|threshold|=2.0,

p=0.02; |threshold|=2.1, p<0.001, Sidak's multiple comparisons test, n=7, Fig. 6D).

Finally, we investigated the influence of fluctuations of intrinsic VTA DAergic

activity on the waiting performance of mice in the delayed gratification task. There

were certain trials in which DAergic activity across the whole duration of waiting was

significantly higher (red, high-ramping) or lower (green, low-ramping) than the mean

DAergic activity (see Methods). We then got two groups of trials and found that the

cumulative distribution of waiting durations in high-ramping trials shifted to the right,

with significantly higher normalized waiting durations (1.03±0.01) compared to those

low-ramping trials (0.98±0.01, p=0.024, n=7, paired Student's t-test, Fig. 6E), but there

was no difference between the normalized waiting durations for the trials immediately

following the high-ramping and low-ramping trials (next trial of H-R, 1.01±0.01; next

trial of L-R,0.99±0.01; p=0.290, n=7, paired Student's t-test, Fig. 6F). These results

accorded with our optogenetic manipulation experiment (Figs. 4D-E), which indicated

that optogenetically manipulated VTA DAergic activity transiently influences the

duration of waiting in the delayed gratification task.

DISCUSSION

Here we reported a novel behavioral task in which the mice were trained to learn

a foraging task with a delayed gratification paradigm. Mice learned to wait for bigger

rewards that they received after waiting for longer (Figs. 1F-H). On a neural level, we found that the calcium signal of VTA DAergic neurons ramped up consistently when mice waited in place before taking action to fetch the expected reward (Figs. 2G-H). Further data analysis showed that the ramping VTA DA activity indeed influenced the behavioral performance of waiting (Figs. 6B-E), which was confirmed with bi-directional optogenetic manipulations of VTA DAergic activity (Figs. 4D-E). Finally, we developed a RL model that predicted our experimental observations well and consolidated the conclusion that the ramping up of VTA DAergic activity signaled the value of waiting in the delayed gratification task, which involved real-time deliberation (Figs. 5B-G).

DA release in the nucleus accumbens (NAc) was previously conjectured to sustain or motivate the goal-directed behavior as well as resistance to distractions (*13, 14*). Here, we explicitly implemented continuous less-optimal options during the delayed gratification process, in which, to achieve better performance, mice needed to sustain waiting as well as prevent/control impulsivity (*3, 6, 36, 37*). We found remarkable and sustained DAergic activation when mice managed to wait longer, and further demonstrated a causal link between DAergic activation and the increase in transient waiting probability. Furthermore, we found DAergic activity ramps up in a consistent manner during waiting, mimicking the value of waiting along with a series of states in our *Continuous Deliberation* RL model, both of which presumably contributed to pursuing a more valuable future goal and resisting the distraction of the less-optimal immediate options in our task. Intriguingly, the momentary DAergic activity was found

to correlate positively with the momentary waiting probability, which also suggested DAergic activity may be involved in the continuous deliberation process. Therefore, we not only for the first time to our knowledge demonstrated the behavioral significance of DAergic activity in delayed gratification, but also depicted "a continuous deliberation" framework in which DAergic activity may participate to help achieve more flexible and sophisticated performance.

Numerous works have used Pavlovian conditioning in studying DAergic activity(*10, 12, 38-40*). Some studies paired the reward with a cue (or cues), such that the animals don't need to perform effortful work to obtain rewards. It is well known that this kind of DAergic activity signals the RPE via phasic firing. In the studies using operant conditioning or goal-directed behavior, the animals have to perform actions and need effortful work to obtain outcomes, and a ramping up of DAergic activity was reported to emerge while the animals were approaching the reward (*13, 14, 41, 42*). This ramping activity was suggested to signal the value of work (*13*) or a distant reward (*14*), but key evidence is lacking because the change of sensory input flow dramatically alters the DAergic activity over time. Under such mutual influence, it is impossible to identify the RPE or the value of work from external cues. The RPE model of ramping activity assumes that the value increases exponentially (or at least in a convex curve) as the reward is approached. Under this model, sensory feedback is suggested to result in the RPE signal ramping up (*41, 43, 44*), while a lack of sensory feedback is predicted to make a flat RPE signal. In contrast, the ramping up of DAergic activity is well isolated from the external sensory inputs when performing the delayed gratification task

in our model. Mice continuously deliberate about the current state and future rewards without any external sensory inputs while waiting in place. Despite the lack of external sensory inputs, we still observed the calcium signal of VTA DAergic neurons ramping up in a consistent manner, functionally mimicking an inner variable of the evolving value of waiting. This observation is consistent with the hypothesis of dopamine signaling the value that is related to time and effort investment under certain circumstances (*13*), but cannot be immediately explained by an RPE response to external sensory inputs.

Midbrain DAergic neurons play an important role in reinforcement learning(*9, 11, 12, 45, 46*), where activation of DAergic neurons usually produces a reinforcement effect on an associated action, stimulus, or place. But in our delayed gratification task, optogenetic manipulation of DAergic activity substantially influenced the ongoing behavior during the current trial without a visible reinforcement effect on later trials. Notably, this optogenetic manipulation was not sufficient to induce a reinforcement effect in the random place performance test. These results revealed the distinct and potent instantaneous effect of DAergic activity during delayed gratification. By simulating transient manipulations of variables in the RL model, we showed that manipulating dopamine activity was equivalent to manipulating the value of waiting in the model. In contrast, manipulating the value of leaving, or the RPE signal itself caused dramatically different effects on behavior. The observations and analysis in our experiments suggest that the value of waiting is represented in VTA DAergic neurons during a delayed gratification task. This significantly updates the understanding of the

coding mechanisms and fundamental functions of the DAergic system in delayed

gratification. Our results suggested that DAergic neuron stimulation during the RPPT

test is not rewarding, but does lead to a shift in wait time. This control experiment was

performed to exclude the rewarding effect of our DAergic stimulation paradigm during

the DG task. However, such a test may be not strictly comparable with stimulation

during the delayed gratification task due to the differences in the behavioral contexts.

The previous finding suggested that reasonable behavior in the face of instant

gratification requires suppression of reflexive reward desiring. Human brain imaging

results demonstrated hemodynamic responses to conditioned (rewarding) stimuli in

both the NAc and the VTA were significantly attenuated during the desire–reason

dilemma(*47*). Such discrepancies with our results may be the consequence of different

behavioral strategies, in which they measured the reward-related activation during a

desire–reason dilemma, and we measured the DA activity during the waiting time. Also,

the different temporal and spatial resolution of human brain imaging and fiber

photometry and electrophysiology may lead to the discrepancies, and our ramping

pattern of DAergic activity also indicated there are inhibitory tones in the NAc and

VTA during the beginning phase of wait.

The design of our delayed gratification task recapitulates the realistic situation

where immediate less-valuable choices lie in the way of pursuing a later but possibly

larger benefit. A deficit in the ability to resist immediate reward for the delayed but

possibly larger reward is closely related to a variety of disorders like obesity, gambling,

and addiction(*1, 48*). The ramping VTA DAergic activity accords with a model of

410   NOW vs LATER decisions in which DAergic signals have a strong influence on the

411   prefrontal cortex in favoring LATER rewards(*2*). We propose that the sustained phasic

412   VTA DAergic activity during the delay period could serve as a neural basis for the

413   power to resist a temptation close at hand, and improve reward rate or goal pursuit in

414   the long run.

MATERIALS AND METHODS

Mice. Animal care and use strictly followed institutional guidelines and governmental regulations. All experimental procedures were approved by the IACUC at the Chinese Institute for Brain Research (Beijing) and ShanghaiTech University. Adult (8-10 weeks) DAT-IRES-Cre knock-in mice (Jax stock# 006660) were trained and recorded. Mice were housed under a reversed 12/12 day/night cycle at 22–25°C with free access to ad libitum rodent food.

Stereotaxic viral injection and optical fiber implantation. After deep anesthesia with isoflurane in oxygen, mice were placed on the stereoscopic positioning instrument. Anesthesia remained constant at 1~1.5% isoflurane supplied per anesthesia nosepiece. The eyes were coated with aureomycin eye cream. The scalp was cut open, and the fascia over the skull was removed with 3% hydrogen peroxide in saline. The Bregma and Lambda points were used to level the mouse head. A small window of 300~500μm in diameter was drilled just above the VTA (AP: -3.10 mm, ML: ±1.15mm, and DV: -4.20 mm) for viral injection and fiber implantation. 300 nl of AAV2/9-hSyn-DIO-GCamp6m (10^12) solution was slowly injected at 30nl /min unilaterally for fiber photometry recording. Either 300 nl of AAV2/9-EF1a-DIO-hChR2(H134R)-mCherry (10^12) or 300 nl of AAV2/9-EF1a-DIO-eNpHR3.0-mCherry (10^12) was injected bilaterally for optogenetic experiments. The injection glass pipette was tilted at an angle of 8° laterally to avoid the central sinus. After injection, the glass pipette was kept in place for 10 min and then slowly withdrawn. An optical fiber (200 μm O.D., 0.37 NA; Anilab) hold in a ceramic ferrule was slowly

inserted into the brain tissue with the tip slightly above the viral injection sites. The

fiber was sealed to the skull with dental cement. Mice were transferred to a warm

blanket for recovery, then housed individually in a new home until all experiments

were done.

Behavioral tasks. One week after surgery, mice started a water restriction schedule to

maintain 85–90% of free-drinking bodyweight for 5 days. The experimenter petted

the mice 5 minutes per day for 3 days in a row and then started task training. All

behavioral tasks were conducted during the dark period of the light/dark cycle.

The foraging task shuttle box had two chambers (10×10×15 cm) connected by a

narrow corridor (45×5×15 cm, Fig. 1A). A water port (1.2 mm O.D. steel tube, 3 cm

above the floor) was attached to the end of one chamber, defined as the reward zone,

with the other as the waiting zone. The position of the mouse in the shuttle box was

tracked online with a custom MATLAB (2016b, MathWorks) program through an

overhead camera (XiangHaoDa, XHD-890B). The experimental procedure control and

behavioral event acquisition were implemented with a custom MATLAB program and

an IC board (Arduino UNO R3).

One-arm foraging task (pre-training)*:* A water-restricted mouse was put in the

shuttle box for free exploration for up to 1 hour. When the animal traveled from the

waiting zone through the corridor to the reward zone to lick the water port, 10 μl water

was delivered by a step motor in 100 ms as a reward. A capacitor sensor monitored the

timing and duration of licking. The animal's return to the waiting zone to initiate the

next trial. Exiting from the waiting zone triggered an auditory cue (200 ms at 4 kHz

sine wave with 90 dB) to signal this exit from the waiting zone. The time spent in the

waiting zone was defined as the waiting duration. The training was conducted every

day for a week. All mice learned to move quickly back and forth between the two

chambers to maximize the reward rate within one week.

Delayed gratification task. From the second week, the volume of water reward was

changed to a function proportional to the waiting time: a wait time of 0-2 s for 0 μl; 2-

4 s triggered delivery of, 2 μl; 4-6 s, 6 μl; 6-8 s, 18 μl; and >8 s, 30 μl as shown in Fig.

1A. The training was conducted five days a week, from Monday to Friday.

P_w Calculation. We divided all trials into two groups, Waiting Trials and Leaving

Trials according to whether the animal remained to wait or left during a given time

interval, such as 1 sec after each behavioral period. Then, we calculated the probability

of waiting (P_w) in this given time interval by the number of 'Waiting Trials' ($N_{w(n)}$) and

the number of 'Leaving Trials' ($N_{L(n)}$) in the time window n:

$$P_{w(n)} = \frac{N_{w(n)}}{N_{w(n)} + N_{L(n)}}$$

Then we could calculate the P_w for a given time duration:

$$P_{w(n)} = \frac{\sum_0^9 N_{w(n)}}{\sum_0^9 N_{w(n)} + \sum_0^9 N_{L(n)}}$$

Linear Mixed Model. We implemented the Linear Mixed Model Analysis using the

open-source   Python   package   "statsmodels"

(https://www.statsmodels.org/stable/mixed_linear.html). The binary value of waiting

or leaving during a specific behavioral period t_{beh} was set as the dependent factor

(t_{beh}=[1, 2), [2, 3), [3, 4), or [4, 5), unit: second); the fluctuation of momentary DA

signal from its mean during a preceding period t_{DA} was set as a fixed-effects (t_{DA}=[0,

1), [1, 2), [2, 3), [3, 4), unit: second. Note that t_{DA} is always smaller than t_{beh}); the animal

identity and session numbers were set as a random effect (n=5 for each animal from the

third week). The parameters of the model were estimated by restricted maximum

likelihood estimation (REML).

Optogenetic stimulation. Lasers, 473 nm wavelength for activation and 589 nm for

inhibition, were coupled to the common end of a patchcord (200 μm O.D., 1-m long,

0.37 NA). The patchcord split through an integrated rotary joint into two ends

connected to optical fibers implanted as described above (200 μm O.D., 0.37 NA) for

bilateral light delivery. First, the mice were trained for 3 weeks to learn the delayed

gratification task. Optical stimulation was delivered pseudo-randomly in ~20% of

behavioral trials in the test experiment. Square pulses of 20 ms at 10 Hz for activation

or a continuous stimulation for inhibition were delivered. The laser was set to ON when

the animal entered the reward zone and to OFF upon exiting from the reward zone. The

maximal laser stimulation was no longer than 16 seconds, even if in the case a mouse

stayed in the waiting zone longer than this time. Continuous laser power at the tip of

the splitting patchcord was about 10 mW for the 473 nm laser and 8 mW for the 589

nm laser, respectively.

Random place performance test. After finishing optogenetic tests for delayed

gratification, all mice took a random place performance test (RPPT). The RPPT was

carried on in a rectangular apparatus consisting of two chambers (30×30×30 cm)

separated by an acrylic board. With an 8 cm wide door open, the mice could move

freely between the two chambers. Before testing, each mouse was placed into the

apparatus for 5-min free exploration. The RPPT consisted of two rounds of 10-min tests.

First, we randomly assigned one chamber as a test chamber. Laser pulses were delivered

with 20% possibility (in accord with the setting of laser pulses delivering in delayed

gratification task) while the mouse entered the test chamber. The delivery of light, no

longer than 16 sec, stopped while the mouse exited the test chamber. Next, we switched

the chamber in which laser pulses were delivered. The laser output power and pulse

length were set the same as in the optogenetic manipulations in the delayed gratification

task. In this task, we analyzed the time in each chamber with or without laser delivered.

Fiber photometry recording. During the behavioral task training and test, we

recorded the fluorescence signal of VTA dopaminergic (DAergic) neurons. The signal

was acquired with a fiber photometry system equipped with a 488 nm excitation laser

and a 505~544 nm emission filter. The GCaMP6m signal was focused on a

photomultiplier tube (R3896 & C7319, Hamamatsu) and then digitalized at 1 kHz and

recorded with a 1401 digitizer and Spike2 software (CED, Cambridge, UK). An

optical fiber (200μm O.D., 0.37 NA, 1.5-m long, Thorlabs) was used to transfer the

excitation and emission light between recording and brain tissue. The laser power

output at the fiber tip was adjusted to 5~10 μW to minimize bleaching.

All data were analyzed with custom programs written in MATLAB (MathWorks).

First, we sorted the continuously recorded data by behavioral trials. For each trial, the

data spanned the range between 1 s before the waiting onset and 2 s after the reward.

Before hooking up the fiber to the mouse, we recorded 20 s of data and averaged as F_b

as the ground baseline. For each trial, we averaged 1-sec of data before the waiting

onset as the baseline F_0 and then calculated its calcium transient as:

526                         $$\Delta F/F\ (\%) = (F-F_0)/(F_0-F_b)\ \times\ 100\ (\%)$$

In the correlation analysis between VTA DAergic activity before waiting and the

waiting duration of mice, we used averaged 1-sec data before the waiting onset as the

DAergic activity before waiting.

In the analysis of high-ramping and low-ramping DAergic activity, we compared

the whole calcium signal of every trial with the average curve (the same length as the

analyzed calcium signal) of all trials from one mouse in a single training day with paired

a t-test and then separated their waiting times into high-ramping group and low-ramping

group.

To facilitate presenting the data, we divided each trial data into four segments,

including 1 s before waiting onset, waiting, running, and 2 s after rewarding. For

comparing the rising trends, we resampled the data segments at 100, 100, 50, and 100

data points, respectively. In the delayed gratification task, the trial data were aligned to

the waiting onset and presented as the mean plots with a shadow area indicating SEM

of fluctuations.

In vivo electrophysiological recording. A custom-made head plate was placed on the

skull of each mouse and affixed in place with dental cement. We removed the skull and

dural carefully above the recording window before the implantation. Stereotrodes were

twined from 12.7 mm Ni-Cr-Fe wires (Stablohm 675, California Fine Wire, CA, USA).

Then 8 stereotrodes were glued together and gold plated to reduce impedance to 250–

500 kΩ. The stereotrodes were gradually lowered to a depth of 0.7 mm above the VTA.

A silver wire (127 mm dia, A-M System) was attached to one of the four skull-

penetrating M1 screws to serve as ground(*19*). Mice were allowed a recovery time of 7

days. Extracellular spiking signals were detected with 8 stereotrodes and amplified

(1000×) through a custom-made 16-channel amplifier with built-in band-pass filters

(0.5–6 kHz). We selected one channel that did not show spike signals and defined it as

a reference ground to reduce movement artifacts. Analog signals were digitized at 25

kHz and sampled by a Power1401 digitizer and Spike2 software (Cambridge Electronic

Design). Spikes recorded by the stereotrode were sorted offline using Spike2 software

(Cambridge Electronic Design). Classified single units should have a high signal-to-

noise ratio (>3:1), reasonable refractory period (interspike interval >1 ms), and

relatively clear boundaries among different PCA clusters. The spike frequency and

waveform were used to determine cell type as the DA or GABA neurons. The putative

DA neurons were identified by their relatively low firing rate (the mean firing

rate<15Hz), and a broad initial positive phase of >1 ms. Then, the spike trains were

aligned with the waiting onset in delayed gratification task. Peri-stimulus time

histograms (PSTHs) (bin width = 100 ms) for each trial were calculated and presented

with averages in plots.

Calcium signal simulation. We performed a simple simulation of the "calcium

fluorescence signal" contributed by each recorded unit using kernel convolution with

Reinforcement learning model. We investigated two potential scenarios. One was

that the mouse decided on a waiting duration before entering the waiting area, and

then waited according to the decided goal. The other scenario was that the mouse

entered the waiting zone, and determined whether to wait or leave as an ongoing

process throughout the whole waiting period. We called these two scenarios

"*Decision Ahead*" and "*Continuous Deliberation*", respectively, and formulated

corresponding reinforcement learning based models for simulation using Python

(Python Software Foundation, version 2.7. Available at https://www.python.org/).

Decision Ahead. Inspired by animal behavior, we simply set three optional "actions"

with different expected waiting durations that could empirically cover the main range

of animals' waiting durations seen during training (T_{a1} = 1.65 sec for action1, T_{a2} = 2.72

sec for action2, T_{a3} = 4.48 sec for action3). These waiting durations were equally spaced

on the log-time axis, consistent with Weber's law (that is, $\ln(T_{a1})$ = 0.5, $\ln(T_{a2})$ = 1,

$\ln(T_{a3})$ = 1.5). During the execution of action a_i, we imposed additional noise on the

timing so that the actual waiting time τ_{ai} for action a_i followed a Gaussian distribution

on the log-time axis centered at the T_{ai}, $\ln \sim \mathcal{N}(\ln(T_{ai}), 0.4^2), i = 1, 2, 3$. These

settings allowed us to best capture the animals' waiting performance in the model. For

each trial, the agent chose an action randomly based on the three action values and a

Boltzmann distribution (Softmax):

$$P_{a_i} = \frac{e^{\beta Q_{a_i}}}{\sum_{j=1,2,3} e^{\beta Q_{a_j}}}$$

Where P_{a_i} was the probability of choosing action a_i and waiting for τ_{ai}. Q_{a_i} was the

value for a_i. β was the inverse temperature constant tuned to 5 to best fit the animal

experimental data. After waiting, the agent would get a reward according to the same

reward schedule used in our experiment. Each action value was updated separately

during the reward delivery:

$$\delta = r - Q_a$$

$$r = R/(\tau + 1)$$

$$Q_a \leftarrow Q_a + \alpha * \delta$$

Where the reward prediction error δ was calculated by the difference between the

hyperbolically discounted reward r (or "reward rate", given by the absolute reward R

dividing total time $\tau+1$ for obtaining the reward, where τ was the waiting duration and

the additional 1sec was the estimated delay of running between the two zones) and the

chosen action value Q_a. The reward prediction error was then used to update the value

of the chosen action. We tuned the learning rate α to 0.002 to fit the animal behavioral

data.

Continuous Deliberation. In each trial, the agent would go through a series of hidden

states, each lasting for 0~2sec randomly according to a Gaussian distribution (mean at

1 sec). At each hidden state, the agent had two action options, either to keep waiting or

to leave. If it chose to keep waiting, the agent would transition to the next hidden state,

with the past time of the previous state cumulated to the whole waiting duration. If the

choice was to leave, the cumulation would cease and a virtual reward dependent on the

duration was delivered; a new trial would then begin from the initial state. The reward

schedule was identical to that used for the animals during the experiments.

The action choice for the future was determined randomly by a Boltzmann

distribution (SoftMax) and action values:

616
$$P_{a_w}^{(T+1)} = \frac{e^{\beta Q_{a_w}^{(T+1)}}}{e^{\beta Q_{a_w}^{(T+1)}} + e^{\beta Q_{a_L}^{(T+1)}}}$$

617     $P_{a_w}^{(T+1)}$ was the probability of choosing to wait for the next state $T + 1$. $Q_{a_w}^{(T+1)}$

618   and $Q_{a_L}^{(T+1)}$ were the value of waiting and leaving, respectively, for state $T + 1$. β

619   was the inverse temperature constant tuned to 5 to best fit the animal experimental

620   data. The action values for each hidden state T were updated by a temporal

621   difference learning algorithm (SARSA):

622
$$\delta = r + \gamma * Q_{a'}^{(T+1)} - Q_a^{(T)}$$

623
$$r = R/(\tau + 1)$$

624
$$Q_a^{(T)} \leftarrow Q_a^{(T)} + \alpha * \delta$$

Where the future action a' was determined by the Boltzmann distribution in the

previous step. The current action a and the future action a' could both be either

waiting or leaving. The prediction error δ was calculated by the sum of reward rate r

(r remained zero until the reward R was delivered. $\tau+1$ was the total time for obtaining

the reward, where τ was the waiting duration and the additional 1sec was the

estimated delay of running between the two zones) and the future action value $\gamma *$

$Q_{a\prime}^{(T+1)}$ discounted by γ ($\gamma = 0.9$), minus the current action value $Q_a^{(T)}$. When a was

leaving, the future action value $Q_{a\prime}^{(T+1)}$ would always be zero. This error signal δ

was used to update $Q_a^{(T)}$ with the learning rate $\alpha = 0.001$.

As a Markovian process, each state would be identical to the agent no matter

how the state was reached or what the following actions might be. So, we extracted the

learned value of waiting as a time series along all the hidden states to compare with the

averaged curve of VTA DAergic activity. For each trial, we also extracted the time

series of the transient waiting value for a trial-wise analysis. Apart from the value of

waiting, we could also extract the time series of RPE for each trial.

We simulated optogenetic manipulation in the model after normal training was

accomplished as in the animal experiments.

Value manipulation. In 20% of trials in the simulation session, the future waiting value

throughout the whole waiting period was manipulated. The optogenetics activation was

simulated as an extra positive value added onto the future waiting value, and the

optogenetics inhibition corresponded to a proportional decrease of the future waiting

value as follows:

647
$$Q_{a_w}^{(T+1)} \leftarrow \tilde{Q}_{a_w}^{(T+1)}, \quad for\ the\ current\ trial$$

$$where \; \tilde{Q}_{a_w}^{(T+1)} = \begin{cases} Q_{a_w}^{(T+1)} + \Delta_{value-ext}, & if \; "ChR2 - lighton" \\ \kappa_{value-inh} Q_{a_w}^{(T+1)}, & if \; "eNPHR - lighton" \end{cases}$$

and, $\delta = r + \gamma * \tilde{Q}_{a_w}^{(T+1)} - Q_a^{(T)}, \quad if \; a' = a_w$

Here we set $\Delta_{value-ext} = 0.15$, and $\kappa_{value-inh} = 0.9$, so that the change in

averaged waiting duration in the simulated "light-on" trials could capture the magnitude

of the instantaneous effect of optogenetic stimulations on the current trials. Using these

parameters "calibrated" by the current trial effect, we were able to compare the

stimulation effect on the light-off or the following trials in both real and simulated

situations. Also note that if the future action was chosen as waiting, the manipulated

value of waiting would be used in the RPE calculation, and thus current action value

updating as well.

RPE manipulation. Under this situation, in 20% of trials in the stimulation session,

instead of the future waiting value, RPE (δ) was manipulated throughout the whole

waiting period as follows:

$$\tilde{\delta} = \begin{cases} \delta + \Delta_{RPE-ext}, & if \; "ChR2 - lighton" \\ \delta - \Delta_{RPE-inh}, & if \; "eNPHR - lighton" \end{cases}$$

and, $Q_a^{(T)} \leftarrow Q_a^{(T)} + \alpha * \tilde{\delta}$

we set $\Delta_{RPE-ext} = 15$, and $\Delta_{RPE-inh} = 20$, which was calibrated by the current trial

effect of real light stimulation.

To simulate the fluctuation in real DAergic signal, we simply multiplied the future

waiting value during each state by a factor of $\sigma \sim \mathcal{N}(1, 0.3^2)$ (determined by the

averaged signal-dependent noise magnitude / relative standard deviation for all

momentary DAergic amplitudes), as an addition to the original model (this is only

implemented for Figs. S8E~F).

Electrophysiological recordings. Adult (8-10 weeks) DAT-IRES-Cre knock-in male

mice 4 weeks after injection with AAV2/9-EF1a-DIO-ChR2(H134R)-mCherry or

AAV-DIO-eNpHR3.0-mCherry were anesthetized with an intraperitoneal injection of

pentobarbital (100 mg kg^{-1}) and then perfused transcardially with ice-cold oxygenated

(95% O_2/5% CO_2) NMDG ACSF solution (93 mM NMDG, 93 mM HCl, 2.5 mM

KCl, 1.25 mM NaH_2PO_4, 10 mM $MgSO_4 \cdot 7H_2O$, 30 mM $NaHCO_3$, 25 mM glucose,

20 mM HEPES, 5 mM sodium ascorbate, 3 mM sodium pyruvate, and 2 mM

thiourea, pH 7.4, 295-305 mOsm). After perfusion, the brain was rapidly dissected out

and immediately transferred into an ice-cold oxygenated NMDG ACSF solution.

Then the brain tissue was sectioned into slices horizontally at 280 mm in the same

buffer with a vibratome (VT-1200 S, Leica). The brain slices containing the VTA

were incubated in oxygenated NMDG ACSF at 32℃ for 10~15 min, then transferred

to a normal oxygenated solution of ACSF (126 mM NaCl, 2.5 mM KCl, 1.25 mM

NaH_2PO_4, 2 mM $MgSO_4 \cdot 7H_2O$, 10 mM Glucose, 26 mM $NaHCO_3$, 2 mM $CaCl_2$) at

room temperature for 1h. A slice was then transferred to the recording chamber, then

submerged and superfused with ACSF at a rate of 3 ml/min at 28℃. Cells were

visualized using infrared DIC and fluorescence microscopy (BX51, Olympus). VTA

DAergic neurons were identified by their fluorescence and other electrophysiological

characteristics. Whole-cell current-clamp recordings of VTA DAergic neurons were

made using a MultiClamp 700B amplifier and Digidata 1440A interface (Molecular

Devices). Patch electrodes (3-5 MΩ) were backfilled with internal solution containing

(in mM): 130 K-gluconate, 8 NaCl, 10 HEPES, 1 EGTA, 2 Mg·ATP and 0.2

Na$_3$·GTP (pH:7.2, 280 mOsm). Series resistance was monitored throughout the

experiments. For optogenetic activation, blue light was delivered onto the slice

through a 200-μm optical fiber attached to a 470 nm LED light source (Thorlabs,

USA). The functional potency of the ChR2-expressing virus was validated by

measuring the number of action potentials elicited in VTA DAergic neurons using

blue light stimulation (20 ms, 10 Hz, 2.7 mW) in VTA slices. For optogenetic

inhibition, yellow light (0.7 mW) was generated by a 590 nm LED light source

(Thorlabs, USA) and delivered to VTA DAergic neurons expressing eNpHR3.0

through a 200-μm optical fiber. To assure eNpHR-induced neuronal inhibition,

whole-cell recordings were carried out in current-clamp mode and spikes were

induced by current injection (200 pA) with the presence of yellow light. Data were

filtered at 2 kHz, digitized at 10 kHz, and acquired using pClamp10 software

(Molecular Devices).

Immunostaining. Mice were deeply anesthetized with pentobarbital (100 mg/kg, i.p.),

followed by saline perfusion through the heart. After blood was drained out, 4%

paraformaldehyde (PFA) was used for fixation. Upon decapitation, the head was

soaked in 4% PFA at room temperature overnight. The brain was harvested the next

day, post-fixed overnight in 4% PFA at 4°C, and transferred to 30% sucrose in 0.1 M

PBS, pH 7.4 for 24~48 h. Coronal sections (20 μm) containing the VTA were cut on a

cryostat (Leica CM3050 S). The slides were washed with 0.1 M PBS, pH 7.4,

incubated in blocking buffer (0.3% Triton X-100, 5% bovine serum albumin in 0.1 M

PBS, pH 7.4) for an hour, and then transferred into the primary antibody (rabbit anti-

tyrosine hydroxylase antibody, 1:1,000; Invitrogen) in blocking buffer overnight at

4°C. The sections were washed three times in 0.1 M PBS, then incubated with donkey

anti-rabbit IgG H&L secondary antibody (conjugated to fluor-488 or fluor-594,

1:1,000; Jackson ImmunoResearch) at room temperature for 2 h. The nucleus was

stained with DAPI (4',6-diamidine-2-phenylindole). Sections were mounted in

glycerine and covered with coverslips sealed in place. Fluorescent images were

collected using a Zeiss confocal microscope (LSM 880).

Quantification and statistics. All statistics were performed by MATLAB (R2016b,

MathWorks) and Python (V2.7, Python Software Foundation) routines. Data were

judged to be statistically significant when the P-values were less than 0.05. Asterisks

denote statistical significance $*p < 0.05$; $**p < 0.01$; $***p < 0.001$. Unless stated

otherwise, values were presented as Mean $\pm$ s.e.m..

REFERENCES AND NOTES

1.   D. Tomasi, N. D. Volkow, Striatocortical pathway dysfunction in addiction and
obesity: differences and similarities. *Crit Rev Biochem Mol Biol* **48**, 1-19 (2013).
2.   N. D. Volkow, R. D. Baler, NOW vs LATER brain circuits: implications for obesity
and addiction. *Trends Neurosci* **38**, 345-352 (2015).
3.   W. Mischel, Y. Shoda, M. I. Rodriguez, Delay of gratification in children. *Science*
244, 933-938 (1989).
4.   J. E. Maddux, J. P. Tangney, *Social psychological foundations of clinical*
psychology (Guilford Press, New York, 2010), pp. xv, 555 p.
5.   J. Grosch, A. Neuringer, Self-control in pigeons under the Mischel paradigm. *J*
Exp Anal Behav **35**, 3-21 (1981).
6.   B. Reynolds, H. de Wit, J. B. Richards, Delay of gratification and delay
discounting in rats. *Behav Process* **59**, 157-168 (2002).

7.   B. Engelhard, J. Finkelstein, J. Cox, W. Fleming, H. J. Jang, S. Ornelas, S. A. Koay,
740   S. Y. Thiberge, N. D. Daw, D. W. Tank, I. B. Witten, Specialized coding of sensory,
741   motor and cognitive variables in VTA dopamine neurons. *Nature*,   (2019).
742   8.   C. K. Starkweather, B. M. Babayan, N. Uchida, S. J. Gershman, Dopamine reward
743   prediction errors reflect hidden-state inference across time. *Nat Neurosci* **20**,
744   581-589 (2017).
745   9.   P. W. Glimcher, Understanding dopamine and reinforcement learning: the
746   dopamine reward prediction error hypothesis. *Proc Natl Acad Sci U S A* **108**
747   **Suppl 3**, 15647-15654 (2011).
748   10.   G. Morris, A. Nevet, D. Arkadir, E. Vaadia, H. Bergman, Midbrain dopamine
749   neurons encode decisions for future action. *Nat Neurosci* **9**, 1057-1063 (2006).
750   11.   J. R. Hollerman, W. Schultz, Dopamine neurons report an error in the temporal
751   prediction of reward during learning. *Nat Neurosci* **1**, 304-309 (1998).
752   12.   W. Schultz, P. Dayan, P. R. Montague, A neural substrate of prediction and
753   reward. *Science* **275**, 1593-1599 (1997).
754   13.   A. A. Hamid, J. R. Pettibone, O. S. Mabrouk, V. L. Hetrick, R. Schmidt, C. M.
755   Vander Weele, R. T. Kennedy, B. J. Aragona, J. D. Berke, Mesolimbic dopamine
756   signals the value of work. *Nat Neurosci* **19**, 117-126 (2016).
757   14.   M. W. Howe, P. L. Tierney, S. G. Sandberg, P. E. M. Phillips, A. M. Graybiel,
758   Prolonged dopamine signalling in striatum signals proximity and value of distant
759   rewards. *Nature* **500**, 575-+ (2013).
760   15.   S. Soares, B. V. Atallah, J. J. Paton, Midbrain dopamine neurons control
761   judgment of time. *Science* **354**, 1273-1277 (2016).
762   16.   M. Guitart-Masip, U. R. Beierholm, R. Dolan, E. Duzel, P. Dayan, Vigor in the face
763   of fluctuating rates of reward: an experimental examination. *J Cogn Neurosci* **23**,
764   3933-3938 (2011).
765   17.   Y. Niv, Cost, benefit, tonic, phasic: what do response rates tell us about
766   dopamine and motivation? *Ann N Y Acad Sci* **1104**, 357-376 (2007).
767   18.   Y. Niv, N. D. Daw, P. Dayan, Choice values. *Nat Neurosci* **9**, 987-988 (2006).
768   19.   Y. Li, W. Zhong, D. Wang, Q. Feng, Z. Liu, J. Zhou, C. Jia, F. Hu, J. Zeng, Q. Guo,
769   L. Fu, M. Luo, Serotonin neurons in the dorsal raphe nucleus encode reward
770   signals. *Nat Commun* **7**, 10503 (2016).
771   20.   G. A. Rummery, M. Niranjan, On-Line Q-Learning   Using   Connectionist
772   Systems. *Technical Report   CUED/F-Infeng/TR 166*,   (1994).
773   21.   R. S. Sutton, A. G. Barto, *Reinforcement learning : an introduction* (Adaptive
774   computation and machine learning, MIT Press, Cambridge, Mass., 1998), pp.
775   xviii, 322 p.
776   22.   M. Treisman, Temporal discrimination and the indifference interval. Implications
777   for a model of the "internal clock". *Psychol Monogr* **77**, 1-31 (1963).
778   23.   P. R. Killeen, J. G. Fetterman, A behavioral theory of timing. *Psychol Rev* **95**, 274-
779   295 (1988).
780   24.   W. H. Meck, Neuropharmacology of timing and time perception. *Brain Res Cogn*
781   *Brain Res* **3**, 227-242 (1996).

782   25.   M. Jazayeri, M. N. Shadlen, A Neural Mechanism for Sensing and Reproducing a
783   Time Interval. *Curr Biol* **25**, 2599-2609 (2015).

784   26.   Y. Niv, N. D. Daw, D. Joel, P. Dayan, Tonic dopamine: opportunity costs and the
785   control of response vigor. *Psychopharmacology (Berl)* **191**, 507-520 (2007).

786   27.   R. S. Sutton, A. G. Barto, *Reinforcement learning : an introduction* (Adaptive
787   computation and machine learning series, The MIT Press, Cambridge,
788   Massachusetts, ed. Second edition., 2018), pp. xxii, 526 pages.

789   28.   B. A. Bari, C. D. Grossman, E. E. Lubin, A. E. Rajagopalan, J. I. Cressy, J. Y. Cohen,
790   Stable Representations of Decision Variables for Flexible Behavior. *Neuron*,
791   (2019).

792   29.   P. Simen, F. Balci, L. de Souza, J. D. Cohen, P. Holmes, A model of interval timing
793   by neural integration. *J Neurosci* **31**, 9238-9253 (2011).

794   30.   D. Durstewitz, Self-Organizing Neural Integrator Predicts Interval Times through
795   Climbing Activity. **23**, 5342-5353 (2003).

796   31.   F. Balcı, P. Simen, A decision model of timing. *Current Opinion in Behavioral*
797   *Sciences* **8**, 94-101 (2016).

798   32.   S. I. Gass, C. M. Harris, "Markov propertyMarkov property" in *Encyclopedia of*
799   *Operations Research and Management Science,* S. I. Gass, C. M. Harris, Eds.
800   (Springer US, New York, NY, 2001), pp. 490-490.

801   33.   T. K. Koerner, Y. Zhang, Application of Linear Mixed-Effects Models in Human
802   Neuroscience Research: A Comparison with Pearson Correlation in Two Auditory
803   Electrophysiology Studies. *Brain Sci* **7**, (2017).

804   34.   M. P. Boisgontier, B. Cheval, The anova to mixed model transition. *Neurosci*
805   *Biobehav Rev* **68**, 1004-1005 (2016).

806   35.   S. N. Chettih, C. D. Harvey, Single-neuron perturbations reveal feature-specific
807   competition in V1. *Nature* **567**, 334-340 (2019).

808   36.   K. Jimura, M. S. Chushak, T. S. Braver, Impulsivity and self-control during
809   intertemporal decision making linked to the neural dynamics of reward value
810   representation. *J Neurosci* **33**, 344-357 (2013).

811   37.   B. Schmidt, C. B. Holroyd, S. Debener, J. Hewig, Why Is It So Hard to Wait? Brain
812   Responses to Delayed Gratification Predict Impulsivity and Self-Control.
813   *Psychophysiology* **52**, S42-S42 (2015).

814   38.   C. D. Fiorillo, W. T. Newsome, W. Schultz, The temporal precision of reward
815   prediction in dopamine neurons. *Nat Neurosci* **11**, 966-973 (2008).

816   39.   B. M. Babayan, N. Uchida, S. J. Gershman, Belief state representation in the
817   dopamine system. *Nat Commun* **9**, 1891 (2018).

818   40.   J. Y. Cohen, S. Haesler, L. Vong, B. B. Lowell, N. Uchida, Neuron-type-specific
819   signals for reward and punishment in the ventral tegmental area. *Nature* **482**,
820   85-88 (2012).

821   41.   J. G. Mikhael, H. R. Kim, N. Uchida, S. J. Gershman, Ramping and State
822   Uncertainty in the Dopamine Signal. 805366 (2019).

823   42.   A. Guru, C. Seo, R. J. Post, D. S. Kullakanda, J. A. Schaffer, M. R. Warden,
824   Ramping activity in midbrain dopamine neurons signifies the use of a cognitive
825   map. 2020.2005.2021.108886 (2020).

43. S. J. Gershman, Dopamine ramps are a consequence of reward prediction errors. *Neural Comput* **26**, 467-471 (2014).

44. H. R. Kim, A. N. Malik, J. G. Mikhael, P. Bech, I. Tsutsui-Kimura, F. Sun, Y. Zhang, Y. Li, M. Watabe-Uchida, S. J. Gershman, N. Uchida, A Unified Framework for Dopamine Signals across Timescales. *Cell* **183**, 1600-1616 e1625 (2020).

45. W. X. Pan, R. Schmidt, J. R. Wickens, B. I. Hyland, Dopamine cells respond to predicted events during classical conditioning: evidence for eligibility traces in the reward-learning network. *J Neurosci* **25**, 6235-6242 (2005).

46. H. C. Tsai, F. Zhang, A. Adamantidis, G. D. Stuber, A. Bonci, L. de Lecea, K. Deisseroth, Phasic firing in dopaminergic neurons is sufficient for behavioral conditioning. *Science* **324**, 1080-1084 (2009).

47. E. K. Diekhof, O. Gruber, When Desire Collides with Reason: Functional Interactions between Anteroventral Prefrontal Cortex and Nucleus Accumbens Underlie the Human Ability to Resist Impulsive Desires. *Journal of Neuroscience* **30**, 1488-1493 (2010).

48. A. E. Goudriaan, M. Yucel, R. J. van Holst, Getting a grip on problem gambling: what can neuroscience tell us? *Front Behav Neurosci* **8**, 141 (2014).

49. M. Pachitariu, C. Stringer, K. D. Harris, Robustness of Spike Deconvolution for Neuronal Calcium Imaging. *J Neurosci* **38**, 7976-7985 (2018).

50. T. W. Chen, T. J. Wardill, Y. Sun, S. R. Pulver, S. L. Renninger, A. Baohan, E. R. Schreiter, R. A. Kerr, M. B. Orger, V. Jayaraman, L. L. Looger, K. Svoboda, D. S. Kim, Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature* **499**, 295-300 (2013).

Acknowledgments: We thank Drs. M. Lou, W. Ge, Y. Rao, W Zhou, and B Min for

comments on the manuscript. We thank Dr. L. Lu for the help of in vivo electrophysiology

recording. We thank the Molecular Imaging Core Facility (MICF) at the School of Life

Science and Technology, ShanghaiTech University for providing technical support. This

work was supported by the National Natural Science Foundation of China (grant nos.

31922029, 31671086, 61890951, and 61890950 to J.H.), and a Shanghai Pujiang Talent

Award (grant no. 2018X0302-101-01 to W.S.). **Author contributions:** W.S. and J. H.

oversaw the whole project. W.S., J. H. & Z. G. designed the experiments. Z. G., C. L. & T.

L.. performed all animal experiments. Z. G. & H. W. analyzed the data. H. W. & S. F.

performed the computational modeling under the supervision of X. J. W.. M. C. performed

the electrophysiological recordings. W.S., J. H., Z. G. & H. W. wrote the paper with the

participation of all other authors. **Competing interests:** The authors declare no competing

interests. **Data and Materials Availability:** All data needed to evaluate the conclusions in

the paper are present in the paper and/or the Supplementary Materials.

Fig. 1. The behavioral performance of mice during learning of a delayed gratification task learning. (**A**) Left panel, schematic of the delayed gratification task. Right panel, the relationship between reward volumes and waiting durations in the two behavioral tasks. (**B**) A plot of Transistor-Transistor Logic (TTL) signals for the chronological sequence of behavioral events in the tasks. (**C-E**). The waiting duration and running duration both decreased during the training process in the pre-training phase (p<0.001). (**F**) The distribution of waiting durations from the behavioral session on the last analyzed day (Day 15, light red), revealing significantly longer waiting durations compared to those from Day 1 (Day 1, gray, n=7 mice). (**G**) The distribution

875   of running durations on Day1 and Day15 did not differ with training. **(H)** Continuous

876   training significantly increased the average waiting duration (p<0.001), whereas the

877   training did not change the average running duration (p = 0.97).

Fig. 2. VTA DAergic activity ramps up consistently while the mice are waiting for the reward. (**A**) Schematic of stereotaxic virus injection procedures. (**B**) Confocal images illustrating GCaMP6m expression in VTA TH[+] neurons. (**C**) A live-recording trace (magenta) of Ca^{2+} signal in VTA DAergic neurons and running speed (black) when the mouse was performing the delayed gratification task. Task events over time (top): the dashed vertical lines indicate waiting onset (blue), waiting termination (green), and reward onset (red). (**D**) The scaled Ca^{2+} signals (magenta) and GFP signals (green) curves of VTA DAergic neurons from the last day of pre-training and Days in the

delayed gratification task training (black, speed). (**E**) Waiting duration sorted ramping

Ca^{2+} signal data from one mouse during the delayed gratification task training (150

trials). (**F**) Z-scored $\Delta F/F$ values at 0.5s before exit were significantly different when

the reward volumes were different ($p<0.01$). (**G**) Averaged Ca^{2+} signal curves with

different outcomes from (**E**). (**H**) Shown separately for different trial stages (DAergic

ramping periods), during the last week of training. There were no differences in the

slope of the Ca^{2+} signals between trials with different reward outcomes.

Fig. 3. Single unit recordings reveal that high-frequency spiking of VTA DAergic

neurons sustains waiting in the delayed gratification task. (**A**) Schematic of single

unit electrical recordings in the delayed gratification task. (**B**) An example image showing the placement of electrode tips in VTA with an electrolytic lesion. **(C)**. An example recording trace when a mouse performed a whole trial of the delayed gratification task. The waiting period was noted with a solid magenta line. **(D)** Raster plot of spikes trial by trial sorted in the waiting duration of the delayed gratification task. **(E)** A 3D PSTH plot for D. **(F)** Average response curve during the waiting period (n = 17 cells, each dashed line represents one cell). **(G)** Predicted calcium responses based on the convolution of the spike rate from (E) with a 0.1s time bin. **(H)** The predicted calcium response curve (black) is based on the spiking response curves in (F), compared to the measured calcium response curve (blue line) (r=0.982, Pearson correlation).

Fig. 4. Optogenetic manipulation of VTA DAergic activity altered the waiting

durations. (**A**) Left: schematic of stereotaxic virus injection and surgical procedure.

Right: the behavioral events and optogenetic manipulation protocol. (**B**) Images

showing ChR2-mCherry expression (Top),. And whole-cell recordings showing action

potentials evoked by 10-Hz laser flash sequences (50 flashes) in VTA TH+ neurons

(Bottom). (**C**) Images showing eNpHR3.0-mCherry expression (Top), and whole-cell

recordings showing that evoked action potentials were inhibited by continued 589nm

laser in VTA TH+ neurons (Bottom). (**D**) Cumulative probabilities of waiting durations.

The waiting durations of optogenetically activated trials were significantly increased

(p=0.002, blue) relative to those of the previous day's trials (gray). Inset: a bar graph of

the normalized waiting durations showing that the average waiting duration of

unstimulated trials (p=0.96, magenta) did not differ from those of the previous day's

trials, or the next trials following photoactivation (p=0.63, green). (**E**) The same

experimental configuration as in (**D**), but VAT TH⁺ neurons were optogenetically

inhibited by a yellow laser. Optogenetic inhibition decreased the waiting duration

(p=0.008, yellow), whereas there were no differences between other trials (p>0.5).

Fig. 5. Behavioral performances and ramping VTA DAergic activity are explained

by the RL model. (**A**) Two reinforcement learning (RL) models: *Decision Ahead* and

Continuous Deliberation. $Q(a_n)$: value for action a(n); $P(T_{an})$: probability of action a(n).

$P_{wait}^{(n)}$: the probability of waiting; $Q_{wait}^{(n)}$: waiting action value; $P_{leave}^{(n)}$: probability of

leaving; $Q_{leave}^{(n)}$: leaving action value; $R^{(n)}$: received reward. (**B-C**) The distributions of

waiting durations from the early session and late session were simulated in the *Decision*

Ahead model (**B**) and *Continuous Deliberation* model (**C**). (**D**) The Kullback-Leibler

(KL) divergence between the distribution of early and late experimental data was

significantly larger than the KL divergences between the distributions of simulated

behavioral performances and experimental data (p = 0.005), but no difference (p > 0.99)

between the two RL models. (**E**) Plots of Z-scored ΔF/F values (DA$_w$, blue), scaled

Q$_{wait}$ (purple), and Q$_{leave}$ (green) from the *Continuous Deliberation* model. The Q$_{wait}$

and Q$_{leave}$ both predicted the experimental calcium activity well. (**F-G**) Computational

RL model (continuous deliberation)-simulated data, dependent on manipulating Q$_{wait}$

in delayed gratification task. The simulation showed that manipulating Q$_{wait}$ (i.e.,

simulated dopamine activity) only influenced the waiting durations of Q$_{wait}$

manipulated trials (**(F-G)**, p<0.001).

Fig. 6. VTA DAergic activity during waiting predicts the behavioral performance

in the delayed gratification task. (**A**) Schematic of waiting probability (P_{wait}) in

waiting periods after momentary DAergic periods. (**B**) Relationship between

momentary VTA DAergic activity (Ca^{2+} signals) and its waiting probability. For each

momentary DAergic period, its DAergic activity is only highly correlated ($p<0.001$)

with P_{wait} in the adjacent waiting period (the left bar of each cluster). (**C**) The

distribution of Z-scored mean $\Delta F/F$ of momentary DAergic periods. Three colors are

used to illustrate: high dopamine activity (High DA: red,), low dopamine activity (Low

DA: green), and other activity (gray). (**D**) The P_{wait} of High DA and Low DA activity

trials for the adjacent period after the momentary DA periods. The difference between

the P_w of the High DA trials and Low DA trials increases with the increase of threshold

(Th). (**E-F**) Cumulative probabilities of waiting durations for the high-DA-ramping

trials (**E**, H-R, red), the lower-DA-ramping trials (**E**, L-R, green) and their "next-in-

series" trials (**F**). Bar graph showing that the normalized waiting durations of the H-R

trials are significantly longer than those of the L-R trials ((**E**), p=0.024), but did not

differ significantly between their "next-in-series" trials ((**F**), p=0.290).

Fig. S1. The change of behavioral performance and reward rate during behavioral training. (**A**) The KL divergence of the distribution of waiting duration and running duration between the first day and the last day in pre-task and delayed gratification (DG). The KL divergence of waiting duration have no difference between pre-task and delayed gratification ($p = 0.53$, paired t-test), but KL divergence of running duration in pre-task was significantly ($p = 0.002$, paired t-test) different with the KL divergence of running duration from (**B**) The slopes of curves (Fig.1 E and Fig.1 H) that waiting duration and running duration changed during behavioral training. The waiting duration and running duration in pre-task were both changed more quickly ($p = 0.004$, waiting duration; $p<0.001$, running duration, paired t-test) than those from delayed gratification. (**C-D**) The reward rate both increased in pre-task training (**C**) and delayed gratification training (**D**). All error bars represent the s.e.m..

Fig. S2. The histology and tip positions of optical fibre in GCaMP6m (green, n = 7), GFP (blue, n = 4), ChR2 (red, n = 12), ENpHR (yellow, n = 12), mCherry (magenta, n = 7) and electrolytic lesion (black, n = 5) were shown as coordinates in the mouse brain atlas. The circle markers showed the fiber placements from the mouse with optogenetic manipulation in the early training process. Scale bar 500um.

Fig. S3. The calcium and GFP signals of VTA DA neurons in behavioral tasks. (**A**) The GFP (green), calcium signals (magenta) of VTA DA neurons, and mouse running speed (black) in 7 days of pre-training. (**B**) The calcium signals of VTA DA neurons were changing with pre-training progress. The max calcium signals after the mouse received water rewards significantly decreased on days 6&7 (p<0.01, Friedman test). All error bars represent the s.e.m..

Fig. S4. The ramping dopamine activity became more stable with delayed gratification training. (**A**) The Frequency distribution of the slope of Z-scored ΔF/F during waiting (gray: day 1, red: day 15, n = 7). (**B**) The slope of ΔF/F had no difference between day 1 and day 15 (p=0.63, paired Student's t-test). (**C**) The standard deviation(σ) of the slope of ΔF/F in day 15 (0.32±0.07) was significantly decreased than that in day 1 (0.17±0.05, p=0.03, paired Student's t-test). (**D-E**) The z-scored σ of ΔF/F and z-scored waiting durations were negatively correlated both in the experimental data (d, blue, r = -0.80, p<0.001) and RL model (e, red, r = -0.76, p<0.001). W.D. is for waiting duration. All error bars represent the s.e.m..

Fig. S5. Optical activation or inhibition of VTA DA activity didn't change the motivation of mouse in the delayed gratification tasks, the waiting duration of mouse in the early process of delayed gratification training, or the duration the mouse stayed in the given box in RPPT. (**A**) The running duration didn't change while the VTA DA neurons were optically activated (F=0.20, p=0.82, one-way ANOVA). (**B**) The result was the same as Figure A while optical inhibiting the VTA DA neurons (F=0.12, p=0.88, one-way ANOVA). (**C**) The waiting duration didn't change (p = 0.37, laser-on vs laser-off) while the VTA DA neurons were optically activated in the early training process (F=1.31, p=0.30, one-way ANOVA). (**D**) The result was the same (p = 0.59, laser-on vs laser-off) as Figure C while optical inhibiting the VTA DA neurons (F=4.97, p=0.22, one-way ANOVA). (**E**) The heatmap of mouse traces in RPPT in which the VTA DA neurons were optically activated pseudo-randomly in 20% probability while the mouse entered into the given box (red rectangle). (**F**) The Z-scored duration that the mouse stayed in the given box while the VTA DA neurons were activated (Laser-On In) had no significant difference (F=0.75, p=0.44, one-way ANOVA, n=6) with the un-inhibited durations (Laser-Off In) and durations in another box (Out). (**G**) The heatmap of mouse traces as shown in Figure C while inhibiting the VTA DA neurons in the given box (red rectangle). (**H**) Optical inhibiting the VTA DA neurons also didn't change the duration the mouse stayed in the given box (F = 0.17, p = 0.73, one-way ANOVA, n=6). All error bars represent the s.e.m..

Fig. S6. Optogenetic manipulation of DAT-Cre mouse expressed mCherry in the delayed gratification tasks and RPPT.
(**A**) Waiting durations in 473nm laser delivered trials (blue) are not different compared with those of all other trials (p=0.17, Friedman test, n=7). (**B**) Waiting durations of 589nm laser un-delivered trials (magenta) slightly increased compared with the waiting duration of the previous day (p=0.02, Friedman test, n=7). (**C**) Heat-map of mouse traces in RPPT in which the VTA of the mouse was delivered 473nm laser pseudo-randomly in 20% probability while the mouse entered into a randomly chosen box (red rectangle). (**D**) Mean durations that the mouse stayed in the chosen box while the laser delivered (Laser-On In), laser off (Laser-Off In) and the other box. There is no significant difference in waiting duration between Laser-On In, and Laser-Off In (F=3.54, p=0.09, one-way ANOVA, n=7). (**E**) Heatmap of mouse traces same as shown in (c) while the mouse was delivered 589nm laser in a randomly chosen box (red rectangle). (**F**) Mean durations that the mouse stayed in the chosen box while 589nm laser delivered (Laser-On In), laser off (Laser-Off In), and in the other box. 589nm laser delivering to mCherry mouse didn't alter waiting duration mouse stayed in any boxes under all experimental conditions (F=2.64, p=0.14, one-way ANOVA, n=7). All error bars represent the s.e.m..

Fig. S7. Reward prediction error is not significantly correlated with ramping dopaminergic activity. (**A**) The correlation coefficient of mean DA activity 1 s before waiting (DA_b), the value of action (Q_b), and RPE of action (RPE_b) before waiting in the Decision Ahead model, with waiting durations. The correlation of V_b (r = 0.35±0.01, p=0.001, n=10, Pearson correlation) and RPE_b (r = 0.34±0.01, p=0.001, n=10, Pearson correlation) with waiting duration were significantly (p<0.001, Kruskal-Wallish test) higher than the CC of DA_b (r=0.01±0.02, p=0.36, n=7) with waiting durations. (**B**) Plots of Z-scored ΔF/F values (DA_w, light blue) at 0.5s before the waiting ended and RPE of waiting (RPE_w). There were no significant correlation between DA_w and RPE_w (r = 0.34, p = 0.41, Pearson correlation). (**C-D**) Value of waiting (Q_w) (**C**) and RPE (**D**) changed in the Continuous Deliberation model. All error bars represent the s.e.m..

Fig. S8. Manipulation of the value of leaving and RPE in RL model. (**A**-**B**) Either increasing or decreasing the value of leaving (Q_{leave}) in the continuous deliberation model as with the manipulation of Q_{wait} induced the opposite results compared with the optogenetics manipulating DAergic activity (increasing Q_{leave} in **A**: p<0.001, Friedman test, n=10; decreasing Q_{leave} in **B**: p <0.001, Friedman test, n=10) and had no influences on other trials (**A**-**B**, p>0.999, Friedman test, n=10). (**C**-**D**) Either increasing (**C**) or decreasing (**D**) the RPE in the continuous deliberation model as with the experimental data, alters the waiting durations in the same direction in all trials, whether or not the RPEs-manipulation (increasing RPEs in c: p<0.001, Friedman test, n=10; decreasing RPE in (**D**): p <0.001, Friedman test, n=10). (**E**) The value of waiting is only positively correlated (0.23±0.02, p =0.03±0.01, n =20) with the adjacent behavior in the *Continuous Deliberation* RL model. (**F**) The p values of correlation coefficients in **E**. All error bars represent the s.e.m..