Corticosterone differentially modulates time-dependent fear generalization following mild or moderate fear conditioning training in rats

Moisés dos Santos Corrêa[1], Barbara dos Santos Vaz[1], Beatriz Scazufca Menezes[1], Tatiana Lima Ferreira[1], Paula Ayako Tiba[1], Raquel Vecchio Fornari[1]

[1] Center for Mathematics, Computing and Cognition (CMCC), Universidade Federal do ABC (UFABC), São Bernardo do Campo, SP, Brazil

Authors' e-mails and ORCID:

Moisés dos Santos Corrêa: mscorrea.86@gmail.com - 0000-0002-4488-0922

Barbara dos Santos Vaz: barbara.vaz@aluno.ufabc.edu.br - 0000-0002-1634-5279

Beatriz Scazufca Menezes: beatrizscazufca@gmail.com - 0000-0002-1227-3570

Tatiana L. Ferreira: tatiana.ferreira@ufabc.edu.br - 0000-0002-2805-513X

Paula A. Tiba: paula.tiba@ufabc.edu.br - 0000-0001-8030-9232

Raquel V. Fornari (correspondence): raquel.fornari@ufabc.edu.br - Orcid: 0000-0003-1330-6263

Color version only on online version, in print black and white

HIGHLIGHTS:

- Rats were trained in a contextual fear task and tested for fear generalization.
- Corticosterone (CORT) injected after a mild training promoted accuracy up to 28 days.
- CORT injected after a moderate training facilitated time-dependent fear generalization.

ABSTRACT:

Stressful and emotionally arousing experiences create strong memories that seem to lose specificity

over time. It is uncertain, however, how the stress system contributes to the phenomenon of time-

dependent fear generalization. Here, we investigated whether post-training corticosterone (CORT-

HBC) injections, given after different training intensities, affect contextual fear memory specificity at

several time points. We trained male Wistar rats on the contextual fear conditioning (CFC) task using

two footshock intensities (mild CFC, 3 footshocks of 0.3 mA, or moderate CFC, 3x 0.6 mA) and

immediately after the training session we administered CORT-HBC systemically. We first tested the

animals in a novel context and then in the training context at different intervals following training (2,

14, 28 or 42 days). By measuring freezing in the novel context and then contrasting freezing times

shown in both contexts, we inferred contextual fear generalization for each rat, classifying them into

Generalizers or Discriminators. Following mild CFC training, the glucocorticoid injection promoted

an accurate contextual memory at the recent time point (2 days), and increase the contextual memory

accuracy 28 days after training. In contrast, after the moderate CFC training, CORT-HBC facilitated

contextual generalization at 14 days, compared to the control group that maintained contextual

discrimination at this timepoint. For this training intensity, however, CORT-HBC did not have any

effect on recent memory specificity. These findings indicate that treatment with CORT-HBC

immediately after the encoding of mild or moderately arousing experiences may differentially

modulate memory consolidation and time-dependent fear generalization.

KEYWORDS:

INTRODUCTION

Stressful events elicit the formation of lasting and intense memories and their effects on

memory is thought to be exerted by temporally contiguous activation of the noradrenergic and

glucocorticoid systems (Joëls et al., 2011; Vogel et al., 2016). Even though these two hormonal

systems have been long associated with the modulation of the strength of fear memories (de

Quervain et al., 2016; Roozendaal et al., 2009), their role on memory specificity has only

recently become a new focus of investigation. Moreover, overgeneralization of fear is associated

with anxious and phobic behavior and is one of the main symptoms of several stress-related

disorders, such as post-traumatic stress disorder (PTSD), generalized anxiety disorder and

phobias (Lenaert et al., 2015; Lissek et al., 2014b, 2014a). Considering the widespread use of

corticosteroids for the treatment of a myriad of diseases (Rice et al., 2017), understanding how

stress modifies not only memory strength but also memory specificity has deep implications in

the prevention of these disorders and the development of better treatments for them.

A growing body of evidence suggests that noradrenaline and corticosterone (CORT)

modulate memory acquisition and consolidation by shifting from cognitive detailed contextual

learning towards a more rigid memory system, associated with habit and stimulus response

memories (Schwabe, 2017). Such evidence suggests that generalized fear is associated with this

rigid, habit-forming memory system, favoring rapid reactions toward threatening stimuli

(Quaedflieg and Schwabe, 2018). Likewise, several other studies in humans found association

between high levels of cortisol, during or immediately after memory acquisition, and a loss of

contextual dependency in episodic memories (van Ast et al., 2013). Furthermore, Roozendaal

and Mirone have found, in rats, that acute CORT treatment after training in a modified version

of the inhibitory avoidance task elicited higher fear generalization in episodic-like memories

(Roozendaal and Mirone, 2020). Accumulated evidence have shown that, in fact, high plasma

CORT levels after an aversive situation increase the interaction between the basolateral nucleus

of the amygdala and the prefrontal cortex, eliciting the consolidation of a more 'generalizable'

memory trace (Bahtiyar et al., 2020). On the other hand, in humans, cortisol seemed to impair

generalization in low-arousing types of associative learning (Dandolo and Schwabe, 2016) and

enhance contextualization of emotionally charged pictures (Van Ast et al., 2014). Hence, it is

possible that the glucocorticoid system modifies the specificity of emotionally charged

memories, but the orientation of this modulation is still controversial.

Parallel to the investigation of how stress modulates recent memory generalization,

decades of research have shown that the process of systems consolidation -- in which the

retrieval of contextual memory traces supposedly become more dependent of neocortical

regions over time -- induces a loss of contextual details in episodic memories with the passage

of time (Moscovitch et al., 2016). This loss of contextual specificity is associated with time-

dependent fear generalization across different contexts (Tonegawa et al., 2018). Even though

there is evidence associating stress with generalized contextual memories, few studies

investigated whether stress or high arousal levels during training alter systems consolidation and

the phenomenon of time-dependent fear generalization (Atucha et al., 2017; Bahtiyar et al.,

2020; Pedraza et al., 2016).

To elucidate this issue, a recent study showed that a mildly arousing contextual fear

conditioning (CFC) training in rats (with 3 footshocks of 0.3 mA) induced low levels of post-

training plasma CORT and persistent contextual discrimination at 14 and 28 days after training.

On the other hand, moderate CFC training (3x 0.6 mA footshocks) elicited higher levels of post-

training plasma CORT and a generalized contextual fear response 28 days later. Interestingly,

animals trained with moderate CFC and tested 14 days later showed a generalized fear memory

in the first experiment, and a discriminative fear response in the second (replication)

experiment. In addition, freezing behavior of rats tested in the novel context at this time point

was linearly associated with individual post-training plasma CORT levels, suggesting that there

could be an interaction between training arousal level and post-training glucocorticoid

activation that facilitates fear generalization, depending on individual variability (dos Santos

Corrêa et al., 2019). This evidence suggests that systems consolidation is a complex process that

expresses itself differently after mild or highly arousing CFC, but it also raises the question of

whether this process can be further tempered by artificially activating the glucocorticoid system

via systemic treatment with CORT. The present study, therefore, investigated the modulation of

time-dependent fear generalization exerted by CORT administration following mild and
moderate CFC trainings.

METHODS

2.1 Subjects

Three-month old male Wistar rats, obtained from *Centro de Desenvolvimento de*
Modelos Experimentais para Biologia e Medicina (CEDEME-UNIFESP, total n = 224;
weighing between 275-385g at time of training), were kept in controlled conditions of
temperature ($23 \pm 2°C$), humidity and light/darkness cycle of 12:12 hours (light phase starting at
7am). Rats were randomly housed in groups of 4 per home-cage (30 cm $\times$ 16 cm $\times$ 28 cm) and
provided with food and water *ad libitum*. Wooden flakes were used as bedding in the cages. The
rats were adapted to the vivarium for at least one week before the beginning of the experiments.
A paper towel in each cage was used as environmental enrichment. Sanitation was provided
twice a week by the vivarium technicians. Training and testing were performed during the light
phase of the cycle (between 10:00am-03:00pm), at the rat's nadir of the diurnal rhythm for
corticosterone. The principal experimenter was a male. All procedures were conducted
according to the guidelines and standards of CONCEA - *Conselho Nacional de Controle de*
Experimentação Animal (Brazilian Council of Animal Experimentation) and were previously
approved by the Ethics Committee on Animal Use - UFABC (CEUA - protocol numbers
5553080618 and 7734310719).

2.2 Systemic Drug administration

A preformed water-soluble complex of corticosterone and 2-hydroxypropyl-β-
cyclodextrin (CORT-HBC, 0.25, 0.5, 1.0, 2.0, 4.0 or 8.0 mg/kg, C174 Sigma-Aldrich,
Germany) dissolved in sterile saline, in a volume of 2 ml/kg body weight, was administered
subcutaneously immediately after the training session. We chose the drug, solubility and other
parameters based on previous studies (Kaouane et al., 2012; Rusu et al., 2018; Timmer and
Sandi, 2010). Drug administration took place in the training room, opposite from the

conditioning chambers. Doses of CORT-HBC were selected based on a previous study that

showed an effect on contextual fear memory in mice with this specific molecular compound

(Kaouane et al., 2012) and a standardization experiment was done to choose the best doses for

our experimental design with Wistar rats. Drug solutions were freshly prepared before each

experiment.

2.3 Apparatus

Behavioral experiments were conducted in two identical automated fear conditioning

chambers (Med-Associates, Inc., St. Albans, VT) connected to a computer interface enabling

video recording, analysis, and automated measurement of the rat's freezing behavior in real

time. The chambers and software used were previously described in detail (de Paiva et al., 2021;

dos Santos Corrêa et al., 2019). Freezing measure was automated (VideoFreeze Version

1.12.0.0, MedAssociates) and set for less than 20 movement units for at least 30 frames (1s)

(Bueno et al., 2017; Moreira-Silva et al., 2018).

The training context (context A) was characterized by a grid floor composed of 20

stainless steel rods (diameter: 4.8mm), top and front walls made of transparent polycarbonate, a

back wall made of white acrylic covered with a 5-cubed white-on-black pattern, stainless-steel

sidewalls, and a drop pan below the grid floor. The light in the conditioning box remained on

and only the exhaust fan noise was emitted during the training and test sessions. Context A was

cleaned with alcohol 10% before and after each session.

The novel context (Context B) consisted of the conditioning chamber and stainless-steel

drop pan personalized with a grid floor (20 interleaved rods of either 4.8 or 9.5mm of diameter)

and white Plexiglas curved sidewalls extending across the back wall. The light in the box

remained turned off and a 90-dB white noise was emitted in addition to the fan noise during the

test. Context B was cleaned with a 5% acetic acid solution before and after each session.

2.4. Contextual Fear Conditioning task and experimental designs

In all experiments, rats from each home-cage were randomly assigned to one of the

experimental or control groups, and all groups were matched according to the average body

weight. Rats were handled for 3 days prior to the CFC training, each for 3 min, in the training

room. They were habituated to an adjacent room for 40 to 90 minutes before the handling sessions and both training and testing. To minimize potential confounders, the same experimenter always handled the rats.

On the training session, rats were individually transported by the experimenter from the adjacent room to the training room in a white acrylic, individual cage (to which they were accustomed during habituation days) and placed in context A. After 2 minutes of free exploration, three footshocks (0.3 or 0.6 mA, 1s each, depending on the experiment) were delivered with intervals of 30s between them. One minute after the third shock, the animal was removed from the apparatus and immediately received a subcutaneous (s.c) injection of saline or different doses of CORT-HBC. The same experimenter that transported the animals to the training chamber removed them from the apparatus and administered the injections. Each animal was then returned to its individual cage and transported back to the adjoining room where it stayed for at least 90 minutes before being placed back in it home-cage and returned to the vivarium.

For test sessions, rats were transported from the adjacent room to the training room in a wooden box (which was novel to the animals), by a different experimenter from the training session, unaware of the animal's treatment group, and placed in context B for 4 minutes. Afterwards, each animal was removed from this conditioning chamber by the principal experimenter, placed into the white acrylic individual cage for up to 1 minute, and then exposed to context A, for 4 minutes. In both contexts, no shock was delivered during the test and animal's freezing time was automatically measured by the software. Both the blindness of the different experimenter and the automated freezing measurement reduced the potential risk of bias. Figure 1 shows an overview of the experimental procedures.

2.4.1 Dose selection experiment

A dose-response curve was performed to verify the best doses of CORT-HBC for our experimental design with Wistar rats. Rats were trained with three footshocks (0.3 mA) and, immediately after training, received a s.c injection of saline or CORT-HBC (0.25, 0.5, 1.0, 2.0, 4.0 or 8.0 mg/kg). The test sessions were performed 2 days after training, as described above.

2.4.2 Corticosterone modulation of recent and remote contextual fear memory specificity

In order to investigate whether the treatment with CORT-HBC modulates contextual

fear specificity over time following mild or moderate CFC training, different testing timepoints

were chosen based on the study of dos Santos Corrêa et al. (2019). In this previous study, mild

CFC (0.3mA) elicited discriminative fear memory up to 28 days, whereas moderate CFC

(0.6mA) resulted in contextual generalization or discrimination 14 days after training, in two

different experiments. These divergent results on the 14th day may be explained by the different

levels of post-training plasma CORT observed between both groups of rats (see Figure S1 in the

supplementary material). For the present study, we hypothesized that injections of CORT-HBC

immediately after a mild CFC training would promote fear generalization at remote timepoints

(28 or 42 days after training), whereas the same treatment after a moderate CFC training would

facilitate fear generalization at the 14th day timepoint. Therefore, rats were subject to the CFC

training with one of two shock intensities (3x of 0.3 or 0.6mA) and immediately after training

received a s.c. injection of saline or CORT-HBC (4.0 or 8.0 mg/kg). Rats trained with 0.3mA

footshocks were tested 28 or 42 days after training, whereas those trained with the 0.6mA

intensity were tested 2 or 14 days after training. Different groups of rats were used for each

footshock intensity and timepoint.

2.5 Statistical analysis

For each experiment, the experimental unit was the individual rat. All descriptive and

inferential statistical analysis was done using JAMOVI (version 1.63). Fear response to

footshocks was quantified as the time the animal spent in freezing during the last minute of the

training session. The conditioned fear response to the training context was quantified as the time

the animal spent in freezing in context A, whereas fear generalization was considered as the

time of freezing in context B during the recall tests. Behavioral results are expressed as the

group mean percent freezing time ± standard error of the mean (S.E.M). An "index of

generalization" was calculated using the formula ($\frac{Freezing\ time\ in\ context\ B}{Sum\ of\ freezing\ times\ in\ contexts\ A\ and\ B}$) and

are expressed as group mean generalization (± S.E.M). A generalization threshold of 0.3 was set

to distinguish rats between 'discriminator' (< 0.30) and 'generalizer' animals (≥ 0.30), a level

that is equivalent to the one reported by Wiltgen and colleagues (Wiltgen et al., 2010). When

needed to satisfy the requirements for the use of the ANOVA, total freezing times were

transformed using the square root transform function (dos Santos Corrêa et al., 2019) to make

data normally distributed and homogeneous, according to the tests of Kolmogorov-Smirnov and

Levene. When not stated otherwise, data used in the inferential tests was normally distributed

and homogenous. All statistical analysis are stored together with the behavioral data at

https://data.mendeley.com/datasets/7xvshh2s59/.

Freezing data from the last minute of the training sessions (post-shock freezing times)

were analyzed with one-way ANOVAs with Treatment as the independent factor. Freezing data

from the test sessions were analyzed with one-way or repeated measures ANOVAs

(RMANOVAs, with freezing time in Contexts as within-subject dependent variables and

Treatment as an independent factor). Generalization index data were analyzed with one factor

ANOVAs with Treatment as the independent factor. The Tukey *post hoc* test was further used

to identify significant differences when applicable. Bonferroni corrections were used for

multiple comparisons when needed. For the non-homogenous Generalization Index data, the

Welch corrected ANOVA was used to analyze means and the Games-Howell Tukey corrected

post hoc tests were used to compare differences between groups. The tests 'linear tendency of

Cochran-Armitage' (experiment 1 only) and 'likelihood-ratio chi-square' were used to test the

difference in the distribution of 'generalizer' and 'discriminator' animals across all treatment

groups.

Significance for all inferential tests was set at $p \leq 0.05$. Effect sizes for ANOVAs and

RMANOVAs ($\eta^2 p$ or $\eta^2 G$, respectively) are reported only when the test was found significant

("$\eta^2 p$" or "$\eta^2 G$" values above 0.14 are considered large effects; values between 0.06 and 0.14

are considered moderate; and below 0.06, small). For post hoc tests, the Cohen's d value is

reported as a measure of effect size when the test was found significant. All four measures

(statistical difference in mean freezing times in the novel context, lack of difference between the

training and the novel contexts, difference in mean generalization index and difference in

proportion of generalizers per group) were used to consider group level generalization.

Significant results in at least two of these measures were considered as an evidence of fear

generalization in any particular group.

Training Session   Test Session

Figure 1: Schematic representation of the experimental designs. *Rats were exposed to the*
training context for two minutes, when they were free to explore. Thereafter, they received three
footshocks of one intensity, 0.3 or 0.6mA (1s each, 30s between them). Following the last
footshock, each rat stayed in the chamber for another minute and was then removed.
Immediately after training, the rats received an injection of CORT-HBC or saline, according to
their experimental group. After 2, 14, 28 or 42 days, rats were tested for long-term memory.
They were first exposed to a novel context (Context B) for 4 minutes and then returned to their
individual cage, where they stayed for 1 minute. Then, they were exposed to the training context
(Context A) for 4 minutes.

RESULTS

3.1 CORT-HBC dosage selection

In order to verify and select which CORT-HBC doses would be effective in modulating

contextual fear memory consolidation, different groups of rats received a subcutaneous injection

of saline or CORT-HBC (0.25, 0.5, 1.0, 2.0, 4,0 and 8.0 mg/kg) immediately after training and

were tested 2 days later as described above. Results indicate that higher doses of CORT-HBC

seem to promote recent memory accuracy (Figure 2). The RMANOVA showed a significant

Context effect [$F_{(1, 40)} = 39.73$, $p < 0.01$, $\eta^2G = 0.24$], but no Treatment effect [$F_{(6, 40)} = 0.44$, p

$= 0.85$] or interaction effect between Context * Treatment [$F_{(6, 40)} = 1.12$, $p = 0.37$, Fig. 2a]. The

same result was confirmed by analyzing the mean Generalization Index [$F_{(6, 40)} = 1.58$, $p = 0.18$,

Fig. 2b]. Nonetheless, a trend toward a differential distribution in the number of animals that

showed Generalization Indices lower than 0.3 (Discriminators), as the dose of CORT-HBC

increased, was confirmed by the test of linear tendency of Cochran-Armitage [$\chi^2_{(1,\ 47)} = 9.4$, $p <$

0.01] and the likelihood-ratio chi-square test [$\chi^2_{(6,\ 47)} = 12.68$, $p = 0.048$, Fig. 2c].

Figure 2: Post-training corticosterone-HBC (CORT-HBC) modulates recent (2 days) memory
specificity. *(a) Mean freezing percentage ± standard error of the mean (S.E.M) shown in the*
novel context (B, diamond symbol) and then in the training context (A, square symbol) of groups
injected with Saline (n = 7), CORT-HBC 0.25 mg/kg (6), 0.5 mg/kg (6), 1 mg/kg (6), 2 mg/kg
(7), 4 mg/kg (7) and 8 mg/kg (7), immediately after mild CFC training (3 footshocks of 0.3 mA).
The dotted lines and gray area represent the lower error bar in context B and the higher error
bar in context A for the saline group. (#) p < 0.01 in the main effect of the factor Context. (b)
Mean (± S.E.M.) Generalization Index [B/(A+B)] of each treatment group. The dotted line
represents the threshold point between discriminators (0 – 0.3) and generalizers (0.31-1). (c)
Percentage of animals that were considered "discriminators" (below) or "generalizers"
(above). The percentage of discriminators shows a linear increase in response to increasing
doses of CORT-HBC (*) p < 0.01 in the Cochran-Armitage test for linear association).

3.2 Corticosterone differentially modulates fear memory specificity for mild and moderate CFC

training intensities over time

3.2.1 Mild CFC training intensity

240   To allow comparison of the treatment effect found above with all the other experiments,

241 a subset of the data presented at 3.1 was used to further analyze fear generalization of rats

242 treated with the highest doses of CORT-HBC (4.0 or 8.0 mg/kg, s.c.) or saline, immediately

243 after the mild intensity CFC training, and tested 2 days later (Fig. 3a-c). There was no difference

244 in post-shock freezing levels between these groups during training (Treatment [$F_{(2, 19)} = 0.09$, p

245 $= 0.92$], Table S1). For the test session, CORT-HBC treatment increased the difference in rat's

246 freezing percentage between contexts B and A in the recent memory time point. RMANOVA

247 indicated a significant effect for Context [$F_{(1, 19)} = 26.96$, $p < 0.001$, $\eta^2G = 0.26$] and for the

248 interaction Context * Treatment [$F_{(2, 19)} = 3.82$, $p = 0.04$, $\eta^2G = 0.09$], but not for Treatment [F

249 $_{(2, 19)} = 0.38$, $p = 0.69$]. Interaction post hoc tests confirmed that rats treated with Saline

250 expressed similar levels of freezing in both contexts ($t_{(30.11)} = -0.85$, $p = 0.95$) whereas CORT-

251 HBC treated animals showed lower freezing times when exposed to context B than when

252 exposed to context A (CORT 4.0mg/kg, $t_{(30.11)} = -3.46$, $p = 0.03$, $d=-1,60$; CORT 8.0mg/kg, t

253 $_{(30.11)} = -4.72$, $p = 0.002$, $d=-2.16$, Fig. 3a). For the Generalization Index, the one-way ANOVA

254 showed a significant Treatment effect [$F_{(2, 19)} = 4.67$, $p = 0.02$, $\eta^2p = 0.33$] and the post hoc tests

255 revealed that the 8.0 mg/kg-CORT-HBC group showed lower generalization levels than the

256 Saline group ($t_{(19)} = -2.86$, $p = 0.03$, $d=-1.31$). Other comparisons were not statistically

257 significant ($p > 0.07$, Fig. 3b). The likelihood-ratio chi-square test has further confirmed a

258 difference in the percentage of discriminators' rats in CORT-HBC treated groups (N=6

259 discriminators for both 4,0mg/kg and 8.0mg/kg doses) when compared to the saline-treated

260 animals [N = 1 discriminator, $\chi^2_{(2, 22)} = 9.29$, p = 0.01, Fig. 3c].

261   Following the recent memory test, we chose two time points *a priori*, based on the

262 results found by dos Santos Corrêa and colleagues (2019), to test our hypothesis that CORT-

263 HBC would elicit generalized fear at remote timepoints. Two additional sets of rats were trained

264 with the mild CFC intensity (0.3 mA), received a s.c. injection of CORT-HBC (4.0 or 8.0

265 mg/kg) or saline immediately after training and were tested 28 or 42 days later. For the first set

266 of animals (that were tested at the 28[th] day), there was no difference in post-shock freezing

267 levels during training (Treatment [$F_{(2, 21)} = 0.50$, $p = 0.61$], Table S1). During the test session,

contrary to our hypothesis, rats injected with the highest dose of CORT-HBC (8.0 mg/kg)

showed a lower generalization across contexts, when compared to the control group (Fig. 3d-f).

As freezing data in contexts B and A did not have homogenous variance (p < 0.05 for both),

they were transformed using the square root transform and the RMANOVA was calculated

using the transformed data. The RMANOVA revealed significant effects for Context [$F_{(1, 21)}$ =

59.92, $p < 0.001$, $\eta^2G = 0.44$] and Treatment [$F_{(2, 21)}$ = 5.67, $p = 0.01$, $\eta^2G = 0.28$]. The

interaction between Context and Treatment showed a nearly significant effect [$F_{(2, 21)}$ = 3.34, p

= 0.055, $\eta^2G = 0.08$], but as the generalized eta squared of 0.08 indicates a moderate effect size,

unprotected post hoc tests were calculated. Post hoc tests revealed that groups injected with 8.0

mg/kg of CORT-HBC ($t_{(21)}$ = -6.36, $p < 0.001$, d=-2.77) or saline ($t_{(21)}$ = -4.35, $p = 0.003$, d=-

1.90) showed higher freezing levels in context A than in context B. In contrast, rats injected

with 4.0 mg/kg of CORT-HBC displayed similar freezing levels in both contexts ($t_{(21)}$ = -2.71, p

= 0.12). Besides, rats in the 4.0 mg/kg-CORT-HBC group showed lower freezing times in

context A when compared to the Saline group in the same context ($t_{(34.7)}$ = -3.48, $p = 0.02$, d=-

1.18). Other post hoc comparisons were not significant ($p > 0.09$). To further assess between-

group differences, additional Bonferroni corrected ANOVAs were used to compare freezing in

B and A separately (Table 1). These ANOVAs confirmed that rats in the 4.0 mg/kg-CORT-

HBC group showed lower freezing times when compared to the Saline group in both contexts (p

= 0.01 *and p = 0*.047 in context A and B, respectively) whereas the 8.0 mg/kg-CORT-HBC

group showed a non-significant trend towards a decrease in freezing percentage in the novel

context, when compared to the saline group in this same context ($p = 0.06$, Table 1; for all other

one-way ANOVAs see Table S2). For the Generalization Index, the Welch corrected ANOVA

was used due to non-homogeneity of variance ($p < 0.05$). There was a significant Treatment

effect [$F_{(2, 10.92)}$ = 5.50, $p = 0.02$, $\eta^2p = 0.19$], and the post hoc tests revealed that animals

injected with 8.0 mg/kg of CORT-HBC presented lower generalization levels in comparison to

the Saline group ($t_{(9.1)}$ = -3.13, $p = 0.03$, d=-2.08), whereas the 4.0 mg/kg-CORT-HBC group

showed a mean generalization index similar to the Saline ($t_{(11.3)}$ = -0.26, $p = 0.97$) and to the 8.0

mg/Kg-CORT-HBC groups ($t_{(7.7)}$ = 1.63, $p = 0.29$, Fig. 3e) and below the threshold of

generalization. The likelihood-ratio chi-square test, however, did not show differences in the

proportion of discriminators across all groups [$\chi^2_{(2, 24)} = 4.98$, p = 0.08, Fig. 3f].

All rats tested on the 42nd day after training, on the other hand, showed higher freezing

levels in Context A than in Context B and similar generalization levels (Fig. 3g-i). For this set

of animals, there was no difference in post-shock freezing levels during training (Treatment [F

$_{(2, 20)} = 0.27$, $p = 0.77$], Table S1). During the test session at this time point, freezing time data in

context A was non-homogenous ($p < 0.05$) and therefore the transformed data was analyzed via

RMANOVA. There was a significant effect for Context [$F_{(1, 20)} = 13.93$, $p = 0.001$, $\eta^2 G = 0.26$]

but not for Treatment [$F_{(2, 20)} = 0.29$, $p = 0.75$] or the interaction between Context and

Treatment [$F_{(2, 20)} = 0.26$, $p = 0.77$]. Mean generalization index was similar in all Treatment

groups [$F_{(2, 20)} = 0.37$, $p = 0.70$, Fig. 3h] and this result was further confirmed by the likelihood-

ratio chi-square test [$\chi^2_{(2, 23)} = 0.26$, p = 0.88, Fig. 3i].

In summary, for the mild CFC training intensity, the post-training treatment with the

highest dose of CORT-HBC (8.0 mg/kg) promoted contextual discrimination at 2 days and

induced an enhancement of contextual memory accuracy at 28, but not 42, days after training.

Additionally, rats injected with 4.0 mg/kg of CORT-HBC showed less freezing in both contexts

at the 28[th] day when compared to Saline-treated rats, but they preserved the generalization index

under the threshold of generalization and this group showed a similar proportion of generalizer

animals when compared to the other two groups.

Figure 3: Corticosterone administration after mild intensity CFC training increases contextual memory specificity up to 28 days. (a, d, g) Mean freezing percentage ± Standard error of the mean (S.E.M) shown by rats exposed to the novel context (B) and then to the training context (A) after (a) 2, (d) 28 or (g) 42 days after training. Lines represent individual freezing data for each rat in context B and then in A. (#) p ≤ 0.001 when compared to within-subject freezing expressed in Context A. (*) p = 0.02 when compared to the mean freezing times expressed by the Saline group in Context A. (b, e, h) Mean Generalization Index ± S.E.M. [B/(A+B)] after (b) 2, (e) 28 or (h) 42 days after CFC. The dotted line represents the threshold between "discriminators" and "generalizers". Filled symbols represent the animals considered discriminators and the unfilled symbols represent the ones considered generalizers. (*) p < 0.05 when compared to the Saline group. (c, f, i) Percentage of generalizer and discriminator animals per group after (c) 2, (f) 28 or (i) 42 days after CFC. (*) p < 0.05 in the proportion test. Number of rats for each group: 2 days, Saline (N = 7), CORT-HBC 4 mg/kg (N = 8), 8 mg/kg (N = 7). 28 days, Saline (N = 8), CORT-HBC 4 mg/kg (N = 8), 8 mg/kg (N = 8). 42 days, Saline (N = 8), CORT-HBC 4 mg/kg (N = 8), 8 mg/kg (N = 7).

3.2.2 Moderate CFC training intensity

To verify whether CORT injections after a moderate CFC would facilitate or accelerate contextual fear generalization , different sets of rats were trained with 3 footshocks of 0.6mA, received s.c. injections of CORT-HBC (4.0 or 8.0 mg/kg) or saline immediately after training and were tested 2 or 14 days later. The results from the first set of animals, tested for the recent memory (2 days after training), indicate that all groups of rats showed higher freezing response

in the training context than in the novel context, unabated by treatment (Fig. 4a). Again, there

was no difference in post-shock freezing times during training (Treatment [$F_{(2, 21)}$ = 1.64, p =

0.22], Table S1). For the test data, The RMANOVA showed a significant effect for Context [F

$_{(1, 21)}$ = 26.45, $p < 0.001$, η^2G = 0.28] but not for Treatment [$F_{(2, 21)}$ = 0.07, p = 0.93] or Context

* Treatment interaction [$F_{(2, 21)}$ = 0.35, p = 0.71, Fig. 4a]. The mean Generalization Index were

similar between all groups [$F_{(2, 21)}$ = 0.29, p = 0.75, Fig. 4b] and this lack of effect was further

confirmed by the likelihood-ratio chi-square test [$\chi^2_{(2, 24)}$ = 0.00, p = 1.00, Fig. 4c].

On the other hand, results from the second set of animals, tested 14 days after training,

indicate that rats injected with 4.0 mg/kg of CORT-HBC showed increased freezing in Context

B when compared to the other groups, while maintaining the same level of freezing in Context

A (Fig. 4d). Freezing data from the training session did not show any difference between groups

(Treatment [$F_{(2, 21)}$ = 1.61, p = 0.22], Table S1). During the test session, freezing data from

Context B did not show homogenous variance ($p < 0.05$); therefore, they were transformed

using the square root function. The RMANOVA showed a significant effect for Context [$F_{(1, 21)}$

= 6.83, p = 0.02, η^2G = 0.11] and Treatment [$F_{(2, 21)}$ = 5.27, p = 0.01, η^2G = 0.24]. However, the

interaction between Context and Treatment was not significant [$F_{(2, 21)}$ = 1.63, p = 0.22]. Post

hoc tests of the Treatment main effect showed that rats treated with 4.0 mg/kg of CORT-HBC

showed higher overall freezing times than the ones treated with Saline ($t_{(21)}$ = 2.54, p = 0.048, d

= 1.11) or 8.0 mg/kg of CORT-HBC ($t_{(21)}$ = 3.02, p = 0.02, d = 1.32). Overall freezing times of

Saline and 8.0 mg/kg-CORT-HBC groups were not statistically different from each other ($t_{(21)}$ =

-0.48, p = 0.88; Fig. 4d). As there was no interaction effect, it is not possible to perform pos-hoc

tests to compare the freezing behavior in contexts A and B of each group, and the significant

effect for Context indicates that all animals spent in average more time in freezing in the

training context than in the novel context. To further assess between-group differences,

additional Bonferroni corrected one-way ANOVAs were used to compare freezing in Context A

or B, separately (Table 1 – 0.6mA / 14 days). These ANOVAs showed no significant effect

between groups tested in the training context (p = 0.36), but a significant effect between groups

tested in the novel context [$F_{(2, 21)}$ = 7.23, $p < 0.01$, η^2p = 0.41]. The post-hoc analyses

confirmed that, in tests performed in context B, rats in the 4.0 mg/kg-CORT-HBC group showed higher freezing times when compared to Saline ($p = 0.02$) and 8.0 mg/kg-CORT-HBC ($p < 0.01$) groups, while these two groups showed no difference between them ($p = 0.86$). Moreover, although the comparison between generalization indices did not show any differences [$F_{(2, 21)} = 1.58$, $p = 0.23$, Fig. 4e], the likelihood-ratio chi-square test showed a significant effect in the distribution of generalizers in the 4.0mg/kg-CORT-HBC group (N = 8) when compared the 8.0mg/kg-CORT-HBC (N = 4) and Saline groups [N = 5, $\chi^2_{(2, 24)} = 7.30$, $p = 0.03$, Fig. 4f].

In summary, according to two of the pre-established measures (increased freezing levels in Context B and a higher proportion of generalizing animals), we may infer that treatment with CORT-HBC at the dose of 4.0 mg/kg, but not 8.0 mg/kg, immediately after the moderate CFC training, enhanced fear memory generalization 14 days later, without interfering in recent memory accuracy.

Table 1: Bonferroni corrected One-way ANOVAs for the freezing times in the training (A) or novel (B) contexts and post hoc tests

0.3mA|28 days					
ANOVAs	**F**	**df1**	**df2**	**P**	**η²p**
Freezing B	7.23	2	21	**<0.01***	0.41
Freezing A	5.60	2	21	**0.01***	0.35

Post hoc tests Freezing A		**Mdiff**	**t**	**df**	**P**	**D**
Saline (FreezA)	CORT-HBC 4.0 (Freez A)	4.71	3.17	21	**0.01***	1.58
Saline (FreezA)	CORT-HBC 8.0 (Freez A)	0.97	0.65	21	0.79	0.33
CORT-HBC 4.0 (Freez A)	CORT-HBC 8.0 (Freez A)	-3.74	-2.52	21	**0.05***	-1.26

Post hoc tests Freezing B		**Mdiff**	**t**	**df**	**P**	**D**
Saline (FreezB)	CORT-HBC 4.0 (Freez B)	3.08	2.55	21	**0.047***	1.28
Saline (FreezB)	CORT-HBC 8.0 (Freez B)	2.97	2.46	21	0.06	1.23

| CORT-HBC 4.0 (Freez B) | CORT-HBC 8.0 (Freez B) | -0.11 | -0.09 | 21 | 1.00 | -0.04 |

0.6mA|14 days						
ANOVAs		**F**	**df1**	**df2**	**P**	**η²p**
Freezing B		7.23	2	21	**<0.01***	0.41
Freezing A		1.07	2	21	0.36	0.09

Post hoc tests Freezing B		**Mdiff**	**t**	**df**	**P**	**d**
Saline (FreezB)	CORT-HBC 4.0 (Freez B)	-3.64	-3.0	21	**0.02***	-1.5
Saline (FreezB)	CORT-HBC 8.0 (Freez B)	0.64	0.52	21	0.86	0.26
CORT-HBC 4.0 (Freez B)	CORT-HBC 8.0 (Freez B)	-4.3	-3.5	21	**<0.01***	-1.76

Bonferroni corrected ANOVAs significance is 0.025. (*) indicates a p-value below the level of significance set for the test.

Figure 4: Corticosterone administration after moderate CFC training intensity increases
contextual fear generalization at 14 days. (a, d) Mean freezing percentage ± Standard error of
the mean (S.E.M) shown by rats exposed to the novel context (B) and then to the training
context (A) (a) 2 or (d) 14 days after training. Lines represent individual freezing data for each
rat in context B and then A. (#) p < 0.001 when compared to within-subject freezing expressed
in Context A. (*) p = 0.01 when compared to the 8.0mg/kg and Saline groups. (b, e) Mean
Generalization Index ± S.E.M. [B/(A+B)] (b) 2 or (d) 28 days after CFC. The dotted line
represents the threshold between "discriminators" and "generalizers". Filled symbols
represent the animals considered discriminators and the symbols unfilled represent the ones
considered generalizers. (c, f) Percentage of generalizer and discriminator animals per group
after (c) 2 or (f) 14 days after CFC. (*) p < 0.05 in the proportion test. Number of rats for each

group: 2 days, Saline (N = 8), CORT-HBC 4 mg/kg (N = 8), 8 mg/kg (N = 8). 14 days, Saline
(N = 8), CORT-HBC 4 mg/kg (N = 8), 8 mg/kg (N = 8).

DISCUSSION

The results reported here indicate that CORT-HBC modulates contextual fear memory

specificity differently for mild and moderate training intensities. Our findings in the Saline

groups replicate previous findings that mild CFC (3x 0.3mA) elicit similar levels of freezing

during recent retrieval tests but discriminate between contexts at remote timepoints (dos Santos

Corrêa et al., 2019). Contrary to our hypothesis, the treatment with CORT-HBC after this

training intensity promoted a contextual memory accuracy in the recent (2 days) retrieval test,

and increased the discriminative fear response of animals at the 28th day, compared to the

control rats (even though only one of the variables measured was significant and only with the

higher dose). Additionally, on the 28th day retrieval test, the 4.0 mg/kg dose seemed to lower

overall freezing levels while still maintaining contextual discrimination. It is possible that this

dose of CORT-HBC, given immediately after a mild CFC training, is impairing the strength or

the persistence of the contextual memory, without interfering in its specificity (according to the

index of generalization and the proportion of "generalizer" animals). Curiously, freezing levels

of this group (4 mg/kg-CORT-HBC) at the 28th day was very similar to that observed in all

groups at the 42 day time point. Therefore, another possibility is that post-training injections of

4 mg/kg of CORT-HBC, immediately after a mild CFC training, is anticipating the freezing

levels that would be expected at more remote time points.

Dos Santos Corrêa and colleagues (2019) also showed that rats trained with the

moderate CFC intensity (3x 0.6mA) showed contextual discrimination in the recent retrieval

tests (2 days after training) and then generalized fear between the training context and a new

context at the 28th day after training. It is important to note that another group of rats tested 14

days after training showed a generalized memory in the first experiment. However, this result

was not replicated in a second experiment reported in the same study in which another group of

rats trained under the same parameters discriminated across contexts at this intermediate time

point (dos Santos Corrêa et al., 2019). We did additional *post hoc* analysis of that data and the

results thereof revealed that the replication group had lower mean CORT levels when compared

to the rats from the original experiment (Fig S1, supplementary material) which raised the

question whether this moderate training intensity and time point of testing combined are more

susceptible to CORT's modulation on memory accuracy. In the present study, we hypothesized

that animals trained with this footshock intensity and treated with CORT-HBC would show a

facilitation to generalize contextual fear. Our results indicate that treatment with CORT-HBC at

the dose of 4.0 mg/kg, but not 8.0 mg/Kg, increased generalized freezing to the novel context,

without increasing freezing in the training context, 14 days after the moderate CFC training.

These data suggest that treatment with CORT-HBC may 'tip the balance' towards fear

generalization at the 14^{th} day after training following an inverted U-curve dose-response

function, with the dose of 4.0 mg/kg eliciting fear generalization and the dose of 8.0 mg/kg

maintaining some contextual discrimination similarly to the saline group. This result suggests

that there is an optimal glucocorticoid activation after a highly arousing experience that

modulates the consolidation of generalized memory traces, which agrees with previous data

reported in rats (Pedraza et al., 2016; Roozendaal and Mirone, 2020). However, it is important

to emphasize that all comparisons across timepoints are done at the group level because

different sets of animals were used for each time interval and footshock intensity, therefore this

inference is only correlational.

Our results indicate, therefore, that acute post-training CORT treatment elicits different

behavioral outcomes, contingent on the CFC training intensity. Likewise, studies in humans

suggest that stress may enhance memory accuracy (Kensinger et al., 2007; Segal et al., 2012),

but there are divergent results on the literature showing an increase of false content in episodic

memories and reduced accuracy after learning under stress (Rimmele et al., 2011; Schwabe et

al., 2009; Van Ast et al., 2014). We emphasized, however, that our study used a post-training

CORT treatment that is not directly translatable to some of the studies in humans cited above,

since their emotional or pharmacological interventions were performed at different phases of

memory encoding, not only after training. Regardless of the intervention's timepoint, these

studies indicate that stress at the time of encoding can act either enhancing or impairing memory

accuracy. As supporting evidence in rodents to our results, that suggest that acute CORT

treatment immediately after a highly arousing CFC training temporally modulates contextual

specificity, Pedraza and colleagues reported that pre-training CORT inhibition by metyrapone

maintained the accuracy of contextual fear memory up to 25 days after a strong CFC training in

rats, in contrast to vehicle treated rats that showed generalized fear at this timepoint. Roozendaal

and Mirone (2020) showed that post-training CORT treatment induced a generalized

strengthening of memory in a modified version of the inhibitory avoidance task during recent

retrieval, while noradrenergic activation after training induced an increase in the accuracy of the

association between context and shock, which indicates that both hormones act synergistically

to consolidate contextual features of an aversive event. These findings together point to an

important role played by the level of arousal during acquisition on CORT's modulation of

contextual fear memories. It is important to note that the latter results diverge slightly from our

results with mild training intensity, in which CORT treatment seems to enhance memory

specificity up to 28 days, but it can be rationalized by differences in rat lineage, quantity and

intensity of footshocks received, as well as by the different behavioral outcomes quantified

during the test and the different intrinsic features of the CFC and inhibitory avoidance tasks. On

the other hand, our results with the moderate CFC training agree with this hypothesis that

CORT increases fear generalization. More studies investigating the effects of blocking the

glucocorticoid and noradrenergic systems are needed to better understand the role of these

hormones in the specificity of contextual fear memory over time.

Additional evidences in humans showed that increasing stress before training elicited

reorientation of attention toward salient contextual features (Clewett and Murty, 2019;

Schwabe, 2017). These evidences suggest that more arousing experiences elicit increased

noradrenergic activity (Roozendaal and Hermans, 2017) that, followed by appropriate CORT

expression, elicit optimal consolidation of the memory trace. Therefore, our results reported

here can be explained by the increasing levels of arousal during mild or moderate CFC sessions

that interact with circulating CORT in response to the aversive situation, prompting differential

processes of systems consolidation (Bahtiyar et al., 2020; dos Santos Corrêa et al., 2019). Mild

intensity CFC may induce a process of systems consolidation in which exogenous CORT

sustains an increased memory specificity at remote timepoints, whereas higher intensity CFC

possibly recruits brain regions associated with more rigid learning, such as the striatum, and

induces a different process of systems consolidation in which exogenous CORT facilitates time-

dependent fear generalization.

Our data raise the question of how CORT would prompt different routes of systems

consolidation after low or highly arousing events. It is possible to conjecture that memory

acquisition under stress shifts the focus of attention to central features of the aversive event and

less on its contextual details (Christianson and Loftus, 1987). This shift of focus is associated

with a shift of functional activity from areas involved in contextual consolidation, such as the

hippocampus (Addis and Schacter, 2008), to cortical areas implicated in semantic

representations, such as the parahippocampal cortex (Binder et al., 2009). The hippocampal

memory system is thought to encode separate representation of events and allow the acquisition

of flexible memory representations (Shohamy and Wagner, 2008). Rapid glucocorticoid activity

would act in association with the noradrenergic system in the amygdala, prompting brain-wide

recruitment of brain regions associated with semantic representations and the habit memory

system, favoring the association of threat-related signs to the memory trace and faster

incorporation of the trace into preexisting semantic schemas (Quaedflieg and Schwabe, 2018;

Schwabe, 2017; Schwabe and Wolf, 2012). If stress induces a shift from cognitive to habit

learning, which lacks this flexibility for consolidating contextual representations (Myers et al.,

2003), it could also increase generalization, which agrees with studies done in humans

(McGlade et al., 2019; Rimmele et al., 2011; Schwabe et al., 2009). In contrast, low arousing

events would maintain memory dependence on the hippocampal system.

Our results agree with this supposition. In one hand, mild CFC possibly did not increase

rapid noradrenergic and glucocorticoid activation to levels at which there would be a shift of

memory systems towards striatal learning (Quirarte et al., 1998; Roozendaal and Hermans,

2017), maintaining memory accuracy up to 42 days. Additional treatment with exogenous

CORT further affected memory specificity, increasing contextual discrimination up to 28 days,

as shown by the rats trained with 0.3mA footshocks and treated with CORT-HBC. On the other

hand, moderate CFC possibly reached the necessary stress levels to induce the shift of memory

systems towards reflexive learning. Exogenous CORT for this training intensity acted

facilitating the consolidation of a rigid and contextually poor memory trace (Vogel et al., 2016)

that became generalized over time.

Surprisingly, we did not find an enhancement in fear memory strength after post-

training treatment with CORT when rats were exposed to the training context, as it was

previously reported (Abrari et al., 2009; Cordero and Sandi, 1998; Kaouane et al., 2012; Pugh et

al., 1997; Roozendaal and Mirone, 2020). Even though there are other studies that also did not

report this enhancement after treatment with CORT (Bueno et al., 2017; Marchand et al., 2007),

we hypothesize that the lack of effect in our experiment is due to differences in training

parameters, experimental design or task-related features. It is possible that the order of context

exposure during test sessions in the present study, i.e., first in the novel context and then in the

conditioning context, may have prevented the observation of the CORT-induced enhancement

in contextual memory strength. Some of the studies that have investigated stress hormones

effects on episodic-like memory strength and accuracy used a counterbalanced exposure to

different contexts in the inhibitory avoidance task and reported no order effect (Atucha and

Roozendaal, 2015; Roozendaal and Mirone, 2020). However, pilot experiments done in our

laboratory using a counterbalanced testing after a moderate CFC training (3 footshocks of 0.6

mA) showed an order effect: animals tested first in the training context and then in the novel

context showed an accurate memory either 2 or 28 days after training, whereas rats tested first

in the novel context showed an accurate memory only 2 days after training, and a generalized

memory at the 28th day (unpublished results). These results suggest that, for our experimental

parameters, the exposure to the conditioning context first, during testing sessions, influences

remote memory generalization in the CFC task. As our main interest was to verify whether

CORT administration would modulate memory generalization over time, we chose to test

animals first in the new context.

In conclusion, this study strengthens the hypothesis that CORT modifies time-

dependent memory generalization, associated with the process of systems consolidation. In the

experimental design described here, post-training exogenous CORT after mild CFC seemed to

increase recent and remote memory accuracy and to accelerate freezing levels expected only in

more remote timepoints. On the other hand, CORT injections after moderate CFC seemed to

facilitate time-dependent generalization. This differential modulation of CORT on fear memory

specificity, contingent to training intensity, conveys strong support to the hypothesis of arousal-

induced shift in memory systems during CFC acquisition and consolidation.

Acknowledgements

We thank the valuable support from Vitor Farhat Fernandes, Lorena Vido Lopes and

Gabriela Iramina Gomes for helping in some of the experiments. We also thank the work done

by the technicians that manage the vivarium. The authors would like to thank the members of

the Research group in Neurobiology of Learning and Memory at UFABC (MANAs) for

scientific advice and discussion.

Declaration of Competing Interests

The authors report no conflicts of interest in this work.

Research data for this article

The data used, collected and analyzed for all results here presented can be found at

([dataset] Dos Santos Corrêa, M.; Fornari, 2020) http://dx.doi.org/10.17632/7xvshh2s59.1

Funding

This work was financially supported by the São Paulo Research Foundation (FAPESP)

grant #2017/24012-9 (MdSC), São Paulo Research Foundation (FAPESP) grant #2017/03820-0

(RVF) and *Conselho Nacional de Desenvolvimento Científico e Tecnológico* (CNPQ) grant

#429894/2016-3 (TLF). Students BSM and BdSV were partially funded by grants from

Universidade Federal do ABC (UFABC).

563   **Author Statement**

MdSC helped conceptualizing, curating data, doing formal analysis and experiments,

planning methodology and writing the original draft. BdSV and BSM helped doing experiments

and reviewing the manuscript. PAT and TLF helped reviewing the manuscript. RVF helped

conceptualizing, acquiring funds, planning methodology, administrating and supervising the

project and reviewing the manuscript.

References

Abrari, K., Rashidy-Pour, A., Semnanian, S., Fathollahi, Y., 2009. Post-training administration
of corticosterone enhances consolidation of contextual fear memory and hippocampal
long-term potentiation in rats. Neurobiol. Learn. Mem. 91, 260–265.
https://doi.org/10.1016/j.nlm.2008.10.008

Addis, D.R., Schacter, D.L., 2008. Constructive episodic simulation: temporal distance and
detail of past and future events modulate hippocampal engagement. Hippocampus 18,
227–37. https://doi.org/10.1002/hipo.20405

Atucha, E., Roozendaal, B., 2015. The inhibitory avoidance discrimination task to investigate
accuracy of memory. Front. Behav. Neurosci. 9, 60.
https://doi.org/10.3389/fnbeh.2015.00060

Atucha, E., Vukojevic, V., Fornari, R. V., Ronzoni, G., Demougin, P., Peter, F., Atsak, P.,
Coolen, M.W., Papassotiropoulos, A., McGaugh, J.L., de Quervain, D.J.-F., Roozendaal,
B., 2017. Noradrenergic activation of the basolateral amygdala maintains hippocampus-
dependent accuracy of remote memory. Proc. Natl. Acad. Sci. 201710819.
https://doi.org/10.1073/pnas.1710819114

Bahtiyar, S., Karaca, K.G., Henckens, M.J.A.G.A.G., Roozendaal, B., 2020. Norepinephrine
and glucocorticoid effects on the brain mechanisms underlying memory accuracy and
generalization. Mol. Cell. Neurosci. 108, 103537.
https://doi.org/10.1016/j.mcn.2020.103537

Binder, J.R., Desai, R.H., Graves, W.W., Conant, L.L., 2009. Where is the semantic system? A
critical review and meta-analysis of 120 functional neuroimaging studies. Cereb. Cortex
19, 2767–2796. https://doi.org/10.1093/cercor/bhp055

Bueno, A.P.A., de Paiva, J.P.Q., Corrêa, M. dos S., Tiba, P.A., Fornari, R.V., 2017.
Corticosterone administration after a single-trial contextual fear conditioning does not
influence the strength and specificity of recent and remote memory in rats. Physiol. Behav.
171, 175–180. https://doi.org/10.1016/j.physbeh.2017.01.011

Christianson, S., Loftus, E.F., 1987. Memory for traumatic events. Appl. Cogn. Psychol. 1,
225–239. https://doi.org/10.1002/acp.2350010402

Clewett, D., Murty, V., 2019. Echoes of emotions past: How neuromodulators determine what
we recollect. Eneuro 6, ENEURO.0108-18.2019. https://doi.org/10.1523/ENEURO.0108-
18.2019

Cordero, M.I., Sandi, C., 1998. A role for brain glucocorticoid receptors in contextual fear
conditioning: dependence upon training intensity. Brain Res. 786, 11–17.
https://doi.org/S0006-8993(97)01420-0 [pii]

Dandolo, L.C., Schwabe, L., 2016. Stress-induced cortisol hampers memory generalization.
Learn. Mem. 23, 679–683. https://doi.org/10.1101/lm.042929.116

de Paiva, J.P.Q., Bueno, A.P.A., Dos Santos Corrêa, M., Oliveira, M.G.M., Ferreira, T.L.,
Fornari, R. V, 2021. The posterior insular cortex is necessary for the consolidation of tone
fear conditioning. Neurobiol. Learn. Mem. 179, 107402.
https://doi.org/10.1016/j.nlm.2021.107402

de Quervain, D., Schwabe, L., Roozendaal, B., 2016. Stress, glucocorticoids and memory:
Implications for treating fear-related disorders. Nat. Rev. Neurosci. 18, 7–19.
https://doi.org/10.1038/nrn.2016.155

dos Santos Corrêa, M., Vaz, B. dos S., Grisanti, G.D.V., de Paiva, J.P.Q., Tiba, P.A., Fornari,
R.V., 2019. Relationship between footshock intensity, post-training corticosterone release
and contextual fear memory specificity over time. Psychoneuroendocrinology 110,
104447. https://doi.org/10.1016/j.psyneuen.2019.104447

Joëls, M., Fernandez, G., Roozendaal, B., 2011. Stress and emotional memory: A matter of
timing. Trends Cogn. Sci. 15, 280–288. https://doi.org/10.1016/j.tics.2011.04.004

Kaouane, N., Porte, Y., Vallée, M., Brayda-Bruno, L., Mons, N., Calandreau, L., Marighetto,
A., Piazza, P.V., Desmedt, A., 2012. Glucocorticoids can induce PTSD-like memory
impairments in mice. Science 335, 1510–3. https://doi.org/10.1126/science.1207615

Kensinger, E.A., Gutchess, A.H., Schacter, D.L., 2007. Effects of Aging and Encoding
Instructions on Emotion-Induced Memory Trade-Offs. Psychol. Aging 22, 781–795.
https://doi.org/10.1037/0882-7974.22.4.781

Lenaert, B., Boddez, Y., Vervliet, B., Schruers, K., Hermans, D., 2015. Reduced
Autobiographical Memory Specificity is Associated With Impaired Discrimination
Learning in Anxiety Disorder Patients. Front. Psychol. 6, 1–12.
https://doi.org/10.3389/fpsyg.2015.00889

Lissek, S., Bradford, D.E., Alvarez, R.P., Burton, P., Espensen-Sturges, T., Reynolds, R.C.,
Grillon, C., 2014a. Neural substrates of classically conditioned fear-generalization in
humans: A parametric fMRI study. Soc. Cogn. Affect. Neurosci. 9, 1134–1142.
https://doi.org/10.1093/scan/nst096

Lissek, S., Kaczkurkin, A.N., Rabin, S., Geraci, M., Pine, D.S., Grillon, C., 2014b. Generalized
anxiety disorder is associated with overgeneralization of classically conditioned fear. Biol.
Psychiatry 75, 909–915. https://doi.org/10.1016/j.biopsych.2013.07.025

Marchand, A.R., Barbelivien, A., Seillier, A., Herbeaux, K., Sarrieau, A., Majchrzak, M., 2007.
Contribution of corticosterone to cued versus contextual fear in rats. Behav. Brain Res.
183, 101–110. https://doi.org/10.1016/j.bbr.2007.05.034

McGlade, A.L., Zbozinek, T.D., Treanor, M., Craske, M.G., 2019. Pilot for novel context
generalization paradigm. J. Behav. Ther. Exp. Psychiatry 62, 49–56.
https://doi.org/10.1016/j.jbtep.2018.08.009

Moreira-Silva, D., Carrettiero, D.C., Oliveira, A.S.A., Rodrigues, S., Dos Santos-Lopes, J.,
Canas, P.M., Cunha, R.A., Almeida, M.C., Ferreira, T.L., 2018. Anandamide Effects in a
Streptozotocin-Induced Alzheimer's Disease-Like Sporadic Dementia in Rats. Front.
Neurosci. 12, 653. https://doi.org/10.3389/fnins.2018.00653

Moscovitch, M., Cabeza, R., Winocur, G., Nadel, L., 2016. Episodic memory and beyond: The
hippocampus and neocortex in transformation. Annu. Rev. Psychol. 67, 105–134.
https://doi.org/10.1146/annurev-psych-113011-143733

Myers, C.E., Shohamy, D., Gluck, M.A., Grossman, S., Onlaor, S., Kapur, N., 2003.
Dissociating medial temporal and basal ganglia memory systems with a latent learning
task. Neuropsychologia 41, 1919–1928. https://doi.org/10.1016/S0028-3932(03)00127-1

Pedraza, L.K., Sierra, R.O., Boos, F.Z., Haubrich, J., Quillfeldt, J.A., de Oliveira Alvares, L.,
2016. The dynamic nature of systems consolidation: Stress during learning as a switch
guiding the rate of the hippocampal dependency and memory quality. Hippocampus 26,
362–371. https://doi.org/10.1002/hipo.22527

Pugh, C.R., Tremblay, D., Fleshner, M., Rudy, J.W., 1997. A selective role for corticosterone in
contextual-fear conditioning. Behav. Neurosci. 111, 503–511.

https://doi.org/10.1037/0735-7044.111.3.503

Quaedflieg, C.W.E.M., Schwabe, L., 2018. Memory dynamics under stress. Memory 26, 364–
376. https://doi.org/10.1080/09658211.2017.1338299

Quirarte, G.L., Galvez, R., Roozendaal, B., McGaugh, J.L., 1998. Norepinephrine release in the
amygdala in response to footshock and opioid peptidergic drugs. Brain Res. 808, 134–40.
https://doi.org/10.1016/s0006-8993(98)00795-1

Rice, J.B., White, A.G., Scarpati, L.M., Wan, G., Nelson, W.W., 2017. Long-term Systemic
Corticosteroid Exposure: A Systematic Literature Review. Clin. Ther. 39, 2216–2229.
https://doi.org/10.1016/j.clinthera.2017.09.011

Rimmele, U., Davachi, L., Petrov, R., Dougal, S., Phelps, E.A., 2011. Emotion Enhances the
Subjective Feeling of Remembering, Despite Lower Accuracy for Contextual Details.
Emotion 11, 553–562. https://doi.org/10.1037/a0024246

Roozendaal, B., Hermans, E.J., 2017. Norepinephrine effects on the encoding and consolidation
of emotional memory: improving synergy between animal and human studies. Curr. Opin.
Behav. Sci. 14, 115–122. https://doi.org/10.1016/j.cobeha.2017.02.001

Roozendaal, B., McEwen, B.S., Chattarji, S., 2009. Stress, memory and the amygdala. Nat. Rev.
Neurosci. 10, 423–433. https://doi.org/10.1038/nrn2651

Roozendaal, B., Mirone, G., 2020. Opposite Effects of Noradrenergic and Glucocorticoid
Activation on Accuracy of Episodic-like Memory. Psychoneuroendocrinology 104588.
https://doi.org/10.1016/j.psyneuen.2020.104588

Rusu, S.I., Lankelma, J. V., Jackson, J.C., Van Mourik-Donga, L.A., Joëls, M., Pennartz,
C.M.A., 2018. Corticosterone impairs flexible adjustment of spatial navigation in an
associative place-reward learning task. Behav. Pharmacol. 29, 351–364.
https://doi.org/10.1097/FBP.0000000000000370

Schwabe, L., 2017. Memory under stress: from single systems to network changes. Eur. J.
Neurosci. 45, 478–489. https://doi.org/10.1111/ejn.13478

Schwabe, L., Böhringer, A., Wolf, O.T., 2009. Stress disrupts context-dependent memory.
Learn. Mem. 16, 110–113. https://doi.org/10.1101/lm.1257509

Schwabe, L., Wolf, O.T., 2012. Stress modulates the engagement of multiple memory systems
in classification learning. J. Neurosci. 32, 11042–11049.
https://doi.org/10.1523/JNEUROSCI.1484-12.2012

Segal, S.K., Stark, S.M., Kattan, D., Stark, C.E., Yassa, M.A., 2012. Norepinephrine-mediated
emotional arousal facilitates subsequent pattern separation. Neurobiol. Learn. Mem. 97,
465–469. https://doi.org/10.1016/j.nlm.2012.03.010

Shohamy, D., Wagner, A.D., 2008. Integrating Memories in the Human Brain: Hippocampal-
Midbrain Encoding of Overlapping Events. Neuron 60, 378–389.
https://doi.org/10.1016/j.neuron.2008.09.023

Timmer, M., Sandi, C., 2010. A role for glucocorticoids in the long-term establishment of a
social hierarchy. Psychoneuroendocrinology 35, 1543–1552.
https://doi.org/10.1016/j.psyneuen.2010.05.011

Tonegawa, S., Morrissey, M.D., Kitamura, T., 2018. The role of engram cells in the systems
consolidation of memory. Nat. Rev. Neurosci. 19, 485–498.
https://doi.org/10.1038/s41583-018-0031-2

van Ast, V.A., Cornelisse, S., Marin, M.F., Ackermann, S., Garfinkel, S.N., Abercrombie, H.C.,
2013. Modulatory mechanisms of cortisol effects on emotional learning and memory:
Novel perspectives. Psychoneuroendocrinology 38, 1874–1882.
https://doi.org/10.1016/j.psyneuen.2013.06.012

Van Ast, V.A., Cornelisse, S., Meeter, M., Kindt, M., 2014. Cortisol mediates the effects of
stress on the contextual dependency of memories. Psychoneuroendocrinology.
https://doi.org/10.1016/j.psyneuen.2013.12.007

Vogel, S., Fernández, G., Joëls, M., Schwabe, L., 2016. Cognitive Adaptation under Stress: A
Case for the Mineralocorticoid Receptor. Trends Cogn. Sci. 20, 192–203.
https://doi.org/10.1016/j.tics.2015.12.003

Wiltgen, B.J., Zhou, M., Cai, Y., Balaji, J., Karlsson, M.G., Parivash, S.N., Li, W., Silva, A.J.,
2010. The hippocampus plays a selective role in the retrieval of detailed contextual
memories. Curr. Biol. 20, 1336–1344. https://doi.org/10.1016/j.cub.2010.06.068

CORT-HBC
or Saline

Priming Session

Test Session

a

Freezing %

- Context B
- Context A

#

Saline | CORT-HBC 0.25 mg/kg | CORT-HBC 0.5 mg/kg | CORT-HBC 1 mg/kg | CORT-HBC 2 mg/kg | CORT-HBC 4 mg/kg | CORT-HBC 8 mg/kg

b

Generalization Index

$\chi^2(6,47) = 12.68$, $p = 0.048*$

Saline | CORT-HBC | CORT-HBC | CORT-HBC | CORT-HBC | CORT-HBC | CORT-HBC

c

% of Animals

Generalizers | Discriminators *

Saline | CORT-HBC 0.25 mg/kg | CORT-HBC 0.5 mg/kg | CORT-HBC 1 mg/kg | CORT-HBC 2 mg/kg | CORT-HBC 4 mg/kg | CORT-HBC 8 mg/kg