Chimeric spike mRNA vaccines protect against sarbecovirus challenge in mice

David R. Martinez [1, *,] Alexandra Schaefer [1,] Sarah R. Leist [1], Gabriela De la Cruz [2], Ande West[1], Elena N. Atochina-Vasserman [4], Robert Parks [5], Maggie Barr [5], Dapeng Li [5], Boyd Yount [1], Kevin O. Saunders [5], Drew Weissman [4], Barton F. Haynes [5], Stephanie A. Montgomery [3], Ralph S. Baric [1, *].

[1] Department of Epidemiology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

[2] Lineberger Comprehensive Cancer Center, University of North Carolina School of Medicine, Chapel Hill, NC, USA

[3] Department of Laboratory Medicine and Pathology, University of North Carolina School of Medicine, Chapel Hill, NC, USA

[4] Infectious Disease Division, Department of Medicine, Perelman School of Medicine, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, USA

[5] Duke Human Vaccine Institute, Duke University School of Medicine, Durham, NC, USA

[*]Corresponding authors.

David R. Martinez, D.R.M., david.rafael.martinez@gmail.com
Ralph S. Baric, R.S.B., rbaric@email.unc.edu

Keywords: SARS-CoV-2, SARS-like virus, *Sarbecovirus*, mRNA vaccine, universal coronavirus vaccine.

Abstract

The emergence of SARS-CoV and SARS-CoV-2 in the 21st century highlights the need to develop universal vaccination strategies against the SARS-related *Sarbecovirus* subgenus. Using structure-guided chimeric spike designs and multiplexed immunizations, we demonstrate protection against SARS-CoV, SARS-CoV-2, and bat CoV (BtCoV) RsSHC014 challenge in highly vulnerable aged mice. Chimeric spike mRNAs containing N-terminal domain (NTD), and

receptor binding domains (RBD) induced high levels of broadly protective neutralizing

antibodies against three high-risk sarbecoviruses: SARS-CoV, RsSHC014, and WIV-1. In

contrast, SARS-CoV-2 mRNA vaccination not only showed a 10 to >500-fold reduction in

neutralizing titers against heterologous sarbecovirus strains, but SARS-CoV challenge in mice

resulted in breakthrough infection including measurable lung pathology. Importantly, chimeric

spike mRNA vaccines efficiently neutralized both the D614G and the South African B.1.351

variants of concern despite some reduction in neutralization activity. Thus, multiplexed-chimeric

spikes may provide a novel strategy to prevent pandemic and SARS-like zoonotic coronavirus

infections, while revealing the limited efficacy of SARS-CoV-2 spike vaccines against other

sarbecoviruses.

Introduction

A novel coronavirus that caused severe acute respiratory syndrome (SARS-CoV)

emerged in 2003 and caused more than 8,000 infections and ~800 deaths worldwide (*1*). Less

than a decade later, another coronavirus that caused Middle East Respiratory Syndrome (MERS-

CoV) emerged in Saudi Arabia in 2012 (*2*), resulting in an ongoing outbreak with at least ~2,600

cases and 900 deaths (*3*). In December 2019, another novel human SARS-like virus from the

genus *Betacoronavirus* and subgenus *Sarbecocovirus* emerged in Wuhan China, designated

SARS-CoV-2 (*4, 5*). The emergence and spread of SARS-CoV-2, the causative agent of

coronavirus disease 2019 (COVID-19), was explosive. By March 2020, the World Health

Organization (WHO) had declared COVID-19 a global pandemic. By March 2021, more than

117 million COVID-19 human infections had resulted in more than 2.6 million deaths globally in

an ever-expanding pandemic (*6*).

Zoonotic transmission of animal coronaviruses of probable bat origin into naïve animal

and human populations remains a clear One Health threat (*7-10*). Broadly protective vaccines

and therapeutic monoclonal antibodies are desperately needed to mitigate the risk of animal

coronavirus zoonosis into naïve species including humans. Bats are known reservoirs of SARS-

like coronaviruses (CoVs) and harbor high-risk "pre-emergent" SARS-like variant strains, such

as WIV-1 and RsSHC014, which are able to utilize human ACE2 receptors for entry, replicate

efficiently in primary airway epithelial cells, and may escape existing countermeasures (*9-13*)

Given the high pandemic potential of zoonotic and epidemic sarbecoviruses (*13*), the

development of broadly effective countermeasures, such as universal vaccination strategies,

antibodies and drugs is a global health priority (*14-17*). However, a major hurdle to develop

vaccines that can neutralize diverse sarbecoviruses is the vast genetic diversity that exists within

critical immunodominant epitope landscapes, like the receptor binding domain (RBD) (*13*). As

COVID-19 convalescent patients exhibit little cross-neutralization of other human pathogenic

viruses including SARS-CoV and MERS-CoV (*18, 19*), it is uncertain whether natural immunity

to SARS-CoV-2 will protect against other sarbecoviruses of zoonotic origin that may emerge in

the future. As double-inactivated SARS-CoV vaccines in aged mice reported increased

eosinophilic infiltration in the lung following heterologous sarbecovirus challenge, it is critical to

examine heterologous challenge in the context of sarbecovirus vaccination (*20, 21*). This

particularly important as high levels of neutralizing antibodies are required for protecting against

vaccine-associated enhanced respiratory disease (VAERD) following heterologous SARS-CoV

challenge in aged mice.

The RBD is a target for neutralizing antibodies elicited in the context of natural SARS-

CoV-2 and MERS-CoV infections (*22-25*). In addition to the RBD, the N-terminal domain

(NTD) is also a target for SARS-CoV-2 and MERS-CoV neutralizing antibodies (*23, 26, 27*).

Passive immunization with SARS-CoV-2 NTD-specific antibodies can protect naïve mice from

challenge, demonstrating that the NTD is a target of protective immunity (*28*). However, it

remains unclear if vaccine-elicited neutralizing antibodies induced by vaccines can protect

against *in vivo* challenge with epidemic and bat coronaviruses. Here, we provide the proof-of-

concept that nucleoside-modified mRNA-lipid nanoparticle (LNP) vaccines expressing chimeric

spikes containing admixtures of RBD and NTD domains from epidemic and pandemic

sarbecoviruses can: 1) elicit robust neutralizing antibody responses against multiclade

sarbecoviruses, and 2) protect against viral replication in upper and lower respiratory airways of

both epidemic and pandemic human coronaviruses in aged mice, and 3) attenuate VAERD

responses associated with homologous virus challenge. Our findings suggest that mRNA-LNP

vaccination with chimeric CoV spikes is a viable strategy to protect against contemporary and

high-risk sarbecovirus emergence events.

Results

Design and expression of chimeric spike constructs to cover pandemic and zoonotic SARS-

related coronaviruses

Sarbecoviruses exhibit considerable genetic diversity (Fig. 1A) and several bat CoVs (Bt-

CoVs) that are SARS-like exist in nature (*13*). As previous studies have demonstrated that Bt-

CoVs, including WIV-1 and RsSHC014, are poised for emergence (*9, 10*), we sought to design

vaccination strategies that can protect against antigenically diverse, epidemic, pandemic, and

pre-pandemic CoVs isolated from humans or *Rhinolophus* bats. Harnessing the modular design

of CoV spikes (*29*), we designed chimeric spikes to create within-spike "bivalent" and

"trivalent" vaccines that have the potential to elicit protective antibody responses against more

than one virus within a subgenus (e.g., *Sarbecovirus*). As the NTD and RBD are both the target

of binding and neutralizing antibodies, we used domain-focused immunogen design, which

admixes major neutralizing antigenic sites (e.g., NTD, RBD, S2) into new spike modules from

clade I, II, and III sarbecoviruses designed to maximize breadth while focusing immunity against

broadly conserved high-risk spike domains. To achieve vaccination strategies that can overcome

the broad genetic diversity of SARS-like viruses, we designed four sets of chimeric spike

constructs that contained unique admixtures of the RBD and/or NTD neutralizing domains from

various pandemic and zoonotic SARS-like viruses. Chimera 1 included the NTD from Bt-CoV

Hong Kong University 3-1 (HKU3-1), the SARS-CoV RBD, and the SARS-CoV-2 subunit 2

(S2) (Fig. S1A). Chimera 2 included SARS-CoV-2 RBD and SARS-CoV NTD and S2. Chimera

3 included the SARS-CoV RBD, SARS-CoV-2 NTD and S2. Finally, chimera 4 included the

RsSHC014 RBD, SARS-CoV-2 NTD and S2. We also generated a monovalent SARS-CoV-2

spike furin knock out (KO) vaccine, partially phenocopying the Johnson & Johnson, Moderna,

and Pfizer vaccines that recently received emergency use authorization by the U.S. FDA, and a

negative control norovirus GII capsid vaccine (Fig. S1A to F). We generated these chimeric

spikes and control spikes as lipid nanoparticle-encapsulated, nucleoside-modified mRNA

(mRNA-LNP) vaccines as described previously (*30*), and verified their expression in HEK cells

(Fig. S1G). To confirm that coronavirus spikes are modular in design allowing for rationale

admixing of component parts, we also designed and recovered a panel of high titer recombinant

live viruses between RsSHC014 nanoluciferase ΔORF7&8 that harbored motifs from the SARS-

CoV-2 S1, NTD, RBD and S2 domains. After infection *in vitro*, these high-titer recombinant

viruses expressed high levels of nanoluciferase, demonstrating that the chimeric spikes promoted

efficient entry and replication (Fig. S1H).

Immunogenicity of mRNAs expressing chimeric spike constructs against coronaviruses

As previous studies demonstrated the successful delivery of more than one HSV-2

glycoprotein to broaden antigen coverage via mRNA vaccination (*31-33*), we sought to

determine if simultaneous immunization with mRNA-LNP expressing the chimeric spikes of

diverse sarbecoviruses was a feasible strategy to elicit broadly binding and neutralizing

antibodies. As aged human populations are the most susceptible to severe COVID-19 disease and

death, we sought to examine the protective efficacy of our vaccines in aged mouse models that

develop severe disease and death following sarbecovirus infection. We immunized aged mice

with the chimeric spikes formulated to induce cross reactive responses to the NTD and RBD

against multiple divergent clade I-III sarbecoviruses, and/or focused boosting of antibody

responses against epidemic strains, a SARS-CoV-2 furin KO spike, and a GII.4 norovirus capsid

negative control. Group 1 was primed and boosted with chimeric spikes 1, 2, 3, and 4 (Fig. 1B).

Group 2 was primed with chimeric spikes 1 and 2 and boosted with chimeric spikes 3 and 4.

Group 3 was primed and boosted with chimeric spike 4. Group 4 was primed and boosted with

the monovalent SARS-CoV-2 furin knockout spike. Finally, group 5 was primed and boosted

with a norovirus capsid GII.4 Sydney 2011 strain (Fig. 1B). To examine the magnitude and cross

binding responses of the chimeric spikes compared to the SARS-CoV-2 monovalent vaccine, we

examined the binding antibody responses by ELISA against a diverse panel of CoV spike

proteins that included contemporary, epidemic, pandemic, and zoonotic coronaviruses. While

pre-immunization mouse serum samples did not bind to any of the coronavirus ELISA panel

antigens, we observed high binding to SARS-CoV-2 and SARS-like spike antigens in several of

the vaccinated groups (Fig 2 and Fig. S2). Mice immunized with chimeric spikes 1-4 either in

combination or separately in the prime/boost generated the highest magnitude responses to

SARS-CoV Toronto Canada isolate (Tor2) and HKU3-1 spike compared to mice immunized

with chimera 4 and a SARS-CoV-2 furin KO spike (Fig 2A and 2H), demonstrating that

immunization with multiplexed chimeric spikes elicits higher magnitude and more cross-reactive

antibody profiles than immunization with a monovalent SARS-CoV-2 spike mRNA-LNP. While

mice primed with chimeras 1-2 and boosted with chimeras 3-4 generated lower magnitude

binding responses to both SARS-CoV-2 RBD (Fig. 2C) and SARS-CoV-2 NTD (Fig 2D), mice

immunized with chimeras 1-4 in the prime and boost generated similar magnitude binding

antibodies to SARS-CoV-2 D614G compared to mice immunized with the SARS-CoV-2 furin

KO spike mRNA-LNP (Fig 2B). Mice immunized with the chimeric spikes 1-4 all together or

separately in the prime and boost generated similar magnitude binding antibody responses

against SARS-CoV-2 D614G, Pangolin GXP4L, and RaTG13 spikes (Fig. 2B, 2E, and 2F)

compared to mice immunized with SARS-CoV-2 spike, underlining that the chimeric spikes not

only generate indistinguishable binding responses against SARS-CoV-2-like viruses compared to

monovalent SARS-CoV-2 vaccines but also elicit higher responses against SARS-like viruses at

high risk of emergence. Mice immunized with chimeras 1-4 in the prime and boost generated the

highest magnitude binding antibodies to RsSHC014 spike (Fig. 2G). Importantly, mice

immunized with the norovirus capsid mRNA-LNP vaccine did not generate binding IgG

antibodies to any of the tested CoV protein antigens (Fig 2 and Fig. S2). In agreement with the

binding ELISA data, mice immunized with chimeras 1-4 in the prime and boost elicited

indistinguishable magnitude levels of hACE2 blocking responses to the SARS-CoV-2 spike

immunized mice (Fig. 2J). In contrast, mice from groups 2 and 3 elicited lower magnitude

blocking responses against hACE2 compared to mice from groups 1 and 4 (Fig. 2J). Finally, we

did not observe cross-binding antibodies against common-cold CoV spike antigens from HCoV-

HKU1, HCoV-NL63, and HCoV-229E, in most of the vaccine groups (Fig. S2A-2D), but we did

observe low binding levels against more distant group 2C MERS-CoV (Fig. 2I) and

Betacoronaviruses like group 2A HCoV-OC43 in vaccinated mice from groups 1 and 2 (Fig.

S2B). Altogether, these results suggest that chimeric spike mRNA vaccines elicit more broad and

higher magnitude binding responses against pandemic and bat SARS-like viruses compared to

monovalent SARS-CoV-2 spike furin KO mRNA-LNP vaccines.

Neutralizing antibody responses against live sarbecoviruses and variants of concern

We then tested the neutralizing antibody responses against SARS-CoV, SARS-CoV-2,

Bt-CoV RsSHC014, and BtCoV WIV-1, using live viruses expressing nanoluciferase as

previously described (Fig 3A-3D) (*18*). Group 4 SARS-CoV-2 S mRNA vaccinated animals

mounted a robust response against SARS-CoV-2, however responses against SARS-CoV,

RsSHC014, and WIV-1 were 18-, >500- or 116-fold more resistant, respectively, demonstrating

the importance of including different high-risk pre-emergent Bt-sarbecovirus strains in these

studies (Fig 3A-3D and Fig. S3G-H). In contrast, aged mice that were primed with chimeras 1-2

and boosted with chimeras 3-4 showed a 42- and 2-fold increase in neutralizing titer against

SARS-CoV and WIV1, and less than 1-fold decrease against RsSHC014 relative to SARS-CoV-

2 neutralizing titers (Fig 3A-3D and Fig. S3C-D). The chimera 4 vaccine, which contained the

RsSHC014 RBD/SARS-CoV-2 NTD and S2 domains, elicited 3-, 7-fold higher neutralizing

titers against SARS-CoV and RsSHC014 yet showed a 3-fold reduction in WIV-1 neutralizing

titles relative to its SARS-CoV-2 neutralizing activity (Fig 3A-3D and Fig. S3E-F). Finally, mice

immunized with chimeras 1-4 as prime and boost together generated the most balanced and

highest neutralizing titers that were 13- and 1.2-fold higher against SARS-CoV and WIV-1 and

only less than 1-fold lower against RsSHC014 relative to the SARS-CoV-2 neutralizing titers in

this group (Fig 3A-3D and Fig. S3A-B). We also examined the neutralizing activity of mice

primed and boosted with chimeras 1-4 vs. the monovalent SARS-CoV-2 vaccine against two

variants of concern (VOC): the predominant D614G variant and the B.1.351 South African

variant. The serum of aged mice immunized with the multiplexed chimeras and the monovalent

SARS-CoV-2 vaccine neutralized the dominant D614G variant with similar potency as the wild

type D614 non-predominant variant (Fig. 3E). Mouse sera from group 1 primed and boosted with

chimeras 1-4 only had a 2-fold reduction in neutralizing activity against the B.1.351 South

African variant of concern, whereas a 4-fold reduction was observed in the SARS-CoV-2 furin

KO immunized mouse sera (Fig. 3F). Despite the significant but small reduction in neutralizing

activity against the B.1.351 variant, we did not observe a complete ablation in neutralizing

activity in either group. Thus, both monovalent SARS-CoV-2 vaccines and multiplexed chimeric

spikes elicit neutralizing antibodies against newly emerged SARS-CoV-2 variants and

multiplexed chimeric spike vaccines outperform the monovalent SARS-CoV-2 vaccines in terms

of breadth of potency against multiclade sarbecoviruses.

In vivo **protection against epidemic and pandemic coronavirus challenge**

To assess the ability of the mRNA-LNP vaccines in mediating protection against

previously epidemic SARS-CoV, pandemic SARS-CoV-2, and Bt-CoVs, we challenged the

different groups and observed the mice for signs of clinical disease. Group 1 mice (chimeras 1-4

prime/boost) or group 2 (chimeras 1-2 prime, 3-4 boost) were completely protected from weight

loss, lower, and upper airway virus replication as measured by infectious virus plaque assays

following SARS-CoV mouse-adapted (MA15) challenge (Fig. 4A, 4B and 4C). Similarly, these

two vaccine groups were also protected against SARS-CoV-2 mouse-adapted (MA10). In

contrast, groups 3 (chimera 4) showed some protection against SARS-CoV MA15 induced

weight loss, but not against viral replication in the lung or nasal turbinates, and full protection

against SARS-CoV-2 MA10. In contrast, norovirus capsid control mice developed severe disease

including mortality in both SARS-CoV MA15 and SARS-CoV-2 MA10 infections (Fig. S5).

Monovalent SARS-CoV-2 mRNA vaccines were highly efficacious against SARS-CoV-2 MA10

challenge but failed to protect against SARS-CoV MA15-induced weight loss, and replication in

the lower and upper airway (Fig. 4A, 4B, and 4C), suggesting that SARS-CoV-2 mRNA-LNP

vaccines currently being administered during the COVID-19 pandemic are not likely to protect

against future SARS-CoV emergence events in vulnerable populations. Mice from all chimeric

spike immunization groups, including group 4 which received the SARS-CoV-2

NTD/RsSHC014 RBD chimeric spike, and SARS-CoV-2 furin KO vaccine groups were

completely protected from weight loss and lower airway SARS-CoV-2 MA10 replication (Fig.

4D, 4E, and 4F). In addition, using the BtCoV RsSHC014 replication model in mice, we also

demonstrated protection against RsSHC014 replication in the lung and nasal turbinates (Fig. S4)

in mice that received the multiplexed chimeras 1-4 as prime and boost and in mice that

immunized with the SARS-CoV-2 NTD/RsSHC014 RBD chimeric spike, demonstrating the

breadth of the universal spike vaccine formulations and the ability to introduce "bivalency" into

CoV spike to increase protection. Our findings also suggest that SARS-CoV-2 mRNA-LNP

vaccines do not protect against clinical disease of SARS-CoV MA15 infection in mice.

Lung pathology and cytokines in mRNA-LNP vaccinated mice challenged with epidemic

and pandemic coronaviruses

Lung discoloration is the gross manifestation of various processes of acute lung damage,

including congestion, edema, hyperemia, inflammation, and protein exudation. We used this

macroscopic scoring scheme to visually score mouse lungs at the time of harvest. To quantify the

pathological features of acute lung injury (ALI) in mice, we used a tool from the American

Thoracic Society (ATS - Matute-Bello lung pathology score). We have previously used this tool

to describe the pulmonary pathogenesis in BALB/c mice infected with SARS-CoV-2 MA10 (*34,*

35). To quantify microscopic differences in lung pathology, we used this tool on three random

diseased fields in lung tissue sections per mouse and tissues were blindly evaluated by a board-

certified veterinary pathologist (Fig. 5B). With a complementary histological quantitation tool,

we similarly scored lung tissue sections for diffuse alveolar damage (DAD), the pathological

hallmark of ALI (cellular sloughing, necrosis, hyaline membranes, etc.) (*36, 37*) and found these

data were generally consistent with those from the lung discoloration scores. We observed

significant lung pathology by both the Matute-Bello and DAD scoring tools in groups 4 and 5

vaccinated animals, consistent with the weight loss, lung titer, and lung cytokine data after

heterologous SARS-CoV MA15 challenge. In contrast, multiplexed chimeric spike vaccine

formulations in groups 1 and 2 provided complete protection from lung pathology after SARS-

CoV MA15 challenge (Fig. 5A and 5B). Concerningly, mice immunized with the SARS-CoV-2

spike that showed breakthrough infection with SARS-CoV MA15 developed severe lung

inflammation, potentially suggesting that future outbreaks of SARS-CoV may lead to lung

pathology in individuals vaccinated with SARS-CoV-2. In contrast to the heterologous SARS-

CoV MA15 challenge, all groups challenged with SARS-CoV-2 MA10 were protected against

lung pathology compared to the norovirus capsid-immunized control group. Mice immunized

with the multiplexed chimeric spikes alone or in combination, and also with the SARS-CoV-2

NTD/RsSHC014 chimera, and with SARS-CoV-2 were also completely protected against both

macroscopic and microscopic lung damage (Fig 5C and 5D).

To examine the level of protection in the lung of SARS-CoV and SARS-CoV-2

challenged mice in more detail, we measured lung proinflammatory cytokines and chemokines in

the different vaccination groups. Group 1 mice (chimeras 1-4 prime/boost) and group 2

(chimeras 1-2 prime, 3-4 boost) mice following SARS-CoV MA15 challenge had baseline levels

of macrophage activating cytokines and chemokines including, IL-6, CCL2, IL-1α, G-SCF, and

CCL4, compared to group 5 (norovirus prime/boost) (Fig. S6A). In contrast, group 3 (chimera 4

prime/boost) and group 4 (monovalent SARS-CoV-2 prime/boost) mice following SARS-CoV

MA15 challenge, showed high and indistinguishable levels of IL-6, CCL2, IL-1α, G-SCF, and

CCL4 compared to group 5. Following SARS-CoV-2 MA10 challenge, group 4 and group 1

showed the lowest levels of IL-6, and G-SCF relative to group 5 controls (Fig. S6B), and we

only observed significant reductions in CCL2, IL-1α, CCL4 lung levels in groups 3 and 4

compared to the group 5 control despite full protection from weight loss, lower, and some level

of upper airway replication in groups 1-4. These results suggest that SARS-CoV-2 vaccination

does not protect against SARS-CoV challenge and underlines that universal vaccination

strategies that can protect against heterologous sarbecovirus strains at high risk for human

emergence are needed.

Discussion

The mRNA-1273 vaccine developed by Moderna Inc. demonstrated protection against

SARS-CoV-2 challenge both in mice and in non-human primates (*38, 39*), but there is a growing

concern due to the emergence of VOCs like South African B1.351 which is 2-6 fold more

resistant to vaccine-elicited polyclonal neutralizing antibodies (*40*). Both the Moderna and

Pfizer/BioNTech mRNA-LNP vaccines were safe and efficacious against SARS-CoV-2

infections in large Phase 3 efficacy human clinical trials (*41-43*), and the Moderna mRNA-1273

vaccine was also immunogenic in aged adults (*44*), which are highly vulnerable for severe

COVID-19 symptoms and death. Given the rapid development and success of the mRNA-LNP

vaccine platform, we sought to replicate this existing platform to establish the proof-of-concept

of vaccine protection against sarbecoviruses in aged mouse models that recapitulate extreme

vulnerability of aged human populations. Consistent with previous studies that measured efficacy

of mRNA-LNP vaccines for SARS-CoV-2 vaccines in mice (*38*), our monovalent SARS-CoV-2

vaccine elicited robust neutralizing antibody titers to the SARS-CoV-2 D614G variant. We

observed a 2-fold reduction in neutralizing antibodies in mice immunized with the multiplexed-

chimeric spikes and a 4-fold reduction in neutralizing antibodies in SARS-CoV-2 monovalent

vaccinated mice against the South African B.1.351 VOC. While monovalent SARS-CoV-2

vaccinated mice showed an 18 to >500-fold reduction in neutralization antibody activity against

clade I and clade III sarbecoviruses, multiplexed chimeric spike immunizations showed

improved neutralizing antibody activity against these epidemic and zoonotic sarbecoviruses (Fig.

3 and Fig. S3). The lack of protection against SARS-CoV challenge in SARS-CoV-2 immunized

mice underlines the need for the development of universal vaccination strategies that can achieve

broader coverage against pre-emergent bat SARS-CoV-like and SARS-CoV-2-like viruses. One

such strategy, as described herein, is to admix key antigenic sites (e.g., NTD and RBD) from

different sarbecoviruses into new chimeric spikes that are designed to enhance breadth while

focusing the neutralization responses on key antigenic sites conserved across multiple strains

(e.g. S2). A caveat of including multiple chimeric spikes in a single shot is the potential

formation of heterotrimers not present in the intended vaccine formulation. While it remains

unknown if our chimeric mRNA-LNP vaccines generate heterotrimers *in vivo*, the robustness of

the cross-neutralizing titers against sarbecoviruses and protection against SARS-CoV and SARS-

CoV-2 in groups 1 and 2 in aged mice lends support to this strategy as a way to elicit broadly

cross-reactive neutralizing antibodies against Group 2B coronaviruses. In agreement with this

notion, chimera 4, which contains the RsSHC014 RBD and SARS-CoV-2 NTD and S2, elicited

binding and neutralizing antibodies and also fully protected mice from BtCoV RsSHC014 and

SARS-CoV-2 challenge, suggesting that CoV spikes vaccines can be designed to maximize their

display of neutralizing and protective epitopes that can cover more than one

pandemic/epidemic/pre-emergent CoV that are at high risk for emergence into naïve human

populations. While other strategies also exist, such as multiplexing mosaic sarbecovirus RBDs

(*19*), S1 or spike glycoproteins and RBDs on nanoparticles (*45*), chimeric spike mRNA-LNP

vaccination can clearly achieve broad protection, using existing manufacturing technologies, and

are highly portable to other high-risk emerging coronaviruses like group 2C MERS-CoV-related

strains. It is notable that our chimeric spike vaccines and the SARS-CoV-2 furin KO, all of

which lacked the two proline stabilizing mutations (S-2P) vaccines protected aged mice from

challenge. This suggests that the two proline stabilization mutations (S2P) are not required for

eliciting protective levels of neutralizing antibodies in aged mice, although such mutations may

enhance the neutralization titers after immunization (*38, 46*).

As previously reported with RNA recombinant viruses, our chimeric spike live viruses

containing SARS-CoV-2 antigenic domains not only demonstrate the known interchangeability

and functional plasticity of CoV spike glycoprotein structural motifs (*29, 47, 48*), but also could

serve as live-attenuated vaccines. Our demonstration of cross-protection against sarbecoviruses

in mice lends support to the notion that universal vaccines against group 2B CoVs is likely

achievable. Moving forward it will be important to determine if these chimeric mRNA-LNP

vaccines can also protect in large pre-clinical animal models, such as rhesus macaques, and if the

S2P mutations will improve the novel chimeric spikes described herein. We conclude that

chimeric mRNA spikes vaccines comprised of zoonotic, epidemic, and pandemic coronaviruses

are a feasible strategy to protect against high-risk, pre-emergent, and pandemic sarbecovirus

infections.

Methods

Chimeric spike vaccine design and formulation

Chimeric spike vaccines were designed with RBD and NTD swaps to increase coverage

of epidemic (SARS-CoV), pandemic (SARS-CoV-2), and high-risk pre-emergent bat CoVs (bat

SARS-like HKU3-1, and bat SARS-like RsSHC014). Chimeric and monovalent spike mRNA-

LNP vaccines were designed based on SARS-CoV-2 spike (S) protein sequence (Wuhan-Hu-1,

GenBank: MN908947.3), SARS-CoV (urbani GenBank: AY278741), bat SARS-like CoV

HKU3-1 (GenBank: DQ022305), and Bat SARS-like RsSHC014 (GenBank: KC881005).

Coding sequences of full-length SARS-CoV-2 furin knockout (RRAR furin cleavage site

abolished between amino acids 682-685), the four chimeric spikes, and the norovirus capsid

negative control were codon-optimized, synthesized and cloned into the mRNA production

plasmid mRNAs were encapsulated with LNP (*49*). Briefly, mRNAs were transcribed to contain

101 nucleotide-long poly(A) tails. mRNAs were modified with m1Ψ-5′-triphosphate (TriLink

#N-1081) instead of UTP and the *in vitro* transcribed mRNAs capped using the trinucleotide

cap1 analog, CleanCap (TriLink #N-7413). mRNA was purified by cellulose (Sigma-Aldrich #

11363-250G) purification. All mRNAs were analyzed by agarose gel electrophoresis and were

stored at -20°C. Cellulose-purified m1Ψ-containing RNAs were encapsulated in proprietary

LNPs using a self-assembly process as previously described wherein an ethanolic lipid mixture

of ionizable cationic lipid, phosphatidylcholine, cholesterol and polyethylene glycol-lipid was

rapidly mixed with an aqueous solution containing mRNA at acidic pH. The RNA-loaded

particles were characterized and subsequently stored at -80°C at a concentration of 1 mg/ml. The

mean hydrodynamic diameter of these mRNA-LNP was ~80 nm with a polydispersity index of

0.02-0.06 and an encapsulation efficiency of ~95%.

Animals, immunizations, and challenge viruses

Eleven-month-old female BALB/c mice were purchased from Envigo (#047) and were

used for all experiments. The study was carried out in accordance with the recommendations for

care and use of animals by the Office of Laboratory Animal Welfare (OLAW), National

Institutes of Health and the Institutional Animal Care and Use Committee (IACUC) of

University of North Carolina (UNC permit no. A-3410-01). mRNA-LNP vaccines were kept

frozen until right before the vaccination. Mice were immunized with 1μg of each vaccine diluted

in sterile 1XPBS in a 50μl volume and were given 25μl intramuscularly in each hind leg. Prime

and boost immunizations were given three weeks apart. Three weeks post boost, mice were bled,

sera was collected for analysis, and mice were moved into the BSL3 facility for challenge

experiments. Animals were housed in groups of five and fed standard chow diets. Virus

inoculations were performed under anesthesia and all efforts were made to minimize animal

suffering. All mice were anesthetized and infected intranasally with 1×10^4 PFU/ml of SARS-

CoV MA15, 1×10^4 PFU/ml of SARS-CoV-2 MA10, or 1×10^4 PFU/ml RsSHC014 which have

been described previously (*9, 34, 50*). Mice were weighted daily and monitored for signs of

clinical disease. Each challenge virus challenge experiment encompassed 50 mice with 10 mice

per vaccine group to obtain statistical power.

Measurement of CoV spike binding by ELISA

A binding ELISA panel that included SARS-CoV spike Protein DeltaTM,

SARS-CoV-2 (2019-nCoV) spike Protein (S1+S2 ECD, His tag), MERS-CoV, Coronavirus

spike S1+S2 (Baculovirus-Insect Cells, His), HKU1 (isolate N5) spike Protein (S1+S2 ECD, His

Tag), OC43 spike Protein (S1+S2 ECD, His Tag), 229E spike Protein (S1+S2 ECD, His tag)

Human coronavirus (HCoV-NL63) spike Protein (S1+S2 ECD, His Tag), Pangolin

CoV_GXP4L_spikeEcto2P_3C8HtS2/293F, bat CoV RsSHC014_spikeEcto2P_3C8HtS2/293F,

RaTG13_spikeEcto2P_3C8HtS2/293F, and bat CoV HKU3-1 spike were tested. Indirect binding

ELISAs were conducted in 384 well ELISA plates (Costar #3700) coated with 2μg/ml antigen in

0.1M sodium bicarbonate overnight at 4°C, washed and blocked with assay diluent (1XPBS

containing 4% (w/v) whey protein/ 15% Normal Goat Serum/ 0.5% Tween-20/ 0.05% Sodium

Azide). Serum samples were incubated for 60 minutes in three-fold serial dilutions beginning at

1:30 followed by washing with PBS/0.1% Tween-20. HRP conjugated goat anti-mouse IgG

secondary antibody (SouthernBiotech 1030-05) was diluted to 1:10,000 in assay diluent without

azide, incubated at for 1 hour at room temperature, washed and detected with 20μl SureBlue

Reserve (KPL 53-00-03) for 15 minutes. Reactions were stopped via the addition of 20μl HCL

stop solution. Plates were read at 450nm. Area under the curve (AUC) measurements were

determined from binding of serial dilutions.

ACE2 blocking ELISAs.

Plates were coated with 2μg/ml recombinant ACE2 protein, then washed and blocked

with 3% BSA in PBS. While assay plates blocked, and sera was diluted 1:25 in 1%BSA/0.05%

Tween-20. Then SARS-CoV-2 spike protein was mixed with equal volumes of each sample at a

final spike concentration equal to the EC_{50} at which it binds to ACE2. The mixture was allowed

to incubate at room temperature for 1 hour. Blocked assay plates were washed, and the serum-

spike mixture was added to the assay plates for a period of 1 hour at room temperature. Plates

were washed and Strep-Tactin HRP, (IBA GmbH, Cat# 2-1502-001) was added at a dilution of

1:5000 followed by TMB substrate. The extent to which antibodies were able to block the

binding of spike protein to ACE2 was determined by comparing the OD of antibody samples at

450nm to the OD of samples containing spike protein only with no antibody. The following

formula was used to calculate percent blocking (100-(OD sample/OD of spike only) *100).

Measurement of neutralizing antibodies against live viruses

Full-length SARS-CoV-2 Seattle, SARS-CoV-2 D614G, SARS-CoV-2 B.1.351, SARS-

CoV, WIV-1, and RsSHC014 viruses were designed to express nanoluciferase (nLuc) and were

recovered via reverse genetics as described previously (*18*). Virus titers were measured in Vero

E6 USAMRIID cells, as defined by plaque forming units (PFU) per ml, in a 6-well plate format

in quadruplicate biological replicates for accuracy. For the 96-well neutralization assay, Vero E6

USAMRID cells were plated at 20,000 cells per well the day prior in clear bottom black walled

plates. Cells were inspected to ensure confluency on the day of assay. Serum samples were tested

at a starting dilution of 1:20 and were serially diluted 3-fold up to nine dilution spots. Serially

diluted serum samples were mixed in equal volume with diluted virus. Antibody-virus and virus

only mixtures were then incubated at 37°C with 5% CO_2 for one hour. Following incubation,

serially diluted sera and virus only controls were added in duplicate to the cells at 75 PFU at

37°C with 5% CO_2. After 24 hours, cells were lysed, and luciferase activity was measured via

Nano-Glo Luciferase Assay System (Promega) according to the manufacturer specifications.

Luminescence was measured by a Spectramax M3 plate reader (Molecular Devices, San Jose,

CA). Virus neutralization titers were defined as the sample dilution at which a 50% reduction in

RLU was observed relative to the average of the virus control wells.

Lung pathology scoring

Acute lung injury was quantified via two separate lung pathology scoring scales: Matute-

Bello and Diffuse Alveolar Damage (DAD) scoring systems. Analyses and scoring were

performed by a board vertified veterinary pathologist who was blinded to the treatment groups as

described previously (*36*). Lung pathology slides were read and scored at 600X total

magnification.

The lung injury scoring system used is from the American Thoracic Society (Matute-

Bello) in order to help quantitate histological features of ALI observed in mouse models to relate

this injury to human settings. In a blinded manner, three random fields of lung tissue were

chosen and scored for the following: (A) neutrophils in the alveolar space (none = 0, 1–5 cells =

1, > 5 cells = 2), (B) neutrophils in the interstitial septa (none = 0, 1–5 cells = 1, > 5 cells = 2),

(C) hyaline membranes (none = 0, one membrane = 1, > 1 membrane = 2), (D) Proteinaceous

debris in air spaces (none = 0, one instance = 1, > 1 instance = 2), (E) alveolar septal thickening

(< 2x mock thickness = 0, 2–4x mock thickness = 1, > 4x mock thickness = 2). To obtain a lung

injury score per field, A–E scores were put into the following formula score = [(20x A) + (14 x

B) + (7 x C) + (7 x D) + (2 x E)]/100. This formula contains multipliers that assign varying

levels of importance for each phenotype of the disease state. The scores for the three fields per

mouse were averaged to obtain a final score ranging from 0 to and including 1.

The second histology scoring scale to quantify acute lung injury was adopted from a lung

pathology scoring system from lung RSV infection in mice (*37*). This lung histology scoring

scale measures diffuse alveolar damage (DAD). Similar to the implementation of the ATS

histology scoring scale, three random fields of lung tissue were scored for the following in a

blinded manner: 1= absence of cellular sloughing and necrosis, 2=Uncommon solitary cell

sloughing and necrosis (1–2 foci/field), 3=multifocal (3+foci) cellular sloughing and necrosis

with uncommon septal wall hyalinization, or 4=multifocal (>75% of field) cellular sloughing

and necrosis with common and/or prominent hyaline membranes. The scores for the three fields

per mouse were averaged to get a final DAD score per mouse. The microscope images were

generated using an Olympus Bx43 light microscope and CellSense Entry v3.1 software.

Measurement of lung cytokines

Lung tissue was homogenized, spun down at 13,000g, and supernantant was used to

measure lung cytokines using Mouse Cytokine 23-plex Assay (BioRad). Briefly, 50µl of lung

homogenate supernatant was added to each well and the protocol was followed according to the

manufacturer specifications. Plates were read using a MAGPIX multiplex reader (Luminex

Corporation).

Biocontainment and biosafety

Studies were approved by the UNC Institutional Biosafety Committee approved by

animal and experimental protocols in the Baric laboratory. All work described here was

performed with approved standard operating procedures for SARS-CoV-2 in a biosafety level 3

(BSL-3) facility conforming to requirements recommended in the Microbiological and

Biomedical Laboratories, by the U.S. Department of Health and Human Service, the U.S. Public

Health Service, and the U.S. Center for Disease Control and Prevention (CDC), and the National

Institutes of Health (NIH).

Statistics

All statistical analyses were performed using GraphPad Prism 9. Statistical tests used in

each figure are denoted in the corresponding figure legend.

REFERENCES AND NOTES

1. J. D. Cherry, P. Krogstad, SARS: The First Pandemic of the 21st Century. *Pediatric Research* **56**, 1-5 (2004).
2. A. M. Zaki, S. van Boheemen, T. M. Bestebroer, A. D. Osterhaus, R. A. Fouchier, Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med* **367**, 1814-1820 (2012).
3. C. I. Paules, H. D. Marston, A. S. Fauci, Coronavirus Infections-More Than Just the Common Cold. *Jama* **323**, 707-708 (2020).
4. P. Zhou *et al.*, A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **579**, 270-273 (2020).
5. The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nat Microbiol* **5**, 536-544 (2020).
6. P. Zhou, Z.-L. Shi, SARS-CoV-2 spillover events. *Science* **371**, 120-122 (2021).
7. P. Zhou *et al.*, Fatal swine acute diarrhoea syndrome caused by an HKU2-related coronavirus of bat origin. *Nature* **556**, 255-258 (2018).
8. C. E. Edwards *et al.*, Swine acute diarrhea syndrome coronavirus replication in primary human cells reveals potential susceptibility to infection. *Proc Natl Acad Sci U S A* **117**, 26915-26925 (2020).
9. V. D. Menachery *et al.*, A SARS-like cluster of circulating bat coronaviruses shows potential for human emergence. *Nat Med* **21**, 1508-1513 (2015).
10. V. D. Menachery *et al.*, SARS-like WIV1-CoV poised for human emergence. *Proc Natl Acad Sci U S A* **113**, 3048-3053 (2016).
11. W. Li *et al.*, Bats Are Natural Reservoirs of SARS-Like Coronaviruses. *Science* **310**, 676-679 (2005).
12. X. Y. Ge *et al.*, Isolation and characterization of a bat SARS-like coronavirus that uses the ACE2 receptor. *Nature* **503**, 535-538 (2013).
13. B. Hu *et al.*, Discovery of a rich gene pool of bat SARS-related coronaviruses provides new insights into the origin of SARS coronavirus. *PLoS Pathog* **13**, e1006698 (2017).
14. W. C. Koff, S. F. Berkley, A universal coronavirus vaccine. *Science* **371**, 759-759 (2021).
15. T. P. Sheahan *et al.*, Broad-spectrum antiviral GS-5734 inhibits both epidemic and zoonotic coronaviruses. *Sci Transl Med* **9**, (2017).
16. T. P. Sheahan *et al.*, An orally bioavailable broad-spectrum antiviral inhibits SARS-CoV-2 in human airway epithelial cell cultures and multiple coronaviruses in mice. *Sci Transl Med* **12**, (2020).
17. C. G. Rappazzo *et al.*, Broad and potent activity against SARS-like viruses by an engineered human monoclonal antibody. *Science* **371**, 823-829 (2021).
18. Y. J. Hou *et al.*, SARS-CoV-2 Reverse Genetics Reveals a Variable Infection Gradient in the Respiratory Tract. *Cell*.
19. A. A. Cohen *et al.*, Mosaic nanoparticles elicit cross-reactive immune responses to zoonotic coronaviruses in mice. *Science*, eabf6840 (2021).
20. Y. Honda-Okubo *et al.*, Severe acute respiratory syndrome-associated coronavirus vaccines formulated with delta inulin adjuvants provide enhanced protection while ameliorating lung eosinophilic immunopathology. *J Virol* **89**, 2995-3007 (2015).

21. M. Bolles *et al.*, A double-inactivated severe acute respiratory syndrome coronavirus
vaccine provides incomplete protection in mice and induces increased eosinophilic
proinflammatory pulmonary response upon challenge. *J Virol* **85**, 12201-12215 (2011).
22. L. Premkumar *et al.*, The receptor binding domain of the viral spike protein is an
immunodominant and highly specific target of antibodies in SARS-CoV-2 patients. *Sci*
Immunol **5**, (2020).
23. L. Liu *et al.*, Potent neutralizing antibodies against multiple epitopes on SARS-CoV-2
spike. *Nature* **584**, 450-456 (2020).
24. L. Dai *et al.*, A Universal Design of Betacoronavirus Vaccines against COVID-19,
MERS, and SARS. *Cell* **182**, 722-733.e711 (2020).
25. D. Li *et al.*, The functions of SARS-CoV-2 neutralizing and infection-enhancing
antibodies in vitro and in mice and nonhuman primates. *bioRxiv*, (2021).
26. X. Chi *et al.*, A neutralizing human antibody binds to the N-terminal domain of the Spike
protein of SARS-CoV-2. *Science* **369**, 650-655 (2020).
27. N. Wang *et al.*, Structural Definition of a Neutralization-Sensitive Epitope on the MERS-
CoV S1-NTD. *Cell Rep* **28**, 3395-3405.e3396 (2019).
28. N. Suryadevara *et al.*, Neutralizing and protective human monoclonal antibodies
recognizing the N-terminal domain of the SARS-CoV-2 spike protein. *bioRxiv*, (2021).
29. M. M. Becker *et al.*, Synthetic recombinant bat SARS-like coronavirus is infectious in
cultured cells and in mice. *Proc Natl Acad Sci U S A* **105**, 19944-19949 (2008).
30. D. Laczkó *et al.*, A Single Immunization with Nucleoside-Modified mRNA Vaccines
Elicits Strong Cellular and Humoral Immune Responses against SARS-CoV-2 in Mice.
Immunity **53**, 724-732.e727 (2020).
31. S. Awasthi *et al.*, Nucleoside-modified mRNA encoding HSV-2 glycoproteins C, D, and
E prevents clinical and subclinical genital herpes. *Sci Immunol* **4**, (2019).
32. K. P. Egan *et al.*, An HSV-2 nucleoside-modified mRNA genital herpes vaccine
containing glycoproteins gC, gD, and gE protects mice against HSV-1 genital lesions and
latent infection. *PLoS Pathog* **16**, e1008795 (2020).
33. P. C. LaTourette, 2nd *et al.*, Protection against herpes simplex virus type 2 infection in a
neonatal murine model using a trivalent nucleoside-modified mRNA in lipid nanoparticle
vaccine. *Vaccine* **38**, 7409-7413 (2020).
34. S. R. Leist *et al.*, A Mouse-Adapted SARS-CoV-2 Induces Acute Lung Injury and
Mortality in Standard Laboratory Mice. *Cell* **183**, 1070-1085.e1012 (2020).
35. G. Matute-Bello *et al.*, An official American Thoracic Society workshop report: features
and measurements of experimental acute lung injury in animals. *Am J Respir Cell Mol*
Biol **44**, 725-738 (2011).
36. T. P. Sheahan *et al.*, Comparative therapeutic efficacy of remdesivir and combination
lopinavir, ritonavir, and interferon beta against MERS-CoV. *Nature Communications* **11**,
222 (2020).
37. M. E. Schmidt *et al.*, Memory CD8 T cells mediate severe immunopathology following
respiratory syncytial virus infection. *PLoS Pathog* **14**, e1006810 (2018).
38. K. S. Corbett *et al.*, SARS-CoV-2 mRNA vaccine design enabled by prototype pathogen
preparedness. *Nature*, (2020).
39. K. S. Corbett *et al.*, Evaluation of the mRNA-1273 Vaccine against SARS-CoV-2 in
Nonhuman Primates. *N Engl J Med*, (2020).

40.   K. Wu *et al.*, Serum Neutralizing Activity Elicited by mRNA-1273 Vaccine —
607   Preliminary Report. *New England Journal of Medicine*, (2021).
608   41.   L. A. Jackson *et al.*, An mRNA Vaccine against SARS-CoV-2 — Preliminary Report.
609   *New England Journal of Medicine* **383**, 1920-1931 (2020).
610   42.   E. E. Walsh *et al.*, Safety and Immunogenicity of Two RNA-Based Covid-19 Vaccine
611   Candidates. *New England Journal of Medicine* **383**, 2439-2450 (2020).
612   43.   L. R. Baden *et al.*, Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N*
613   *Engl J Med*, (2020).
614   44.   E. J. Anderson *et al.*, Safety and Immunogenicity of SARS-CoV-2 mRNA-1273 Vaccine
615   in Older Adults. *N Engl J Med* **383**, 2427-2438 (2020).
616   45.   K. O. Saunders *et al.*, SARS-CoV-2 vaccination induces neutralizing antibodies against
617   pandemic and pre-emergent SARS-related coronaviruses in monkeys. *bioRxiv*, (2021).
618   46.   N. B. Mercado *et al.*, Single-shot Ad26 vaccine protects against SARS-CoV-2 in rhesus
619   macaques. *Nature*, (2020).
620   47.   L. R. Banner, J. G. Keck, M. M. Lai, A clustering of RNA recombination sites adjacent
621   to a hypervariable region of the peplomer gene of murine coronavirus. *Virology* **175**, 548-
622   555 (1990).
623   48.   J. G. Keck, L. H. Soe, S. Makino, S. A. Stohlman, M. M. Lai, RNA recombination of
624   murine coronaviruses: recombination between fusion-positive mouse hepatitis virus A59
625   and fusion-negative mouse hepatitis virus 2. *J Virol* **62**, 1989-1998 (1988).
626   49.   A. W. Freyn *et al.*, A Multi-Targeting, Nucleoside-Modified mRNA Influenza Virus
627   Vaccine Provides Broad Protection in Mice. *Mol Ther* **28**, 1569-1584 (2020).
628   50.   A. Roberts *et al.*, A mouse-adapted SARS-coronavirus causes disease and mortality in
629   BALB/c mice. *PLoS Pathog* **3**, e5 (2007).

ACKNOWLEDGEMENTS

Funding: David R. Martinez is currently supported by a Burroughs Wellcome Fund Postdoctoral

Enrichment Program Award and a Hanna H. Gray Fellowship from the Howard Hugues Medical

Institute and was supported by an NIH NIAID T32 AI007151 and an NIAID F32 AI152296.

This project was supported by the North Carolina Policy Collaboratory at the University of North

Carolina at Chapel Hill with funding from the North Carolina Coronavirus Relief Fund

established and appropriated by the North Carolina General Assembly. This project was funded

in part by the National Institute of Allergy and Infectious Diseases, NIH, U.S. Department of

Health and Human Services award 1U19 AI142759 (Antiviral Drug Discovery and Development

Center awarded to R.S.B), 5R01AI132178 (partnership grant awarded to R.S.B), AI100625, and

AI108197 to R.S.B., AI124429 and a BioNTech SRA to D.W., and E.A.V., as well as an animal

models contract from the NIH (HHSN272201700036I) and U54CA260543 sponsored by NCI.

Animal histopathology services were performed by the Animal Histopathology & Laboratory

Medicine Core at the University of North Carolina, which is supported in part by an NCI Center

Core Support Grant (5P30CA016086-41) to the UNC Lineberger Comprehensive Cancer Center.

Author contributions: Conceived the study: D.R.M. and R.S.B. designed experiments: D.R.M,

R.S.B., performed laboratory experiments: D.R.M, A.S., S.R.L., A.W.; Provided critical

reagents: K.O.S. Analyzed data and provided critical insight: D.R.M, A.S., S.R.L., G.D.l.C.,

A.W., E.A.V., R.P., M.B, D.L., B.Y., K.O.S., D.W., B.F.H., S.M.; Wrote the first draft of the

paper: D.R.M; Read and edited the paper: D.R.M, A.S., S.R.L., G.D.l.C., A.W., E.A.V., R.P.,

M.B, D.L., B.Y., K.O.S., D.W., B.F.H., S.M., R.S.B. Funding acquisition: D.R.M., R.S.B. All

authors reviewed and approved the manuscript. **Competing interests:** The University of North

Carolina at Chapel Hill has filed provisional patents for which D.R.M. and R.S.B are co-

inventors (U.S. Provisional Application No. 63/106,247 filed on October 27[th], 2020) for the

chimeric vaccine constructs and their applications described in this study.

Figures

Figure 1. Genetic relationships among sarbecoviruses and mouse mRNA-LNP vaccination

strategy. (**A**) Genetic diversity of pandemic and bat zoonotic coronaviruses. SARS-CoV is

shown in light blue, RsSHC014 is shown in purple, and SARS-CoV-2 is shown in red. (**B**)

Mouse vaccination strategy using mRNA-LNPs: group 1 received chimeric spike 1, 2, 3, and 4

as the prime and boost, group 2 received chimeric spike 1, 2 as the prime and chimeric spikes 3

and 4 as the boost, group 3 received chimeric spike 4 as the prime and boost, group 4 received

SARS-CoV-2 furin KO prime and boost, and group 5 received a norovirus capsid prime and

boost. Different vaccine groups were separately challenged with SARS-CoV, SARS-CoV-2, and

RsSHC014.

Figure 2. Human pathogenic coronavirus spike binding and hACE2-blocking responses in

chimeric and monovalent SARS-CoV-2 spike-vaccinated mice. ELISA binding responses in

the five different vaccination groups. Both pre-immunization and post-boost binding responses

were evaluated against sarbecovirus, MERS-CoV, and common-cold CoV antigens including:

(**A**) SARS-CoV Toronto Canada (Tor2) S2P, (**B**) SARS-CoV-2 S2P D614G, (**C**) SARS-CoV-2

RBD, (**D**) SARS-CoV-2 NTD, (**E**) Pangolin GXP4L spike, (**F**) RaTG13 spike, (**G**) RsSHC014

S2P spike, (**H**) HKU3-1 spike, (**I**) MERS-CoV spike, (**J**) hACE2 blocking responses against

SARS-CoV-2 spike in the distinct immunization groups. Blue squares represent mice from group

1, orange triangles represent mice from group 2, green triangles represent mice from group 3, red

rhombuses represent mice from group 4, and upside-down triangle represent mice from group 5.

Statistical significance for the binding and blocking responses is reported from a Kruskal-Wallis

test after Dunnett's multiple comparison correction. $^*p < 0.05$, $^{**}p < 0.01$, $^{***}p < 0.001$, and

$^{****}p < 0.0001$.

Figure 3. Live sarbecovirus neutralizing antibody responses. Neutralizing antibody responses

in mice from the five different vaccination groups against nanoluciferase-expressing infectious

molecular clones. (**A**) SARS-CoV neutralizing antibody responses from baseline and post boost

in the distinct vaccine groups. (**B**) SARS-CoV-2 neutralizing antibody responses from baseline

and post boost. (**C**) RsSHC014 neutralizing antibody responses from baseline and post boost. (**D**)

WIV-1 neutralizing antibody responses from baseline and post boost. (**E**) SARS-CoV-2 D614G

and South African B.1.351 variant of concern neutralizing activity in groups 1 and 4. (**F**)

Neutralization comparison of SARS-CoV-2 D614G vs. South African B.1.351. Statistical

significance for the live-virus neutralizing antibody responses is reported from a Kruskal-Wallis

test after Dunnett's multiple comparison correction. $^*p < 0.05$, $^{**}p < 0.01$, $^{***}p < 0.001$, and

$^{****}p < 0.0001$.

Figure 4. *In vivo* protection against sarbecovirus challenge by chimeric spikes mRNA-

vaccines. (**A**) Percent starting weight from the different vaccine groups of mice challenged with

SARS-CoV MA15. (**B**) SARS-CoV MA15 lung viral titers in mice from the distinct vaccine

groups. (**C**) SARS-CoV MA15 nasal turbinate titers. (**D**) Percent starting weight from the

different vaccine groups of mice challenged with SARS-CoV-2 MA10. (**E**) SARS-CoV-2 MA10

lung viral titers in mice from the distinct vaccine groups. (**F**) SARS-CoV-2 MA10 nasal

turbinate titers. Figure legend at the bottom right depicts the vaccines utilized in the different

groups. Statistical significance for weight loss is reported from a two-way ANOVA after

Dunnett's multiple comparison correction. For lung and nasal turbinate titers, statistical

significance is reported from a one-way ANOVA after Tukey's multiple comparison correction.

$^*p < 0.05$, $^{**}p < 0.01$, $^{***}p < 0.001$, and $^{****}p < 0.0001$.

Figure 5. Lung pathology in protected vs. infected mice challenged with SARS-CoV and

SARS-CoV-2. (**A**) Hematoxylin and eosin 4 days post infection lung analysis of SARS-CoV

MA15 challenged mice from the different groups: group 1: chimeras 1-4 prime and boost, group

2: chimeras 1-2 prime and 3-4, group 3: chimera 4 prime and boost, SARS-CoV-2 furin KO

prime and boost, and norovirus capsid prime and boost. (**B**) Lung pathology quantitation in

SARS-CoV MA15 challenged mice from the different groups. Macroscopic lung discoloration

score, microscopic acute lung injury (ALI) score, and diffuse alveolar damage (DAD) in day 4

post infection lung tissues are shown. (**C**) Hematoxylin and eosin 4 days post infection lung

analysis of SARS-CoV-2 MA10 challenged mice from the different groups: group 1: chimeras 1-

4 prime and boost, group 2: chimeras 1-2 prime and 3-4, group 3: chimera 4 prime and boost,

SARS-CoV-2 furin KO prime and boost, and norovirus capsid prime and boost. (**D**) Lung

pathology measurements in SARS-CoV-2 MA10 challenged mice from the different groups.

Macroscopic lung discoloration score, microscopic acute lung injury (ALI) score, and diffuse

alveolar damage (DAD) in day 4 post infection lung tissues are shown. Statistical significance is

reported from a one-way ANOVA after Dunnet's multiple comparison correction. $^*p <$

0.05, $^{**}p < 0.01$, $^{***}p < 0.001$, and $^{****}p < 0.0001$.

 Supplemental figures

 Figure S1. Chimeric spike constructs from sarbecoviruses. (**A**) Spike chimera 1 includes the

 NTD from HKU3-1, the RBD from SARS-CoV, and the rest of the spike from SARS-CoV-2.

 (**B**) Spike chimera 2 includes the RBD from SARS-CoV-2 and the NTD and S2 from SARS-

 CoV. (**C**) Spike chimera 3 includes the RBD from SARS-CoV and the NTD and S2 SARS-CoV-

 2. (**D**) Spike chimera 4 includes the RBD from RsSHC014 and the rest of the spike from SARS-

 CoV-2. (**E**) SARS-CoV-2 furin KO spike vaccine and (**F**) is the norovirus capsid vaccine. (**G**).

 Protein expression of chimeric spikes, SARS-CoV-2 furin KO, and norovirus mRNA vaccines.

 GAPDH was used as the loading control. (**H**) Nanoluciferase expression of RsSHC014/SARS-

 CoV-2 chimeric spike live viruses.

 Figure S2. Human common-cold CoV ELISA binding responses in chimeric and

 monovalent SARS-CoV-2 spike mRNA-LNP-vaccinated mice. Pre-immunization and post

 boost binding to (**A**) HCoV-HKU1 spike, (**B**) HCoV-OC43 spike, (**C**) HCoV-229E spike, and

 (**D**) HCoV-NL63 spike. Statistical significance for the binding and blocking responses is

 reported from a Kruskal-Wallis test after Dunnett's multiple comparison correction. $^*p <$

 0.05, $^{**}p < 0.01$, $^{***}p < 0.001$, and $^{****}p < 0.0001$.

 Figure S3. Comparison of neutralizing antibody activity of CoV mRNA-LNP vaccines

 against sarbecoviruses. (**A**) Group 1 neutralizing antibody responses against SARS-CoV-2,

 SARS-CoV, RsSHC014, and WIV-1 and (**B**) fold-change of SARS-CoV, RsSHC014, and WIV-

 1 neutralizing antibodies relative to SARS-CoV-2. (**C**) Group 2 neutralizing antibody responses

 against SARS-CoV-2, SARS-CoV, RsSHC014, and WIV-1 and (**D**) fold-change of SARS-CoV,

RsSHC014, and WIV-1 neutralizing antibodies relative to SARS-CoV-2. (**E**) Group 3

neutralizing antibody responses against SARS-CoV-2, SARS-CoV, RsSHC014, and WIV-1 and

(**F**) fold-change of SARS-CoV, RsSHC014, and WIV-1 neutralizing antibodies relative to

SARS-CoV-2. (**G**) Group 4 neutralizing antibody responses against SARS-CoV-2, SARS-CoV,

RsSHC014, and WIV-1 and (**H**) fold-change of SARS-CoV, RsSHC014, and WIV-1

neutralizing antibodies relative to SARS-CoV-2.

Figure S4. *In vivo* protection against BtCoV challenge by chimeric spikes mRNA-vaccines.

(**A**) Percent starting weight from the different vaccine groups of mice challenged with

RsSHC014. (**B**) RsSHC014 lung viral titers in mice from the distinct vaccine groups. (**C**)

RsSHC014 nasal turbinate titers in mice from the different immunization groups. Statistical

significance is reported from a one-way ANOVA after Tukey's multiple comparison correction.

$^*p < 0.05$, $^{**}p < 0.01$, $^{***}p < 0.001$, and $^{****}p < 0.0001$.

Figure S5. Survival analysis of immunized mice challenged with sarbecoviruses. (**A**)

Survival analysis from immunized mice infected with SARS-CoV MA15 and (**B**) SARS-CoV-2

MA10. Statistical significance is reported from a Mantel-Cox test.

Figure S6. Lung cytokine analysis in protected vs. sarbecovirus-infected vaccinated mice.

CCL2, IL-1α, G-SCF, and CCL4 in (**A**) SARS-CoV-infected mice and in (**B**) SARS-CoV-2-

infected mice. Statistical significance for the binding and blocking responses is reported from a

Kruskal-Wallis test after Dunnett's multiple comparison correction. $^*p < 0.05$, $^{**}p < 0.01$, $^{***}p <$

0.001, and $^{****}p < 0.0001$.

Figure 1

A

BtCoV/273/2005
Rf1
BtRs-BetaCoV/YN2018A
HKU3-1
BtCoV/279/2005
Rm1
SARS-CoV-2
bat-SL-CoVZC45
bat-SL-CoVZXC21
Rs4247
Rs4237
BtRs-YN2018C
Rs4255
As6526
Rs4081
BtRs-BetaCoV/YN2018D
Rf4092
Rp3
BtRl-BetaCoV/SC2018
RsSHC014
Rs4231
WIV-1
BtRs-BetaCoV/YN2018B
Rs7327
Rs9401
Rs4874
HC/SZ/61/03
CUHK-W1
SARS-CoV Urbani
GD03T0013
0.120

B

Group 1
Chimeric spikes 1, 2, 3, 4
day 0
Chimeric spikes 1, 2, 3, 4
day 21
day 55 challenge
1. SARS-CoV
2. SARS-CoV-2
3. RsSHC014
day -1
pre-prime
bleed
day 49
post boost
bleed

Group 2
Chimeric spikes 1, 2
day 0
Chimeric spikes 3, 4
day 21
day 55 challenge
1. SARS-CoV
2. SARS-CoV-2
3. RsSHC014
day -1
pre-prime
bleed
day 49
post boost
bleed

Group 3
Chimeric spike 4
day 0
Chimeric spike 4
day 21
day 55 challenge
1. SARS-CoV
2. SARS-CoV-2
3. RsSHC014
day -1
pre-prime
bleed
day 49
post boost
bleed

Group 4
SARS-CoV-2 wild type
day 0
SARS-CoV-2 wild type
day 21
day 55 challenge
1. SARS-CoV
2. SARS-CoV-2
3. RsSHC014
day -1
pre-prime
bleed
day 49
post boost
bleed

Group 5
Norovirus capsid
day 0
Norovirus capsid
day 21
day 55 challenge
1. SARS-CoV
2. SARS-CoV-2
3. RsSHC014
day -1
pre-prime
bleed
day 49
post boost
bleed

Figure 2

Figure 3

Figure 4

mRNA vaccine group
- Group 1: chimeras 1-4 prime/boost
- Group 2: chimeras 1-2 prime and 3-4 boost
- Group 3: chimera 4 prime/boost
- Group 4: SARS-CoV-2 spike furin KO prime/boost
- Group 5: Norovirus capsid prime/boost

Figure 5

Figure S1

A Spike chimera 1
HKU3-1 NTD/SARS-CoV RBD/SARS-CoV2 S2

B Spike chimera 2
SARS-CoV-2 RBD/SARS-CoV S1 and S2

C Spike chimera 3
SARS-CoV RBD/SARS-CoV-2 S1 and S2

D Spike chimera 4
RsSHC014 RBD/SARS-CoV-2 S1 and S2

E SARS-CoV-2 wild type furin knockout

F Norovirus capsid

G

spike

GAPDH

Chimera 1 | Chimera 2 | Chimera 3 | Chimera 4 | SARS-CoV-2 spike furin KO | Norovirus capsid

H Luciferase

Log_{10} RLU

10^7
10^6
10^5
10^4
10^3
10^2
10^1
10^0

10^0 10^5 10^{10} 10^{15}

dilution

○ RsSHC014nLuc wild type
● RsSHC014/SARS-CoV-2 RBD nLuc
● RsSHC014/SARS-CoV-2 NTD nLuc
● RsSHC014/SARS-CoV-2 NTD + RBD nLuc
● RsSHC014/SARS-CoV-2 RBD+ nLuc
● RsSHC014/SARS-CoV-2 S2 nLuc

Figure S2

Figure S3

Figure S4

A RsSHC014 challenge

B Virus Lung Titer
4dpi RsSHC014

C Nasal Turbinate Titer
4dpi RsSHC014

<u>mRNA vaccine group</u>

- Group 1: chimeras 1-4 prime/boost
- Group 2: chimeras 1-2 prime and 3-4 boost
- Group 3: chimera 4 prime/boost
- Group 4: SARS-CoV-2 spike furin KO prime/boost
- Group 5: Norovirus capsid prime/boost

Figure S5

A   Survival SARS-CoV challenge

B   Survival SARS-CoV-2 challenge

mRNA vaccine group

- Group 1: chimeras 1-4 prime/boost
- Group 2: chimeras 1-2 prime and 3-4 boost
- Group 3: chimera 4 prime/boost
- Group 4: SARS-CoV-2 spike furin KO prime/boost
- Group 5: Norovirus capsid prime/boost

Figure S6