1

PBN-PVT projection modulates negative affective states in mice

Ya-Bing Zhu[1,4], Yan Wang[1,4], Xiao-Xiao Hua[2,4], Ling Xu[1], Ming-Zhe Liu[3], Rui Zhang[1], Peng-Fei Liu[1], Jin-Bao Li[1], Ling Zhang[2,*], Di Mu[1,5,*]

[1]Department of Anesthesiology, Shanghai General Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, China, 201620, China

[2]The First Rehabilitation Hospital of Shanghai, Tongji University School of Medicine, Shanghai 200090, China

[3]Department of Respiratory, The First Affiliated Hospital of Guangzhou Medical University, Guangzhou 510120, China

[4]These authors contributed equally

[5]Lead Contact

*Correspondence to:

Ling Zhang, Ph.D., The First Rehabilitation Hospital of Shanghai, Tongji University School of Medicine, Shanghai 200090, China; Email: lzhang0808@tongji.edu.cn.

Di Mu, Ph.D., Department of Anesthesiology, Shanghai General Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai 201620, China; Email: damonmu@163.com or dimu08207@ustc.edu

Abstract

Long-lasting negative affections dampen enthusiasm for life, and dealing with negative affective states is essential for individual survival. The parabrachial nucleus (PBN) and the thalamic paraventricular nucleus (PVT) are critical for modulating affective states in mice. However, the functional role of the PBN-PVT projection in modulating affective states remains elusive. Here, we show that the PBN neurons send dense projection fibers to the PVT and form direct excitatory synapses with the PVT neurons. Activation of the PBN-PVT projection or PVT-projecting PBN neurons induces robust anxiety-like, aversion-like, and fear-like behaviors without affecting nociceptive behaviors. Inhibition of the PBN-PVT projection or PVT-projecting PBN neurons reduces fear-like and aversion-like behaviors. Furthermore, the PVT neurons innervated by the PBN are activated by aversive stimulation, and activation of PBN-PVT projection enhances the neuronal activity of PVT neurons in response to the aversive stimulus. Activation of these downstream PVT neurons induces anxiety-like behavior behaviors. Thus, our study indicates that the PBN-PVT projection modulates negative affective states in mice.

Keywords: Parabrachial nucleus (PBN); thalamic paraventricular nucleus (PVT); affective states; anxiety; aversion.

Introduction

Threat and injury often induce defensive reactions, such as flight, freezing, hiding *(Öhman and Mineka, 2001)*, and negative affective states, such as fear and anxiety *(Jimenez et al., 2018)*. Such behavioral adaptations and psychological responses are essential for survival, and understanding the mechanisms is of fundamental interest. It is worth noting that the parabrachial nucleus (PBN) in the brainstem plays a critical role in encoding danger signals and promoting affective behavior states to limit harm in response to potential threats *(Campos et al., 2018)*.

The PBN receives the majority of the ascending inputs from the spinal cord

(Todd, 2010), and the PBN neurons respond robustly to nociception, food

neophobia, hypercapnia, and threat for maintaining homeostasis under

stressful circumstances *(Campos et al., 2018; Kaur et al., 2013)*. The PBN

relays this information (visceral malaise, taste, temperature, pain, itch) to brain

areas, such as the hypothalamus, the central of the amygdala (CeA), thalamus,

insular cortex (IC), and periaqueductal gray (PAG), to participate in diverse

physiology process *(Chiang et al., 2019; Palmiter, 2018; Saper, 2016)*. A

recent study has found that the subpopulations of PBN have distinct projection

patterns and functions *(Chiang et al., 2020)*. The dorsal division PBN neurons

projecting to the ventromedial hypothalamus (VMH) and PAG mediate

escaping behaviors. In contrast, the external lateral division PBN neurons

projecting to the bed nucleus of the stria terminalis (BNST) and the CeA

mediate aversion and avoidance memory *(Chiang et al., 2020)*. Optogenetic

manipulation of specific outputs from PBN generates a specific function

(Bowen et al., 2020). In the thalamus, the intralaminar thalamus nucleus (ILN)

is the downstream target of PBN neurons that receive spinal cord inputs, and

the ILN pathway participates in nociception processing *(Deng et al., 2020)*.

Besides the ILN nucleus, the thalamic paraventricular nucleus (PVT) is

another primary target of the PBN nucleus in the thalamus *(Chiang et al.,*

2020).

The PVT nucleus locates in the dorsal part of the midline thalamus *(Vertes*

et al., 2015), and has been heavily implicated in a range of affective behaviors

(Hsu et al., 2014). The functional roles of the PVT include diverse processes

such as arousal (*Ren et al.,* 2018), drug addiction (*Zhu et al.,* 2016),

reward-seeking (*Do-Monte et al.,* 2017; *Engelke et al.,* 2021), stress (*Beas et*

al., 2018; *Gao et al., 2020*), associative learning and memory retrieval (*Penzo*

et al., 2015; *Do-Monte et al.,* 2015; *Zhu et al.,* 2018; *Keyes et al.,* 2020). The

PVT receives a significant amount of inputs from the brainstem (locus

coeruleus, PBN, PAG), hypothalamus, prefrontal cortical areas and projects to

the infralimbic cortex, nucleus accumbens (NAc), BNST, and CeA *(Kirouac,*

2015). The convergent signals included the arousal from the hypothalamus

(Ren et al., 2018), the emotional saliency from the prefrontal cortex

(Yamamuro et al., 2020), and the stress responsivity from locus coeruleus (LC)

(Beas et al., 2018) might help to promote the appropriate behavioral

responses to environmental challenges. However, despite the substantial

improvements in our understanding of the PVT nucleus's neurocircuitry, the

functional role of the PBN-PVT projection remains mostly unknown.

In this study, we used virus tracing and electrophysiology to dissect the

anatomical and functional connection between the PBN nucleus and the PVT

nucleus. By using optogenetic and pharmacogenetic approaches, we then

demonstrated that the PBN-PVT projection modulates negative affective

states in mice.

Results

Functional connectivity pattern of the PBN-PVT projection

Previous studies have reported that the PVT could receive input from the PBN

nucleus *(Chiang et al., 2020; Li and Kirouac, 2012)*. The detailed morphology

of the PBN-PVT projection and whether these two nuclei form direct functional

synapses remain unknown. We first injected the AAV2/8-hSyn-ChR2-mCherry

virus into the PBN and found that there were dense projection fibers in the PVT

(*Figure 1A–C*). We employed the whole-cell patch-clamp recording to examine

the synaptic connectivity between the PBN and the PVT. Precisely time-locked

action potentials were induced by a train of brief laser pulses (5 Hz, 10 Hz, and

20 Hz, *Figure 1G*). We found that optogenetic activation of the PBN projection

fibers evoked excitatory postsynaptic currents (EPSCs) in 34 of 52 PVT

neurons. The medial PVT showed higher connectivity (bregma -0.94 to -1.82

mm, 27 of 37 cells, 72.97%) than anterior PVT (bregma -0.22 to -0.94 mm, 2 of

6 cells, 33.33%) or posterior PVT (bregma -1.82 to -2.3 mm, 5 of 9 cells,

55.56%, *Figure 1D–F*). The average amplitude of the light-evoked EPSCs was

103.4 ± 11.93 pA (*Figure 1H*). Moreover, the latency of EPSCs was 3.13 ± 0.29 ms with a jitter of 0.16 ± 0.02 ms (*Figure 1I-J*), indicating monosynaptic connections between the PBN and the PVT nuclei. The EPSCs were sensitive to the Na^+ channel blocker tetrodotoxin (TTX, 1 μM) and were rescued by the K^+ channel blocker 4-aminopyridine (4-AP, 100 μM). The EPSCs were further blocked by the AMPA receptor antagonist NBQX (10 μM), confirming the monosynaptic glutamatergic innervation of the PVT neurons by the PBN neurons (*Figure 1K–L*). In addition, we also observed there were light-evoked inhibitory postsynaptic currents (IPSCs) in only 2 of 52 PVT neurons (less than 30 pA).

Next, we injected the retroAAV2/2-hSyn-Cre virus into the PVT nucleus on *Rosa26-tdTomato* mice which could retrogradely label projection neurons in the PBN (*Figure 1M–P*). We found that $tdTomato^+$ neurons were bilaterally located in the lateral PBN (55 ± 5.82 neurons, *n* = 4 mice) and rarely in the medial PBN (7.75 ± 1.03 neurons, *Figure 1O–Q*). These results indicate that bilateral PBN project to the PVT. We then performed tdTomato staining with *type 2 vesicular glutamate transporter* (*VgluT2*) mRNA in situ hybridization and found that about 94.4% of tdTomato neurons express *VgluT2 mRNA* (*Figure 1R-U*). These results indicate that the majority of PVT-projecting PBN neurons are glutamatergic. We also examined several markers for subpopulations of PBN neurons, including tachykinin 1 receptor (Tacr1), tachykinin 1 (Tac1), prodynorphin (Pdyn), calcitonin gene-related peptide (CGRP). And we found that tdTomato neurons were only partially co-labeled with *Tacr1, Tac1,* or *Pdyn* mRNA, but not with CGRP (*Figure 1–figure supplement 1*).

Then, we used the *VgluT2-ires-Cre* mice combined with the AAV2/8-EF1a-DIO-EGFP virus to specifically label the glutamatergic neurons of the PBN nucleus (*Figure 1–figure supplement 2A*). Robust expression of AAV2/8-EF1a-DIO-EGFP was found in both the lateral and medial PBN nuclei (*Figure1–figure supplement 2B–D*). It is worth noting that the density of $EGFP^+$ fibers was higher in the middle and posterior PVT (*Figure1–figure supplement*

2E–H), considering that the notion of the posterior PVT being a particularly

aversive region of the PVT (*Gao et al.,* 2020). We also examined the collateral

projections from PVT-projecting PBN neurons (*Figure 1–figure supplement 3*).

The collateral projections were also found in the BNST, lateral hypothalamus

(LH), paraventricular nucleus of the hypothalamus (PVN), PAG, but not CeA or

VMH.

Optogenetic activation of PBN-PVT projection induces anxiety-like

behaviors and aversion-like behaviors

We   injected   the   AAV2/9-EF1a-DIO-ChR2-mCherry   virus   or

AAV2/9-EF1a-DIO-mCherry virus into the bilateral PBN of *VgluT2-ires-Cre*

mice and implanted optic fibers above the PVT to activate the PBN-PVT

projection selectively (*Figure 2A*). Four weeks after surgery, we found robust

expression of ChR2-mCherry (*Figure 2B–C, Figure 2–figure supplement 1A*)

or mCherry (*Figure 2–figure supplement 1C*) in bilateral PBN neurons and the

axon terminals in the PVT (*Figure 2D, Figure 2–figure supplement 1B, Figure*

2–figure supplement 1D).

We performed a 15 minutes optogenetic manipulation open field test (OFT,

0–5 minutes laser OFF, 5–10 minutes laser ON, 10–15 minutes laser OFF,

Figure 2E). Optogenetic activation (473 nm, 20 Hz, 5 mW, 5ms) of the

efferents from the PBN to the PVT elicited instant running behavior along the

chamber wall with a significantly increased velocity in ChR2 group mice

(*Figure 2F–H, Figure 2–video 1*). The ChR2 injected mice rarely entered into

the center area of the chamber, represented as a decrease in center time than

that of the control group (*Figure 2I*). It is worth noting that the velocity returned

to normal once the laser was off, but the time spent in the center was still lower

than the control group in the 5 minutes after stimulation. These results

indicated that the anxiety could last for at least several minutes after acute

activation. Although the speed increased during the laser ON period, the

unmoving time of the ChR2 mice during the laser ON period was also

increased (*Figure 2–figure supplement 2A*). Therefore, the distance during the

laser ON period and the total distance in 15 minutes were not changed (*Figure*

2–figure supplement 2B).

To dissect a more detailed profile of the behaviors in the OFT, we further

divided the laser ON period (5–10 minutes) into 5 one-minute periods and

analyzed the velocity, unmoving time, center time, distance, and jumping

(*Figure 2–figure supplement 2C–G*). We found that the velocity and unmoving

time were increased, and the center time was decreased in the ChR2 mice

during most periods (*Figure 2–figure supplement 2C–E*). Furthermore, we

observed that the distance and jumping behaviors were increased mainly in

the first one-minute period in ChR2 mice (*Figure 2–figure supplement 2F–G*).

This detailed analysis indicates that optogenetic activation induces brief and

robust running, jumping behaviors, and persistent anxiety-like behaviors, such

as unmoving and less time spent in the center.

Besides anxiety, another critical component of negative affective states is

aversion. Therefore, we used the real-time placed aversion test (RTPA) to

explore the function of optogenetic activation of the PBN-PVT projection

(*Figure 2J*). We found that the ChR2 mice spent less time in the laser-paired

chamber, and the aversion disappeared when the laser was off (*Figure 2K–M,*

Figure 2–video 2). We also used a prolonged conditioning protocol that mimics

drug-induced conditioned place aversion (*Figure 2–figure supplement 2H*). We

found that the ChR2 mice did not display aversion in the post-conditioning test

(*Figure 2-figure supplement 2I*). These results indicate that activation of the

PBN-PVT projection is sufficient to induce aversion but could not enable

associative aversive memory formation.

To further confirm this instant aversion phenomenon, we subjected mice to

the cue-dependent optogenetic conditioning test (*Figure 2N*). A 30 seconds

auditory conditioning stimulus (CS) co-terminated with 30 seconds of

synchronous optogenetic activation of the PBN-PVT projection (laser stimulus,

LS) in this test. The ChR2 expressing mice generated significant freezing

behavior during 6 CS-LS pairings (*Figure 2O*). However, the freezing behavior

to the same context or to the auditory cue in a novel context was disappeared

on the second day (*Figure 2P–Q*). These results demonstrate that optogenetic

activation of the PBN-PVT projection induces instant aversion and freezing but

does not drive associative fear memory formation.

Pharmacogenetic activation of the PVT-projecting PBN neurons induces

anxiety-like behaviors and freezing behaviors

We also used pharmacogenetic manipulation and retrograde tracing to confirm

the effects of activating the PVT-projecting PBN neurons. We first injected the

retroAAV2/2-hSyn-Cre virus into the PVT, and the

AAV2/9-hSyn-DIO-hM3Dq-mCherry virus or control virus into bilateral PBN to

specifically transduce the PVT-projecting PBN neurons with a designer

receptor exclusively activated by designer drugs (DREADDs, *Figure 3A*)

(Armbruster et al., 2007). The PVT-projecting PBN neurons could be activated

by intraperitoneal injection of clozapine N-oxide dihydrochloride (CNO, *Figure*

3B–D). The region of the virus expression in the PBN nucleus is shown in

Figure 3–figure supplement 1. Consistent with the optogenetic activation

results, the hM3Dq mice spent less time in the center, had more unmoving

time, and traveled fewer distances than the mCherry mice in the OFT (*Figure*

3E–I). At the same time, the velocities were not significantly different (*Figure*

3J). We also found that activation of the PVT-projecting PBN neurons did not

affect motor ability in the rotarod test (*Figure 3–figure supplement 2G*).

Besides, the hM3Dq mice showed decreased exploration time of open

quadrants in the EZM (*Figure 3K–L*), further suggesting that activation of

PVT-projecting PBN neurons could induce anxiety-like behaviors.

We further evaluated freezing behaviors in the fear conditioning chamber

and found that the hM3Dq mice displayed more freezing behaviors after

injection of CNO than control mice (*Figure 3M–N*). Although activation of

PVT-projecting PBN neurons induced significant anxiety-like behavior, it did

not affect the depressive-like behaviors evaluated by the tail suspension test (TST, *Figure 3–figure supplement 2A*) and the forced swimming test (FST, *Figure 3–figure supplement 2B*). Previous studies have revealed that the PBN receives direct projections from the spinal cord and plays a vital role in pain processing *(Deng et al., 2020; Sun et al., 2020)*. We then assessed whether pharmacogenetic activation of the PVT-projecting PBN neurons affects the nociceptive behaviors. By performing the von Frey test and Hargreaves test, we found that the basal nociceptive thresholds were not affected after pharmacogenetic activation of PVT-projecting PBN neurons (*Figure 3–figure supplement 2C–D*). Given that the distinct mechanisms between the reflexive and coping responses induced by nociceptive stimulation *(Huang et al., 2019)*, we injected formalin into the paw to induce inflammatory pain. We found that activation of the PVT-projecting PBN neurons did not affect the formalin-evoked licking behaviors (*Figure 3–figure supplement 2E–F*). These results indicate that the PBN-PVT projection might not participate in the pain signal processing.

Inhibition of the PBN-PVT projection reduces the 2-MT-induced aversive behaviors and footshock-induced fear behaviors

The activation manipulation results prompted us to investigate whether inhibition of the PBN-PVT projection could modulate the negative affective states. We first injected the AAV2/9-EF1a-DIO-NpHR3.0-EYFP virus or the AAV2/8-EF1a-DIO-EGFP virus into the PBN and implanted the optic fibers into the PVT of *VgluT2-ires-Cre* mice (*Figure 4A–C, Figure 4–figure supplement 1*). We used 2-methyl-2-thiazoline (2-MT), a widely-used odorant molecule that could generate innate fear-like freezing responses in rodents *(Isosaka et al., 2015)*, to induce the fear-like state. We found that 589 nm laser-induced inhibition of the PBN-PVT projection reduced the aversion caused by the 2-MT (*Figure 4D–F*) and increased the moving duration (*Figure 4G*). We also observed that inhibition of the PBN-PVT projection increased the time in the

2-MT zone in the OFT (*Figure 4H–I*). Besides the 2-MT, footshock is another

paradigm that induces robust freezing behaviors. We found that constant

inhibition of the PBN-PVT projection reduced the footshock-induced freezing

behaviors (*Figure 4J–K*).

We also examined whether inhibition of the PBN-PVT projection affects

aversive memory acquisition or retrieval (*Figure 4–figure supplement 2A*). We

briefly suppressed the activity of the PBN-PVT projection during footshock

stimulation and found that freezing levels were not changed (*Figure 4–figure*

supplement 2B). We further compared the freezing levels in contextual and

cue tests without or with laser and found that aversive memory retrieval was

not affected either (*Figure 4–figure supplement 2C–D*). In addition, we

performed optogenetic inhibition of the PVT-projecting PBN neurons and

observed similar phenomena (*Figure 4–figure supplement 3*).

PBN input shapes PVT neuronal responses to aversive stimulation

By using *in vivo* fiber photometry, we found that the calcium signals of the PVT

neurons were increased after aversive stimuli, such as footshock and air puff

(*Figure 5–figure supplement 1*), indicating that the calcium signal of the PVT

neurons is increased in response to aversive stimuli. Besides, we injected the

AAV2/1-hSyn-Cre virus, which could anterogradely label the downstream

neurons *(Zingg et al., 2017)*, into the PBN of *Rosa26-tdTomato* mice (*Figure*

5A). The distribution pattern of tdTomato$^+$ neurons in PVT (hereafter referred to

as PVT$_{PBN}$ neurons) was shown in *Figure 5B–D*. We used Fos as a marker to

assess the activity change in 2-MT treated mice and footshock treated mice.

The percentage of Fos$^+$tdTomato$^+$ neurons/tdTomato$^+$ neurons in the PVT was

significantly increased in the aversive stimuli-treated mice than that of control

mice (*Figure 5E–H*), confirming that the PVT$_{PBN}$ neurons could be activated by

aversive stimuli.

The next question is whether the PBN-PVT projection modulates the

neuronal activity of the PVT neurons in response to aversive stimuli. We first

injected the AAV2/9-EF1a-DIO-ChR2-mCherry virus into the PBN and

performed a dual Fos staining (*Seike et al.,* 2020), detecting *fos* mRNA and

Fos protein induced by two episodes of stimulation (*Figure 6–figure*

supplement 1A). We found that there was a broad overlap between

optogenetic stimulation-activated neurons (expressing the Fos protein) and

footshock-activated neurons (expressing the *fos* mRNA) (*Figure 6–figure*

supplement 1B–E). Then we injected the AAV2/9-EF1a-DIO-ChR2-mCherry

virus into the PBN and implanted the optoelectrode into the PVT of

VgluT2-ires-Cre mice (*Figure 6A*). We first recorded the spiking signals in

response to 10 sweeps of 2 s laser pulse trains (20 Hz, 5 mW, 5 ms). Then we

recorded the spiking signals in response to 20 sweeps of 2 s footshock (0.5

mA) without laser in odd number sweeps or with laser in even number sweeps

(*Figure 6A*). We found that laser or footshock (without laser) increased firing

rates in 22 or 28 of 40 units (*Figure 6B–C*). And there was also a broad overlap

between laser-activated and footshock-activated units (*Figure 6D*). It was

consistent with the dual Fos staining result, suggesting that PVT_{PBN} neurons

are activated by aversive stimulation. Next, we analyzed the firing rates of PVT

neurons in footshock with laser sweeps and footshock without laser sweeps

(*Figure 6E-G*). We found that the footshock stimulus with laser activated 30 of

40 units (*Figure 6H*) and increased the overall firing rates of neurons

compared with the footshock without laser result (*Figure 6I*). These results

indicate that activation of the PBN-PVT projection could enhance the PVT

neuronal responses to aversive stimulation.

Pharmacogenetic activation of the PVT_{PBN} neurons induces anxiety-like

behaviors

To further investigate the specific role of PVT_{PBN} neurons in modulating

negative affective states, we injected the AAV2/1-hSyn-Cre virus into the

bilateral PBN and injected the AAV2/9-hSyn-DIO-hM3Dq-mCherry virus or

control virus into the PVT to activate the PVT_{PBN} neurons *(Figure 7A)*. The

majority of the PVT$_{PBN}$ neurons could be activated by CNO in the hM3Dq

expressing mice but not the control mice (*Figure 7B–D*). We found that

pharmacogenetic activation of the PVT$_{PBN}$ neurons reduced the center time

(*Figure 7E–F*). Similarly, the time spent in open quadrants was decreased in

the EZM of hM3Dq expressing mice (*Figure 7G–H*). And the unmoving time in

the EZM test had an increased tendency in hM3Dq expressing mice (*Figure 7I*).

We did not observe obvious nociception-related behaviors, such as forelimb

wiping, hindlimb flinching, licking, or bitting during the experiments. These

results indicate that pharmacogenetic activation of PVT$_{PBN}$ neurons induces

anxiety-like behaviors.

Furthermore, we examined the anatomic distribution of terminals of

PVT$_{PBN}$ neurons. We labeled the PVT$_{PBN}$ neurons in WT mice by injecting the

AAV2/1-hSyn-Cre virus into the PBN and AAV2/8-EF1a-DIO-EGFP virus into

the PVT (*Figure 7–figure supplement 1A–B*). We found that the PVT$_{PBN}$

neurons sent projections to several brain areas, in particular the nucleus

accumbens core (NAc), BNST, and CeA (*Figure 7–figure supplement 1B–H*),

which was similar to the early tracing research of PVT efferent projections

(Kirouac, 2015).

Discussion

In this study, we employed viral tracing and electrophysiology to confirm the

monosynaptic excitatory connectivity between the PBN and the PVT.

Optogenetic or pharmacogenetic activation of the PBN-PVT projection or the

PVT-projecting PBN neurons induced anxiety-like, aversion-like, and fear-like

behaviors. Optogenetic inhibition of the PBN-PVT projection or the

PVT-projecting PBN neurons could partially reduce 2-MT induced aversive

behaviors as well as footshock-induced freezing behaviors. The activity of

PVT$_{PBN}$ neurons was increased in several aversive stimuli and could be further

increased by activation of PBN-PVT projection. Besides, activation of PVT$_{PBN}$

neurons induced anxiety-like behaviors. Taken together, our results reveal the

functional role of the PBN-PVT projection in modulating negative affective
states in mice.

PBN efferents and PBN-PVT monosynaptic excitatory projection

The PBN is a critical hub receiving sensory information from the spinal
cord *(Todd, 2010)*. The widespread distribution of PBN efferents contributes to
different aspects of behavioral and physiological responses. Previous studies
showed that the CGRP-expressing neurons in the PBN project to the CeA
contribute to the affective dimension of pain. In contrast, non-CGRP neurons
may transmit sensory pain information *(Han et al., 2015)*. The projections from
the PBN to the VMH or PAG are involved in producing escape behaviors to
avoid injury, while the projections from the PBN to the BNST or CeA participate
in facilitating aversive memory *(Chiang et al., 2020)*. The PBN neurons, which
receive projections from the spinal cord, form strong functional synaptic
connections with the ILN neurons but not the CeA neurons to process the
nociceptive signals *(Deng et al., 2020)*. The PVT nucleus located in the middle
line of the brain is an important area that participates in affective states
processing *(Kirouac, 2015)*. Although recent research has reported that the
projecting fibers from PBN were found in the PVT *(Chiang et al., 2020)*,
remarkably little is known about the connectivity information and function of the
PBN-PVT projection.

Since we injected the constitutively expressed ChR2 virus into the PBN,
few neurons in the LC (which is medial to the PBN) might be infected. The LC
neurons express the VgluT1 and also project to the PVT *(Beas et al., 2018)*.
Although the PBN-PVT projection comprises the major portion of the
projections, there is still potential contamination from the LC-PVT projections.
We also observed a small portion of inhibitory connections between the PBN
and the PVT. It is consistent with the previous study showing GABAergic
neurons in the PBN also send sparse projections to the PVT *(Chiang et al.,*
2020). Further in situ hybridization results confirmed that the PVT-projecting

PBN neurons are mainly glutamatergic neurons expressing *Vglut2* mRNA.

These results suggest that the majority of the PBN-PVT projection appeared to

be excitatory.

We also found that the density of PBN glutamatergic is higher in the middle

and posterior PVT (pPVT). These results are consistent with the various

studies supporting that pPVT is a particularly aversive region of the PVT (*Gao*

et al., 2020; *Beas et al.,* 2018; *Barson et al.,* 2020).

395 PBN-PVT projection modulates negative affective states

We found that activation of PBN-PVT projection or the PVT-projecting PBN

neurons induced anxiety-like behaviors and fear-like behaviors in the OFT and

EZM. We observed that mice displayed robust running and jumping behaviors

mainly in the first minute in optogenetic manipulation, and these phenomena

were not observed in the pharmacogenetic experiment. These might be

caused by an instantly increased activity of PBN-PVT projection induced by

optogenetic manipulation. Mice might display "fight or flight" during sudden

affective state transitions. And the pharmacogenetic approach takes several

minutes to gradually enhance neural activity, and the affective state changes in

a relatively mild way. We also observed that the anxiety-like behaviors in the

OFT still existed several minutes after optogenetic activation of the PBN-PVT

projection. However, in the RTPA test, the aversion appeared when the laser

was on and disappeared when the laser was off, indicating the aversion was

transient and could not be translated to associative learning. It was further

confirmed by the prolonged condition place aversion test and cue-dependent

optogenetic conditioning test. These results suggest that activation of the

PBN-PVT induces instant negative affective states but does not drive

associative fear memory formation.

Selectively optogenetic inhibition of PBN-PVT projection or PVT-projecting

PBN neurons could reduce aversion-like and fear-like behaviors. To better

examine the behavioral changes, we performed 10 minutes test in 2-MT and

footshock experiments. So we used a relatively long-term protocol in

optogenetic inhibition experiments (10 minutes constant laser). Such long-term

inhibition protocols were used in other studies (*Zhou et al.,* 2019; *Sun et al.,*

2020). We also performed the classical fear conditioning test and found that

inhibition of the PBN-PVT projection did not affect associative fear memory

formation or retrieval, suggesting that the PBN-PVT projection mainly

promotes aversion but does not facilitate negative association.

Our calcium imaging and Fos staining results indicated PVT neurons were

activated after exposure to aversive stimuli, consistent with a previous study

(Zhu et al., 2018). The dual Fos staining experiment and optoelectrode

experiments confirmed a broad overlap between laser-activated and

footshock-activated neurons. Further analysis showed that activation of the

PBN-PVT projection enhanced the overall firing rates of PVT neurons in

response to footshock. These results suggest that the activation of the

PBN-PVT could enhance the neuronal activity in response to aversive

stimulation.

The activation of PBN innervated PVT neurons induced anxiety-like

behaviors, suggesting the PVT_{PBN} neurons are involved in modulating

negative affective states. Previous studies have reported that activation of the

PBN-CeA pathway is sufficient to drive a series of negative affective states

behaviors *(Bowen et al., 2020; Cai et al., 2018; Han et al., 2015)*, enable

associative learning, and generate aversive memory *(Chiang et al., 2020)*.

Distinct from the PBN-CeA projection, we found that activation of the PBN-PVT

projection only induced transient aversion-related behaviors, and inhibition of

the PBN-PVT projection did not affect fear memory acquisition or retrieval. A

study reported that only a few Fluoro-gold (FG)/tetramethylrhodamine-dextran

(TMR) double-labeled neurons were sparsely distributed in the PBN of the

mice injected with FG into the PVT and TMR into the CeA *(Liang et al., 2016)*.

Our results also showed few collateral projecting fibers in the CeA or VMH

from the PVT-projecting PBN neurons. These results suggested that the

PBN-PVT pathway and the PBN-CeA pathway are two parallel pathways

originating from distinct efferent neurons within the PBN to perform distinct

functions. However, we also observed collateral projection fibers in BNST, LH,

PVN, PAG, but not CEA or VMH. The PBN-PAG projection is suggested to

mediate escaping behaviors *(Chiang et al., 2020)*. The possibility of antidromic

effects following photoactivation of PBN terminals in PVT should be reminded.

The tracing results showed the PVT_{PBN} neurons projected to multiple brain

areas, particularly the NAc, BNST, and CeA. The BNST and CeA have been

previously implicated in negative affective behaviors *(Jennings et al., 2013;*

Tye et al., 2011). Previous studies showed that the activation of the PVT-CeA

projections induces place aversion, and the effect persists on the next day in

the absence of photostimulation (*Do Monte et al.,* 2017). Similarly, long-term

depression (LTD)-like the stimulation of PVT-CeA projections or inhibition of

the same circuit induces a persistent attenuation of fear responses (*Chen and*

Bi., 2019; *Do Monte et al.,* 2015; *Penzo et al.,* 2015). These results revealed a

critical role of the PVT-CeA projection in aversive memory formation. In our

study, we found that PBN-PVT is not crucial for aversive memory formation.

The possible reason might be that manipulation of the PVT-CeA induces direct

excitatory inputs to the CeA, and the inputs are strong enough for aversive

memory formation. However, activation of the PBN-PVT projection might not

induce enough excitatory inputs to the CeA via the disynaptic connection.

A study also found that PVT mediates descending pain facilitation

underlying persistent pain conditions via the PVT-CeA-PAG circuit *(Liang et al.,*

2020). Different downstream pathways of PVT_{PBN} neurons might have different

functions and deciphering the circuit mechanisms needs further examination.

The potential role of PBN-PVT projection in depression and pain

It is worth noting that although the pharmacogenetic activation of the

PVT-projecting PBN neurons induced anxiety-like behaviors and fear-like

behaviors in the hM3Dq group mice, no depression-like symptoms were

observed in TST and FST. On the other side, chronic pain models, such as the

partial sciatic nerve ligation model, the spared nerve injury model, and

complete Freund's adjuvant model, generally induce anxiety and depression at

least 3–4 weeks after the surgery in mice *(Dimitrov et al., 2014; Zhou et al.,*

2019). Our study collected the behavioral data 30 minutes after a single dose

of CNO injection. Different behavioral tests were performed at least three days

apart to eliminate the residual CNO effects. We hypothesized that

depression-like behaviors might be observed if we repeatedly activate the

PBN-PVT projection for weeks. However, whether the PBN-PVT is involved in

depression is still unknown.

A recent study revealed that the PBN neurons convey nociception

information from the spinal cord to the ILN, which is relatively closed to the

PVT *(Deng et al., 2020).* In our results, we carefully checked the virus

expression and optic fiber locations of mice. We found that pharmacogenetic

activation of PVT-projecting PBN neurons did not affect the basal nociceptive

thresholds or formalin-induced licking behaviors. Moreover, no obvious

nociception-related behaviors (such as forelimb wiping, hindlimb flinching,

licking, or biting) were found through specific manipulation of the PBN

innervated PVT neurons, which suggests that the PBN-PVT projection might

be not involved in nociceptive information processing.

In summary, we identified the functional role of the PBN-PVT projection in

modulating negative affective states. Our study paves the way for further

deciphering the distinct roles of the PBN neural circuit in affective behaviors.

Acknowledgements

We thank Dr. Hua-Tai Xu for providing *Rosa26-tdTomato* mice. We thank Dr.

Yan-Gang Sun for providing *VgluT2-ires-Cre* mice. We thank all the lab

members of D.M. for their helpful discussion. This work was supported by the

National Natural Science Foundation of China (No. 31900717, 31571086), the

Shanghai Sailing Program (19YF1438700 to D.M.), the Young Elite Scientists

Author contributions

Y.B.Z. performed the virus injection experiments and behavioral experiments.

Y.W. performed the dual Fos staining experiments. X.X.H. and Y.W. performed

the optoelectrode experiments. R.Z. performed the electrophysiological

experiments. Y.B.Z., Y.W., M.Z.L., and P.F.L. performed the histological

experiments. J.B.L., L.Z., and D.M. designed the experiments. Y.B.Z. and D.M.

wrote the manuscript.

Materials and methods

Animals

Male C57Bl/6J wild-type mice, *Rosa26-tdTomato* mice (Jax Stock No: 007909,

gifted from Dr. Hua-Tai Xu, Institutes of Neuroscience, Chinese Academic of

Sciences), *VgluT2-ires-Cre* mice (Jax Stock No: 016963, gifted from Dr.

Yan-Gang Sun, Institutes of Neuroscience, Chinese Academic of Sciences)

were used. Animals were housed in standard laboratory cages in a

temperature (23-25°C)-controlled vivarium with a 12:12 light/dark cycle, free to

food and water. For tracing and behavioral experiments, the mice were

injected with the virus at 7–8 weeks old and performed the behavioral tests at

11–12 weeks old. For the electrophysiological experiments, the mice were

injected with the virus at 4–6 weeks old to accomplish the electrophysiological

experiments at 7–9 weeks old. For *in vivo* fiber photometry and optoelectrode

experiments, the mice were injected with the virus at 7–8 weeks old to

accomplish the experiments at 10–11 weeks old. All animal experiment

procedures were approved by the Animal Care and Use Committee of

Shanghai General Hospital (2019AW008).

Stereotaxic surgery

Mice were anesthetized by vaporized sevoflurane (induction, 3%; maintenance,

1.5%) and head-fixed in a mouse stereotaxic apparatus (RWD Life Science

Co.).

For electrophysiological experiments, the AAV2/8-hSyn-ChR2-mCherry

virus (300 nl, 4×10^{12} v.g./ml, AG26976, Obio Technology) was injected into

the PBN nucleus of WT mice in the stereotaxic coordinate: anteroposterior (AP)

–5.2 mm, mediolateral (ML) +1.3 mm, and dorsoventral (DV) –3.4 mm.

For tracing studies, the AAV2/8-EF1a-DIO-EGFP virus (300 nl, S0270,

Taitool Bioscience) was injected into the PBN (mentioned above) of

VgluT2-ires-Cre mice.

For the retrovirus injection surgery, the retrograde transport Cre

recombinase retroAAV2/2-hSyn-Cre virus (150 nl, 4×10^{12} v.g./ml,

S0278-2RP-H20, Taitool Bioscience) was injected in the *Rosa26-tdTomato*

mice at two locations of PVT respectively: (1) AP –1.22 mm, ML 0 mm, DV

–2.9 mm; (2) AP –1.46 mm, ML 0 mm, DV –2.9 mm.

For optogenetic activation of PVT-projecting PBN fibers, the

AAV2/9-EF1a-DIO-ChR2-mCherry virus (300 nl, 4×10^{12} v.g./ml, S0170,

Taitool Bioscience) or the AAV2/9-EF1a-DIO-mCherry virus (300 nl, 4×10^{12}

v.g./ml, AG20299, Obio Technology) were bilaterally injected into the PBN

(mentioned above) of *VgluT2-ires-Cre* mice, and a 200 μm diameter optic fiber

was implanted over the PVT (AP –1.46 mm, ML 0 mm, DV –2.9 mm) with a 20°

angle towards the midline.

For the pharmacogenetic activation of PVT-projecting PBN neurons, the

retroAAV2/2-hSyn-Cre virus (150 nl, 4×10^{12} v.g./ml, S0278-2RP-H20, Taitool

Bioscience) was injected into the PVT (AP –1.46 mm, ML 0 mm, DV –2.9 mm),

the AAV2/9-hSyn-DIO-hM3Dq-mCherry virus (300 nl, 4×10^{12} v.g./ml, PT-0019,

BrainVTA) or the control AAV2/9-EF1a-DIO-mCherry virus were bilateral

injected into the PBN (mentioned above) of the WT mice.

For optogenetic inhibition of PVT-projecting PBN fibers,

AAV2/9-EF1a-DIO-NpHR3.0-EYFP virus (300 nl, 4×10^{12} v.g./ml, AG26966,

Obio Technology) or the AAV2/8-EF1a-DIO-EGFP virus were bilaterally

injected into the PBN (mentioned above) of *VgluT2-ires-Cre* mice, and a 200

μm diameter optic fiber was implanted over the PVT (AP −1.46 mm, ML 0 mm,

DV −2.9 mm) with a 20° angle towards the midline.

For optogenetic inhibition of PVT-projecting PBN neurons,

retroAAV2/2-hSyn-Cre was injected into the PVT(AP −1.46 mm, ML 0 mm, DV

−2.9 mm), AAV2/9-EF1a-DIO-NpHR3.0-EYFP virus (300 nl, 4 x 10^{12} v.g./ml,

AG26966, Obio Technology) or the AAV2/8-EF1a-DIO-EGFP virus was

injected into the PBN of WT mice, the left optic fiber was implanted over the

PBN vertically and the right one were placed over the PBN with a 20° angle

towards the midline.

For *in vivo* fiber photometry experiments, the AAV2/8-hSyn-GCaMP6s

virus (200 nl, 4 x 10^{12} v.g./ml, S0225-8, Taitool Bioscience) was injected into

the PVT nucleus (AP −1.46 mm, ML 0 mm, DV −2.90 mm) of the WT mice, the

optic fiber was implanted above the PVT with a 20° angle towards the midline.

For optoelectrode experiments, the AAV2/9-EF1a-DIO-ChR2-mCherry

virus (300 nl, 4 x 10^{12} v.g./ml, AAV2/9-S0170, Taitool Bioscience) were

bilaterally injected into the PBN (mentioned above) of *VgluT2-ires-Cre* mice.

Three weeks later, the homemade optoelectrode was implanted into the PVT

nucleus (AP −1.46 mm, ML 0 mm, DV −2.90 mm).

For pharmacogenetic activation of PVT_{PBN} neurons, the AAV2/1-hSyn-Cre

virus (300 nl, 1.5 x 10^{13} v.g./ml, S0278-1-H50, Taitool Bioscience) was

bilaterally injected into the PBN nucleus, the

AAV2/9-hSyn-DIO-hM3Dq-mCherry virus or the control

AAV2/9-EF1a-DIO-mCherry virus was injected into the PVT (AP −1.46 mm, ML

0 mm, DV −2.9 mm) of the WT mice.

The virus was infused through a glass pipette (10–20 μm in diameter at the

tip) at the rate of 50–100 nl/minute. The injection pipette was left in place for

additional 8 minutes. After the surgeries, the skin was closed by the sutures,

and the optic fiber was secured through the dental acrylic. Generally, tracing,

electrophysiological or behavioral experiments were performed at least three
weeks later. After experiments, histological analysis was used to verify the
location of viral transduction and the optic fiber. The mice without correct
transduction of virus or correct site of optic fiber were excluded for analysis.

Histology

Animals were deeply anesthetized with vaporized sevoflurane and
transcardially perfused with 20 ml saline, followed by 20 ml paraformaldehyde
(PFA, 4% in PBS). Brains were extracted and soaked in 4% PFA at 4°C for a
minimum of 4 hours and subsequently cryoprotected by transferring to a 30%
sucrose solution (4°C, dissolved in PBS) until brains were saturated (for 36–48
hours). Coronal brain sections (40 μm) were cut using a freezing microtome
(CM1950, Leica). The slices were collected and stored in PBS at 4°C until
immunohistochemical processing. Nuclei were stained with DAPI (Beyotime,
1:10000) and washed three times with PBS.

The brain sections undergoing immunohistochemical staining were
washed in PBS 3 times (10 minutes each time) and incubated in a blocking
solution containing 0.3% TritonX-100 and 5% normal donkey serum (Jackson
ImmunoResearch, USA) in PBS for 1 hour at 37°C. Sections were then
incubated (4°C, 24 hours) with primary antibodies dissolved in 1% normal
donkey serum solution. Afterward, sections were washed in PBS 4 times (15
minutes each time), then incubated with secondary antibodies for 2 hours at
room temperature. After DAPI staining and washing with PBS, sections were
mounted on glass microscope slides, dried, and covered with 50% glycerin
(ThermoFisher). The images were taken by the Leica DMi8 microscope and
the Leica SP8 confocal microscopy. The images were further processed by Fiji
and Photoshop.

RNAscope in situ hybridization

Mice were anesthetized with isoflurane and rapidly decapitated. Brains were
roughly dissected from perfused mice and post-fixed in 4% PFA at 4 °C
overnight, dehydrated in 30% sucrose 1×PBS at 4 °C for 2 days. Mouse brains

were embedded in OCT compound, cryosectioned in 15 µm coronal slices,

and mounted on SuperFrost Plus Gold slides (Fisher Scientific). In

situ hybridization was performed according to the protocol of the RNAscope

Multiplex Fluorescent Reagent Kit v2 (Cat. No. 320293). Probes were

purchased from Advanced Cell Diagnostics: *c-Fos* (Cat. No. 316921-C2), *Tac1*

(Cat. No. 410351-C2), *Tacr1* (Cat. No. 428781-C2), *Pdyn* (Cat. No. 318771),

and *VgluT2* (Cat. No. 319171-C2). Primary antibodies include rabbit anti-c-Fos

(Abcam, cat. No. ab190289, 1:4000), goat anti-CGRP (Abcam, ab36001,

1:1000), and rabbit anti-DsRed (Clontech, Cat. No. 632496, 1:500). All

secondary antibodies were purchased from Jackson ImmunoResearch and

used at 1:400 dilution. Secondary antibodies include Alexa 488 donkey

anti-rabbit (Cat. No. 711-545-152), Cy3 donkey anti-rabbit (Cat. No.

711-165-152), and Alexa 488 donkey anti-goat (Cat. No. 705-546-147).

Images were collected on a Leica fluorescence microscope and Leica LAS

Software.

Fos induction

The mice were habituated for three days and performed gentle grabbing and

holding for 1 minute, five times every day to minimize background Fos

expression.

To study the effect of pharmacogenetic manipulations on PVT-projecting

PBN neurons, we intraperitoneally injected 0.5 mg/kg clozapine N-oxide (CNO,

Sigma). Ninety minutes later, the brain tissues were processed. To assess

2-MT evoked Fos expression in the PVT, the mice were kept in a chamber with

a floor covered with cotton containing 100 ml, 1:1000 2-MT volatilized the

predator odor for 90 minutes. To assess footshock-induced Fos expression in

the PVT, we placed the mice into the chamber and delivered 30 times

inevitable footshock (0.5 mA, 1 second) with a variable interval (averaging 60

seconds). After stimulation, animals were kept in the same apparatus for

another 60 minutes, and brain tissues were then processed.

For the dual Fos experiments, we first delivered 20 minutes 473 nm laser

pulses (20 Hz, 5 mW, 5ms) and left the mice to rest in the homecage for 60

minutes. Then we delivered the 20 minutes footshock stimulus (0.5 mA, 1 s, 30

times) and perfused the mice.

Electrophysiology

The electrophysiological experiment was performed as previously described

(Mu et al., 2017). Mice were anesthetized with sevoflurane and perfused by

the ice-cold solution containing (in mM) sucrose 213, KCl 2.5, NaH_2PO_4 1.25,

$MgSO_4$ 10, $CaCl_2$ 0.5, $NaHCO_3$ 26, glucose 11 (300–305 mOsm). Brains were

quickly dissected, and the coronal slice (250 µm) containing the PBN or PVT

were chilled in ice-cold dissection buffer using a vibratome (V1200S, Leica) at

a speed of 0.12 mm/second. The coronal sections were subsequently

transferred to a chamber and incubated in the artificial cerebrospinal fluid

(ACSF, 34°C) containing (in mM): NaCl 126, KCl 2.5, NaH_2PO_4 1.25, $MgCl_2$ 2,

$CaCl_2$ 2, $NaHCO_3$ 26, glucose 10 (300–305 mOsm) to recover for at least 40

minutes, then kept at room temperature before recording. All solutions were

continuously bubbled with 95% O_2/5% CO_2.

All experiments were performed at near-physiological temperatures

(30–32°C) using an in-line heater (Warner Instruments) while perfusing the

recording chamber with ACSF at 3 ml/minute using a pump (HL-1, Shanghai

Huxi). Whole-cell patch-clamp recordings were made from the target neurons

under IR-DIC visualization and a CCD camera (Retiga ELECTRO, QIMAGING)

using a fluorescent Olympus BX51WI microscope. Recording pipettes (2–5

MΩ; Borosilicate Glass BF 150-86-10; Sutter Instrument) were prepared by a

micropipette puller (P97; Sutter Instrument) and backfilled with

potassium-based internal solution containing (in mM) K-gluconate 130, $MgCl_2$

1, $CaCl_2$ 1, KCl 1, HEPES 10, EGTA 11, Mg-ATP 2, Na-GTP 0.3 (pH 7.3, 290

mOsm) or cesium-based internal solution contained (in mM) $CsMeSO_3$ 130,

$MgCl_2$ 1, $CaCl_2$ 1, HEPES 10, QX-314 2, EGTA 11, Mg-ATP 2, Na-GTP 0.3 (pH

7.3, 295 mOsm). Biocytin (0.2%) was included in the internal solution.

In PBN-PVT ChR2 experiments, whole-cell recordings of PBN neurons

with current-clamp (I = 0 pA) were obtained with pipettes filled with the

potassium-based internal solution. The 473 nm laser (5 Hz, 10 Hz, 20 Hz

pulses, 0.5 ms duration, 2 mW/mm^2) was used to activate PBN ChR2 positive

neurons. Light-evoked EPSCs and IPSCs of PVT neurons recorded with

voltage-clamp (holding voltage of -70 mV or 0 mV) were obtained with pipettes

filled with the cesium-based internal solution. The 473 nm laser (20 Hz paired

pulses, 1 ms duration, 4 mW/mm^2) was used to activate ChR2 positive fibers.

The light-evoked EPSCs were completely blocked by 1 μM TTX (tetrodotoxin),

rescued by 100 μM 4-AP (4-Aminopyridine), and blocked by 10 μM NBQX

(6-nitro-7-sulphamoylbenzo(f)quinoxaline-2,3-dione). NBQX and TTX were

purchased from Tocris Bioscience. All other chemicals were obtained from

Sigma.

Voltage-clamp and current-clamp recordings were carried out using a

computer-controlled amplifier (MultiClamp 700B; Molecular Devices, USA).

During recordings, traces were low-pass filtered at 4 kHz and digitized at 10

kHz (DigiData 1550B1; Molecular Devices). Data were acquired by Clampex

10.6 and filtered using a low-pass-Gaussian algorithm (-3 dB cut-off frequency

= 1000 Hz) in Clampfit 10.6 (Molecular Devices).

Optogenetic manipulation

For activating the PBN-PVT projection, a 473 nm laser (20 Hz, 5 ms pulse

duration, 5 mW) was delivered. For inhibition of the PBN-PVT projection and

the PVT-projecting PBN neurons, a constant laser (589 nm, 10 mW) was

delivered.

Pharmacogenetic manipulation

All behavioral tests were performed 30 minutes after intraperitoneal injection of

0.5 mg/kg CNO in pharmacogenetic manipulation. Different behavior tests

were performed at least three days apart.

Open field test

The open field test (OFT) was used to assess locomotor activity and

anxiety-related behavior in an open field arena (40 x 40 x 60 cm) with opaque

plexiglass walls. The mouse was placed in the center of the box and recorded by a camera attached to a computer. The movement was automatically tracked and analyzed by AniLab software (Ningbo AnLai, China). The total distance traveled, the total velocity, the total unmoving time (the mice were considered to be unmoving if unmoving time lasts more than 1 s), and time spent in the center area (20 x 20 cm) were measured. The box was cleaned with 70% ethanol after each trial.

To assess the effect of optogenetic activation of the PBN-PVT projection, 15 minutes sessions consisting of 5 minutes pre-test (laser OFF), 5 minutes laser on test (laser ON), and 5 minutes post-test (laser OFF) periods. Laser (473 nm, 20 Hz, 5 ms, 5 mW) was delivered during the laser on phase.

To assess the effect of pharmacogenetic manipulations of PVT-projecting PBN neurons on locomotor activity and affective behaviors, we recorded the the movement 30 minutes after intraperitoneal (i.p.) injection with CNO.

To assess the effect of inhibition of the PBN-PVT projection on the aversive behaviors induced by 2-MT. One cotton ball containing 5 ml 2-MT (1:1000) solution was placed on the center of the upper left quadrant to disseminate fear-odor, then a constant laser (589 nm, 10 mW) was delivered during the 10 minutes test. The time spent in the 2-MT paired quadrant was calculated.

Elevated zero maze (EZM)

The EZM was an opaque plastic circle (60 cm diameter), which consisted of four sections with two opened and two closed quadrants. Each quadrant had a path width of 6 cm. The maze was elevated 50 cm above the floor. The animals were placed into an open section facing a closed quadrant and freely explored the maze for 5 minutes.

Real-time place aversion (RTPA) test

Mice were habituated to a custom-made 20 x 30 x 40 cm two-chamber apparatus (distinct wall colors and stripe patterns) before the test. Each mouse was placed in the center and allowed to explore both chambers without laser

stimulation for 10 minutes on Day 1. The movement was recorded for 10 minutes as a baseline. The mice performed a slight preference for the black chamber according to the fact the mice have innate aversive to brightly illuminated areas. On Day 2, 473 nm laser stimulation (20 Hz, 5 ms, 5 mW) was automatically delivered when the mouse entered or stayed in the black chamber and turned off when the mouse exited the black chamber for 10 minutes. Finally, the mouse was allowed to freely explore both chambers without laser stimulation for another 10 minutes. The RTPA location plots and total time on the stimulated side were recorded and counted with the AniLab software.

Condition place aversion (CPA)

After habituation, mice were placed in the center of the two-chamber apparatus and allowed to explore either chamber for 15 minutes on Day 1. On Day 2, mice were restricted to one chamber (laser paired chamber) with photostimulation (473 nm, 20 Hz, 5 ms, 5 mW) for 30 minutes in the morning and restricted to the other chamber (unpaired chamber) without photostimulation in the afternoon. On Day 3, mice were restricted to the unpaired chamber without photostimulation in the morning and restricted to the laser paired chamebr with photostimulation in the afternoon. On Day 4, mice were allowed to explore both chambers without laser stimulation for another 15 minutes. The time in the laser-paired chamber was calculated on Day 1 and Day 4.

2-MT-induced aversion

To assess the effect of optogenetic inhibition of the PBN-PVT projection or PVT-projecting PBN neurons on the aversive state, three cotton balls containing 15 ml 2-MT (1:1000) solution were placed in the black chamber. A constant laser (589 nm, 10 mW) was delivered during the 10 minutes test.

Cue-dependent optogenetic conditioning test

Video Freeze fear conditioning system with optogenetic equipment (MED Associates, MED-VFC-OPTO-USB-M) and Video Freeze software were used.

On Day 1, mice were habituated to the fear conditioning chambers and
allowed to explore for 2 minutes freely, then three tones (75 dB, 4 kHz, 30
seconds duration) separated by a variable interval with a range of 60–120
seconds and the average of 90 seconds were delivered.

On Day 2, mice were trained with the sound cue (75 dB, 4 kHz, 30
seconds) paired with a simultaneous 30 seconds laser pulse train (20 Hz, 5 ms,
5 mW) for six times separated by a variable interval (averaging 90 seconds).
The mice were kept in the conditioning chamber for another 60 seconds before
returning to the home cages.

On Day 3, mice were placed back into the original training chamber for 3
minutes to perform the contextual test. After 2–3 hours, the conditioning
chamber was modified by changing its metal floor and sidewalls. Mice were
placed in the altered chamber for 3 minutes to measure the freezing level in
the altered context. A tone (75 dB, 4 kHz) was delivered for 30 seconds to
perform the cue test.

The behavior of the mice was recorded and analyzed with the Video
Freeze software. Freezing was defined as the complete absence of movement
for at least 0.5 seconds. On the conditioning day, the freezing percentages
were calculated for 30 seconds after each tone/laser stimulus. For the
contextual test, the freezing percentages were calculated for three minutes.
For the cue test, the freezing percentages were calculated for 30 seconds
during tone.

Auditory fear conditioning test

On Day one, mice were habituated to the fear conditioning chambers. On Day
two, mice were conditioned by seven trials of sound tone (75 dB, 4 kHz, 30 s)
co-terminated with footshock (0.6 mA, 2 s) averagely separated by 90 seconds.
Laser (589 nm, 10 mW) was delivered 1 second before the footshock and
lasted for 4 seconds at each trial. On Day 3, mice were placed back into the
original training chamber for 3 minutes to perform the contextual test, and the
laser was delivered during the second minute. After 2–3 hours, the mice were

placed into a modified chamber to perform the cue test. Three tones were

given averagely separated by 90 seconds. The laser was delivered during the

second tone.

The behavior of the mice was recorded and analyzed with the Video

Freeze software. The freezing percentages of the 27 seconds tone before

laser (to avoid the influence of laser) for each trial were summarized to indicate

fear memory acquisition in the conditioning test. For the contextual test, the

freezing percentages were calculated for every minute. For the cue test, the

freezing percentages were calculated for 30 seconds during tone.

Freezing behavior

For analyses of freezing behavior induced by pharmacogenetic activation

of PVT-projecting PBN neurons, we injected CNO and recorded the mouse

behavior using the Video Freeze fear conditioning system 30 minutes later.

The Video Freeze fear conditioning system (MED Associates,

MED-VFC-OPTO-USB-M) was also used to assess the effect of optogenetic

inhibition of PBN-PVT projection and the PVT-projecting PBN neurons on the

fear-like behavior induced by footshock. After free exploration of the chamber

for 2 minutes, 15 times footshocks (0.6 mA,1 second) were delivered within 10

minutes with a constant 589 nm laser (10 mW). The freezing percentages

during 10 minutes were analyzed.

The Video Freeze fear conditioning system was also used to assess the

effect of optogenetic inhibition of the PVT-projecting PBN neurons on the

fear-like behavior induced by 2-MT. 10 ml 2-MT (1:1000) dissolved in the

ddH$_2$O was soaked into the cotton ball on the bottom of the training box. A

constant laser (589 nm, 10 mW) was delivered during the tests.

Tail suspension test (TST)

Mice were individually suspended by an adhesive tape placed roughly 2 cm

from the tip of the tail and videotaped for 6 minutes. Mice were considered

immobile without initiated movements, and the immobility time was scored in

the last 3 minutes by an observer unknown of the treatments.

Forced swim test (FST)

Mice were individually placed for 6 minutes in clear cylinders (45 cm height, 20 cm internal diameter) containing freshwater (25°C, 15 cm depth). The swimming activity was videotaped, and immobility time in the last 3 minutes was counted manually by an investigator unaware of animal grouping. The mice were considered immobile when they stopped swimming/struggling or only slightly moved to keep the nose above the surface.

von Frey test

The von Frey test was used to assess the mechanical sensitivity *(Mu et al., 2017)*. The mice were acclimated to the observation chambers for two days (2 hours for each day) before the test. A series of von Frey hairs with logarithmically incrementing stiffness (0.16–2.0 grams) were used to stimulate the mouse hind paw perpendicularly. The 50% paw withdrawal threshold was determined using the up-down method.

Hargreaves test

Hargreaves tests were performed as described previously *(Mu et al., 2017)*. Mice were placed in an individual plexiglass box with a glass floor. A radiant heat beam was exposed directly to the hind paw until the paw was withdrawn. The trials were repeated three times with an interval of at least 15 minutes. To avoid potential damage, the test was executed with a 20 seconds cut-off time.

Formalin test

In the formalin test, the mice received an intraplantar injection of formalin (5%, 20 µl/mouse) and were placed into a plexiglass box (width: 10 cm, length: 10 cm, height: 15 cm) individually to record the pain-related licking behaviors for 1 hour. All videos were analyzed by trained investigators blinded to the experimental treatment of the animals.

Rotarod test

Mice were trained twice on a rotarod apparatus (MED Associates) with a rod accelerated 5–20 revolutions per minute (r.p.m.) for 5 minutes before the experimental day. On the second day, each mouse underwent three trials with

a rod was programmed to accelerate from 0 to 40 rpm over 300 seconds, then the average rpm at the point of falling was recorded.

Fiber photometry

In vivo fiber photometry experiments were performed as previously described *(Zhu et al., 2020)*. After two weeks for virus expression, the mice were gently handled to be familiar with the calcium signal recording experiments (Thinker-Biotech). A signal (for synchronization) was manually tagged with the shock and air puff to evaluate the activity of PVT neurons. The calcium transient was recorded at 50 Hz. The fluorescence values change (ΔF/F) was calculated from the formula of (F−F0)/F0 and the F0 represented the median of the fluorescence values in the baseline period (−1 to −0.5 seconds relative to the stimulation onset). To precisely quantify the change of the fluorescence values across the shock or air-puff stimulation, we defined 0.5 to 1.0 seconds after the onset as the post-stimulus period.

Optoelectrode recording and analysis

The homemade optoelectrode consisted of an optic fiber (200 mm in diameter) glued to 16 individually insulated nichrome wires (35 µm internal diameter, 300-900 Kohm impedance, Stablohm 675, California Fine Wire). The 16 microwire arrays were arranged in a 4 + 4 + 4 + 4 pattern and soldered to an 18-pin connector (Mil-Max). Three weeks after virus injection, the optoelectrode was implanted to the PVT nucleus (AP −1.46 mm, ML 0 mm, DV −2.90 mm). After one week of recovery, two trials were performed continuously. Trial 1 contained ten sweeps of 2 s laser pulse trains (473 nm, 5 ms, 20 Hz, 8 mW). The interval of sweeps was 60 s. Trial 2 contained twenty sweeps of 2 s footshock (0.5 mA). The interval of sweeps was 60 s. In the even time sweeps (2, 4, 6, 8,10, 12, 14, 16, 18, 20), 2 s laser pulse trains were delivered spontaneously with the 2 s footshock. Neuronal signals were recorded using a Zeus system (Zeus, Bio-Signal Technologies: McKinney, TX, USA), and Spike signals were filtered online at 300 Hz. At the end of the experiment, all animals were perfused to confirm the optical fiber sites. Only the data of animals with

correct optical fiber sites and virus expression regions were analyzed.

The spikes were sorted by the valley-seeking method with Offline Sorter

software (Plexon, USA) and analyzed with NeuroExplorer (Nex Technologies:

Boston, MA, U.S.A.). Firing rates of the neurons and timestamps were

exported for further analysis using customized scripts in MATLAB. The

Kolmogorov-Smirnov (K-S) test was used to compare the spike firing rate of

PVT during 2 s baseline (before stimulus) and 2 s after each stimulus. $p <$

0.001 indicated statistical significance. Z-score normalization maps were

constructed from normalized firing rates.

Quantification of the fiber intensity

For quantification of fluorescence of PVT_{PBN} efferents, the downstream targets

of PVT_{PBN} neurons were taken photofluorograms with identical exposure time,

the mean fluorescence value in each ROI (400 x 400 pixels) of each brain

region was analyzed by Fiji. The fiber intensity was calculated as the

fluorescence value of each brain region divided by that of the NAc. All data

came from at least three different mice and were presented as mean ± SEM.

Analysis

Statistical detection methods include unpaired student's *t*-test, paired student's

t-test, one-way ANOVA with Bonferroni's correction for multiple comparisons,

two-way ANOVA with Bonferroni's correction for multiple comparisons. A value

of $p < 0.05$ was considered statistically significant. All data were represented

as mean ± SEM.

REFERENCES

Armbruster BN, Li X, Pausch MH, Herlitze S, Roth BL. (2007). Evolving the

lock to fit the key to create a family of G protein-coupled receptors potently

activated by an inert ligand. *PNAS,* **104**:5163-5168. DOI:

https://doi.org/10.1073/pnas.0700293104, PMID:17360345

Barson JR, Mack NR, Gao WJ. The paraventricular nucleus of the thalamus is

an important node in the emotional processing network. (2020). *Front Behav*

Neurosci, **14**:598469. DOI: 10.3389/fnbeh.2020.598469. PMID: 33192373

Beas BS, Wright BJ, Skirzewski M, Leng Y, Hyun JH, Koita O, Ringelberg N,

Kwon HB, Buonanno A, Penzo MA. (2018). The locus coeruleus drives

disinhibition in the midline thalamus via a dopaminergic mechanism. *Nature*

Neuroscience, **21**:963-973. DOI: https://doi.org/10.1038/s41593-018-0167-4,

PMID:29915192

Bowen AJ, Chen JY, Huang YW, Baertsch NA, Park S, Palmiter RD. (2020).

Dissociable control of unconditioned responses and associative fear learning

by parabrachial CGRP neurons. *eLife,* **9**. DOI:

https://doi.org/10.7554/eLife.59799, PMID:32856589

Cai YQ, Wang W, Paulucci-Holthauzen A, Pan ZZ. (2018). Brain circuits

mediating opposing effects on emotion and pain. *Journal of Neuroscience,*

38:6340-6349. DOI: https://doi.org/10.1523/JNEUROSCI.2780-17.2018,

PMID:29941444

Campos CA, Bowen AJ, Roman CW, Palmiter RD. (2018). Encoding of danger

by parabrachial CGRP neurons*. Nature,* **555**:617-622. DOI:

https://doi.org/10.1038/nature25511, PMID:29562230

Chen M, Bi LL. (2019). Optogenetic long-term depression induction in the

PVT-CeL circuitry mediates decreased fear memory. *Mol Neurobiol,*

56:4855-4865. DOI: 10.1007/s12035-018-1407-z. PMID: 30406427

Chiang MC, Bowen A, Schier LA, Tupone D, Uddin O, Heinricher MM. (2019).

Parabrachial complex: a hub for pain and aversion. *Journal of Neuroscience,*

39:8225-8230. DOI: https://doi.org/10.1523/JNEUROSCI.1162-19.2019,

PMID:31619491

Chiang MC, Nguyen EK, Canto-Bustos M, Papale AE, Oswald AM, Ross SE.

(2020). Divergent neural pathways emanating from the lateral parabrachial

nucleus mediate distinct components of the pain response*. Neuron,*

106:927-939. DOI: https://doi.org/10.1016/j.neuron.2020.03.014,

PMID:32289251

Deng J, Zhou H, Lin JK, Wei YC, Xu XH, Sun YG. (2020). The parabrachial

nucleus directly channels spinal nociceptive signals to the intralaminar thalamic nuclei, but not the amygdala. *Neuron,* **107**:909-923. DOI: https://doi.org/10.1016/j.neuron.2020.06.017, PMID:32649865

Dimitrov EL, Tsuda MC, Cameron HA, Usdin TB. (2014). Anxiety- and depression-like behavior and impaired neurogenesis evoked by peripheral neuropathy persist following resolution of prolonged tactile hypersensitivity. *Journal of neuroscience,* **34**:12304-12312. DOI: https://doi.org/10.1523/JNEUROSCI.0312-14.2014, PMID:25209272

Do-Monte FH, Minier-Toribio A, Quiñones-Laracuente K, Medina-Colón EM, Quirk GJ. (2017). Thalamic regulation of sucrose seeking during unexpected reward omission. *Neuron*, **94**:388-400. DOI: 10.1016/j.neuron.2017.03.036, PMID:28426970

Do-Monte FH, Quinones-Laracuente K, Quirk GJ. (2015). A temporal shift in the circuits mediating retrieval of fear memory. *Nature,* **519**:460-463. DOI: https://doi.org/10.1038/nature14030, PMID:25600268

Engelke DS, Zhang XO, O'Malley JJ, Fernandez-Leon JA, Li S, Kirouac GJ, Beierlein M, Do-Monte FH.(2021). A hypothalamic-thalamostriatal circuit that controls approach-avoidance conflict in rats. Nat Commun, 12:2517. DOI: 10.1038/s41467-021-22730-y. PMID: 33947849

Gao C, Leng Y, Ma J, Rooke V, Rodriguez-Gonzalez S, Ramakrishnan C, Deisseroth K, Penzo MA. (2020). Two genetically, anatomically and functionally distinct cell types segregate across anteroposterior axis of paraventricular thalamus. *Nat Neurosci,* **23**:217-228. DOI: 10.1038/s41593-019-0572-3. PMID: 31932767

Han S, Soleiman MT, Soden ME, Zweifel LS, Palmiter RD. (2015). Elucidating an affective pain circuit that creates a threat memory. *Cell,* **162**:363-374. DOI: https://doi.org/10.1016/j.cell.2015.05.057, PMID:26186190

Hsu DT, Kirouac GJ, Zubieta JK, Bhatnagar S. (2014). Contributions of the paraventricular thalamic nucleus in the regulation of stress, motivation, and mood. *Frontiers in Behavioral Neuroscience,* **8**:73. DOI:

https://doi.org/10.3389/fnbeh.2014.00073, PMID:24653686

Huang TW, Lin SH, Malewicz NM, Zhang Y, Zhang Y, Goulding M, LaMotte RH,

Ma QF. (2019). Identifying the pathways required for coping behaviours

associated with sustained pain. *Nature,* **565**:86-90. DOI:

https://doi.org/10.1038/s41586-018-0793-8, PMID:30532001

Isosaka T, Matsuo T, Yamaguchi T, Funabiki K, Nakanishi S, Kobayakawa R,

Kobayakawa K. (2015). Htr2a-expressing cells in the central amygdala

control the hierarchy between innate and learned fear. *Cell,* **163**:1153-1164.

DOI: https://doi.org/10.1016/j.cell.2015.10.047, PMID:26590419

Jennings JH, Sparta DR, Stamatakis AM, Ung RL, Pleil KE, Kash TL, Stuber

GD. (2013). Distinct extended amygdala circuits for divergent motivational

states. *Nature,* **496**:224-228. DOI: https://doi.org/10.1038/nature12041,

PMID:23515155

Jimenez JC, Su K, Goldberg AR, Luna VM, Biane JS, Ordek G, Zhou P, Ong

SK, Wright MA, Zweifel L, Paninski L, Hen R, Kheirbek MA. (2018). Anxiety

cells in a hippocampal-hypothalamic circuit. *Neuron,* **97**:670-683. DOI:

https://doi.org/10.1016/j.neuron.2018.01.016, PMID:29397273

Kaur S, Pedersen NP, Yokota S, Hur EE, Fuller PM, Lazarus M, Chamberlin

NL, Saper CB. (2013). Glutamatergic Signaling from the Parabrachial

Nucleus Plays a Critical Role in Hypercapnic Arousal. *Journal of*

Neuroscience, **33**:7627-7640. DOI:

https://doi.org/10.1523/Jneurosci.0173-13.2013,

PMID:WOS:000318420400003

Keyes PC, Adams EL, Chen Z, Bi L, Nachtrab G, Wang VJ, Tessier-Lavigne M,

Zhu Y, Chen X. (2020). Orchestrating opiate-associated memories in

thalamic circuits. *Neuron,* **107**:1113-1123.e4. DOI:

10.1016/j.neuron.2020.06.028. PMID: 32679036

Kirouac GJ. (2015). Placing the paraventricular nucleus of the thalamus within

the brain circuits that control behavior. *Neuroscience and Biobehavioral*

Reviews, **56**:315-329. DOI: https://doi.org/10.1016/j.neubiorev.2015.08.005,

PMID:26255593

Li S, Kirouac GJ. (2012). Sources of inputs to the anterior and posterior aspects of the paraventricular nucleus of the thalamus. *Brain Structure & Function,* **217**:257-273. DOI: https://doi.org/10.1007/s00429-011-0360-7, PMID:22086160

Liang SH, Yin JB, Sun Y, Bai Y, Zhou KX, Zhao WJ, Wang W, Dong YL, Li YQ. (2016). Collateral projections from the lateral parabrachial nucleus to the paraventricular thalamic nucleus and the central amygdaloid nucleus in the rat. *Neuroscience Letters,* **629**:245-250. DOI: https://doi.org/10.1016/j.neulet.2016.07.017, PMID:27423318

Liang SH, Zhao WJ, Yin JB, Chen YB, Li JN, Feng B, Lu YC, Wang J, Dong YL, Li YQ. (2020). A neural circuit from thalamic paraventricular nucleus to central amygdala for the facilitation of neuropathic pain. *Journal of Neuroscience,* **40**:7837-7854. DOI: https://doi.org/10.1523/JNEUROSCI.2487-19.2020, PMID:32958568

Mu D, Deng J, Liu KF, Wu ZY, Shi YF, Guo WM, Mao QQ, Liu XJ, Li H, Sun YG. (2017). A central neural circuit for itch sensation. *Science,* **357**:695-699. DOI: https://doi.org/10.1126/science.aaf4918, PMID:28818946

Nakahara TS, Carvalho VMA, Souza MAA, Trintinalia GZ, Papes F. (2020). Detection of activated mouse neurons with temporal resolution via dual c-Fos staining. *STAR Protoc,* **1**:100153. DOI: 10.1016/j.xpro.2020.100153. PMID: 33377047

Öhman A, Mineka S. (2001). Fears, phobias, and preparedness: Toward an evolved module of fear and fear learning. *Psychological Review,* **108**:483-522. DOI: https://doi.org/10.1037/0033-295x.108.3.483, PMID:11488376

Palmiter RD. (2018). The parabrachial nucleus: CGRP neurons function as a general alarm. *Trends in Neurosciences,* **41**:280-293. DOI: https://doi.org/10.1016/j.tins.2018.03.007, PMID:29703377

Penzo MA, Robert V, Tucciarone J, De Bundel D, Wang M, Van Aelst L, Darvas M, Parada LF, Palmiter RD, He M, Huang ZJ, Li B. (2015). The

paraventricular thalamus controls a central amygdala fear circuit. *Nature,* **519**:455-459. DOI: https://doi.org/10.1038/nature13978, PMID:25600269

Ren SC, Wang YL, Yue FG, Cheng XF, Dang RZ, Qiao QC, Sun XQ, Li X, Jiang Q, Yao JW, Qin H, Wang GZ, Liao X, Gao D, Xia JX, Zhang J, Hu B, Yan JA, Wang YJ, Xu M, et al. (2018). The paraventricular thalamus is a critical thalamic area for wakefulness. *Science,* **362**:429-434. DOI: https://doi.org/10.1126/science.aat2512, PMID:30361367

Saper CB. (2016). The house alarm. *Cell Metabolism,* **23**:754-755. DOI: https://doi.org/10.1016/j.cmet.2016.04.021, PMID:27166934

Sun L, Liu R, Guo F, Wen MQ, Ma XL, Li KY, Sun H, Xu CL, Li YY, Wu MY, Zhu ZG, Li XJ, Yu YQ, Chen Z, Li XY, Duan SM. (2020). Parabrachial nucleus circuit governs neuropathic pain-like behavior. *Nature Communications,* **11**. DOI: https://doi.org/10.1038/s41467-020-19767-w, PMID:33239627

Todd AJ. (2010). Neuronal circuitry for pain processing in the dorsal horn. *Nature Review Neuroscience,* **11**:823-836. DOI: https://doi.org/10.1038/nrn2947, PMID:21068766

Tye KM, Prakash R, Kim SY, Fenno LE, Grosenick L, Zarabi H, Thompson KR, Gradinaru V, Ramakrishnan C, Deisseroth K. (2011). Amygdala circuitry mediating reversible and bidirectional control of anxiety. *Nature,* **471**:358-362. DOI: https://doi.org/10.1038/nature09820, PMID:21389985

Vertes RP, Linley SB, Hoover WB. (2015). Limbic circuitry of the midline thalamus. *Neuroscience and Biobehavioral Review,* **54**:89-107. DOI: https://doi.org/10.1016/j.neubiorev.2015.01.014, PMID:25616182

Yamamuro K, Bicks LK, Leventhal MB, Kato D, Im S, Flanigan ME, Garkun Y, Norman KJ, Caro K, Sadahiro M, Kullander K, Akbarian S, Russo SJ, Morishita H. (2020). A prefrontal-paraventricular thalamus circuit requires juvenile social experience to regulate adult sociability in mice. *Nature Neuroscience,* **23**:1240-1252. DOI: https://doi.org/10.1038/s41593-020-0695-6, PMID:32868932

Zhou WJ, Jin Y, Meng Q, Zhu X, Bai TJ, Tian YH, Mao Y, Wang LK, Xie W,

Zhong N, Luo M, H., Tao WJ, Wang HT, Li J, Qiu BS, Zhou JN, Li XY, Xu H,

Wang K, Zhang XC, et al. (2019). A neural circuit for comorbid depressive

symptoms in chronic pain. *Nature Neuroscience,* **22**:1649-1658. DOI:

https://doi.org/10.1038/s41593-019-0468-2, PMID:31451801

Zhu YJ, Nachtrab G, Keyes PC, Allen WE, Luo LQ, Chen XK. (2018). Dynamic

salience processing in paraventricular thalamus gates associative learning.

Science, **362**:423-429. DOI: https://doi.org/10.1126/science.aat0481,

PMID:30361366

Zhu Y, Wienecke CF, Nachtrab G, Chen X. (2016). A thalamic input to the

nucleus accumbens mediates opiate dependence. *Nature,* **530**:219-222. DOI:

https://doi.org/10.1038/nature16954, PMID:26840481

Zhu YB, Xu L, Wang Y, Zhang Rui, Wang YC, Li JB, Mu D. (2020). Posterior

thalamic nucleus mediates facial histaminergic itch. *Neuroscience,* **444**:54-63.

DOI: https://doi.org/10.1016/j.neuroscience.2020.07.048, PMID:32750381

Zingg B, Chou XL, Zhang ZG, Mesik L, Liang F, Tao HW, Zhang LI. (2017).

AAV-Mediated anterograde transsynaptic tagging: mapping corticocollicular

input-defined neural pathways for defense behaviors. *Neuron,* **93**:33-47. DOI:

https://doi.org/10.1016/j.neuron.2016.11.045, PMID:27989459

1095

1096

 Figures and figure legends

1098

Figure 1. Functional connectivity pattern of the PBN-PVT projection. (A) The

schematic for virus injection of AAV2/8-hSyn-ChR2-mCherry into the PBN

nucleus and the slice recording with 473 nm laser stimulation. (B) The

projection fibers in the PVT nucleus. Scale bar: 100 µm. (C) The

AAV2/8-hSyn-ChR2-mCherry virus expression in the PBN nucleus. Scale bar:

200 µm. (D–F) The locations of the recorded cells in the anterior PVT (D), the

middle PVT (E), and the posterior PVT (F). Red circles indicated neurons with

excitatory postsynaptic currents (EPSCs), and blue circles indicated neurons

without EPSCs. (G) The 473 nm laser-induced time-locked action potential

firing at 5 Hz (top), 10 Hz (middle), and 20 Hz (bottom) in the ChR2-expressing

neuron in the PBN. Scale bars: 100 ms, 40 mV. (H–J) The amplitude of

light-evoked EPSCs (H), the latency of EPSCs (I), and the latency jitter of

EPSCs (J) from all 34 responsive neurons in the PVT. (K) Amplitudes of

light-evoked EPSCs recorded from a PVT neuron (right panel). (L) The

light-evoked EPSC was completely blocked by 1 µM tetrodotoxin (TTX),

rescued by 100 µM 4-aminopyridine (4-AP), and blocked by 10 µM NBQX

(AMPA/kainate receptor antagonist). Scale bars: 5 ms, 10 pA. (M) Schematic

shows retroAAV2/2-hSyn-Cre injection into the PVT nucleus on

Rosa26-tdTomato mice. (N) The injection site in the PVT nucleus. Scale bar:

200 µm. (O and P) The distribution of the tdTomato positive neurons in the left

PBN (O) and the right PBN (P). (Q) The quantification of the tdTomato positive

neurons in the lateral PBN (LPBN) and the media PBN (MPBN). $n = 4$ mice.

Scale bar: 200 µm. (R–T) Double staining of tdTomato with *VgluT2* mRNA by in

situ hybridization. Scale bar: 50 µm, the scale bar in the quadrangle was 25

µm. (U) Quantification of the double-positive neurons over the total number of

tdTomato positive neurons, $n = 6$ sections from 3 mice. ***p < 0.001, data were

represented as mean ± SEM. Paired student's *t*-test for Q.

1126

The following figure supplement is available for figure 1:

Figure 1–figure supplement 1. Characterization of PVT-projecting neurons in

the PBN nucleus.

Figure 1–figure supplement 2. The distribution pattern of the PBN-PVT

glutamatergic projection.

Figure 1–figure supplement 3. The distribution pattern of collateral projection

fibers from PVT-projecting PBN neurons.

1134

Figure 2. Optogenetic activation of the PBN-PVT projection induced negative affective states. (A) The illustration showed the injection of the AAV2/9-EF1a-DIO-ChR2-mCherry virus into the PBN nucleus and the optic fiber above the PVT on the *VgluT2-ires-Cre* mice. (B and C) The virus injection sites of the left PBN (B) and the right PBN (C). Scale bar: 200 μm. (D) The projection axons from the PBN and the location of the optic fiber (rectangle) in the PVT. Scale bar: 200 μm. (E) The schematic of the open field test (OFT) with optogenetic activation via a 473 nm laser. (F and G) The example traces of the 15 minutes optogenetic manipulation OFT from an AAV2/9-EF1a-DIO-mCherry virus injected mouse (F) or an AAV2/9-EF1a-DIO-ChR2-mCherry virus injected mouse (G). (H and I) Quantification of the velocity (H) and the center time (I) in the OFT, mCherry

group: $n = 7$ mice; ChR2 group: $n = 8$ mice. (J) The illustration of the real-time place aversion test (RTPA) with optogenetic activation via a 473 nm laser. The right side was paired with the laser. (K and L) The example traces of the RTPA and post-test from the mice injected with AAV2/9-EF1a-DIO-mCherry (K) or AAV2/9-EF1a-DIO-ChR2-mCherry (L). (M) Quantification of the time spent in the laser-paired chamber in the pre-test (Pre), RTPA, and post-test (Post), mCherry group: $n = 7$ mice; ChR2 group: $n = 8$ mice. (N) Schematic timeline of cue-dependent optogenetic conditioning. (O) Conditioned-freezing responses to sound cue paired with optogenetic activation of the PBN-PVT projection during training, mCherry group: $n = 4$ mice; ChR2 group: $n = 5$ mice. (P and Q) Optogenetic activation of the projection fibers from the PBN in the PVT did not induce context-dependent fear (P) and cue-dependent fear (Q), mCherry group: $n = 4$ mice; ChR2 group: $n = 5$ mice. $^*p < 0.05$, $^{**}p < 0.01$, $^{***}p < 0.001$, all data were represented as mean ± SEM. Two-way ANOVA followed by Bonferroni test for H, I, M, and O. Unpaired student's *t*-test for P and Q.

The following figure supplement is available for figure 2:

Figure 2–figure supplement 1. The virus expression in the PBN and the optic fiber position in the PVT of *VgluT2-ires-Cre* mice injected with AAV2/9-EF1a-DIO-ChR2-mCherry virus or AAV2/9-EF1a-DIO-mCherry virus.

Figure 2–figure supplement 2. Effects of optogenetic activation of PBN-PVT projection in the OFT and the CPA.

Figure 2–video 1. Optogenetic activation of PBN-PVT projection in OFT. The 473 nm laser (20 Hz, 5 ms, 5 mW) was delivered from 00:10 to 05:10 in the video.

Figure 2–video 2. Optogenetic activation of PBN-PVT projection in RTPA. The 473 nm laser (20 Hz, 5 ms, 5 mW) was delivered when the mouse entered the laser-paired chamber and withdrew when the mouse exited the laser-paired chamber during the 10 minutes. The video was played with 4x speed.

Figure 3. Pharmacogenetic activation of the PVT-projecting PBN neurons induced anxiety-like behaviors and fear-like behaviors. (A) The illustration showed virus injection of retroAAV2/2-hSyn-Cre into the PVT nucleus and bilateral injection of AAV2/9-hSyn-DIO-hM3Dq-mCherry into the PBN nucleus. (B and C) CNO administration evokes Fos expression in AAV2/9-hSyn-DIO-hM3Dq-mCherry injected mice (B) but not in AAV2/9-EF1a-DIO-mCherry injected mice (C). Scale bar: 200 μm. (D) Percentage of co-labeled neurons in the PBN, mCherry group: n = 3 mice; hM3Dq group: n = 4 mice. (E) The illustration of the OFT test with pharmacogenetic activation. (F) Example of the OFT traces from the mice infected with AAV2/9-EF1a-DIO-mCherry or AAV2/9-hSyn-DIO-hM3Dq-mCherry. (G–I) Quantification of the center time (G), the unmoving time (H), the total distance (I) in the OFT, mCherry group: n = 7 mice; hM3Dq group: n = 8 mice. (J) Quantification of the velocity in the OFT, mCherry group: n = 7 mice; hM3Dq group: n = 5 mice. (K) Example elevated zero maze (EZM) traces from the mice infected with AAV2/9-EF1a-DIO-mCherry and AAV2/9-hSyn-DIO-hM3Dq-mCherry. (L)

Quantification of the time spent in open quadrants in the EZM test, $n = 8$ mice

per group. (M) The illustration of pharmacogenetic activation-induced fear-like

freezing behavior. (N) Pharmacogenetic activation of PVT-projecting PBN

neurons induced fear-like freezing behaviors, mCherry group: $n = 8$ mice;

hM3Dq group: $n = 7$ mice. *p < 0.05, ***p < 0.001, all data were represented as

mean ± SEM. Unpaired student's t-test for D, G, H, I, and J. Two-way ANOVA

followed by Bonferroni test for L and N.

1204

The following figure supplements are available for figure 3:

Figure 3–figure supplement 1. The virus expression in the PBN of mice

injected with AAV2/9-hSyn-DIO-hM3Dq-mCherry or

AAV2/9-EF1a-DIO-mCherry in the pharmacogenetic manipulation.

Figure 3–figure supplement 2. Pharmacogenetic activation of PVT-projecting

PBN neurons did not affect depressive-like behaviors, basal nociceptive

thresholds, formalin-induced licking behavior, or motor function.

1212

1213

1214 **Figure 4.** Optogenetic inhibition of the PBN-PVT projection reduced

aversion-like behavior and fear-like behaviors. (A) The illustration showed the bilateral injection of AAV2/9-EF1a-DIO-NpHR3.0-EYFP virus into the PBN and placement of optic fiber above the PVT on *VgluT2-ires-Cre* mice. (B and C) Examples of AAV2/9-EF1a-DIO-NpHR3.0-EYFP expression in the PVT (B) and PBN (C). The cyan rectangle represented the position of the optic fiber. Scale bar: 200 μm. (D) Schematic of 2-MT induced aversion test with optogenetic inhibition via the 589 nm laser. (E) Representative traces of the mice infected with AAV2/8-EF1a-DIO-EGFP or AAV2/9-EF1a-DIO-NpHR3.0-EYFP in two chambers. (F and G) Quantification of the time spent in the 2-MT paired chamber (F) and the total moving duration (G), $n = 7$ mice per group. *p < 0.05, **p < 0.01, (H) Representative traces of the mice infected with AAV2/8-EF1a-DIO-EGFP or AAV2/9-EF1a-DIO-NpHR3.0-EYFP in the OFT chamber. (I) Quantification of the time spent in the 2-MT zone, $n = 7$ mice per group. (J) Illustration of footshock-induced freezing behavior with optogenetic inhibition via a 589 nm laser. (K) Quantification of the freezing behavior, $n = 7$ mice per group. *p < 0.05, all data were represented as mean ± SEM. Unpaired student's *t*-test for F, G, I, and K.

The following figure supplements are available for figure 4:

Figure 4–figure supplement 1. The virus expression in the PBN and the optic fiber position in the PVT of *VgluT2-ires-Cre* mice injected with AAV2/9-EF1a-DIO-NpHR3.0-EYFP or AAV2/8-EF1a-DIO-EGFP.

Figure 4–figure supplement 2. Optogenetic inhibition of the PBN-PVT projection did not affect associative fear memory acquisition and retrieval.

Figure 4–figure supplement 3. Optogenetic inhibition of the PVT-projecting PBN neurons reduced the aversion-like behavior and fear-like freezing behavior.

1244

Figure 5. Activation of PVT$_{PBN}$ by diverse aversive stimuli. (A) The illustration showed the injection of AAV2/1-hSyn-Cre into the PBN of *Rosa26-tdTomato* mice. (B–D) The distribution of the neurons in the PVT at bregma -1.06 mm(B), bregma -1.46 mm (C), and bregma -1.70 mm (D). Scale bar: 200 µm. (E–G) Fos induced by habituation control (E), footshock (F), or 2-MT (G) co-labeled with the tdTomato positive neurons in the PVT. Scale bar: 200 µm. (H) Quantification of the co-labeled neurons, *n* = 3 mice per group. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, all data were represented as mean ± SEM, one-way ANOVA followed by Bonferroni test for H.

The following figure supplements are available for figure 5:

Figure 5–figure supplement 1. Calcium signals of PVT neurons in response to aversive stimuli.

Figure 6. Neuronal activity of the PVT neurons in response to the footshock was modulated by the PBN-PVT projection. (A) Top: Schematic showed injection of AAV2/9-EF1a-DIO-ChR2-mCherry into the PBN and placement of the optoelectrode above the PVT of *VgluT2-ires-Cre* mice. Bottom: The protocol of 10 sweeps of laser stimuli (Trial 1) and 20 sweeps of footshock stimuli without or with laser (Trial 2). (B) Firing rates (z-score) of 40 units during laser stimuli (20 Hz, 5 mW, 5 ms, 2 s). Inserted: percentages of different groups of neurons according to z-score. (C) Firing rates (z-score) of 40 units during footshock (0.5 mA, 2 s) without laser stimuli. (D) Percentage of laser-activated, footshock-activated, and both stimuli-activated units. (E-G) Rastergrams and firing rates showed the spiking activity of one PVT neuron during laser stimulus (E), footshock without laser stimulus (F), and footshock with laser stimulus (G). (H) Firing rates (z-score) of 40 units during footshock (0.5 mA, 2 s) with laser stimuli (20 Hz, 5 mW, 5 ms, 2 s). (I) Quantification of the firing rates of 40 units before and during footshock without and with laser, *n* = 40 units. **$p < 0.01$, ***$p < 0.001$, all data were represented as mean ± SEM, two-way ANOVA followed by Bonferroni test for I.

The following figure supplements are available for figure 6:

Figure 6–figure supplement 1. Dual Fos staining detecting Fos protein and *fos* mRNA induced by laser stimulation and footshock.

1281

1282

Figure 7. Activation of PVT_{PBN} neurons induced anxiety-like behaviors. (A)

The illustration showed injection of AAV2/1-hSyn-Cre into the PBN and

AAV2/9-hSyn-DIO-hM3Dq-mCherry into the PVT. (B and C) CNO

administration evoked Fos expression in AAV2/9-hSyn-DIO-hM3Dq-mCherry

injected mice (B) but not in AAV2/9-EF1a-DIO-mCherry injected mice (C).

Scale bar: 200 μm. (D) Percentage of co-labeled neurons in the PVT, $n = 4$

mice per group. (E) The illustration of the OFT test with pharmacogenetic

activation. (F) Quantification of center time in the OFT, $n = 7$ mice per group.

(G) Example of elevated zero maze (EZM) traces from the mice injected with

AAV2/9-EF1a-DIO-mCherry or AAV2/9-hSyn-DIO-hM3Dq-mCherry. (H and I)

Quantification of the time spent in open quadrants (H) and the unmoving time

in the EZM test (I), $n = 7$ mice per group. *$p < 0.05$, ***$p < 0.001$, all data were

presented as mean ± SEM. Unpaired student's t-test for D, F, H, and I.

1296

1297 The following figure supplements are available for figure 7:

1298 **Figure 7–figure supplement 1.** Distribution pattern of projection fibers of

1299 PVT_{PBN} neurons.

1300

1301

Figure 1–figure supplement 1. Characterization of PVT-projecting neurons in

the PBN nucleus. (A–O) Double staining of tdTomato signals (red) with *Tacr1*

mRNA (A–C), *Tac1* mRNA (E–G), *Pdyn* mRNA (I–K), and CGRP protein

(M–O). Scale bar: 100 µm. (D, H and L) The proportions of co-expressing

neurons of tdTomato positive neurons, *n* = 6 sections from 3 mice. Tacr1,

tachykinin 1 receptor; Tac1, tachykinin 1; Pdyn, prodynorphin; CGRP,

calcitonin gene-related peptide.

1309

1310

Figure 1–figure supplement 2. The distribution pattern of the PBN-PVT

glutamatergic projection. (A) The illustration for virus injection of

AAV2/8-EF1a-DIO-EGFP into the PBN nucleus on *VgluT2-ires-Cre* mice. (B–D)

The virus expression in the anterior (B), the middle (C), and the posterior PBN

(D). Scale bar: 200 μm. (E–H) The distribution pattern of PBN glutamatergic

projection fibers in the anterior (E and F), the middle (G), and the posterior

PVT (H). PVA, anterior paraventricular thalamus. Scale bar: 200 μm.

1318

1319

Figure 1–figure supplement 3. The distribution pattern of collateral projection fibers from PVT-projecting PBN neurons. (A) The illustration showed the injection of retroAAV2/2-hSyn-Cre into the PVT and AAV2/8-EF1a-DIO-EGFP into the PBN to label the PVT-projecting PBN neurons. (B) Examples of AAV2/8-EF1a-DIO-EGFP expression in the PBN. Scale bar: 200 µm. (C–N) The efferents from the PVT-projecting PBN neurons could be found in the MnPo (C), BNST (D), LH (E), PVN (F), PVT (G), ILN (H), DM (I), PSTh (J), VTA (K) and PAG (L) but not in the CeA (M) and VMH (N). (O) Schematic showing summary of the distribution pattern of fibers from PVT-projecting PBN neurons. MnPo, Median preoptic nucleus; BNST, bed nucleus of the stria terminalis; LH, lateral hypothalamic area; PVN, paraventricular nucleus of the hypothalamus; ILN, intralaminar thalamic nucleus; DM, dorsomedial hypothalamic nucleus;

 PSTh, parasubthalamic nucleus; VTA, ventral tegmental areas; PAG,
periaqueductal gray; CeA, central nucleus of the amygdala; VMH,
ventromedial hypothalamic nucleus.

Figure 2–figure supplement 1. The virus expression in the PBN and the optic
fiber position in the PVT of *VgluT2-ires-Cre* mice injected with
AAV2/9-EF1a-DIO-ChR2-mCherry virus or AAV2/9-EF1a-DIO-mCherry virus.
(A) Histological map showing area of ChR2 expression in the PBN at bregma
–5.20 mm in 8 mice. Scale bar: 400 μm. (B) The position of optic fiber
(rectangle) in the PVT in the AAV2/9-EF1a-DIO-ChR2-mCherry injected mice.
Scale bar: 200 μm. (C) The area of AAV2/9-EF1a-DIO-mCherry virus
expression in the PBN at bregma –5.20 mm in 7 mice. Scale bar: 400 μm. (D)

1347

1348

Figure 2–figure supplement 2. Effects of optogenetic activation of PBN-PVT

projection fibers in the OFT and the CPA. (A and B) Quantification of the

unmoving time (A) and the distance (B) in the OFT (mCherry group: n = 7 mice;

ChR2 group: n = 8 mice). (C–F) Quantification of the velocity (C), the

unmoving time (D), the center time (E), the distance (F) and the number of

jumps (G) during the 5–10 minutes laser ON period in the OFT test (mCherry

group: n = 7 mice; ChR2 group: n = 8 mice). (H) Protocol for the prolonged

conditioned place aversion (CPA). (I) Photostimulation of PBN-PVT projection

did not induce CPA (n = 7 mice per group). *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$,

all data were represented as mean ± SEM. Two-way ANOVA followed by

Bonferroni test for A, B, C, D, E, F, G, and I.

1360

Figure 3–figure supplement 1. The virus expression in the PBN of mice injected with AAV2/9-hSyn-DIO-hM3Dq-mCherry or AAV2/9-EF1a-DIO-mCherry in the pharmacogenetic manipulation. (A) Representative histological images of hM3Dq expression in an AAV2/9-hSyn-DIO-hM3Dq-mCherry injected mouse at brain level from bregma –5.02 mm to bregma –5.34 mm. Scale bar: 1 mm. (B) Depiction of virus

infection area according to the histological images in (A). (C) Superimposed

depiction of virus transduction from 8 mice. (D) Representative histological

images of mCherry expression in an AAV2/9-EF1a-DIO-mCherry injected mice

at brain level from bregma –5.02 mm to bregma –5.34 mm. Scale bar: 1 mm.

(E) Depiction of virus infection area according to the histological images in (D).

(F) Superimposed depiction of virus transduction from 8 mice.

1374

1375

Figure 3–figure supplement 2. Pharmacogenetic activation of PVT-projecting

PBN neurons did not affect depressive-like behaviors, basal nociceptive

thresholds, formalin-induced licking behavior, or motor function. (A) Immobility

time in the tail suspension test (TST), $n = 7$ mice per group. (B) Immobility time

in the forced swimming test (FST), mCherry group: $n = 7$ mice; hM3Dq group:

$n = 8$ mice. (C and D) Effects of pharmacogenetic activation of PVT-projecting

PBN neurons on the nociceptive response tested by von Frey (C) and

Hargreaves (D), $n = 8$ mice per group. (E and F) Duration of licking behaviors

in the formalin-induced inflammatory pain test, mCherry group: $n = 7$ mice;

hM3Dq group: $n = 8$ mice. Phase I: 0–10 minutes, Phase II: 10–60 minutes. (G)

The latency to fall in the rotarod test, $n = 8$ mice per group. Data were

represented as mean ± SEM. Unpaired student's t-test for A, B and G.

Two-way ANOVA followed by Bonferroni test for C, D, E, and F.

1389

1390

1391

Figure 4–figure supplement 1. The virus expression in the PBN and the optic

fiber position in the PVT of *VgluT2-ires-Cre* mice injected with

AAV2/9-EF1a-DIO-NpHR3.0-EYFP or AAV2/8-EF1a-DIO-EGFP. (A)

Histological map showing area of NpHR3.0 expression in the PBN at bregma

–5.20 mm in 7 mice. Scale bar: 400 μm. (B) The position of optic fiber

(rectangle) in the PVT in the AAV2/9-EF1a-DIO-NpHR3.0-EYFP injected mice.

Scale bar: 200 μm. (C) The area of AAV2/8-EF1a-DIO-EGFP expression in the

PBN at bregma –5.20 mm in 7 mice. Scale bar: 400 μm. (D) Position of the

optic fiber tip from 7 mice injected with AAV2/9-EF1a-DIO-mCherry. Scale bar:

200 μm.

1402

1403

Figure 4–figure supplement 2. Optogenetic inhibition of the PBN-PVT

projection did not affect associative fear memory acquisition and retrieval. (A)

The protocol of auditory fear conditioning experiments with optogenetic

inhibition of the PBN-PVT projection. (B–D) Quantification of freezing levels

during condition trials (B), contextual test (C), and cue test (D), $n = 7$ mice per

group. The yellow box indicated optogenetic inhibition. All data were

represented as mean ± SEM, two-way ANOVA followed by Bonferroni test for

B, C and D.

1412

1413

Figure 4–figure supplement 3. Optogenetic inhibition of the PVT-projecting

PBN neurons reduced the aversion-like behavior and fear-like freezing

behavior. (A) The illustration showed virus injection of retroAAV2/2-hSyn-Cre

into the PVT, bilateral injection of AAV2/9-EF1a-DIO-NpHR3.0-EYFP into the

PBN, and bilateral placement of optic fiber above the PBN on WT mice. (B and

C) Examples of AAV2/9-EF1a-DIO-NpHR3.0-EYFP (B) and

AAV2/8-EF1a-DIO-EGFP (C) expression in the PBN, the rectangle

represented the position of the optic fiber. Scale bar: 200 μm. (D) Schematic of

2-MT induced aversion test with optogenetic inhibition via a 589 nm laser. (E)

Representative traces of the mice infected with AAV2/8-EF1a-DIO-EGFP or

AAV2/9-EF1a-DIO-NpHR3.0-EYFP in the chamber. (F and G) Quantification of

the time spent in the 2-MT paired chamber (F) and the moving duration (G), n

= 5 mice per group. (H) Schematic of 2-MT induced fear-like freezing behavior

with optogenetic inhibition via a 589 nm laser. (I) Quantification of the freezing

behavior, n = 5 mice per group. (J) Illustration of footshock-induced freezing

behavior with optogenetic inhibition via a 589 nm laser. (K) Quantification of

the freezing behavior, n = 5 mice per group.*p < 0.05, **p < 0.01, all data were

1431 represented as mean ± SEM. Unpaired student's *t*-test for F, G, I, and K.

1432

1433

Figure 5–figure supplement 1. Calcium signals of PVT neurons in response

to aversive stimuli. (A) Schematic showed injection of AAV2/8-hSyn-GCaMP6s

into the PVT and placement of the optic fiber above the PVT. (B)

Representative of GCaMP6s expression and the position of optic fiber in the

PVT. Scale bar: 400 µm. (C and D) The calcium signal of the PVT neurons (C)

and the quantification of the average Ca^{2+} signal before and after footshock

(D). The black bar represented the baseline period (B.s, -1 to -0.5 s), and the

red bar represented the post-stimulus period (Post, 0.5 to 1 s), *n* = 5 mice. (E

and F) The calcium signal of the PVT neurons (E) and the quantification of

average Ca^{2+} signal before and after air puff (F), *n* = 5 mice. **p < 0.01, all data

were represented as mean ± SEM. Paired student's *t*-test for D and F.

1445

1446

Figure 6–figure supplement 1. Dual Fos staining detecting Fos protein and

fos mRNA induced by laser stimulation and footshock. (A) Top: Schematic

showed injection of AAV2/9-EF1a-DIO-ChR2-mCherry into the PBN and

placement of the optic fiber above the PVT of *VgluT2-ires-Cre* mice. Bottom:

Time windows containing laser (20 Hz, 5 mW, 5ms) and shock stimuli (0.5 mA,

1 s, 30 times), separated by 60 minutes of the rest period. (B–D) Example of

Fos protein and *fos* mRNA expression in the PVT. Red fluorescence

represents Fos protein induced by laser stimulus, and green fluorescence

indicates *fos* mRNA detected by *in situ* hybridizations. Scale bar: 100 μm. (E)

The proportion of co-expression neurons over Fos protein-expressing cells

and *fos* mRNA-expressing cells, $n = 5$ mice.

1458

1459

Figure 7–figure supplement 1. Distribution pattern of projection fibers of

PVT_{PBN} neurons. (A) The illustration showed injection of AAV2/1-hSyn-Cre into

the PBN and AAV2/8-EF1a-DIO-EGFP into the PVT of *Rosa26-tdTomato* mice.

(B) The representative image of EGFP and tdTomato-transduced neurons in

the PVT. Scale bar: 200 μm. (C–G) Distribution patterns of EGFP fibers in the

PFC (C), CI (D), NAc (E), BNST (F), and CeA (G). PFC, prefrontal cortex; IC,

insular cortex; CI, claustrum; NAc, nucleus accumbens core; BNST, bed

nucleus of the stria terminalis; CeA, central nucleus of the amygdala. Scale bar:

200 μm. (H) Quantification of the fiber intensity in these brain regions, $n = 3$

mice.