A single synonymous nucleotide change impacts the male-killing phenotype of prophage WO

gene *wmk*

Jessamyn I. Perlmutter[a,b,c#], Jane E. Meyers[a,c], & Seth R. Bordenstein[a,c,d,e#]

[a]Department of Biological Sciences, Vanderbilt University, Nashville, Tennessee, USA

[b]Department of Molecular Biosciences, University of Kansas, Lawrence, Kansas, USA

[c]Vanderbilt Microbiome Innovation Center, Vanderbilt University, Nashville, Tennessee, USA

[d]Department of Pathology, Microbiology, and Immunology, Vanderbilt University, Nashville,

Tennessee, USA

[e]Vanderbilt Institute for Infection, Immunology, and Inflammation, Vanderbilt University,

Nashville, Tennessee, USA

#Address correspondence to jessamyn.perlmutter@outlook.com and

s.bordenstein@vanderbilt.edu

Abstract

Wolbachia are the most widespread bacterial endosymbionts in animals. Within arthropods, these maternally-transmitted bacteria can selfishly hijack host reproductive processes to increase the relative fitness of their transmitting females. One such form of reproductive parasitism called male killing, or the selective killing of infected males, is recapitulated to degrees by transgenic expression of the *WO-mediated killing* (*wmk*) gene. Here, we characterize the genotype-phenotype landscape of *wmk*-induced male killing in *D. melanogaster* using transgenic expression. While phylogenetically distant *wmk* homologs induce no sex-ratio bias, closely-related homologs exhibit complex phenotypes spanning no death, male death, or death of all hosts. We demonstrate that alternative start codons, synonymous codons, and notably a single synonymous nucleotide in *wmk* can ablate killing. These findings reveal previously unrecognized features of transgenic *wmk*-induced killing and establish new hypotheses for the impacts of post-transcriptional processes in male killing variation. We conclude that synonymous sequence changes are not necessarily silent in nested endosymbiotic interactions with life-or-death consequences.

Introduction

Wolbachia are maternally-transmitted, obligate intracellular bacteria primarily residing in the cells of germline tissues in many arthropod species worldwide[1,2]. To facilitate their spread through the host matriline, these bacteria hijack sex ratios, sex determination systems, and embryonic viability to cause various reproductive parasitism phenotypes[3-6]. One such phenotype is male killing, whereby sons of infected females are selectively killed[4,7,8]. Male killing selfishly drives the spread of the bacteria when, for instance, brothers and sisters compete for limited resources, and male lethality yields females (the transmitters) a relative fitness benefit due to reduced competition with fewer siblings[9-12]. This form of reproductive parasitism can have profound impacts on host behavior and evolution, including a sex role reversal[13], host nuclear genome changes to resist the phenotype[14,15], and potentially host population extinction[16]. Population modelling also suggests male killing may be deployed as a control method to crash the population size of arthropod pests and disease vectors[17].

Many lines of evidence suggest a complex relationship between male killing genotype and phenotype. For example, male killing can be suppressed by host background[14,15,18,19], where male hosts exhibit resistance to the phenotype even in the presence Wolbachia. In addition, Wolbachia that do not cause male killing in one host species can cause male killing in a second host upon introgression or transfer to a naïve host[18,20]. Furthermore, some strains induce death at different host developmental stages (embryo vs. larvae) or across different sex determination systems (ZW lepidopterans, XY dipterans, XO arachnids), and the number of surviving males may vary widely[21-23]. Such findings specify that the expression of male killing, and particularly the genotype-phenotype relationship in these symbioses, is more complex than simply the presence or absence of a male-killing gene.

We recently identified a male-killing candidate gene, *wmk* (*WO-mediated killing*), in the

*w*Mel *Wolbachia* strain of *Drosophila melanogaster* based on comparative genomic, transgenic,

and cytological approaches[24,25]. It is a putative transcription factor with two predicted helix-turn-

helix (HTH) DNA-binding domains, and transgenic expression in *D. melanogaster* embryos

recapitulates many cytological and molecular aspects of male killing including accumulation of

DNA damage in male embryos that overlaps with sites of dosage compensation activity[21,25,26]. The

wmk gene and two cytoplasmic incompatibility factor (*cif*) genes that underlie cytoplasmic

incompatibility (a parasitism phenotype whereby offspring die in crosses between infected males

and uninfected females) occur in the eukaryotic association module (EAM) of prophage WO[24,25,27],

which refers to the phage WO genome that is inserted into the bacterial chromosome. The EAM

is common in WO phages across several *Wolbachia* strains[28,29] and is rich in genes that are

homologous to eukaryotic genes or annotated with eukaryotic functions[24]. As such, the expression

of reproductive parasitism genes from the EAM and tripartite interactions between phage WO,

Wolbachia, and eukaryotic hosts are central to *Wolbachia*'s ability to interact with and modify

host reproduction.

The discovery of *wmk* has now enabled investigation of the impacts of genetic variation on

the transgenic male-killing phenotype. Although the role of *wmk* in male killing has not yet been

assessed in nature, evidence based on native *wmk* genotypes and transgenic *wmk* expression in

flies suggest refined intricacy to the interactions between genotype and phenotype. For example,

the native *Wolbachia wmk* homologs in three closely-related strains (*w*Mel of *D. melanogaster*,

*w*Rec of *D. recens*, and *w*Suzi of *D. suzukii*) have nearly identical nucleotide sequences that vary

by one to three SNPs[25], and yet these strains induce different forms of reproductive parasitism.

*w*Mel and *w*Rec induce cytoplasmic incompatibility, *w*Rec also causes male killing in some strains

of a sister species, and *w*Suzi is not known to induce any reproductive phenotype[18,30,31]. Previous

transgenic testing also demonstrated that addition of nine additional amino acids to the N-terminal

region of the Wmk protein at the site of a putative alternative start codon ablates the phenotype[32].

Therefore, the Wmk N-terminus is particularly sensitive to genetic alterations, which is notable

since small modifications such as protein tags in this location are often common additions to

proteins that do not typically interfere with function. Furthermore, *D. melanogaster* does not

harbor *Wolbachia* known to cause male killing, though transgenic *wmk* expression recapitulates

the phenotype in this host[25], and the *w*Mel strain is most closely related to the *w*Rec male-killing

strain[33]. In addition, some *D. melanogaster* contain male-killing *Spiroplasma* symbionts whose

plasmid gene *Spaid* can also transgenically recapitulate male killing[34]. In summary, the evidence

for *Wolbachia* and *wmk* transgenic male killing suggests an intricate and multifaceted genotype-

phenotype landscape.

Building on this background, several key questions emerge: Do closely-related or distantly-

related homologs induce male killing? How sensitive is the *wmk* phenotype to small or large

genetic changes? And how adapted are *wmk* homologs to their arthropod hosts? Here, we evaluate

codon-optimized *wmk* homologs that span a spectrum of genetic divergence (Fig. 1), including

homologs (i) from distantly-related hosts, (ii) with putative upstream alternative start codons, (iii)

with a single amino acid change, or (iv) with one or more synonymous codon changes, all in the

D. melanogaster host. In this way, we aimed to investigate how a variety of genetic alterations to

wmk, from a single synonymous nucleotide to the gene level, affect the phenotype. In particular,

we hypothesized that divergent strains would not induce the phenotype due to co-adaptation with

a distantly-related host. We also anticipated that the alternative start codons would inhibit function

based on previous results[32]. In contrast, we hypothesized that neither a single amino acid change

nor synonymous codon changes would alter the male-killing phenotype. We report that while

distant homologs do not cause a male-killing phenotype in this host, single amino acid and

synonymous nucleotide changes remarkably do alter the phenotype. Thus, *wmk* male killing is

sensitive to the full spectrum of genetic alterations at a fine-scale level not previously recognized.

Notably, synonymous sequence changes and post-transcriptional processes appear to play a role

in controlling the genotype-phenotype relationships that underpin *wmk* male killing.

Figure 1. Overview of experimental design. To investigate the genotype-phenotype landscape,
we transgenically expressed *wmk* homologs with varying degrees of genetic changes. These
sequences are codon-optimized based on different codon biases due to different tRNA abundances
in the divergent bacterial source and eukaryotic destination species[35,36]. Transgenic *wmk* in
Drosophila melanogaster embryos results in three different phenotypes: no killing, male killing,
and killing of males and females. Compared to *w*Mel *wmk*, these transgenes were either divergent
homologs from other *Wolbachia* strains, a homolog with a single amino acid change, homologs
with an additional 9 codons at the 5' ends of the genes starting at an alternative upstream start
codon, or variants with a single synonymous codon or nucleotide difference. These genotypes
resulted in varying degrees of RNA sequence- and amino acid-level changes. Created with
BioRender.com.
122

Results

***Closely-related homologs of* wmk.** We tested homologs from strains related to *w*Mel that

occur in several groups of species in the *Drosophila* genus, along with two from Order Lepidoptera

(Fig. 1- figure supplement 1). The phylogenetic clades of both the native *Wolbachia* genes and

their transgenes are similar to those of the host species. Roughly divided, the transgenes group into

two clusters that offer a range of genetic divergence for evaluating genotype-phenotype

relationships (Fig. 1- figure supplement 1) - those distantly related to the *w*Mel *wmk* transgene

(less than 90% codon-optimized nucleotide sequence similarity to transgenic *w*Mel *wmk*) and those

closely related (greater than 90% identity, from *w*Rec, *w*Suzi, and an HA-tagged *w*Mel homolog).

Host Species

Native *wmk*   **Transgene *wmk***

Transgene Nucleotide Alignment

Figure 1- figure supplement 1. Homologs of *wmk* tested in this study include variation in
native gene and transgene sequence identity as well as host species. (Top) Bayesian nucleotide
phylogeny of insect hosts based on 652 bp of the cytochrome oxidase subunit 1 (COI) gene from
D. melanogaster, *H. bolina*, *D. bifasciata*, *C. cautella*, *D. innubila*, *D. borealis*, *D. suzukii*, and *D.*
recens. Branch labels and scale bar indicate posterior probability. Species names are colored either
by group within the *Drosophila* genus or by Order Lepidoptera. Underlines indicate hosts in which
male-killing strains have been reported. Colored circles indicate host species group. (Middle)
Bayesian nucleotide phylogenies of native (middle left, non-transgenic) or transgene (middle right)
wmk sequences based on 690 or 686 bp, respectively. Label colors reflect groups from host
phylogeny for comparison. Colored circles indicate host species group. Branch labels and scale
bar indicate posterior probability. *w*Bif *wmk* branches distantly due to a highly divergent sequence.
*w*Bor and *w*Inn *wmk* share a branch because they share the same transgenic sequence. (Bottom)
Nucleotide alignment of transgenes tested in this study as compared to the previously tested *w*Mel
wmk, with the regions encompassing the Helix-turn-helix (HTH) protein domains marked with
black lines. Black ticks indicate sequence differences with the *w*Mel *wmk* reference strain, light
grey indicates sequence matches to the reference sequence, and white indicates an indel in at least
one strain. In parentheses under strain names, percentages refer to nucleotide similarity compared
to the *w*Mel *wmk* transgene. MK indicates a male killing strain in its native host, transgenic MK
indicates the ability to induce a biased sex ratio transgenically only, and MK in sister species
indicates the ability to kill males in a sister host species but not in the native host species.
Alignment excludes the HA tag in the tagged strain, which is located between the two HTH
domains.

The closely-related *wmk* homologs to *w*Mel include those from *Wolbachia* strains (i) *w*Suzi

of the fruit pest species *D. suzukii* that has no confirmed reproductive phenotype but notably occurs

in populations with female-biased sex ratios[30,37] and (ii) *w*Rec of *D. recens* that kills males when

introgressed into its sister species, *D. subquinaria*[18]. The natural *w*Suzi *wmk* homolog has one

synonymous single nucleotide polymorphism (SNP) compared to the *w*Mel *wmk* reference, thus

yielding the same amino acid sequence. The natural *w*Rec *wmk* homolog has three SNPs, one of

which is non-synonymous relative to *w*Mel *wmk*, located in the first HTH DNA-binding domain

(Fig. 2A). These transgenes were codon optimized for expression in *D. melanogaster*. We

hypothesized that both transgenes would induce a biased sex ratio comparable to *wmk* when

expressed in *D. melanogaster*. However, transgenic expression of both homologs unexpectedly

resulted in death of all expressing flies, both male and female (Fig. 2B). In addition, we

simultaneously tested a transgene of *w*Mel *wmk* with an internal 3X HA tag epitope in the linker

region between the two HTH DNA-binding domains. This transgene, HA-*wmk*, exhibited a sex

ratio bias comparable to *w*Mel *wmk*, as expected (Fig. 2B). Therefore, two transgenes of *w*Mel

wmk (one with a tag) resulted in the biased sex ratio previously reported[25], while two closely-

related transgenes from strains that at least associate with female-biased host sex ratios yielded an

all-killing phenotype.

Figure 2. Transgenic expression of closely-related *wmk* homologs causes male killing and all killing phenotypes in *D. melanogaster*. (A) Schematic of *w*Mel, *w*Suzi, and *w*Rec *wmk* native nucleotide sequences. The blue tick mark indicates a non-synonymous nucleotide difference. Black tick marks indicate synonymous nucleotide changes. Numbers indicate nucleotide position across the entire 912 nucleotide sequence. (B) Sex ratios of adult flies are shown for expressing (*Act5c*-Gal4) and non-expressing (CyO) embryonic offspring. Each sample point represents the adult offspring (N=50-157, mean 86) produced by a replicate family of ten mothers and two fathers, with expressing and non-expressing flies of a given genotype being siblings. Bars represent the mean sex ratio. Statistics are based on a Kruskal-Wallis, one-way ANOVA followed by Dunn's correction across either expressing or non-expressing flies. *w*Rec and *w*Suzi *wmk* have no points in the expressing category due to death of most or all males and females. HA-*wmk* contains a 3X HA tag in the linker region between the two helix-turn-helix domains. This experiment was performed twice. Data and statistical outputs are available in Figure 2 - Source Data 1 and Figure 2- Source Data 2, respectively. (C) Gene expression in embryos 4-5 h AED of each indicated *wmk* transgene from (B), relative to *Drosophila* housekeeping gene, *rp49*. There is no significant difference in expression based on a Kruskal-Wallis one-way ANOVA followed by Dunn's correction. Data and statistical outputs are available in Figure 2 - Source Data 3 and Figure 2- Source Data 4, respectively. (D) Gene expression in embryos 4-5 h AED of the host *msl-2* dosage compensation gene relative to *rp49* under simultaneous expression of the indicated transgene. There is no significant difference in expression based on a Kruskal-Wallis one-way ANOVA followed by Dunn's correction. Data and statistical outputs are available in Figure 2 - Source Data 5 and Figure 2- Source Data 6, respectively. (E) Predicted RNA secondary structures of native *w*Mel *wmk* and several transgene strains. Black arrows point to the location of the start codon within each structure.

Gene expression similarities. To assess if variation in gene expression underpins the

phenotypic variation, we measured the transcript levels of the transgenes in embryos 4-5 h AED

(after egg deposition) when *wmk* kills males[25]. Gene expression levels are not significantly

different across *w*Suzi, *w*Mel, *w*Rec, or HA tag *w*Mel *wmk* transgenes, indicating that transcript

levels do not account for the phenotypic differences (Fig. 2C). An alternative hypothesis is that

transgenic expression of the *w*Suzi and *w*Rec homologs impacts native expression of a host gene

in *D. melanogaster* that causes male and female lethality. For instance, transgenic expression of

the DNA-binding dosage compensation gene, *male-specific lethal 2* (*msl-2*), can induce total

lethality with male-killing *Spiroplasma*[38]. Based on this, we quantified *msl-2* transcript levels in

embryos expressing *w*Mel *wmk* (sex ratio bias), *w*Suzi *wmk* (all expressing hosts die), and *w*Bif

wmk (no killing or sex ratio bias) and found that *msl-2* levels were comparable across all genotypes

and phenotypes (Fig. 2D).

RNA structural model variation. We next considered mRNA secondary structural

differences of the transgene transcripts as a factor explaining the observed phenotypic variation.

Modelling RNA structures showed they were substantially different, even in the case of few to no

amino acid level changes from one sequence to another. Notably, some RNA structural features

grouped by phenotype, such as the location of the start codon (Fig. 2E). Two transgenic transcripts,

*w*Mel *wmk* and HA-*wmk*, cause a sex ratio bias and have start codons in the middle of the structure,

while the other transgenes, *w*Suzi *wmk* and *w*Rec *wmk*, that kill all flies have start codons on outer

loops or hairpins. Although caution is warranted with this predicted structural analysis, mRNA

secondary structure could explain some phenotypic outcomes of closely-related *wmk* homologs.

Native *w*Mel *wmk* also has a start codon on an outer loop but does not induce an "all killing"

phenotype like the *w*Suzi and *w*Rec *wmk* transgenes, which may be due to lower expression of

native genes[25], and/or other structural differences.

***Distantly-related homologs of* wmk.** To determine if *wmk* homologs from distantly-related

strains induce a biased sex ratio in *D. melanogaster*, we transgenically expressed four codon-

optimized homologs from known male-killing strains of *Wolbachia*: the *w*Bol1b strain from

Hypolimnas bolina butterflies[39], the *w*Bif strain from *Drosophila bifasciata* flies[21], the *w*Caub

strain from *Cadra cautella* moths[22], and the *w*Inn strain from *D. innubila* flies[40] (same gene

sequence as the *w*Bor male-killing strain from *D. borealis* flies[41]) (Fig. 1 – figure supplement 1).

While transgene expression of *w*Mel *wmk* induces a biased sex ratio (~ one third of expressing

males die), none of the more distantly-related *w*Bol1b, *w*Bif, *w*Caub, or *w*Inn/*w*Bor *wmk* homologs

yield a biased sex ratio, demonstrating that they do not recapitulate male killing when

transgenically expressed in *D. melanogaster* under the conditions tested (Fig. 3).

Figure 3. Divergent homologs of *wmk* from male-killing strains do not induce a biased sex ratio in *D. melanogaster*. Sex ratios of adult flies are shown from either expressing (*Act5c*-Gal4) or non-expressing (CyO) offspring. WT refers to the background insertion line and Control gene refers to the WD0034 control transgene that induces no sex ratio bias. Each sample point represents the adult offspring (N=50-132, mean 69) produced by a replicate family of ten mothers and two fathers, with expressing and non-expressing flies of a given genotype being siblings. Bars represent the mean sex ratio. Statistics are based on a Kruskal-Wallis one-way ANOVA followed by Dunn's correction across either expressing or non-expressing flies. This experiment was performed twice. Data and statistical outputs are available in Figure 3 - Source Data 1 and Figure 3- Source Data 2, respectively.

Alternative start codon variation and male killing. Relevant to the studies here, we

previously provided evidence that some strains contain alternative start codons upstream of the

annotated start for *wmk* and these upstream regions are expressed in the *w*Mel strain[32]. When *w*Mel

wmk was transgenically expressed with the most likely upstream start codon, the phenotype was

lost, and no biased sex ratio resulted. We also showed that some non-male-killing strains tended

to have more of the alternative start codons[32]. To determine if *w*Rec and *w*Suzi *wmk* transgene

phenotypes are similarly sensitive to transcript changes, we expressed them with upstream codons

that are native to each of their genetic sequences. As previously observed with other homologs,

they lost their killing phenotype with only nine amino acids added to the 5' end of the gene, despite

being smaller than many commonly used protein tags (Fig. 2- figure supplement 1). All expressing

flies survived with no sex ratio bias. Returning to the RNA structure models, we find that simply

adding the corresponding nucleotides at the 5' end of each homolog resulted in several predicted

differences in RNA secondary structure for each transgene compared to the structures without the

additional 5' nucleotides (Fig. 2- figure supplement 1). This includes additional loops and different

predicted placement of the start codon.

Figure 2- figure supplement 1. *w*Rec and *w*Suzi transgenes expressed with an alternative start codon lose their transgenic phenotypes. (Top) Sex ratios of adult flies are shown for expressing (*Act5c*-Gal4) and non-expressing (CyO) offspring. Each sample point represents the adult offspring (N=50-120, mean 84) produced by a replicate family of ten mothers and two fathers, with expressing and non-expressing flies of a given genotype being siblings. Bars represent the mean sex ratio. Statistics are based on a Kruskal-Wallis one-way ANOVA followed by Dunn's

correction across either expressing or non-expressing flies. This experiment was performed twice.
Data and statistical outputs are available in Figure 2 - Source Data 7 and Figure 2- Source Data 8,
respectively. (Bottom) Predicted RNA secondary structures of the *w*Rec and *w*Suzi *wmk*
transgenes with the additional 5' sequence exhibit slight structural changes compared to their non-
lengthened counterparts (included again from Fig. 2 for ease of comparison). Black arrows point
to the location of the start codon within each structure.
274
Silent site variation and male-killing. Finally, to identify particular nucleotides that may

account for phenotypic variation among the homologs, we aligned the sequences of the four

transgenes in Fig. 2 and investigated codons that clustered by phenotype (sex ratio bias for *w*Mel

wmk and HA-*wmk*, or all killing for *w*Rec and *w*Suzi *wmk*). Across the length of the genes (and

excluding the HA tag), there were only two codon differences: one at the sixteenth amino acid

position and another near the end of the gene. As previous work demonstrated that changes at the

5' end of this gene affect phenotype[32] and since approximately the first ten codons in model

prokaryotic genes are known to substantially affect mRNA structure and resulting translation

rate[42], we focused on the earlier codon at site 16. This codon, which codes for Serine in all

homologs, segregates among sequences by phenotype. The HA-*wmk* and *w*Mel *wmk* transgenes,

which recapitulate male killing, have a TCG codon, while the all-killing transgenes *w*Suzi and

*w*Rec *wmk* have TCC and AGC, respectively (Fig. 4A).

Figure 4. Synonymous nucleotide changes in the 16th codon position of *wmk* alters resulting phenotype. (A) Sequence alignment of transgenic *wmk* homologs. The codon farthest on the left is the fourth codon in the sequence, and the highlighted codon is the 16th, with the farthest right representing the 23rd codon, and ellipses indicating codons continuing on either side. The red box outlines where the genotypes cluster by phenotype. The "HA-*wmk*" and "*w*Mel *wmk*" genotypes share the same codon in this position, and both induce male-specific death. The "*w*Rec *wmk*" and "*w*Suzi *wmk*" genotypes both exhibit different codons from the previous two and exhibit an all-killing phenotype. Colors correlate with amino acid identity. (B) Sequence alignment of transgenes with either the *w*Mel *wmk* sequence made anew (*w*Mel *wmk* new), or with the 16th codon (red box) replaced with the synonymous codons from the *w*Rec and *w*Suzi *wmk* transgenes. The colors and symbols reflect those in (A). (C) Sex ratios of adult transgenic flies are shown for expressing (*Act5c*-Gal4) and non-expressing (CyO) offspring that include the original transgene *w*Mel *wmk* strain used in previous figures, along with the newly created identical *w*Mel *wmk* (new) transgene

and the additional transgenes with the single codon swapped out for the indicated codons noted in
(A). *w*Suzi codon and *w*Rec codon refer to the strains that have the same sequence as the *w*Mel
wmk, but with one or three silent sites changed in the single codon at the 16th amino acid position.
Each sample point represents the adult offspring (N=50-161, mean 73) produced by a replicate
family of ten mothers and two fathers, with expressing and non-expressing flies of a given
genotype being siblings. Bars represent the mean sex ratio. Statistics are based on a Kruskal-Wallis
one-way ANOVA followed by Dunn's correction across either expressing or non-expressing flies.
This experiment was performed twice. Data and statistical outputs are available in Figure 4 -
Source Data 1 and Figure 4- Source Data 2, respectively. (D) Gene expression in embryos 4-5 h
AED denotes expression of each transgene relative to that of *rp49*. There is no significant
difference in expression based on a Kruskal-Wallis one-way ANOVA followed by Dunn's
correction. Data and statistical outputs are available in Figure 4 - Source Data 3 and Figure 4-
Source Data 4, respectively. (E) Predicted RNA secondary structures of the *w*Mel *wmk* transcript
compared to both of the *w*Rec or *w*Suzi codon transgenes exhibiting slight structural differences.
The structure for transgene *w*Mel *wmk* is included again from Fig. 3 for ease of comparison. Black
arrows point to the location of the start codon within each structure. Black circles highlight a key
area of difference between the structures, with a stem absent in the *w*Suzi codon strain, and
different base pair match probabilities calculated for each as indicated by color (blue to red, low
to high probability). Within the black circle, the *w*Suzi codon transgene structure is missing a
predicted stem that the others have. The stem in the *w*Rec codon line, while present, has a weaker
prediction as noted by the cooler colors, so there may be structural differences compared to the
*w*Mel *wmk* model.
323
To functionally test if the silent changes in this Serine codon accounted for the phenotype

differences, we generated three new transgenes (Fig. 4B): (i) *w*Mel *wmk* control with no changes

compared to the previously tested transgene (TCG codon, labelled "*w*Mel *wmk* (new)"), (ii) *w*Mel

wmk with three nucleotide changes in the codon that reconstitutes the AGC present in the *w*Rec

transgene line (labelled "*w*Rec codon"), and (iii) *w*Mel *wmk* with one nucleotide change in the

codon that reconstitutes the TCC present in the *w*Suzi transgene line (labelled "*w*Suzi codon").

When these three otherwise identical genes were transgenically expressed, the wMel *wmk* (new)

transgene with no changes caused a biased sex ratio as expected; however, and remarkably,

expression of transgenes with two different Serine codons ablated the phenotype and resulted in a

non-biased sex ratio with normal numbers of expressing flies (Fig. 4C). This ablation occurs even

though transcript levels remain similar across all transgenes and despite sequencing confirmation

of the single codon differences (Fig. 4D). Thus, a minimum of one single synonymous site change

in the 5' region was sufficient to alter the sex ratio phenotype. However, while the nucleotide

changes in the codon changed the phenotype, they did not recapitulate the all-killing phenotypes

of their corresponding homologs. The predicted RNA secondary structures from the transgenes

with the single codon changes are similar to the original *w*Mel *wmk* transgene, but they differ in

some aspects such as presence or absence of stems and loops and the probability score of the base

pair match as indicated by color (scale of red to blue, warmer colors indicate high probability,

cooler colors indicate low probability). (Fig. 4E).

Discussion

345   Linking mutations to function is crucial in resolving the evolutionary dynamics of

adaptations, especially when the functions emerge in a nested system of tripartite phage-bacteria-

animal interactions. In a simple case, the direct impact of genetic divergence is functional

divergence with increasing numbers of mutations leading to increased functional divergence. Here,

we investigated evolutionary and molecular hypotheses related to the genotype-phenotype

relationships between the prophage *WO-mediated killing* gene, *wmk*[25,32], and male killing in *D.*

melanogaster. Sequences among *wmk* homologs in nature can differ across divergent hosts,

leading to the hypothesis that highly divergent alleles are functionally fine-tuned. Using a variety

of sequences, we report three key results: (i) mutations even at the synonymous codon and single

nucleotide levels can alter the male-killing phenotype, (ii) phenotype, genotype, and RNA

structural variation exhibit some correlation with each other, and (iii) distantly-related homologs

do not induce a male-killing phenotype in transgenic *D. melanogaster*, while some closely-related

homologs from *Drosophila* species/populations with female-biased sex ratios induce all-killing

phenotypes. We discuss how these findings expand and support mechanistic models of *wmk* male

killing, emphasize important implications for transgenic assays in endosymbiont studies, and relate

these findings to aspects of male killing such as phenotype switching and host resistance.

The most surprising and unanticipated result was that transgene expression of highly

similar homologs and even single synonymous site changes alter phenotype, and they make the

difference between life and death for some males. We tested *wmk* homologs with high native

sequence identity compared to native *w*Mel *wmk*: *w*Rec *wmk* from the mushroom-feeding *D.*

recens (native sequence has 2 synonymous and 1 non-synonymous nucleotide differences) and

*w*Suzi *wmk* from the fruit crop pest *D. suzukii* (native sequence has 1 synonymous nucleotide

difference). Although we anticipated similar results to *w*Mel *wmk* expression, we found that

transgenic expression of these genes killed all flies (Fig. 2), even though the *w*Suzi *wmk* transgene

produces an identical protein to transgenic *w*Mel Wmk. Of particular note, transgenic *w*Suzi *wmk*

(all killing) and the original transgenic *w*Mel *wmk* sequence (male killing) produce proteins of the

exact same amino acid sequence, but they shared only a 91% sequence identity in the codon-

optimized transgene, largely due to an updated codon optimization algorithm. Codon optimization

is the norm in transgenic symbiosis research under the common assumption that synonymous

codons are functionally redundant. However, codon optimization algorithms often choose a codon

based on factors including codon adaptation, mRNA folding, regulatory motifs, nucleotide bias,

or codon correlations and biases[35]. With even a few different codons input into the algorithm as

well as algorithm updates over the years, the tested transgenes had different nucleotide sequences.

Based on the aforementioned results, we next sought to assess the sensitivity of male killing

to *wmk* transcriptional and post-transcriptional changes and asked if one synonymous codon or site

change was sufficient to alter phenotype. We created three transgenic lines, each with different

DNA sequences coding for the same Serine at position 16. With only these minor changes that

encode an identical protein sequence to *w*Mel Wmk, the transgenic male-killing phenotype was

ablated (Fig. 4). Thus, one single codon and even a single nucleotide, remarkably determined male

viability. Importantly, this key result implies at the molecular level that endosymbiont phenotypes

of reproductive parasites may not simply be governed by DNA sequence alone.

A key question related to our findings is how synonymous changes cause vastly different

phenotypes. It is often assumed in endosymbiont transgenic experiments that synonymous changes

do not affect function, and codons for the same amino acid are functionally redundant. However,

several decades of research have uncovered mounting evidence that the functional redundancy

hypothesis is not always accurate[35]. Indeed, codon bias varies across species (hence the need for

codon optimization)[44,45], and there are several mechanisms through which single or few codons

can influence transcription or translation. When there is a rare codon, there are fewer tRNAs

available, and the translation rate may be correspondingly lowered[46]. Notably, there is a bias for

rare codons in N-terminal regions of bacterial genes, which is likely due to their influence on

mRNA structure[47]. Lower 5' RNA stability or weaker structure can also be correlated with ease of

initiation and faster translation[48], and stem loop structures may affect translation rate in varying

ways. Indeed, strong 5' RNA secondary structure may inhibit ribosomal initiation or translational

efficiency[49,50], and there can be selection against SNPs that alter RNA secondary structure[51]. There

is also evidence for selection across the domains of life on codon usage in the first 30-60

nucleotides of a gene, likely due to their impact on mRNA structure near the site of initiation[48].

Additionally, early codons with low frequencies can be beneficial by slowing elongation, thus

preventing ribosome collisions, or potentially helping to recruit chaperones to the emerging

peptide for proper folding[52,53]. In addition, some specific codons across the sequence are favored

for their lower likelihood of mistranslation[54]. Indeed, synonymous codon differences in genes can

result, for instance, in altered gene expression[55], lower organism fitness[56], or disease[57].

Beyond codon sequence, gene expression levels across representatives of all phenotypes

are similar (Fig. 2), supporting post-transcriptional differences as the source of functional variation

and refuting the functional redundancy hypothesis in this context. Examining the three

synonymous transgenes more closely, we find that codon usage indeed correlates with phenotype.

According to the codon usage table for *D. melanogaster* on the Genscript website, the original

*w*Mel *wmk* codon at this position (TCG) has a frequency 16.7 per 1000 codons. In contrast, the

*w*Suzi codon (TCC) has a frequency of 19.5/1000, and the *w*Rec codon (AGC) has a frequency of

20.5/1000[43]. Therefore, there is a pattern where the more common codons are present in the all-

killing transgenes, and at similar frequencies, while the strain expressing partial male killing has a

codon at a lower frequency. Thus, some of the phenomena described in studies on synonymous

codons and function likely underlie the phenotypic variation observed here.

Given that transgenic *wmk* expression with silent site changes results in different

phenotypes, we used software that estimated RNA secondary structures for the tested transgenes.

Indeed, we found that the predicted structures correlate somewhat with phenotype, so synonymous

codons may change transcript structure or other post-transcriptional features of the transcript or

protein (Fig. 2). Other small changes including adding a 9-codon N-terminal sequence ablated the

phenotype and reversed it to no death, despite the additional sequence being smaller than many

common gene tags that typically do not interfere with function. This small change also alters

predicted RNA structure (Fig. 2- supplement 1). We were unable to assess protein levels to

determine translational differences due to lack of an antibody. However, the data demonstrates

that the *wmk* transgene is functionally sensitive to some post-transcriptional changes, at least in

the N-terminal region.

What, then, do these results illuminate about transgenic *wmk* function and the utility of

transgenic research in light of the potential for marked influences of synonymous changes on

phenotype? First, as discussed, the *wmk* transgenic phenotypes are likely sensitive to post-

transcriptional processes. This has important implications for understanding *wmk* in a natural male-

killing context as soon as feasible techniques are available, since heterologous expression and its

reliance on codon optimization may obscure our understanding of the gene's biology. Second,

since three different transgenic phenotypes (sex ratio bias, all killing, no killing) have been found

so far with only a few sequences analyzed, it is possible that further testing of new codons may

increase the partial male killing to a full male-killing phenotype. Therefore, it would be fruitful

and may be possible to continue to uncover a transgenic sequence that fully recapitulates the

phenotype to refine this system as a study tool for *Wolbachia* male killing. Third and critically,

although the results show that there is some relationship between synonymous codons and

phenotype, several points remain for further testing. For example, we cannot conclude that the

particular codon tested here is responsible for phenotype alterations in other host genetic

backgrounds or species. It is possible that this codon plays a functional role only in a singular host

genetic context. Here, we changed *wmk* sequences while holding the host genetic background

fixed, but the reverse is required to conclude whether or not the particular codon plays a general

role in other genotypes or natural contexts. Second, due to possible coevolution, various codons

may or may not yield similar functional effects across different host backgrounds, and additional

synonymous sites may contribute to the male killing phenotype. Thus, the results here illuminate

a previously unrecognized need for future research on the functional impacts of synonymous

substitutions in endosymbionts. Future work may focus on determining if there is one specific

synonymous codon that affects the male killing function in all cases, if a more general feature

exists where alteration of any or a subset of N-terminal or other *wmk* codons affects function, or if

the effect of synonymous changes is specific to this background.

In addition, these findings are informative with regards to the more general study of

phenotypes induced by endosymbiont transgenes. It is standard practice to codon optimize genes

for maximizing host expression when testing endosymbiont gene function[25,27,34], with the

assumption that synonymous codons will not alter function. However, if codon optimization

potentially changes the interpretation of transgenic findings, then phenotypes should be

corroborated in natural contexts once tools such as genetic editing are available in the relevant

organism. Specifically, *wmk* should be knocked out in native contexts once it is more technically

feasible. In addition, codon optimization algorithms are updated with new information periodically

with the assumption that they yield improved results, although it is unclear in practice whether an

algorithm is better optimized to produce results that reflect the true biology of a transgene. Future

work is necessary to explore these concepts further. For example, comparisons of alleles may need

to be performed with alleles identical in sequence except for any engineered differences, and the

algorithm should remain constant across all transgenes that are compared to each other. Further,

careful analysis and comparison of transgenic phenotypes produced by different algorithms may

be necessary in some cases where the phenotype is known or expected. This approach could ensure

the algorithm produces a transgenic phenotype that most closely resembles the natural phenotype.

In addition, the reliability of certain molecular evolutionary signatures, such as dN/dS, may be

compromised since synonymous mutations are assumed to be neutral in these analyses. These

principles are particularly important for research on endosymbionts that increasingly relies on

heterologous gene expression for functional studies. The smaller genomes of endosymbionts tend

to have low GC content overall[58], and GC content is the strongest driver of codon usage bias[59,60]

and contributes to the strength of mRNA secondary structure[35], so careful attention to the effects

of synonymous changes may be of acute interest to the endosymbiont research community.

We also find interesting support for the hypothesis that there is co-adaptation between *wmk*

homologs and their hosts since male-killing genes may be evolutionary matched for the host sex

determination and molecular machinery that they manipulate. We first analyzed *wmk* homologs

that are more distantly related to *w*Mel *wmk*, including from male-killing strains *w*Bol1b, *w*Bif,

*w*Caub, and *w*Inn/*w*Bor[22,39,40,61,62]. They did not recapitulate male killing when transgenically

expressed in *D. melanogaster*. The lack of a sex ratio bias is expected if either *wmk* does not

underpin male killing in these systems or divergent homologs are required to be closely co-adapted

to their hosts. This latter hypothesis is based on observations that resistance to male killing is

common (evidence of a potential host-microbe arms race)[8,14,18,19,63], and some strains can cause

male killing anew when transferred to another, usually closely-related host[18,20,22] while not causing

male killing in all recipient host species[64,65]. It is possible, then, that the *wmk* homologs tested

induce a sex ratio bias in their natural hosts, but not in *D. melanogaster*.

If we make the assumption that *wmk* is involved in male killing in nature, which requires

confirmation beyond transgenic recapitulation of the phenotype, then the results here give the basis

for additional hypotheses that require further testing. First, the killing of both males and females

by the closely-related *w*Rec and *w*Suzi *wmk* homologs could indicate that the target is something

shared in both males and females, but functions differently within each sex (assuming no off-target

effects). Indeed, previous studies on *wmk* and *Wolbachia* male killing demonstrate a positive

correlation with between dosage compensation complex (DCC) activity and DNA defects in male

embryos[21,25,26]. Notably, four of the five protein components of the DCC are expressed in both

males and females, and it is only *msl-2* that is male-specific and catalyzes formation of the

complex[66]. Thus, there are several non-sex-specific *wmk* target candidates that the gene product

may interact with to cause lethality in males and females. Second, protein divergence and resultant

conformational changes may impact the specificity between host target and the Wmk toxin, and

could underlie development of host resistance. Male-killing genes are expected to evolve rapidly

within hosts in order to counteradapt host resistance mechanisms and keep evolutionary pace with

the rapid evolution of sex-related genes that may be the target of a male-killing toxin[67,68]. As such,

major or minor divergence in protein or transcript sequence of either the host target or the microbial

toxin may underpin changes that lead to common host resistance, *wmk*-host coadaptation, and

functional transgene differences in the foreign *D. melanogaster* host. Third, these findings leave

open the possibility of a variety of functionally-relevant *wmk* protein or transcript conformations

in nature, which could contribute to the marked diversity of *Wolbachia* male killing in terms of

host species and sex determination systems[4,8,22,23,40,61,69,70].

Taken together, his work reports previously unrecognized relationships for *wmk*-induced

killing and establishes new hypotheses for the impacts of RNA structure and post-transcriptional

processes in *wmk*-induced male killing. It also highlights several critical features for the research

community regarding assumptions about the broad use of transgenes and the role of synonymous

mutations in gene function. If *wmk* is involved in natural male killing, then this work could indicate

how silent sequence changes may relate to known male-killing phenomena such as frequent host

resistance or male-killing function in diverse hosts.

Materials and Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
gene (*Wolbachia pipientis*)	WD0626	NCBI	NCBI:WD_RS02815	Also known as *wmk* (*WO-mediated killing*)
genetic reagent (*D. melanogaster*)	Act5c-Gal4/CyO	Bloomington Drosophila Stock Center	BDSC:3953; FlyBase FBti0012290	P{AyGAL4}25
genetic reagent (*D. melanogaster*)	WT (y^{1}w^{67c23}; P[CaryP]P2)	Bloomington Drosophila Stock Center	BDSC:8622; FlyBase FBti0040535	WT strain used in this study; P{CaryP}attP2
genetic reagent (*D. melanogaster*)	*w*Mel *wmk*	This paper; Perlmutter *et al.,* 2019 PMID 31504075		Expresses codon-optimized transgene; UAS promoter
genetic reagent (*D. melanogaster*)	*w*Bol1b *wmk*	This paper		Expresses codon-optimized transgene; UAS promoter
genetic reagent (*D. melanogaster*)	*w*Bif *wmk*	This paper		Expresses codon-optimized transgene; UAS promoter

genetic reagent (*D. melanogaster*)	*w*Caub *wmk*	This paper		Expresses codon-optimized transgene; UAS promoter
genetic reagent (*D. melanogaster*)	*w*Inn/*w*Bor *wmk*	This paper		Expresses codon-optimized transgene; UAS promoter; *w*Inn and *w*Bor *wmk* have same exact sequence
genetic reagent (*D. melanogaster*)	*w*Suzi *wmk*	This paper		Expresses codon-optimized transgene; UAS promoter
genetic reagent (*D. melanogaster*)	*w*Rec *wmk*	This paper		Expresses codon-optimized transgene; UAS promoter
genetic reagent (*D. melanogaster*)	HA-*wmk*	This paper		Expresses codon-optimized transgene; UAS promoter; 3X HA tag epitope in linker between HTH domains of *w*Mel *wmk*

genetic reagent (*D. melanogaster*)	5' *w*Rec *wmk*	This paper		Expresses codon-optimized transgene; UAS promoter; Sequence has additional 9 amino acids starting at upstream alternative start codon
genetic reagent (*D. melanogaster*)	5' *w*Suzi *wmk*	This paper		Expresses codon-optimized transgene; UAS promoter; Sequence has additional 9 amino acids starting at upstream alternative start codon
genetic reagent (*D. melanogaster*)	*w*Mel *wmk* (new)	This paper		Expresses codon-optimized transgene; UAS promoter; Same exact sequence as *w*Mel *wmk*, newly transformed strain
genetic reagent (*D. melanogaster*)	*w*Suzi codon	This paper		Expresses codon-optimized transgene; UAS

				promoter; Same as *w*Mel *wmk*, but with 16th amino acid position using TCC Serine codon from *w*Suzi *wmk* strain
genetic reagent (*D. melanogaster*)	*w*Rec codon	This paper		Expresses codon-optimized transgene; UAS promoter; Same as *w*Mel *wmk*, but with 16th amino acid position using AGC Serine codon from *w*Rec *wmk* strain
recombinant DNA reagent	pTIGER (plasmid)	Ferguson *et al.,* 2012; PMID 22328499		Modified pUASp plasmid for enhanced germline expression under Gal4/UAS control
sequence-based reagent	Rp49_F	This paper	PCR primers	CGGTTACG GATCGAA CAAGC
sequenced-based reagent	Rp49_R	This paper	PCR primers	CTTGCGCT TCTTGGAG GAGA

sequenced-based reagent	wmk_homologs_opt_F	This paper	PCR primers	CTGTATGC CATTGCCG AGACCCT
sequenced-based reagent	wmk_homologs_opt_R	This paper	PCR primers	TCACCAG ATCCTTGG CGATCTTC ATC
sequenced-based reagent	Msl-2_F	This paper	PCR primers	GGATTAA CGCGGTCT AAGCATG TGTAACTG
sequenced-based reagent	Msl-2_R	This paper	PCR primers	GTATGCCG TCTGGGCC ATGATG
commercial assay or kit	Direct-zol RNA MiniPrep Kit	Zymo	R2051	
commercial assay or kit	Superscript VILO cDNA Synthesis Kit	ThermoFisher	11754050	
chemical compound, drug	DNase, RNase-free	Ambion, Life Technologies	AM2222	
chemical compound, drug	iTaq Universal SYBR Green Mix	Bio-Rad	1725120	
software, algorithm	GraphPad Prism 8	GraphPad Prism 8	RRID:SCR_0 02798	
software, algorithm	Geneious Pro v.2019.2; Geneious Pro v.2020.2.4	Geneious	RRID:SCR_0 10519	

software, algorithm	jModelTest	jModelTest	RRID:SCR_0 15244	
software, algorithm	RNAfold WebServer	University of Vienna	Gruber *et al.*, 2008, PMID 18424795; Lorenz *et al.*, 2011, PMID 22115189	http://rna.tbi. univie.ac.at/c gi- bin/RNAWe bSuite/RNAf old.cgi

**Reagents source from this paper may be obtained from Bordenstein lab

Drosophila strains and maintenance

D. melanogaster strains used in this study include *Act5c*-Gal4/CyO (BDSC 3953,

ubiquitously-expressing zygotic driver), the WT background line of genotype y^1w^{67c23};

P[CaryP]P2 (BDSC 8622), the WD0626 (*wmk*) and WD0034 (control gene) transgene constructs

previously described[25], and several new transgene constructs. Briefly, each *Wolbachia* gene of

interest was codon-optimized for optimal *D. melanogaster* expression using algorithms developed

by GenScript Biotech (Piscataway, NJ). These sequences were then synthesized as DNA

nucleotides and cloned by GenScript into the pTIGER plasmid[71]. The pTIGER vector is a UASp-

based plasmid optimized for germline expression and uses PhiC31 integrase[72] for targeted

integration into the *D. melanogaster* genome. It also includes UAS promoters for inducible

expression with a Gal4 driver[73] as well as a w^+ red eye marker for transformant screening. The

plasmids were then sent to BestGene (Chino Hills, CA), which performed injections of the vectors

into embryos of the BDSC 8622 background line with PhiC31 integrase to integrate the vector into

the attP2 insertion site in the genome. Successful transformants were selected based on the red eye

marker, and each transgene line is descended from the offspring of a single transformant

(isofemale).

D. melanogaster were reared on standard cornmeal, molasses, and yeast (CMY) media (5%

w/v cornmeal (Quaker, Chicago IL), 1.875% w/v yeast (Red Star Yeast, Milwaukee WI), 7.8%

v/v molasses (Sweet Harvest Foods, Rosemount MN), 0.5% w/v type II *Drosophila* agar (Genesee

Scientific, San Diego CA), 0.056% w/v tegosept (Genesee Scientific, San Diego CA), and 0.39%

v/v propionic acid (Sigma Aldrich)). Stocks were maintained at 25°C with virgin flies stored at

room temperature. During virgin collections, stocks were kept at 18°C overnight and 25°C during

the day. All flies were kept on a 12-hour light/dark cycle.

Sex ratio assays

To assess the effect of transgene expression on adult sex ratios (measurement of male

killing), sex ratio assays were performed as previously described[25]. Briefly, twenty biological

replicates of 10 uninfected, 4- to 7-day-old virgin, female *Act5c*-Gal4/CyO driver flies and 2

uninfected, 1- to 2-day-old virgin, male transgene flies were each set up in vials with CMY media.

Individuals were randomly allocated to each vial after all females or males of a given genotype

were mixed together. They were left on the media to lay eggs for 4 days at 25°C with a 12 h

light/dark cycle, at which point adults were discarded. The vials are then left at 25°C until the

offspring are counted. After 9 days of adult offspring emergence, they were scored for both sex

and expression (red eye color from *Act5c*-Gal4 chromosome) or non-expression (curly wings from

CyO balancer chromosome). The number of adult offspring per vial across all experiments ranges

from 50-170, with a mean of 120 (ranges and means per experiment are included in figure

captions). Any vials with fewer than 50 adult offspring were removed from the analysis, as this

indicates either abnormally poor egg laying or hatching (typically 0-2 vials per group). In addition,

vials with no adult emergence, while others of the same genotype had typical levels of offspring,

were also excluded (typically 2-3 vials per group). Results were graphed in GraphPad Prism 8.4.0,

which applies a "Standard Plot Appearance" correction for visibility of data distribution where the

width of distribution of points is proportional to the number of points at that y-value.

RNA secondary structures

RNA secondary structures were generated by uploading the nucleotide sequences of the

indicated gene to the RNA fold web server[74,75]. The structures shown are the graphical outputs of

the MFE (minimum free energy) secondary structures. Colors indicate base pair probabilities, from

blue to red, with blue indicating a probability of 0 and red indicating a probability of 1.

Gene expression

Gene expression was measured in *Drosophila* embryos aged 4-5 h AED. Each point

represents a biological replicate with the RNA of 30 pooled embryos from crosses between a

unique set of 60 uninfected, 4- to 7-day-old virgin, female *Act5c*-Gal4/CyO driver flies and 12

uninfected, 1- to 2-day-old virgin, male transgene flies of the indicated genotype. Each point

represents a biological replicate from different bottles. Individuals were randomly allocated to each

bottle after all females or males of a given genotype were mixed together. Each collection chamber

consisted of a grape juice agar plate with yeast in an 8 oz round bottom bottle, and flies. These

were placed in a 25°C incubator overnight (16 h). Then, the plates were swapped with fresh ones.

The flies were allowed to lay eggs for 1 h. The plates were then left at 25°C for an additional 4 h

to age them to be 4–5 h old (the estimated time of male death in *wmk* crosses). Embryos were then

gathered in groups of 30 (each group from a unique bottle/biological replicate) and flash frozen in

liquid nitrogen. RNA was extracted using the Direct-zol RNA MiniPrep Kit (Zymo), RNase-free

583 DNase (Ambion, Life Technologies), cDNA was generated with SuperScript VILO (Invitrogen),

584 and RT-qPCR was run using iTaq Universal SYBR Green Mix (Bio-Rad). qPCR was performed

585 on a Bio-Rad CFX-96 Real-Time System. Primers are listed in the Key Resources Table.

586 Conditions were as follows: 50°C 10 min, 95°C 5 min, 40x (95°C 10 s, 55°C 30 s), 95°C 30 s.

587 Differences in gene expression were done by calculating $2^{-\Delta ct}$ (difference in ct values of two genes

588 of interest). Data points were excluded if a sample had low quality cDNA that did not amplify in

589 qPCR. Data points of each biological replicate are measured as the mean of two technical replicates

590 from each sample.

592 Phylogenetic trees

593  The nucleotide phylogenetic trees of host COI genes and *wmk* native gene or transgene

594 sequences were inferred based on a MUSCLE alignment in Geneious Prime 2020.2.4 followed by

595 stripping all sites with gaps. The resulting 652, 690, 686 nucleotide base pair alignments

596 (respectively) were then analyzed via jModelTest 2.1.10 v20160303[76,77]. The AICc-corrected best

597 model, JC, was predicted for all three alignments and was used to build the trees using the

598 MrBayes[78,79] Geneious plugin with the JC69 model[80] and equal rate variation.

600 Transgene sequence alignments

601  The gene sequence alignments of different *wmk* transgenes in Figs. 2 and 6 were conducted

602 in Geneious Pro v.2019.2 using a MUSCLE alignment. Black bars in Fig. 2 indicate sequence

603 mismatches compared to the *w*Mel *wmk* transgene reference sequence with any gaps stripped.

604 Codons in Fig. 6 are colored by amino acid.

Statistical analyses

Sample sizes for experiments were based on previous publications, which demonstrated

repeatability in relative differences between treatment groups. Each experiment was completed

twice, and statistical tests were applied to both to confirm repetition in differences between

treatment groups. In all plots, the first experiment is the representative one shown. For sex ratios,

we tested different sample sizes for reliability in sex ratio measurements, and found that 20

biological samples per group, with 10 females and 2 males per sample, resulted in consistently

replicable data, which is the standard we apply here. qPCR data was approached similarly, by

previous work in the lab demonstrating that with embryos 4-5 h AED, we are able to get consistent,

high quality, replicable data with 30 embryos per sample, and at least 8 samples per group.

All statistical analyses for sex ratios and were performed using GraphPad Prism 8 software.

For sex ratios, a non-parametric Kruskal-Wallis one-way ANOVA followed by Dunn's multiple

comparisons test was applied to all gene-expressing categories, followed by the same test but on

all non-expressing categories. For gene expression, groups were compared using the same

Kruskal-Wallis one-way ANOVA with Dunn's multiple comparisons test. Significant results are

indicated with * symbols in figures, with accompanying p values in captions. Any comparisons

with no symbol are nonsignificant. Full statistical information and outputs for all sex ratio and

qPCR data is available in the Source Data file.

Acknowledgements

This work was supported by National Institutes of Health (NIH) grant R21 AI133522 to

S.R.B, the Vanderbilt Microbiome Innovation Center, and NIH grants F31 AI143152, K-INBRE

P20 GM103418, and National Science Foundation (NSF) Postdoctoral Research Fellowship

(PRFB) DBI 2109772 to J.I.P. We would also like to thank R. L. Unckless for helpful comments

on phylogenies.

632   Competing Interests

633   Both J.I.P. and S.R.B. are inventors on a patent related to the transgenic use of *wmk* in

634   vector control: US Patent 20210000092 16/982708.

References

1   Hurst, G. D. & Frost, C. L. Reproductive parasitism: maternally inherited symbionts in a
biparental world. *CSH Perspect Biol* **7**, a017699 (2015).
2   Taylor, M., Bordenstein, S. & Slatko, B. Microbe Profile: *Wolbachia*: a sex selector, a viral
protector and a target to treat filarial nematodes. *Microbiology* **164**, 1345-1347 (2018).
3   Yen, J. H. & Barr, A. R. New hypothesis of the cause of cytoplasmic incompatibility in
Culex pipiens L. *Nature* **232**, 657-658 (1971).
4   Hurst, G. D. D., Jiggins, F. M., von der Schulenburg, J. H. G., Bertrand, D., West, S. A.,
Goriacheva, I. I., Zakharov, I. A., Werren, J. H. Stouthamer, R. & Majerus, M. E. Male–
killing *Wolbachia* in two species of insect. *Proc Royal Soc B* **266**, 735-740 (1999).
5   Bouchon, D., Rigaud, T. & Juchault, P. Evidence for widespread *Wolbachia* infection in
isopod crustaceans: molecular identification and host feminization. *Proc Royal Soc B*
265, 1081-1090 (1998).
6   Hunter, M. S. The influence of parthenogenesis-inducing *Wolbachia* on the oviposition
behaviour and sex-specific developmental requirements of autoparasitoid wasps. *J Evol*
Biol **12**, 735-741 (1999).
7   Hurst, G. D. & Jiggins, F. M. Male-killing bacteria in insects: mechanisms, incidence, and
implications. *Emerg Infect Dis* **6**, 329-336 (2000).
8   Charlat, S., Hornett, E. A., Dyson, E. A., Ho, P. P., Loc, N. T., Schilthuizen, M., Davies, N.,
Roderick, G. K. & Hurst G. D. Prevalence and penetrance variation of male-killing
Wolbachia across Indo-Pacific populations of the butterfly *Hypolimnas bolina*. *Mol Ecol*
14, 3525-3530 (2005).
9   Jaenike, J., Dyer, K. A. & Reed, L. K. Within-population structure of competition and the
dynamics of male-killing *Wolbachia*. *Evol Ecol Res* **5**, 1023-1036 (2003).
10   Unckless, R. L. & Jaenike, J. Maintenance of a male-killing *Wolbachia* in *Drosophila*
innubila by male-killing dependent and male-killing independent mechanisms. *Evolution*
66, 678-689 (2012).
11   Skinner, S. W. Son-killer: a third extrachromosomal factor affecting the sex ratio in the
parasitoid wasp, *Nasonia* (=Mormoniella) *vitripennis*. *Genetics* **109**, 745-759 (1985).
12   Hurst, L. D. The incidences and evolution of cytoplasmic male killers. *Proc Royal Soc B*
244, 91-99 (1991).
13   Jiggins, F. M., Hurst, G. D. & Majerus, M. E. Sex-ratio-distorting *Wolbachia* causes sex-
role reversal in its butterfly host. *Proc Royal Soc B* **267**, 69-73 (2000).
14   Majerus, T. M. & Majerus, M. E. Intergenomic arms races: detection of a nuclear rescue
gene of male-killing in a ladybird. *PLoS Pathog* **6**, e1000987 (2010).
15   Hornett, E. A., Charlat, S., Duplouy, A. M. R., Davies, N., Roderick, G. K., Wedell, N., &
Hurst, G. D. D. Evolution of male-killer suppression in a natural population. *PLoS Biol* **4**,
e283 (2006).
16   Groenenboom, M. A. & Hogeweg, P. Space and the persistence of male–killing
endosymbionts in insect populations. *Proc Royal Soc B* **269**, 2509-2518 (2002).
17   Berec, L., Maxin, D. & Bernhauerova, V. Male-killing bacteria as agents of insect pest
control. *J Appl Ecol* (2016).

18   Jaenike, J. Spontaneous emergence of a new *Wolbachia* phenotype. *Evolution* **61**, 2244-
2252 (2007).

19   Mitsuhashi, W., Ikeda, H. & Muraji, M. Fifty-year trend towards suppression of
Wolbachia-induced male-killing by its butterfly host, *Hypolimnas bolina*. *J Insect Sci* **11**,
92 (2011).

20   Fujii, Y., Kageyama, D., Hoshizaki, S., Ishikawa, H. & Sasaki, T. Transfection of *Wolbachia*
in Lepidoptera: the feminizer of the adzuki bean borer *Ostrinia scapulalis* causes male
killing in the Mediterranean flour moth *Ephestia kuehniella*. *Proc Royal Soc B* **268**, 855-
859 (2001).

21   Riparbelli, M. G., Giordano, R., Ueyama, M. & Callaini, G. *Wolbachia*-mediated male
killing is associated with defective chromatin remodeling. *PLoS One* **7**, e30045 (2012).

22   Sasaki, T., Kubo, T. & Ishikawa, H. Interspecific transfer of *Wolbachia* between two
lepidopteran insects expressing cytoplasmic incompatibility: a *Wolbachia* variant
naturally infecting *Cadra cautella* causes male killing in *Ephestia kuehniella*. *Genetics*
162, 1313-1319 (2002).

23   Zeh, D. W., Zeh, J. A. & Bonilla, M. M. *Wolbachia*, sex ratio bias and apparent male killing
in the harlequin beetle riding pseudoscorpion. *Heredity* **95**, 41-49 (2005).

24   Bordenstein, S. R. & Bordenstein, S. R. Eukaryotic association module in phage WO
genomes from *Wolbachia*. *Nat Commun* **7**, 13155 (2016).

25   Perlmutter, J. I., Bordenstein, S. R., Unckless, R. L., LePage, D. P., Metcalf, J. A., Hill, T.,
Martinez, J., Jiggins, F. M. & Bordenstein, S. R. The phage gene *wmk* is a candidate for
male killing by a bacterial endosymbiont. *PLoS Pathog* **15**, e1007936 (2019).

26   Harumoto, T., Fukatsu, T. & Lemaitre, B. Common and unique strategies of male killing
evolved in two distinct *Drosophila* symbionts. *Proc Royal Soc B* **285**, 20172167 (2018).

27   LePage, D. P., Metcalf, J. A., Bordenstein, S. R., On, J., Perlmutter, J. I., Shropshire, J. D.,
Layton, E. M., Funkhouser-Jones, L. J., Beckmann, J. F. & Bordenstein, S. R. Prophage WO
genes recapitulate and enhance *Wolbachia*-induced cytoplasmic incompatibility. *Nature*
543, 243-247 (2017).

28   Masui, S., Kuroiwa, H., Sasaki, T., Inui, M., Kuroiwa, T. & Ishikawa H. Bacteriophage WO
and virus-like particles in *Wolbachia*, an endosymbiont of arthropods. *Biochem Biophys*
Res Commun **283**, 1099-1104 (2001).

29   Wu, M., Sun, L. V., Vamathevan J., Riegler, M., Deboy, R., Brownlie, J. C., McGraw, E. A.,
Martin, W., Esser, C., Ahmadinejad, N. & Wiegand C. Phylogenomics of the reproductive
parasite *Wolbachia pipientis w*Mel: a streamlined genome overrun by mobile genetic
elements. *PLoS Biol* **2**, E69 (2004).

30   Hamm, C. A. *et al.* Wolbachia do not live by reproductive manipulation alone: infection
polymorphism in Drosophila suzukii and D. subpulchrella. *Molecular ecology* **23**, 4871-
4885 (2014).

31   Hoffmann, A. A. Partial cytoplasmic incompatibility between two Australian populations
of *Drosophila melanogaster*. *Entomol Exp Appl* **48**, 61-67 (1988).

32   Perlmutter, J. I., Meyers, J. E. & Bordenstein, S. R. Transgenic testing does not support a
role for additional candidate genes in *Wolbachia* male killing or cytoplasmic
incompatibility. *mSystems* **5**, e00658-00619 (2020).

33   Metcalf, J. A., Jo, M., Bordenstein, S. R., Jaenike, J. & Bordenstein, S. R. Recent genome
reduction of *Wolbachia* in *Drosophila recens* targets phage WO and narrows candidates
for reproductive parasitism. *PeerJ* **2**, e529 (2014).
34   Harumoto, T. & Lemaitre, B. Male-killing toxin in a bacterial symbiont of *Drosophila*.
Nature **557**, 252 (2018).
35   Plotkin, J. B. & Kudla, G. Synonymous but not the same: the causes and consequences of
codon bias. *Nat Rev Genet* **12**, 32-42 (2011).
36   Gustafsson, C., Govindarajan, S. & Minshull, J. Codon bias and heterologous protein
expression. *Trends Biotechnol* **22**, 346-353 (2004).
37   Drummond, F., Ballman, E. & Collins, J. Population dynamics of spotted wing *Drosophila*
(*Drosophila suzukii* (Matsumura)) in Maine wild blueberry (*Vaccinium angustifolium*
Aiton). *Insects* **10**, 205 (2019).
38   Cheng, B., Kuppanda, N., Aldrich, J. C., Akbari, O. S. & Ferree, P. M. Male-killing
Spiroplasma alters behavior of the dosage compensation complex during *Drosophila*
melanogaster embryogenesis. *Current Biol* **26**, 1339-1345 (2016).
39   Dyson, E., Kamath, M. & Hurst, G. *Wolbachia* infection associated with all-female broods
in *Hypolimnas bolina* (Lepidoptera: Nymphalidae): evidence for horizontal transmission
of a butterfly male killer. *Heredity* **88**, 166 (2002).
40   Dyer, K. A. & Jaenike, J. Evolutionarily stable infection by a male-killing endosymbiont in
Drosophila innubila: molecular evidence from the host and parasite genomes. *Genetics*
168, 1443-1455 (2004).
41   Carson, H. A female producing strain of *D. borealis* Patterson. *Drosoph Inf Serv* **30**, 109-
110 (1956).
42   Bentele, K., Saffert, P., Rauscher, R., Ignatova, Z. & Blüthgen, N. Efficient translation
initiation dictates codon usage at gene start. *Mol Syst Biol* **9** (2013).
43   GenScript. *GenScript Codon Usage Frequency Table (chart) Tool*,
<https://www.genscript.com/tools/codon-frequency-table> (2020).
44   Sharp, P. M., Averof, M., Lloyd, A. T., Matassi, G. & Peden, J. F. DNA sequence evolution:
the sounds of silence. *Proc Royal Soc B* **349**, 241-247 (1995).
45   Andersson, S. & Kurland, C. Codon preferences in free-living microorganisms. *Microbiol*
Mol Biol Rev **54**, 198-210 (1990).
46   Li, G.-W., Oh, E. & Weissman, J. S. The anti-Shine–Dalgarno sequence drives
translational pausing and codon choice in bacteria. *Nature* **484**, 538-541 (2012).
47   Goodman, D. B., Church, G. M. & Kosuri, S. Causes and effects of N-terminal codon bias
in bacterial genes. *Science* **342**, 475-479 (2013).
48   Gu, W., Zhou, T. & Wilke, C. O. A universal trend of reduced mRNA stability near the
translation-initiation site in prokaryotes and eukaryotes. *PLoS Comp Biol* **6** (2010).
49   de Smit, M. H. & Van Duin, J. Secondary structure of the ribosome binding site
determines translational efficiency: a quantitative analysis. *PNAS* **87**, 7668-7672 (1990).
50   Bettany, A. J., Moore, P. A., Cafferkey, R., Bell, L. D., Goodey, A. R., Carter, B. L. & Brown,
A. J. 5'-Secondary structure formation, in constrast to a short string of non-preferred
codons, inhibits the translation of the pyruvate kinase mRNA in yeast. *Yeast* **5**, 187-198
(1989).

51   Salari, R., Kimchi-Sarfaty, C., Gottesman, M. M. & Przytycka, T. M. Sensitive
764   measurement of single-nucleotide polymorphism-induced changes of RNA
765   conformation: application to disease studies. *Nucleic Acids Res* **41**, 44-53 (2013).
766   52   Tuller, T., Carmi, A., Vestsigian, K., Navon, S., Dorfan, Y., Zaborske, J., Pan, T., Dahan, O.,
767   Furman, I. & Pilpel, Y. An evolutionarily conserved mechanism for controlling the
768   efficiency of protein translation. *Cell* **141**, 344-354 (2010).
769   53   Fredrick, K. & Ibba, M. How the sequence of a gene can tune its translation. *Cell* **141**,
770   227-229 (2010).
771   54   Qin, H., Wu, W. B., Comeron, J. M., Kreitman, M. & Li, W.-H. Intragenic spatial patterns
772   of codon usage bias in prokaryotic and eukaryotic genomes. *Genetics* **168**, 2245-2260
773   (2004).
774   55   Kudla, G., Murray, A. W., Tollervey, D. & Plotkin, J. B. Coding-sequence determinants of
775   gene expression in *Escherichia coli*. *Science* **324**, 255-258 (2009).
776   56   Agashe, D., Martinez-Gomez, N. C., Drummond, D. A. & Marx, C. J. Good codons, bad
777   transcript: large reductions in gene expression and fitness arising from synonymous
778   mutations in a key enzyme. *Mol Biol Evol* **30**, 549-560 (2013).
779   57   Sauna, Z. E. & Kimchi-Sarfaty, C. Understanding the contribution of synonymous
780   mutations to human disease. *Nat Rev Genetics* **12**, 683-691 (2011).
781   58   McCutcheon, J. P., McDonald, B. R. & Moran, N. A. Origin of an alternative genetic code
782   in the extremely small and GC–rich genome of a bacterial symbiont. *PLoS Genet* **5**
783   (2009).
784   59   Knight, R. D., Freeland, S. J. & Landweber, L. F. A simple model based on mutation and
785   selection explains trends in codon and amino-acid usage and GC composition within and
786   across genomes. *Genome Biol* **2**, 1-13 (2001).
787   60   Chen, S. L., Lee, W., Hottes, A. K., Shapiro, L. & McAdams, H. H. Codon usage between
788   genomes is constrained by genome-wide mutational processes. *PNAS* **101**, 3480-3485
789   (2004).
790   61   Hurst, G. D., Johnson, A. P., Schulenburg, J. H. & Fuyama, Y. Male-killing *Wolbachia* in
791   *Drosophila*: a temperature-sensitive trait with a threshold bacterial density. *Genetics*
792   **156**, 699-709 (2000).
793   62   Sheeley, S. & McAllister, B. Mobile male-killer: similar *Wolbachia* strains kill males of
794   divergent *Drosophila* hosts. *Heredity* **102**, 286 (2009).
795   63   Hornett, E. A., Duplouy, A. M., Davies, N., Roderick, G. K., Wedell, N., Hurst, G. D. &
796   Charlat, S. You can't keep a good parasite down: evolution of a male-killer suppressor
797   uncovers cytoplasmic incompatibility. *Evolution* **62**, 1258-1263 (2008).
798   64   Veneti, Z., Zabalou, S., Papafotiou, G., Paraskevopoulos, C., Pattas, S., Livadaras, I.,
799   Markakis, G., Herren, J. K., Jaenike, J. & Bourtzis, K. Loss of reproductive parasitism
800   following transfer of male-killing *Wolbachia* to *Drosophila melanogaster* and *Drosophila*
801   *simulans*. *Heredity* **109**, 306-312 (2012).
802   65   Hughes, G. L. & Rasgon, J. L. Transinfection: a method to investigate *Wolbachia*-host
803   interactions and control arthropod-borne disease. *Insect Mol Biol* **23**, 141-151 (2014).
804   66   Lucchesi, J. C. & Kuroda, M. I. Dosage compensation in *Drosophila*. *CSH Perspect Biol* **7**,
805   a019398 (2015).

67 Marın, I. & Baker, B. S. The evolutionary dynamics of sex determination. *Science* **281**,
1990-1994 (1998).
68 Rodriguez, M. A., Vermaak, D., Bayes, J. J. & Malik, H. S. Species-specific positive
selection of the male-specific lethal complex that participates in dosage compensation
in *Drosophila*. *PNAS* **104**, 15412-15417 (2007).
69 Fialho, R. F. & Stevens, L. Male-killing *Wolbachia* in a flour beetle. *Proc Royal Soc B* **267**,
1469-1473 (2000).
70 Van Borm, S., Wenseleers, T., Billen, J. & Boomsma, J. J. *Wolbachia* in leafcutter ants: a
widespread symbiont that may induce male killing or incompatible matings. *J Evol Biol*
14, 805-814 (2001).
71 Ferguson, S. B., Blundon, M. A., Klovstad, M. S. & Schupbach, T. Modulation of gurken
translation by insulin and TOR signaling in *Drosophila*. *J Cell Sci* **125**, 1407-1419 (2012).
72 Groth, A. C., Fish, M., Nusse, R. & Calos, M. P. Construction of transgenic *Drosophila* by
using the site-specific integrase from phage phiC31. *Genetics* **166**, 1775-1782 (2004).
73 Southall, T. D., Elliott, D. A. & Brand, A. H. The GAL4 system: A versatile toolkit for gene
expression in *Drosophila*. *CSH Protoc* **7**, pdb-top49 (2008).
74 Gruber, A. R., Lorenz, R., Bernhart, S. H., Neuböck, R. & Hofacker, I. L. The vienna RNA
websuite. *Nucleic Acids Res* **36**, W70-W74 (2008).
75 Lorenz, R/. Bernhart, S. H., Zu Siederdissen, C. H., Tafer, H., Flamm, C., Stadler, P. F. &
Hofacker, I. L. ViennaRNA Package 2.0. *Algorithms Mol Biol* **6**, 26 (2011).
76 Darriba, D., Taboada, G. L., Doallo, R. & Posada, D. jModelTest 2: more models, new
heuristics and parallel computing. *Nat Methods* **9**, 772 (2012).
77 Guindon, S. & Gascuel, O. A simple, fast, and accurate algorithm to estimate large
phylogenies by maximum likelihood. *Syst Biol* **52**, 696-704 (2003).
78 Huelsenbeck, J. P. & Ronquist, F. MRBAYES: Bayesian inference of phylogenetic trees.
Bioinformatics **17**, 754-755 (2001).
79 Ronquist, F. & Huelsenbeck, J. P. MrBayes 3: Bayesian phylogenetic inference under
mixed models. *Bioinformatics* **19**, 1572-1574 (2003).
80 Jukes, T. H. & Cantor, C. R. Evolution of protein molecules. *Mammal Prot Metabol* **3**, 21-
132 (1969).
836
837