1 **Intergenerational Effects of Early Life Starvation on Life-History, Consumption,**

2 **and Transcriptomes of a Holometabolous Insect**

4 Sarah Catherine Paul[1,a], Pragya Singh[1,a], Alice B. Dennis[2], Caroline Müller[1*]

6 [1]Chemical Ecology, Bielefeld University, Universitätsstr. 25, 33615 Bielefeld,

7 Germany

8 [2]Evolutionary Biology & Systematic Zoology, University of Potsdam Karl-Liebknecht-

9 Strasse 24-25, 14476 Potsdam

10 [a] shared first authorship

11 *corresponding author: caroline.mueller@uni-bielefeld.de

13 **Running title:** Intergenerational effects on sawfly

15 **Keywords**

16 Intergenerational effects, match-mismatch, sawfly, transcriptome, parental effects,

17 compensatory growth.

ABSTRACT: Intergenerational effects, also known as parental effects in which the offspring phenotype is influenced by the parental phenotype, can occur in response to parental early life food-limitation and adult reproductive environment. However, little is known about how these parental life stage-specific environments interact with each other and with the offspring environment to influence offspring phenotypes, particularly in organisms that realize distinct niches across ontogeny. We examined the effects of parental early life starvation and adult reproductive environment on offspring traits under matching or mismatching offspring early life starvation conditions using the insect *Athalia rosae* (turnip sawfly). The parental early life starvation treatment had context-dependent intergenerational effects on life-history and consumption traits of offspring larvae, partly in interaction with offspring conditions (intragenerational effects) and sex, while there was no significant effect of parental adult reproductive environment. In addition, parental starvation led to differential expression of only few genes, while offspring larval starvation caused a pronounced differential gene expression and modulation of multiple pathways. Our findings reveal that parental starvation leads to intergenerational effects on offspring life-history traits, consumption patterns as well as gene expression, although the effects are less pronounced than those of offspring starvation.

Introduction

Intergenerational effects, more commonly known as parental effects, are defined as causal influences of parental phenotype on offspring phenotype that are likely mediated by non-DNA sequence-based inheritance (Wolf and Wade 2009; Perez and Lehner 2019). Together with transgenerational effects, which occur across several generations, they play a key role in the ecology and evolution of organisms (Badyaev and Uller 2009) and impact the responses of individuals to changing environmental conditions (Sánchez-Tójar et al. 2020). One important environmental factor that can rapidly change during an organism's lifetime and across generations is the food availability. Periods of starvation are commonly encountered by insects in the wild (Jiang et al. 2019) and, when experienced early in life, often have long-lasting effects on an individual's phenotype, affecting life-history, consumption, and behavior (Miyatake 2001; Wang et al. 2016; McCue et al. 2017). Intergenerational effects of starvation events on offspring developmental trajectories have been found to differ depending on the exact starvation regime, sex, and traits that are considered (Saastamoinen et al. 2013; McCue et al. 2017; Paul et al. 2019). Moreover, to fully elucidate the adaptive nature of intergenerational effects, studies should encompass different phases in development during the parental life cycle and how such experience affects individual offspring phenotypes (English and Barreaux 2020).

The early life environment can shape the phenotype of an individual and its offspring via distinct routes (Monaghan 2008; Burton and Metcalfe 2014; Taborsky 2017). Individuals experiencing a benign environment in early life may pass on their superior condition to their offspring (van de Pol et al. 2006), often called silver spoon or carry-over effects (Monaghan 2008), resulting in offspring that are better able to deal with stressful conditions (Franzke and Reinhold 2013). Similarly, negative effects of a poor start in life might also be passed on, resulting in offspring less able to cope with stressful conditions (Naguib et al. 2006). Alternatively, such stressed

individuals may produce offspring that are buffered against stressors, for example,

through parental provisioning (Valtonen et al. 2012; Pilakouta et al. 2015; Hibshman

et al. 2016). Finally, identical environmental conditions experienced in a certain stage

(i.e. a match between parental and offspring environment) may be predicted to be

optimal for the offspring (match-mismatch scenario). Evidence for such predictive

adaptive effects has been found in several species (Raveh et al. 2016; Le Roy et al.

2017), but is weak in others (Uller et al. 2013). Each of these avenues of

intergenerational effects are not mutually exclusive. For example, offspring in

mismatched environments may do relatively better if their parents were of high

quality, due to silver spoon effects (Engqvist and Reinhold 2016). Fully reciprocal

designs are needed to determine the relative importance of the experience in a

certain life-stage or generation on such developmental plasticity (Groothuis and

Taborsky 2015).

In addition, parental investment in offspring can vary depending on the condition of

the parent's mate (Ratikainen and Kokko 2010). Both males and females have been

shown to enhance investment in response to perceived increases in mate quality or

attractiveness (Cunningham and Russell 2000; Cornwallis and O'Connor 2009).

Changes in parental investment based on mate cues may potentially counteract or

augment the effects of the parental early life environment on offspring phenotype.

These effects may act in concert; for example, poor early life conditions can

contribute to poor quality of mating partners and therefore affect mating success and

reproductive investment (Hebets et al. 2008; Vega-Trejo et al. 2016). As niches often

shift across an individual's lifetime, both parental early and later life environments

and their interaction must be considered.

With discrete phases during development, insects, particularly holometabolous

insects, present ideal organisms in which to investigate how intra- and

intergenerational environmental cues, such as food limitation and mate quality, might

influence individual phenotypes (English and Barreaux 2020). Periods of low

resource availability during larval development are known to influence developmental

trajectories with knock-on effects on adult size, reproductive success and adult

starvation resistance (Boggs and Niitepõld 2016; Wang et al. 2016). Individuals may

respond to periods of starvation with increased food uptake (Regalado et al. 2017) as

well as compensatory growth (i.e. a faster growth rate) or catch-up growth (i.e.

attaining a minimum size by prolonging the larval development) (Hector and

Nakagawa 2012). Moreover, starvation leads to substantial shifts in gene expression

within individuals (Moskalev et al. 2015; Jiang et al. 2019; Etebari et al. 2020;

Farahani et al. 2020). However, little is known how long these effects may persist

(McCue et al. 2017) and to which extent parental starvation may affect gene

expression of offspring experiencing starvation.

In the present study, we investigated the effects of parental early life starvation

and parental reproductive environment on offspring (i.e. intergenerational effects),

under matching or mismatching larval starvation conditions, using the turnip sawfly,

Athalia rosae (Hymenoptera: Tenthredinidae). Adult females oviposit on various

species of Brassicaceae, including crop species, on which *A. rosae* can become a

pest (Riggert 1939; Sawa et al. 1989). The leaf-consuming larvae can readily

experience periods of starvation when their hosts are overexploited (Riggert 1939).

The adults are nectar-feeding and in addition collect neo-clerodanoid-like compounds

(hereafter called 'clerodanoids') from non-Brassicaceae plants, which improve their

mating probability (Amano et al. 1999) and affect the social interactions between

adults of both sexes (preprint Paul et al. 2021; preprint Paul and Müller 2021). We

measured various key life-history traits to assess the combined and interactive

effects of parental and offspring larval resource availability (starvation/non-

starvation), and parental clerodanoid exposure on the offspring. Moreover, we

measured effects of different early life starvation regimes on consumption and gene

expression of offspring. We predicted that 1) there will be stronger intragenerational

than intergenerational effects of larval starvation treatment on all parameters

measured; 2) matching conditions between parental and offspring starvation

treatment are beneficial, while a mismatch leads to a reduced performance; 3)

offspring phenotype will also be influenced via parental effects such as silver spoon

or parental buffering effects that may augment or dampen potential mismatch effects;

4) clerodanoid exposure will increase parental investment in offspring due to an

increased mate attractiveness, leading to positive effects for the offspring.

128   **Materials and Methods**

129   *Set-Up of Insect Rearing and Plant Cultivation*

Adults of *A. rosae* were collected in May 2019 from meadows adjacent to farmland in

Germany at two locations (population A 52°02'48.0"N 8°29'17.7"E, population B:

52°03'54.9"N 8°32'22.2"E). Adults from each population were split between two

cages (total of 4 cages) and maintained for one generation at room temperature and

16 h: 8 h light:dark on potted plants of white mustard (*Sinapis alba*) and Chinese

cabbage (*Brassica rapa* var. *pekinensis*). F1 adults were used to set up a total of six

cages with males and females from different populations in four cages and two cages

with virgin females only (Fig. 1). This sawfly species is haplodiploid, i.e. virgin

females produce male offspring (Naito and Suzuki 1991). Larvae of the final instar,

the eonymphs, were placed into individual cups containing soil for pupation. Emerged

adults were kept individually in Petri dishes and provided with a honey:water mixture

(1:50). For mating, pairs of non-sib F2 females and males (N = 20 per treatment

level) were placed together and allowed to mate at least once. Afterwards, mated as

well as virgin females were placed individually into boxes (25 x 15 x 10 cm) with a

middle-aged leaf of 6-8 week old cabbage plants supplied with water for oviposition

and a honey:water mixture in a climate chamber (20 °C:16 °C, 16 h: 8 h light:dark,

70% r.h.). The boxes were checked and topped up with honey water daily and an

additional leaf was added if the first leaf showed signs of wilting. Females were

removed from the boxes after one week and their offspring used to set up the

experimental generations (Fig. 1).

Plants of *B. rapa* and *S. alba* were grown from seeds (Kiepenkerl, Bruno Nebelung

GmbH, Konken, Germany) in a greenhouse (20 °C, 16 h: 8 h light:dark, 70% r.h.) and

a climate chamber (20 °C, 16 h: 8 h light:dark, 70% r.h.). Plants of *Ajuga reptans*

used for clerodanoid supply were grown from seeds (RHS Enterprise Ltd, London,

UK) in the greenhouse and transferred outside in late spring once ~2 months old.

Medium aged leaves were picked from non-flowering plants when plants were

approximately 8 months old.

Experimental Set-Up and Measurements of Life-History Data

The leaves of *B. rapa* offered to F2 females were checked daily for hatching larvae.

Hatched larvae were individually placed into Petri dishes (5.5 cm diameter) with

moistened filter paper and *B. rapa* leaf discs cut from 7-10 week old plants. Leaf

discs were provided *ad libitum* (except during starvation periods) and replaced daily.

Each dish was checked daily for the presence of exuviae to track the larval instar.

Larvae from each box (parental pair) in both the F3 and F4 generation were split

equally between one of two treatments, no starvation (N) or starvation (S). Starvation

individuals were twice starved for 24 h, first the day after moulting into 2nd instar (L2)

and second on the day of moulting into 4th instar (L4) to minimize early mortality

whilst mimicking the food deprivation larvae may experience when occurring in high

densities (Riggert 1939).

Eonymphs were placed in individual cups with soil for pupation. On the day of

emergence, adults were weighed (Sartorius AZ64, M-POWER Series Analytical

Balance, Göttingen, Germany), kept individually in Petri dishes and provided with

honey:water mixture. For the F3 adults, pairs of non-sib females and males reared

under the same larval starvation treatments (N or S) were assigned to one of two

reproductive environment treatments (C- or C+). C+ individuals were exposed to a

leaf section (1 cm^2) of *A. reptans* for 48 h prior to mating, giving individuals time to

take up clerodanoids (preprint Paul et al. 2021). Mated F3 females (2-9 days old)

from each of these four treatments (NC-, NC+, SC-, SC+) and C- virgin females (NC-,

SC-) were then placed in individual breeding boxes, as described above for the F2.

The experimental generations (F3-F4) were reared to test the effects of parental (F3)

and offspring (F4) larval starvation, and their interaction with the adult reproductive

environment on the offspring phenotype (F4). In both generations, larval, pupal and

total development time (from larva to adult) as well as the adult body mass at the day

of emergence were taken for each individual. In total 358 F3 larvae (177 females,

181 males) and 486 F4 larvae (282 females, 202 males) reached adulthood, out of

the 607 and 688 larvae, respectively, that were reared in each generation. Sample

sizes for the F3 and F4 generation are given in Fig. 1. Experimental rearing and

consumption assays were all carried out in a climate chamber (20 °C:16 °C, 16 h: 8 h

light:dark, 70% r.h.).

191   *Consumption Assays*

Consumption assays were performed with F4 larvae at the start of L3 to measure the

relative growth rate (RGR), relative consumption rate (RCR), and efficiency of

conversion of ingested food (ECI). Each L3 larva was weighed (= initial body mass)

(ME36S, accuracy 0.001 mg; Sartorius, Göttingen, Germany) and provided with four

fresh leaf discs (surface area of 230.87 mm^2 per disc). After 24 hours, larvae were

weighed again (= final body mass) and the leaf disc remains were scanned

(Samsung SAMS M3375FD, resolution 640 x 480). The total area of leaf consumed

(mm^2) was then calculated as the difference between the average leaf area and the

remaining leaf area.

Statistical Analyses

All data were analyzed using R 4.0.2 (2020-06-22) -- "Taking Off Again". We set α =

0.05 for all tests and model residuals were checked for normality and variance

homogeneity. Stepwise backwards deletion using Chi-squared ratio tests

(package:MASS) was employed to reach the minimum adequate model (Crawley

2012). Posthoc analyses were carried out using the package 'multcomp' (Hothorn et

al. 2008). Post data entry, raw data were visually inspected thrice, all variables

plotted and outliers and possible anomalies in the data (e.g. strings of similar values)

interrogated (package:pointblank). Intragenerational effects of starvation on life-

history traits of F3 individuals were tested as described in S1.

Due to observations made during the experiment (no *a priori* hypothesis), we tested

in the F4 generation whether the likelihood of an additional larval instar (7 for females

and 6 for males) differed based on offspring larval starvation treatment (OS: N or S)

using a binomial generalized linear mixed model (glmm), where the predictor was OS

and parental pair was included as a random effect. The effects of the parental larval

starvation treatment (PS: N or S), parental clerodanoid exposure (PC: C- or C+), OS

and their interaction on larval, pupal and total developmental time as well as adult

mass of F4 individuals were assessed in separate linear mixed models (lmm), with

parental pair included as a random effect (controlling for non-independence of sibling

larvae). Female and male data were analyzed separately to enable model

convergence. Male adult mass was log-transformed to improve model fit.

RGR, RCR, and ECI (Castagneyrol et al. 2018) of F4 larvae were analyzed using

glmm. We excluded PC as a predictor variable from the consumption assay analysis

due to the low number of individuals in certain treatments (Fig. 1) and analyzed male

and female data separately as above. To assess variation in RGR, the change in larval body mass [final mass — initial mass] was used as the response variable and initial larval body mass, PS, OS, and both their three-way and two-way interactions as the predictors. To assess variation in RCR, we used the total area of consumed leaf material as the response variable and initial larval body mass, PS, OS, and both their three-way and two-way interactions as the predictors. To assess variation in ECI, we used the change in larval body mass as the response variable and the total area of consumed leaf material, PS, OS, and both their three-way and two-way interactions as the predictors. Parental pair was included in all three models as a random effect.

Sample Collection, Library Preparation, Sequencing and Alignment for RNAseq

A total of 24 male larvae (L4, 9 d old), comprising six biological replicates per treatment level (F3/F4: N/N, N/S, S/N, S/S, all C-, Fig. 2), were collected to investigate the effects of parental and offspring starvation treatment on gene expression in larvae of the offspring generation (F4). Individuals were chosen in a way that maximized the equal spread of siblings across treatments and frozen at -80 °C prior to extraction. RNA was extracted with an Invitrogen PureLink™ RNA Mini Kit (ThermoFisher Scientific, Germany), including a DNase step (innuPREP DNase I Kit, analyticJena, Jena, Germany). RNA quality was assessed on a bioanalyzer 2100 (Agilent, CA, United States) and Xpose (PLT SCIENTIFIC SDN. BHD, Malaysia). Library preparation (Ribo-Zero for rRNA removal) and sequencing (NovaSeq6000 and S4 Flowcells, Illumina, CA, United States) were provided by Novogene (Cambridge, UK). Sequence quality before and after trimming was assessed using FastQC (v. 0.11.9; Andrews 2010). After short (< 75 bp), low quality (Q < 4, 25 bp sliding window) and adapter sequences were removed using Trimmomatic (Bolger et al. 2014), more than 98% of reads remained. Reads were mapped to the annotated

genome of *A. rosae*, version AROS v.2.0 (GCA_000344095.2) with RSEM v1.3.1 (Li

and Dewey 2011), which implemented mapping with STAR v2.7.1a (Dobin et al.

2013).

Differential expression analysis was conducted with DESeq2 (version 1.28.1; Love

et al. 2014). RSEM results were passed to DESeq2 for gene-level analyses using

Tximport (version 1.16.1; Soneson et al. 2015). Prior to analysis, genes with zero

counts in all samples and those with low counts (< 10) in less than a quarter of

samples (6) were excluded. Model fitting was assessed by plotting dispersion

estimates and outliers were inspected using principle component analysis of all

expressed genes and pairwise-distance matrices between samples. DESeq2 allows

for both the normalization of read counts (accounting for differences in library

depth/sequencing depth, gene length, and RNA composition) and the shrinkage of

gene-wise dispersion (accounting for differences in dispersion estimates between

genes) whilst modelling a negative binomial glm of predictors for each gene to correct

for overdispersion (Love et al. 2014). Expression was modelled based on the entire

dataset, with the four levels representing the combination of parental and offspring

starvation treatments (F3/F4): N/N, N/S, S/N, and S/S. Differential expression was

based on four pairwise comparisons between the treatments: 1) N/N vs N/S and 2)

S/N vs S/S were used to assess the effects of offspring starvation treatment for

individuals whose parents experienced the same starvation treatment, whereas 3)

N/N vs S/N and 4) N/S vs S/S were used to assess the effects of differing parental

starvation treatment on individuals that experienced the same offspring starvation

treatment. Significance was based on a Wald Test and shrunken LFC values with

apeglm (version 1.10.1; Zhu et al. 2019). P-values were adjusted for multiple testing

using Benjamini-Hochberg (Benjamini and Hochberg 1995) procedure with a false

discovery rate (FDR) of 0.05. Afterwards, significantly differentially expressed genes

(relatively up- or downregulated) were extracted and filtered with a p-adjust of < 0.05.

Among these significantly differentially expressed genes we searched for the terms

linked to genes with known roles in stress and/or starvation response, using the

keywords: "heat shock protein" (or "hsp"), "cytochrome P450", "octopamine" and

"tyramine" in the putative gene names. Venn diagrams (Venn.Diagram, version

1.6.20) were used to depict the relationship between the differentially expressed

genes for each pairwise comparison. The expression of the significantly expressed

genes of each comparison was visualized in a heatmap using normalized counts

scaled per gene (scale="row") (pheatmap, version 1.0.12).

Unmapped reads were extracted to check for the differential expression of genes

not present in the reference genome from the RSEM-produced BAM file using

samtools view (Li et al. 2009) and converted to fastq (using bamtools bamtofastq;

Barnett et al. 2011). The unmapped reads were mapped back to this reference and

expected read counts extracted using eXpress (Roberts et al. 2011). Transcripts with

>10,000 mapped reads were identified using BLASTN (default settings, limited to one

match per gene and $1e^{-1}$) against the NCBI nucleotide database. This process was

done jointly with reads from additional samples (preprint Paul et al. 2021) to optimize

coverage for the assembly. Differential expression of unmapped reads in the larval

samples was carried out as for mapped reads (see above).

For characterization of differential expression at the pathways level, KEGG (Kyoto

Encyclopedia of Genes and Genomes) terms were assigned to the predicted genes

in the *A. rosae* genome using the KEGG Automatic Annotation Server (KAAS; Moriya

et al. 2007) (S2). Of the annotated genes for which we had read counts, 65% were

assigned to at least one KEGG pathway. A gene set enrichment analysis was then

performed on the normalized counts using GAGE (Luo et al. 2009) and pathview

(Luo and Brouwer 2013) in R, applying an FDR-adjusted *P*-value cut-off of < 0.05 to

identify differentially expressed pathways. We derived the non-redundant significant

gene set lists, meaning those that did not overlap heavily in their core genes, using

esset.grp and a *P*-value cut-off for the overlap between gene sets of $10e^{-10}$.

309   **Results**

310   *Life-History Traits*

In the parental generation (F3), larval starvation had trait- and sex-specific effects on

life-history, leading to a prolonged total development time for both sexes and a lower

body mass in females (for details see S1). In the offspring generation (F4), both

females ($X^2{}_1$ = 11.06, *P* < 0.001) and males ($X^2{}_1$ = 29.25, *P* < 0.001) were

significantly more likely to have an additional larval instar prior to the eonymph stage

if they were starved than if they were not starved (Fig. 2A,B). The exposure of

parents to clerodanoids (PC) had no influence on any of the life-history parameters

measured in the offspring, whereas the effect of parental (PS) and offspring larval

starvation treatment (OS) were trait- and sex-specific. There was an interactive effect

of PS and OS on both female (PS * OS: $X^2{}_1$ = 10.37, *P* = 0.001) and male (PS * OS:

$X^2{}_1$ = 4.89, *P* = 0.027) total development time, such that intra-generational starvation

led to a longer development time for both sexes (Fig. 2C,D; S3A, S3B). Larval

development of F4 females was only affected by OS (OS: $X^2{}_1$ = 319.40, *P* < 0.001;

S4A), while there was an interactive effect of PS and OS on male larval development

time (PS * OS: $X^2{}_1$ = 10.37, *P* = 0.001; S3B, S4B). The reverse occurred for pupal

developmental time with an interactive effect of PS and OS on females (PS * OS: $X^2{}_1$

= 24.57, *P* < 0.001, S3A, S4C), but only a significant effect of OS on males (OS: $X^2{}_1$

= 17.23, *P* < 0.001; S4D). Both PS ($X^2{}_1$ = 4.02, *P* = 0.045) and OS ($X^2{}_1$ = 12.95, *P* <

0.001) independently influenced female adult mass (PS * OS: n.s.), with F4 females

having a lower mass if they had experienced starvation during development, but

those females whose parents were starved had a higher overall mass than when

parents had not starved (Fig. 2E). In contrast, F4 male adult mass was significantly

affected by the interaction between PS and OS (PS *OS: $X^2_1 = 4.58$, $P = 0.030$, Fig. 2F). Male offspring of starved parents that themselves were not starved during development (S/N) had a higher mass than offspring that were starved (S/S) (pairwise comparison: $z = -3.47$, $P = 0.003$, Table S3B).

Consumption

OS had a significant effect on the relationship between change in mass and initial body mass (i.e. RGR) in offspring larvae in both females (initial body mass * OS: $F_{1,60.75} = 5.96$, $P = 0.017$) and males (initial body mass * OS: $F_{1,39.54} = 7.32$, $P = 0.009$). The change in body mass of starved larvae increased more steeply with an increase in initial larval body mass compared to non-starved larvae for both sexes (Fig. 3A,B), indicating a higher growth rate (RGR) in starved larvae. There was no effect of PS on RGR either independently or interactively for both sexes. In contrast to RGR, OS did not have a significant effect on RCR either independently or interactively for both sexes. For females, there was a significant interaction effect of initial larval body mass and PS on the area of leaf consumed (initial body mass * PS: $F_{1,62} = 6.15$, $P = 0.015$), such that the area increased with initial body mass for individuals from non-starved parents but did not change for individuals from starved parents (Fig. 3C); i.e. larger larvae consumed more than smaller larvae when their parents were not starved but not when their parents were starved. For males, initial body mass had a significant positive effect on the area of leaf consumed (initial body mass: $F_{1,41.45} = 21.97$, $P < 0.001$) across all treatments (Fig. 3D); i.e. there was no effect of either OS or PS on RCR in males. For ECI (change in mass per unit leaf area consumed), leaf area consumed and OS independently influenced change in body mass for both females (leaf area consumed * $F_{1,54.12} = 7.17$, $P = 0.009$; OS: $F_{1,54.77} = 11.36$, $P = 0.001$) and males (leaf area consumed * $F_{1,44} = 31.55$, $P < 0.001$; OS: $F_{1,44} = 6.92$, $P = 0.011$). The ECI was higher when larvae were not starved than

360   when they were starved, but under both conditions consuming more led to a higher

361   body mass increase for larvae (Fig. 3E, F).

363   *Gene Expression*

Sequencing of the 24 larvae resulted in a total of 1.4 billion reads with an average of

58 million reads per sample (SE: ±2 million reads) and an average GC content of

43%. Prior to normalization, the average number of mapped reads per sample was

33,675,600, equating to an average of 91% reads per sample aligned to the

reference genome (range 85-93%).

Offspring starvation had the strongest effect on gene expression (Figs 4, 5, S5).

The two comparisons between starved and non-starved larvae revealed 4727 (N/S vs

N/N) and 5089 (S/N vs S/S) significantly differentially expressed genes (LFC > 0,

adjusted p < 0.05) and a large number of these overlapped between the two

comparisons (4002). In contrast, there were far fewer significantly differentially

expressed genes in the two comparisons in which the larvae had the same starvation

treatment, namely 155 (S/N vs N/N) and 75 (N/S vs S/S). In terms of genes known to

be associated with stress response, in comparison to non-starved larvae when larvae

from non-starved parents were starved, there was significant downregulation of

putative heatshock proteins (8 downregulated and 1 upregulated) and upregulation of

putative cytochrome P450 genes (21 genes upregulated, 6 downregulated) as well

as upregulation of 1 octopamine and 1 tyramine gene. When parental starvation

conditions but not offspring starvation treatments differed, there was far less

differential expression of such genes with only one or two cytochrome P450 genes

being differentially expressed (S6).

While the majority of reads mapped to the reference genome, 247,997,896 reads

did not map; we assembled these into 334,717 transcripts using Trinity (default

settings) (Grabherr et al. 2011). Of those genes that we assembled *de novo*, 803

were putatively annotated against the NCBI nt database and of these a large portion were mitochondrial (Table 1). DE analysis of unmapped reads showed that a lower number of genes were differential expressed between the treatment pairs [1523 (N/S vs N/N), 1860 (S/N vs S/S), 149 (S/N vs N/N) and 34 (N/S vs S/S)]. Many of these were *A. rosae* genes linked to the mitochondria, particularly in the comparisons between starved and non-starved offspring individuals.

The KEGG pathway level analysis showed that across all four comparisons the ribosomal pathway was the most consistently differentially regulated, being upregulated when offspring were starved and downregulated when they were not. Interestingly, ribosomal regulation was affected not only by differences in offspring starvation treatment but also by differences in parental starvation treatment. The pathway was upregulated in starved larvae when compared to larvae that were not starved (offspring treatment differed) and to larvae that were but whose parents were also starved (parental treatment differed), with the reverse trend for non-starved larvae (Table 1). The other significantly differentially expressed KEGG pathways only occurred between individuals that differed in offspring but not parental larval starvation treatments, mirroring the gene expression results. These pathways belong to the four main categories: metabolic breakdown and protein processing (pancreatic secretion, proteasome and protein processing in the endoplasmic reticulum, ER), immune response (phagosome and antigen processing), energy production and central metabolic processes (TCA/citrate cycle, glycolysis, thyroid hormone signalling pathway), and ECM-receptor interaction. There was generally a downregulation of these processes in larvae that were starved compared to non-starved individuals.

Discussion

We studied the influence of intergenerational effects on offspring phenotype in the holometabolous insect *A. rosae*, which realizes distinct niches during its life-cycle.

Our results revealed that offspring starvation interacted with parental starvation in a

trait-and sex-specific manner for life-history traits. There was an interactive effect of

intergenerational (parental) and intragenerational (offspring) starvation treatment on

total development, male larval development, and female pupal development time.

The trend for faster development times in offspring that experienced a similar

environment as their parents was at least significant for female pupal development

time and may be suggestive of positive effects of matching conditions, i.e. a match-

mismatch scenario (Monaghan 2008). A matching of offspring phenotypes to

environments experienced by their parents is possible due to developmental plasticity

and potentially based on epigenetic effects (Le Roy et al. 2017). Such parental

effects have been found in a number of species in response to food availability

(Hibshman et al. 2016; Raveh et al. 2016) and can be sex-specific (Le Roy et al.

2017). However, other life history data of *A. rosae* did not reveal this match-mismatch

pattern, indicating a complex interplay between different intergenerational effects.

Adult mass showed a different pattern than development time, with female

offspring of *A. rosae* having a higher adult mass when their parents were starved as

larvae irrespective of their own starvation conditions, indicative of enhanced parental

provisioning to offspring of starved parents, i.e. parental buffering. Similar effects

have also been observed in the vinegar fly, *Drosophila melanogaster*, where females

reared on poor food had larger offspring than females reared on standard food

(Valtonen et al. 2012). This finding is consistent with life-history theory prediction that

under hostile conditions there is a shift towards fewer but better provisioned offspring

(Smith and Fretwell 1974; Roff 1992), although we did not examine the number of

offspring produced in our study. As neither larval developmental time for female *A.*

rosae nor their consumption (RGR and ECI) were affected by the parental starvation

treatment, other physiological effects (discussed below for the gene expression) may

mediate this kind of buffering, visible in the higher adult body mass of individuals

whose parents were starved. In contrast, for male *A. rosae*, only offspring of starved

parents differed significantly in adult mass, with lower mass in starved sons. Thus,

investment in body mass may be particularly low in males under repeatedly (across

generations) experienced poor environmental conditions, indicating "paternal

suffering" rather than buffering. Such trait-specific and sex-specific effects of

intergenerational treatment and interaction with intragenerational treatment have

been seen in previous studies (Zizzari et al. 2016; Le Roy et al. 2017; Wilson et al.

2019; Yin et al. 2019), suggesting that context-dependent effects of intergenerational

starvation may be common. We can particularly expect sex-specific differences as a

result of sex-specific directional selection on life-history traits (Tarka et al. 2018).

Overall, there was no clear indication of silver spoon effects for any of the life-history

traits, but such effects cannot be entirely ruled out (Engqvist and Reinhold 2016).

Consumption assays revealed that starved larvae had a marginally higher growth

rate (RGR) but a lower consumption efficiency (ECI) than non-starved larvae,

irrespective of their parental starvation treatment. Since we conducted the

consumption assays for all larvae at the same developmental stage, we were able to

disentangle the effect of starvation from effects due to differences in ontogeny

(Nicieza and Álvarez 2009), in contrast to earlier studies on this sawfly (Paul et al.

2019). Offspring that experienced starvation exhibited a steeper RGR than non-

starved larvae, suggesting compensatory growth (Hector and Nakagawa 2012).

Combining this compensatory growth with a longer development period potentially

allows starved individuals to become larger and to overcome size-based selection,

which can be an important determinant of individual fitness for many insect species

(Beukeboom 2018). While such compensatory growth has been seen in many

species, it can also pose consequent costs to individuals (Arendt et al. 2001;

Dmitriew and Rowe 2007; Auer et al. 2010). We found that individuals that were

starved had a lower ECI than non-starved individuals, which may suggest a

physiological cost of compensatory growth. For the RCR we found differences in

responses between sexes, with female offspring consuming different amounts of leaf

material depending on their parental starvation treatment, while male RCR was solely

influenced by body mass. Considering the larger body mass of female offspring from

starved parents, we would have expected such individuals to consume more leaf

material, although the opposite pattern was observed. Adjustments in consumption

patterns to different starvation regimes have been found to be expressed in the next

generation (McCue et al. 2017). However, patterns may differ depending on the

developmental stage in which an individual is facing starvation, which needs to be

further explored in *A. rosae*.

Larval starvation led to an additional larval instar in *A. rosae* of both sexes. For this

sawfly species, usually 6 instars for female and 5 instars for males are found (Sawa

et al. 1989). Increasing the number of larval instars is potentially an adaptive strategy

that allows individuals to recover from starvation via the prolongation of

developmental time, e.g. by catch-up growth (Hector and Nakagawa 2012). Such

intraspecific variation in the number of larval instars is common in insect species and

can occur in response to various biotic and abiotic factors such as injuries, food

availability and quality, temperature, and humidity (Esperk et al. 2007). Intraspecific

variability in instar numbers in response to food availability and quality can be in the

range of one to three instars, including tenthredinid species (Esperk et al. 2007), but

has not been described, to our knowledge, in the tenthredinid species *A. rosae*.

In contrast to our expectation, clerodanoid uptake, which is known to enhance

mate attractiveness in *A. rosae* (Amano et al. 1999; preprint Paul and Müller 2021),

had no effect on the measured traits in offspring of both sexes. This is surprising,

because partner attractiveness has been shown to affect investment into offspring

or/and offspring traits across a wide variety of species (Robart and Sinervo 2019),

including also multigenerational consequences (Gilbert et al. 2012; Wilson et al.

2019). In other species, mating with more attractive partners has mostly direct effects

for the partner, but not necessarily for the offspring. For example, in the field cricket,

Gryllus firmus, mating with more attractive males led to a higher number of eggs laid

by females but offspring did not show any fitness benefits (Kelly and Adam-Granger

2020). Instead other partner traits, such as body size, may influence offspring

(Booksmythe et al. 2017; Young and Long 2020), which needs further testing in *A.*

rosae. Alternatively, parental clerodanoid exposure may affect other traits, such as

immunity (Bozov et al. 2015), lifespan (Zanchi et al. 2021), or traits exhibited in

adulthood, e.g. mating success (Amano et al. 1999; preprint Paul and Müller 2021).

Parental effects that occur via differential allocation may also be context-dependent.

Transcriptome analysis of male larvae of the F4 generation also revealed stronger

intra- than intergenerational effects of larval starvation on gene expression, in line

with our hypothesis. If F4 larvae were starved, about half of the genes were

differentially expressed compared to non-starved larvae. As typical stress response

indicators, genes potentially encoding heat shock proteins and cytochrome P450s

were differentially expressed, being partly up- and partly downregulated in *A. rosae* in

response to starvation. Regulation of heat shock proteins in response to starvation

has been reported elsewhere, including larvae of Lepidoptera and Hymenoptera

(Farahani et al. 2020; Wang et al. 2012). These proteins act to stabilize and protect

other proteins in the face of both abiotic and biotic stresses (Sørensen et al. 2003;

Farahani et al. 2020). Cytochrome P450s are best known for their role in metabolism

of xenobiotics by insects (Feyereisen 2012). However, they were also found to be

more highly expressed in *D. melanogaster* flies selected for starvation resistance

compared to controls (Doroszuk et al. 2012). In addition, genes putatively encoding

the monoamines octopamine and tyramine were upregulated in starved larvae of *A.*

rosae, both of which play a key role in modulating behavioral and physiological

processes in invertebrates and can be involved in starvation resistance (Li et al.

2016).

Furthermore, starvation of the offspring led to differential gene expression of

pathways involved in metabolism, including pancreatic secretion, which is also found

in starved larvae of the cotton bollworm, *Helicoverpa armigera* (Lepidoptera:

Noctuidae) (Jiang et al. 2019). In starved larvae of *A. rosae*, differential regulation of

metabolic pathways may also explain why these larvae showed an overall lower ECI

compared to non-starved larvae. Expression of metabolic genes likely also changes

with the duration of starvation. For example, in *D. melanogaster* a 16 h starvation

period led to a downregulation particularly of the endopeptidase gene family

(Moskalev et al. 2015), but extended periods of starvation may also induce an

activation of proteolysis genes, as found in different microorgansims (Nakashima et

al. 2006; Michalik et al. 2009). However, little is known about long-term effects,

including cross-generational, on gene expression patterns in response to starvation.

Moreover, many of the genes differentially expressed in starved vs non-starved

individuals of *A. rosae* linked to the mitochondria, reflecting wholescale disruption of

homeostasis caused by starvation (Gilbert 2012).

We found that parental starvation also caused changes in differential gene

expression of about 1 % of the genes of their offspring, indicating that subtle

intergenerational imprints on gene expression can occur. Ribosomal pathway genes

were enriched among downregulated genes when parents were starved vs non-

starved but offspring was not starved and upregulated when offspring were starved.

These results indicate that physiological buffering differs in dependence of the

parental as well as offspring starvation treatment. Larvae that were starved but

whose parents were not starved may have suffered more than starved larvae from

starved parents, as indicated by the upregulation of putative ribosomal pathway

genes. Differential expression of genes of the ribosomal pathway has been observed

in a wide variety of taxa in response to thermal stress (Paraskevopoulou et al. 2020;

Schwanz et al. 2020; Srikanth et al. 2020), demonstrating the broad role of this

pathway in stress response. Differential expression of genes involved in stress and

other physiological and metabolic responses may have energetic costs, which may

lead to a prolonged development time of starved larvae in *A. rosae*.

Alteration in ribosomal expression may also be an important mechanism mediating

intergenerational effects of starvation (Aldrich and Maggert 2015; Bughio and

Maggert 2019). In *Caenorhabditis elegans* starvation led to the generation of small

RNAs, and small RNA-induced gene silencing persisted over up to three generations

(Rechavi et al. 2014). Such mechanisms may also explain the intergenerational

changes in gene expression observed in *A. rosae* in the present study; future work

comparing small RNA expression in relation to starvation could address this. Further

work could also utilize tissue-specific gene expression to determine, where gene

expression changes are most prominent. For example, larvae of the soldier fly,

Inopus falvus (Diptera: Stratiomyidae), differentially express numerous genes in the

salivary glands in response to starvation stress (Etebari et al. 2020). Other tissues of

interest may be different sections of the gut.

In summary, our findings highlight that intragenerational starvation effects were

somewhat stronger than the intergenerational effects of starvation on various traits

investigated here, including life-history, consumption, and gene expression patterns.

Shaping of offspring occurred via parental effects including match-mismatch and

parental buffering, while there was no clear evidence for silver spoon effects. The

parental reproductive environment left no signature on the measured offspring traits.

Overall, due to different trajectories and environments experienced during different

stages, distinct niches are realized, which are expressed in a diversity of phenotypes.

In terms of wider consequences these results suggest that periods of starvation may

make individuals even more robust when facing another food shortage in the next

generation and may thus lead to enhanced damage of crops.

Acknowledgments

This study was funded by the German Research Foundation (DFG) as part of the

SFB TRR 212 (NC³), project number 396777467 (granted to CM). We also thank

University of Potsdam AG Genetics and AG Evolutionary Adaptive Genomics for use

of computational resources and to Dr Tobias Busche and Katrin Lehmann for

assistance with RNA extraction.

Data availability

All data in this manuscript will be deposited online on Zenodo.

Author contributions

Conceptualization and funding acquisition: CM; methods development/experimental

design: SCP, CM; data collection: SCP, data validation and analysis: life history data:

SCP, consumption assay data: SCP, PS, gene expression data: SCP, ABD; data

visualization: SCP, PS; writing original draft: SCP, CM, PS; reviewing and editing:

SCP, PS, ABD, CM.

Supplemental Material

S1. Effects of Starvation on Life-History Traits in F3 Individuals

S2. KEGG Term Analysis

S3. Results of Posthoc Analyses for F4 Generation

S4: Influence of Parental and Offspring Larval Starvation Treatments on

Developmental Times of *Athalia rosae* in F4 Generation

S5. Percentage of Differentially Expressed Genes.

S6. Numbers of Differentially Expressed Genes.

Literature Cited

Aldrich, J. C., and K. A. Maggert. 2015. Transgenerational inheritance of diet-induced
genome rearrangements in *Drosophila*. Plos Genetics 11:e1005148.
Amano, T., R. Nishida, Y. Kuwahara, and H. Fukami. 1999. Pharmacophagous acquisition of
cleodendrins by the turnip sawfly (*Athalia rosae ruficornis*) and their role in the mating
behavior. Chemoecology 9:145-150.
Andrews, S. 2010. FastQC: A Quality Control Tool for High Throughput Sequence Data.
Available online at http://www.bioinformatics.babraham.ac.uk/projects/fastqc.
Arendt, J., D. S. Wilson, and E. Stark. 2001. Scale strength as a cost of rapid growth in
sunfish. Oikos 93:95-100.
Auer, S. K., J. D. Arendt, R. Chandramouli, and D. N. Reznick. 2010. Juvenile compensatory
growth has negative consequences for reproduction in Trinidadian guppies (*Poecilia*
reticulata). Ecology Letters 13:998-1007.
Badyaev, A. V., and T. Uller. 2009. Parental effects in ecology and evolution: mechanisms,
processes and implications. Philosophical Transactions of the Royal Society B-Biological
Sciences 364:1169-1177.
Barnett, D. W., E. K. Garrison, A. R. Quinlan, M. P. Stromberg, and G. T. Marth. 2011.
BamTools: a C++ API and toolkit for analyzing and managing BAM files. Bioinformatics
27:1691-1692.
Benjamini, Y., and Y. Hochberg. 1995. Controlling the false discovery rate: a practical and
powerful approach to multiple testing. Journal of the Royal Statistical Society B 57:289-
300.
Beukeboom, L. W. 2018. Size matters in insects - an introduction. Entomologia
Experimentalis et Applicata 166:2-3.
Boggs, C. L., and K. Niitepõld. 2016. Effects of larval dietary restriction on adult morphology,
with implications for flight and life history. Entomologia Experimentalis et Applicata
159:189-196.
Bolger, A. M., M. Lohse, and B. Usadel. 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 30:2114-2120.
Booksmythe, I., B. Mautz, J. Davis, S. Nakagawa, and M. D. Jennions. 2017. Facultative
adjustment of the offspring sex ratio and male attractiveness: a systematic review and
meta-analysis. Biological Reviews 92:108-134.
Bozov, P., T. Girova, N. Prisadova, Y. Hristova, and V. Gochev. 2015. Antimicrobial activity
of neo-clerodane diterpenoids isolated from Lamiaceae species against pathogenic and
food spoilage microorganisms. Natural Product Communications 10:1797-1800.
Bughio, F., and K. A. Maggert. 2019. The peculiar genetics of the ribosomal DNA blurs the
boundaries of transgenerational epigenetic inheritance. Chromosome Research 27:19-30.
Burton, T., and N. B. Metcalfe. 2014. Can environmental conditions experienced in early life
influence future generations? Proceedings of the Royal Society B-Biological Sciences
281:20140311.
Castagneyrol, B., X. Moreira, and H. Jactel. 2018. Drought and plant neighbourhood
interactively determine herbivore consumption and performance. Scientific Reports
8:5930.
Cornwallis, C. K., and E. A. O'Connor. 2009. Sperm: seminal fluid interactions and the
adjustment of sperm quality in relation to female attractiveness. Proceedings of the Royal
Society B-Biological Sciences 276:3467-3475.
Crawley, M. J. 2012, The R book, John Wiley & Sons.
Cunningham, E. J. A., and A. F. Russell. 2000. Egg investment is influenced by male
attractiveness in the mallard. Nature 404:74-77.

Dmitriew, C., and L. Rowe. 2007. Effects of early resource limitation and compensatory
growth on lifetime fitness in the ladybird beetle *(Harmonia axyridis)*. Journal of
Evolutionary Biology 20:1298-1310.
Dobin, A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut et al. 2013.
STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29:15-21.
Doroszuk, A., M. J. Jonker, N. Pul, T. M. Breit, and B. J. Zwaan. 2012. Transcriptome
analysis of a long-lived natural *Drosophila* variant: a prominent role of stress- and
reproduction-genes in lifespan extension. BMC Genomics 13:167.
English, S., and A. M. G. Barreaux. 2020. The evolution of sensitive periods in development:
insights from insects. Current Opinion in Behavioral Sciences 36:71-78.
Engqvist, L., and K. Reinhold. 2016. Adaptive trans-generational phenotypic plasticity and
the lack of an experimental control in reciprocal match/mismatch-experiments. Methods in
Ecology & Evolution 7:1482-1488.
Esperk, T., T. Tammaru, and S. Nylin. 2007. Intraspecific variability in number of larval
instars in insects. Journal of Economic Entomology 100:627-645.
Etebari, K., K. R. Lindsay, A. L. Ward, and M. J. Furlong. 2020. Australian sugarcane soldier
fly's salivary gland transcriptome in response to starvation and feeding on sugarcane
crops. Insect Science 27:708-720.
Farahani, S., A. R. Bandani, H. Alizadeh, S. H. Goldansaz, and S. Whyard. 2020. Differential
expression of heat shock proteins and antioxidant enzymes in response to temperature,
starvation, and parasitism in the Carob moth larvae, *Ectomyelois ceratoniae* (Lepidoptera:
Pyralidae). Plos One 15:e0228104.
Feyereisen, R. 2012. Insect CYP genes and P450 enzymes, Pages 236–316 *in* I. G.
Lawrence, ed. Insect Molecular Biology and Biochemistry. San Diego, USA, Academic
Press.
Franzke, A., and K. Reinhold. 2013. Transgenerational effects of diet environment on life-
history and acoustic signals of a grasshopper. Behavioral Ecology 24:734-739.
Gilbert, L. I. 2012, Insect Molecular Biology and Biochemistry. Amsterdam, Elsevier.
Gilbert, L., K. A. Williamson, and J. A. Graves. 2012. Male attractiveness regulates daughter
fecundity non-genetically via maternal investment. Proceedings of the Royal Society B-
Biological Sciences 279:523-528.
Grabherr, M. G., B. J. Haas, M. Yassour, J. Z. Levin, D. A. Thompson, I. Amit, X. Adiconis et
al. 2011. Full-length transcriptome assembly from RNA-Seq data without a reference
genome. Nature Biotechnology 29:644-U130.
Groothuis, T. G. G., and B. Taborsky. 2015. Introducing biological realism into the study of
developmental plasticity in behaviour. Frontiers in Zoology 12:(Suppl 1): S6.
Hebets, E. A., J. Wesson, and P. S. Shamble. 2008. Diet influences mate choice selectivity
in adult female wolf spiders. Animal Behaviour 76:355-363.
Hector, K. L., and S. Nakagawa. 2012. Quantitative analysis of compensatory and catch-up
growth in diverse taxa. Journal of Animal Ecology 81:583-593.
Hibshman, J. D., A. Hung, and L. R. Baugh. 2016. Maternal diet and insulin-like signaling
control intergenerational plasticity of progeny size and starvation resistance. Plos
Genetics 12:e1006396.
Hothorn, T., F. Bretz, and P. Westfall. 2008. Simultaneous inference in general parametric
models. Biometrical Journal 50:346-363.
Jiang, T., L. Ma, X. Y. Liu, H. J. Xiao, and W. N. Zhang. 2019. Effects of starvation on
respiratory metabolism and energy metabolism in the cotton bollworm *Helicoverpa*
armigera (Hubner) (Lepidoptera: Noctuidae). Journal of Insect Physiology 119:103951.
Kelly, C. D., and É. Adam-Granger. 2020. Mating with sexually attractive males provides
female *Gryllus firmus* field crickets with direct but not indirect fitness benefits. Behavioral
Ecology and Sociobiology 74:80.
Le Roy, A., I. Loughland, and F. Seebacher. 2017. Differential effects of developmental
thermal plasticity across three generations of guppies (*Poecilia reticulata*): canalization
and anticipatory matching. Scientific Reports 7:4313.
Li, B., and C. N. Dewey. 2011. RSEM: accurate transcript quantification from RNA-Seq data
with or without a reference genome. BMC Bioinformatics 12:323.
Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth et al. 2009. The
Sequence Alignment/Map format and SAMtools. Bioinformatics 25:2078-2079.

Li, Y., J. Hoffmann, F. Stephano, I. Bruchhaus, C. Fink, and T. Roeder. 2016. Octopamine
controls starvation resistance, life span and metabolic traits in *Drosophila*. Scientific
Reports 6:35359.
Love, M. I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biology 15:550.
Luo, W. J., and C. Brouwer. 2013. Pathview: an R/Bioconductor package for pathway-based
data integration and visualization. Bioinformatics 29:1830-1831.
Luo, W. J., M. S. Friedman, K. Shedden, K. D. Hankenson, and P. J. Woolf. 2009. GAGE:
generally applicable gene set enrichment for pathway analysis. BMC Bioinformatics
10:161.
McCue, M. D., J. S. Terblanche, and J. B. Benoit. 2017. Learning to starve: impacts of food
limitation beyond the stress period. Journal of Experimental Biology 220:4330-4338.
Michalik, S., M. Liebeke, D. Zuhlke, M. Lalk, J. Bernhardt, U. Gerth, and M. Hecker. 2009.
Proteolysis during long-term glucose starvation in *Staphylococcus aureus* COL.
Proteomics 9:4468-4477.
Miyatake, T. 2001. Effects of starvation on death-feigning in adults of *Cylas formicarius*
(Coleoptera: Brentidae). Annals of the Entomological Society of America 94:612-616.
Monaghan, P. 2008. Early growth conditions, phenotypic development and environmental
change. Philosophical Transactions of the Royal Society B-Biological Sciences 363:1635-
1645.
Moriya, Y., M. Itoh, S. Okuda, A. C. Yoshizawa, and M. Kanehisa. 2007. KAAS: an automatic
genome annotation and pathway reconstruction server. Nucleic Acids Research 35:W182-
W185.
Moskalev, A., S. Zhikrivetskaya, G. Krasnov, M. Shaposhnikov, E. Proshkina, D.
Borisoglebsky, A. Danilov et al. 2015. A comparison of the transcriptome of *Drosophila*
melanogaster in response to entomopathogenic fungus, ionizing radiation, starvation and
cold shock. BMC Genomics 16(Supple 13):S8.
Naguib, M., A. Nemitz, and D. Gil. 2006. Maternal developmental stress reduces
reproductive success of female offspring in zebra finches. Proceedings of the Royal
Society B-Biological Sciences 273:1901-1905.
Naito, T., and H. Suzuki. 1991. Sex determination in the sawfly, *Athalia rosae* Ruficornis
(Hymenoptera) - occurence of triploid males. Journal of Heredity 82:101-104.
Nakashima, A., T. Hasegawa, S. Mori, M. Ueno, S. Tanaka, T. Ushimaru, S. Sato et al. 2006.
A starvation-specific serine protease gene, isp6(+), is involved in both autophagy and
sexual development in *Schizosaccharomyces pombe*. Current Genetics 49:403-413.
Nicieza, A. G., and D. Álvarez. 2009. Statistical analysis of structural compensatory growth:
how can we reduce the rate of false detection? Oecologia 159:27-39.
Paraskevopoulou, S., A. B. Dennis, G. Weithoff, and R. Tiedemann. 2020. Temperature-
dependent life history and transcriptomic responses in heat-tolerant versus heat-sensitive
Brachionus rotifers. Scientific Reports 10:13281.
Paul, S. C., A. B. Dennis, L. J. Tewes, J. Friedrichs, and C. Müller. 2021. Consequences of
pharmacophagous uptake from plants and conspecifics in a sawfly elucidated using
chemical and molecular techniques. bioRxiv preprint. doi: 10.1101/2021.02.09.430406
Paul, S. C., and C. Müller. 2021. Fighting over defence chemicals disrupts mating behaviour.
bioRxiv preprint. doi: 10.1101/2021.02.12.430958
Paul, S. C., R. Putra, and C. Müller. 2019. Early life starvation has stronger intra-generational
than transgenerational effects on key life-history traits and consumption measures in a
sawfly. Plos One 14:e0226519.
Perez, M. F., and B. Lehner. 2019. Intergenerational and transgenerational epigenetic
inheritance in animals. Nature Cell Biology 21:143-151.
Pilakouta, N., S. Jamieson, J. A. Moorad, and P. T. Smiseth. 2015. Parental care buffers
against inbreeding depression in burying beetles. Proceedings of the National Academy of
Sciences of the United States of America 112:8031-8035.
Ratikainen, II, and H. Kokko. 2010. Differential allocation and compensation: who deserves
the silver spoon? Behavioral Ecology 21:195-200.
Raveh, S., D. Vogt, and M. Kolliker. 2016. Maternal programming of offspring in relation to
food availability in an insect (*Forficula auricularia*). Proceedings of the Royal Society B-
Biological Sciences 283:20152936.

Rechavi, O., L. Houri-Ze'evi, S. Anava, W. S. S. Goh, S. Y. Kerk, G. J. Hannon, and O. Hobert. 2014. Starvation-induced transgenerational inheritance of small RNAs in *C. elegans*. Cell 158:277-287.

Regalado, J. M., M. B. Cortez, J. Grubbs, J. A. Link, A. van der Linden, and Y. Zhang. 2017. Increased food intake after starvation enhances sleep in *Drosophila melanogaster*. Journal of Genetics and Genomics 44:319-326.

Riggert, E. 1939. Untersuchungen über die Rübenblattwespe *Athalia colobri* Christ (*A. spinarum* F.). Zeitschrift für angewandte Entomologie 26:462-516.

Robart, A. R., and B. Sinervo. 2019. Females increase parental care, but not fecundity, when mated to high-quality males in a biparental fish. Animal Behaviour 148:9-18.

Roberts, A., C. Trapnell, J. Donaghey, J. L. Rinn, and L. Pachter. 2011. Improving RNA-Seq expression estimates by correcting for fragment bias. Genome Biology 12:R22.

Roff, D. A. 1992, The evolution of life histories: Theory and analysis. New York, Chapman & Hall.

Saastamoinen, M., N. Hirai, and S. van Nouhuys. 2013. Direct and trans-generational responses to food deprivation during development in the Glanville fritillary butterfly. Oecologia 171:93-104.

Sánchez-Tójar, A., M. Lagisz, N. P. Moran, S. Nakagawa, D. W. A. Noble, and K. Reinhold. 2020. The jury is still out regarding the generality of adaptive 'transgenerational' effects. Ecology Letters 23:1715-1718.

Sawa, M., A. Fukunaga, T. Naito, and K. Oishi. 1989. Studies on the sawfly, *Athalia rosae* (Insecta, Hymenoptera, Tenthredinidae) I. General biology. Zoological Science 6:541-547.

Schwanz, L. E., J. Crawford-Ash, and T. Gale. 2020. Context dependence of transgenerational plasticity: the influence of parental temperature depends on offspring environment and sex. Oecologia 194:391-401.

Smith, C. C., and S. D. Fretwell. 1974. Optimal balance between size and number of offspring. American Naturalist 108:499-506.

Soneson, C., M. Love, and M. Robinson. 2015. Differential analyses for RNA-seq: transcript-level estimates improve gene level inferences F1000Research 4:1521.

Sørensen, J. G., T. N. Kristensen, and V. Loeschcke. 2003. The evolutionary and ecological role of heat shock proteins. Ecology Letters 6:1025-1037.

Srikanth, K., J.-E. Park, S. Y. Ji, K. H. Kim, Y. K. Lee, H. Kumar, M. Kim et al. 2020. Genome-wide transcriptome and metabolome analyses provide novel insights and suggest a sex-specific response to heat stress in pigs. Genes 11:540.

Taborsky, B. 2017. Developmental plasticity: preparing for life in a complex world. Advances in the Study of Behavior 49:49-99.

Tarka, M., A. Guenther, P. T. Niemelä, S. Nakagawa, and D. W. A. Noble. 2018. Sex differences in life history, behavior, and physiology along a slow-fast continuum: a meta-analysis. Behavioral Ecology and Sociobiology 72:132.

Uller, T., S. Nakagawa, and S. English. 2013. Weak evidence for anticipatory parental effects in plants and animals. Journal of Evolutionary Biology 26:2161-2170.

Valtonen, T. M., K. Kangassalo, M. Pölkki, and M. J. Rantala. 2012. Transgenerational effects of parental larval diet on offspring development time, adult body size and pathogen resistance in *Drosophila melanogaster*. PloS One 7:e31611.

van de Pol, M., L. W. Bruinzeel, D. Heg, H. P. van der Jeugd, and S. Verhulst. 2006. A silver spoon for a golden future: long-term effects of natal origin on fitness prospects of oystercatchers (*Haematopus ostralegus*). Journal of Animal Ecology 75:616-626.

Vega-Trejo, R., M. D. Jennions, and M. L. Head. 2016. Are sexually selected traits affected by a poor environment early in life? BMC Evolutionary Biology 16:263.

Wang, H., K. Li, J. Y. Zhu, Q. Fang, G. Y. Ye, H. Wang, K. Li et al. 2012. Cloning and expression pattern of heat shock protein genes from the endoparasitoid wasp, *Pteromalus puparum* in response to environmental stresses. Archives of Insect Biochemistry and Physiology 79:247-263.

Wang, Y., O. Kaftanoglu, C. S. Brent, R. E. Page, and G. V. Amdam. 2016. Starvation stress during larval development facilitates an adaptive response in adult worker honey bees (*Apis mellifera* L.). Journal of Experimental Biology 219:949-959.

Wilson, K. M., A. Tatarenkov, and N. T. Burley. 2019. Early life and transgenerational stressors impact secondary sexual traits and fitness. Behavioral Ecology 30:830-842.

Wolf, J. B., and M. J. Wade. 2009. What are maternal effects (and what are they not)?
Philosophical Transactions of the Royal Society B-Biological Sciences 364:1107-1115.
Yin, J. J., M. Zhou, Z. R. Lin, Q. S. Q. Li, and Y. Y. Zhang. 2019. Transgenerational effects
benefit offspring across diverse environments: a meta-analysis in plants and animals.
Ecology Letters 22:1976-1986.
Young, Y., and T. A. F. Long. 2020. Nutritional developmental history and its consequences
for reproductive success in *Drosophila suzukii* (Diptera: Drosophilidae). Canadian
Entomologist 152:490-515.
Zanchi, C., L. K. Lo, R. R, I. Moritz, J. Kurtz, and C. Müller. 2021. Survival of the sawfly
Athalia rosae upon infection by an entomopathogenic fungus and in relation to
clerodanoid uptake. Frontiers in Physiology.
Zhu, A. Q., J. G. Ibrahim, and M. I. Love. 2019. Heavy-tailed prior distributions for sequence
count data: removing the noise and preserving large differences. Bioinformatics 35:2084-
2092.
Zizzari, Z. V., N. M. van Straalen, and J. Ellers. 2016. Transgenerational effects of nutrition
are different for sons and daughters. Journal of Evolutionary Biology 29:1317-1327.

Figures

Figure 1:

Figure 1: Full experimental design showing outline of breeding structure and

experimental treatments from F0 to F4 generation in *Athalia rosae*; * indicates

individuals taken for RNASeq analysis. Sample sizes split by sex are given in boxes

for F3 and F4 generation (numbers in brackets refer to sample sizes for consumption

assay).

Figure 2

Figure 2: Number of larval instars in F4 in dependence of offspring starvation

treatment (blue: non starved, red: starved) and influence of offspring larval starvation

treatments (N = no starvation, S = starvation) and parental larval starvation

treatments (blue solid line = no starvation, red dashed line = starvation) on offspring

total development time (C, D) and adult body mass (E, F) of *Athalia rosae*. Data are

860   plotted separately for females (A, C, E) and males (B, D, F); individuals were sexed

861   on emergence. Points are model predictions with associated confidence intervals and

862   colors of points and lines correspond to parental starvation treatment. Raw data are

863   plotted in transparent colors in the background.

Figure 3

Figure 3: Influence of offspring (blue line = no starvation, red line = starvation) and

parental larval starvation treatments (solid line = no starvation, dashed line =

starvation) in F4 on relative growth rate, RGR (A, B); relative consumption rate, RCR

(C, D); and efficiency of conversion of ingested food, ECI (E, F) of *Athalia rosae*.

Note that neither offspring starvation treatment nor parental starvation treatment had

an effect on RCR in males. Data are plotted separately for females (A, C, E) and

males (B, D, F); individuals were sexed on emergence. Lines depict model

predictions with shaded regions showing the 95% confidence interval range. Raw

data representing offspring larval starvation treatment (blue = no starvation, red =

starvation) and parental larval starvation treatment (open circles = no starvation, filled

circles = starvation) are plotted in the background.

Figure 4

Figure 4: Venn diagram illustrating the number of unique and shared differentially expressed genes in male offspring larvae (F4) of *Athalia rosae* resulting from each of the four pairwise comparisons, (N = no starvation and S = starvation; left box F3, right box F4).

Figure 5

Figure 5: Heatmaps showing the expression (normalized counts) of genes in male

offspring larvae (F4) of *Athalia rosae* across all samples for those genes that were

differentially expressed when offspring starvation treatment differed (A, B) or parental

starvation treatment differed (C, D), (N = no starvation and S = starvation; left box F3,

right box F4). Plotted values are *z*-scores computed from normalized counts post

clustering.

Table 1. Significantly differentially expressed KEGG gene sets/pathways male offspring larvae (F4) of *Athalia rosae* that differed in either their own or their parent's (F3) larval starvation regime (N = no starvation and S = two periods of starvation for 24 hours during larval development; left box F3, right box F4). The combination listed first (on top) is the 'treatment' the one listed second (underneath) is the 'control' in each comparison, meaning that pathways are up- or downregulated in 'treatment' relative to 'control'. Larvae undergoing starvation (S) were subjected to two periods of starvation for 24 hours during larval development, one in L2 and one in L4.

KEGG pathway	Offspring starvation treatment differs			Parental starvation treatment differs	
	N S vs N N	S N vs S S		S N vs N N	N S vs S S
Ribosome	↑ up	↓ down		↓ down	↑ up
Pancreatic secretion	-	↓ down		-	-
Protein processing in endoplasmic reticulum	↓ down	↑ up		-	-
Proteasome	↓ down	↑ up		-	-
Phagosome	-	↑ up		-	-
Citrate (TCA) cycle	↓ down	-		-	-
Glycolysis / Gluconeogenesis	↓ down	-		-	-
Antigen processing and presentation	↓ down	↑ up		-	-
Protein export	↓ down	-		-	-
ECM-receptor interaction	-	↑ up		-	-
Thyroid hormone synthesis	-	↑ up		-	-