Dosage sensitivity in Pumilio1-related phenotypes reflects distinct disease mechanisms

Salvatore Botta[1,2], Nicola de Prisco[1], Alexei Chemiakine[1], Maximilian Cabaj[1], Purvi Patel[3], Ella Doron-

Mandel[4], Colton J. Treadway[5], Marko Jovanovic[4], Nicholas G. Brown[5], Rajesh K. Soni[3], and Vincenzo A.

Gennarino[1,6,7,8,*]

[1]Department of Genetics and Development, Columbia University Irving Medical Center, New York, NY.

[2]Department of Translational Medical Science, University of Campania Luigi Vanvitelli, Caserta, Italy.

[3]Proteomics and Macromolecular Crystallography Shared Resource, Herbert Irving Comprehensive

Cancer Center, Columbia University Irving Medical Center, New York, NY, USA.

[4]Department of Biological Sciences, Columbia University, New York, NY, USA.

[5]Department of Pharmacology and Lineberger Comprehensive Cancer Center, University of North

Carolina School of Medicine, Chapel Hill, NC, USA.

[6]Departments of Pediatrics and Neurology, Columbia University Irving Medical Center, New York, NY.

[7]Columbia Stem Cell Initiative, Columbia University Irving Medical Center, New York, NY.

[8]Initiative for Columbia Ataxia and Tremor, Columbia University Irving Medical Center, New York, NY.

[*]Corresponding author: Tel: +1 212-305-7863; Email: vag2138@cumc.columbia.edu

Abstract

Mutations in the RNA-binding protein (RBP) Pumilio1 (PUM1) can cause dramatically different

phenotypes. We previously noted that phenotypic severity tracked with protein dosage: a mild mutation

that reduces PUM1 levels by 25% causes late-onset ataxia, whereas PUM1 haploinsufficiency causes

developmental delay and seizures. Why this difference in expression should cause such different

phenotypes has been unclear: PUM1 targets are de-repressed to equal degrees in both cases, and the more

severe mutation does not hinder PUM1's RNA-binding ability. We therefore developed a PUM1

interactome in the murine brain. We find that mild PUM1 loss de-represses PUM1-specific targets, but

PUM1 haploinsufficiency disrupts several interactors and regulation of their targets. We validated these

phenomena in patient-derived cell lines and show that normalizing PUM1 levels restores interactors and

their targets to proper levels. We therefore propose that dosage sensitivity does not necessarily reflect a

linear change in protein abundance but can involve distinct mechanisms. Studying the interactors of RBPs

in vivo will be necessary to understand their functions in neurological diseases.

Keywords: Pumilio/PADDAS/PRCA/protein interactome/RNA neurobiology

Introduction

35
RNA-binding proteins (RBPs) modify the proteome at the post-transcriptional level, regulating RNA

localization, transport, translation, splicing, and decay; they have been found to orchestrate hundreds of

pathways that are responsible for proper biological functions (Hentze *et al*, 2018; Keene, 2007b). As our

understanding of RBPs has grown over the past decade, it has become clear that their functions are

particularly important in neurons, whose synaptic plasticity demands rapid local responses to stimuli

(Mauger *et al*, 2016). Several complex neurological disorders—e.g., Fragile X syndrome, Fragile X-

associated tremor and ataxia syndrome, and amyotrophic lateral sclerosis—are known to involve

disruptions in the function of RBPs, which has understandably led to considerable interest in mapping

their neuronal targets (Darnell & Richter, 2012; Khalil *et al*, 2018; Ravanidis *et al*, 2018).

Our work on Pumilio1 (PUM1), however, has revealed that the targets may not provide a full picture

of the RBP's functioning. After discovering that PUM1 is important for mouse neurobiology (Gennarino

et al, 2015), we searched human databases for individuals with mutations in *PUM1*. We initially found 13

individuals with deletions or missense mutations, and their phenotypes tracked with protein dosage. A

mild mutation (T1035S) that reduces PUM1 levels by 25% causes a slowly progressive, pure ataxia with

onset in mid-life, which we called PUM1-related cerebellar ataxia or PRCA. Deletions or the most severe

missense mutation (R1147W) reduce PUM1 levels by ~50% and cause a neurodevelopmental syndrome

we called PUM1-associated developmental delay and seizures, or PADDAS (Gennarino *et al*, 2018). Both

the infantile and adult-onset phenotypes are now considered to be forms of spinocerebellar ataxia type 47

(SCA47). T1035S lies within a highly conserved RNA-binding domain and impairs RNA binding, so it

was not surprising to find known PUM1 targets upregulated in patient cells. The puzzle came with

R1147W, which lies outside this domain and does not impair target regulation, yet deregulates the same

targets to the same degree as T1035S (Gennarino *et al.*, 2018). Moreover, the R1147W phenotype is quite

severe, closer to that of the null mice than the heterozygous mice. Given that RBPs can interact with,

influence, or compete with each other (Dassi, 2017), we hypothesized that R1147W disrupts PUM1's

ability to interact with its native partners, which would then lead to de-repression of the targets of these

complexes and not just the direct targets of PUM1.

Testing this hypothesis required us to identify PUM1 interactors in the mouse brain. Although a great

deal is known about the PUMILIO/FBF (PUF) family of RBPs (Bohn *et al*, 2018; Friend *et al*, 2012;

Goldstrohm *et al*, 2018; Kedde *et al*, 2010; Miles *et al*, 2012; Temme *et al*, 2014; Uyhazi *et al*, 2020; Van

Etten *et al*, 2012; Weidmann *et al*, 2014), of which PUM1 and its homolog PUM2 are members, little is

known about PUM1 function in the postnatal mammalian brain. Protein interactions in general, and those

of PUF (Pumilio and FBF) family members in particular, can be organism-, transcript-, and even

condition-specific (Marrero *et al*, 2011). We therefore took an unbiased approach by using *in vivo*

proteomics to identify PUM1's native partners in the mouse brain. We then studied the effect of PUM1

insufficiency on a subset of RBP interactors in *PUM1* heterozygous and null mice and cell lines from

patients bearing either the T1035S or R1147W mutation. The results support our hypothesis that two

different mechanisms underlie PRCA and PADDAS, underscoring the need to examine an RBP's

interactions as well as identify its targets.

Results

Establishing the PUM1 interactome in the adult mouse brain

We performed co-immunoprecipitation (IP) on brains from 10-week-old wild-type (WT) mice

followed by liquid chromatography with tandem mass spectrometry (LC-MS/MS); we used IgG as a

negative control (**Figure S1A**). Principal component analysis (PCA) separated IP-IgG from IP-Pum1

(**Figure S1B**). No residual Pum1 was detected in brain tissue by post-IP western blot (**Figure S1C**). (In

this paper, we will differentiate between human and mouse proteins by using all-caps only for the

former.) Considering only those candidates that had at least two unique peptides in at least five out of six

IP-Pum1 samples, this approach identified 234 putative interactors (**Figure S1C** and **Table S1**) that we

clustered into 10 functional groups using protein-protein interaction data from CORUM and the Human

Protein Atlas (Raudvere *et al*, 2019) (**Figure 1**).

Our list included several proteins that had been previously found to interact with Pum1, as well as

new members of protein complexes or families that interact with Pum1 in other contexts. For example,

our LC-MS/MS identified Fmrp (cluster 7), which associates with Pum1 in neural progenitor cells (Zhang

et al, 2017). We also identified Cnot1, the central scaffold of the CCR4-NOT complex (Enwerem *et al*,

2021; Van Etten *et al.*, 2012), which is recruited by Pum1 to shorten poly(A) tails and promote mRNA

degradation (Temme *et al.*, 2014; Van Etten *et al.*, 2012; Weidmann *et al.*, 2014). Translation initiation

factors (cluster 6) cooperate with PUF proteins in invertebrates (Blewett & Goldstrohm, 2012).

Drosophila studies showed that Pumilio partners with different proteins in different neuronal types

(Muraro *et al*, 2008), so we repeated the LC-MS/MS experiments in three brain regions that express

abundant Pum1: the cerebellum, hippocampus, and cortex (Gennarino *et al.*, 2015). PCA readily

separated Pum1 and IgG samples (**Figure S2A**). This analysis identified 854 putative Pum1 interactors in

the cerebellum, 423 in the hippocampus, and 598 in the cortex (**Figure 2A**, **Figure S2B-D** and **Table S1**).

489 were unique to the cerebellum, 145 to the hippocampus, 247 to the cortex, and 48 unique to the rest

of the brain (i.e., excluding these three regions). Only 82 candidates appeared in all three brain regions

and the whole brain (**Figure 2A**, *yellow dots*).

To determine whether these results merely reflect regional differences in protein abundance, we

performed quantitative proteomics in 10-week-old whole brain, cerebellum, cortex, and hippocampus.

Pum1 interactors in general were not the most abundant proteins in these samples (**Figure S3A, B**). For

example, Pum2, which turned up in only 3 out of 6 of the whole-brain samples, was one of the strongest

Pum1 interactors in the cortex (**Figure 2A, S3C,** and **Table S1**). It is expressed at low levels in all brain

regions examined (**Figure S3D**). These findings suggest that the candidate interactors represent biological

information rather than post-lysis effects or high abundance.

Components of the APC/C and mTOR pathways were consistent across the three brain regions

(**Figure 2A, B**, and **Table S1**), but the list of Pum1 interactors in several other pathways was expanded by

this regional analysis (**Figure 2**). For example, although Cnot1 turned up in all three brain regions

(**Figure S3C**), Cnot2 appeared only in cortex and cerebellum, and Cnot10 only in the cortex (**Figure 2A**).

There were many putative interactors involved in translation initiation, with Eif3b showing up in both the

hippocampus and the whole brain (**Figure 2A**). The splicing factors showed regional specificity in

interactions: Rbfox1 was restricted to the cortex, while Rbfox2 and Rbfox3 were in the cerebellum,

hippocampus, and whole brain (**Figure 2A**). This is consistent with previous work showing that Rbfox1

mediates cell-type-specific splicing in cortical interneurons (Wamsley *et al*, 2018) and that Rbfox2 is

needed for cerebellar development (Gehman *et al*, 2012).

DAVID Gene Ontology analysis revealed that the main functional categories across the three regions

were ubiquitin ligases (anaphase-promoting complex [APC/C], E2/E3 and Kll-linked ubiquitin) and RBPs

involved in various aspects of RNA metabolism (RNA silencing, 3'UTR binding, mRNA stability,

transport, and splicing) (**Figure 2B** and **Table S2**). To test some of these interactions biochemically, we

tried to purify recombinant proteins, but many of the proteins or their complexes are either quite large

(>100kD) or strongly tended to aggregate. We were able pull down PUM1 by recombinant GST-tagged

FMRP (**Figure S4A**) and GST-tagged PUM2 (**Figure S4B**), confirming their direct interactions (**Figure**

S4). We also used IP and co-IP to spot-check key candidates from the anaphase promoting complex or

cyclosome (APC/C, cluster 1: Anapc1, Fzr1) and mTOR (cluster 10: Rptor, Cad) pathways (**Figure S3C,**

S4C, and **D**).

Realizing we would need to select certain proteins for further *in vivo* study, we decided to prioritize

the RBPs in cluster 7 (**Figure 1**): Fmrp and Ago2 (involved in RNA silencing), Cnot1 (mRNA

deadenylase protein), Rbfox3 (alternative splicing factor), and Pum2. We reasoned as follows. First, RBP-

RBP interactions are biologically important, and RNA-related categories were prominent in the gene

ontology analyses for both whole brain and all three brain regions. Second, this cluster was also the most

highly interconnected with other clusters and likely to influence their activities. Third, these RBPs have

been well studied, which would enable us to more readily test the consequences of PUM1 loss.

Nevertheless, these proteins have been studied mostly *in vitro* and, with the exception of Fmrp and Pum2,

have not been previously associated with Pum1 in the murine brain, so they might shed more light on

Pum1 biology.

Pum1 associates with Fmrp, Ago2, Cnot1, and Pum2 in the absence of mRNA

We confirmed Pum1 associations with Pum2, Fmrp, Ago2, Rbfox3, and Cnot1 by co-IP followed by

western blot (**Figure 3A**, *left panel*). We also tested for murine Mov10, which binds both Fmrp and Ago2

in vitro (Kenny *et al*, 2020; Kenny *et al*, 2014), and found it pulled down with Pum1, likely in concert

with Fmrp (**Figure 3A**). We co-IP'd Pum1 and blotted for all six RBPs in *Pum1$^{-/-}$* mouse brains; none

were detected (**Figure S5A**). We then tested other proteins associated with the RNA silencing machinery

that did not appear in our LC-MS/MS data, such as Ago1 and Ago3, and we found no interactions

(**Figure S5B**).

To exclude the possibility that the co-IP experiments were recovering proteins that co-bound target

RNAs but are not part of the same complex as the protein of interest, we treated mouse brain samples

with RNase and verified that no detectable RNA remained (see Materials and Methods). Pum1 still

associated with Pum2, Fmrp, Ago2, and Cnot1 in the absence of mRNA, but not with Rbfox3 or Mov10

(**Figure 3A**, *right panel*, and **S5C**). We repeated the RNase experiments in HEK293T cells, which

confirmed our results (except for Rbfox3, which was not detectable in these cells) (**Figure 3B**). These

data suggest that Pum1 interacts with Pum2, Fmrp, Ago2, and Cnot1 prior to binding RNA.

Pum1 loss alters RBP interactors and miRNA machinery in mouse cerebella by sex

If Pum1 is an important interactor for these six RBPs, loss of Pum1 should affect their stability or

abundance in mouse brain. *Pum1* heterozygous and null mice showed changes in the quantities of Pum2,

Ago2, and Mov10 proteins across the brain (**Figure S6A**), but only *Pum2* showed changes in mRNA

levels (**Figure S6B**). Ago2 and Mov10 levels fell only in male mice (**Figure S6A**), which led us to

quantify Pum2, Fmrp, Ago2, Rbfox3, Cnot1 and Mov10 mRNA and protein in the cerebellum, cortex,

and hippocampus of male and female mice.

As previously reported, Fmrp protein expression was upregulated in all three brain regions in male

null mice, but in female null cerebella Fmrp was almost 70% lower (**Figure 4A**) (Singh *et al*, 2007; Singh

& Prasad, 2008). Pum2 protein levels rose in all three brain regions, as did its mRNA (**Figure 4A-C**, and

Figure S7). Ago2, Rbfox3, and Cnot1 also showed divergent responses to Pum1 loss according to sex

and brain region (**Figure 4A-C**). None of these proteins showed any changes in mRNA levels (**Figure**

S7), despite the fact that *Fmr1* and *Cnot1*, like *Pum2*, have a Pumilio Response Element (PRE) (Zamore

et al, 1997) in their 3'UTR.

Because Pum1 loss has deleterious effects on the cerebellum in both mice (Gennarino *et al.*, 2015)

and humans (Gennarino *et al.*, 2018), and Pum1 interacts closely with the miRNA machinery (Kedde *et*

al., 2010), we decided to investigate the physiological relevance of these sex-specific observations by

examining the effects of divergent Ago2 protein levels on cerebellar miRNAs in male and female mice.

Our miRNA-seq identified 701 expressed miRNAs, many of which diverged in expression between the

two sexes (**Figure S8A, B**). Heatmap clustering of significant miRNA expression in *Pum1^{-/-}* and WT

male and female cerebella at 10 weeks of age revealed that the expression of 166 miRNAs diverged

between the two sexes in parallel with Ago2 expression (**Table S3, Figure 4D**).

To study the expression of downstream targets that are co-bound by those miRNAs in 10-week-old

WT and *Pum1^{-/-}* male and female cerebella, we selected the 49 miRNAs that showed >25% change in

expression in either direction. Using TargetScan and CoMeTa (Agarwal *et al*, 2015; Gennarino *et al*,

2012) we identified 6832 putative targets that are co-bound by at least two miRNA . To reduce this list to

manageable size, we prioritized targets that are co-bound by at least eight miRNAs, for a total of,

coincidentally, 49 putative targets. *Pum1^{-/-}* male and female cerebella showed gene expression changes

for 44 out of these 49 targets that correlated with the sex differences in Ago2 levels (**Figure 4E**, **Figure**

S9 and **Table S4**).

To elucidate the biological pathways in which these miRNAs play an essential role, we performed

David Gene Ontology (GO) with all the non-redundant targets predicted by CoMeTa (Gennarino *et al.*,

2012) and TargetScan 7.1 (Agarwal *et al.*, 2015) that are co-bound by at least four miRNAs (GO needs a

larger pool of genes than 49 but fewer than 3000), 2127 targets (**Figure S10A-C**). Among 2127 targets

analyzed, we found enrichment for multiple categories having to do with synaptic function under "cellular

components." The most enriched categories under "biological processes" were organ growth and post-

embryonic development. Under KEGG pathways, there was a particular enrichment in Wnt signaling,

dopaminergic and cholinergic pathways, cancers, and protein ubiquitination.

We then analyzed the same targets by SynGO (Koopmans *et al*, 2019), which uses single-cell data to

identify genes that are expressed in specific neurons. SynGO pinpointed 117 presynpatic targets and 124

postsynaptic (**Figure S10D**). Among the 166 miRNAs inversely expressed between sexes were the entire

miR-200 family (miR-200a, miR-220b, miR-200c, miR-141, and miR-429), which regulates targets

involved in neurogenesis, glioma, and neurodegenerative diseases (Fu *et al*, 2019; Trumbach & Prakash,

2015). These results are consonant with SCA47 cerebellar deficits and suggest an intimate relation

between Pum1 and Ago2 in mouse cerebellum.

Pum1, Pum2, Fmrp, Ago2, and Rbfox3 share their top targets

If the complexes Pum1 forms with these RBPs are physiologically relevant, as seen for Ago2 in

cerebellum, then they should co-regulate at least some of the same mRNA targets. Indeed, one corollary

of the "regulon theory," which posits that mRNA targets in the same pathway are co-regulated

(Blackinton & Keene, 2014; Keene, 2007a, b; Keene & Lager, 2005), is that there should be a discernible

set of RBPs that do the co-regulating.

To test this hypothesis, we analyzed all the available high-throughput sequencing UV-crosslinking

and immunoprecipitation (HITS-CLIP) data available for the murine brain. These data exist for Fmrp

(Maurin *et al*, 2018), Ago2 (Chi *et al*, 2009), Rbfox3 (Weyn-Vanhentenryck *et al*, 2014), Pum1, and

Pum2 (Zhang *et al.*, 2017)). We then performed gene set enrichment analysis (GSEA) (Subramanian *et al*,

2005) using Fmrp as the basis for comparison (because it has the largest dataset). As negative controls,

we used HITS-CLIP data from mouse brain for the other four available RBPs that did not show up as

Pum1 interactors in our LC-MS/MS: Mbnl2 (Charizanis *et al*, 2012), Nova (Zhang *et al*, 2010), Apc

(Preitner *et al*, 2014), and Ptpb2 (Licatalosi *et al*, 2012).

Although the datasets are not perfectly matched (the mice were different ages in some cases; see

Methods), this analysis nonetheless revealed that Pum1 targets were concentrated in the top 5th percentile

of all Fmrp targets, with an enrichment score (ES) of 0.93 (the maximum is 1) and FDR of 0 (**Figure 5A,**

blue line represents ES). Pum2, Ago2, and Rbfox3 showed nearly identical patterns. There was no

significant overlap between the targets of Fmrp and those of any negative control (Nova had the highest

ES, but this was only 0.36 with a rank max of 45[th] percentile and FDR=1; **Figure 5B**). Neither Pum1 nor

its partner RBP targets were enriched among any of the ranked target lists of the negative controls

(**Figure S11A**).

To ascertain where highest-ranking Fmrp targets fell among the genes with the highest probability of

being Pum1 targets, we divided the Fmrp ranked target list into 10 equal bins according to percentile. We

then repeated GSEA of Pum1 HITS-CLIP data for each bin and found that 648 of the 1194 identified

Pum1 targets (54%) are in the top 10[th] percentile of Fmrp targets, with an ES of 0.8 (**Figure 5C**). This

was also true for Pum2, Ago2, and Rbfox3 (**Figure 5C**).

We then performed the same analysis but instead used the Pum1 target list as the basis for

comparison. We ran GSEA on each of the four Pum1 partners against the list of Pum1 target genes, and

each partner's targets were within the top 20% of the Pum1 list (**Figure 5D**). Specifically, Fmrp's targets

reside in the top 10[th] percentile (with an ES of 0.81), Pum2's targets within the 16[th] percentile (ES=0.9),

Ago2's targets within the 18th percentile (ES=0.76), and Rbofx3's targets within the 19th percentile

(ES=0.67). The four RBPs used here as negative controls have a minimum rank at the 37th percentile, and

the best ES was 0.26 for Apc; none of the five reached statistical significance (**Figure 5D**). These

analyses demonstrate that there is substantial overlap among the highest-ranked targets of Pum1, Pum2,

Fmrp, Ago2, and Rbfox3.

We also studied the targets shared by Pum1, Pum2, Ago2, and Rbfox3 to determine how they

distribute within Fmrp. We found an ES of 0.93 falling within the top 5th percentile (**Figure 5E**); 141 out

of 175 common targets were within the top 10th percentile (bin1) of Fmrp targets, with 99 within the top

5th (**Figure 5F**). This contrasts with the negative controls, for which the best ES was 0.41 within the top

40th-60th percentile (**Figure 5G**). DAVID gene ontology analysis of those 175 common targets between

Ago2, Pum1, Pum2, Fmrp, and Rbfox3 revealed pathways enriched in neurons and axonal projections

(**Figure S11B, C**). Previous studies have shown that Pum1 and Pum2 cooperate with the miRNA

machinery to suppress certain targets (Gennarino *et al.*, 2015; Kedde *et al.*, 2010). Among Fmrp HITS-

CLIP targets, there were almost 300 microRNAs. Pum1 HITS-CLIP had 60 miRNAs, only four of which

were not shared with Fmrp; Pum2 HITS-CLIP had no miRNAs that were not shared with either Pum1 or

Fmrp (**Figure S11D** and **Table S6**). The Pum1 complexes with these fellow RBPs are therefore

physiologically relevant.

PUM1 interactors are destabilized in cell lines from infantile-onset but not adult-onset patients

Having identified Pum1 interactors and shared targets, we were finally in a position to determine

whether either the mildest (T1035S) or most severe (R1147W) missense mutations in PUM1 destabilize

these key RBP interactors. We compared patient-derived cell lines (lymphoblasts for T1035S, fibroblasts

for R1147W) with their respective age-, sex-, and cell type matched healthy controls. IP (**Figure 6A, B**)

and post-IP (**Figure S12A, B**) against PUM1 followed by western blot show that our protocol pulled

down 100% of PUM1 protein from both patient-derived cell lines. We then confirmed that R1147W,

which lies outside the RNA-binding domain, retains RNA-binding activity, while T1035S does not

(**Figure S12C**).

Co-IP confirmed that PUM1 associates with FMRP, AGO2, CNOT1, and MOV10 in patient cell lines

(**Figure 6A, B**). (RBFOX3 and PUM2 were expressed at too low levels in the patient-derived cells.) The

mild T1035S variant had no effect on any of these proteins (**Figure 6B**), but R1147W impaired PUM1

association with AGO2, MOV10, and CNOT1 (**Figure 6A**).

In order to compare the effects of the mutants in the same cell type, we turned to HEK293T cells.

GST-AGO2 associated with Myc-PUM1-R1147W 72% less than it did with Myc-PUM1-WT (**Figure**

S13A), in alignment with our observations in the PADDAS cell lines (**Figure 6A, B**). CNOT1

interactions were diminished by ~35% (**Figure S13B**). We observed no decrease in PUM1 association

with FMRP (**Figure S13C**), again in accord with our findings in patient-derived cells.

We next asked whether the R1147W mutation might be impaired in binding to WT PUM1. Myc-

PUM1-R1147W pulled down 51% of total GST-PUM1-WT, but the interaction between Myc-PUM1-

T1035S and GST-PUM1-WT remained unchanged (**Figure S13D**). The same results were observed after

RNase treatment (**Figure S13E**). Moreover, the combination of lower protein levels and marked protein

instability explains why the R1147W human phenotype is closer to that of the *Pum1* null mice than to the

heterozygous mice (Gennarino *et al.*, 2018).

To confirm that R1147W destabilizes these PUM1 interactors, we measured these RBPs in the

patient-derived cell lines. The proteins that lose their association with the R1147W variant also are

reduced in their expression (**Figure 6C, D**). Notably, MOV10's association with R1147W was greatly

reduced (**Figure 6A**) even though its protein levels were unchanged (**Figure 6C, D**). AGO2 and CNOT1

levels were unchanged in the T1035S cell line but were ~50% lower in the R1147W cell line (**Figure 6C,**

D). The mRNA levels of *PUM1*, *AGO2*, *CNOT1*, and *MOV10* did not change (**Figure S13F**), confirming

that the reductions in their respective protein levels were due to the loss of interaction with PUM1-

R1147W. Collectively, these data suggest that the R1147W variant exerts a dominant-negative effect on

PUM1-RBP interactors by destabilizing them.

Shared targets are upregulated in PADDAS but not PRCA

We next tested the effects of the T1035S and R1147W mutations on both shared targets and validated

PUM1-specific targets (Chen *et al*, 2012; Gennarino *et al.*, 2015; Zhang *et al.*, 2017) that are not in the

HITS-CLIP data for the other RBPs but are expressed in both fibroblasts and lymphoblasts. PUM1-

specific mRNA were dysregulated to very similar extents in PRCA and PADDAS patient cells, with only

a few targets being upregulated by as much as 20% in the adult-onset disease (**Figure S14A**).

Of the 175 targets shared between PUM1, PUM2, AGO2, FMRP, and RBFOX3 (**Table S5**), 54 were

expressed in both R1147W fibroblasts and T1035S lymphoblastoid cells. Fifty-one of those were

upregulated in the former but not the latter (**Figure 7A**), by an average of two-fold (ranging from a low of

121% for *IDS* to 347% for *TLK1*). There was little or no change in most of these targets in T1035S cells,

though levels of *CALM1*, *ATP2A2*, *CREB1*, and *GNAQ* fell by ~40%, and *CALM2*, *TAOK1*, and *UBE2A*

by ~20% (**Figure 7A**).

To determine whether restoring PUM1 levels would normalize expression of these shared targets, we

transfected Myc-PUM1-WT in R1147W cells, using transfection with an empty vector as a negative

control (**Figure 7B**, **Figure S14B**). WT PUM1 rescued AGO2 and CNOT1 protein levels compared to

age- and sex-matched healthy controls and this reduced the levels of the top 15 upregulated shared targets

(**Figure 7B, C**). These data confirm that the effects of the R1147W mutation, which does not impair

PUM1 binding to mRNA (Gennarino *et al.*, 2018), result from loss of interactions with RBPs that repress

the same mRNA targets. These results also support the hypothesis that the symptoms observed in adult-

onset SCA47 are attributable to the dysregulation of PUM1-specific target genes, while infantile-onset

SCA47 involves both protein partner destabilization and dysregulation of the partner proteins' targets.

Discussion

Since our initial study describing PUM1-related phenotypes (Gennarino *et al.*, 2018), we and others

have identified additional SCA47 patients (Bonnemason-Carrere *et al*, 2019; Gennarino *et al.*, 2018;

Imaizumi *et al*, 2019; Lai *et al*, 2019). In our cohort, heterozygous deletion and the R1147W mutation

account for the majority of infantile-onset patients, and T1035S for the majority of adult-onset cases. The

question that drove the present study is: why should the additional 25% drop in PUM1 levels from PRCA

to PADDAS produce such different phenotypes, especially when R1147W is not impaired in binding to

mRNA? Our data support the hypothesis that the difference is not due to a linear increase in the de-

repression of mRNA targets but is rather a function of an additional mechanism coming into play: the

destabilization of numerous interactors and the de-repression of their downstream targets. This conclusion

relies on five lines of evidence. First, loss of Pum1 in heterozygous and knockout mice changes the levels

of associated proteins, with unexpected differences emerging between brain regions and between male

and female mice. The odds of consistently identifying specific proteins in different brain regions and

sexes as false positives, across as many mice as these experiments required, are extremely low. Second,

we observed diminished function of the RBP interactors in the absence of Pum1, insofar as their targets

are dysregulated in Pum1-KO mice; moreover, the dysregulation of miRNA showed opposite patterns in

male and female cerebella that correlated with the sex-specific patterns of Ago2 expression. Third, the

levels of these proteins were reduced 40-70% in PADDAS patient cell lines, despite unaltered mRNA

levels, but not in PRCA patient cells; we also found that 55 shared targets expressed in both lymphoblasts

and fibroblasts were derepressed in infantile-onset SCA47, but not adult-onset SCA47, cells. Fourth, our

in vitro studies showed that AGO2, CNOT1, and WT PUM1 lose their interaction with PUM1-R1147W.

Fifth, re-expression of PUM1 in PADDAS cell lines restored the abundance of its interactors and

suppression of downstream targets. These results underscore the importance of examining RBP

interactions *in vivo*, in specific contexts (different sex or brain regions), with and without RNase

treatment.

There are other dosage-sensitive proteins that produce different phenotypes depending on their

expression level (Tan & Zoghbi, 2019), and our results raise the possibility that interacting complexes

may be disrupted once expression falls below a certain threshold, which for PUM1 appears to be around

50%. It is worth noting that a recent study found that, below a threshold of ~70% of normal levels of

FMRP, there were steep decreases in IQ for each further decrement in FMRP levels, even as small as 5-

10% (Kim *et al*, 2019). The amount of loss that can be sustained for a given protein would likely depend

on its usual abundance, and it is possible that for some proteins, the difference in phenotype between

greater and lesser abundance reflects a linear progression from mild to severe. For example, in

proteopathies such as Alzheimer's or Parkinson's disease, genetic duplications of *APP* or *SNCA* cause an

earlier onset of what is recognizably the same disease (Chartier-Harlin *et al*, 2004; Rovelet-Lecrux *et al*,

2006). Similarly, a mutation in *MECP2* that reduces its protein levels by only 16% is still sufficient to

cause a mild form of Rett syndrome (Takeguchi *et al*, 2020).

In contrast, there are diseases in which the phenotypes do not simply range from mild to severe

versions of the same symptoms, but seem to take on a different character. In the polyglutaminopathies,

the disease-causing protein bears an abnormally expanded CAG tract that tends to expand upon

intergenerational transmission. Although the range of normal and pathogenic repeat tract lengths differs

from one polyglutamine disease to another, larger expansions are more unstable, cause earlier onset, and

affect far more tissues than smaller expansions (Orr & Zoghbi, 2007). For example, adult-onset SCA7

presents as ataxia, but infantile SCA7 affects the entire nervous system, the heart, and the kidneys, and

leads to death by two years of age (Bah *et al*, 2020). Another example is Huntington's disease (HD),

where the juvenile form frequently lacks the classic chorea yet produces seizures, which are not a feature

of the adult-onset disease; brain morphometry is also quite different in adult- and juvenile-onset cases

(Tereshchenko *et al*, 2019). In this family of diseases, therefore, the mechanism is the same (repeat

expansion), but different tissues have different thresholds for the CAG repeats. Moreover, the brain

regions most vulnerable to HD show dramatic levels of somatic instability that correlate better with

clinical outcomes than the germline polyglutamine expansion (Genetic Modifiers of Huntington's

Disease, 2015; Kacher *et al*, 2021).

In the case of PUM1-related disease, it seems that an additional mechanism comes into play for the

more severe phenotypes, beyond upregulation of mRNA targets. Interestingly, FMRP, which harbors a

dynamic CGG repeat, is also associated with two very different phenotypes, through two different

mechanisms: very large expansions silence the gene and produce Fragile X syndrome, whereas

premutations cause the adult-onset Fragile-X-associated tremor/ataxia syndrome through RNA toxicity

(FXTAS) (Salcedo-Arellano *et al*, 2020). Interestingly, the clinical presentation of FXTAS differs by sex.

Our data suggest that the three mechanisms of repression that have been proposed for PUM1—

collaborating with the miRNA machinery (Friend *et al.*, 2012; Kedde *et al.*, 2010; Miles *et al.*, 2012),

recruiting the CCR4-NOT deadenylase complex to trigger degradation (Temme *et al.*, 2014; Van Etten *et*

al., 2012; Weidmann *et al.*, 2014), and antagonizing poly(A)-binding proteins to repress translation

(Goldstrohm *et al.*, 2018)—might be coordinated in neurons, insofar as PUM1, PUM2, FMRP, AGO2,

MOV10, CNOT1 and RBFOX3 (and related proteins in specific brain regions) either interact or are so

close to each other within the ribonucleosome that the loss of Pum1 or RNA can change the composition

of the complexes that are identified by co-IP, in ways that are specific to brain region and sex. In this

context it is worth noting that a very recent study found alternative splicing is altered in hippocampal

slices from Fmrp-deficient mice; this observation was attributed to changes in H3K36me3 levels (Shah *et*

al, 2020), but our data suggest that FMRP has a closer relationship with the RBFOX protein family and

alternative splicing machinery than previously imagined. Indeed, recent work has provided tantalizing

glimpses of close interactions among various kinds of RNA metabolism. For example, members of the

RBFOX family of proteins may, depending on their interactors (and perhaps cell type, sex, age, and

species), be involved in microRNA processing in the nucleus and translation in the cytoplasm (Conboy,

2017). The FMRP/MOV10 complex appears to be involved in regulating translation through miRNA,

with evidence that this role may change according to cell type (Kenny *et al.*, 2020). Another study used

quantitative mass spectroscopy to examine how Fmrp expression levels change with age in the wild-type

rat dentate gyrus and found differences in the levels of myriad proteins; among the 153 proteins with the

most significant changes in levels were Pum1, Pum2 and Papbc1 (Smidak *et al*, 2017). The region-, sex-,

and age-specificity of certain interactions indicates that unraveling RBP interactomes *in vivo* will require

considerable finesse. But creating such interactomes should lead to a more complex yet realistic picture of

RBP roles in neuronal function and in neurological disease.

Acknowledgments

We thank the patients and their families for participating in our study. We thank Mingyu Cao for help

with statistical analyses, U.N. Wasko for starting the cloning procedure with Myc and GST tag sequences

in vitro, Wayne Miles for sharing the RIP protocol and send us the HCT116-*PUM1*[-/-] cell lines, and V. L.

Brandt for editing the manuscript and suggesting we examine the targets of PUM1 interactors. We also

thank G. Struhl, V. Ambros, T. Duchaine, G. Karsenty, H. Zoghbi, M. Jovanovic, Y. Giardina, and

members of the Gennarino laboratory for helpful discussions and support. This work was supported by

the National Institute of Neurological Disorders and Stroke (NINDS; R01NS109858 to V.A.G.); the

National Ataxia Foundation/Young Investigator Research Grant (V.A.G.); the National Institute of

406   Author contributions

416   Declaration of interests

417   The authors declare not competing interests

419   Figure legends

Figure 1. A brain-specific Pum1 interactome. Network of putative Pum1 interactors in 10-week-old

mouse brain (circles connected to Pum1 by gray lines). Interactions between interactors (purple lines)

were inferred by g:GOSt from Corum and the Human Protein Atlas (see Materials and Methods). The

proteins in each of the 10 clusters are listed to the right. We combined and homogenized whole brains

from two 10-week-old wild-type mice per sample (1 female and 1 male), aliquoting half of each sample

for IP against either Pum1 or IgG, then performed six biological replicates (six samples, 12 mice total) for

each LC-MS/MS experiment against IP-Pum1 and IP-IgG. All putative Pum1 interactors are listed in

Table S1.

Figure 2. A brain-region specific Pum1 interactome. (**A**) Pum1 interactome from 10-week-old mouse

cerebellum (n=8 mice, 4 male and 4 female), hippocampus (n=10, 5/5), cortex (n=8, 4/4) and the rest of

the brain (i.e., excluding those three regions) for a total of 1,500 proteins (**Table S1**). Node colors

represent different brain regions or the overlap between two or more brain regions as noted. All

experiments were performed at least in triplicate. IP against IgG was used as a negative control. (**B**)

Bubble plots show the top categories from gene ontology analyses of Pum1 interactors from whole brain,

hippocampus, cortex, and cerebellum. Only the categories with fold enrichment >1.5 and FDR<0.05 are

shown; not all are labeled because of space limitations. The full list of gene ontology categories is

available in **Table S2**.

Figure 3. Validation of Pum1 associations with post-transcriptional RNA-binding proteins in mouse

brain and HEK293T cells. (**A**) Representative western blot of proteins pulled down by IP against Pum1

compared to IgG from wild-type mice brain without (*left*) and with (*right*) RNase treatment. In this panel,

after IP-Pum1, we immunoblotted for Pum1 (positive control), Pum2, Fmrp, Ago2, Rbfox3, Cnot1, and

Mov10. (see Materials and Methods.). (**B**) Representative western blots of the same proteins validated in

panels **A** after IP against PUM1 with or without RNase treatment from HEK293T cell lines. In panels **A**

and **B**, IP against IgG was used as a negative control and Input (1% from the initial protein lysate) as a

loading control. The numbers on the right are the respective molecular weights expressed in kilodaltons

(kDa). All the experiments were repeated at least four times. All mice were sacrificed at 10 weeks of age.

Figure 4. Pum1 interactors and the microRNA machinery show brain region- and sex-specific

responses to Pum1 loss. Representative western blots of Pum1, Pum2, Fmrp, Ago2, Rbfox3, Cnot1, and

Mov10 in (**A**) cerebellum, (**B**) hippocampus, and (**C**) cortex in both male (left panel) and female (right

panel) WT, *Pum1$^{+/-}$*, and *Pum1$^{-/-}$* mice. All the experiments were conducted with equal numbers of 10-

week-old male and female mice per genotype, for a total of at least 12 mice per genotype (data represent

mean ± SEM). Graphs below show quantification for each protein by brain region, sex, and genotype. All

data were normalized to Gapdh protein levels. Data represent mean ± SEM. P values were calculated by

two-tailed Student's t test. *p < 0.05, **p < 0.01, ***p < 0.001. P1 indicates Pum1. See Figure S8 for

mRNA quantification for each interactor, brain region, and sex. (**D**) Heatmap showing 166 microRNAs

from cerebella of *Pum1$^{-/-}$* male and female mice that were dysregulated (fold change -3 to +3) relative to

wild-type cerebellum. The full list of miRNA names and fold changes are available in Table S3. See

Figure S9 for male and female miRNA scatter plots. (**E**) Heatmap showing mRNA quantification by

qPCR for 49 targets co-bound by a minimum of eight dysregulated miRNAs (>25% change) from panel

D. Statistical significance and magnitude of dysregulation are illustrated for both male and female in

Figure S10. The entire list of targets predicted to be co-bound by at least two miRNAs is presented in

Table S4.

Figure 5. Pum1 and its RNA-binding protein interactors share many neuronal mRNA targets. (A)

Enrichment plots generated by Gene Set Enrichment Analysis (GSEA) of Pum1, Pum2, Rbfox3, and

Ago2 HITS-CLIP targets plotted again Fmrp ranked HITS-CLIP data. Pum1, Pum2, Rbfox3, and Ago2

targets are enriched at the top 10^{th} percentile of the Fmrp targets with FDR=0. (**B**) GSEA analysis scores

of HITS-CLIP data from each negative control (Apc, Nova, Ptpb2, and Mbnl2) plotted against Fmrp

ranked HITS-CLIP data. The negative controls have a maximum enrichment score of 0.36 for Apc

ranking at the top 44.7% with FDR=1. (**C**) GSEA analysis scores of Pum1, Pum2, Rbfox3, and Ago2

HITS-CLIP data plotted against Fmrp HITS-CLIP data divided into 10-percentile ranked bins shows the

shared targets are among the top percentiles of targets for each protein. (**D**) GSEA analysis scores of the

HITS-CLIP data for Fmrp, Pum2, Ago2, Rbfox3 and four negative controls (Apc, Nova, Ptpb2, and

Mbnl2) against Pum1 ranked HITS-CLIP data. The targets of Fmrp, Ago2, Pum2, and Rbfox3 are

enriched at the top 5^{th} to 18^{th} percentile of Pum1 targets. (**E**) GSEA analysis of the shared targets between

Pum1, Pum2, Ago2, and Rbfox3 against Fmrp showing that they are enriched in the top 5^{th} percentile of

Fmrp ranked targets. (**F**) Pum1, Pum2, Ago2, and Rbfox3 shared targets plotted against Fmrp ranked

HITS-CLIP targets and divided into 10-percentile bins shows that all of their respective targets are

enriched at the top 10^{th} percentile of the Fmrp ranked targets. (**G**) GSEA analysis scores of the targets

shared by Pum1, Pum2, Ago2, and Rbfox3 and the four negative controls (Apc, Nova, Ptpb2, and Mbnl2)

plotted against Fmrp. At best the negative controls are enriched at the top 40% with a maximum ES of

0.41. For all the GSEA analyses, the False Discovery Rate (FDR) was provided by GSEA: **FDR<0.05

and ***FDR<0.01. ES=Enrichment score (blue line). Note that lowest rank at max percentage indicates

stronger targets in the rank (see Materials and Methods). HITS-CLIP data, and the respective rank, were

obtained from the literature and were initially acquired as follows: Pum1 and Pum2 (Zhang *et al.*, 2017),

Fmrp (Maurin *et al.*, 2018), Ago2 (Chi *et al.*, 2009), Rbfox3 (Weyn-Vanhentenryck *et al.*, 2014), Nova

(Zhang *et al.*, 2010), Ptpb2 (Licatalosi *et al.*, 2012), Mbnl2 (Charizanis *et al.*, 2012), and Apc (Preitner *et*

al., 2014) (see Materials and Methods for more details). The full list of shared targets is reported in Table

S5.

Figure 6. The R1147W mutation, but not T1035S, destabilizes PUM1 interactors. (A-B) IP against

PUM1 from Subject 1 (S1, R1147W) (**A**) and Subject 2 (S2, T1035S) (**B**) compared to their respective

age-, sex-, and cell type- matched healthy controls confirm the interactions between PUM1 (used here as

a positive control), and PUM2, FMRP, AGO2, CNOT1, and MOV10. Input (1%) was used as a loading

control and IP against IgG was used as a negative control. (**C**) Representative western blots (*left panels*)

and relative quantification (*bar graphs to the right*) of protein levels for PUM1, PUM2, FMRP, AGO2,

CNOT1, and MOV10 in PADDAS patient-derived and PRCA patient-derived cells compared to their

respective controls. Data were normalized to Gapdh protein levels. From **A** to **C,** all the experiments were

performed at least three times. Data represent mean ± SEM. P values were calculated by two-tailed

Student's t test. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$.

Figure 7. Shared targets are upregulated only in PADDAS, not in PRCA. (**A**) mRNA level

quantification of PUM1 neuronal targets in common with FMRP, PUM2, AGO2, and RBFOX3 (Figure

5E and Table S5) in S1 (R1147W, *blue bars*) and S2 and 3 (T1035S, *orange bars*) cell lines compared to

their respective healthy controls. Only genes expressed in both R1147W and T1035S cell lines are

represented here, for a total of 54 genes. (**B**) Representative western blots (*right panel*) and relative

quantifications (*left panel*) of PUM1 and its interactors (AGO2, CNOT1, FMRP, and MOV10) in

R1147W patient-derived cell lines after Myc-PUM1-WT expression. (**C**) mRNA quantification of the top

15 shared target genes from panel **A** in R1147W patient-derived cell lines after Myc-PUM1-WT

expression. All data were normalized to GAPDH mRNA or protein levels and experiments were

performed at least three times. Data represent mean ± SEM. P values were calculated by two-tailed

Student's t test. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$. The full list of shared targets expressed in fibroblast

and lymphoblast cell lines is reported in Table S5.

Supplemental figures and legends

Figure S1. Pumilio1 antibody efficiency for IP LC-MS/MS.

(**A**) Volcano plot analysis showing all the proteins pulled down by IP against IgG and Pum1 from mouse

brain. (**B**) Principal component analysis (PCA) of IP-Pum1 followed by LC-MS/MS in WT adult mouse

brains compared with IP against IgG; each dot represents one sample for a total of 12 samples processed

by LC-MS/MS. (**C**) Pre-IP, IP, and post-IP against Pum1 and IgG from wild-type (WT) mouse brain.

Even at very long exposure, the post-IP (10%) Pum1 lane has no residual band at 120 kDa even though

10 times more protein is loaded than Input (1%). This demonstrates the specificify of the Pum1 antibody,

which makes it suitable for IP LC-MS/MS. The numbers on the right show molecular weight in

kilodaltons (kDa).

Figure S2. Volcano plot and PCA analyses of IP:Pum1 followed by LC-MS/MS in cerebellum,

hippocampus, and cortex. (A) Principal component analysis (PCA) of IP-Pum1 followed by LC-MS/MS in cortex, hippocampus, and cerebellum from WT mice. **(B-D)** Volcano plots show all the proteins pulled down by IP against IgG and Pum1 from **(B)** cerebellum, **(C)** cortex, and **(D)** hippocampus at 10 weeks of age. IP against IgG was used as a negative control. Each dot represents a total of 3 samples processed by MS for each brain region. All putative Pum1 interactors are listed in **Table S1**.

Figure S3. Many Pum1 interactors are specific to certain brain regions and not determined by expression level. (A) Proteomic analysis in whole brain, cerebellum, hippocampus, and cortex at 10 weeks of age shows that Pum1 interactors are not the most highly expressed proteins. Dotted line represents the mean expression of Pum1 interactors from each brain region. Proteomics was performed in duplicate (one male and one female) for each brain region. **(B)** Absolute expression value of the validated Pum1 interactors. Pum2 is expressed a low levels in all three brain regions but was still the strongest Pum1 interactor in the cortex, suggesting a specific interaction. **(C)** Immunoblot for Pum1 (positive control), Anapc1, Cnot1 and Pum2. **(D)** Western blot analysis at 10 weeks of age to evaluate Pum2 expression levels in eight different brain regions as well as whole brain. Pum2 is highly expressed in olfactory bulbs and amygdala, and expressed at similar levels in hippocampus, cerebellum, and cortex. All experiments performed in triplicate. Cerebellar and cortical tissues: n=8 wild-type mice (4 male and 4 female), for a total of 24 mice. Hippocampus: n=10 wild-type mice (5 female and 5 male), for a total of 30 mice. All mice were 10 weeks of age. IP against IgG was used as a negative control. Molecular protein weights are expressed in kilodaltons (kDa). HP: hippocampus; CB: cerebellum; CX: cortex; MB: midbrain; OB: olfactory bulbs; AM: amygdala; SN: substantia nigra pars compacta; BS: brain stem; WB: whole brain. All the experiments were repeated at least three times.

Figure S4. Fmrp and Pum2 IP recombinant proteins and validation of APC/C and mTOR proteins. (A-B) Representative western blot of PUM1 pulled down by recombinant **(A)** GST-FMRP or **(B)** GST-PUM2. **(C)** Representative western blot of Rptor, Cad, Anapc1 and Fzr1 proteins pulled down by IP against Pum1 (used here as positive control) from WT mice. **(D)** Representative western blot of the reciprocal co-IPs against Anapc1, Fzr1, Rptor, and Cad. Each co-IP was immunoblotted against Pum1 and the pulled down protein used here as a reference protein to confirm the respective protein-protein interaction. For **C** and **D**, IP against IgG was used as a negative control, and Input (1% from the initial protein lysate) as a loading control. Molecular weights are expressed in kilodaltons (kDa) to the right. All the experiments were repeated at least four times. All mice were sacrificed at 10 weeks of age with an equal number or male and female.

Figure S5. IP and co-IP against Pum1 and the interactors from whole brain in *Pum1*^{-/-} and WT

mice, with and without RNase. (**A**) IP against Pum1 in *Pum1*^{-/-} mouse demonstrates the complete

absence of Pum1 and thus the specificity of the anti-Pum1 antibody. IP against IgG was used as a

negative control, and Input (1% from the initial protein lysate) as a loading control. (**B**) IP against Pum1

(with or without RNase treatment) shows no interaction with Ago1 or Ago3 in the mouse brain. (**C**)

Representative western blots of Pum1 pulled down by Pum2, Fmrp, Ago2, Cnot1 from WT and *Pum1*^{-/-}

mouse brains, with and without RNase. Molecular weights at the right are in kilodaltons (kDa).

All the experiments were conducted with one female and one male mouse brain at 10 weeks of age, in

triplicate, for a total of 6 mice. IP against IgG was used as a negative control, and Input (1% from the

initial protein lysate) as a loading control.

Figure S6. Protein and mRNA quantification from WT, *Pum1*^{+/-} and *Pum1*^{-/-} mouse brains. (**A**)

Representative western blot with relative quantifications of Pum1, Pum2, Fmrp, Ago2, Rbfox3, Cnot1,

and Mov10 from whole brains of WT, *Pum1*^{+/-} and *Pum1*^{-/-} mice. All data were normalized to Gapdh

protein levels. The numbers on the right are the respective molecular weights expressed in kilodaltons

(kDa). (**B**) mRNA level quantification by qPCR of *Pum1*, *Pum2*, *Fmrp*, *Ago2*, *Rbfox3*, *Cnot1*, and *Mov10*

from whole brains of WT, *Pum1*^{+/-} and *Pum1*^{-/-} mice. Again, all data were normalized to *Gapdh* mRNA

levels. All the experiments were conducted with equal number of male (at least 6 per genotype) and

female (at least 6 per genotype) mice at 10 weeks of age, for a total of at least 12 mice per genotype (data

represent mean ± SEM). The *p* values were calculated by the Student's t test. *p < 0.05, **p < 0.01, ***p

< 0.001.

Figure S7. mRNA quantification of Pum1 interactors by brain region and sex in WT, *Pum1*^{+/-} and

***Pum1*^{-/-} mice.** mRNA levels in cerebellum, hippocampus, and cortex in male and female for all the

validated Pum1 interactors. The same number of mice were used here as in **Figure 4A-C** for a total of at

least 12 mice per genotype and sex at 10 weeks of age. All data were normalized to *Gapdh* mRNA levels.

All the experiments were performed at least six times (data represent mean ± SEM). The *p* values were

calculated by the Student's t test. *p < 0.05, **p < 0.01, ***p < 0.001.

Figure S8. Volcano plots representing all the miRNAs sequenced by miRNAseq in male and female.

(**A**) Principal component analysis (PCA) of miRNAseq in wild-type and *Pum1*^{-/-} male and female mice

cerebella at 10 weeks of age. (**B**) Volcano plots show the expression profile for all the miRNAs in male

and female *Pum1$^{-/-}$* mice compared to WT at 10 weeks of age. The orange dots represent the miRNAs

upregulated in female and downregulated in males; the blue dots represent the miRNAs downregulated in

female and upregulated in males (see Material and Methods).

Figure S9. mRNA quantification of the 49 targets co-bound by at least eight dysregulated miRNAs

in mouse cerebellum. qPCR in cerebellum of male (*left*, red) and female (*right*, blue) mice at 10 weeks

of age for the 49 targets co-bound by at least eight dysregulated miRNAs (with minimum 25% change in

expression) from **Figure 4E** and **Table S4**. All the experiments were performed in triplicate for both male

and female (data represent mean ± SEM). The *p* values were calculated by two-tailed Student's t test. *p

< 0.05, **p < 0.01, ***p < 0.001.

Figure S10. Gene ontology analysis for all targets predicted by CoMeTa and TargetScan that are

co-bound by at least four miRNAs. (**A**-**C**) David Gene Ontology representing the enriched (**A**)

biological processes, (**B**) cellular components, and (**C**) KEGG pathways for all the targets co-bound by at

least four miRNAs. For this analysis we set FDR<0.01 and a fold-enrichment >2. (**D**) Synaptic Gene

Ontology (SynGO) predicts that 117 targets are presynaptic and 124 are postsynaptic with a log10Q

value≥5.

Figure S11. GSEA and Gene Ontology data pertaining to Figure 5. (**A**) Gene Set Enrichment

Analysis (GSEA) of Pum1 HITS-CLIP data plotted against HITS-CLIP data from the negative controls

(RBPs that did not show up in the Pum1 interactome: Ptpb2, Nova, Apc, and Mbnl2) reveals no

significant enrichment. (**B**-**C**) Gene ontology analysis of the HITS-CLIP targets shared between Pum1,

Pum2, Fmrp, Ago2, and Rbfox3 reveals enrichment for certain (**B**) biological functions and (**C**) cellular

localization. Only categories with FDR<0.05 and fold enrichment > 5 were plotted in **B** and **C**. (**D**) Venn

diagram of miRNAs identified by Pum1 and Pum2 shows almost 100% overlap with the miRNAs pulled

down by Fmrp HITS-CLIP. For full list of shared miRNAs see **Table S6**. For all GSEA analyses the

False Discovery Rate (FDR) was provided by GSEA, ***FDR < 0.01. ES=Enrichment score (blue line).

Figure S12. PUM1 R1147W retains RNA-binding activity. (**A**-**B**) Pre-IP, IP, and post-IP against

PUM1 and IgG from (**A**) R1147W (infantile-onset SCA7, or PADDAS) fibroblasts and

(**B**) T1035S (adult-onset SCA47 or PRCA) lymphoblastoid cells. In both cell lines we were able to pull

down 100% of PUM1. Pre-IP represents 1% from the initial protein lysate as a loading control, while

10% of the protein lysate was loaded as post-IP. Molecular weights provided at right in kilodaltons (kDa).

(**C**) RNA-immunoprecipitation assay followed by qRT-PCR to show that T1035S is impaired in RNA-

binding while R1147W is not. We used *E2F3* and *CDK1* as validated PUM1 targets and E2F4 as a

negative control. We transfected WT and PUM1-KO HCT116 cells with 250ng of either Myc-PUM1-

WT, Myc-PUM1-R1147W, or Myc-PUM1-T105S. RIP was performed in triplicate.

Figure S13. PUM1 validation experiments in patient-derived cells. (**A-D**) Representative western

blots and relative IP quantification (*bar graphs*) of IP against Myc-PUM1-WT, Myc-PUM1-T1035S

(PRCA), or Myc-PUM1-R1147W (PADDAS) followed by immunoblotting for: (**A**) GST-AGO2, (**B**)

GST-CNOT1, (**C**) GST-FMRP, and (**D**) GST-PUM1-WT. Myc- and GST-tagged proteins were co-

transfected in HEK293T cells in equal quantities (250ng each). The molecular weights were expressed in

kilodaltons (kDa). The amount of protein pulled down compared to IP-PUM1 was quantified as

$[IP^X/(Input^X/GAPDH^X)]/ [IP^{PUM1}/(Input^{PUM1}/ GAPDH^{PUM1})]$, where X is the protein of interest. (**E**)

Representative western blots of IP with RNase treatment against Myc-PUM1-WT, Myc-PUM1-T1035S

(PRCA), and Myc-PUM1-R1147W (PADDAS) followed by immunoblotting to test binding between

PUM1 proteins without the RNA. The numbers on the right are the respective molecular weights

expressed in kilodaltons (kDa). All the IPs were repeated at least three times. (**F**) mRNA quantification

for all of the immunoblotted proteins in **Figure 6C** in PADDAS and PRCA patient-derived cell lines

compared to their respective age-, sex-, and cell-type-matched controls. In all the experiments data

represent mean ± SEM. P values were calculated by two-tailed Student's t test. *$p < 0.05$, **$p < 0.01$,

***$p < 0.001$.

Figure S14. mRNA quantification of PUM1-specific targets. (**A**) qPCR analysis of validated PUM1-

specific targets from PADDAS (R1147W) patient-derived fibroblasts (blue bars) compared to three age-

and sex-matched fibroblast control cell lines, and PRCA (T1035S) patient-derived lymphoblastoid cell

lines (orange bars) compared to three age- and sex-matched lymphoblastoid control cell lines. Only genes

expressed in both fibroblasts and lymphoblasts are represented here, for a total of 10 genes. (**B**) mRNA

quantification of PUM1 from PADDAS patient-derived fibroblasts transfected with empty and Myc-

PUM1-WT vectors, compared to three age- and sex-matched fibroblast control cell lines. All data were

normalized to *GAPDH* mRNA levels and experiments performed at least three times. Data represent

mean ± SEM. P values were calculated by two-tailed Student's t test. *$p < 0.05$, **$p < 0.01$, ***$p <$

0.001.

Materials and Methods

Immunoprecipitation (IP) experiments using mouse brain tissue. Mouse brain tissues were gathered from an equal number of 10-week-old male and female mice. For whole brain experiments, we combined and homogenized two 10-week-old wild-type mouse brains per sample (1 female and 1 male), aliquoting half of each sample for IP against either Pum1 or IgG, then performed six biological replicates (12 mice total) for each LC-MS/MS experiment against IP-Pum1 and IP-IgG. For experiments on the hippocampus, cerebellum, and cortex, we needed much larger numbers of mice: we combined cerebellar and cortical tissues from eight wild-type mice (4 male and 4 female) and performed the experiment in triplicate (total of 24 mice), while for hippocampus we combined tissues from ten wild-type mice (5 female and 5 male) for three experiments (a total of 30 mice).

Samples were processed with a dounce homogenizer using a lysis buffer consisting of 200mM $NaCl_2$, 100mM $NaPO_4$, 20mM Hepes pH 7.4, 1% Triton X (which should disrupt all but the strongest protein-protein interactions) and complemented by 1X of Xpert Protease and 1X of Phosphatase Inhibitor Cocktail Solutions (GenDepot, #P3100-100, #P3200-020). Following homogenization, the samples were placed on ice for 15 minutes then centrifuged at 14,800 rpm at 4°C for 25 minutes to remove the debris from the supernatant. The supernatant was then moved to 1.5 ml tubes (Beckman microfuge tube #357448) and spun down in a Beckman ultra-centrifuge (Optima Max XP) at 4°C for 25 minutes at 44,000 rpm. 10% of the protein lysate was stored as input and only 1% was loaded for western blot. The protein extract was later divided into two aliquots, one for IP against the protein of interest (antibodies listed below) and the other for IP against IgG, and was then incubated with 30 μL of Dynabeads[TM] Protein G (Invitrogen, #10004D) and 5 μg of antibody overnight at 4°C on a rotisserie tube rotator. The next day, the beads were washed four times with the same lysis buffer used for IP and resuspended in 40μL of elution buffer (consisting of lysis buffer, NuPAGE 10X Reducing Agent [Invitrogen, #NP0009], NuPAGE LDS sample buffer at 1X final concentration [Invitrogen, #NP0007]) and boiled at 95°C for 10 minutes before the samples were loaded in the NuPAGE 4%–12% Bis-Tris Gels (Invitrogen, #NP0335BOX & #NP0336BOX) for further resolution and western blot analysis.

For the IP with RNase treatment, the beads were resuspended in 400 μL of lysis buffer after the three final washes and divided into two separate 1.5 ml tubes of 200 μL each. To establish the dose required to remove all RNA, we tested different amounts of RNase I (Invitrogen, #EN0602) and found that 4 μL was enough to render RNA undetectable both by denaturing gel and cDNA amplification. This sample and the negative control (i.e., one without RNase treatment) were incubated at 37°C for 15 min on a rotisserie tube rotator. After incubation, all the samples were washed one last time with 500 μL of lysis buffer and then eluted in 20 μL of elution buffer. We used the same protocol for all the IP processed by LC-MS/MS.

The antibodies used for IP were: goat α-PUM1 (Bethyl Laboratories, #A300-201A), rabbit α-PUM2

(Bethyl Laboratories, #A300-202A), rabbit α-FMRP (Abcam Cambridge, #ab17722), rabbit α-AGO2

(Abcam Cambridge, #ab32381), rabbit α-NeuN (Thermo Fisher Scientific, #PA5-37407), rabbit α-

CNOT1 (Cell Signaling Technology, #44613), rabbit α-MOV10 (Bethyl Laboratories, #A301-571A), and

rabbit α-ANAPC1 (Bethyl Laboratories, #A301-653A).

Please note that *in vivo* IPs from brain lysates present certain challenges that are not encountered *in*

vitro. Whereas the total lysate from cells is usually 200μl-300μl, the brain lysate is made in a large

volume, usually 1.5 to 2.4 ml, depending on the size of the brain or brain region. This means that in a

normal western blot that accommodates 30-40μl total volume, including reducing buffer and loading blue,

we cannot load more than 1%-3% from the total brain lysate as input. Therefore, when we pull down a

protein of interest (Pum1) and immunoblot for the same protein compared to a standard input (loading the

entire IP in one gel), the resulting IP band will be much darker than the input. We then need to expose the

Input from the same membrane much longer to visualize it—this is common practice when working with

in vivo tissues (De Maio *et al*, 2018; Di Grazia *et al*, 2021; Jin *et al*, 2020; Lee *et al*, 2020; Lee *et al*,

2008; Rousseaux *et al*, 2018).

Immunoprecipitation experiments from HEK293T and patient-derived cell lines. HEK293T cells

and patient-derived fibroblasts or lymphoblastoid cells were lysed by pipetting up and down with a

1000μl tip in a lysis buffer consisting of 200mM NaCl$_2$, 100mM NaPO$_4$, 20mM Hepes pH 7.4, 1% Triton

X and complemented by 1X of Xpert Protease and 1X of Phosphatase Inhibitor Cocktail (GenDepot,

#P3100-100, #P3200-020). The rest of the protocol is the same as described above for mouse brain tissue,

except that we used 2.5 μg of primary antibody for IP.

Co-Immunoprecipitation in-gel digestion for LC-MS/MS. Immunoprecipitated samples were separated

on NuPAGE 4-12% gradient SDS-PAGE (Invitrogen, #NP0335BOX & #NP0336BOX) and stained with

SimplyBlue (Invitrogen, #LC6060). Protein gel slices were excised and *in-gel* digestion performed as

previously described (Shevchenko *et al*, 2006), with minor modifications. Gel slices were washed with

1:1 Acetonitrile and 100mM ammonium bicarbonate for 30 min then dehydrated with 100% acetonitrile

for 10 min until shrunk. The excess acetonitrile was then removed and the slices dried in a speed-vacuum

at room temperature for 10 minutes. Gel slices were reduced with 5 mM DTT for 30 min at 56°C in an air

thermostat, cooled down to room temperature, and alkylated with 11 mM IAA for 30 min with no light.

Gel slices were then washed with 100 mM of ammonium bicarbonate and 100% acetonitrile for 10 min

each. Excess acetonitrile was removed and dried in a speed-vacuum for 10 min at room temperature and

the gel slices were re-hydrated in a solution of 25 ng/µl trypsin in 50 mM ammonium bicarbonate for 30

min on ice and digested overnight at 37 °C in an air thermostat. Digested peptides were collected and

further extracted from gel slices in extraction buffer (1:2 ratio by volume of 5% formic acid: acetonitrile)

at high speed, shaking in an air thermostat. The supernatants from both extractions were combined and

dried in a speed-vacuum. Peptides were dissolved in 3% acetonitrile/0.1% formic acid.

Liquid chromatography with tandem mass spectrometry (LC-MS/MS). The Thermo Scientific

Orbitrap Fusion Tribrid mass spectrometer was used for peptide tandem mass spectroscopy (MS/MS).

Desalted peptides were injected in an EASY-Spray™ PepMap™ RSLC C18 50cm X 75cm ID column

(Thermo Scientific) connected to the Orbitrap Fusion™ Tribrid™. Peptide elution and separation were

achieved at a non-linear flow rate of 250 nl/min using a gradient of 5%-30% of buffer B (0.1% (v/v)

formic acid, 100% acetonitrile) for 110 minutes, maintaining the temperature of the column at 50 °C

during the entire experiment. Survey scans of peptide precursors are performed from 400 to 1500 *m/z* at

120K full width at half maximum (FWHM) resolution (at 200 *m/z*) with a 2×10^5 ion count target and a

maximum injection time of 50 ms. The instrument was set to run in top speed mode with 3-second cycles

for the survey and the MS/MS scans. After a survey scan, MS/MS was performed on the most abundant

precursors, i.e., those ions that had a charge state between 2 and 6, and an intensity of at least 5000, by

isolating them in the quadrupole at 1.6 Th. We used collision-induced dissociation (CID) with 35%

collision energy and detected the resulting fragments with the rapid scan rate in the ion trap. The

automatic gain control (AGC) target for MS/MS was set to 1×10^4 and the maximum injection time was

limited to 35ms. The dynamic exclusion was set to 45s with a 10ppm mass tolerance around the precursor

and its isotopes. Monoisotopic precursor selection was enabled.

LC-MS/MS data analysis. Raw mass spectrometric data were analyzed using the MaxQuant

environment v.1.6.1.0 (Cox & Mann, 2008) and Andromeda for database searches (Cox *et al*, 2011) at

default settings with a few modifications. The default was used for first search tolerance and main search

tolerance (20 ppm and 6 ppm, respectively). MaxQuant was set up to search with the reference mouse

proteome database downloaded from Uniprot (https://www.uniprot.org/proteomes/UP000000589).

MaxQuant searched for trypsin digestion with up to 2 missed cleavages. Peptide, site and protein false

discovery rates (FDR) were all set to 1% with a minimum of 1 peptide needed for identification; label-

free quantitation (LFQ) was performed with a minimum ratio count of 1. The following modifications

were used for protein quantification: oxidation of methionine (M), acetylation of the protein N-terminus,

and deamination for asparagine or glutamine (NQ). Results obtained from MaxQuant were further

analyzed using the Perseus statistical package (Tyanova *et al*, 2016) that is part of the MaxQuant

distribution. Protein identifications were filtered for common contaminants. Proteins were considered for

quantification only if they were found in at least two replicate groups. Significant alterations in protein

abundance were determined by ANOVA with a threshold for significance of $P < 0.05$ (permutation-based

FDR correction). Pum1 protein interactors were later considered if they were found in at least five out of

six LC-MS/MS experiments for whole brain and in at least two out of three experiments for each

respective brain region with a fold-change of >1.5 between LFQ-PUM1-WT and LFQ-IgG-WT (see

Table S1).

Protein-protein interaction map. The protein-protein interaction map for the whole brain (Figure 1) was

generated by Cytoscape (https://cytoscape.org/) (Otasek *et al*, 2019) and interactions or functional

relationships (clusters) were inferred from Corum (Giurgiu *et al*, 2019) and the Human Protein Atlas

(Thul *et al*, 2017) by g:GOSt, which is a package of g:Profiler (https://biit.cs.ut.ee/gprofiler/gost)

(Raudvere *et al.*, 2019). The brain region-specific map (**Figure 2A**) was generated by Cytoscape

(Shannon *et al*, 2003).

Quantitative proteomics from mouse brain regions.

Tissue processing for LC-MS/MS measurements

Frozen brain tissues were weighted and 25-50mg (dry weight) per sample were cryopulverized and lysed

in Urea Buffer (8M Urea, 75mM NaCl, 50mM Tris/HCl pH 8.0, 1mM EDTA) in a final 1:10 (mg/μL)

tissue:buffer ratio. Protein concentration was determined by BCA assay (Pierce). 40μg of total protein per

sample were processed further. Disulfide bonds were reduced with 5mM dithiothreitol and cysteines were

subsequently alkylated with 10mM iodoacetamide. Samples were diluted 1:4 with 50mM Tris/HCl (pH

8.0) and sequencing grade modified trypsin (Promega) was added in an enzyme-to-substrate ratio of 1:50.

After 16h of digestion, samples were acidified with 1% formic acid (final concentration). Tryptic peptides

were desalted on C18 StageTips according to (Rappsilber *et al*, 2007) and evaporated to dryness in a

vacuum concentrator. Dried peptides were then reconstituted in 3% ACN / 0.2% Formic acid to a final

concentration of 0.5 μg/μL.

LC-MS/MS analysis on a Q-Exactive HF

About 1 μg of total peptides were analyzed on a Waters M-Class UPLC using a 25cm Thermo Scientific

PepMap RSLC C18 2 μm 25cm column coupled to a benchtop Thermo Fisher Scientific Orbitrap Q

Exactive HF mass spectrometer. Peptides were separated at a flow rate of 400 nL/min with a 160 min

gradient, including sample loading and column equilibration times. Data was acquired in data

independent mode using Xcalibur software. MS1 Spectra were measured with a resolution of 120,000, an

AGC target of 5e6 and a mass range from 300 to 1800m/z. MS2 spectra were measured in 47 segment

windows per MS1, each with an isolation window width of 32 m/z (0.5 m/z overlap with the neighboring

window), a resolution of 30,000, an AGC target of 3e6, and a stepped collision energy of 22.5, 25, 27.5.

All raw data were analyzed with SpectroNaut software version 15.6.211220 (Biognosys) using a

directDIA method based on a UniProt mouse database (release 2014_07, Mus musculus), performed with

the "BGS factory settings" including the following parameters: Oxidation of methionine and protein N-

terminal acetylation as variable modifications; carbamidomethylation as fixed modification; Trypsin/P as

the digestion enzyme; For identification, we applied a maximum FDR of 1% separately on protein and

peptide level. "Cross run normalization" was activated. This gave intensity values for a total of 5699

protein groups across all samples and replicates. "PG.Quantity" (normalized across samples) values were

used for all subsequent analyses.

Protein purification of recombinant proteins. Recombinant FMRP, PUM2, and PUM1 were purified

from *Escherichia coli* BL21-Codon Plus (DE3)-RIL cells. Both FMRP and PUM2 were expressed with

an N-terminal GST tag and isolated from the cell lysate using Glutathione Sepharose 4B (GS4B) resin.

The proteins were further subjected to size exclusion chromatography using 20mM HEPES pH 8.0,

200mM NaCl, and 1mM DTT as the buffer. MBP-SNAP-PUM1-His$_6$ was expressed and purified similar

to the previously described protocol (Elguindy & Mendell, 2021). After nickel affinity chromatography,

MBP-SNAP-PUM1-His$_6$ was buffer exchanged into 20mM HEPES pH 8.0, 200mM NaCl, and 1mM

DTT using PD-10 desalting columns.

GST pull-down assay. 500mM of purified GST-PUM2 or GST-FMRP were incubated with 1μM of HIS-

PUM1 with 30μl of Glutathione Sepharose 4B GST-tagged protein purification resin (Cytiva, #17075601)

in a final volume of binding buffer (20mM HEPES pH 8.0, 200mM NaCl, 0.5% Tween-20) for 1 hour at

4°C on the rocker. Resins were washed 3 times in binding buffer and centrifuged at 4000g at 4°C; resins

were resuspended in 30μl of elution buffer—which consisted of binding buffer, NuPAGE 10X Reducing

Agent (Invitrogen, #NP0009), and NuPAGE LDS sample buffer at 1X final concentration (Invitrogen,

#NP0007)—and boiled at 95°C for 10 minutes before the samples were loaded in the NuPAGE 4%–12%

Bis-Tris Gels (Invitrogen, #NP0335BOX & #NP0336BOX) for further resolution and western blot

analysis. 500mM of purified GST-PUM2, GST-FMRP and HIS-PUM1 were used as input controls; GST-

PUM2 and GST-FMRP were visualized using rabbit α-GST antibody (1:1000 [Cell Signaling

Technologies, #2625]), whereas for HIS-PUM1 we used rabbit α-HIS Tag antibody (1:3000, [Cell

Signaling Technologies, #2365]).

Protein quantification and western blot analysis. Patient-derived lymphoblastoid, fibroblast cell lines,

and control cell lines were collected at 6×10^6 cell confluence and processed for protein extraction. For

mouse tissues, we processed either half of the whole brain (the other half was processed for RNA

extraction, see below) or the entire hippocampus, cortex, or cerebellum for protein extraction. Mouse

tissues or cell pellets were subsequently lysed with modified RIPA buffer consisting of 25 mM Tris-HCL,

pH 7.6, 150 mM NaCl, 1.0% Tween 20, 1.0% sodium deoxycholate, 0.1% SDS, completed with 1X Xpert

Protease and 1X Phosphatase Inhibitor Cocktail Solutions (GenDepot, #P3100-100 & #P3200-020). Cells

were lysed by pipetting them up and down with a p1000 tip and then placed on ice for 20 min followed by

centrifugation at 14,800 rpm at 4°C for 25 minutes. Mouse brain tissues were pipetted up and down by

syringe needles—starting from an 18G 1½" (Becton Dickson, #305196), moving to 21G 1½" (Becton

Dickson, #305167) and finally to a 26G 1½" (Becton Dickson, #305111) needle—until the lysate passed

through the needle smoothly. Proteins were quantified by Pierce BCA Protein Assay Kit (Thermo

Scientific, # PI23225) and their absorbance measured by NanoDrop OneC (Thermo Scientific). Proteins

were resolved by high resolution NuPAGE 4%–12% Bis-Tris Gel (Invitrogen, #NP0335BOX &

#NP0336BOX) according to the manufacturer's instructions. All the blots were acquired on the G:BOX

Chemi XX9 machine (Syngene; Frederick, MD) using GeneSys software 1.6.5.0. Gel exposures were

determined by the software.

Antibodies used for western blot experiments were: goat α-PUM1 [1:2500, (Bethyl Laboratories,

#A300-201A)], rabbit α-PUM1 [1:2000, (Abcam Cambridge, #ab92545)], rabbit α-PUM2 [1:2000,

(Bethyl Laboratories, # A300-202A)], rabbit α-FMRP [1:1000, (Abcam Cambridge, #ab17722)], rabbit

α-AGO2 [1:1000, (Abcam Cambridge, #ab32381)], rabbit α-NeuN (Rbfox3) [1:1000, (Thermo

Scientific, #PA5-37407)], rabbit α-CNOT1 [1:1000, (Cell Signaling Technology, #44613)], rabbit α-

MOV10 [1:2000, (Bethyl Laboratories, #A301-571A)], and mouse α-GAPDH [1:10000, (Millipore,

#CB1001)].

HEK293T cell culture and maintenance. Human embryonic kidney immortalized 293T (HEK293T)

cells were grown in DMEM (GenDepot, #CM002-320) supplemented with 10% of heat-inactivated fetal

bovine serum (FBS [GenDepot, #F0901-050) and 1% penicillin/streptomycin (GenDepot, #CA005-010).

All cells were incubated at 37 °C in a humidified chamber supplemented with 5% CO_2. HEK293T cells

were later processed according to the needs of specific experiments (described below).

RNA extraction and quantitative real-time PCR (qPCR). Human fibroblast, lymphoblastoid, and

respective control cell lines were harvested at 6 X 10^6 confluence prior to RNA extraction. For mouse

tissues, half of the whole brain (the other half was processed for protein extraction, see above) or the

entire hippocampus, cortex, or cerebellum were processed for RNA extraction. The RNA was collected

for both human cells, mouse brain and brain region tissues using the miRNeasy kit (QIAGEN, # 217004)

according to the manufacturer's instructions. RNA was quantified using NanoDrop OneC (Thermo Fisher

Scientific). cDNA was synthesized using Quantitect Reverse Transcription kit (QIAGEN, # 205313)

starting from 1 μg of RNA. Quantitative RT-polymerase chain reaction (qRT-PCR) experiments were

performed using the CFX96 Touch Real-Time PCR Detection System (Bio-Rad Laboratories, Hercules)

with PowerUP SYBR Green Master Mix (Applied Biosystems, #A25743). Real-time PCR runs were

analyzed using the comparative C_T method normalized against the housekeeping human gene *GAPDH* or

mouse *Gapdh*, depending on the experiment (Vandesompele *et al*, 2002).

RNA Immunoprecipitation (RIP). HCT116 wild-type and PUM1-KO were obtained from (Lee *et al*,

2016). HCT116-PUM1-KO cells were grown in McCoy's 5A Media (Merck, # M4892) supplemented

with 10% of heat-inactivated fetal bovine serum (FBS [GenDepot, #F0901-050]) and 1%

penicillin/streptomycin (GenDepot, #CA005-010) for 16 hours in a 10-cm plate (80% confluency) and

transfected with 6μg of Myc-PUM1-WT, Myc-PUM1-R1147W, and Myc-PUM1-T1035S. After 48 hours

cells were collected and lysed in Polysome Lysis Buffer (PLB) (10mM HEPES-KOH pH 7, 100mM KCl,

5mM MgCl$_2$, 25 mM EDTA, 0.5% NP40, 2mM Dithiothreitol, 0.2 mg/ml Heparin) supplemented with

50U/ml of RNAse OUT^{TM} recombinant ribonuclease inhibitor (Thermo Fischer Scientific, # 10777019),

50U/ml of SuperaseIN^{TM} RNAse inhibitor (Thermo Fischer Scientific, #AM2694) and 1X of Xpert

Protease Inhibitor Cocktail Solutions (GenDepot, #P3100-100) for 20 minutes in ice; cells were

centrifuged at maximum speed at 4°C to remove cellular debris. 5μg of goat α-PUM1 (Bethyl

Laboratories, #A300-201A) or α-Goat-IgG with 50μl of Dynabeads^{TM} Protein G (Invitrogen, #10004D)

were incubated overnight at 4°C on a rotisserie tube rotator; 100μl of lysate were removed to serve as

input. 3mg of lysate with 10μl of Dynabeads^{TM} Protein G (Invitrogen, #10004D) were precleared for 30

minutes at 4°C on a rotisserie tube rotator; the pre-cleared lysate was incubated with antibody-conjugated

beads overnight at 4°C on a rotisserie tube rotator. Beads were washed 3 times with NT2 buffer (50mM

of Tris-HCl, 300mM NaCl, 1mM EDTA, 1% NP40, 0.1% SDS, 0.5% Na-deoxycholate supplemented

with 1X of Xpert Protease Inhibitor Cocktail Solutions). Beads and input were resuspended in 100μl of

NT2 buffer supplemented with 80U of RNAse OUT and 30μg of proteinase K and incubated for 30

minutes at 50°C to eliminate the proteins. RNA was extracted from the immunoprecipitated samples and

their input; 200μl of RLT buffer was added using RNeasy Mini Kit (Qiagen, #74104) according to the

manufacturer's instruction. cDNA was synthetized using SuperScript^{TM} II Reverse Transcripatse (Thermo

Fisher Scientific, #18064014) according to the manufacturer's instructions. Quantitative RT-polymerase

chain reaction (qRT-PCR) experiments were performed using the CFX96 Touch Real-Time PCR

Detection System (Bio-Rad Laboratories, Hercules) with PowerUP SYBR Green Master Mix (Applied

Biosystems, #A25743). The oligos used for qPCR are listed in **Table S7**.

MicroRNA library construction and sequencing. Library preparation and microRNA sequencing were

performed by LC Sciences (Houston, TX) according to the following criteria. Total RNA was extracted

from cerebellum of WT and *Pum1$^{-/-}$* male and female at 10 weeks of age using the miRNeasy kit

(QIAGEN, # 217004) according to the manufacturer's instructions. The total RNA quality and quantity

were assessed with Bioanalyzer 2100 (Agilent Technologies, Santa Clara) with RIN number > 7.0.

Approximately 1 μg of total RNA were used to prepare the small RNA library according to the protocol

of TruSeq Small RNA Sample Prep Kits (Illumina, San Diego). Then the single-end sequencing 50bp was

performed on an Illumina Hiseq 2500 at LC Sciences (Hangzhou, China) following the vendor's

recommended protocol.

Labelling scheme:

Sample	Gender	Genotype
LCS7846_GV_KO1_F	Female	KO
LCS7846_GV_KO1_M	Male	KO
LCS7846_GV_KO2_M	Male	KO
LCS7846_GV_KO3_F	Female	KO
LCS7846_GV_KO3_M	Male	KO
LCS7846_GV_KO4_F	Female	KO
LCS7846_GV_WT1_M	Male	WT
LCS7846_GV_WT2_F	Female	WT
LCS7846_GV_WT2_M	Male	WT
LCS7846_GV_WT3_F	Female	WT
LCS7846_GV_WT3_M	Male	WT

MicroRNA sequencing bioinformatic analysis. Raw reads were subjected to an in-house program,

ACGT101-miR (LC Sciences, Houston), to remove adapter dimers, junk, common RNA families (rRNA,

tRNA, snRNA, snoRNA), and repeats. Subsequently, unique sequences of 18–26 nucleotides in length

were mapped to specific species precursors in miRBase 22.0 (http://www.mirbase.org/) by BLAST search

to identify known miRNAs and novel 3p- and 5p-derived miRNAs. Length variation at both 3' and 5'

ends and one mismatch inside of the sequence were allowed in the alignment. The unique sequences

mapping to specific species of mature miRNAs in hairpin arms were identified as known miRNAs. The

unique sequences mapping to the other arm of known specific species precursor hairpins opposite the

annotated mature miRNA-containing arm were considered to be novel 5p- or 3p-derived miRNA

candidates. The remaining sequences were mapped to other selected species precursors (with the

exclusion of specific species) in miRBase 22.0 by BLAST search, and the mapped pre-miRNAs were

further BLASTed against the specific species genomes to determine their genomic locations. The last two

were also defined as known miRNAs. The unmapped sequences were BLASTed against the specific

genomes, and the hairpin RNA structures containing sequences were predicted from the flank 80 nt

sequences using RNAfold software (http://rna.tbi.univie.ac.at/cgi-bin/RNAWebSuite/RNAfold.cgi). The

criteria for secondary structure prediction were: (1) number of nucleotides in one bulge in stem ($\leq$12), (2)

number of base pairs in the stem region of the predicted hairpin ($\geq$16), (3) cutoff of free energy (kCal/mol

$\leq$-15), (4) length of hairpin (up and down stems + terminal loop $\geq$50), (5) length of hairpin loop ($\leq$20), (6)

number of nucleotides in one bulge in mature region ($\leq$8), (7) number of biased errors in one bulge in

mature region ($\leq$4), (8) number of biased bulges in mature region ($\leq$2), (9) number of errors in mature

region ($\leq$7), (10) number of base pairs in the mature region of the predicted hairpin ($\geq$12), (11) percent of

mature region in stem ($\geq$80).

Gene Set Enrichment Analysis (GSEA). GSEA was performed as previously described (Subramanian *et*

al., 2005). The cumulative distribution function was conducted by performing 1000 random gene-set

membership assignments. A nominal p-value < 0.01 and an FDR < 0.25 were used to assess the

significance of the enrichment score (ES). HITS-CLIP data, and the respective rank, were obtained from

the literature and were initially acquired as follows: Pum1 and Pum2 from neonatal murine brains (Zhang

et al., 2017), Fmrp from cerebellum, cortex, and hippocampus together (Maurin *et al.*, 2018), Ago2 from

neocortex at embryonic day 13 (Chi *et al.*, 2009), Rbfox3 from mouse brain (age not specified) (Weyn-

Vanhentenryck *et al.*, 2014), Nova from mouse brain (age not specified) (Zhang *et al.*, 2010), Ptpb2 from

neocortex at embryonic day 18.5 (Licatalosi *et al.*, 2012), Mbnl2 from hippocampus at 8-12 weeks of age

(Charizanis *et al.*, 2012), and Apc from mouse brain at embryonic day 14 (Preitner *et al.*, 2014).

Gene ontology analyses. Gene ontology analyses were performed with David Gene Ontology (GO). For

Figure 2B, Figure S11B and **C** only categories with FDR<0.05 were considered; while for **Figure S10**

only categories with FDR<0.01 were considered. David GO for the Pum1 interactome in **Figure 2B**

considered the entire interactome as background. For the GO regarding the HITS-CLIP targets shared

among Pum1, Pum2, Fmrp, Ago2, and Rbfox3 (**Figure S11B** and **C**), we considered the entire set of all

targetomes together as background. Regarding the Synaptic (Syn) GO analysis (**Figure S10D**), brain-

expressed genes were used as background (Koopmans *et al.*, 2019).

Myc and GST cloning procedure with in vitro immunoprecipitation (IP) assays. Human *PUM1* full-

length cDNA was amplified by PCR and subcloned in a pRK5 plasmid containing the Myc tag sequence

(Addgene, pRK5-Myc-Parkin #17612) at the N-terminal by using SalI (New England Biolabs, # R3138S)

and NotI (New England Biolabs, #R0189S) restriction enzymes to replace *Parkin* with *PUM1*. For GST,

the human full-length *PUM1* cDNA was, again, subcloned first in the pRK5 plasmid containing the GST

tag sequence (Addgene, pRK5-HA GST RagC wt, #19304) at the N-terminal by using SalI and NotI

restriction enzymes to replace *RagC* with *PUM1*. Human *FMRP*, *AGO2* and *CNOT1*, full-length cDNA

were cloned and contain the GST tag sequence at the N-terminal, as described for GST-*PUM1*.

To introduce the T1035S or R1147W mutations we used the QuikChange II XL Multi Site-Directed

Mutagenesis kit (Agilent Technologies, #200521). The primers for the single mutagenesis experiments

were designed by QuikChange software (Stratagene, San Diego, https://www.genomics.agilent.com/

primerDesignProgram.jsp).

For IP, HEK293T cells were seeded in 6-well plates for 24 h and then transfected with 250 ng of

either WT or mutant PUM1 plasmid with one of the interactors using the jetPRIME Transfection Reagent

(Polyplus transfection, #55-132) as per the manufacturer's protocol. pRK5-Myc empty plasmid (no

cDNA) was used as a negative control. After 48 h, the cells were collected and processed for

immunoprecipitation. Protein lysates were incubated overnight at 4°C with mouse α-Myc antibody

(1:400, [Cell Signaling Technologies, #2276]) on a rotisserie tube rotator. The next day, the beads were

washed four times with an IP lysis buffer and resuspended in 40 μL elution buffer (lysis buffer, NuPAGE

10X Reducing Agent [Invitrogen, #NP0009], NuPAGE LDS sample buffer 4X [Invitrogen, #NP0007])

and boiled at 95°C for 10 minutes before loading the samples in NuPAGE 4%–12% Bis-Tris Gels

(Invitrogen, #NP0335BOX & #NP0336BOX) for further resolution and western blot analysis. Antibodies:

mouse α-Myc antibody [1:2000, (Cell Signaling Technologies, #2276)], rabbit α-GST antibody [1:1000,

(Cell Signaling Technologies, #2625)], mouse α-GAPDH [1:10000, (Millipore, #CB1001)].

Patient-derived cell lines. Primary fibroblasts from the *PUM1* PADDAS patient (9-year-old female) and

the age- and sex-matched controls (three different 9-year-old female) were generated as previously

described (Gennarino *et al.*, 2018). Briefly, cells were isolated from skin biopsies taken from the patient

or age-matched controls using standard methodology (Barch and Association of Cytogenetic Technology,

1991) and placed in a transport medium (Ham's F10, Thermo Scientific, #11550043). The skin specimen

was later removed from the transport medium using a sterile technique (in a Class II biohazard cabinet)

and transferred to a sterile Petri dish where it was cut into small pieces (< 0.5 mm) using sterile scalpel

blades. These pieces were transferred to the lower surface of a 25 cm^2 culture flask (6-8 pieces per flask)

which had been pre-moistened with 1-2 mL of AmnioMAX Complete Medium (Thermo Scientific,

#11269016) supplemented with 1% penicillin/streptomycin (GenDepot, #CA005-010). Cell cultures were

maintained at 37 °C in a humidified incubator supplemented with 5% CO_2. When cell growth was

observed around the edges of the tissue, usually 3 to 5 days later, 2 to 3 mL of AmnioMAX Complete

Medium were added. Once growth was established and the tissue was anchored to the flask, another 8 mL

of AmnioMAX Complete Medium was added. Thereafter, the medium was renewed every 3 to 4 days

until ready for sub-culturing.

Lymphoblastoid cells from *PUM1* PRCA patients (female 59 and 58 years old, respectively) and the

age- and sex-matched controls (three different 58-year-old female) were generated as previously

described (Gennarino *et al.*, 2018). Briefly, lymphoblastoid suspension cell cultures were grown in RPMI

1640 medium (Invitrogen, #11875093) supplemented with 10% heat-inactivated fetal bovine serum

(Atlanta Biological, Flowery Branch, #S11195H) and 1% penicillin/streptomycin (GenDepot, #CA005-

010). Cell cultures were maintained at 37°C in a humidified incubator supplemented with 5% CO_2.

Medium was renewed every 2 to 3 days.

1000

Fibroblast patient-derived cell lines transfection. Fibroblasts from age- and sex-matched healthy

controls and from a female PADDAS patient were seeded at 80% of confluency in 6-well plates

(~150.000 cells/well). The day after, 500ng of pRK5-CMV-Myc-Pum1 or pRK5-CMV-Myc-Empty

plasmids were transfected in antibiotic-free DMEM (GenDepot, #CM002-320) using Lipofectamine LTX

with Plus Reagent (Thermo Fisher, #15338030) according to the manufacturer's protocol. After 5 hours

we replaced the media with new complete DMEM supplemented with 10% of heat-inactivated fetal

bovine serum (FBS [GenDepot, #F0901-050]) and 1% penicillin/streptomycin (GenDepot, #CA005-010).

Cells were incubated at 37 °C in a humidified chamber supplemented with 5% CO2 and collected after 72

hours for RNA and protein extraction.

1010

Primers. For the qPCR analysis to unambiguously distinguish spliced cDNA from genomic DNA

contamination, specific exon primers were designed to amplify across introns of the gene tested. The

primers for all genes tested were designed with Primer3 (Koressaar & Remm, 2007; Untergasser *et al*,

2012). Cloning primers were manually designed to amplify the longest spliced gene isoform tested; if

there was more than one isoform according to the UCSC Genome Browser (https://genome.ucsc.edu/), we

chose the longest. See Table S7 for primer sequences.

1017

Ethical statement and mouse strains. All animal procedures were approved by the Institutional Animal

Care and Use Committee at Columbia University, New York under the protocol AC-AAAU8490. Mice

were maintained on a 12-hr light, 12-hr dark cycle with regular chow and water ad libitum. Pum1 knock-

out mice were generated as previously described (Chen *et al.*, 2012). C57BL/6J wild-type mice were

purchased from Jackson Laboratory and maintained as described above. For brain dissection, mice were

anesthetized with isoflurane, and the brain rapidly removed from the skull and lysed in the appropriate

buffer according to the experiment (see Materials and Methods Details).

1025

Experimental design. For protein and RNA quantification from patient-derived cell lines, we used values

from at least six independent experiments with three biological replicates for each experiment. At every

stage of the study, the experimenter was blinded to the identity of control and patient-derived cell lines.

For example, for the data regarding both human patient-derived cell lines and mice, Experimenter #1

made a list of samples and controls to be tested, and Experimenter #2 randomized this list and re-labeled

the tubes; Experimenter #2 was the only person with the key to identify the samples. These samples were

then distributed to Experimenter #3 to culture the cells, then to Experimenter #1 to perform western blots

and qRT-PCR, and lastly Experimenters #1 and #4 analyzed the data. Only then was the key applied to

identify the samples.

For mouse experiments, the experimenters were randomized and blinded as described above. The

number of animals used and sex, and the specific statistical tests used, are indicated for each experiment

in the figure legends. Sample size was based on previous experience using the same mice (Gennarino *et*

al., 2015).

1039

Software and statistical analysis. Statistical significance was analyzed using GraphPad Prism 8

(https://www.graphpad.com/ scientific-software/prism/) and Excel Software (Microsoft). All data are

presented as mean ± SEM. Statistical details for each experiment can be found in the figures and the

legends. The range of expression levels in qPCR was determined from at least six independent

experiments with three biological replicates by calculating the standard deviation of the ΔCt (Pfaffl,

2001). The range of expression levels in western blots was determined from at least six independent

experiments with at least six biological replicates. P values were calculated by Student's T-test or analysis

of variance with Tukey's post hoc analysis. For the IP in **Figure 6A** and protein quantification in patient

cell lines in **Figure 6C**, we had only one PADDAS patient, so the repeated experiments were technical

replicates rather than biological replicates. We therefore calculated the statistical significance based on

these technical replicates in comparison to the three biological replicates (i.e., healthy controls).

1051

1052 **Study approval.** PADDAS and PRCA patient cell lines are the same as those reported previously

1053 (Gennarino *et al.*, 2018). The consent form for each subject specifically allows for sharing of medical

1054 information and physical exam findings; the sharing of cell lines from the PADDAS and PRCA subjects

1055 and the controls was approved under the Columbia University Medical Center IRB-AAAS7401

1056 (Y01M00) and the Baylor College of Medicine IRB H-34578.

1057

1058 **Data availability.**

1059 IP mass spec and quantitative proteomics raw data are available at: "pending".

1060 microRNA sequencing raw data are available at "pending"

1061

References
Agarwal V, Bell GW, Nam JW, Bartel DP (2015) Predicting effective microRNA target sites in
mammalian mRNAs. *Elife* 4
Bah MG, Rodriguez D, Cazeneuve C, Mochel F, Devos D, Suppiej A, Roubertie A, Meunier I, Gitiaux C,
Curie A *et al* (2020) Deciphering the natural history of SCA7 in children. *Eur J Neurol* 27: 2267-
2276
Blackinton JG, Keene JD (2014) Post-transcriptional RNA regulons affecting cell cycle and proliferation.
Semin Cell Dev Biol 34: 44-54
Blewett NH, Goldstrohm AC (2012) A eukaryotic translation initiation factor 4E-binding protein
promotes mRNA decapping and is required for PUF repression. *Mol Cell Biol* 32: 4181-4194
Bohn JA, Van Etten JL, Schagat TL, Bowman BM, McEachin RC, Freddolino PL, Goldstrohm AC
(2018) Identification of diverse target RNAs that are functionally regulated by human Pumilio
proteins. *Nucleic Acids Res* 46: 362-386
Bonnemason-Carrere P, Morice-Picard F, Pennamen P, Arveiler B, Fergelot P, Goizet C, Hellegouarch
M, Lacombe D, Plaisant C, Raclet V *et al* (2019) PADDAS syndrome associated with hair dysplasia
caused by a de novo missense variant of PUM1. *Am J Med Genet A* 179: 1030-1033
Charizanis K, Lee KY, Batra R, Goodwin M, Zhang C, Yuan Y, Shiue L, Cline M, Scotti MM, Xia G *et*
al (2012) Muscleblind-like 2-mediated alternative splicing in the developing brain and dysregulation
in myotonic dystrophy. *Neuron* 75: 437-450
Chartier-Harlin MC, Kachergus J, Roumier C, Mouroux V, Douay X, Lincoln S, Levecque C, Larvor L,
Andrieux J, Hulihan M *et al* (2004) Alpha-synuclein locus duplication as a cause of familial
Parkinson's disease. *Lancet* 364: 1167-1169

Chen D, Zheng W, Lin A, Uyhazi K, Zhao H, Lin H (2012) Pumilio 1 suppresses multiple activators of
p53 to safeguard spermatogenesis. *Curr Biol* 22: 420-425

Chi SW, Zang JB, Mele A, Darnell RB (2009) Argonaute HITS-CLIP decodes microRNA-mRNA
interaction maps. *Nature* 460: 479-486

Conboy JG (2017) Developmental regulation of RNA processing by Rbfox proteins. *Wiley Interdiscip*
Rev RNA 8

Cox J, Mann M (2008) MaxQuant enables high peptide identification rates, individualized p.p.b.-range
mass accuracies and proteome-wide protein quantification. *Nat Biotechnol* 26: 1367-1372

Cox J, Neuhauser N, Michalski A, Scheltema RA, Olsen JV, Mann M (2011) Andromeda: a peptide
search engine integrated into the MaxQuant environment. *J Proteome Res* 10: 1794-1805

Darnell JC, Richter JD (2012) Cytoplasmic RNA-binding proteins and the control of complex brain
function. *Cold Spring Harb Perspect Biol* 4: a012344

Dassi E (2017) Handshakes and Fights: The Regulatory Interplay of RNA-Binding Proteins. *Front Mol*
Biosci 4: 67

De Maio A, Yalamanchili HK, Adamski CJ, Gennarino VA, Liu Z, Qin J, Jung SY, Richman R, Orr H,
Zoghbi HY (2018) RBM17 Interacts with U2SURP and CHERP to Regulate Expression and Splicing
of RNA-Processing Proteins. *Cell Rep* 25: 726-736 e727

Di Grazia A, Marafini I, Pedini G, Di Fusco D, Laudisi F, Dinallo V, Rosina E, Stolfi C, Franze E, Sileri
P *et al* (2021) The Fragile X Mental Retardation Protein Regulates RIPK1 and Colorectal Cancer
Resistance to Necroptosis. *Cell Mol Gastroenterol Hepatol* 11: 639-658

Elguindy MM, Mendell JT (2021) NORAD-induced Pumilio phase separation is required for genome
stability. *Nature* 595: 303-308

Enwerem, III, Elrod ND, Chang CT, Lin A, Ji P, Bohn JA, Levdansky Y, Wagner EJ, Valkov E,
Goldstrohm AC (2021) Human Pumilio proteins directly bind the CCR4-NOT deadenylase complex
to regulate the transcriptome. *RNA* 27: 445-464

Friend K, Campbell ZT, Cooke A, Kroll-Conner P, Wickens MP, Kimble J (2012) A conserved PUF-
Ago-eEF1A complex attenuates translation elongation. *Nat Struct Mol Biol* 19: 176-183

Fu J, Peng L, Tao T, Chen Y, Li Z, Li J (2019) Regulatory roles of the miR-200 family in
neurodegenerative diseases. *Biomed Pharmacother* 119: 109409

Gehman LT, Meera P, Stoilov P, Shiue L, O'Brien JE, Meisler MH, Ares M, Jr., Otis TS, Black DL
(2012) The splicing regulator Rbfox2 is required for both cerebellar development and mature motor
function. *Genes Dev* 26: 445-460

Genetic Modifiers of Huntington's Disease C (2015) Identification of Genetic Factors that Modify
Clinical Onset of Huntington's Disease. *Cell* 162: 516-526

Gennarino VA, D'Angelo G, Dharmalingam G, Fernandez S, Russolillo G, Sanges R, Mutarelli M,
Belcastro V, Ballabio A, Verde P *et al* (2012) Identification of microRNA-regulated gene networks
by expression analysis of target genes. *Genome Res* 22: 1163-1172

Gennarino VA, Palmer EE, McDonell LM, Wang L, Adamski CJ, Koire A, See L, Chen CA, Schaaf CP,
Rosenfeld JA *et al* (2018) A mild PUM1 mutation is associated with adult-onset ataxia, whereas
haploinsufficiency causes developmental delay and seizures. *Cell* 172: 924-936 e911

Gennarino VA, Singh RK, White JJ, De Maio A, Han K, Kim JY, Jafar-Nejad P, di Ronza A, Kang H,
Sayegh LS *et al* (2015) Pumilio1 haploinsufficiency leads to SCA1-like neurodegeneration by
increasing wild-type Ataxin1 levels. *Cell* 160: 1087-1098

Giurgiu M, Reinhard J, Brauner B, Dunger-Kaltenbach I, Fobo G, Frishman G, Montrone C, Ruepp A (2019) CORUM: the comprehensive resource of mammalian protein complexes-2019. *Nucleic Acids Res* 47: D559-D563

Goldstrohm AC, Hall TMT, McKenney KM (2018) Post-transcriptional Regulatory Functions of Mammalian Pumilio Proteins. *Trends Genet* 34: 972-990

Hentze MW, Castello A, Schwarzl T, Preiss T (2018) A brave new world of RNA-binding proteins. *Nat Rev Mol Cell Biol* 19: 327-341

Imaizumi T, Mogami Y, Okamoto N, Yamamoto-Shimojima K, Yamamoto T (2019) De novo 1p35.2 microdeletion including PUM1 identified in a patient with sporadic West syndrome. *Congenit Anom (Kyoto)* 59: 193-194

Jin Z, Feng H, Liang J, Jing X, Zhao Q, Zhan L, Shen B, Cheng X, Su L, Qiu W (2020) FGFR3 big up tri, open7-9 promotes tumor progression via the phosphorylation and destabilization of ten-eleven translocation-2 in human hepatocellular carcinoma. *Cell Death Dis* 11: 903

Kacher R, Lejeune FX, Noel S, Cazeneuve C, Brice A, Humbert S, Durr A (2021) Propensity for somatic expansion increases over the course of life in Huntington disease. *Elife* 10

Kedde M, van Kouwenhove M, Zwart W, Oude Vrielink JA, Elkon R, Agami R (2010) A Pumilio-induced RNA structure switch in p27-3' UTR controls miR-221 and miR-222 accessibility. *Nat Cell Biol* 12: 1014-1020

Keene JD (2007a) Biological clocks and the coordination theory of RNA operons and regulons. *Cold Spring Harb Symp Quant Biol* 72: 157-165

Keene JD (2007b) RNA regulons: coordination of post-transcriptional events. *Nat Rev Genet* 8: 533-543

Keene JD, Lager PJ (2005) Post-transcriptional operons and regulons co-ordinating gene expression. *Chromosome Res* 13: 327-337

Kenny PJ, Kim M, Skariah G, Nielsen J, Lannom MC, Ceman S (2020) The FMRP-MOV10 complex: a translational regulatory switch modulated by G-Quadruplexes. *Nucleic Acids Res* 48: 862-878

Kenny PJ, Zhou H, Kim M, Skariah G, Khetani RS, Drnevich J, Arcila ML, Kosik KS, Ceman S (2014) MOV10 and FMRP regulate AGO2 association with microRNA recognition elements. *Cell Rep* 9: 1729-1741

Khalil B, Morderer D, Price PL, Liu F, Rossoll W (2018) mRNP assembly, axonal transport, and local translation in neurodegenerative diseases. *Brain Res* 1693: 75-91

Kim K, Hessl D, Randol JL, Espinal GM, Schneider A, Protic D, Aydin EY, Hagerman RJ, Hagerman PJ (2019) Association between IQ and FMR1 protein (FMRP) across the spectrum of CGG repeat expansions. *PLoS One* 14: e0226811

Koopmans F, van Nierop P, Andres-Alonso M, Byrnes A, Cijsouw T, Coba MP, Cornelisse LN, Farrell RJ, Goldschmidt HL, Howrigan DP *et al* (2019) SynGO: An Evidence-Based, Expert-Curated Knowledge Base for the Synapse. *Neuron* 103: 217-234 e214

Koressaar T, Remm M (2007) Enhancements and modifications of primer design program Primer3. *Bioinformatics* 23: 1289-1291

Lai KL, Liao YC, Tsai PC, Hsiao CT, Soong BW, Lee YC (2019) Investigating PUM1 mutations in a Taiwanese cohort with cerebellar ataxia. *Parkinsonism Relat Disord* 66: 220-223

Lee EJ, Seo E, Kim JW, Nam SA, Lee JY, Jun J, Oh S, Park M, Jho EH, Yoo KH *et al* (2020) TAZ/Wnt-beta-catenin/c-MYC axis regulates cystogenesis in polycystic kidney disease. *Proc Natl Acad Sci U S A* 117: 29001-29012

Lee S, Kopp F, Chang TC, Sataluri A, Chen B, Sivakumar S, Yu H, Xie Y, Mendell JT (2016) Noncoding RNA NORAD Regulates Genomic Stability by Sequestering PUMILIO Proteins. *Cell* 164: 69-80

Lee Y, Samaco RC, Gatchel JR, Thaller C, Orr HT, Zoghbi HY (2008) miR-19, miR-101 and miR-130 co-regulate ATXN1 levels to potentially modulate SCA1 pathogenesis. *Nat Neurosci* 11: 1137-1139

Licatalosi DD, Yano M, Fak JJ, Mele A, Grabinski SE, Zhang C, Darnell RB (2012) Ptbp2 represses adult-specific splicing to regulate the generation of neuronal precursors in the embryonic brain. *Genes Dev* 26: 1626-1642

Marrero E, Rossi SG, Darr A, Tsoulfas P, Rotundo RL (2011) Translational regulation of acetylcholinesterase by the RNA-binding protein Pumilio-2 at the neuromuscular synapse. *J Biol Chem* 286: 36492-36499

Mauger O, Lemoine F, Scheiffele P (2016) Targeted Intron Retention and Excision for Rapid Gene Regulation in Response to Neuronal Activity. *Neuron* 92: 1266-1278

Maurin T, Lebrigand K, Castagnola S, Paquet A, Jarjat M, Popa A, Grossi M, Rage F, Bardoni B (2018) HITS-CLIP in various brain areas reveals new targets and new modalities of RNA binding by fragile X mental retardation protein. *Nucleic Acids Res* 46: 6344-6355

Miles WO, Tschop K, Herr A, Ji JY, Dyson NJ (2012) Pumilio facilitates miRNA regulation of the E2F3 oncogene. *Genes Dev* 26: 356-368

Muraro NI, Weston AJ, Gerber AP, Luschnig S, Moffat KG, Baines RA (2008) Pumilio binds para mRNA and requires Nanos and Brat to regulate sodium current in Drosophila motoneurons. *J Neurosci* 28: 2099-2109

Orr HT, Zoghbi HY (2007) Trinucleotide repeat disorders. *Annu Rev Neurosci* 30: 575-621

Otasek D, Morris JH, Boucas J, Pico AR, Demchak B (2019) Cytoscape Automation: empowering workflow-based network analysis. *Genome Biol* 20: 185

Pfaffl MW (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* 29: e45

Preitner N, Quan J, Nowakowski DW, Hancock ML, Shi J, Tcherkezian J, Young-Pearse TL, Flanagan JG (2014) APC is an RNA-binding protein, and its interactome provides a link to neural development and microtubule assembly. *Cell* 158: 368-382

Rappsilber J, Mann M, Ishihama Y (2007) Protocol for micro-purification, enrichment, pre-fractionation and storage of peptides for proteomics using StageTips. *Nat Protoc* 2: 1896-1906

Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, Vilo J (2019) g:Profiler: a web server for functional enrichment analysis and conversions of gene lists (2019 update). *Nucleic Acids Res* 47: W191-W198

Ravanidis S, Kattan FG, Doxakis E (2018) Unraveling the Pathways to Neuronal Homeostasis and Disease: Mechanistic Insights into the Role of RNA-Binding Proteins and Associated Factors. *Int J Mol Sci* 19

Rousseaux MWC, Tschumperlin T, Lu HC, Lackey EP, Bondar VV, Wan YW, Tan Q, Adamski CJ, Friedrich J, Twaroski K *et al* (2018) ATXN1-CIC Complex Is the Primary Driver of Cerebellar Pathology in Spinocerebellar Ataxia Type 1 through a Gain-of-Function Mechanism. *Neuron* 97: 1235-1243 e1235

Rovelet-Lecrux A, Hannequin D, Raux G, Le Meur N, Laquerriere A, Vital A, Dumanchin C, Feuillette S, Brice A, Vercelletto M *et al* (2006) APP locus duplication causes autosomal dominant early-onset Alzheimer disease with cerebral amyloid angiopathy. *Nat Genet* 38: 24-26

Salcedo-Arellano MJ, Dufour B, McLennan Y, Martinez-Cerdeno V, Hagerman R (2020) Fragile X
syndrome and associated disorders: Clinical aspects and pathology. *Neurobiol Dis* 136: 104740
Shah S, Molinaro G, Liu B, Wang R, Huber KM, Richter JD (2020) FMRP Control of Ribosome
Translocation Promotes Chromatin Modifications and Alternative Splicing of Neuronal Genes Linked
to Autism. *Cell Rep* 30: 4459-4472 e4456
Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T
(2003) Cytoscape: a software environment for integrated models of biomolecular interaction
networks. *Genome Res* 13: 2498-2504
Shevchenko A, Tomas H, Havlis J, Olsen JV, Mann M (2006) In-gel digestion for mass spectrometric
characterization of proteins and proteomes. *Nat Protoc* 1: 2856-2860
Singh K, Gaur P, Prasad S (2007) Fragile x mental retardation (Fmr-1) gene expression is down regulated
in brain of mice during aging. *Mol Biol Rep* 34: 173-181
Singh K, Prasad S (2008) Differential expression of Fmr-1 mRNA and FMRP in female mice brain
during aging. *Mol Biol Rep* 35: 677-684
Smidak R, Sialana FJ, Kristofova M, Stojanovic T, Rajcic D, Malikovic J, Feyissa DD, Korz V, Hoeger
H, Wackerlig J *et al* (2017) Reduced Levels of the Synaptic Functional Regulator FMRP in Dentate
Gyrus of the Aging Sprague-Dawley Rat. *Front Aging Neurosci* 9: 384
Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy
SL, Golub TR, Lander ES *et al* (2005) Gene set enrichment analysis: a knowledge-based approach for
interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* 102: 15545-15550
Takeguchi R, Takahashi S, Kuroda M, Tanaka R, Suzuki N, Tomonoh Y, Ihara Y, Sugiyama N, Itoh M
(2020) MeCP2_e2 partially compensates for lack of MeCP2_e1: A male case of Rett syndrome. *Mol
Genet Genomic Med* 8: e1088
Tan Q, Zoghbi HY (2019) Mouse models as a tool for discovering new neurological diseases. *Neurobiol
Learn Mem* 165: 106902
Temme C, Simonelig M, Wahle E (2014) Deadenylation of mRNA by the CCR4-NOT complex in
Drosophila: molecular and developmental aspects. *Front Genet* 5: 143
Tereshchenko A, Magnotta V, Epping E, Mathews K, Espe-Pfeifer P, Martin E, Dawson J, Duan W,
Nopoulos P (2019) Brain structure in juvenile-onset Huntington disease. *Neurology* 92: e1939-e1947
Thul PJ, Akesson L, Wiking M, Mahdessian D, Geladaki A, Ait Blal H, Alm T, Asplund A, Bjork L,
Breckels LM *et al* (2017) A subcellular map of the human proteome. *Science* 356
Trumbach D, Prakash N (2015) The conserved miR-8/miR-200 microRNA family and their role in
invertebrate and vertebrate neurogenesis. *Cell Tissue Res* 359: 161-177
Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, Mann M, Cox J (2016) The Perseus
computational platform for comprehensive analysis of (prote)omics data. *Nat Methods* 13: 731-740
Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, Rozen SG (2012) Primer3--new
capabilities and interfaces. *Nucleic Acids Res* 40: e115
Uyhazi KE, Yang Y, Liu N, Qi H, Huang XA, Mak W, Weatherbee SD, de Prisco N, Gennarino VA,
Song X *et al* (2020) Pumilio proteins utilize distinct regulatory mechanisms to achieve
complementary functions required for pluripotency and embryogenesis. *Proc Natl Acad Sci U S A*
117: 7851-7862

Van Etten J, Schagat TL, Hrit J, Weidmann CA, Brumbaugh J, Coon JJ, Goldstrohm AC (2012) Human
Pumilio proteins recruit multiple deadenylases to efficiently repress messenger RNAs. *J Biol Chem*
287: 36370-36383
Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F (2002) Accurate
normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal
control genes. *Genome Biol* 3
Wamsley B, Jaglin XH, Favuzzi E, Quattrocolo G, Nigro MJ, Yusuf N, Khodadadi-Jamayran A, Rudy B,
Fishell G (2018) Rbfox1 Mediates Cell-type-Specific Splicing in Cortical Interneurons. *Neuron* 100:
846-859 e847
Weidmann CA, Raynard NA, Blewett NH, Van Etten J, Goldstrohm AC (2014) The RNA binding
domain of Pumilio antagonizes poly-adenosine binding protein and accelerates deadenylation. *RNA*
20: 1298-1319
Weyn-Vanhentenryck SM, Mele A, Yan Q, Sun S, Farny N, Zhang Z, Xue C, Herre M, Silver PA, Zhang
MQ *et al* (2014) HITS-CLIP and integrative modeling define the Rbfox splicing-regulatory network
linked to brain development and autism. *Cell Rep* 6: 1139-1152
Zamore PD, Williamson JR, Lehmann R (1997) The Pumilio protein binds RNA through a conserved
domain that defines a new class of RNA-binding proteins. *RNA* 3: 1421-1433
Zhang C, Frias MA, Mele A, Ruggiu M, Eom T, Marney CB, Wang H, Licatalosi DD, Fak JJ, Darnell RB
(2010) Integrative modeling defines the Nova splicing-regulatory network and its combinatorial
controls. *Science* 329: 439-443
Zhang M, Chen D, Xia J, Han W, Cui X, Neuenkirchen N, Hermes G, Sestan N, Lin H (2017) Post-
transcriptional regulation of mouse neurogenesis by Pumilio proteins. *Genes Dev* 31: 1354-1369

1276

Figure 1

Cluster 1
Cdc16
Cdc23
Cdc27
Anapc4
Anapc1
Anapc7
Anapc5
Anapc2
Fzr1
Sh3kbp1

Cluster 2
Ppp1r12a
Ppp1r12b
Ppp1r12c
Ppp6r2
Ppp1r21
Ppp1r16b

Cluster 3
Kdm1a
Zmym2
Hdac2

Cluster 4
Mllt1
Mllt3
Aff4

Cluster 5
Hnrnpul1
Dhx15
Wdr20
Acin1
Rbm39
Snrnp200
Ddx3x
Ddx3y
Ddx17
Sf3b3
Wdr48

Cluster 6
Eif3l
Eif2s3x/y
Eif3b
Eif2s2
Ctif
Eif5
Eif3f
Eif4g3

Cluster 7
Ago2
Cnot1
Fmrp
Rbfox3

Cluster 8
Matr3
Pfkp
Pfkm
Pfkl
Pfkfb2
Pspc1

Cluster 9
Pex5l
Atg7
Snx2
Snx4
Madd
Fga
Fgb
Fgg
Rab11fip5
Pck2

Cluster 10
Rictor
Rptor
Tsc1
Tsc2
Camkk2
Cad
Lrpprc
Lppr4

Figure 2

A

Legend:
- ● Cerebellar tissue
- ● Cortical tissue
- ● Hippocampal tissue
- ● Rest of brain
- ● Whole brain (WB)
- ● WB, Cerebellum
- ● WB, Cortex
- ● WB, Hippocampus
- ● Cerebellum, Cortex
- ● Cerebellum, Hippocampus
- ● Cortex & Hippocampus
- ● WB, Cerebellum, Cortex
- ● WB, Cerebellum, Hippocampus
- ● WB, Cortex, Hippocampus
- ● Cerebellum, Cortex, Hippocampus

B

DAVID Gene Ontology analysis

Whole brain: mRNA metabolism, K-11 ubiquit., E2/E3 ubiquit., RNA silencing, RNA transport, APC/C

Hippocampus: mRNA metabolism, K-11 ubiquit., E2/E3 ubiquit., APC/C, RNA silencing

Cortex: mRNA splicing, mRNA 3'UTR binding, E2/E3 ubiquit., APC/C, RNA silencing, K-11 ubiquit.

Cerebellum: mRNA 3'UTR binding, E2/E3 ubiquit., APC/C, RNA silencing, ribosomal proteins, K-11 ubiquit.

fold enrichment

Gene count: 50 40 30 20 10

Figure 3

Figure 4

A Cerebellum   **B** Hippocampus   **C** Cortex

Figure 5

Figure 6

Figure 7

A

mRNA levels of HITS-CLIP targets shared by PUM1, PUM2, FMRP, AGO2, and RBFOX3 in PADDAS and PRCA patient-derived cells normalized against their respective age- and sex-matched controls

■ R1147W PADDAS (S1) ■ T1035S PRCA (S2 & S3)

----- Controls   *p<0.05; **p<0.01; ***p<0.001

B

Protein quantification after Myc-PUM1 overexpression in PADDS patient-derived cell lines

■ Controls
■ R1147W (S1) Empty
■ R1147W (S1) Myc-*PUM1*

**p<0.01; ***p<0.001

C

mRNA quantification of top 15 dysregulated shared targets (panel A) after Myc-PUM1 overexpression in PADDS patient-derived cell lines

■ R1147W (S1) Empty ■ R1147W (S1) Myc-*PUM1*

----- Controls   *p<0.05; **p<0.01; ***p<0.001

Figure S1

A

Mass Spec scatter plot - mouse brain

IP:IgG enriched interactors   IP:Pum1 enriched interactors

Pum1
Anapc1
Fzr1
Cnot1
Rptor
Fmr1
Ago2

-Log Student's T-test p-value IP:Pum1 – IP:IgG

Student's T-test difference IP:Pum1 – IP:IgG

B

Principal component analysis mouse brain

IP:IgG
IP:Pum1

PCA 2 (2.3%)

PCA 1 (38.2%)

C

IP:Pum1 from WT mouse brain
compared to IgG at 10 weeks of age

Post-IP (10%)

Input (1%) IP:IgG IP:Pum1 IgG Pum1

Pum1   120

Pum1   120

Longer exposure time to visualize
the Input and the post-IPs

Figure S2

Figure S3

A

Brain region specific Quantitative proteomics
- - - mean Pum1 interactors

C

IP:Pum1 from mouse brain regions – Input (1%)

D

Western blot different brain regions at 10 weeks of age

B

Pum1 interactors from different brain regions Quantitative proteomics

Figure S4

A

Input | GST resin pull down | Input | GST resin pull down

Beads | FMRP | PUM1 | FMRP | PUM1 | FMRP+PMU1 | Beads | FMRP | PUM1 | FMRP | PUM1 | FMRP+PMU1

Pum1

Fmrp

—190

—97

B

Input | GST resin pull down | Input | GST resin pull down

Beads | PUM2 | PUM1 | PUM2 | PUM1 | PUM2+PMU1 | Beads | PUM2 | PUM1 | PUM2 | PUM1 | PUM2+PUM1

Pum1
Pum2

—190
—146

C

IP:Pum1 from mouse brain

Input (1%) | IP:IgG | IP:Pum1

Pum1 — 120
Rptor — 150
Cad — 243
Anapc1 — 240
Fzr1 — 50

D

Co-IPs from mouse brain

Input (1%) | IP:IgG | IP:Anapc1
Anapc1 — 240
Pum1 — 120

Input (1%) | IP:IgG | IP:Fzr1
Fzr1 — 50
Pum1 — 120
IP:IgG

Input (1%) | IP:IgG | IP:Rptor
Rptor — 150
Pum1 — 120

Input (1%) | IP:IgG | IP:Cad
Cad — 243
Pum1 — 120
IP:IgG

Figure S5

A

	Input (1%)	IP:IgG	IP:Pum1	
Pum1				120
Pum2				120
Fmrp				71
Ago2				97
Rbfox3				50
Cnot1				240
Mov10				107

Pum1⁻/⁻

B

IP against IgG and Pum1
mouse brain

	1% Input	IgG −	Pum1 −	IgG +	Pum1 +	
RNase:						
Pum1						120
Ago1						97
Ago3						97

C

without RNase

WT — Input | IP:IgG | IP:Pum2
Pum1⁻/⁻ — Input | IP:IgG | IP:Pum2

Pum1 — 120
Pum2 — 120

WT — Input | IP:IgG | IP:Ago2
Pum1⁻/⁻ — Input | IP:IgG | IP:Ago2

Pum1
Ago2

WT — Input | IP:IgG | IP:Fmrp
Pum1⁻/⁻ — Input | IP:IgG | IP:Fmrp

Pum1
Fmrp

WT — Input | IP:IgG | IP:Cnot1
Pum1⁻/⁻ — Input | IP:IgG | IP:Cnot1

Pum1
Cnot1

with RNase

WT — Input | IP:IgG | IP:Pum2
Pum1⁻/⁻ — Input | IP:IgG | IP:Pum2

Pum1 — 120
Pum2 — 120

WT — Input | IP:IgG | IP:Ago2
Pum1⁻/⁻ — Input | IP:IgG | IP:Ago2

Pum1 — 120
Ago2 — 97

WT — Input | IP:IgG | IP:Fmrp
Pum1⁻/⁻ — Input | IP:IgG | IP:Fmrp

Pum1 — 120
Fmrp — 71

WT — Input | IP:IgG | IP:Cnot1
Pum1⁻/⁻ — Input | IP:IgG | IP:Cnot1

Pum1 — 120
Cnot1 — 240

Figure S6

Figure S7

mRNA levels – 10 weeks of age

Figure S8

A

PCA miRNAseq

- wild-type male
- wild-type female
- *Pum1*⁻/⁻ male
- *Pum1*⁻/⁻ female

PC1

PC2

B

● miRNAs up in females and down in males   ● miRNAs down in females and up in males

Volcano plot Female

-p value (t-test)

-log2 expression

Volcano plot Male

-p value (t-test)

-log2 expression

Figure S9

RNA quantification from male and female cerebellum at 10 weeks of age.
Genes plotted here sorted by number of co-bound miRNAs from **Figure 4D** with at least 25% expression change

Table showing the order of gene represented in the figure above starting from
***Gapdh* as represented in Figure 4E**

Position in the graph	Gene	Male	Female	Error_Male	Error_Female	Co-bound miRNAs
1	*Gapdh*	1.000	1.000	0.065	0.188	11
2	*Aak1*	1.222	0.714	0.099	0.096	11
3	*Plekhm3*	1.283	0.764	0.132	0.102	11
4	*Zbtb20*	1.268	0.636	0.185	0.093	10
5	*Ago3*	1.485	0.686	0.122	0.102	10
6	*Dgkh*	1.511	0.751	0.106	0.103	10
7	*Fto*	1.407	0.618	0.132	0.097	10
8	*Grin2b*	1.517	0.749	0.071	0.117	10
9	*Lpp*	1.353	0.689	0.124	0.123	10
10	*Stx17*	1.266	0.682	0.108	0.098	10
11	*Acvr2b*	1.359	0.668	0.108	0.092	9
12	*Ap5m1*	2.366	0.456	0.282	0.101	9
13	*Chl1*	1.506	0.691	0.121	0.094	9
14	*Gpr161*	1.457	0.735	0.118	0.113	9
15	*Kcnn3*	1.477	0.694	0.145	0.094	9
16	*Klf7*	1.262	0.670	0.103	0.103	9
17	*Nfat5*	1.228	0.598	0.099	0.086	9
18	*Plxna4*	0.712	0.680	0.051	0.092	9
19	*Slc1a2*	1.344	0.713	0.125	0.111	9
20	*Slc8a1*	0.904	0.417	0.094	0.056	9
21	*Ston2*	1.669	0.690	0.148	0.114	9
22	*Tsc22d2*	1.319	0.743	0.146	0.137	9
23	*Xkr4*	1.823	0.573	0.093	0.105	9
24	*A1cf*	2.821	0.232	0.314	0.031	8
25	*Aff4*	1.053	1.304	0.116	0.238	8
26	*Clstn2*	1.340	0.768	0.062	0.109	8
27	*Cnnm2*	1.475	0.605	0.097	0.099	8
28	*Csnk1a1*	1.698	0.641	0.115	0.087	8
29	*Ctdspl2*	1.749	0.624	0.108	0.108	8
30	*Dcaf7*	1.685	0.750	0.141	0.132	8
31	*Fmnl3*	1.705	0.620	0.246	0.106	8
32	*Frmd4a*	1.394	0.654	0.085	0.123	8
33	*Grin2a*	1.459	0.755	0.148	0.112	8
34	*Hipk2*	1.381	0.662	0.094	0.088	8
35	*Kif26b*	1.588	0.627	0.181	0.094	8
36	*Klf12*	2.030	0.668	0.202	0.101	8
37	*Lmln*	1.435	0.595	0.143	0.091	8
38	*Lrrc40*	1.262	0.679	0.126	0.113	8
39	*Myo5a*	0.958	0.707	0.089	0.102	8
40	*Nav2*	1.377	0.658	0.135	0.097	8
41	*Psd3*	1.493	0.728	0.076	0.110	8
42	*Ptbp2*	1.463	0.177	0.068	0.177	8
43	*Rimkla*	1.368	0.779	0.092	0.112	8
44	*Snx30*	1.833	0.781	0.147	0.112	8
45	*Taok1*	1.290	0.660	0.078	0.088	8
46	*Zbtb10*	1.624	0.686	0.102	0.092	8
47	*Apbb2*	1.278	0.660	0.094	0.097	8
48	*Acap2*	1.374	0.697	0.107	0.093	8
49	*Acer2*	1.544	0.610	0.187	0.083	8
50	*Aebp2*	1.219	1.904	0.085	0.269	8

Figure S10

Figure S11

A

Pum1 against Ptpb2
ES = 0.36
FDR = 0.73
Rank at max 4153 (41.8th)

Pum1 against Nova
ES = 0.41
FDR = 0.71
Rank at max 3476 (31.8th)

Pum1 against Apc
ES = 0.24
FDR = 1
Rank at max 160 (61.3th)

Pum1 against Mbnl2
ES = 0.29
FDR = 1
Rank at max 2247 (46.8th)

Enrichment score (ES)

Negative control ranked target lists

B

Biological functions
all 175 shared targets

fold enrichment

cardiac contraction
cell recognition
synaptic transmission
spinal cord development
neurotransmitter secretion
regulation of heart rate
axon extension
regulation of sensory pain
learning and memory
visual learning
adult locomotor behavior
neuromuscular balance
embryo development
chemical synaptic transmission
axon guidance
axonogenesis
nervous system development
neuron projection
brain development

C

Cellular localization
all 175 shared targets

fold enrichment

astrocyte projection
catenin complex
axonal growth cone
neuron projection
presynaptic membrane
synaptic vesicle
dendritic shaft
growth cone
terminal button
synaptic vesicle
axon
dendritic spine
neuron projection
postsynaptic density
neuronal cell body

D

Shared miRNAs

291
Fmrp

38
Pum1

17
Pum2

1
1
3

Figure S12

A IP:PUM1 from fibroblast patient-derived cell lines

Pre-IP ctrl
Pre-IP S1
IP:IgG ctrl
IP:PUM1 ctrl
IP:IgG S1
IP:PUM1 S1
Post-IP:IgG ctrl
Post-IP:PUM1 S1

PUM1 — 120

B IP:PUM1 from lymphoblastoid patient-derived cell lines

Pre-IP ctrl
IP:IgG ctrl
IP:PUM1 ctrl
Post-IP:IgG ctrl

PUM1 — 120

C

RNA immunoprecipitation

Fold change on site occupancy

WT
$PUM1^{-/-}$
$PUM1^{-/-}$ + MYC-PUM1
$PUM1^{-/-}$ + MYC-PUM1 (T1035S)
$PUM1^{-/-}$ + MYC-PUM1 (R1147W)

- E2F3 (blue)
- CDK1 (yellow)
- E2F4 (grey)

$*p<0.05; **p<0.01$

Figure S13

Fold Change (FC) IP = $\dfrac{[\text{IP}^{X}/(\text{Input}^{X}/\text{GAPDH}^{X})]}{[\text{IP}^{\text{PUM1}}/(\text{Input}^{\text{PUM1}}/\text{GAPDH}^{\text{PUM1}})]}$

WT or Empty (E)   R1147W PADDAS (S1)   T1035S PRCA (S2 & S3)

*$p<0.05$; **$p<0.01$; ***$p<0.001$

Figure S14

A

mRNA quantification PUM1-specific targets
compared to their respective age- and sex-matched controls

B

PUM1 mRNA levels after Myc-*PUM1*
overexpression in PADDS patient-derived cell lines