Two distinct mechanisms underlie dosage sensitivity in Pumilio1-associated diseases

Salvatore Botta[1,2], Nicola de Prisco[1], Alexei Chemiakine[1], Maximilian Cabaj[1], Vicky L. Brandt[1], Purvi Patel[3], Rajesh K. Soni[3], and Vincenzo A. Gennarino[1,4,5,6,7]

[1]Department of Genetics and Development, Columbia University Irving Medical Center, New York, NY.

[2]Department of Translational Medical Science, University of Campania Luigi Vanvitelli, Caserta, Italy.

[3]Proteomics and Macromolecular Crystallography Shared Resource, Herbert Irving Comprehensive Cancer Center, Columbia University Irving Medical Center, New York, NY.

[4]Departments of Pediatrics and Neurology, Columbia University Irving Medical Center, New York, NY.

[5]Columbia Stem Cell Initiative, Columbia University Irving Medical Center, New York, NY.

[6]Initiative for Columbia Ataxia and Tremor, Columbia University Irving Medical Center, New York, NY.

[7]Corresponding author: vag2138@cumc.columbia.edu

Keywords

Abstract

The RNA-binding protein Pumilio1 (PUM1) is associated with distinct diseases: a mild, late-onset ataxia and a severe, wide-ranging neurodevelopmental syndrome. The ataxia patients retain 75% of normal PUM1 levels, the syndromic patients ~50%, but why this should cause such different phenotypes was unclear. Target dysregulation did not explain the difference, so we developed a PUM1 interactome for the mouse brain. This revealed RNA-binding proteins involved in aspects of RNA metabolism not previously associated with PUM1 (e.g., alternative splicing), as well as proteins in unexpected pathways (e.g., cyclosome). We focused on a subset of RNA-binding proteins and found that the effects of PUM1 loss on these proteins and their targets varied with brain region and sex. In patient cells, mice, and *in vitro*, mild PUM1 loss affected only PUM1-specific targets, whereas PUM1 haploinsufficiency destabilized these native complexes and altered expression of shared targets. We propose that dosage sensitivity can involve distinct mechanisms, and that there may be more intimate coordination among RNA-binding proteins than is readily evinced by more traditional *in vitro* approaches.

Impact statement

Most studies of RNA-binding proteins focus on identifying their targets, but identifying their protein

partners *in vivo* provides important insights into their functions and loss of function.

Introduction

To respond with agility to a specific perturbation, cells must modify their protein repertoire, much

of which is controlled at the post-transcriptional level by RNA-binding proteins (RBPs). RBPs regulate

RNA localization, transport, translation, splicing, and decay, and they orchestrate hundreds of pathways

that are responsible for proper biological functions (Hentze *et al*, 2018; Keene, 2007b). RBPs can

coordinate or compete with each other or exert mutual influence (Dassi, 2017), and they also interact with

microRNAs (miRNAs), which are small non-coding RNAs that suppress the expression of their target

genes by binding to a short complementary seed region in the 3' UTRs of mRNAs. The RBP and

microRNA machinery are particularly important in neurons, whose plasticity demands a rapid local

(axonal) response to stimuli despite distance from the nucleus (Mauger *et al*, 2016). Disruptions in RBP

function underlie several complex neurological disorders: for example, large CAG expansions in FMRP

cause Fragile X syndrome, but milder "premutations" in FMRP cause adult-onset Fragile X-associated

tremor and ataxia syndrome (FXTAS) (Lukong *et al*, 2008). The RBPs TDP43 and FUS are both

involved in amyotrophic lateral sclerosis, with TDP43 mutations also linked to frontotemporal dementia

(Humphrey *et al*, 2020; Ravanidis *et al*, 2018). Despite recent interest in these and other RBPs implicated

in neuronal function, we still know relatively little about RBP interactions and regulation (Darnell &

Richter, 2012; Khalil *et al*, 2018; Ravanidis *et al.*, 2018).

We recently discovered that the RBP Pumilio1 (PUM1) is involved in two distinct human diseases

(Gennarino *et al*, 2018; Gennarino *et al*, 2015). PUM1-associated developmental delay and seizures

(PADDAS) is a wide-ranging neurodevelopmental disorder, while PUM1-related cerebellar ataxia

(PRCA) is a slowly progressive, pure ataxia with onset in mid-life. Although the severity of the PRCA

and PADDAS phenotypes tracks with the levels of functional PUM1 (75% of wild-type levels in

PRCA, 40-60% in PADDAS), precisely what is happening at the molecular level was unclear. PUM1

comprises a highly conserved RNA-binding domain composed of eight tandem repeats known as Puf

homology domains (HDs), and the mutation that produces mild disease (PRCA), T1035S, lies within this

domain and impairs RNA binding—but the most severe PADDAS mutation (R1147W) is outside this

domain and does not seem to impair RNA binding (Gennarino *et al.*, 2018; Gennarino *et al.*, 2015).

Moreover, the PADDAS phenotype is wide-ranging and variable. The most severe mutation, R1147W,

causes small stature, physical dysmorphologies, cognitive and language delays, motor impairments, and

seizures starting in infancy. The enormous difference in phenotypic outcomes between mutations in the

RNA-binding domain and outside of it, plus the fact that the six then-known PUM1 targets were

upregulated to similar extents in the PRCA and PADDAS patient cell lines (Gennarino *et al.*, 2018), led

us to hypothesize that PRCA is caused by deregulation of PUM1 targets, but that PADDAS results from

disruption of PUM1's interactions with its protein partners along with de-repression of the targets of these

complexes.

To test this hypothesis would require identifying PUM1 interactors in the mouse brain. Although a

great deal is known about the PUMILIO/FBF (PUF) family of RBPs, of which PUM1 and its homolog

PUM2 are members, surprisingly little is known about PUM1 function in the postnatal mammalian brain.

PUM1 was initially discovered in *Drosophila* for its contributions to embryonic patterning (Lin &

Spradling, 1997) and stem cell maintenance (Leeb *et al*, 2014). Studies in fly, worm, and germ cells show

that Pumilio proteins regulate mRNA targets through at least three distinct mechanisms: they can

collaborate with the miRNA machinery to repress targets (Friend *et al*, 2012; Kedde *et al*, 2010; Miles *et*

al, 2012); they can recruit the CCR4-NOT deadenylase complex to shorten poly(A) tails and promote

mRNA degradation (Temme *et al*, 2014; Van Etten *et al*, 2012; Weidmann *et al*, 2014); and they can

antagonize poly(A)-binding proteins to repress translation (Goldstrohm *et al*, 2018). PUM1/2 can also

activate certain mRNAs (Bohn *et al*, 2018; Uyhazi *et al*, 2020).

Protein interactions in general, and those of PUF family members specifically, can be organism-,

transcript-, and even condition-specific (Miller & Olivas, 2011). We therefore took an unbiased approach

by using *in vivo* proteomics to identify PUM1's native partners. Because *Drosophila* studies show that

Pumilio can change its protein partners in different neuronal types (Muraro *et al*, 2008), we first examined

the brain as a whole and then repeated our analyses in three brain regions that most highly express PUM1.

We then determined the effect of PUM1 loss on a focused subset of RBP interactors that were the most

highly connected within the interactome, using *PUM1* heterozygous and knockout mice and patient-

derived cell lines. We also identified targets shared among these interactors and examined their responses

to PUM1 insufficiency in mice and cell lines from patients bearing the T1035S and R1147W mutations.

Our results support our hypothesis and underscore the need to examine all the interactions an RBP

engages in, not just its targets. (Note that, because we deal with both mouse and human proteins in this

paper, we will refer to the mouse proteins with lower case (e.g., Pum1) and the human proteins in all caps

(PUM1).)

Results

Establishing the Pumilio1 interactome across the adult mouse brain

To identify Pum1 protein partners, we performed co-immunoprecipitation (IP) for Pum1 on whole

brains from 10-week-old wild-type mice, followed by liquid chromatography with tandem mass

spectrometry (LC-MS/MS, or simply mass spec); we used IP against IgG (IP-IgG) as a negative control

(see Materials and Methods). Because of the intrinsic variability of the mass spec technique, we combined

and homogenized whole brains from two 10-week-old wild-type mice per sample (1 female and 1 male),

aliquoting half of each sample for IP against either Pum1 or IgG, then performed six biological replicates

(six samples, 12 mice total) for each mass spec experiment against IP-Pum1 and IP-IgG. Post-IP western

blot detected no residual Pum1 from brain tissue, confirming that our protocol recovers virtually all of

Pum1 (**Figure 1–figure supplement 1A**).

Principal component analysis (PCA) readily separated the IP-Pum1 interactors from the control IP-

IgG (**Figure 1–figure supplement 1B**). To increase the likelihood that our candidates would prove to be

genuine interactors, the proteins had to have at least two unique peptides present in at least five out of six

IP-Pum1 samples and also show at least an average >1.5-fold difference between IP-Pum1 and IP-IgG in

label-free quantification intensity (LFQ-intensity, see Materials and Methods). Our analysis yielded 412

putative Pum1 interactors (**Figure 1A** and **Supplementary Table 1**). We incorporated mammalian

protein-protein interaction data from CORUM and the Human Protein Atlas by using g:GOSt (g:Profiler;

see Materials and Methods) (Raudvere *et al*, 2019) to reveal 22 clusters of functionally related proteins

(**Figure 1A**). Among the putative interactors were those that have been previously identified, supporting

the validity of our approach. For example, our mass spec identified Nanos1, the mammalian homologue

of *Drosophila* Nanos, which was one of the first proteins found to interact with Pumilio during fly

development (Wang *et al*, 2001); it also identified Pum2 and Fmrp, which associate with Pum1 in neural

progenitor cells (Zhang *et al*, 2017). Some of the identified proteins belong to families previously found

to interact with PUM1 *in vitro*, such as Cnot1, which is the central scaffold of the CCR4-NOT complex

(Enwerem *et al*, 2021; Van Etten *et al.*, 2012). Translation initiation factors (cluster 3) have been found to

cooperate with Puf proteins in other species (Blewett & Goldstrohm, 2012), and human PUM2-mediated

repression was found to require PABPC1, while our mass spec yielded Pabpc4 (Weidmann *et al.*, 2014).

Nanos1 and Pabpc4 have numerous connections among the clusters. Cluster 8, which includes proteins

that have been associated with RNA fate regulation, was the most strongly interconnected with other

clusters.

Most of the proteins, of course, have never been associated with PUM1. Among the candidate

interactors with the greatest fold-change were many members of the anaphase-promoting complex

(cluster 21). This E3 ubiquitin ligase complex targets substrates for degradation and is essential for

neuronal survival (Almeida *et al*, 2005; Gieffers *et al*, 1999; Sivakumar & Gorbsky, 2015). We performed

preliminary validation by reciprocal co-IP between Pum1 and Anapc1, which scaffolds the APC/C

complex, and between Pum1 and Fzr1, which forms its catalytic pocket (Sivakumar & Gorbsky, 2015)

(**Figure 1–figure supplement 1C, D**). Cluster 11 was also unexpected, as it contains members of both

mTORC1 (Rptor, Cad, Tsc1, Tsc2) and mTORC2 (Rictor) pathways, which are involved in a variety of

cellular functions from promoting cell growth to autophagy (Liu & Sabatini, 2020). Reciprocal co-IP with

Rptor and Cad confirmed interactions with Pum1 (**Figure 1–figure supplement 1C, D**).

Given the plethora of putative interactors to investigate, we repeated the mass spec experiments on

the cerebellum, hippocampus, and cortex, where Pum1 is most highly expressed (Gennarino *et al.*, 2015),

to see which pathways emerged as predominant. Because our mass spec protocol requires at least 5 mg of

protein, we combined cerebellar tissue from eight 10-week-old wild-type mice (4 female and 4 male),

performing the experiment in triplicate (for a total of 24 mice); we did the same for cortical tissue. For the

hippocampus, we combined tissues from ten wild-type mice (5 of each sex) for three experiments (30

mice; see Materials and Methods). As before, we selected proteins with at least two unique peptides and

an average of >1.5 fold-change between IP-Pum1 and the IP-IgG. PCA again showed a clear separation

of Pum1 samples from the IgG and indicated that the Pum1 hippocampal and cortical interactomes are

more similar to each other than to that of the cerebellum (**Figure 1–figure supplement 2A**). We

identified 854 putative Pum1 interactors in the cerebellum, 423 in the hippocampus, and 598 in the cortex

(**Supplementary Table 1**). 467 were unique to the cerebellum, 140 to the hippocampus, 229 to the

cortex, and 154 unique to the rest of the brain (i.e., excluding these three regions). Only 88 candidates

were shared among these three brain regions and the whole brain (**Figure 1–figure supplement 2B**,

yellow dots; **Figure 1–figure supplement 2C**). To our surprise, the only brain region to show interaction

between Pum1 and Pum2 was the cortex (**Figure 1–figure supplement 2C**).

The region-specific proteins expanded the list of Pum1 interactors in the pathways described initially

(**Figure 1–figure supplement 2D-F**). For example, Cnot1 and Cnot2 turned up in all three brain regions,

but Cnot10 appears to be cortex-specific (**Figure 1–figure supplement 2F**). There were many proteins

involved in translation initiation, with Eif3b being specific to the hippocampus (**Figure 1–figure**

supplement 2B). Rbfox1 was specific to the cortex, and Rbfox2 to the cerebellum (**Figure 1–figure**

supplement 2B), consistent with previous work showing that Rbfox1 mediates cell-type-specific splicing

in cortical interneurons (Wamsley *et al*, 2018) and that Rbfox2 is needed for cerebellar development

(Gehman *et al*, 2012). The brain-region-specific analysis did not yield many additional interactors in the

APC/C (**Figure 1–figure supplement 2D**) or mTOR pathways (**Figure 1–figure supplement 2E**). It did,

however, reveal many pathways that have never been associated with Pum1 (**Supplementary Table 1**).

Taken as a whole, these data suggest that the RNA-silencing, CCR4-NOT, translational initiation, polyA

regulation, and splicing machineries are fine-tuned according to brain region or neuronal cell types,

whereas the APC/C and mTOR machineries remain fairly consistent, at least across these three brain

regions.

This is not to say that certain interactions are necessarily limited to these brain regions. They may

occur in regions not tested; they may occur across the whole brain, but at levels that did not reach the

threshold we set. It could also be that some interactors are expressed more highly in specific brain

regions, though this is not the determining factor for the interaction with Pum2, which is expressed to an

equivalent degree in all three brain regions (**Figure 1–figure supplement 2G**).

We then performed DAVID Gene Ontology analysis for the hits from both the whole brain and each

brain region (**Figure 1B** and **Supplementary Tables 2**). The dominant categories were ubiquitin ligases

(anaphase-promoting complex [APC/C], E2/E3 and Kll-linked ubiquitin) and RBPs involved in various

aspects of RNA metabolism (RNA silencing, 3'UTR binding, mRNA stability, transport, and splicing).

We decided to prioritize the RBPs in cluster 8 (**Figure 1A**)—Fmrp and Ago2 (involved in RNA

silencing), Pum2, Cnot1, and Rbfox3 (an alternative splicing factor)—for three reasons. First, this cluster

was the most highly interconnected with other clusters; second, RNA-related categories were prominent

in the gene ontology analyses for both whole brain and all three brain regions; and third, these RBPs have

been well studied and would allow us to more readily test the consequences of PUM1 loss.

Pum1 associates with Pum2, Fmrp, Ago2, and Cnot1 in the absence of mRNA

We first performed co-IP followed by western blot to confirm Pum1 association with Pum2, Fmrp,

Ago2, Rbfox3 and Cnot1 (**Figure 2A**, *left panel*). Because Fmrp and Ago2 both bind Mov10 *in vitro*

(Kenny *et al*, 2020; Kenny *et al*, 2014), we also blotted for murine Mov10, even though it did not appear

among our mass spec results. Mov10 was pulled down with Pum1, likely in concert with Fmrp (**Figure**

2A). Reciprocal co-IP using antibodies against Pum2, Ago2, Fmrp, Rbfox3, Cnot1, and Mov10 followed

by western blot confirmed these associations (**Figure 2B**). To verify the specificity of the antibodies, we

co-IPed Pum1 and blotted for all six RBPs in *Pum1$^{-/-}$* mouse brain and detected none of them (**Figure 2–**

figure supplement 1A). As negative controls, we tested other proteins associated with the RISC complex

(Ago1 and Ago3) that did not appear in our mass spec data, and our co-IP experiments found no

interactions (data not shown).

To exclude the possibility that the co-IPs simply recovered other proteins that are co-bound to target

RNAs but are not part of the same complex as the protein of interest, we treated the mouse brain samples

with RNase and verified that no detectable RNA remained (see Materials and Methods). Pum1 still

associated with Pum2, Fmrp, Ago2, and Cnot1 in the absence of mRNA, but not Rbfox3 or Mov10

(**Figure 2A**, *right panel*). Interestingly, a previous *in vitro* study observed PUM1 interaction with AGO2

only in the presence of RNA in T-REx 293 cells (Sternburg et al., 2018). We repeated the RNase

experiments in HEK293T cells, which confirmed our results (except for RBFOX3, which was not
detectable in these cells) (**Figure 2C**).

201
Some interactions among these RBPs require PUM1
We next performed reciprocal co-IP in wild-type and *Pum1*$^{-/-}$ 10-week-old mouse brains for Pum2,
Ago2, Fmrp, Cnot1, Rbfox3, and Mov10. Pum2 interacted with Cnot1 only in the presence of Pum1
(**Figure 3A**). Unexpectedly, Fmrp associated with Mov10 and Rbfox3 only in the presence of Pum1
(**Figure 3B-D**). Fmrp and Cnot1 did not associate in the wild-type brain (**Figure 3B**, *left*) but they did so
in the absence of Pum1 (**Figure 3B**, *right*). Ago2 associated with Pum1, Fmrp, Cnot1 and Mov10 in wild-
type brain, but in the absence of Pum1 it no longer interacted with Fmrp or Mov10 (**Figure 3E**). Mov10
associated with Fmrp but not with Ago2 (**Figure 3D** and **E**). Rbfox3 associated with Cnot1 (**Figure 3C**)
but not *vice versa* (**Figure 3F**). IPs against Pum2, Ago2, Fmrp, Cnot1, Rbfox3, and Mov10 with and
without RNase treatment in wild-type and *Pum1*$^{-/-}$ mouse brains (**Figure 3–figure supplement 1A-F**)
confirmed that, in the absence of Pum1, these interactions require RNA.
These results underscore the importance of examining RBP interactions *in vivo*, in specific contexts,
with and without RNase treatment. For example, previous work showed that AGO2, PABPC1, GW182
and CNOT1 interact with nuclear EGFR to regulate VEGF signaling in tumor cell lines (Dittmann *et al*,
2017), but neither GW182 nor EGFR turned up in either our whole-brain or regional interactomes. In fact,
none of GW182's three mammalian paralogues (Tnrc6A, Tnrc6B, or Tnrc6C), which interact with all
four human AGO proteins, was pulled down in our mass spec against Pum1 (**Supplementary Table 1**).
In murine neurons, therefore, Pum1 associates with Pum2, Fmrp, Ago2, and Cnot1, with or without
RNA (**Figure 3A-F** and **Figure 3–figure supplement 1A-F**). Pum1 seems to be required for association
between Fmrp, Mov10 and Rbfox3, and between Ago2 and Fmrp, Mov10, and Cnot1. In the absence of
Pum1, the associations between Pum2, Fmrp, Ago2 and Cnot1 require RNA (**Figure 3–figure**
supplement 1A-F).

224
Pum1 loss alters RBP levels by brain region and sex
If these are *bona fide* interactors of Pum1, then we would expect loss of Pum1 to alter their
abundance. *Pum1* heterozygous and knockout mice showed changes in the quantities of Pum2, Ago2, and
Mov10 proteins across the brain (**Figure 4–figure supplement 1A**), but only *Pum2* showed changes in
mRNA levels (**Figure 4–figure supplement 1B**). Interestingly, Ago2 and Mov10 levels fell only in male
mice (**Figure 4–figure supplement 1A**), so we decided to pursue this sex difference by brain region. We
quantified Pum2, Fmrp, Ago2, Rbfox3, Cnot1 and Mov10 in the cerebellum, cortex, and hippocampus.

We first measured Pum1 mRNA and protein levels to confirm the reduction of Pum1 in our *Pum1*$^{+/-}$ or

Pum1$^{-/-}$ mice in each brain region (**Figure 4A-C,** and **Figure 4–figure supplement 2A**). Pum2 protein

levels rose in all three brain regions, as did its mRNA (**Figure 4A-C,** and **Figure 4–figure supplement**

2A).

Fmrp protein expression was upregulated in all three brain regions in male KO mice, but in female

KO cerebella Fmrp was reduced by almost 70% (**Figure 4A**). (We should note that sex differences in

Fmrp expression have been previously reported (Singh *et al*, 2007; Singh & Prasad, 2008).) Three other

proteins also showed divergent responses to Pum1 loss according to sex and brain region: Ago2 (up in

female but down in male cerebella, with the opposite pattern in hippocampus and cortex), Rbfox3 (in the

hippocampus), and Cnot1 (in the cerebellum) (**Figure 4A-C**). None of them showed any changes in

mRNA levels (**Figure 4–figure supplement 2A**), despite the fact that *Fmr1* and *Cnot1*, like *Pum2*, have a

Pumilio Response Element (PRE) (Zamore *et al*, 1997)—sometimes known as a Pumilio Recognition

Element (PRE) or Pumilio Binding Element (PBE)—in their 3'UTR. It may be that the loss of Pum1

causes sex- and region-specific reorganization of these complexes, or that there are additional sex-specific

Pum1 interactors.

Pum1 loss dysregulates Ago2 and the microRNA machinery in mouse cerebella by sex

Since loss of Pum1 function has deleterious effects on the cerebellum in both mice (Gennarino *et al.*,

2015) and humans (Gennarino *et al.*, 2018), we were curious to know whether the divergence of Ago2

protein levels in males and females extended to cerebellar miRNAs. We performed miRNAseq.

Hierarchical heatmap clustering of miRNA expression in *Pum1*$^{-/-}$ and WT male and female cerebella at 10

weeks of age revealed that the expression of 166 miRNAs (**Supplementary Table 3**) diverged between

the two sexes in parallel with Ago2 expression (**Figure 4D**).

To examine the functional consequences of this Ago2/miRNA dysregulation, we looked at the

expression of downstream targets that are co-bound by those miRNAs in 10-week-old wild-type and

Pum1$^{-/-}$ male and female cerebella. We first selected all the miRNAs with at least a 25% change in

expression in either direction, for a total of 19 miRNAs. Using TargetScan and CoMeTa (Agarwal *et al*,

2015; Gennarino *et al*, 2012) we identified 6832 putative targets that are co-bound by at least 2 out of 19

possible miRNAs. To reduce this list to a more manageable size for testing expression changes by qPCR,

we prioritized targets that are co-bound by at least 8 miRNAs, for a total of 49 putative targets. *Pum1*$^{-/-}$

male and female cerebella showed gene expression changes for 44 out of these 49 targets, which

correlated with the sex-dependent differences in Ago2 levels (**Figure 4E**, **Figure 4–figure supplement**

2B and **Supplementary Table 4**).

The top targets of Pum1, Pum2, Fmrp, Ago2, and Rbfox3 are shared

So far our *in vivo* proteomics results seem to be borne out by changes in the levels of protein interactors upon loss of Pum1. Ideally, we would purify recombinant proteins to perform gel filtration assays to get a better handle on the nature of the interactions, but Pum1 is difficult to purify because it is prone to aggregation (Miller & Olivas, 2011). If the complexes Pum1 forms with these RBPs are physiologically relevant, however, then they should co-regulate at least some of the same mRNA targets. Indeed, one corollary of the "regulon theory," which posits that mRNA targets in the same pathway are co-regulated (Blackinton & Keene, 2014; Keene, 2007a, b; Keene & Lager, 2005), is that there should be a discernible set of RBPs that do the co-regulating.

To test this hypothesis, we downloaded and reanalyzed all the high-throughput sequencing UV-crosslinking and immunoprecipitation (HITS-CLIP) data that is available from the murine brain. Fortunately, this data exist for Fmrp (Maurin *et al*, 2018), Ago2 (Chi *et al*, 2009), Rbfox3 (Weyn-Vanhentenryck *et al*, 2014), Pum1, and Pum2 (Zhang *et al.*, 2017). Then, to determine whether and how much the targets overlap, we performed gene set enrichment analysis (GSEA) (Subramanian *et al*, 2005). GSEA is a method for determining whether members of a particular class of genes from set *S,* such as the targets of a particular RBP like Pum1, are randomly distributed throughout another gene set list *L*, which is ranked (in this case) according to reads per kilobase of transcript, per million mapped reads (RPKM). Genes from *S* that are enriched (over-represented) at the top of *L* are expressed more than would be expected; the converse is true for genes over-represented at the bottom of *L*. Here, we performed pairwise GSEA analysis using Fmrp as the basis for comparison (because it has the largest dataset) and Pum1, Pum2, Ago2 and Rbfox3. As negative controls, we used HITS-CLIP data from mouse brain for four RBPs that did not show up as Pum1 interactors in our mass spec: Mbnl2 (Charizanis *et al*, 2012), Nova (Zhang *et al*, 2010), Apc (Preitner *et al*, 2014), and Ptpb2 (Licatalosi *et al*, 2012).

Strikingly, Pum1 HITS-CLIP targets were preferentially distributed in the top 5th percentile of all Fmrp targets, with an enrichment score (ES) of 0.93 (the maximum is 1) and a FDR of 0 (**Figure 5A,** *blue line represents ES*). Pum2, Ago2, and Rbfox3 showed nearly identical patterns (**Figure 5A**). There was no significant overlap between the targets of Fmrp and those of any negative control (Nova had the highest ES, but this was only 0.36 with a rank max of 45[th] percentile and FDR=1; **Figure 5B**).

To ascertain whether the highest-ranking Fmrp targets correspond to the genes with the highest probability of being Pum1 targets, we performed a more detailed GSEA. We first divided the Fmrp ranked target list into 10 equal bins according to percentile. We then repeated GSEA of Pum1 HITS-CLIP data for each bin and found that 648 of the 1194 identified Pum1 targets (54%) are in the top 10[th]

percentile of Fmrp targets, with an ES of 0.8; moreover, this was the only statistically significant bin

(**Figure 5C**). This was also true for Pum2, Ago2, and Rbfox3 (**Figure 5C**). Neither Pum1, Pum2, Rbfox3,

Fmrp, nor Ago2 targets were enriched among any of the target lists of the negative controls ranked by

RPKM (**Figure 5–figure supplement 1A,** and data not shown).

We then performed the corresponding analysis using the Pum1 target list as the basis for comparison

instead of the Fmrp list. We ran GSEA on each of the four Pum1 partners against the list of Pum1 target

genes (again ranked by RPKM), and each partner's targets are within the top 20% of the Pum1 list

(**Figure 5D**). Specifically, Fmrp's targets reside in the top 10^{th} percentile (with an ES of 0.81), Pum2's

targets within the 16^{th} percentile (ES=0.9), Ago2's targets within the 18^{th} percentile (ES=0.76), and

Rbofx3's targets within the 19^{th} percentile (ES=0.67). The four RBPs used here as negative controls have

a minimum rank at the 37^{th} percentile, and the best ES was 0.26 for Apc; none of the five reached

statistical significance (**Figure 5D**). These analyses demonstrate that there is substantial overlap among

the highest-ranked targets of Pum1, Pum2, Fmrp, Ago2, and Rbfox3.

We then decided to look at the targets shared by Pum1, Pum2, Ago2, and Rbfox3 and see how they

distribute within Fmrp. The result was striking, with an ES of 0.93 falling within the top 5^{th} percentile

(**Figure 5E**); 141 out of 175 common targets were within the top 10^{th} percentile (bin 1) of Fmrp targets,

with 99 within the top 5^{th} (**Figure 5F** and **Supplementary Table 5**). This contrasts with the negative

controls, for which the best ES was 0.41 within the top 40^{th}-60^{th} percentile (**Figure 5G**). DAVID gene

ontology analysis of those 175 common targets between Ago2, Pum1, Pum2, Fmrp, and Rbfox3 revealed

pathways enriched in neurons and axonal projections (**Figure 5–figure supplement 1B** and **C**). Previous

studies have shown that Pum1 and Pum2 cooperate with the microRNA machinery to suppress certain

targets (Gennarino *et al.*, 2015; Kedde *et al.*, 2010). Among Fmrp HITS-CLIP targets, there were almost

300 microRNAs. Pum1 HITS-CLIP has 60 miRNAs, only four of which are not shared with Fmrp; Pum2

HITS-CLIP has no miRNAs that are not shared with either Pum1 or Fmrp (**Figure 5–figure supplement**

1D and **Supplementary Table 6**).

Because neuronal genes tend to be long (Zylka *et al*, 2015) and RPKM can introduce bias toward

longer genes (Mandelboum *et al*, 2019), we wanted to ensure that the targets were actually more highly

expressed. Since the published HITS-CLIPs data from other studies were acquired from mice at different

ages (but most at 3 weeks of age) and these studies did not typically perform RNAseq in parallel with

HITS-CLIP from the same tissue and age, we decided to perform RNAseq from WT mice at 3 weeks of

age in triplicate to determine a baseline level of gene expression in mouse brain expressed as DESeq2-

normalized counts (Love *et al*, 2014) (see Materials and Methods and **Supplementary Table 7**). We then

normalized RPKM values and repeated the GSEA enrichment analysis. The enrichment between the

targets of Pum1 and those of Pum2, Ago2, Fmrp, and Rbfox3 was retained even after gene expression normalization (**Figure 5–figure supplement 1E-G**), with a minimum ES of 0.78, while the maximum score of the negative controls was 0.38 (**Figure 5–figure supplement 1E, F**). The shared targets of Pum1, Pum2, Rbfox3, and Ago2 were even more enriched, with an ES of 0.84 (**Figure 5–figure supplement 1H**).

PUM1 interactors are destabilized in cell lines from PADDAS, but not PRCA, patients

Having identified PUM1 interactors and shared targets, we were now in a position to test our two-pronged hypothesis about how *PUM1* mutations can lead to different diseases. To determine whether either mutation destabilizes PUM1 interactors, we compared their levels in patient-derived cell lines with those in controls. For PADDAS, we compared fibroblasts from one 9-year-old female patient (R1147W; subject 1, or S1) with fibroblasts from three different 9-year-old female healthy controls. For PRCA, we compared lymphoblastoid cells from two female PRCA patients (both with the T1035S mutation; subjects 2 and 3, or S2 and S3, 59 and 58 years old, respectively) with lymphoblastoid cells from three different 58-year-old female healthy controls (Gennarino *et al.*, 2018). We first measured how much PUM1 we could pull down from the patient-derived cell lines compared to their respective controls. Post-IP did not detect any residual PUM1 from PADDAS or PRCA cell lines, or their controls, confirming that our protocol was able to pull down virtually all of the PUM1 protein from patient-derived cell lines (**Figure 6–figure supplement 1A, B**). IP against PUM1 followed by western blot showed that PADDAS cells had 49%, and PRCA cells 76%, of the amount of PUM1 found in healthy controls (**Figure 6A** and **B**), consistent with our previous report (Gennarino *et al.*, 2018).

Co-IP confirmed that PUM1 associates with PUM2, FMRP, AGO2, CNOT1, and MOV10 in patient cell lines (**Figure 6A** and **B**). The mild T1035S variant reduced PUM1 binding to AGO2 by ~21% (**Figure 6B**). The more severe R1147W variant, however, reduced PUM1 association with AGO2, CNOT1, and MOV10 by ~84%, ~59%, and ~90%, respectively (**Figure 6A**). Interaction with FMRP did not appear to be affected. (We could not examine the effect of PUM1 mutations on RBFOX3, which is not expressed in fibroblasts or lymphoblastoid cells.)

Western blot analysis revealed that the proteins that lose their association with the R1147W variant undergo reduction in their expression levels (**Figure 6C**). Note that MOV10's association with R1147W was greatly reduced (**Figure 6A**) even though its protein levels were unchanged (**Figure 6C**). AGO2 and CNOT1 levels were unchanged in the PRCA cell line but were ~50% lower in the PADDAS cell line (**Figure 6C**). The mRNA levels of *PUM1*, *AGO2*, *CNOT1*, and *MOV10* did not change (**Figure 6–figure supplement 1C**), confirming that the reductions in their respective protein levels were due to the loss of

interaction with PUM1-R1147W.

One limitation of these studies is that we are implicitly comparing different cell types. Since we

haven't been able to procure lymphoblasts from PADDAS patients or fibroblasts from PRCA patients, we

sought to confirm our findings *in vitro*. We inserted a GST-tag at the N-terminal of *AGO2*, *CNOT1*, and

FMRP (see Materials and Methods). (AGO2 and CNOT1 were the only proteins whose levels changed in

fibroblasts, so FMRP is used here as a control.) We also cloned a Myc-tag polypeptide at the N-terminal

of *PUM1*-WT, *PUM1*-R1147W (PADDAS) and *PUM1*-T1035S (PRCA) (see Materials and Methods).

We then co-transfected the same amount of Myc-tagged PUM1 proteins with the GST-tagged interactors,

followed by western blot to quantify the amount of GST-tagged protein retrieved relative to the quantity

of PUM1 proteins pulled down in the IP. GST-AGO2 associated with Myc-PUM1-R1147W 76% less

than it did with Myc-PUM1-WT (**Figure 7A**), in alignment with our observations in the PADDAS cell

lines (**Figure 6A, B**). We also found ~35% less interaction with CNOT1 (**Figure 7B**) but no decrease in

PUM1 association with FMRP (**Figure 7C**), again in accord with our findings in patient-derived cells

(**Figure 6A**, **B**).

We also examined the effects of the PUM1 mutant on PABPC1, PABPC4, EIF4A3, and EIF4G3

(**Figure 1A**), which various studies have indicated interact with each other (Cheng & Gallie, 2007; Gallie,

1998; Le *et al*, 1997) and seems to be important for Pumilio activities but not necessarily as its interactors

(Goldstrohm *et al.*, 2018; Menon *et al*, 2004). We cloned a GST-tag at the respective N-terminals and

performed the experiment as described above. While the R1147W mutation reduced interaction with

PABPC1 by roughly 40%, it actually increased interaction with PABPC4 by 350% (**Figure 7–figure**

supplement 1A and **B**). Strikingly, PUM1-R1147W increased interaction with EIF4A3 and EIF4G3 by

1600% to 2760% (**Figure 7–figure supplement 1C** and **D**).

These data suggest that the effects of the R1147W mutation, which does not impair PUM1 binding to

mRNA (Gennarino *et al.*, 2018), are due to the loss of interactions with RBPs that repress mRNA with a

concomitant gain of function with translational initiation proteins.

R1147W mutation dramatically destabilizes PUM1

We next tested the PUM1 variants in a protein stability assay. We measured the half-life of the three

Myc-tagged PUM1 proteins by separately transfecting 125ng of each. After cycloheximide treatment (T0)

to block translation, we quantified the amount of protein at different time points: 30min, 1h, 2h, 4h, and

5h, compared to T0 (**Figure 7–figure supplement 1E**). The half-life of PUM1-WT was roughly 5.5

hours, the half-life of

PUM1-T1035S was ~2 hours, and that of PUM1-R1147W slightly less than 1 hour (**Figure 7–figure**

supplement 1E).

We also wondered whether the R1147W mutation might be impaired in binding with WT PUM1. It

has been thought that Puf proteins do not dimerize (Edwards *et al*, 2001; Zamore *et al*, 1999), but recent

work supports this possibility (Qiu *et al*, 2019). We therefore co-transfected each of the Myc-tagged

PUM1 variants with the same amount of GST-tagged *PUM1*-WT and used an empty vector as a negative

control. First, we confirmed that IP against Myc-PUM1-WT pulled down GST-PUM1-WT. IP against

Myc-PUM1-R1147W pulled down 51% of the total GST-PUM1-WT, while the interaction between Myc-

PUM1-T1035S and GST-PUM1-WT remained unchanged (**Figure 7D**). The same interaction was

observed after RNase treatment, suggesting that mammalian PUM1 dimerizes without RNA (**Figure 7–**

figure supplement 1F). These data suggest that the R1147W mutation might exert a dominant-negative

effect on wild-type PUM1.

The combination of lower protein levels and marked protein instability helps explain why the

R1147W phenotype is better represented by the *Pum1* null mice than by the heterozygotes (Gennarino *et*

al., 2018).

Shared targets are upregulated in PADDAS but not PRCA

We had hypothesized that PRCA would involve dysregulation of PUM1 targets, whereas PADDAS

would involve both destabilization of PUM1 interactors and dysregulation of downstream targets of those

interactors. We therefore tested the effects of the T1035S and R1147W mutations on validated PUM1-

specific targets (Chen *et al*, 2012; Gennarino *et al.*, 2015; Zhang *et al.*, 2017) that are not in the HITS-

CLIP data for the other RBPs but are expressed in both fibroblasts and lymphoblasts. PUM1-specific

mRNA were dysregulated to very similar extents in PRCA and PADDAS patient cells, with only a few

targets being up to 20% more upregulated in PADDAS (**Figure 7E**).

We then examined the mRNA levels of all 175 targets shared between PUM1, PUM2, AGO2, FMRP,

and RBFOX3 (**Figure 5E** and **Supplementary Table 5**) in the same PRCA and PADDAS patient cell

lines. Of these shared targets, 55 were expressed in both PADDAS fibroblasts and PRCA lymphoblastoid

cells. Of these 55, 52 shared targets were upregulated in PADDAS but not in PRCA (**Figure 7F**), by an

average of two-fold (ranging from a low of 121% for IDS to 347% for TLK1). There was little or no

change in most of these targets in PRCA cells, though levels of CALM1, ATP2A2, CREB1, and GNAQ

fell by ~40%, and CALM2, TAOK1, and UBE2A by ~20% (**Figure 7F**).

These results support our hypothesis that the symptoms observed in PRCA are attributable to the

dysregulation of PUM1-specific target genes, while PADDAS involves both protein partner

destabilization and dysregulation of the partner proteins' targets.

Discussion

Since our initial study describing PUM1-related diseases (Gennarino *et al.*, 2018), we and others have identified additional PADDAS and PRCA patients (Bonnemason-Carrere *et al*, 2019; Gennarino *et al.*, 2018; Imaizumi *et al*, 2019; Lai *et al*, 2019). In our sample, R1147W accounts for the majority of PADDAS patients, and T1035S for the majority of PRCA, which supports the value of studying these particular two mutations as causing the most severe and the most mild forms of PUM1-related disease. The question that drove the present study is: why should the 25% drop in PUM1 levels from PRCA to PADDAS produce such different phenotypes, especially when R1147W is not impaired in binding to mRNA? Our data strongly support our hypothesis that the difference is not due to a linear increase in the de-repression of mRNA targets but is rather a function of an additional mechanism coming into play: the destabilization of numerous interactions and the de-repression of their downstream targets. We draw this conclusion from several lines of evidence. First, when Pum1 was reduced by 50% (in heterozygous mice) or 100% (in knockout mice), there were changes in the levels of associated proteins that depended on brain region and sex. Second, we found changes in the levels of these proteins in patient cell lines, despite unchanged mRNA levels. Third, our *in vitro* studies using Myc- and GST-tagged proteins showed that AGO2 and CNOT1 lose their interaction with PUM1-R1147W, which itself binds less to WT PUM1. Fourth, not all interactions were diminished: the responses to R1147W ranged from a loss of 40-70% to a gain of 2760%. Together these data persuade us that a 50% loss of PUM1, whether through heterozygosity or the R1147W mutation, disrupts interactions with native partners and differentiates PADDAS from PRCA.

Diseases involving dosage-sensitive proteins

There are other dosage-sensitive proteins that produce different phenotypes depending on their expression level (Tan & Zoghbi, 2019), and our results raise the possibility that interacting complexes may be disrupted once expression falls below a certain threshold. What that threshold might be, exactly, is not clear, but for PUM1 it seems to be somewhere between the 75% of wild-type levels of PUM1 seen in PRCA and the 60% level estimated for the R1139W mutation that produced a milder form of PADDAS (Gennarino et al., 2018). In this context it is worth noting that a recent study found that, below a threshold of ~70% of normal levels of FMRP, there were steep decreases in IQ for each further decrement in FMRP levels, even as small as 5-10% (Kim *et al*, 2019). The amount of loss that can be sustained for a given protein would likely depend on its usual quantities, and it is possible that for some proteins, the difference in phenotype between greater and lesser abundance may indeed reflect a linear progression from mild to

severe. For example, a mutation in *MECP2* that reduces the protein levels by only 16% causes a mild

form of Rett syndrome: the child developed normally until the age of two years, then regressed, lost

acquired language and manual skills, and developed hand stereotypies (Takeguchi *et al*, 2020). She

eventually regained the ability to repeat words in an echolalic fashion and never developed seizures, but

the phenotype was clearly Rett syndrome. The relationship between protein levels and disease features in

MeCP2-related disorders may be more linear because this protein is so extraordinarily abundant

(Sandweiss *et al*, 2020). Similarly, in proteopathies such as Alzheimer's or Parkinson's disease, genetic

duplications of *APP* or *SNCA* cause earlier onset of what is recognizably the same disease (Chartier-

Harlin *et al*, 2004; Rovelet-Lecrux *et al*, 2006).

In contrast, there are diseases in which the phenotypes do not simply range from mild to severe

versions of the same symptoms, but seem to take on a different character. The polyglutamine diseases,

such as Huntington's disease and the spinocerebellar ataxias, come to mind, perhaps because it was

through studying spinocerebellar ataxia type 1 (SCA1), that PUM1's role in neurological function was

discovered (Gennarino *et al.*, 2015). In the polyglutaminopathies, the disease-causing protein bears an

abnormally expanded CAG tract that tends to expand upon intergenerational transmission. Although the

range of normal and pathogenic repeat tract lengths differs from one polyglutamine disease to another,

larger expansions are more unstable, cause earlier onset, and affect far more tissues than smaller

expansions (Orr & Zoghbi, 2007). For example, adult-onset SCA7 presents as ataxia, but infantile SCA7

affects the entire nervous system, the heart, and the kidneys, and leads to death by two years of age (Bah

et al, 2020). Another example is Huntington's disease, where the juvenile form frequently lacks the

classic chorea yet produces seizures, which are not a feature of the adult-onset disease; brain

morphometry is also quite different in adult- and juvenile-onset cases (Tereshchenko *et al*, 2019). In this

family of diseases, therefore, the mechanism is technically the same—repeat expansion—but different

tissues have different thresholds for the CAG repeats. In the case of PUM1-related disease, it seems that

an additional mechanism comes into play for the more severe phenotypes, beyond upregulation of mRNA

targets. Interestingly, FMRP, which harbors a dynamic CGG repeat, is also associated with very different

diseases, through two different mechanisms. Very large expansions silence the gene and produce Fragile

X syndrome, whereas premutations are thought to cause the adult-onset Fragile-X-associated

tremor/ataxia syndrome through RNA toxicity (Salcedo-Arellano *et al*, 2020).

Implications of the interactome

It has never been clear how the various modes of action attributed to PUM1 or other RBPs relate to

one another, in large part because studies typically focus on one protein or pathway at a time. Our data

suggest that the three mechanisms of repression that have been proposed for PUM1 may be coordinated in

neurons, insofar as PUM1, PUM2, FMRP, AGO2, MOV10, CNOT1 and RBFOX3 (and related proteins

in specific brain regions) either interact or are so close to each other within the ribonucleosome that the

loss of Pum1 or RNA can change the composition of the complexes that are identified by co-IP, in ways

that are specific to brain region and sex. In this context it is worth noting that a very recent study found

alternative splicing is altered in hippocampal slices from Fmrp-deficient mice; this observation was

attributed to changes in H3K36me3 levels (Shah *et al*, 2020). Our data suggest that FMRP has an even

closer relationship with the RBFOX protein family and alternative splicing machinery than suspected.

Indeed, recent work has provided tantalizing glimpses of close interactions among various kinds of RNA

metabolism. For example, members of the RBFOX family of proteins may, depending on their interactors

(and perhaps cell type, sex, age, and species), be involved in microRNA processing in the nucleus and

translation in the cytoplasm (Conboy, 2017). The FMRP/MOV10 complex appears to be involved in

regulating translation through miRNA, with evidence that this role may change according to cell type

(Kenny *et al.*, 2020). Another study used quantitative mass spec to examine how Fmrp expression levels

change with age in the wild-type rat dentate gyrus, and found differences in the levels of myriad proteins;

among the 153 proteins with the most significant changes in levels were Pum1, Pum2 and Papbc1

(Smidak *et al*, 2017). The region-, sex-, and age-specificity of certain interactions indicates that

unraveling RBP interactomes *in vivo* will require considerable finesse.

514   Although we did not pursue the entire interactome here, future studies investigating interactions

between PUM1 and members of the mTOR pathway and anaphase-promoting complex should be very

fruitful. As we were preparing this work, a new study demonstrated that mTORC1 is involved in

postsynaptic responses in hippocampal glutamatergic neurons, whereas mTORC2 modulates presynaptic

release (McCabe *et al*, 2020). Given the importance of proteostasis at the synapse—and its failure in

many neurological diseases (Zoghbi & Bear, 2012)—the notion that different protein-regulating activities

might be coordinated is intuitively appealing.

521   There is still much to be understood about how PUM1 works, and where and how the various

RNA metabolic machineries come into contact. The fact that several associations we tested are RNA-

independent suggests that certain complexes may form prior to binding mRNA. Could different pools of

PUM1 form different complexes that determine whether and where a target is bound? Do the subtle

differences between brain regions in the choice of interacting partners from the same pathways influence

target choice, activity, or both? Given that mRNA is much larger than proteins, is the quantity of these

RBPs so great that they coat the RNA? What other proteins may be involved in 'handing off' an mRNA

from one pathway to another? Although we began this study to better understand Pum1 functions in

neurons, and thus have been rather Pum1-centric, it is by no means certain that Pum1 is the only protein

that links these different molecular machineries. It may be simply the first of several missing links that we

didn't realize were missing. Creating RBP interactomes should lead to a more complex yet realistic

picture of RBP roles in neuronal function and in neurological disease.

Acknowledgments

We thank the patients, their families, and their clinical teams for participating in our study, and the

anonymous reviewers for helpful suggestions. We thank Mingyu Cao for help with statistical analyses

and U.N. Wasko for starting the cloning procedure with Myc and GST tag sequences *in vitro*. We also

thank G. Struhl, V. Ambros, T. Duchaine, G. Karsenty, H. Zoghbi, M. Jovanovic, Y. Giardina, and

members of the Gennarino laboratory for helpful discussions and support. This work was supported by

the National Institute of Neurological Disorders and Stroke (NINDS; R01NS109858 to V.A.G.); the

National Ataxia Foundation/Young Investigator Research Grant (V.A.G.); the Brain & Behavior Research

Foundation Young Investigator Award; and the Paul A. Marks Scholar Program, Columbia University

Vagelos College of Physicians and Surgeons. This project was also supported by the Genomic and RNA

Profiling Core at Baylor College of Medicine and the Bioinformatics Core at the Jan and Dan Duncan

Neurological Research Institute. The authors declare no competing interests.

Author Contributions

S.B. designed and performed molecular experiments, analyzed and interpreted the data, and drafted the

manuscript. N.D.P. performed the qPCR experiments, *in vitro* immunoprecipitation assays, analyzed the

data, and edited the manuscript. A.C. performed all the brain region dissections in mice, performed the

qPCR experiments and maintained the *Pum1* mutant mouse colony. M.C. performed the cloning

experiments. V.L.B. interpreted data and wrote the manuscript. P.P and R.K.S. performed mass

spectrometry and helped analyze the data. V.A.G. conceived the study, analyzed and interpreted the data,

performed molecular experiments, and wrote the manuscript.

Declaration of Interests

The authors declare not competing interests

Materials and Methods

Ethical statement and mouse strains. All animal procedures were approved by the Institutional Animal

Care and Use Committee at Columbia University, New York under the protocol AC-AAAU8490. Mice

were maintained on a 12-hr light, 12-hr dark cycle with regular chow and water ad libitum. Pum1 knock-

out mice were generated as previously described (Chen *et al.*, 2012). C57BL/6J wild-type mice were

purchased from Jackson Laboratory and maintained as described above. For brain dissection, mice were

anesthetized with isoflurane, and the brain rapidly removed from the skull and lysed in the appropriate

buffer according to the experiment (see Materials and Methods Details).

HEK293T cell culture and maintenance. Human embryonic kidney immortalized 293T (HEK293T)

cells were grown in DMEM (GenDepot, #CM002-320) supplemented with 10% of heat-inactivated fetal

bovine serum (FBS [GenDepot, #F0901-050) and 1% penicillin/streptomycin (GenDepot, #CA005-010).

All cells were incubated at 37 °C in a humidified chamber supplemented with 5% CO_2. HEK293T cells

were later processed according to the needs of specific experiments (described below).

Patient-derived cell lines: fibroblasts and lymphoblasts. PADDAS and PRCA patient cell lines are the

same as those reported previously (Gennarino *et al.*, 2018). The consent form for each subject specifically

allows for sharing of medical information and physical exam findings; the sharing of cell lines from the

PADDAS and PRCA subjects and the controls was approved under the Baylor College of Medicine IRB

H-34578 and the Columbia University Medical Center IRB-AAAS7401 (Y01M00).

Primary fibroblasts from the *PUM1* PADDAS patient and the age- and sex-matched controls were

generated as previously described (Gennarino *et al.*, 2018). Briefly, cells were isolated from skin biopsies

taken from the patient or age-matched controls using standard methodology (Barch and Association of

Cytogenetic Technology, 1991) and placed in a transport medium (Ham's F10, Thermo Scientific,

#11550043). The skin specimen was later removed from the transport medium using a sterile technique

(in a Class II biohazard cabinet) and transferred to a sterile Petri dish where it was cut into small pieces (<

0.5 mm) using sterile scalpel blades. These pieces were transferred to the lower surface of a 25 cm^2

culture flask (6-8 pieces per flask) which had been pre-moistened with 1-2 mL of AmnioMAX Complete

Medium (Thermo Scientific, #11269016) supplemented with 1% penicillin/streptomycin (GenDepot,

#CA005-010). Cell cultures were maintained at 37 °C in a humidified incubator supplemented with 5%

CO_2. When cell growth was observed around the edges of the tissue, usually 3 to 5 days later, 2 to 3 mL

of AmnioMAX Complete Medium were added. Once growth was established and the tissue was anchored

to the flask, another 8 mL of AmnioMAX Complete Medium was added. Thereafter, the medium was

renewed every 3 to 4 days until ready for sub-culturing.

Lymphoblastoid cells from *PUM1* PRCA patients and the age- and sex-matched controls were

generated as previously described (Gennarino *et al.*, 2018). Briefly, lymphoblastoid suspension cell

cultures were grown in RPMI 1640 medium (Invitrogen, #11875093) supplemented with 10% heat-

inactivated fetal bovine serum (Atlanta Biological, Flowery Branch, #S11195H) and 1%

penicillin/streptomycin (GenDepot, #CA005-010). Cell cultures were maintained at 37°C in a humidified

incubator supplemented with 5% CO_2. Medium was renewed every 2 to 3 days.

Immunoprecipitation experiments using mouse brain tissue. Mouse brain tissues were gathered from

an equal number of 10-week-old male and female mice as follows: whole brains from two mice;

hippocampi from ten mice; cerebral cortices from eight mice; and cerebella from eight mice. Each

grouping is considered one sample for one experiment. Samples were processed with a dounce

homogenizer using a lysis buffer consisting of 200mM $NaCl_2$, 100mM $NaPO_4$, 20mM Hepes pH 7.4, 1%

Triton X (which should disrupt all but the strongest protein-protein interactions) and complemented by

1X of Xpert Protease and 1X of Phosphatase Inhibitor Cocktail Solutions (GenDepot, #P3100-100,

#P3200-020). Following homogenization, the samples were placed on ice for 15 minutes then centrifuged

at 14,800 rpm at 4°C for 25 minutes to remove the debris from the supernatant. The supernatant was then

moved to 1.5 ml tubes (Beckman microfuge tube #357448) and spun down in a Beckman ultra-centrifuge

(Optima Max XP) at 4°C for 25 minutes at 44,000 rpm. 10% of the protein lysate was stored as input and

only 1% was loaded for western blot. The protein extract was later divided into two aliquots, one for IP

against the protein of interest (antibodies listed below) and the other for IP against IgG, and was then

incubated with 30 μL of Dynabeads™ Protein G (Invitrogen, #10004D) and 5 μg of antibody overnight at

4°C on a rotisserie tube rotator. The next day, the beads were washed four times with the same lysis

buffer used for IP and resuspended in 40μL of elution buffer (consisting of lysis buffer, NuPAGE

10X Reducing Agent [Invitrogen, #NP0009], NuPAGE LDS sample buffer at 1X final concentration

[Invitrogen, #NP0007]) and boiled at 95°C for 10 minutes before the samples were loaded in the

NuPAGE 4%–12% Bis-Tris Gels (Invitrogen, #NP0335BOX & #NP0336BOX) for further resolution and

western blot analysis.

For the IP with RNase treatment, the beads were resuspended in 400 μL of lysis buffer after the three

final washes and divided into two separate 1.5 ml tubes of 200 μL each. To establish the dose required to

remove all RNA, we tested different amounts of RNase I (Invitrogen, #EN0602) and found that 4 μL was

enough to render RNA undetectable both by denaturing gel and cDNA amplification. The negative

control was to go without RNase treatment; both were incubated at 37°C for 15 min on a rotisserie tube

rotator. After incubation, all the samples were washed one last time with 500 μL of lysis buffer and then

eluted in 20 μL of elution buffer. We used the same protocol for all the IP processed by mass

spectrometry.

The antibodies used for IP were: goat α-PUM1 (Bethyl Laboratories, #A300-201A), rabbit α-PUM2

(Bethyl Laboratories, #A300-202A), rabbit α-FMRP (Abcam Cambridge, #ab17722), rabbit α-AGO2

(Abcam Cambridge, #ab32381), rabbit α-NeuN (Thermo Fisher Scientific, #PA5-37407), rabbit α-

CNOT1 (Cell Signaling Technology, #44613), rabbit α-MOV10 (Bethyl Laboratories, #A301-571A),

rabbit α-RAPTOR (Cell Signaling Technology, #2280), rabbit α-CAD (Bethyl Laboratories, #A301-

374A), rabbit α-ANAPC1 (Bethyl Laboratories, #A301-653A), and rabbit α-FZR1 (Thermo Fisher

Scientific, #PA5-85303).

N.B.: The reason the inputs are separated from the IPs and then collaged is because *in vivo* we load

only 1% of the lysate, which we later run off the same blot of the IPs and expose the membrane to see

either the input (which usually appears first) or the IPs (which for some of them appear with a longer

exposure). This is common practice when working with *in vivo* tissues (De Maio *et al*, 2018) (Lee *et al*,

2008) (Rousseaux *et al*, 2018) (Di Grazia *et al*, 2021) (Jin *et al*, 2020) (Lee *et al*, 2020).

Immunoprecipitation experiments from HEK293T and patient-derived cell lines. HEK293T cells

and patient-derived fibroblasts or lymphoblastoid cells were lysed by pipetting up and down with a

1000μl tip in a lysis buffer consisting of 200mM NaCl₂, 100mM NaPO₄, 20mM Hepes pH 7.4, 1% Triton

X and complemented by 1X of Xpert Protease and 1X of Phosphatase Inhibitor Cocktail (GenDepot,

#P3100-100, #P3200-020). The rest of the protocol is the same as described above for mouse brain tissue,

except that we used 2.5 μg of primary antibody for IP.

Co-Immunoprecipitation in-gel digestion for mass spectrometry. Immunoprecipitated samples were

separated on NuPAGE 4-12% gradient SDS-PAGE (Invitrogen, #NP0335BOX & #NP0336BOX) and

stained with SimplyBlue (Invitrogen, #LC6060). Protein gel slices were excised and *in-gel* digestion

performed as previously described (Shevchenko *et al*, 2006), with minor modifications. Gel slices were

washed with 1:1 Acetonitrile and 100mM ammonium bicarbonate for 30 min then dehydrated with 100%

acetonitrile for 10 min until shrunk. The excess acetonitrile was then removed and the slices dried in a

speed-vacuum at room temperature for 10 minutes. Gel slices were reduced with 5 mM DTT for 30 min

at 56°C in an air thermostat, cooled down to room temperature, and alkylated with 11 mM IAA for 30

min with no light. Gel slices were then washed with 100 mM of ammonium bicarbonate and 100%

acetonitrile for 10 min each. Excess acetonitrile was removed and dried in a speed-vacuum for 10 min at

room temperature and the gel slices were re-hydrated in a solution of 25 ng/μl trypsin in 50 mM

ammonium bicarbonate for 30 min on ice and digested overnight at 37 °C in an air thermostat. Digested

peptides were collected and further extracted from gel slices in extraction buffer (1:2 ratio by volume of

5% formic acid: acetonitrile) at high speed, shaking in an air thermostat. The supernatants from both

extractions were combined and dried in a speed-vacuum. Peptides were dissolved in 3% acetonitrile/0.1%

formic acid.

Liquid chromatography with tandem mass spectrometry (LC-MS/MS). The Thermo Scientific

Orbitrap Fusion Tribrid mass spectrometer was used for peptide tandem mass spectroscopy (MS/MS).

Desalted peptides were injected in an EASY-Spray^{TM} PepMap^{TM} RSLC C18 50cm X 75cm ID column

(Thermo Scientific) connected to the Orbitrap Fusion^{TM} Tribrid^{TM}. Peptide elution and separation were

achieved at a non-linear flow rate of 250 nl/min using a gradient of 5%-30% of buffer B (0.1% (v/v)

formic acid, 100% acetonitrile) for 110 minutes, maintaining the temperature of the column at 50 °C

during the entire experiment. Survey scans of peptide precursors are performed from 400 to 1500 *m/z* at

120K full width at half maximum (FWHM) resolution (at 200 *m/z*) with a 2 x 10^5 ion count target and a

maximum injection time of 50 ms. The instrument was set to run in top speed mode with 3-second cycles

for the survey and the MS/MS scans. After a survey scan, MS/MS was performed on the most abundant

precursors, i.e., those exhibiting a charge state from 2 to 6 of greater than 5 x 10^3 intensity, by isolating

them in the quadrupole at 1.6 Th. We used collision-induced dissociation (CID) with 35% collision

energy and detected the resulting fragments with the rapid scan rate in the ion trap. The automatic gain

control (AGC) target for MS/MS was set to 1 x 10^4 and the maximum injection time was limited to 35ms.

The dynamic exclusion was set to 45s with a 10ppm mass tolerance around the precursor and its isotopes.

Monoisotopic precursor selection was enabled.

LC-MS/MS data analysis. Raw mass spectrometric data were analyzed using the MaxQuant

environment v.1.6.1.0 (Cox & Mann, 2008) and Andromeda for database searches (Cox *et al*, 2011) at

default settings with a few modifications. The default was used for first search tolerance and main search

tolerance (20 ppm and 6 ppm, respectively). MaxQuant was set up to search with the reference mouse

proteome database downloaded from Uniprot (https://www.uniprot.org/proteomes/UP000000589).

MaxQuant performed the search for trypsin digestion with up to 2 missed cleavages. Peptide, site and

protein false discovery rates (FDR) were all set to 1% with a minimum of 1 peptide needed for

identification; label-free quantitation (LFQ) was performed with a minimum ratio count of 1. The

following modifications were used for protein quantification: oxidation of methionine (M), acetylation of

the protein N-terminus, and deamination for asparagine or glutamine (NQ). Results obtained from

MaxQuant were further analyzed using the Perseus statistical package (Tyanova *et al*, 2016) that is part of

the MaxQuant distribution. Protein identifications were filtered for common contaminants. Proteins were

considered for quantification only if they were found in at least two replicate groups. Significant

alterations in protein abundance were determined by ANOVA with a threshold for significance of $P <$

0.05 (permutation-based FDR correction). Pum1 protein interactors were later considered if they were

found in at least five out of six mass spec experiments for whole brain and in at least two out of three

experiments for each respective brain region with a fold-change of >1.5 between LFQ-PUM1-WT and

LFQ-IgG-WT (see **Table S1**).

Protein-protein interaction map. The protein-protein interaction map for the whole brain (**Figure 1A)**

was generated by Cytoscape (https://cytoscape.org/) (Otasek *et al*, 2019) and the interactions among the

Pum1 interactors were inferred from Corum (Giurgiu *et al*, 2019) and the Human Protein Atlas (Thul *et*

al, 2017) by g:GOSt which is a package of g:Profiler (https://biit.cs.ut.ee/gprofiler/gost) (Raudvere *et al.*,

2019). The protein-protein interaction to generate the brain region-specific map (**Figure 1–figure**

supplement 2B) was generated by Cytoscape.

Protein quantification and western blot analysis. Patient-derived lymphoblastoid, fibroblast cell lines,

and control cell lines were collected at 6×10^6 cell confluence and processed for protein extraction. For

mouse tissues, we processed either half brain (the other half was processed for RNA extraction, see

below) or the entire hippocampus, cortex, or cerebellum for protein extraction. Mouse tissues or cell

pellets were subsequently lysed with modified RIPA buffer consisting of 25 mM Tris-HCL, pH 7.6,

150 mM NaCl, 1.0% Tween 20, 1.0% sodium deoxycholate, 0.1% SDS, completed with 1X Xpert

Protease and 1X Phosphatase Inhibitor Cocktail Solutions (GenDepot, #P3100-100 & #P3200-020). Cells

were lysed by pipetting them up and down with a p1000 tip and then placed on ice for 20 min followed by

centrifugation at 14,800 rpm at 4°C for 25 minutes. Mouse brain tissues were lysed starting from a 18G 1

½" (Becton Dickson, #305196) and move down to 21G 1 ½" (Becton Dickson, #305167) and finally to

26G 1 ½" (Becton Dickson, #305111) syringe needles by pipetting them up and down until the lysate

passed through the needle smoothly. Proteins were quantified by Pierce BCA Protein Assay Kit (Thermo

Scientific, # PI23225) and the absorbance was measured by NanoDrop OneC (Thermo Scientific).

Consequently, proteins were resolved by high resolution NuPAGE 4%–12% Bis-Tris Gel (Invitrogen,

#NP0335BOX & #NP0336BOX) according to the manufacturer's instructions. All the blots were

acquired on the G:BOX Chemi XX9 machine (Syngene; Frederick, MD) using GeneSys software 1.6.5.0.

Gel exposures were determine by the software.

Antibodies used for all the western blot experiments were: goat α-PUM1 [1:2500, (Bethyl

Laboratories, #A300-201A)], rabbit α-PUM1 [1:2000, (Abcam Cambridge, #ab92545)], rabbit α-PUM2

[1:2000, (Bethyl Laboratories, # A300-202A)], rabbit α-FMRP [1:1000, (Abcam Cambridge, #ab17722)],

rabbit α-AGO2 [1:1000, (Abcam Cambridge, #ab32381)], rabbit α-NeuN (Rbfox3) [1:1000, (Thermo

Scientific, #PA5-37407)], rabbit α-CNOT1 [1:1000, (Cell Signaling Technology, #44613)], rabbit α-

MOV10 [1:2000, (Bethyl Laboratories, #A301-571A)], and mouse α-GAPDH [1:10000, (Millipore,

#CB1001)].

RNA extraction and quantitative real-time PCR (qPCR). Human fibroblast, lymphoblastoid, and

respective control cell lines were harvested at 6×10^6 confluence prior to RNA extraction. For mouse

tissues, half of the whole brain (the other half was processed for protein extraction, see above) or the

entire hippocampus, cortex, or cerebellum were processed for RNA extraction. The RNA was collected

for both human cells, mouse brain and brain region tissues using the miRNeasy kit (QIAGEN, # 217004)

according to the manufacturer's instructions. RNA was quantified using NanoDrop OneC (Thermo Fisher

Scientific). cDNA was synthesized using Quantitect Reverse Transcription kit (QIAGEN, # 205313)

starting from 1 μg of RNA. Quantitative RT-polymerase chain reaction (qRT-PCR) experiments were

performed using the CFX96 Touch Real-Time PCR Detection System (Bio-Rad Laboratories, Hercules)

with PowerUP SYBR Green Master Mix (Applied Biosystems, #A25743). Real-time PCR runs were

analyzed using the comparative C_T method normalized against the housekeeping human gene *GAPDH* or

mouse *Gapdh*, depending on the experiment (Vandesompele *et al*, 2002).

Protein stability assay. HEK293T cells were seeded in 12-well plates for 24h and then transfected with

125 ng of either *PUM1* WT or mutant cDNA and one of the interactors using the jetPRIME Transfection

Reagent (Polyplus transfection, #55-132) according to the manufacturer's protocol. After 24 h cells were

treated with 0.1 mg/ml of cycloheximide (Fisher Scientific, #AAJ66004XF), which we considered time

zero. Cells were later collected after 30min, 1h, 2h, 4h and 5h of treatment and processed for protein

extraction and western blot analysis. Antibody: mouse α-Myc antibody [1:2000, (Cell Signaling

Technologies, #2276)].

MicroRNA library construction and sequencing. Library preparation and microRNA sequencing was

performed by LC Sciences according to the following criteria. Total RNA was extracted from cerebellum

of WT and *Pum1*$^{-/-}$ male and female at 10 weeks of age in triplicate, for a total of 12 samples using the

miRNeasy kit (QIAGEN, # 217004) according to the manufacturer's instructions. The total RNA quality

and quantity were assessed with Bioanalyzer 2100 (Agilent Technologies, Santa Clara) with RIN number

> 7.0. Approximately 1 μg of total RNA were used to prepare the small RNA library according to the

protocol of TruSeq Small RNA Sample Prep Kits (Illumina, San Diego). Then the single-end sequencing

50bp was performed on an Illumina Hiseq 2500 at LC Sciences (Hangzhou, China) following the

vendor's recommended protocol.

MicroRNA sequencing bioinformatic analysis. Raw reads were subjected to an in-house program,

ACGT101-miR (LC Sciences, Houston) to remove adapter dimers, junk, low complexity, common RNA

families (rRNA, tRNA, snRNA, snoRNA) and repeats. Subsequently, unique sequences of 18–26

nucleotides in length were mapped to specific species precursors in miRBase 22.0

(http://www.mirbase.org/) by BLAST search to identify known miRNAs and novel 3p- and 5p-derived

miRNAs. Length variation at both 3' and 5' ends and one mismatch inside of the sequence were allowed

in the alignment. The unique sequences mapping to specific species of mature miRNAs in hairpin arms

were identified as known miRNAs. The unique sequences mapping to the other arm of known specific

species precursor hairpins opposite the annotated mature miRNA-containing arm were considered to be

novel 5p- or 3p-derived miRNA candidates. The remaining sequences were mapped to other selected

species precursors (with the exclusion of specific species) in miRBase 22.0 by BLAST search, and the

mapped pre-miRNAs were further BLASTed against the specific species genomes to determine their

genomic locations. The last two were also defined as known miRNAs. The unmapped sequences were

BLASTed against the specific genomes, and the hairpin RNA structures containing sequences were

predicted from the flank 80 nt sequences using RNAfold software (http://rna.tbi.univie.ac.at/cgi-

bin/RNAWebSuite/RNAfold.cgi). The criteria for secondary structure prediction were: (1) number of

nucleotides in one bulge in stem (≤12), (2) number of base pairs in the stem region of the predicted

hairpin (≥16), (3) cutoff of free energy (kCal/mol ≤-15), (4) length of hairpin (up and down stems +

terminal loop ≥50), (5) length of hairpin loop (≤20), (6) number of nucleotides in one bulge in mature

region (≤8), (7) number of biased errors in one bulge in mature region (≤4), (8) number of biased bulges

in mature region (≤2), (9) number of errors in mature region (≤7), (10) number of base pairs in the mature

region of the predicted hairpin (≥12), (11) percent of mature region in stem (≥80).

RNA sequencing. The Genomic and RNA Profiling Core at Baylor College of Medicine first conducted

Sample Quality checks using the NanoDrop spectrophotometer and Agilent Bioanalyzer 2100. We then

used Illumina TruSeq Stranded Total RNA library preparation protocol (p/n 15031048, revE, document

#RS-122-9007) along with ThermoFisher's ERCC RNA Spike-In Control Mixes Protocol (publication

number 4455352, rev. D). Our process is as follows:

Illumina TruSeq stranded total RNA sample preparation protocol: Total RNA samples were

normalized to 250ng each, based on picogreen quantitation, and ERCC RNA Spike-In Controls (p/n

4456739) were incorporated into each sample. Using the TruSeq Stranded Total RNA Library Prep Gold

Kit (p/n RS-122-2301), samples then underwent ribosomal RNA reduction and subsequently fragmented

and primed with random hexamers to produce first strand cDNA. During second strand cDNA synthesis,

RNA templates were removed and replaced with cDNA strands containing dUTP. The ds-cDNA was then

purified using Beckman Coulter AMPure XP beads (p/n A63881).

Library preparation: Libraries were created from the cDNA by attaching an adenosine to the 3' end

and ligating unique adapters to the ends (for more information on this process, see below). The ligated

products underwent 15 cycles of amplification and a final bead purification. The resulting libraries were

quantitated using the NanoDrop spectrophotometer and fragment size assessed using the Agilent 2100

Bioanalyzer. A qPCR assay was performed on the libraries to determine the concentration of adapter

ligated fragments using the Applied Biosystems ViiA 7 Quantitative PCR instrument and a KAPA

Library Quant Kit (p/n KK4824). All samples were pooled equimolarly and re-quantitated by qPCR, and

also re-assessed on the Bioanalyzer.

Cluster generation by bridge amplification: HiSeq 2500 HOM. Using the concentration from the

ViiA7 qPCR machine above, 26pM of equimolarly pooled library was loaded onto three lanes of a high

output v4 flowcell (Illumina p/n PE-401-4001) and amplified by bridge amplification using the Illumina

cBot machine (cBot protocol: PE_HiSeq_Cluster_Kit_v4_cBot_recipe_v9.0). PhiX Control v3 adapter-

ligated library (Illumina p/n FC-110-3001) is spiked-in at 2% by weight to ensure balanced diversity and

to monitor clustering and sequencing performance. A paired-end 100 cycle run was used to sequence the

flowcell on a HiSeq 2500 Sequencing System (Illumina p/n FC-401-4003). An average of 129 million

read pairs per sample was sequenced.

End repair converts the overhangs caused by fragmentation into blunt or polished ends. The 3' to 5'

exonuclease activity of the End Repair mix removes the 3' overhangs, and polymerase activity fills in the

5' overhangs.

Adenylation: a single 'A' nucleotide is added to the 3' ends of the blunt fragments.

Adapter ligation: an adapter is ligated to the 3' end of the DNA fragment with a corresponding 'T'

nucleotide.

Absolute expression: The absolute gene expression was calculated as DESeq2-normalized counts as

previously described (Love *et al.*, 2014) (**Table S7**).

Samples: RNAseq was performed in triplicate from whole C57BL/6J mouse brains at 3 weeks of age.

Gene Set Enrichment Analysis (GSEA). GSEA was performed as previously described (Subramanian *et*

al., 2005). The cumulative distribution function was conducted by performing 1000 random gene-set

membership assignments. A nominal p-value < 0.01 and an FDR < 0.25 were used to assess the

significance of the enrichment score (ES). HITS-CLIP data were obtained from the literature and were

initially acquired as follows: Pum1 and Pum2 from neonatal murine brains (Zhang *et al.*, 2017), Fmrp

from cerebellum, cortex, and hippocampus together (Maurin *et al.*, 2018), Ago2 from neocortex at

embryonic day 13 (Chi *et al.*, 2009), Rbfox3 from mouse brain (age not specified) (Weyn-Vanhentenryck

et al., 2014), Nova from mouse brain (age not specified) (Zhang *et al.*, 2010), Ptpb2 from neocortex at

embryonic day 18.5 (Licatalosi *et al.*, 2012), Mbnl2 from hippocampus at 8-12 weeks of age (Charizanis

et al., 2012), and Apc from mouse brain at embryonic day 14 (Preitner *et al.*, 2014). For consistency all

the HITS-CLIP data were ranked according to their RPKM.

For the normalized RPKM-GSEA analysis we performed RNAseq experiments (as described above),

in triplicate, from WT C57BL/6 mouse brain at 3 weeks of age (we chose this age to be comparable to the

other HITS-CLIP datasets) to infer the basal level of gene expression by DESeq2-normalized counts (see

above). We then normalized the RPKM values by the DESeq2-normalized counts and repeated the GSEA

enrichment analysis.

Gene ontology analyses. All gene ontology analyses were performed with David Gene Ontology (GO)

and only categories with FDR<0.05 were considered. David GO for the Pum1 interactome considered the

entire interactome as background. For GSEA HITS-CLIP targets shared among Pum1, Pum2, Fmrp,

Ago2, and Rbfox3, we considered the entire set of all targetomes together as background.

Myc and GST cloning procedure with in vitro immunoprecipitation (IP) assays. Human *PUM1* full-

length cDNA was amplified by PCR and subcloned in a pRK5 plasmid containing the Myc tag sequence

(Addgene, pRK5-Myc-Parkin #17612) at the N-terminal by using SalI (New England Biolabs, # R3138S)

and NotI (New England Biolabs, #R0189S) restriction enzymes to replace *Parkin* with *PUM1*. For GST,

the human full-length *PUM1* cDNA was, again, subcloned first in the pRK5 plasmid containing the GST

tag sequence (Addgene, pRK5-HA GST RagC wt, #19304) at the N-terminal by using SalI and NotI

restriction enzymes to replace *RagC* with *PUM1*. Human *FMRP*, *AGO2* and *CNOT1*, full-length cDNA

were cloned and contain the GST tag sequence at the N-terminal as described for GST-*PUM1*.

To introduce the T1035S or R1147W mutations we used the QuikChange II XL Multi Site-Directed

Mutagenesis kit (Agilent Technologies, #200521). The primers for the single mutagenesis experiments

were designed by QuikChange software (Stratagene, San Diego, https://www.genomics.agilent.com/

primerDesignProgram.jsp).

For IP, HEK293T cells were seeded in 6-well plates for 24 h and then transfected with 250 ng of

either WT or mutant PUM1 plasmid with one of the interactors using the jetPRIME Transfection Reagent

(Polyplus transfection, #55-132) as per the manufacturer's protocol. pRK5-Myc empty plasmid (no

cDNA) was used as a negative control. After 48 h, the cells were collected and processed for

immunoprecipitation. Protein lysates were incubated overnight at 4°C with mouse α-Myc antibody

(1:400, [Cell Signaling Technologies, #2276]) on a rotisserie tube rotator. The next day, the beads were

washed four times with an IP lysis buffer and resuspended in 40 μL elution buffer (lysis buffer, NuPAGE

10X Reducing Agent [Invitrogen, #NP0009], NuPAGE LDS sample buffer 4X [Invitrogen, #NP0007])

and boiled at 95°C for 10 minutes before loading the samples in NuPAGE 4%–12% Bis-Tris Gels

(Invitrogen, #NP0335BOX & #NP0336BOX) for further resolution and western blot analysis. Antibodies:

mouse α-Myc antibody [1:2000, (Cell Signaling Technologies, #2276)], rabbit α-GST antibody [1:1000,

(Cell Signaling Technologies, #2625)], mouse α-GAPDH [1:10000, (Millipore, #CB1001)].

Primers. For the qPCR analysis to unambiguously distinguish spliced cDNA from genomic DNA

contamination, specific exon primers were designed to amplify across introns of the gene tested. The

primers for all genes tested were designed with Primer3 (Koressaar & Remm, 2007; Untergasser *et al*,

2012). Cloning primers were manually designed to amplify the longest spliced gene isoform tested; if

there was more than one isoform according to the UCSC Genome Browser (https://genome.ucsc.edu/), we

chose the longest. See **Supplementary Table 8** for primer sequences.

Quantification and Statistical Analysis

Experimental design

For protein and RNA quantification from patient-derived cell lines, we used values from at least six

independent experiments with three biological replicates for each experiment. At every stage of the study,

the experimenter was blinded to the identity of control and patient-derived cell lines. For example, for the

data regarding both human patient-derived cell lines and mice, Experimenter #1 made a list of samples

and controls to be tested, and Experimenter #2 randomized this list and re-labeled the tubes; Experimenter

#2 was the only person with the key to identify the samples. These samples were then distributed to

Experimenter #3 to culture the cells, then to Experimenter #1 to perform western blots and qRT-PCR, and

lastly Experimenters #1 and #4 analyzed the data. Only then was the key applied to identify the samples.

For mouse experiments, the experimenters were randomized and blinded as described above. The number

of animals used and sex, and the specific statistical tests used, are indicated for each experiment in the

figure legends. Sample size was based on previous experience using the same mice (Gennarino *et al.*,

2015).

Software and statistical analysis. Statistical significance was analyzed using GraphPad Prism 8

(https://www.graphpad.com/ scientific-software/prism/) and Excel Software (Microsoft). All data are

presented as mean ± SEM. Statistical details for each experiment can be found in the figures and the

legends. The range of expression levels in qPCR was determined from at least six independent

experiments with three biological replicates by calculating the standard deviation of the **Δ**Ct (Pfaffl,

2001). The range of expression levels in western blots was determined from at least six independent

experiments with at least six biological replicates. P values were calculated by Student's T-test or analysis

of variance with Tukey's post hoc analysis. For the IP and protein quantification in patient cell lines in

Figure 7, we had only one PADDAS patient, so the six repeated experiments were technical replicates

rather than biological replicates. We therefore calculated the statistical significance based on these

technical replicates in comparison to the three biological replicates (i.e., healthy controls).

Data Availability

Materials and reagents

Further information and requests for resources, reagents, and mouse models used in this manuscript

should be directed to and will be fulfilled by Vincenzo A. Gennarino (vag2138@cumc.columbia.edu).

Code and raw data

No software was generated for this project. All software used in this study is publicly available and links

are provided as appropriate in different sections of the Materials and Methods. Mass spectrometry, RNA

sequencing and microRNA sequencing raw data generated during this study are available at PRIDE

Archive at https://www.ebi.ac.uk/pride/archive, and Gene Expression Omnibus (GEO) at

https://www.ncbi.nlm.nih.gov/geo/ with the accession numbers pending.

 Main figure titles and legends

Figure 1. A brain-specific Pum1 interactome. (**A**) Network of putative Pum1 interactors in 10-week-

old mouse brain (circles connected to Pum1 by gray lines). Interactions between interactors (purple lines)

inferred by g:GOSt from Corum and the Human Protein Atlas (see Materials and Methods). The proteins

in each of the 22 clusters are listed to the right. All putative Pum1 interactors are available in

Supplementary Table 1. (**B**) Bubble plots show the top categories from gene ontology analyses of Pum1

interactors from whole brain, hippocampus, cortex, and cerebellum. Only the categories with fold

enrichment >1.5 and FDR<0.05 are shown; not all are labeled because of space limitations. The full list of

gene ontology categories are available in **Supplementary Table 2**.

Figure 2. Validation of Pum1 associations with post-transcriptional RNA-binding proteins in mouse

brain. (**A**) Representative western blot of proteins pulled down by IP against Pum1 compared to IgG

from wild-type mice brain without (*left*) and with (*right*) RNase treatment. In this panel, after IP-Pum1,

we immunoblotted for Pum1 (positive control), Pum2, Fmrp, Ago2, Rbfox3, Cnot1, and Mov10. (See

Materials and Methods.) (**B**) Representative western blot of the reciprocal IPs against Pum2, Fmrp, Ago2,

Rbfox3, Cnot1 or Mov10. Each IP was immunoblotted against Pum1 and the putative interactor. (**C**)

Representative western blots of the same proteins validated in panels **A** and **B** after IP against PUM1 with

or without RNase treatment from HEK293T cell lines. In panels **A** through **C**, IP against IgG was used as

a negative control and Input (1% from the initial protein lysate) as a loading control. The numbers on the

right are the respective molecular weights expressed in kilodaltons (kDa). All the experiments were

repeated at least six times. All mice were sacrificed at 10 weeks of age. See **Figure 2–source data 1** for

the original blots.

Figure 3. Effects of Pum1 loss on interactions among the six RNA-binding proteins. Representative

western blot of the proteins pulled down by (**A**) Pum2, (**B**) Fmrp, (**C**) Rbfox3, (**D**) Mov10, (**E**) Ago2, and

(**F**) Cnot1 from WT and *Pum1$^{-/-}$* mouse brain at 10 weeks of age. IP against IgG was used as a negative

control, and Input (1% from the initial protein lysate) as a loading control. Molecular weights to the right

are expressed in kilodaltons (kDa). All the experiments were repeated at least four times. Since Rbfox3

and Mov10 interactions with Pum1 are RNA-dependent we did not perform IP from *Pum1$^{-/-}$* mouse brain.

See **Figure 3–source data 1** for the original blots.

Figure 4. Pum1 interactors and the microRNA machinery show brain region- and sex-specific

responses to Pum1 loss. Representative western blots of Pum1, Pum2, Fmrp, Ago2, Rbfox3, Cnot1, and

Mov10 in (**A**) cerebellum, (**B**) hippocampus, and (**C**) cortex in both male (left panel) and female (right

panel) WT, *Pum1*[+/-], and *Pum1*[-/-] mice. The same number of males and females at 10 weeks of age was

used for a total of at least 18 mice per genotype. Graphs below show quantification for each protein by

brain region, sex, and genotype. All data were normalized to Gapdh protein levels. All experiments were

performed at least 9 times per genotype from mice at 10 weeks of age (data represent mean ± SEM). *P*

values were calculated by two-tailed Student's t test. *p < 0.05, **p < 0.01, ***p < 0.001. See **Figure 4–**

figure supplement 2A for mRNA quantification for each interactor, brain region, and sex. P1 indicates

Pumilio1. See **Figure 4–Source data 1** for all western blot quantifications. (**D**) Heatmap showing 166

microRNAs from mouse cerebellum in *Pum1*[-/-] male and female mice at 10 weeks of age that were

dysregulated (fold change -3 to +3) relative to wild-type cerebellum. Three cerebella per genotype and

sex were analyzed. The full list of miRNA names and fold changes are available in **Supplementary**

Table 3. (**E**) Heatmap showing mRNA quantification by qPCR for 49 targets co-bound by a minimum of

eight dysregulated miRNAs (>20% change) from panel **D**. Three cerebella per genotype and sex were

analyzed. Statistical significance and magnitude of dysregulation are illustrated for both male and female

in **Figure 4–figure supplement 2B**. The entire list of targets predicted to be co-bound by at least two

miRNAs are reported in **Supplementary Table 4**. See **Figure 4–source data 1** and **2** for the original

blots and quantifications.

Figure 5. Pum1 and its RNA-binding protein interactors share many neuronal mRNA targets. (**A**)

Enrichment plots generated by Gene Set Enrichment Analysis (GSEA) of Pum1, Pum2, Rbfox3, and

Ago2 HITS-CLIP targets plotted again Fmrp HITS-CLIP data ranked by RPKM. Pum1, Pum2, Rbfox3,

and Ago2 targets are enriched at the top 10[th] percentile of the Fmrp targets with FDR=0. (**B**) GSEA

analysis scores of HITS-CLIP data from each negative control (Apc, Nova, Ptpb2, and Mbnl2) plotted

against Fmrp HITS-CLIP data ranked by RPKM. The negative controls have a maximum enrichment

score of 0.36 for Apc ranking at the top 44.7% with FDR=1. (**C**) GSEA analysis scores of Pum1, Pum2,

Rbfox3, and Ago2 HITS-CLIP data plotted against Fmrp HITS-CLIP data divided into 10-percentile bins

and ranked by RPKM shows the shared targets are among the top percentiles of targets for each protein.

(**D**) GSEA analysis scores of the HITS-CLIP data for Fmrp, Pum2, Ago2, Rbfox3 and four negative

controls (Apc, Nova, Ptpb2, and Mbnl2) against Pum1 HITS-CLIP data ranked by RPKM. The targets of

Fmrp, Ago2, Pum2, and Rbfox3 are enriched at the top 5[th] to 18[th] percentile of Pum1 targets. (**E**) GSEA

analysis of the shared targets between Pum1, Pum2, Ago2, and Rbfox3 against Fmrp showing that they

are enriched in the top 5[th] percentile of Fmrp targets ranked by RPKM. (**F**) Pum1, Pum2, Ago2, and

Rbfox3 shared targets plotted against Fmrp HITS-CLIP targets ranked by RPKM and divided into 10-

percentile bins shows that all of their respective targets are enriched at the top 10th percentile of the Fmrp

targets ranked by RPKM. (**G**) GSEA analysis scores of the targets shared by Pum1, Pum2, Ago2, and

Rbfox3 and the four negative controls (Apc, Nova, Ptpb2, and Mbnl2) plotted against Fmrp. At best the

negative controls are enriched at the top 40% with a maximum ES of 0.41. For all the GSEA analyses, the

False Discovery Rate (FDR) was provided by GSEA: **FDR<0.05 and ***FDR<0.01. ES=Enrichment

score (blue line). Note that lowest rank at max percentage indicates stronger targets in the rank (see

Materials and Methods). The full list of shared targets is reported in **Supplementary Table 5**.

Figure 6. The R1147W mutation, but not T1035S, destabilizes PUM1 interactors. (**A**) IP

against PUM1 from PADDAS (R1147W) patient (Subject 1 or S1)-derived fibroblasts confirms

the interactions between PUM1 (used here as a positive control), and PUM2, FMRP, AGO2,

CNOT1, and MOV10. *Bottom panel*: protein quantification shows loss of interaction between

PUM1-R1147W and AGO2, CNOT1 and MOV10 compared to three age- and sex- matched

control fibroblasts. Input (1%) was used as a loading control and IP against IgG was used as a

negative control. (**B**) IP against PUM1 from two PRCA (T1035S) patient (S2 and S3)-derived

lymphoblastoid cell lines confirm the interaction between PUM1 (used here as positive control),

and PUM2, FMRP, AGO2, CNOT1, and MOV10. *Bottom panel*: protein quantification shows a

slight decrease in interactions with FMRP and AGO2, and a slight increase with CNOT1 and

MOV10, compared to age- and sex- matched lymphoblastoid controls. Input (1%) was used as a

loading control and IP against IgG was used as a negative control. In **A** and **B**, the amount of

protein pulled down compared to IP-PUM1 was quantified as follows: $[IP^X/(Input^X/GAPDH^X)]/$

$[IP^{PUM1}/(Input^{PUM1}/GAPDH^{PUM1})]$, where X is the protein of interest. (**C**) Representative western

blots (*left panels*) and relative quantification (*bar graphs to the right*) of protein levels for

PUM1, PUM2, FMRP, AGO2, CNOT1, and MOV10 in PADDAS patient-derived fibroblasts

and PRCA patient-derived lymphoblastoid cells compared to their respective age- and sex-

matched fibroblast (for PADDAS patients) and lymphoblastoid (for PRCA patients) controls.

Data were normalized to Gapdh protein levels. From **A** to **C**, all the experiments were performed

at least six times. Data represent mean ± SEM. P values were calculated by two-tailed Student's t

test. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. See **Figure 6–source data 1** for the original blots.

1014

Figure 7. RBP complexes with PUM1 R1147W are destabilized *in vitro* and shared neuronal targets

are upregulated only in PADDAS. (**A-D**) Representative western blots and relative IP quantification

(*bar graphs*) of IP against Myc-PUM1-WT, Myc-PUM1-T1035S (PRCA), or Myc-PUM1-R1147W

(PADDAS) followed by immunoblotting for: (**A**) GST-AGO2, (**B**) GST-CNOT1, (**C**) GST-FMRP, and

(**D**) GST-PUM1-WT. Myc- and GST-tagged proteins were co-transfected in HEK293T cells in equal

quantities (250ng each). The molecular weights were expressed in kilodaltons (kDa). The amount of

protein pulled down compared to IP-PUM1 was quantified as $[IP^X/(Input^X/GAPDH^X)]/$

$[IP^{PUM1}/(Input^{PUM1}/ GAPDH^{PUM1})]$, where X is the protein of interest. (**E**) qPCR analysis of validated

PUM1-specific targets from PADDAS patient-derived fibroblasts (S1, blue bars) compared to three age-

and sex-matched fibroblast control cell lines, and PRCA patient-derived lymphoblastoid cell lines (S2 and

S3, orange bars) compared to the three age- and sex-matched lymphoblastoid control cell lines. (**F**)

mRNA level quantification of PUM1 neuronal targets in common with FMRP, PUM2, AGO2, and

RBFOX3 (**Figure 5E** and **Supplementary Table 5**) in fibroblasts from subject 1 (PADDAS patient,

R1147W) compared to the three age- and sex-matched control fibroblast lines (blue bars), and in

lymphoblastoid cells from subjects 2 and 3 (PRCA patients, T1035S) compared to the three age- and sex-

matched control lymphoblastoid cell lines (orange bars). For **E** and **F**, all data were normalized to

GAPDH mRNA levels and experiments performed at least three times. Only genes expressed in both

fibroblasts and lymphoblasts are represented here for a total of 10 for panel **E** and 55 for panel **F**. Data

represent mean ± SEM. P values were calculated by two-tailed Student's t test. *$p < 0.05$, **$p < 0.01$,

***$p < 0.001$. The full list of shared targets expressed in fibroblast and lymphoblast cell lines is reported

in **Supplementary Table 5. See Figure 7–source data 1 for the original blots.**

1036

Supplemental figure titles and legends

Figure 1–figure supplement 1. Immunoprecipitation (IP) antibody specificity against Pum1 from

mouse brain and Pum1 interactor validation. (**A**) Pre-IP, IP, and post-IP against Pum1 and IgG from

wild-type mouse brain show the specificity of the antibody used against Pum1. Pre-IP represents 1% from

the initial protein lysate as a loading control while 10% of the protein lysate after IP-Pum1 was loaded as

post-IP. The numbers on the right show molecular weight in kilodaltons (kDa). (**B**) Principal component

analysis (PCA) of IP-Pum1 followed by mass spectrometry (MS) in WT adult mouse brains compared

with IP against IgG; each dot represents two combined samples for a total of 6 samples processed by mass

spectrometry. (**C**) Representative western blot of Rptor, Cad, Anapc1 and Fzr1 proteins pulled down by

IP against Pum1 (used here as positive control) from WT mice. (**D**) Representative western blot of the

reciprocal co-Ips against Anapc1, Fzr1, Rptor, and Cad. Each co-IP was immunoblotted against Pum1 and

the pulled down protein used here as a reference protein to confirm the respective protein-protein

interaction. For **C** and **D**, IP against IgG was used as a negative control, and Input (1% from the initial

protein lysate) as a loading control. Molecular weights are expressed in kilodaltons (kDa) to the right. All

the experiments were repeated at least four times. All mice were sacrificed at 10 weeks of age with an

equal number or male and female. See **Figure 1–figure supplement 1–source data 1** for the original

blots.

Figure 1–figure supplement 2. Pum1 interactors differ by brain region. (**A**) Principal component

analysis (PCA) of IP-Pum1 followed by mass spectrometry (MS) in cortex, hippocampus, and cerebellum

from WT mice. IP against IgG was used as a negative control. Each dot represents a total of 3 samples

processed by MS for each brain region. (**B**) Pum1 interactome from cerebellum, hippocampus, cortex and

the rest of the brain (i.e., excluding those three regions) showing all Pum1 putative interactors with a fold

change > 1.5 (see also **Supplementary Table 1**), for a total of 1,500 proteins. Node colors represent

different brain regions or the overlap between two or more brain regions as noted. (**C**) After IP against

Pum1 we immunoblotted for Pum1 (positive control), Anapc1, Cnot1 and Pum2. While Cnot1 and

Anapc1 can be pulled down from all three brain regions, Pum2 can only be pulled down from cortex. For

A-C we combined cerebellar and cortical tissues from eight wild-type mice (4 male and 4 female) for a

total of 24 mice in triplicate for cerebellum and cortex, while for hippocampus we combined tissues from

ten wild-type mice (5 female and 5 male) for a total of 30 mice in triplicate. All mice were 10 weeks of

age. IP against IgG was used as a negative control. Molecular protein weights are expressed in kilodaltons

(kDa). (**D-F**) The three most enriched protein complexes among the Pum1 interactors for each brain

region are shown in (**D**) for the anaphase promoting complex/cyclosome (APC/C); (**E**) for the mTOR

pathway; (**F**) for RNA-silencing (RISC), CCR4-NOT, translation initiation and polyA binding. Edge

colors (**D**, **E**, and **F**) represent a specific brain region: red for cortex, green for cerebellum, blue for

hippocampus, and gray for proteins in common between two or more brain regions. (**G**) Western blot

analysis at 10 weeks of age to evaluate Pum2 expression levels in eight different brain regions as well as

whole brain. Pum2 is highly expressed in olfactory bulbs and amygdala, and expressed at similar levels in

hippocampus, cerebellum, and cortex. HP: hippocampus; CB: cerebellum; CX: cortex; MB: midbrain;

OB: olfactory bulbs; AM: amygdala; SN: substantia nigra pars compacta; BS: brain stem; WB: whole

brain. See **Figure 1–figure supplement 2–source data 1** for the original blots.

Figure 2–figure supplement 1. Immunoprecipitation (IP) against Pum1 from whole brain in *Pum1*[-/-]

mice. IP against Pum1 in *Pum1*[-/-] mouse demonstrates the complete absence of Pum1 and thus the

specificity of the anti-Pum1 antibody. IP against IgG was used as a negative control, and Input (1% from

the initial protein lysate) as a loading control. Molecular weights at the right are in kilodaltons (kDa). All

the experiments were repeated at least four times. IP against IgG was used as a negative control, and Input

(1% from the initial protein lysate) as a loading control. Equal numbers of male and female mice were

sacrificed at 10 weeks of age. See **Figure 2–figure supplement 1–source data 1** for the original blots.

Figure 3–figure supplement 1. Immunoprecipitation (IP) against Pum1 interactors from whole

brain in wild-type and *Pum1*$^{-/-}$ mice with and without RNase treatment. (**A-F**) Representative western

blot of the proteins pulled down by (**A**) Pum2, (**B**) Fmrp, (**C**) Ago2, (**D**) Cnot1, (**E**) Mov10, and (**F**)

Rbfox3 from wild-type (WT, *left panel*) and *Pum1*$^{-/-}$ (*right panel*) mouse brain without (*left panel*) and

with (*right panel*) RNase treatment. IP against IgG was used as a negative control, and Input (1% from

the initial protein lysate) as a loading control. Molecular weights at the right are in kilodaltons (kDa). All

the experiments were repeated at least four times. IP against IgG was used as a negative control, and Input

(1% from the initial protein lysate) as a loading control. Equal numbers of male and female mice were

sacrificed at 10 weeks of age. See **Figure 3–figure supplement 1–source data 1** for the original blots.

Figure 4–figure supplement 1. Protein and mRNA quantification from whole brain in WT, *Pum1*$^{+/-}$

and *Pum1*$^{-/-}$ mice. (**A**) Representative western blot with relative quantifications of Pum1, Pum2, Fmrp,

Ago2, Rbfox3, Cnot1, and Mov10 from the whole brain in WT, *Pum1*$^{+/-}$ and *Pum1*$^{-/-}$ mice. All data were

normalized to Gapdh protein levels. The numbers on the right are the respective molecular weights

expressed in kilodaltons (kDa). (**B**) mRNA level quantification by qPCR of *Pum1*, *Pum2*, *Fmrp*, *Ago2*,

Rbfox3, *Cnot1*, and *Mov10* from whole brain in WT, *Pum1*$^{+/-}$ and *Pum1*$^{-/-}$ mice. Again, all data were

normalized to *Gapdh* mRNA levels. All the experiments were conducted with equal number of male (at

least 6 per genotype) and female (at least 6 per genotype) mice at 10 weeks of age, for a total of at least

12 mice per genotype, performed at least six times per genotype (data represent mean ± SEM). The *p*

values were calculated by the Student's t test. *p < 0.05, **p < 0.01, ***p < 0.001. See **Figure 4–figure**

supplement 1–source data 1 for the original blots.

Figure 4–figure supplement 2. mRNA quantification of Pum1 interactors by brain region and sex

and the 49 targets co-bound by at least eight dysregulated miRNAs in mouse cerebellum. (**A**)

mRNA levels in cerebellum, hippocampus, and cortex in male and female for all the validated Pum1

interactors. The same number of mice were used here as in **Figure 4A-C** for a total of at least 9 mice per

genotype and sex at 10 weeks of age. All data were normalized to *Gapdh* mRNA levels. All the

experiments were performed at least six times (data represent mean ± SEM). (**B**) qPCR in cerebellum of

male (*left*) and female (*right*) mice at 10 weeks of age for the 49 targets co-bound by at least eight

dysregulated miRNAs (with minimum 20% change in expression) from **Figure 4E** and **Supplementary**

Table 4. All the experiments were performed in triplicate for three different biological replicates for both

male and female (data represent mean ± SEM). The *p* values were calculated by two-tailed Student's t

test. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 5–figure supplement 1. Pum1, Pum2, Ago2, Fmrp, and Rbfox3 share a number of targets in

the mouse brain. (**A**) Gene Set Enrichment Analysis (GSEA) of Pum1 HITS-CLIP data plotted against

HITS-CLIP data from the negative controls (RNA binding proteins that did not show up in the Pum1

interactome: Ptpb2, Nova, Apc, and Mbnl2) reveals no significant enrichment. (**B-C**) Gene ontology

analysis of the HITS-CLIP targets shared between Pum1, Pum2, Fmrp, Ago2, and Rbfox3 reveals

enrichment for certain (**B**) biological functions and (**C**) cellular localization. Only categories with

FDR<0.05 and fold enrichment > 5 were plotted in **B** and **C**. (**D**) Venn diagram of miRNAs identified by

Pum1, and Pum2 shows almost 100% overlap with the miRNAs pulled down by Fmrp HITS-CLIP. The

full list of shared miRNAs is reported in **Supplementary Table 6**. (**E**) GSEA of HITS-CLIP data

normalized against gene expression in mouse brain at 3 weeks of age (see Materials and Methods) of

Pum1, Pum2, Rbfox3, and Ago2 against Fmrp HITS-CLIP data ranked by normalized RPKM. (**F**) GSEA

of HITS-CLIP data normalized against gene expression in mouse brain at 3 weeks of age (see Materials

and Methods) of Mbnl2, Apc, Nova, and Ptpb2 plotted against Fmrp HITS-CLIP data ranked by

normalized RPKM. Mbnl2, Ptpb2, Apc, and Nova were used as negative controls. (**G**) Fmrp, Pum2,

Ago2, Rbfox3 shared targets, and targets of the four negative controls, plotted against Pum1 HITS-CLIP

data ranked by normalized RPKM. (**H**) Pum1, Pum2, Ago2, Rbfox3 shared targets plotted against Fmrp

HITS-CLIP data ranked by normalized RPKM. The RNAseq gene expression changes are reported in

Supplementary Table 7. For all GSEA analyses the False Discovery Rate (FDR) was provided by

GSEA, ***FDR < 0.01. ES=Enrichment score (blue line). Note that lowest rank at max percentage ES

indicates stronger targets in the rank.

Figure 6–figure supplement 1. PUM1 validation experiments in patient-derived cells. (**A-B**) Pre-IP,

IP, and post-IP against PUM1 and IgG from (**A**) PADDAS fibroblasts and (**B**) PRCA lymphoblastoid

cells. In both cell lines we were able to pull down 100% of PUM1. Pre-IP represents 1% from the initial

protein lysate as a loading control, while 10% of the protein lysate was loaded as post-IP. Molecular

weights provided at right in kilodaltons (kDa). (**C**) mRNA level quantification for all of the

immunoblotted proteins in **Figure 6C** in PADDAS and PRCA patient-derived cell lines compared to their

respective age-, sex-, and cell-type-matched controls. All data were normalized to *GAPDH* mRNA levels,

and experiments performed at least six times. Data represent mean ± SEM. See **Figure 6–figure**

supplement 1–source data 1 for the original blots.

Figure 7–figure supplement 1. PUM1 validation experiments *in vitro*. (**A-D**) Representative western

blots (upper panel) and relative IP quantification (bottom panel) of IP against Myc-PUM1-WT, Myc-

PUM1-T1035S (PRCA), or Myc-PUM1-R1147W (PADDAS) followed by immunoblotting for: (**A**) GST-

PABPC1, (**B**) GST-PABPC4, (**C**) GST-EIF4A3, and (**D**) GST-EIF4G3. Myc- and GST-tagged proteins

were co-transfected in HEK293T cells in equal quantities (250ng each). The molecular weights were

expressed in kilodaltons (kDa). The amount of protein pulled down compared to IP-PUM1 was quantified

as $[IP^X/(Input^X/GAPDH^X)]/[IP^{PUM1}/(Input^{PUM1}/GAPDH^{PUM1})]$, where X is the protein of interest.

(**E**) PUM1 half-life in HEK293T at different time points, from zero (cycloheximide treatment) to 5h total

upon transfection of Myc-PUM1-WT, Myc-PUM1-T1035S, or Myc-PUM1-R1147W. The half-life of the

R1147W mutant is significantly shorter than that of WT and T1035S. All data were normalized to

GAPDH and compared to time zero. The experiments were performed at least four times and the data

represent mean ± SEM. The *p* values were calculated by two-tailed Student's t test. **$p < 0.01$, ***p <

0.001. (**F**) Representative western blots of IP with RNase treatment against Myc-PUM1-WT, Myc-

PUM1-T1035S (PRCA), or Myc-PUM1-R1147W (PADDAS) followed by immunoblotting to

test binding between PUM1 proteins without the RNA. The numbers on the right are the respective

molecular weights expressed in kilodaltons (kDa). See **Figure 7–figure supplement 1–source data 1** for

the original blots.

Supplementary table titles and legends

Supplementary Table 1. Pum1 mass spectrometry results from whole brain and three separate

brain regions in 10-week-old mice. All the proteins have an FDR < 0.05, and only proteins with a Ratio

LFQ-Pum1-WT/LFQ-IgG-WT > 1.5 are reported here.

Supplementary Table 2. Gene Ontology analysis of Pum1 mass spectrometry from whole brain and

the three brain regions. Only categories with fold enrichment of >1.5 and FDR<0.05 are shown.

Supplementary Table 3. MicroRNA names and sequences with log2 fold change as represented in

Figure 4D. PC = Predicted Candidate.

Supplementary Table 4. Predicted target genes for all the differentially expressed miRNA (with at

least 25% change) in cerebellum of male and female in wild-type and *Pum1$^{-/-}$* mice at 10 weeks of

age. The target genes tested to generated Figure 4E and Figure 4–figure supplement 2B have at least 8 co-

bound miRNAs.

Supplementary Table 5. HITS-CLIP targets common to Pum1, Pum2, Ago2, Rbfox3, and Fmrp.

Supplementary Table 6. microRNAs that overlap between Pum1, Pum2, and Fmrp HITS-CLIP

data.

Supplementary Table 7. RNAseq in mouse brain at 3 weeks of age.

1177

1178 **Supplementary Table 8. All primer sequences used in this study.**

References

Agarwal V, Bell GW, Nam JW, Bartel DP (2015) Predicting effective microRNA target sites in mammalian mRNAs. *Elife* 4

Almeida A, Bolanos JP, Moreno S (2005) Cdh1/Hct1-APC is essential for the survival of postmitotic neurons. *The Journal of Neuroscience* 25: 8115-8121

Bah MG, Rodriguez D, Cazeneuve C, Mochel F, Devos D, Suppiej A, Roubertie A, Meunier I, Gitiaux C, Curie A *et al* (2020) Deciphering the natural history of SCA7 in children. *Eur J Neurol* 27: 2267-2276

Blackinton JG, Keene JD (2014) Post-transcriptional RNA regulons affecting cell cycle and proliferation. *Semin Cell Dev Biol* 34: 44-54

Blewett NH, Goldstrohm AC (2012) A eukaryotic translation initiation factor 4E-binding protein promotes mRNA decapping and is required for PUF repression. *Molecular and Cellular Biology* 32: 4181-4194

Bohn JA, Van Etten JL, Schagat TL, Bowman BM, McEachin RC, Freddolino PL, Goldstrohm AC (2018) Identification of diverse target RNAs that are functionally regulated by human Pumilio proteins. *Nucleic Acids Research* 46: 362-386

Bonnemason-Carrere P, Morice-Picard F, Pennamen P, Arveiler B, Fergelot P, Goizet C, Hellegouarch M, Lacombe D, Plaisant C, Raclet V *et al* (2019) PADDAS syndrome associated with hair dysplasia caused by a de novo missense variant of PUM1. *American Journal of Medical Genetics Part A* 179: 1030-1033

Charizanis K, Lee KY, Batra R, Goodwin M, Zhang C, Yuan Y, Shiue L, Cline M, Scotti MM, Xia G *et al* (2012) Muscleblind-like 2-mediated alternative splicing in the developing brain and dysregulation in myotonic dystrophy. *Neuron* 75: 437-450

Chartier-Harlin MC, Kachergus J, Roumier C, Mouroux V, Douay X, Lincoln S, Levecque C, Larvor L, Andrieux J, Hulihan M *et al* (2004) Alpha-synuclein locus duplication as a cause of familial Parkinson's disease. *Lancet* 364: 1167-1169

Chen D, Zheng W, Lin A, Uyhazi K, Zhao H, Lin H (2012) Pumilio 1 suppresses multiple activators of p53 to safeguard spermatogenesis. *Current Biology: CB* 22: 420-425

Cheng S, Gallie DR (2007) eIF4G, eIFiso4G, and eIF4B bind the poly(A)-binding protein through overlapping sites within the RNA recognition motif domains. *The Journal of Biological Chemistry* 282: 25247-25258

Chi SW, Zang JB, Mele A, Darnell RB (2009) Argonaute HITS-CLIP decodes microRNA-mRNA interaction maps. *Nature* 460: 479-486

Conboy JG (2017) Developmental regulation of RNA processing by Rbfox proteins. *Wiley Interdisciplinary Reviews RNA* 8

Cox J, Mann M (2008) MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nature Biotechnology* 26: 1367-1372

1227

1228 Cox J, Neuhauser N, Michalski A, Scheltema RA, Olsen JV, Mann M (2011) Andromeda: a peptide
1229 search engine integrated into the MaxQuant environment. *J Proteome Res* 10: 1794-1805

1230

1231 Darnell JC, Richter JD (2012) Cytoplasmic RNA-binding proteins and the control of complex brain
1232 function. *Cold Spring Harb Perspect Biol* 4: a012344

1233

1234 Dassi E (2017) Handshakes and fights: the regulatory interplay of RNA-binding proteins. *Front Mol*
1235 *Biosci* 4: 67

1236

1237 De Maio A, Yalamanchili HK, Adamski CJ, Gennarino VA, Liu Z, Qin J, Jung SY, Richman R, Orr H,
1238 Zoghbi HY (2018) RBM17 interacts with U2SURP and CHERP to regulate expression and splicing of
1239 RNA-processing proteins. *Cell Reports* 25: 726-736 e727

1240

1241 Di Grazia A, Marafini I, Pedini G, Di Fusco D, Laudisi F, Dinallo V, Rosina E, Stolfi C, Franze E, Sileri
1242 P *et al* (2021) The Fragile X mental retardation protein regulates RIPK1 and colorectal cancer resistance
1243 to necroptosis. *Cell Mol Gastroenterol Hepatol* 11: 639-658

1244

1245 Dittmann K, Mayer C, Czemmel S, Huber SM, Rodemann HP (2017) New roles for nuclear EGFR in
1246 regulating the stability and translation of mRNAs associated with VEGF signaling. *PloS One* 12:
1247 e0189087

1248

1249 Edwards TA, Pyle SE, Wharton RP, Aggarwal AK (2001) Structure of Pumilio reveals similarity between
1250 RNA and peptide binding motifs. *Cell* 105: 281-289

1251

1252 Enwerem, III, Elrod ND, Chang CT, Lin A, Ji P, Bohn JA, Levdansky Y, Wagner EJ, Valkov E,
1253 Goldstrohm A (2021) Human Pumilio proteins directly bind the CCR4-NOT deadenylase complex to
1254 regulate the transcriptome. *RNA* 27: 445-464.

1255

1256 Friend K, Campbell ZT, Cooke A, Kroll-Conner P, Wickens MP, Kimble J (2012) A conserved PUF-
1257 Ago-eEF1A complex attenuates translation elongation. *Nature Structural & Molecular Biology* 19: 176-
1258 183

1259

1260 Gallie DR (1998) A tale of two termini: a functional interaction between the termini of an mRNA is a
1261 prerequisite for efficient translation initiation. *Gene* 216: 1-11

1262 Gehman LT, Meera P, Stoilov P, Shiue L, O'Brien JE, Meisler MH, Ares M, Jr., Otis TS, Black DL
1263 (2012) The splicing regulator Rbfox2 is required for both cerebellar development and mature motor
1264 function. *Genes & Development* 26: 445-460

1265

1266 Gennarino VA, D'Angelo G, Dharmalingam G, Fernandez S, Russolillo G, Sanges R, Mutarelli M,
1267 Belcastro V, Ballabio A, Verde P *et al* (2012) Identification of microRNA-regulated gene networks by
1268 expression analysis of target genes. *Genome Research* 22: 1163-1172

1269

1270 Gennarino VA, Palmer EE, McDonell LM, Wang L, Adamski CJ, Koire A, See L, Chen CA, Schaaf CP,
1271 Rosenfeld JA *et al* (2018) A mild PUM1 mutation is associated with adult-onset ataxia, whereas
1272 haploinsufficiency causes developmental delay and seizures. *Cell* 172: 924-936 e911

1273

Gennarino VA, Singh RK, White JJ, De Maio A, Han K, Kim JY, Jafar-Nejad P, di Ronza A, Kang H,
Sayegh LS *et al* (2015) Pumilio1 haploinsufficiency leads to SCA1-like neurodegeneration by increasing
wild-type Ataxin1 levels. *Cell* 160: 1087-1098

Gieffers C, Peters BH, Kramer ER, Dotti CG, Peters JM (1999) Expression of the CDH1-associated form
of the anaphase-promoting complex in postmitotic neurons. *PNAS* 96: 11317-11322

Giurgiu M, Reinhard J, Brauner B, Dunger-Kaltenbach I, Fobo G, Frishman G, Montrone C, Ruepp A
(2019) CORUM: the comprehensive resource of mammalian protein complexes-2019. *Nucleic Acids*
Research 47: D559-D563

Goldstrohm AC, Hall TMT, McKenney KM (2018) Post-transcriptional regulatory functions of
mammalian pumilio proteins. *Trends in Genetics* 34: 92-990

Hentze MW, Castello A, Schwarzl T, Preiss T (2018) A brave new world of RNA-binding proteins.
Nature reviews Molecular Cell Biology 19: 327-341

Humphrey J, Birsa N, Milioto C, McLaughlin M, Ule AM, Robaldo D, Eberle AB, Krauchi R, Bentham
M, Brown AL *et al* (2020) FUS ALS-causative mutations impair FUS autoregulation and splicing factor
networks through intron retention. *Nucleic Acids Research* 48: 6889-6905

Imaizumi T, Mogami Y, Okamoto N, Yamamoto-Shimojima K, Yamamoto T (2019) De novo 1p35.2
microdeletion including PUM1 identified in a patient with sporadic West syndrome. *Congenit Anom*
(Kyoto) 59: 193-194

Jin Z, Feng H, Liang J, Jing X, Zhao Q, Zhan L, Shen B, Cheng X, Su L, Qiu W (2020) FGFR3 big up tri,
open7-9 promotes tumor progression via the phosphorylation and destabilization of ten-eleven
translocation-2 in human hepatocellular carcinoma. *Cell Death Dis* 11: 903

Kedde M, van Kouwenhove M, Zwart W, Oude Vrielink JA, Elkon R, Agami R (2010) A Pumilio-
induced RNA structure switch in p27-3' UTR controls miR-221 and miR-222 accessibility. *Nature Cell*
Biology 12: 1014-1020

Keene JD (2007a) Biological clocks and the coordination theory of RNA operons and regulons. *Cold*
Spring Harb Symp Quant Biol 72: 157-165

Keene JD (2007b) RNA regulons: coordination of post-transcriptional events. *Nature Reviews Genetics* 8:
533-543

Keene JD, Lager PJ (2005) Post-transcriptional operons and regulons co-ordinating gene expression.
Chromosome Res 13: 327-337

Kenny PJ, Kim M, Skariah G, Nielsen J, Lannom MC, Ceman S (2020) The FMRP-MOV10 complex: a
translational regulatory switch modulated by G-Quadruplexes. *Nucleic Acids Research* 48: 862-878

Kenny PJ, Zhou H, Kim M, Skariah G, Khetani RS, Drnevich J, Arcila ML, Kosik KS, Ceman S (2014)
MOV10 and FMRP regulate AGO2 association with microRNA recognition elements. *Cell Reports* 9:
1729-1741

Khalil B, Morderer D, Price PL, Liu F, Rossoll W (2018) mRNP assembly, axonal transport, and local
translation in neurodegenerative diseases. *Brain Res* 1693: 75-91

1325

1326   Kim K, Hessl D, Randol JL, Espinal GM, Schneider A, Protic D, Aydin EY, Hagerman RJ, Hagerman PJ
1327   (2019) Association between IQ and FMR1 protein (FMRP) across the spectrum of CGG repeat
1328   expansions. *PloS One* 14: e0226811

1329

1330   Koressaar T, Remm M (2007) Enhancements and modifications of primer design program Primer3.
1331   *Bioinformatics* 23: 1289-1291

1332

1333   Lai KL, Liao YC, Tsai PC, Hsiao CT, Soong BW, Lee YC (2019) Investigating PUM1 mutations in a
1334   Taiwanese cohort with cerebellar ataxia. *Parkinsonism Relat Disord* 66: 220-223

1335

1336   Le H, Tanguay RL, Balasta ML, Wei CC, Browning KS, Metz AM, Goss DJ, Gallie DR (1997)
1337   Translation initiation factors eIF-iso4G and eIF-4B interact with the poly(A)-binding protein and increase
1338   its RNA binding activity. *The Journal of Biological Chemistry* 272: 16247-16255

1339   Lee EJ, Seo E, Kim JW, Nam SA, Lee JY, Jun J, Oh S, Park M, Jho EH, Yoo KH *et al* (2020) TAZ/Wnt-
1340   beta-catenin/c-MYC axis regulates cystogenesis in polycystic kidney disease. *PNAS* 117: 29001-29012

1341

1342   Lee JT, Wheeler TC, Li L, Chin LS (2008) Ubiquitination of alpha-synuclein by Siah-1 promotes alpha-
1343   synuclein aggregation and apoptotic cell death. *Human Molecular Genetics* 17: 906-917

1344

1345   Leeb M, Dietmann S, Paramor M, Niwa H, Smith A (2014) Genetic exploration of the exit from self-
1346   renewal using haploid embryonic stem cells. *Cell Stem Cell* 14: 385-393

1347

1348   Licatalosi DD, Yano M, Fak JJ, Mele A, Grabinski SE, Zhang C, Darnell RB (2012) Ptbp2 represses
1349   adult-specific splicing to regulate the generation of neuronal precursors in the embryonic brain. *Genes &*
1350   *Development* 26: 1626-1642

1351

1352   Lin H, Spradling AC (1997) A novel group of pumilio mutations affects the asymmetric division of
1353   germline stem cells in the Drosophila ovary. *Development* 124: 2463-2476

1354

1355   Liu GY, Sabatini DM (2020) mTOR at the nexus of nutrition, growth, ageing and disease. *Nature*
1356   *Reviews Molecular Cell Biology* 21: 183-203

1357

1358   Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq
1359   data with DESeq2. *Genome Biology* 15: 550

1360

1361   Lukong KE, Chang KW, Khandjian EW, Richard S (2008) RNA-binding proteins in human genetic
1362   disease. *Trends in Genetics:* 24: 416-425

1363

1364   Mandelboum S, Manber Z, Elroy-Stein O, Elkon R (2019) Recurrent functional misinterpretation of
1365   RNA-seq data caused by sample-specific gene length bias. *PLoS Biol* 17: e3000481

1366

1367   Mauger O, Lemoine F, Scheiffele P (2016) Targeted intron retention and excision for rapid gene
1368   regulation in response to neuronal activity. *Neuron* 92: 1266-1278

1369

Maurin T, Lebrigand K, Castagnola S, Paquet A, Jarjat M, Popa A, Grossi M, Rage F, Bardoni B (2018)
HITS-CLIP in various brain areas reveals new targets and new modalities of RNA binding by fragile X
mental retardation protein. *Nucleic Acids Research* 46: 6344-6355

McCabe MP, Cullen ER, Barrows CM, Shore AN, Tooke KI, Laprade KA, Stafford JM, Weston MC
(2020) Genetic inactivation of mTORC1 or mTORC2 in neurons reveals distinct functions in
glutamatergic synaptic transmission. *Elife* 9: e51440.

Menon KP, Sanyal S, Habara Y, Sanchez R, Wharton RP, Ramaswami M, Zinn K (2004) The
translational repressor Pumilio regulates presynaptic morphology and controls postsynaptic accumulation
of translation factor eIF-4E. *Neuron* 44: 663-676

Miles WO, Tschop K, Herr A, Ji JY, Dyson NJ (2012) Pumilio facilitates miRNA regulation of the E2F3
oncogene. *Genes & Development* 26: 356-368

Miller MA, Olivas WM (2011) Roles of Puf proteins in mRNA degradation and translation. *Wiley*
Interdisciplinary Reviews RNA 2: 471-492

Muraro NI, Weston AJ, Gerber AP, Luschnig S, Moffat KG, Baines RA (2008) Pumilio binds para
mRNA and requires Nanos and Brat to regulate sodium current in Drosophila motoneurons. *The Journal*
of Neuroscience 28: 2099-2109

Orr HT, Zoghbi HY (2007) Trinucleotide repeat disorders. *Annual review of neuroscience* 30: 575-621
Otasek D, Morris JH, Boucas J, Pico AR, Demchak B (2019) Cytoscape Automation: empowering
workflow-based network analysis. *Genome Biology* 20: 185

Pfaffl MW (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic*
Acids Research 29: e45

Preitner N, Quan J, Nowakowski DW, Hancock ML, Shi J, Tcherkezian J, Young-Pearse TL, Flanagan
JG (2014) APC is an RNA-binding protein, and its interactome provides a link to neural development and
microtubule assembly. *Cell* 158: 368-382

Qiu C, Dutcher RC, Porter DF, Arava Y, Wickens M, Hall TMT (2019) Distinct RNA-binding modules
in a single PUF protein cooperate to determine RNA specificity. *Nucleic Acids Research* 47: 8770-8784

Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, Vilo J (2019) g:Profiler: a web server for
functional enrichment analysis and conversions of gene lists (2019 update). *Nucleic Acids Research* 47:
W191-W198

Ravanidis S, Kattan FG, Doxakis E (2018) Unraveling the pathways to neuronal homeostasis and disease:
mechanistic insights into the role of RNA-binding proteins and associated factors. *Int J Mol Sci* 19: 2280.

Rousseaux MWC, Tschumperlin T, Lu HC, Lackey EP, Bondar VV, Wan YW, Tan Q, Adamski CJ,
Friedrich J, Twaroski K *et al* (2018) ATXN1-CIC Complex is the primary driver of cerebellar pathology
in spinocerebellar ataxia type 1 through a gain-of-function mechanism. *Neuron* 97: 1235-1243 e1235
1415

Rovelet-Lecrux A, Hannequin D, Raux G, Le Meur N, Laquerriere A, Vital A, Dumanchin C, Feuillette S, Brice A, Vercelletto M *et al* (2006) APP locus duplication causes autosomal dominant early-onset Alzheimer disease with cerebral amyloid angiopathy. *Nature Genetics* 38: 24-26

Salcedo-Arellano MJ, Dufour B, McLennan Y, Martinez-Cerdeno V, Hagerman R (2020) Fragile X syndrome and associated disorders: clinical aspects and pathology. *Neurobiology of Disease* 136: 104740

Sandweiss AJ, Brandt VL, Zoghbi HY (2020) Advances in understanding of Rett syndrome and MECP2 duplication syndrome: prospects for future therapies. *Lancet Neurol* 19: 689-698

Shah S, Molinaro G, Liu B, Wang R, Huber KM, Ritcher JD (2020) FMRP control of ribosome translocation promotes chromatin modifications and alternative splicing of neuronal genes linked to autism. *Cell Reports* 30: 4459-4472

Shevchenko A, Tomas H, Havlis J, Olsen JV, Mann M (2006) In-gel digestion for mass spectrometric characterization of proteins and proteomes. *Nature Protocols* 1: 2856-2860

Singh K, Gaur P, Prasad S (2007) Fragile x mental retardation (Fmr-1) gene expression is down regulated in brain of mice during aging. *Mol Biol Rep* 34: 173-181

Singh K, Prasad S (2008) Differential expression of Fmr-1 mRNA and FMRP in female mice brain during aging. *Mol Biol Rep* 35: 677-684

Sivakumar S, Gorbsky GJ (2015) Spatiotemporal regulation of the anaphase-promoting complex in mitosis. *Nature Reviews Molecular Cell Biology* 16: 82-94

Smidak R, Sialana FJ, Kristofova M, Stojanovic T, Rajcic D, Malikovic J, Feyissa DD, Korz V, Hoeger H, Wackerlig J *et al* (2017) Reduced levels of the synaptic functional regulator FMRP in dentate gyrus of the Aging Sprague-Dawley rat. *Front Aging Neurosci* 9: 384

Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES *et al* (2005) Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *PNAS* 102: 15545-15550

Takeguchi R, Takahashi S, Kuroda M, Tanaka R, Suzuki N, Tomonoh Y, Ihara Y, Sugiyama N, Itoh M (2020) MeCP2_e2 partially compensates for lack of MeCP2_e1: a male case of Rett syndrome. *Mol Genet Genomic Med* 8: e1088

Tan Q, Zoghbi HY (2019) Mouse models as a tool for discovering new neurological diseases. *Neurobiol Learn Mem* 165: 106902

Temme C, Simonelig M, Wahle E (2014) Deadenylation of mRNA by the CCR4-NOT complex in Drosophila: molecular and developmental aspects. *Front Genet* 5: 143

Tereshchenko A, Magnotta V, Epping E, Mathews K, Espe-Pfeifer P, Martin E, Dawson J, Duan W, Nopoulos P (2019) Brain structure in juvenile-onset Huntington disease. *Neurology* 92: e1939-e1947

Thul PJ, Akesson L, Wiking M, Mahdessian D, Geladaki A, Ait Blal H, Alm T, Asplund A, Bjork L, Breckels LM *et al* (2017) A subcellular map of the human proteome. *Science* 356

1465

1466 Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, Mann M, Cox J (2016) The Perseus
1467 computational platform for comprehensive analysis of (prote)omics data. *Nat Methods* 13: 731-740
1468

1469 Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, Rozen SG (2012) Primer3--new
1470 capabilities and interfaces. *Nucleic Acids Research* 40: e115
1471

1472 Uyhazi KE, Yang Y, Liu N, Qi H, Huang XA, Mak W, Weatherbee SD, de Prisco N, Gennarino VA,
1473 Song X *et al* (2020) Pumilio proteins utilize distinct regulatory mechanisms to achieve complementary
1474 functions required for pluripotency and embryogenesis. *PNAS* 7:7851-7862.
1475

1476 Van Etten J, Schagat TL, Hrit J, Weidmann CA, Brumbaugh J, Coon JJ, Goldstrohm AC (2012) Human
1477 Pumilio proteins recruit multiple deadenylases to efficiently repress messenger RNAs. *The Journal of*
1478 *Biological Chemistry* 287: 36370-36383
1479

1480 Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F (2002) Accurate
1481 normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control
1482 genes. *Genome Biology* 3(7): RESEARCH0034
1483

1484 Wamsley B, Jaglin XH, Favuzzi E, Quattrocolo G, Nigro MJ, Yusuf N, Khodadadi-Jamayran A, Rudy B,
1485 Fishell G (2018) Rbfox1 mediates cell-type-specific splicing in cortical interneurons. *Neuron* 100: 846-
1486 859 e847
1487

1488 Wang X, Zamore PD, Hall TM (2001) Crystal structure of a Pumilio homology domain. *Molecular Cell*
1489 7: 855-865
1490

1491 Weidmann CA, Raynard NA, Blewett NH, Van Etten J, Goldstrohm AC (2014) The RNA binding
1492 domain of Pumilio antagonizes poly-adenosine binding protein and accelerates deadenylation. *RNA* 20:
1493 1298-1319
1494

1495 Weyn-Vanhentenryck SM, Mele A, Yan Q, Sun S, Farny N, Zhang Z, Xue C, Herre M, Silver PA, Zhang
1496 MQ *et al* (2014) HITS-CLIP and integrative modeling define the Rbfox splicing-regulatory network
1497 linked to brain development and autism. *Cell Reports* 6: 1139-1152
1498

1499 Zamore PD, Bartel DP, Lehmann R, Williamson JR (1999) The PUMILIO-RNA interaction: a single
1500 RNA-binding domain monomer recognizes a bipartite target sequence. *Biochemistry* 38: 596-604
1501

1502 Zamore PD, Williamson JR, Lehmann R (1997) The Pumilio protein binds RNA through a conserved
1503 domain that defines a new class of RNA-binding proteins. *RNA* 3: 1421-1433
1504

1505 Zhang C, Frias MA, Mele A, Ruggiu M, Eom T, Marney CB, Wang H, Licatalosi DD, Fak JJ, Darnell RB
1506 (2010) Integrative modeling defines the Nova splicing-regulatory network and its combinatorial controls.
1507 *Science* 329: 439-443
1508

1509 Zhang M, Chen D, Xia J, Han W, Cui X, Neuenkirchen N, Hermes G, Sestan N, Lin H (2017) Post-
1510 transcriptional regulation of mouse neurogenesis by Pumilio proteins. *Genes & Development* 31: 1354-
1511 1369
1512

1513   Zoghbi HY, Bear MF (2012) Synaptic dysfunction in neurodevelopmental disorders associated with
1514   autism and intellectual disabilities. *Cold Spring Harb Perspect Biol* 4: a009886
1515
1516   Zylka MJ, Simon JM, Philpot BD (2015) Gene length matters in neurons. *Neuron* 86: 353-355

1517

Figure 1, Botta et al.

A

Network diagram centered on **Pum1**, with labeled nodes: Cul4b, Papbc4, Nanos1, Akt3, St13, Grb2, Nae1, Rictor, Camkk2, Anapc5, and clusters numbered 1–22.

Cluster 1	Cluster 6	Cluster 11	Cluster 18
Acin1	D1Pas1	Cad	Eea1
Ddx3x	Khdrbs1	Lppr4	Elp3
Dhx15	Snd1	Lrpprc	Papss1
Eftud2	Ssb	Pten	Pikr4
Hnrnpul1	Ssbp2	Rptor	Prpsap2
Rbm39	Ssbp3	Tsc1	Ruvbl1
Sf3b3		Tsc2	Zyg11b
Snrnp200	**Cluster 7**		
Srrt	Csde1	**Cluster 12**	**Cluster 19**
Usp46	Paip1	Camk1d	Hdac2
Wdr20	Rae1	Clasp1	Kdm1a
Wdr48	Rangap1	Clip2	Ogt
		Rhob	Rbbp7
Cluster 2	**Cluster 8**	Rhot1	Zmym2
Aff4	Ago2		
Mllt1	Cnot1	**Cluster 13**	**Cluster 20**
Mllt3	Fmrp	Prmt5	Ppp1cb
	Pum2	Rpl24	Ppp1r12a
Cluster 3	Rbfox3	Tuba1a	Ppp1r12b
Ctif			Ppp1r12c
Eif2s1	**Cluster 9**	**Cluster 14**	Ppp1r16b
Eif2s2	Atg7	Kcna1	Ppp1r21
Eif2s3x	Brsk2	Kcna2	Ppp2r5c
Eif2s3y	Fga		Ppp2r5d
Eif3b	Fgb	**Cluster 15**	Ppp6r1
Eif3f	Fgg	Actbl2	Ppp6r2
Eif3l	Madd	Actl6b	
Eif4a3	Pck2	Smarce1	**Cluster 21**
Eif4g3	Pex5l		Anapc1
Eif5	Rab11fip5	**Cluster 16**	Anapc2
	Snx2	Cul1	Anapc7
Cluster 4	Snx4	Cul4a	Anapc10
Elavl4	Stxbp5l	Mycbp2	Cdc16
Etf1		Pja1	Cdc23
Upf1	**Cluster 10**	Rlim	Cdc27
Thrap3	Matr3	Rnf14	Cep97
	Pfkfb2	Trim32	Dnaja1
Cluster 5	Pfkl	Uba3	Fzr1
Hnrnpa0	Pfkm	Uba5	Sh3kbp1
Hnrnpl	Pfkp		Znf281
	Pspc1	**Cluster 17**	
		Pi4ka	**Cluster 22**
		Ttc7b	Osbpl1
			Osbpl8

B

DAVID Gene Ontology analysis

Figure 2. Botta et al.

A

IP: Pum1 with RNase – mouse brain

	Input	IgG −	IP:Pum 1 −	IgG +	IP:Pum1 +	
RNase:						
Pum1						120
Pum2						120
Fmrp						71
Ago2						97
Rbfox3						50
Cnot1						240
Mov10						107

B

Reciprocal IP – mouse brain

	Input	IgG	IP:Pum2		Input	IgG	IP:Fmrp
Pum2				Fmrp			
Pum1				Pum1			

	Input	IgG	IP:Ago2		Input	IgG	IP:Rbfox3
Ago2				Rbfox3			
Pum1				Pum1			

	Input	IgG	IP:Cnot1		Input	IgG	IP:Mov10
Cnot1				Mov10			
Pum1				Pum1			

C

IP: Pum1 with RNase – HEK293T cells

	Input	IgG −	IgG +	IP:PUM1 −	IP:PUM1 +	
RNase:						
PUM1						120
PUM2						120
FMRP						71
AGO2						97
RBFOX3	Not detectable in HEK 293T cells					50
CNOT1						240
MOV10						107

Figure 3. Botta et al.

Figure 4. Botta et al.

A Cerebellum
B Hippocampus
C Cortex

○ WT   □ *Pum1⁺ᐟ⁻*   △ *Pum1⁻ᐟ⁻*   ● both sexes   ● male   ● female   *p < 0.05   **p < 0.01   ***p < 0.001

Proteins shown (top to bottom): Pum1 (120), Pum2 (120), Fmrp (71), Ago2 (97), Rbfox3 (50), Cnot1 (240), Mov10 (107), Gapdh (36)

Y-axis: Fold change (Gapdh normal.)

D Cerebellum – *Pum1⁻ᐟ⁻* miRNA sequencing compared to WT mice. Paired t test<1E-05
Y-axis: Fold change. 166 miRNAs

E Cerebellum – *Pum1⁻ᐟ⁻* RNA quantification compared to WT mice. Paired t test<1E-05
Y-axis: Fold change (*Gapdh* norm.). 49 co-regulated targets sorted (left to right) by co-bound miRNAs

Figure 5. Botta et al.

Figure 6. Botta et al.

Figure 7. Botta et al.

Figure 1–figure supplement 1. Botta et al.

Figure 2–figure supplement 1. Botta et al.

Figure 3–figure supplement 1. Botta et al.

Figure 4–figure supplement 1. Botta et al.

Fiigure 4–figure supplement 2. Botta et al.

A

mRNA levels – 10 weeks of age

■ WT ■ *Pum1*⁺/⁻ □ *Pum1*⁻/⁻

Fold change 2⁻ΔΔCt (*Gapdh* normal.)

Cerebellum both sexes · Cerebellum male · Cerebellum female

Hippocampus both sexes · Hippocampus male · Hippocampus female

Cortex both sexes · Cortex male · Cortex female

Pum1 Pum2 Fmr1 Ago2 Rbfox3 Cnot1 Mov10

B

RNA quantification from male and female cerebellum at 10 weeks of age.
Genes plotted here sorted by number of co-bound miRNAs from **Figure 4D** with at least 25% expression change

Male cerebella
Fold change 2⁻ΔΔCt (*Gapdh* normal.)

Female cerebella
Fold change 2⁻ΔΔCt (*Gapdh* normal.)

Figure 6–figure supplement 1. Botta et al.

Figure 7–figure supplement 1. Botta et al.

A

B

C

D

E

F